Sulfate reducing bacteria are active and co-occur with diatoms during a spring

phytoplankton bloom

Siebers, R.[1], Schultz, D.[1], Farza, M. S.[1], Brauer, A.[1], Zühlke, D.[1], Hoff, K.[1], Mücke, P. A[1], Wang,

F.[2], Bernhardt, J.[1], Teeling, H.[2], Becher, D.[1], Riedel, K.[1,3], Urich, T.[1,3], Bengtsson, M. M.[1, 3]

Affiliations

1: Institute of Microbiology, University of Greifswald, Greifswald, Germany

2: Max Planck Institute for Marine Microbiology, Bremen, Germany

3: Institute of Marine Biotechnology, Greifswald, Germany

Corresponding author:

Mia M. Bengtsson

mia.bengtsson@uni-greifswald.de

ABSTRACT

Phytoplankton blooms fuel marine food webs with labile dissolved carbon, but also lead to

the formation of particulate organic matter composed of living and dead algal cells. These

particles are sites of intense algal-bacterial interactions and provide diverse niches for

microbes to thrive. We analyzed co-occurrence networks based on 16S and 18S ribosomal

RNA gene amplicon sequences obtained during a spring phytoplankton bloom in the North

Sea. Particulate fractions larger than 10 µm diameter were collected at almost daily intervals

between early March and late May in 2018. Metaproteomics was used to assess microbial

functional activity at three selected time points during the sampling period. Network analysis

identified two major modules representing bacteria co-occurring with diatoms and with

dinoflagellates, respectively. Bacterial communities associated with these two algal lineages

differed in composition, with diatoms distinctly clustering with known sulfate-reducing

Desulfobacterota as well as with potentially sulfur-oxidizing *Ectothiorhodospiraceae*.

Metaproteomics confirmed the presence of key enzymes involved in dissimilatory sulfate

reduction, a process known to occur in sinking particles at greater depths, but considered of

lesser importance in the photic zone. However, our results suggest presence of sufficiently

anoxic niches in diatom-derived particles, which may have implications for diatom-bacterial

interactions and carbon export during phytoplankton blooms.

INTRODUCTION

Phytoplankton blooms represent primary productivity hotspots, which inject gigatons of

organic carbon into coastal oceans every year. It has been estimated that over 90% of algal-

produced carbon is consumed by heterotrophic bacteria in the immediate vicinity of algal

cells, the phycosphere, during typical bloom situations (Seymour et al. 2017). Some of these

bacteria are directly associated with living algal cells, or with sinking aggregates of senescent

or dead algae, and therefore play an important role in the biological carbon pump. Despite

their importance in global carbon cycling, particle-associated (PA) bacterial communities are

less well understood than their free-living counterparts. In fact, they are often overlooked

due to the common practice of pre-filtration to exclude larger organisms prior to community

composition analyses.

A defining feature of marine particles are the steep chemical and redox gradients that

PA bacterial communities are exposed to compared to free-living, planktonic bacteria. These

gradients have been intensively studied in the context of bathypelagic sinking particles (i.e.,

marine snow), which harbor anoxic micro-niches enabling microbial sulfate reduction and

denitrification (Bianchi et al. 2018, Raven et al. 2021). In the photic zone, these micro-niches

are considered less important, especially for oxygen-producing living algal cells. Aggregates

of living algae are rather known for a diverse microbiome of aerobic, heterotrophic bacteria

that degrade complex algal organic matter, such as polysaccharide-rich exudates. These

bacteria are selected by factors such as host genotype (Ahern et al. 2021), the surrounding

environment (Barreto Filho et al. 2021) and via stochastic processes (Stock et al. 2022). It is

methodologically challenging to study associations between algal and bacterial taxa during

natural phytoplankton blooms by direct observations, (i) due to the transient nature of these

associations, and (ii) due to the innate complexities and dynamics of algal and bacterial

communities during bloom events. Temporal co-occurrence analysis offers an indirect way to

infer algae-bacteria associations that reduces complexity and thereby enables generating

hypotheses about specific interactions and their potential functional implications.

We investigated associations (co-occurrence) of PA bacterial and algal taxa during a
spring phytoplankton bloom in the southern North Sea at the long-term ecological research
site Helgoland Roads (Wiltshire 2010) in the year 2018. We aimed to identify PA bacterial taxa
potentially interacting with diatoms during the bloom. Using 16S and 18S rRNA gene amplicon
data from a well-resolved time series (near daily sampling) collected between early March
and late May in 2018, we constructed co-occurrence networks focusing exclusively on
bacteria-eukaryote co-occurrences (Bengtsson et al. 2017) in particles larger than 10 µm. In
addition, we addressed bacterial functional gene expression by analysis of metaproteomes
(Schultz et al. 2020) from three selected timepoints during the bloom (Fig. 1).
72
RESULTS & DISCUSSION
The spring phytoplankton bloom in 2018 was characterized by an initial dominance of
diatoms (*Bacillariophyceae*), followed by an increase in dinoflagellate (*Dinophyceae*) relative
abundances after the bloom peak (peak in Chlorophyll a concentration, Fig. 1). Co-occurrence
analysis resulted in distinct network modules around diatoms and dinoflagellates. Based on
significant positive correlations (Spearman rho>0.7, corrected p>0.01) between eukaryotic
18S rRNA gene amplicon sequence variants (ASVs) and prokaryotic 16S rRNA gene ASVs,
including all 51 sampled timepoints during the 2018 spring bloom, the network consisted of
three distinct modules (Fig. 2a). Two of these modules were dominated by eukaryotic ASVs
belonging to diatoms and dinoflagellates, respectively, while the third module mostly
contained diatom ASVs and was linked to the dinoflagellate-dominated network module.
These network modules roughly corresponded to the phytoplankton taxa prevalent in the
early stages of the bloom (module I, mainly diatoms), during the late stages of the bloom
(module II, mainly dinoflagellates) and mid-bloom around peak chlorophyll *a* (module III, Fig.
1).
Several ASVs belonging to the *Desulfobacterota* were part of the main diatom -
dominated network module I. Although *Desulfobacterota* have not been identified as
frequent members of diatom microbiomes in either cultures or in the field so far (Helliwell et
al. 2022), a recent global survey of the diatom interactome detected positive correlations
between diatoms and sulfate-reducing bacteria in samples from the Tara Oceans expedition
(Vincent & Bowler 2020). *Desulfobacterota* ASVs were only positively correlated with diatoms
and not with dinoflagellates in our study (Fig. 2b. In addition, ASVs classified as

Ectothiorhodospiraceae also co-occurred with diatoms. These purple sulfur bacteria

(belonging to *Gammaproteobacteria*) are anaerobic to microaerophilic (Imhoff 2022), and can

oxidize H_2S, e.g., produced via dissimilatory sulfate reduction by members of the

Desulfobacterota. This observation is consistent with potential sulfur cycling in close

association with diatoms during this phytoplankton bloom under anoxic to very low oxygen

conditions.

Co-occurrence of diatom- and *Desulfobacterota* rRNA genes does not in itself indicate

that sulfate reduction is taking place in diatom-derived particles in the photic zone. Therefore,

we used metaproteomics to search for actively expressed key functional genes. We detected

4,584 protein groups in > 10 µm filter fractions from three sampled timepoints (see Fig. 1,

Figures S4 and S5) during the bloom. Of these, 22 were classified as belonging to different

Desulfobacterota orders, (Table S1). The relative abundance of predicted *Desulfobacterota*

proteins in the metaproteome was 0.18% (expressed as normalized spectral abundance factor

- NSAF). This order of magnitude corresponded well to the relative abundance of

Desulfobacterota ASVs in the microbiome at the same timepoints (average 0.38%, Fig S1).

Remarkably, no less than 18% of the *Desulfobacterota* metaproteome fraction (in terms of

protein abundance) consisted of key enzymes for dissimilatory sulfate reduction (Fig. 2c).

Both the alpha and the beta subunits of dissimilatory sulfite reductase (DSR) were detected.

Furthermore, adenylyl-sulfate (adenosine-5'-phosphosulfate) reductase (APS reductase), was

also detected in three instances. Only ATP sulfurylase, the third key enzyme of the

dissimilatory sulfate reduction pathway, was not detected. We used a conservative threshold

for protein identification of at least two matching peptides to consider a protein as validly

detected. Therefore, the detection of dissimilatory sulfate reduction key enzymes provides a

strong indication that the diatom-associated *Desulfobacterota* were actively carrying out

sulfate reduction, and thus gaining energy through anaerobic respiration.

Our results highlight the complexity of algal-bacterial interactions during

phytoplankton blooms and corroborate the need to study algae-bacteria particles as spatially

heterogeneous entities. Our observations lead us to hypothesize that living diatom-derived

particles provide sufficiently anoxic niches, at least during parts of the day-night cycle, to

sustain anaerobic sulfur cycling in the photic zone. Diatoms, such as *Thallassiosira* spp. and

Thalassionema spp., which were co-occurring with *Desulfobacterota* in this study (Fig. 2a), are

known to form aggregates (Thornton 2002). For example, several species of *Thalassiosira*

extrude long chitin fibrils, which prevent sinking and bind exopolymeric substances (EPS) also
produced by the algal cells (Den et al. 2023, Herth & Barthlott 1979). This creates a favorable
environment for bacteria to attach, which can in turn stimulate algal EPS production (Gärdes
et al. 2011). EPS makes particles adhesive and thus bacteria can be captured in this sticky EPS
layer. This includes suspended benthic bacteria, which may be abundant in shallow areas such
as the Helgoland sampling site (6-10 m depth depending on tide).

Smaller particles can aggregate to larger particles by collision and adhesion, in
particular during phytoplankon blooms with high particle densities. Such aggregates can
feature high numbers of living, photosynthesizing algal cells (Thornton 2002). These cells
produce ample oxygen during the day, but at night respiration by the algal cells and their
surrounding bacteria may deplete oxygen sufficiently for low oxygen or even anaerobic micro-
niches to form. Our data suggest that these micro-niches can support microbial sulfate
reduction in diatom microbiomes. The latter may thus be as complex as those of animals and
other multicellular organisms, featuring distinct micro-niches with sub-microbiomes
(analogous to e.g., human skin vs. gut microbiomes), whose compositions depend on the
chemical environment on a small spatial scale. Anoxic micro-niches in diatom-derived
aggregates may even affect the biological carbon pump insofar, as sulfate reduction can
chemically alter particulate organic matter, rendering it resistant to microbial degradation
(Raven et al. 2021).

Temporal associations, such as detected by our co-occurrence network analyses, have
to be interpreted with caution as additional variable factors that were not taken into account
may influence the resulting correlations. For example, we cannot completely rule out an
influence of underlying sediment microbiomes at the rather shallow sampling site, which may
become resuspended during times of heavy wind thereby introducing anaerobic microbes to
the pelagic environment. However, we did not find a clear link between wind conditions,
Secchi depth, and *Desulfobacterota* in this study (Fig. S2). Further, the diatom taxa co-
occurring with *Desulfobacteria* were also primarily classified as common pelagic genera such
as *Thalassiosira* and *Thalassionema*. Limitations notwithstanding, co-occurrence network
analysis provides a tool for formulating hypotheses about interactions between taxa that can
be tested experimentally in future investigations.

We showed that there was a temporal association between diatom and bacterial taxa
belonging to the *Desulfobacterota* during a North Sea phytoplankton spring bloom and that

key enzymes involved in sulfate reduction by these bacteria were expressed during the same
bloom. However, a physical association between these bacteria and diatoms was not proven
by our approach and would require *in situ* microscopic investigation using for example taxon-
or gene- specific fluorescent probes. Likewise, it has yet to be investigated, whether such
temporal and physical associations are annually recurrent, under what specific conditions
they occur, and which specific diatom taxa are involved. The presence of anaerobic niches
inside aggregates of living diatoms provides a new facet of algal-bacterial-organic matter
interactions in the photic zone, with potential implications for carbon export from
phytoplankton blooms.

DATA AVAILABILITY

Metagenome raw reads and assembled contigs used for metaproteome assignment are
publicly available at the European nucleotide archive under the project accession number
PRJEB38290. 16S- and 18S-rRNA gene amplicon data are also available via ENA under the
project accession number PRJEB51816. Mass spectrometry proteomics data have been
deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2022),
repository with dataset identifier PXD035982. Original mass spectrometry proteomics data
can be accessed for review (**Username:** reviewer_pxd035982@ebi.ac.uk **Password:**
umim4qCW) and will be made publically available upon manuscript acceptance.

ACKNOWLEDGEMENTS
We thank the team from the Biological Station Helgoland (BAH-AWI, Germany) and the
captains and crews from FS Aade and FS Uthörn for help with sampling, analyses, logistics,
and providing lab space. The Helgoland Time Series is supported by the Helmholtz Association
as an LK-II performance category program. This study was funded by the Deutsche
Forschungsgemeinschaft (DFG) in the framework of the research unit FOR2406
'Proteogenomics of Marine Polysaccharide Utilization (POMPU)' by grants of KR/MB (RI
969/9-2), DB (BE 3869/4-2) and HT (TE 813/2-2).

CONFLICT OF INTEREST

The authors declare no conflicts of interest

REFERENCES

Ahern OM, Whittaker KA, Williams TC, Hunt DE, Rynearson TA. Host genotype structures the
microbiome of a globally dispersed marine phytoplankton. Proc Natl Acad Sci U S A 2021.
doi:10.1073/pnas.2105207118.

Barreto Filho MM, Walker M, Ashworth MP, Morris JJ. Structure and Long-Term Stability of the
Microbiome in Diverse Diatom Cultures. Microbiol Spectr. 2021;9(1):e0026921.
doi:10.1128/Spectrum.00269-21.

Bengtsson MM, Bühler A, Brauer A, Dahlke S, Schubert H, Blindow I. Eelgrass Leaf Surface Microbiomes
Are Locally Variable and Highly Correlated with Epibiotic Eukaryotes. Front Microbiol.
2017;8:1312. doi:10.3389/fmicb.2017.01312.

Bianchi D, Weber TS, Kiko R, Deutsch C. Global niche of marine anaerobic metabolisms expanded by
particle microenvironments. Nature Geosci. 2018;11(4):263–8. doi:10.1038/s41561-018-0081-0.

Den TQ, Neu TR, Sultana S, Giebel HA, Simon M, Billerbeck S. Distinct glycoconjugate cell surface
structures make the pelagic diatom Thalassiosira rotula an attractive habitat for bacteria. Journal
of Phycology 2023. http://doi.org/10.1111/jpy.13308

Gärdes A, Iversen MH, Grossart H-P, Passow U, Ullrich MS. Diatom-associated bacteria are required
for   aggregation   of   *Thalassiosira   weissflogii*.   ISME   J.   2011;5(3):436–45.
doi:10.1038/ismej.2010.145.

Helliwell KE, Shibl AA, Amin SA. The Diatom Microbiome: New Perspectives for Diatom-Bacteria
Symbioses. In: Falciatore A, Mock T, eds. The Molecular Life of Diatoms. Cham: Springer
International Publishing; 2022. p. 679–712. doi:10.1007/978-3-030-92499-7_23.

Herth W, & Barthlott W. The site of beta-chitin fibril formation in centric diatoms. I. Pores and fibril
formation. Journal of Ultrastructure Research, 1979; 68(1), 6–15. http://doi.org/10.1016/s0022-
5320(79)90137-0

Imhoff JF, Kyndt JA, Meyer TE. Genomic Comparison, Phylogeny and Taxonomic Reevaluation of the
Ectothiorhodospiraceae and Description of *Halorhodospiraceae* fam. nov. and *Halochlorospira*
gen. nov. Microorganisms 2022. doi:10.3390/microorganisms10020295.

Raven MR, Keil RG, Webb SM. Microbial sulfate reduction and organic sulfur formation in sinking
marine particles. Science. 2021;371(6525):178–81. doi:10.1126/science.abc6035.

Schultz D, Zühlke D, Bernhardt J, Francis TB, Albrecht D, Hirschfeld C, et al. An optimized
metaproteomics protocol for a holistic taxonomic and functional characterization of microbial

communities from marine particles. Environ Microbiol Rep. 2020;12(4):367–76.
doi:10.1111/1758-2229.12842.

Seymour JR, Amin SA, Raina J-B, Stocker R. Zooming in on the phycosphere: the ecological interface
for phytoplankton-bacteria relationships. Nat Microbiol. 2017;2:17065.
doi:10.1038/nmicrobiol.2017.65.

Stock W, Willems A, Mangelinckx S, Vyverman W, Sabbe K. Selection constrains lottery assembly in
the microbiomes of closely related diatom species. ISME COMMUN. 2022. doi:10.1038/s43705-
022-00091-x.

Thornton DCO. Diatom aggregation in the sea: mechanisms and ecological implications. Euro. J.
Phycol. 2002;37(2):149–61. doi:10.1017/S0967026202003657.

Vincent F, Bowler C. Diatoms Are Selective Segregators in Global Ocean Planktonic Communities.
mSystems 2020. doi:10.1128/mSystems.00444-19.

Wiltshire KH, Kraberg A, Bartsch I, Boersma M, Franke H-D, Freund J, et al. Helgoland Roads, North
Sea: 45 Years of Change. Estuaries and Coasts. 2010;33(2):295–310. doi:10.1007/s12237-009-
9228-y.

FIGURE CAPTIONS

Figure 1: Diatoms and dinoflagellates display opposing dynamics during the spring
phytoplankton bloom 2018.

Particles larger than 10 μm were collected via filtration from 01 of March to 29 of May 2018
and microbial communities were analyzed using 16S and 18S rRNA gene amplicon sequencing
and metaproteomics. Algal biomass (HPLC chlorophyll *a* measurements) peaked around the
26th of April (Julian day 116). Metaproteomics data from three selected timepoints displayed
a dominance of eukaryotic proteins. Diatoms and dinoflagellates were the most abundant
phytoplankton lineages as detected via 18S rRNA gene amplicon sequences and varied along
the course of the bloom with dinoflagellates being more abundant after the chlorophyll *a*
peak.

Figure 2: Diatoms distinctively co-occurred with *Desulfobacterota* and presence of key enzymes of the sulfur cycle was confirmed by metaproteomics.

a) A network analysis of 18S rRNA and 16S rRNA gene ASV co-occurrences resulted in two major network modules containing diatom [I] and dinoflagellate [II] 18S ASVs, respectively, as well as one mixed module [III]. The network was calculated based on Spearman correlations (r >0.7, p < 0.01) between 18S ASVs and 16S ASVs exclusively. Several bacterial groups (circles) were associated with all modules, yet *Desulfobacterota* (yellow) were only associated with the diatom-dominated module. The sizes of the symbols correspond to the number of significant correlations of the nodes (degree). b) The bacterial ASVs temporally associated with diatoms and dinoflagellates represented distinct taxa. *Desulfobacterota* (yellow) as well as potentially sulfur-oxidizing *Ectothiorodospiraceae* were positively associated with diatoms but not with dinoflagellates. c) The pie chart displays relative abundances for all 22 protein groups classified as belonging to *Desulfobacterota* during all three timepoints sampled for metaproteomics. Proteins involved in sulfate reduction made up 18% of the metaproteome fraction (based on NSAF values, or 5 of 22 (23%) of proteins) assigned to *Desulfobacterota*. Of these, five protein groups represented enzymes involved in dissimilatory sulfate reduction (yellow circles). n.d. = not detected.

275 # Supplementary information

276
277

284
285
286

287 Supplementary methods
288
Sampling and filtration
For 16S and 18S rRNA gene amplicon sequencing and metagenome analysis, 40 L of
subsurface seawater (1 m depth) were sampled in the morning at near daily intervals between
beginning of March and end of May in 2018 at the station "Kabeltonne" (50° 11.3' N, 7° 54.0'
E) near Helgoland in the south-eastern North Sea. Plankton biomass from 1 L of seawater was
filtered using 10 μm pore size polycarbonate membrane filters (47 mm diameter, Millipore,
Schwalbach, Germany) to separate PA microbes from smaller bacterial fractions. For
metaproteomics analysis, 100 - 140 L of subsurface seawater were sampled at three selected
timepoints during the bloom (Julian days 107, 128 and 144, representing early-, mid- and late
bloom phases) and filtered onto 10 μm polycarbonate membrane filters (142 mm diameter,
Millipore). In order to maximize biomass harvest while avoiding clogging, filtered volumes
varied between 15 and 30.5 L per filter.
301
rRNA gene amplicon sequencing
For amplicon sequencing, a total of 51 samples were collected and analyzed. DNA was
extracted from the filters using the Qiagen DNeasy Power soil Pro kit (Qiagen, Hilden,
Germany) according to the manufacturer's instructions. Dislocation of microbial cells from
the filters and mechanical lysis were achieved by bead beating in a FastPrep 24 5G (MP
Biomedicals, Irvine, CA, USA). DNA concentrations were measured at a Qubit 3.0 fluorometer
(Invitrogen, Carlsbad, CA, USA). Extracted DNA was amplified with primer pairs targeting the
V4 region of the 16S rRNA gene [515f: 5 -GTGYCAGCMGCCGCGGTAA-3 , 806r: 5 -
GGACTACNVGGGTWTCTAAT-3' (Walters et al., 2016)] and the V7 region of the 18S rRNA gene
[F-1183mod: 5'-AATTTGACTCAACRCGGG-3', R-1443mod: 5'-GRGCATCACAGACCTG-3'] (Ray et
al. 2016) coupled to custom adaptor-barcode constructs. PCR amplification and Illumina
MiSeq (Illumina, San Diego, CA, USA) library preparation and sequencing (V3 chemistry) was
carried out by LGC Genomics (LGC Genomics,Berlin, Germany).
Sequence reads free of adaptor and primer sequence remains were processed using the
DADA2 package in R (version 1.2.0, Callahan et al., 2016). In summary, forward and reverse
Illumina MiSeq reads were truncated to 200 bp, filtered (maxEE = 2, truncQ = 2, minLen =
175), dereplicated and error rates were estimated using the maximum possible error estimate
from the data as an initial guess. Sample sequences were inferred, paired forward and reverse
reads were merged and chimeric sequences were removed using the removeBimeraDenovo
function. The resulting amplicon sequence variants (ASVs) were taxonomically classified using
the Silva database (nr 99 v 138.1) for 16S rRNA and the PR2 database (version 4.13, minboot:
50) for 18S rRNA using the build-in RDP classifier. 16S rRNA gene amplicon reads classified as
chloroplasts and mitochondria, as well as 18S rRNA gene reads classified as *Metazoa*
(Zooplankton) were removed prior to the downstream analyses.
Co-occurrence networks were generated in R using Spearman rank correlation, as
described previously (Bengtsson et al. 2017, correlation coefficient >0.7, adjusted p<0.01).
We considered exclusively correlations between 18S and 16S ASVs and the final network was
plotted using the igraph R package.
330
Construction of a metagenome-based database

Illumina sequencing was performed at the Max Planck Genome Centre Cologne on extracted
DNA using the HiSeq 2500 platform and 2 x 150 bp chemistry. Filtration and trimming of reads
were done as described previously (Francis et al. 2021). Read quality for each sample was
confirmed using FastQC v0.11.9 (Andrews 2014). Assembly was performed with MEGAHIT
v1.2.9 (Li et al. 2015) (kmer length: 21). Contigs below 2.5 kbp were removed using anvi-
script-reformat-fasta from within Anvi'o v6.2 (Eren et al. 2015). Genes were predicted using
Prodigal v2.6.3 (Hyatt et al. 2010) as implemented in the Prokka v1.11 annotation pipeline
(Seemann et al. 2014). These genes sequences were subsequently combined with a list of
common contaminants to generate a database for peptide-spectrum-matching (PSM) after
elimination of redundant sequences (97% redundancy) with CD-Hit (Fu et al. 2012, Li and
Godzik 2006).
343
Metaproteomics
Proteins were extracted from the filters of the 17th of April, 8th of May and 24th of May 2018
(Julian days 107, 128 and 144) as described previously (Schultz et al. 2020, Schultz 2022) and
analyzed with liquid chromatography-tandem mass spectrometry in triplicates. Briefly,
proteins were extracted from the filters via bead-beating followed by acetone precipitation.
The extracts were separated and fractionated by 1D SDS-PAGE and in-gel trypsin digested.
After desalting and concentration of the peptides using C18 Millipore® ZipTip columns,
samples were measured with an Orbitrap Velos™ mass spectrometer (ThermoFisher
Scientific, Waltham, MA, USA). After conversion into mgf file format using MS convert in
ProteoWizard (Palo Alto, CA, USA), spectra were matched against a metagenome-based
database containing 14,764,755 entries. Mascot (Matrix Science, London, UK) and Scaffold
(Proteome Software, Portland, OR, US) were used for peptide-spectrum-matching, protein
identification and protein grouping. Instead of setting an FDR-threshold, identification of
protein groups was based on number of peptide matches with a minimum of two, a protein
threshold of 99% and a peptide threshold of 95%. Identified protein groups were annotated
via Prophane v6.2.3 (Schiebenhoefer et al., 2020), using the Uniprot-TrEMBL (as of September
2021) and NCBI nr (as of February 2022) databases for taxonomic and EggNog v5.0.2 and
Tigrfams 15 databases for functional annotation with Prophane default settings. Relative
abundances were calculated as Normalized Spectral Abundance Factor (NSAF) values using
the quantification method "max_nsaf" integrated in Prophane (Schiebenhoefer et al., 2020).
Briefly, SAF values were calculated by dividing exclusive unique spectrum counts for each
protein group by protein length of the longest sequence in that protein group. For
normalization, SAF values were then divided by the sum of all SAF values of the sample.
367

Fig. S1 SSU rRNA gene amplicon sequencing reveals taxonomic shifts during the course of the bloom. Eukaryote phylum distribution via 18S rRNA gene amplicon sequencing shows dominance of *Bacillariophyta* at the beginning of the phytoplankton bloom, shifting towards dominance of *Dinophyceae* in the last third of the surveillance period. 16S rRNA gene amplicon data revealed *Proteobacteria* and *Bacteroidota* as abundant bacterial phyla during the course of the phytoplankton bloom. Chloroplast and mitochondria sequence variants were removed from the 16S rRNA gene dataset prior to normalization.

379
Fig. S2 Relative abundance of *Desulfobacterota* in relation to wind conditions (a), Secchi
depth and chlorophyll *a* levels (b) during the 2018 spring bloom. Relative abundances of
Desulfobacterota are based on 16S rRNA gene amplicon ASVs. Wind maxima are based on
DWD Helgoland station via wetterkontor.de.

385 **Fig. S3 Association of *Desulfobacterota* and *Ectothiorodospiraceae* with diatoms.**
386 Distribution of bacterial ASVs based on 16S rRNA gene amplicon data. Temporal associations
387 to diatoms and dinoflagellates were taxonomically distinct.

389
Fig. S4 Taxonomic distribution as detected by metaproteome analysis. Annotation based on
the Trembl database of September 2021. Strong dominance of Bacillariophyta shifts towards
increased abundance of bacterial proteins and especially *Proteobacteria* over time.
Dinophyceae could not be identified using NR or Trembl, which reflects an
underrepresentation of annotated dinoflagellate genome sequences in public databases,
making peptide-spectrum-matching challenging.
396

397
Fig. S5 Protein functions as detected by metaproteomics. Annotation was performed using
EggNog v5.0.2 via Prophane. Metaproteome analysis was performed for three timepoints.
400

Supplementary table

Table T1 Five key enzymes associated with the sulfur cycle detected with metaproteomics.
Adenylylsulfate-reductase and dissimilatory sulfite reductase assigned to three
Desulfobacterota species could be detected. Functional annotation was based on EggNog
v5.0.2, and the assignment of taxonomy was based on Swissprot (as of Sept. 2021), Trembl
(as of Sept. 2021), and NCBI nr (as of Feb. 2022) database annnotations

#pg	type	NSAF			Tax_summary
		Julian Day 107	Julian Day 128	Julian Day 144	
112		0	0	0.00016602	*Desulfobublaceae* sp.
140		0	0	0.00049134	*Sorangium cellulosum*
182	Adenylylsulfate reductase	0	0	0.00015253	*Desulfobublaceae* sp.
239		0.0002743	7.6347E-05	0.00030024	*Anaeromyxobacter* sp. (strain Fw109-5)
293		5.7524E-05	7.7272E-05	0	*Desulfosarcina alkanivorans*
411		0	0	0.00025035	*Desulfosarcina alkanivorans*
440	Dissimilatory sulfite reductase	0	0	0.00013727	*Desulfovibrio vulgaris*
540		0.00023951	0	0	*Desulfobulbaceae* sp.
576	Dissimilatory sulfite reductase	0	0	0.00018468	*Desulfovibrio vulgaris*
666		0	0	0.00015508	*Wolinella succinogenes*
743		5.4294E-05	1.8454E-05	8.2017E-05	*Geobacter uraniireducens*
810		0.0002416	0.00015172	0.00011497	*Desulfatibacillum aliphaticivorans*
954		0	0	0.00016176	*Desulfurivibrio alkaliphilus*
960		0	0.00017834	0.00034704	*Desulfobacterales* sp.
1214		0	0	8.941E-05	*Desulfatibacillum aliphaticivorans*
1321		0	0.00010197	0.00015095	*Desulfosarcina alkanivorans*
1513		0	0	0.00011631	*Wolinella succinogenes*
2117		0	0	0.00026687	*Desulfobacterales* sp.
2147	Adenylylsulfate reductase	0	0	0.00037822	*Desulfopila inferna*
2297		0	2.6317E-05	4.2269E-05	*Deltaproteobacteria* sp.
2745		0	0	0.00026719	*Desulfobulbaceae* sp.
2868	Adenylylsulfate reductase	0	0	0.00011735	*Desulfobulbaceae* sp.

Supplementary references

Andrews S, FastQC: a quality control tool for high throughput sequence data. 2014. Available online
at: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (accessed April 03, 2023).

Bengtsson MM, Bühler A, Brauer A, Dahlke S, Schubert H, Blindow I. Eelgrass Leaf Surface
Microbiomes Are Locally Variable and Highly Correlated with Epibiotic Eukaryotes. Front
Microbiol. 2017;8:1312. doi:10.3389/fmicb.2017.01312.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods.
2016;13(7):581–3. doi:10.1038/nmeth.3869.

Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. Anvi'o: an
advanced analysis and visualization platform for 'omics data. PeerJ. 2015;3:e1319.
doi:10.7717/peerj.1319.

Francis TB, Bartosik D, Sura T, Sichert A, Hehemann J-H, Markert S, et al. Changing
expression patterns of TonB-dependent transporters suggest shifts in polysaccharide
consumption over the course of a spring phytoplankton bloom. ISME J.
2021;15(8):2336–50. doi:10.1038/s41396-021-00928-8.

Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012;28(23):3150–2.
doi:10.1093/bioinformatics/bts565.

Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene
recognition and translation initiation site identification. BMC Bioinformatics.
2010;11:119. doi:10.1186/1471-2105-11-119.

Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for
large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.
2015;31(10):1674–6. doi:10.1093/bioinformatics/btv033.

Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. Bioinformatics. 2006;22(13):1658–9.
doi:10.1093/bioinformatics/btl158.

Ray JL, Althammer J, Skaar KS, Simonelli P, Larsen A, Stoecker D, et al. Metabarcoding and
metabolome analyses of copepod grazing reveal feeding preference and linkage to
metabolite classes in dynamic microbial plankton communities. Mol Ecol.
2016;25(21):5585–602. doi:10.1111/mec.13844.

Schiebenhoefer H, Schallert K, Renard BY, Trappe K, Schmid E, Benndorf D, et al. A complete
and flexible workflow for metaproteomics data analysis based on
MetaProteomeAnalyzer and Prophane. Nat Protoc. 2020;15(10):3212–39.
doi:10.1038/s41596-020-0368-7.

Schultz D, Zühlke D, Bernhardt J, Francis TB, Albrecht D, Hirschfeld C, et al. An optimized
metaproteomics protocol for a holistic taxonomic and functional characterization of
microbial communities from marine particles. Environ Microbiol Rep. 2020;12(4):367–
76. doi:10.1111/1758-2229.12842.

Schultz, D. (2022). Mechanisms of polysaccharide degradation of particle-associated
microbial communities. PhD thesis, University of Greifswald.

Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068–9. doi:10.1093/bioinformatics/btu153.

Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM, et al. Substrate-
controlled succession of marine bacterioplankton populations induced by a
phytoplankton bloom. Science. 2012;336(6081):608–11. doi:10.1126/science.1218344.

Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, et al. Improved
Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker
Gene Primers for Microbial Community Surveys. mSystems 2016.
doi:10.1128/mSystems.00009-15.
461
462
463
464