DISCOVERY REPORTS

A remodeled RNA polymerase II complex catalyzing viroid RNA-templated

transcription

Short title: A remodeled Pol II for transcription

Shachinthaka D. Dissanayaka Mudiyanselage[1], Junfei Ma[1], Tibor Pechan[2], Olga

Pechanova[2], Bin Liu[1], Ying Wang[1,*]

[1] Department of Biological Sciences, Mississippi State University, Mississippi State, MS

39762

[2] Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University,

Mississippi State, MS 39762

[*] Address correspondence to wang@biology.msstate.edu

Keywords: viroid, RNA-templated transcription, Pol II, transcription cofactors

Abstract

Viroids, a group of plant pathogens, are mysterious subviral agents composed of

single-stranded circular noncoding RNAs. Nuclear-replicating viroids exploit host RNA

polymerase II (Pol II) activity for transcription from circular RNA genome to minus-strand

intermediates, a classic example illustrating the intrinsic RNA-dependent RNA

polymerase activity of Pol II. The mechanism for Pol II to accept single-stranded RNAs

as templates for transcription remains poorly understood. Here, we reconstituted a robust

in vitro transcription system and demonstrated that Pol II also accepts the minus-strand

viroid RNA template to generate sense-strand RNAs. Based on this reconstituted system,

we purified the Pol II complex on RNA templates for nano-liquid chromatography-tandem

mass spectrometry analysis. We identified a remodeled Pol II missing Rpb5, Rpb6, and

Rpb9, which contrasts to the canonical 12-subunit Pol II or the 10-subunit Pol II core on

DNA templates. This remodeled Pol II is active in transcription with the aid of TFIIIA-7ZF.

Interestingly, this remodeled Pol II appears not to require other canonical general

transcription factors (such as TFIIA, TFIIB, TFIID, TFIIE, TFIIF, TFIIH, and TFIIS),

indicating a different mechanism/machinery regulating RNA-templated transcription.

Transcription elongation factors, such as FACT complex, PAF1 complex, and SPT6, were

also absent in the transcription complex on RNA templates. We further analyzed the

critical zinc finger domains in TFIIIA-7ZF for aiding Pol II activity on RNA template and

revealed the first three zinc finger domains pivotal for template binding. Collectively, our

data illustrated a distinct organization of Pol II complex on viroid RNA templates, providing

new insights into the evolution of transcription machinery, the mechanism of RNA-

templated transcription, as well as viroid replication.

Introduction

Viroids are circular noncoding RNAs that infect crop plants [1, 2]. After five decades of studies, the host machinery for viroid infection has not been fully elucidated [1-3]. There are two viroid families, *Avsunviroidae* and *Popspiviroidae* [4, 5]. Members of *Pospivoirdae* replicate in the nucleus via the rolling circle mechanism (S1 Figure) and rely on the enzymatic activity of RNA polymerase II (Pol II) [1-3]. Specifically, ample evidence support that Pol II activity is critical for the synthesis of oligomer minus-strand or (-) intermediates using viroid circular genomic RNA as templates [1, 6]. However, it remains controversial whether Pol II also uses (-) oligomers as templates for transcription [1, 6].

By and large, RNA polymerases catalyze transcription using DNA templates, which is a fundamental process of life. RNA polymerases are facilitated by a group of general transcription factors to achieve highly regulated transcription, from initiation to elongation and then to termination. Taking Pol II as an example, this 12-subunit complex functions in concert with general transcription factors TFIIA, TFIIB, TFIID, TFIIE, TFIIF, and TFIIH during transcription initiation around DNA promoter regions [7-10]. In general, a minimal system for the promoter-driven transcription requires Pol II and five general transcription factors (TFIIB, TFIID, TFIIE, TFIIF, and TFIIH)[11, 12]. Interestingly, the 10-subunit Pol II core (without Rpb4/Rpb7 heterodimer) is sufficient for transcription elongation [13, 14]. Transcription elongation also requires multiple factors, including TFIIS, TFIIF, DSIF, PAF1 (RNAPII-associated factor 1) complex (PAF1-C), FACT complex (Histone Chaperone), SPT6, etc [15-17].

Since 1974, RNA polymerases have been found to possess intrinsic RNA-dependent RNA polymerase (RdRp) activity to catalyze RNA polymerization using RNA templates

[18]. This intrinsic RdRp activity of RNA polymerases was found in bacteria and

mammalian cells, as well as exploited by subviral pathogens (i.e., viroids and human

hepatitis delta virus (HDV)) for propagation [19-21]. Pol II transcription using viroid or HDV

RNA templates can yield RNA products over 1,000 nt in cells, comparable to some

products from DNA templates. To ensure such efficient transcription, specific factors are

needed. HDV encodes an S-HDAg to promote Pol II activity on its RNA template [22].

Using potato spindle tuber viroid (PSTVd) as a model, we showed that an RNA-specific

transcription factor (TFIIIA-7ZF with seven zinc finger domains) interacts with Pol II and

specifically enhances Pol II activity on circular genomic RNA template [23, 24]. However,

it remains unclear how S-HDAg or TFIIIA-7ZF functions with Pol II for RNA-templated

transcription.

Biochemically reconstituted systems have been successfully used to characterize the

required factors and functional mechanisms underlying DNA-dependent transcription [25].

However, reconstituted transcription systems using RNA templates often exhibited poor

activity. For example, all currently available in vitro transcription (IVT) systems using HDV

template have the premature termination issue generating products less than 100 nt [26,

27], which may not reflect the transcription process in cells. We recently established an

IVT system for PSTVd [23, 24] that can generate longer-than-unit-length products (more

than 360 nt), mimicking the replication process in cells [28].

Using our IVT system, we first confirmed that Pol II and TFIIIA-7ZF function together

in transcribing (-) PSTVd oligomers to (+) oligomers. Interestingly, we found that the Pol

II complex remaining on (-) PSTVd RNA template has a distinct composition as compared

with the 12-subunit Pol II or the 10-subunit Pol II core, via nano-liquid chromatography-

tandem mass spectrometry analysis (nLC-MS/MS). Rpb5, Rpb6, and Rpb9 were absent

in the remodeled Pol II. Interestingly, Rpb9 is responsible for the fidelity of Pol II

transcription. Thus, the absent of Rpb9 may explain the much higher mutation rate of

viroid RNA-templated transcription catalyzed by Pol II. Several critical elongation factors

for DNA templates, such as the PAF1 complex and SPT6, were also absent in the

transcription complex on RNA templates. More importantly, essential general

transcription factors (including TFIIA, TFIIB, TFIID, TFIIE, TFIIF, TFIIH, and TFIIS) were

all absent in the transcription complex on PSTVd RNA template, clearly demonstrating

the distinct regulations between DNA-dependent and RNA-templated transcription. This

distinct Pol II retains the catalytic activity to generate (+)-PSTVd oligomers with the aid of

TFIIIA-7ZF. We further showed that nearly all seven zinc finger domains of TFIIIA-7ZF

are critical for function. In particular, the first three zinc fingers are pivotal for binding with

RNA templates. Our findings provide new insights into the organization of Pol II complex

on RNA templates, which has profound implications for understanding RNA-templated

transcription as well as viroid transcription and its high mutation rates.

Results

Because PSTVd replication from circular genomic RNA to (-) oligomers and then to

plus-strand or (+) oligomers is a continuous process, failure to tease apart each step

resulted in controversial data, as evidenced by previous reports [34,41]. To understand

whether Pol II can catalyze transcription using (-) viroid oligomers, we established a

reconstituted in vitro transcription system using partially purified Pol II from wheat germ

[24, 29] and the (-) PSTVd dimer as RNA template. Based on our previous report [24] and

the rolling circle replication model (S1 Figure), the linear (+) PSTVd (i.e., the product from

IVT assay) cannot serve as a template for further transcription. Therefore, this IVT assay

will only focus on the transcription step from (-) PSTVd dimer to (+) PSTVd dimer. As

shown in Figure 1a, Pol II has a weak activity in transcribing (-) PSTVd template to full

length (+) PSTVd dimer, which is more than 700 nt in length. We then tested the role of

the RNA-specific transcription factor (TFIIIA-7ZF) in this transcription reaction by

supplying various amounts of the protein. As shown in Figure 1a, TFIIIA-7ZF can

significantly increase the RdRp activity (more than 10-fold) of Pol II on the (-) PSTVd

dimer template. Therefore, our results indicate that Pol II can accept (-) PSTVd oligomers

as a template for transcription, expanding the known natural RNA templates for DdRPs.

We then analyzed the composition of Pol II complex on the viroid RNA template using

RNA-based affinity purification. Briefly, a desthiobiotinylated cytidine (Bis)phosphate was

ligated to the 3'end of (-) PSTVd dimer, which was then mounted to magnetic streptavidin

beads. After sequential supplying of TFIIIA-7ZF and partially purified Pol II, the magnetic

beads were washed before elution. Through silver staining, common patterns and distinct

bands can be observed between elution fraction and partially purified Pol II (S2 Figure),

implying that certain factors may be enriched by or removed from RNA templates. We

then performed nLC-MS/MS to reveal the protein factors in partially purified Pol II as well

as the Pol II complex remaining on the RNA template (proteins identified in each replicate

are listed in S1-S6 Datasets). We identified 11 out of 12 Pol II subunits in partially purified

Pol II with high confidence (false discovery rate below 0.01, identified by a minimum of 2

peptides, and 2 peptide-spectrum matches) in all three replicates (Figure 1b and 1c and

S1 Table). The smallest subunit Rpb12 (~6 kDa) was absent, which is possibly caused

by sample loss during the size cut-off enrichment of samples for nLC-MS/MS. This issue

has been seen in another study [17]. Interestingly, only six subunits were confidently

identified in the Pol II complex remaining on the RNA template with high confidence: Rpb1,

Rpb2 and Rpb8 were found in all three replicates while Rpb3, Rpb10, and Rpb11 were

found in two out of three replicates (Figure 1c and S1 Table). Rpb5 can only be detected

in one replicate of Pol II remaining on the RNA template (Figure 1c). Therefore, it is less

likely to participate in the Pol II complex on the RNA template. Rpb1 and Rpb2 form the

catalytic core of Pol II, while Rpb3, Rpb10, Rpb11 form a subassembly group all critical

for Pol II assembly [30]. Since Rpb12 is also a conserved subunit in this subassembly

group, we speculate that Rpb12 is also present in the Pol II complex remaining on RNA

templates. Rpb8 is an auxiliary factor [30]. Given that the Pol II complex remaining on the

RNA template has a distinct composition, we termed it remodeled Pol II hereafter. To test

whether the remodeled Pol II has any catalytic activity, we repeated the RNA-based

affinity purification followed by IVT. As shown in Figure 1d, this remodeled Pol II indeed

possessed transcription activity in generating full-length (+) PSTVd oligomers comparable

to partially purified Pol II. It is noteworthy that the product is more than 700 nt in length,

comparable to the products generated in cells.

Besides the Pol II complex in the partially purified samples, we also found the

presence of several general transcription factors and transcription elongation factors for

DNA-dependent transcription. However, they were all absent in the remodeled Pol II in

the nLC-MS/MS analysis. For example, we found TFIIF in all three repeats of partially

purified Pol II but could not confidently detect it in any of the remodeled Pol II samples

(Figure 2 and S1 Table). In addition, TFIIE was found in two out of three repeats of

partially purified Pol II but could not be confidently detected in any of the remodeled Pol

II samples (S1 Table). Therefore, TFIIE and TFIIF are likely not required for viroid RNA-

templated transcription. Noteworthy is that all the rest of the canonical general

transcription factors (including TFIIA, TFIIB, TFIID, TFIIH, and TFIIS) were absent in our

partially purified or remodeled Pol II. SPT6 is a histone chaperone that interacts with

Rpb4/Rpb7 heterodimer during transcription elongation on DNA templates [16]. All

Rpb4/Rpb7 and SPT6 were absent in the remodeled Pol II sample (Figure 2 and S1

Table). The FACT complex, including SPT16 and SSRP1-B, is also a histone chaperone

assisting Pol II elongation on DNA templates. The FACT complex does interact with Pol

II directly [31], and most of the components also absent in the remodeled Pol II sample

(except for the significantly reduced CTR9 in two out of three replicates)(Figure 2 and S1

Table). The PAF1-C, including PAF1, CTR9, LEO1, and RTF1, regulates transcription-

coupled histone modifications. PAF1-C components extensively interact with Pol II.

PAF1-LEO1 is anchored to the external domains of Rpb2. RTF1 is in proximity to PAF1.

The trestle helix in CTR9 binds to Rpb5 and the surrounding region, while the

tetratricopeptide repeats interact with Pol II around Rpb11 and Rpb8 [16, 32]. Similarly,

the PAF1-C was absent in the remodeled sample (Figure 2 and S1 Table).

Since TFIIIA-7ZF is critical for Pol II to perform transcription using RNA templates, we

attempted to identify the functional domain(s) of TFIIIA-7ZF. TFIIIA-7ZF has seven C2H2

type zinc finger domains. We mutated each zinc finger domain by changing the first

histidine in the C2H2 domain to asparagine, which is commonly used to disrupt the local

structure of a C2H2 motif [33, 34]. We then used those variants for the IVT assay. As

shown in Figure 3, all mutants exhibited greatly reduced activity in aiding Pol II

transcription on viroid RNA templates. Mutants *zf1*, *zf2*, *zf3*, and *zf6* completely lost the

activity, while mutants *zf4*, *zf5*, and *zf7* can increase Pol II activity about two-fold (Figure

3), which is a much weaker activity as compared with more than 10-fold increase

stimulated by wildtype (WT) TFIIIA-7ZF (Figure 1A). We then performed the RNA-based

affinity purification assay using TFIIIA-7ZF mutants. Since Pol II itself has viroid RNA

binding affinity [35], it is not surprising to observe no difference in the amount of the

remodeled Pol II on RNA templates with or without the presence of WT TFIIIA-7ZF (Figure

4). Interestingly, *zf1*, *zf2*, and *zf3* exhibited significantly reduced affinity to (-) PSTVd dimer

RNA, which also led to significantly reduced Pol II remaining on RNA templates. The

amount of the remodeled Pol II was also reduced, to a lesser extent, in the presence of

zf4, *zf5*, or *zf7*, which explains the reduced transcription activity in the corresponding IVT

reactions.

Discussions

Using a robust IVT platform, we found that a remodeled Pol II and TFIIIA-7ZF can

efficiently utilize (-) PSTVd dimer for RNA-templated transcription. TFIIIA-7ZF significantly

enhances Pol II transcription activity on RNA templates. This remodeled Pol II represents

a new organization of functional polymerase complex. In this remodeled Pol II, we

observed the catalytic core (Rpb1 and Rpb2), a subassembly group (Rpb3, Rpb10,

Rpb11, likely Rpb12 as well), and an assembly factor Rpb8. Rpb4/Rpb7 heterodimer,

Rpb6, Rpb9, and likely Rpb5 were absent in the remodeled Pol II. Rpb4/Rpb7

heterodimer is not essential for elongation and is not included in the Pol II core [13, 14].

Rpb6 is involved in contact with TFIIS and TFIIH [36, 37], neither of which were present

in the transcription complex on our viroid RNA template. Furthermore, we showed that

TFIIS is not required for PSTVd replication [24]. Rpb9 is critical for Pol II fidelity by

delaying NTP sequestration [38, 39]. Pol II fidelity is also regulated by TFIIS [40, 41].

Interesting, both Rpb9 and TFIIS were absent in the remodeled Pol II samples, which is

in line with the observation that nuclear replicating viroids have a much higher mutation

rate than cellular Pol II transcripts [42]. Rpb5 is proposed to make contacts with DNA

promoters and coordinate the opening/closing of the Pol II DNA cleft [43-47], which is not

involved in RNA-templated transcription.

It has been proposed that an RNA polymerase may have evolved to transcribe RNA

templates first and then transitioned to use DNA templates in modern life forms [30].

Interestingly, Rpb4/Rpb7 heterodimer is present in all archaeal and eukaryotic cells but

not in bacteria [10, 48], further suggesting that organization changes occurred in RNA

polymerases during the course of evolution. The remodeled Pol II identified here only

retains a reduced set of factors, while most of the missing subunits are absent in bacterial

RNA polymerases (i.e., Rpb4, Rpb5, Rpb7, and Rpb9). Our discovery of a remodeled Pol

II actively transcribing RNA templates may provide a handle to further explore the

functional evolution of transcription machinery.

The high-resolution crystallographic structure of Pol II core-Rpb4/Rpb7-TFIIS showed

that Pol II utilizes the same active site for interacting DNA and RNA templates, revealed

by using chimeric RNA templates [49]. Later, one study using a chimeric RNA template

containing a HDV fragment sequence suggested that multiple general transcription

factors (TFIIA, TFIIB, TFIID, TFIIE, TFIIF, TFIIH, and TFIIS) may be potentially involved

in the initiation of RNA-templated transcription [50]. However, neither experimental

system could yield long RNA products, suggesting that those transcription complexes

might not be optimal for subviral RNA templates. In the transcription complex on PSTVd

RNA template, we could not detect the presence of TFIIE or TFIIF, despite that they were

both identified confidently in at least two out of three replicates in partially purified Pol II

(S1 Table). It is noteworthy that other Pol II-associated general transcription factors (TFIIA,

TFIIB, TFIID, TFIIH, and TFIIS) were also absent in partially purified Pol II and remodeled

Pol II samples. Although we cannot rule out the possibility that minute amount of general

transcription factors were remaining in our samples below the detection capacity of nLC-

MS/MS, it is unlikely for them to play a role in a stoichiometric ratio to Pol II resembling

the transcription complex on DNA templates. Therefore, our observation argues that

those general transcription factors for DNA-dependent transcription are not required in

the transcription complex on RNA templates, at least for viroid RNA templates. Hence,

the results outlined a distinct organization of transcription complex on RNA templates.

During transcription elongation on DNA templates, multiple auxiliary factors play

regulatory roles in cells. For example, the FACT complex and PAF1 complex regulate

transcription elongation. However, those factors may not be needed for in vitro

transcription [13, 14]. In line with the previous observations, PAF1 complex and most of

the FACT components (except for significantly reduced CTR9) were absent in the

transcription complex on PSTVd RNA templates (Figure 2 and S1 Table). Therefore, it

awaits future investigations to understand whether those factors are involved in RNA-

templated transcription.

All seven zinc finger domains of TFIIIA-7ZF are critical for the function in aiding RNA-

templated transcription. The first three zinc finger domains are pivotal for RNA template

binding. Interestingly, Pol II exhibited weaker affinity to RNA templates in the presence of

either *zf1*, *zf2*, or *zf3* mutant. It is intuitive to speculate that free *zf1*, *zf2*, and *zf3*

sequestered Pol II and prevented Pol II binding to RNA templates, leading to greatly

reduced transcription activity. It is unclear why the *zf6* mutant also greatly diminished Pol

II activity in generating full-length product as the amount of *zf6* and Pol II remaining on

the RNA template resembles that in the reaction with WT TFIIIA-7ZF. Since our system

only tested TFIIIA-7ZF and Pol II binding to the RNA templated before reaction initiates,

reasonable speculation is that *zf6* might be critical for transcription initiation or even

elongation.

Our reconstituted IVT system is robust for exploring the factors and functional

mechanism underlying viroid RNA-templated transcription, particularly for studying

transcription initiation and elongation. However, the current system does not have a

regulated termination process as the transcription is terminated by template run-off. A

modified system is needed in the future to serve the purpose of studying transcription

termination on RNA templates. In addition, structural analyses of the remodeled Pol

II/TFIIIA-7ZF complex on viroid RNA templates may provide more insights into the

regulation and mechanism underlying RNA-templated transcription.

Materials and Methods

Molecular constructs

We have previously reported WT TFIIIA-7ZF cloned from *Nicotiana benthamiana* in

bacteria expression vector pTXB1 (New England Biolabs, Ipswich, MA) [23]. The TFIIIA-

7ZF mutants were generated via site-directed mutagenesis using the WT TFIIIA-7ZF in

pTXB1 as the template (See S2 Table for primer information). We have also reported the

pInt95-94(-) and pInt95-94(+) constructs for generating PSTVd probes to detect sense

and antisense PSTVd RNAs, respectively [23]. PSTVd dimer construct was reported

previously [51]. All constructs have been verified by Sanger sequencing.

Protein purification

The protocol for recombinant protein purification was based on our reported protocol

[23]. Various recombinant TFIIIA-7ZF proteins with an intein-chitin binding domain (CBD)

tag were overexpressed using the *Escherichia coli* BL21(DE3) Rosetta strain (EMD

Millipore, Burlington, MA). For each construct, about 500 ml bacterial culture was

collected and re-suspended. After sonication with Bioruptor (Diagenode, Denville, NJ),

samples were subjected to centrifugation at 15,000X *g* for 1 h at 4°C. The cell lysate was

collected and incubated for 1 h with 2 ml of 50% slurry of chitin resin (New England

Biolabs) before loading onto an empty EconoPac gravity-flow column (Bio-Rad

Laboratories, Hercules, CA). After washing, resin was incubated for 18 h at 4°C in a

cleavage buffer [20 mM Tris-HCl (pH 8.5), 500 mM NaCl, 50 mM dithiothreitol, and 5 µM

$ZnSO_4$]. Fractions containing tagless proteins were dialyzed against 20 mM HEPES, pH

7.5, 200 mM NaCl, 50 µM $ZnSO_{4,}$ and 5 mM DTT. Protein concentrations were estimated

by Coomassie Brilliant Blue staining of an SDS-PAGE gel using as reference standards

bovine serum albumin of known concentrations.

Purification of Pol II from wheat germ was carried out following our published protocol

[24]. All operations were performed at 4°C, and all centrifugations were carried out for 15

min. One hundred grams raw wheat germ (Bob's Red Mill, Milwaukie, OR) was ground in

a Waring Blender with 400 mL of buffer A [50 mM Tris-HCl pH 7.9, 0.1 mM EDTA, 1 mM

DTT, and 75 mM $(NH_4)_2SO_4$]. The resulting homogenate was diluted with 100 mL of buffer

A and followed by centrifugation at 15,000× g. Supernatant was filtered through one layer

of Miracloth (MilliporeSigma, Burlington, MA). The resulting crude extract containing Pol

II was precipitated by an addition of 0.075 volume of 10% (v/v) Polymin P with rapid

stirring. The resulting mixture was subject to centrifugation at 10,000X g. The pellet was

washed with 200 mL of buffer A. The insoluble fraction, which contains Pol II, was

resuspended with the buffer B [50 mM Tris-HCl pH 7.9, 0.1 mM EDTA, 1 mM DTT, and

0.2 M $(NH_4)_2SO_4$]. The resulting suspension was centrifuged at 10,000X g to remove

insoluble pellets. $(NH_4)_2SO_4$ precipitation was carried out by slowly adding 20 g of solid

$(NH_4)_2SO_4$ per 100 mL of the above supernatant with stirring. Mixture was centrifuged,

and the pellet was dissolved in buffer C (0.05 M Tris-HCl pH 7.9, 0.1 mM EDTA, 1 mM

DTT, 25% ethylene glycol) plus 0.1% Brij 35 (Thermo Fisher Scientific, Waltham, MA) to

make final $(NH_4)_2SO_4$ concentration 0.15 M. The $(NH_4)_2SO_4$ concentration was

determined by conductivity. The resulting solution was applied to DEAE Sepharose FF

(GE Healthcare Life Sciences, Pittsburgh, PA) equilibrated with buffer C plus 0.15 M

$(NH_4)_2SO_4$. Then column was washed with five bed volume with buffer C containing 0.15

M $(NH_4)_2SO_4$. Finally, bound Pol II was eluted with buffer C containing 0.25 M $(NH_4)_2SO_4$.

Fractions containing Pol II were pooled. The $(NH_4)_2SO_4$ concentration was adjusted to 75

mM by conductivity. Resulting solution was applied to SP Sepharose FF (GE Healthcare

Life Sciences) equilibrated with the buffer C containing 75 mM $(NH_4)_2SO_4$. After washing

the column with the same buffer, Pol II was eluted using the buffer C containing 0.15 M

$(NH_4)_2SO_4$. Eluted fractions containing Pol II were pooled. Ethylene glycol (VWR

Chemicals BDH, Radnor, PA) was added to a final concentration of 50% (v/v) before

storing at −20°C.

In vitro transcription assay

Pol II-catalyzed in vitro transcription was carried out based on our recently developed

protocol [23, 24]. BSA (New England Biolabs) and TFIIIA-7ZF were treated with 1 unit of

Turbo DNase (Thermo Fisher Scientific) for 10 min at 37°C. Then, 0.39 pmol (-) PSTVd

dimer, 0.27 pmol partially purified Pol II, pretreated BSA (4 µM final concentration) and

various amounts of TFIIIA-7ZF were incubated at 28°C for 15 min. The reaction system

was adjusted to contain 50 mM HEPES-KOH pH 7.9, 1 mM $MnCl_2$, 6 mM $MgCl_2$, 40 mM

$(NH_4)_2SO_4$, 10% (v/v) glycerol, 1 unit/µL SuperaseIn RNase inhibitor (Thermo Fisher

Scientific), 0.5 mM rATP, 0.5 mM rCTP, 0.5 mM rGTP, 0.5 mM rUTP. Transcription

reactions (50 µL) were incubated at 28°C for 4 h. About 0.8 U/µL proteinase K (New

England Biolabs) was applied to terminate the reaction by incubation at 37°C for 15 min,

followed by incubation at 95°C for 5 min. The Pol II-catalyzed in vitro transcription assay

was repeated three times for each TFIIIA-7ZF variant. For the IVT assay in Figure 1D,

TFIIIA-7ZF and partially purified Pol II were subsequentially bound to immobilized

desthiobiotinylated RNA templates (see the section below for details). After washing twice,

additional RNA templates without desthiobiotinylation were supplied together with NTPs.

The reaction was then performed the same as abovementioned. This assay was repeated

twice.

RNA-based affinity purification

Using Pierce RNA 3' end desthiobiotinylation kit (Thermo Fisher Scientific, Waltham,

MA, USA), 50 pmol of PSTVd dimer RNA was labeled following the manufacturer's

instructions. Labeled RNA was purified using MEGAclear kit (Thermo Fisher Scientific,

Waltham, MA, USA) and heated at 65°C for 10 min followed by incubation at room

temperature for 12 min. Labeled RNA was bound to the 50 μL of streptavidin magnetic

beads (Thermo Fisher Scientific, Waltham, MA, USA). Magnetic beads were collected

and washed twice with equal volume of 20 mM Tris-HCl, pH 7.5. Beads were

subsequently washed with reaction buffer containing, 50 mM HEPES-KOH pH 8, 2 mM

$MnCl_2$, 6 mM $MgCl_2$, 40 mM $(NH_4)_2SO_4$, 10% glycerol. DNase treated 150 pmol of

recombinant TFIIIA-7ZF was incubated with RNA-bound beads in a 50 μL reaction at

28°C for 15 min. Then, 27 pmol of partially purified Pol II was added to the reaction and

incubated at 28°C for another 15 min. Next, beads were washed twice with wash buffer

(20mM Tris-HCl, pH 7.5, 10mM NaCl, 0.1% Tween-20) and bound proteins were eluted

with 1X SDS-loading buffer by heating 95°C for 5 min.

RNA gel blots

Detailed protocol has been reported previously [52]. Briefly, after electrophoresis in 5%

(w/v) polyacrylamide/8 M urea gel for 1 h at 200 V, RNAs were then transferred to

Hybond-XL nylon membranes (Amersham Biosciences, Little Chalfont, United Kingdom)

by a Bio-Rad semi-dry transfer cassette and were immobilized by a UV-crosslinker (UVP,

Upland, CA). The RNAs were then detected by DIG-labeled UTP probes. PSTVd RNAs

were prepared as described before [23]. *Sma*I-linearized pInt95-94(-) and pInt95-94(+)

were used as templates for generating probes, using the MAXIclear kit (Thermo Fisher

Scientific). The DIG-labeled probes were used for detecting PSTVd RNAs.

Immunoblots

Protein samples were separated on an SDS-PAGE gel, followed by transferring to

nitrocellulose membrane (GE Healthcare Lifesciences) using the Mini-PROTEAN Tetra

Cell (Bio-Rad Laboratories). After 1 h incubation with 1% (w/v) nonfat milk in 1X TBS (50

mM Tris-HCl, pH 7.5, 150 mM NaCl) at room temperature, membranes were incubated

with primary antibodies overnight at 4°C. After three washes with 1X TBST (50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.1% Tween 20), HRP-conjugated secondary antibodies were

added. Membrane was then washed three times with 1X TBST and incubated with HRP

substrates (Li-COR Biosciences, Lincoln, NE). The signals were detected with ChemiDoc

(Bio-Rad Laboratories).

For immunoblotting, polyclonal antibodies against TFIIIA were diluted as 1:2,000 and

the monoclonal 8WG16 antibodies (Thermo Fisher Scientific) were diluted at 1:1,000.

HRP-conjugated anti-mouse serum (Bio-Rad) was diluted at 1:5,000. HRP-conjugated

anti-rabbit serum (Thermo Fisher Scientific) was diluted at 1:3,000. For silver staining, we

followed instructions of Silver BULLit kit (Amresco, Solon, OH).

Nano-liquid chromatography-tandem mass spectrometry analysis (nLC-MS/MS)

Prior mass spectrometry, samples were subjected to in-solution digestion. Briefly,

reduction treatment (100 mM dithiothreitol and 15 min incubation at 65°C) and alkylation

treatment (100 mM iodoacetamide / 45 min incubation at room temperature) were

followed by 16 hr incubation at 37°C with sequencing grade trypsin (Promega, Madison

WI). Tryptic peptides were acidified with formic acid, lyophilized and stored at -80°C. As

described previously [53], two micrograms of digested protein were subjected to nLC-

MS/MS analysis using the LTQ-Orbitrap Velos mass spectrometer (Thermo Fisher

Scientific) directly linked to the Ultimate 3000 UPLC system (Thermo Fisher Scientific),

with following modification: mass spectra of intact and fragmented peptides were

collected in the orbitrap and linear ion trap detector, respectively. All data files were

deposited to PRIDE database [54] (accession PXD033736). The .raw mass spectral files

were searched using the SEQUEST algorithm of the Proteome Discoverer (PD) software

version 2.1 (Thermo Fisher Scientific). Tolerances were set to 10 ppm and 0.8 Da to

match precursor and fragment monoisotopic masses, respectively. The Triticum NCBI

Ref protein database (as of February 2022, with 122,221 entries) and its reversed copy

served as the target and decoy database, respectively, to allow calculation of False

Discovery Rate (FDR). All proteins presented in results were filtered by FDR<1%, and

identified by minimum of 2 peptides and 2 PSMs (peptide-spectrum matches) in each

replicate. The PD result data files showing peptide/protein-ID relevant parameters for

each individual replicate are given in Supplementary Material.

Acknowledgements

This work was supported by US National Science Foundation (MCB-1906060 and

MCB-2145967 to YW), US National Institute of General Medical Sciences

(1R15GM135893 to YW), and NIH MS-IDeA Network of Biomedical Research Excellence

award 5P20GM103476-19. The mass spectrometry proteomics analysis was performed

at the Institute for Genomics, Biocomputing and Biotechnology, Mississippi State

University, with partial support from Mississippi Agricultural and Forestry Experiment

Station. We are grateful for the constructive comments from Donna Gordon at Mississippi

State University.

Competing Interests

The authors declare no competing interests.

Accession Numbers

The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository with the dataset

identifier PXD033736.

Please note, the data will be publicly available upon acceptance of the manuscript.

Reviewers can access the data by using the following login information: Username:

reviewer_pxd033736@ebi.ac.uk ; Password: yThN9ExZ

Author Contributions

Y.W. conceived and designed the experiments. S.D.DM., T.P., B.L., and Y.W. wrote

the manuscript. S.D.DM., T.P., and O.P. performed experiments. S.D.DM., J.M., and Y.W.

prepared key materials. S.D.DM., T.P., B.L., and Y.W. analyzed the data.

References

1.   Wang Y. Current view and perspectives in viroid replication. Curr Opin Virol. 2021;47:32-
7. Epub 2021/01/19. doi: 10.1016/j.coviro.2020.12.004. PubMed PMID: 33460914; PubMed
Central PMCID: PMCPMC8068583.

2.   Navarro B, Flores R, Di Serio F. Advances in Viroid-Host Interactions. Annu Rev Virol.
2021;8(1):305-25. Epub 2021/07/14. doi: 10.1146/annurev-virology-091919-092331. PubMed
PMID: 34255541.
3.   Ma J, Mudiyanselage SDD, Wang Y. Emerging value of the viroid model in molecular
biology and beyond. Virus Res. 2022;313:198730. Epub 2022/03/10. doi:
10.1016/j.virusres.2022.198730. PubMed PMID: 35263622; PubMed Central PMCID:
PMCPMC8976779.
4.   Di Serio F, Owens RA, Li SF, Matousek J, Pallas V, Randles JW, et al. ICTV Virus Taxonomy
Profile: Pospiviroidae. J Gen Virol. 2021;102(2). Epub 2020/12/18. doi: 10.1099/jgv.0.001543.
PubMed PMID: 33331814; PubMed Central PMCID: PMCPMC8116940.
5.   Di Serio F, Li SF, Matousek J, Owens RA, Pallas V, Randles JW, et al. ICTV Virus Taxonomy
Profile: Avsunviroidae. J Gen Virol. 2018;99(5):611-2. Epub 2018/03/28. doi:
10.1099/jgv.0.001045. PubMed PMID: 29580320.
6.   Dissanayaka Mudiyanselage SD, Qu J, Tian N, Jiang J, Wang Y. Potato Spindle Tuber
Viroid RNA-Templated Transcription: Factors and Regulation. Viruses. 2018;10(9). Epub
2018/09/20. doi: 10.3390/v10090503. PubMed PMID: 30227597; PubMed Central PMCID:
PMCPMC6164485.
7.   Roeder RG. Role of general and gene-specific cofactors in the regulation of eukaryotic
transcription. Cold Spring Harb Symp Quant Biol. 1998;63:201-18. Epub 1999/06/29. doi:
10.1101/sqb.1998.63.201. PubMed PMID: 10384284.
8.   Lee TI, Young RA. Transcription of eukaryotic protein-coding genes. Annu Rev Genet.
2000;34:77-137. Epub 2000/11/28. doi: 10.1146/annurev.genet.34.1.77. PubMed PMID:
11092823.
9.   Lemon B, Tjian R. Orchestrated response: a symphony of transcription factors for gene
control. Genes Dev. 2000;14(20):2551-69. Epub 2000/10/21. doi: 10.1101/gad.831000. PubMed
PMID: 11040209.
10.   Armache KJ, Kettenberger H, Cramer P. Architecture of initiation-competent 12-subunit
RNA polymerase II. Proc Natl Acad Sci U S A. 2003;100(12):6964-8. Epub 2003/05/15. doi:
10.1073/pnas.1030608100. PubMed PMID: 12746495; PubMed Central PMCID:
PMCPMC165813.
11.   Liu X, Bushnell DA, Kornberg RD. RNA polymerase II transcription: structure and
mechanism. Biochim Biophys Acta. 2013;1829(1):2-8. Epub 2012/09/25. doi:
10.1016/j.bbagrm.2012.09.003. PubMed PMID: 23000482; PubMed Central PMCID:
PMCPMC4244541.
12.   Bushnell DA, Bamdad C, Kornberg RD. A minimal set of RNA polymerase II transcription
protein interactions. J Biol Chem. 1996;271(33):20170-4. Epub 1996/08/16. doi:
10.1074/jbc.271.33.20170. PubMed PMID: 8702741.
13.   Ruet A, Sentenac A, Fromageot P, Winsor B, Lacroute F. A mutation of the B220 subunit
gene affects the structural and functional properties of yeast RNA polymerase B in vitro. J Biol
Chem. 1980;255(13):6450-5. Epub 1980/07/10. PubMed PMID: 6993472.
14.   Edwards AM, Kane CM, Young RA, Kornberg RD. Two dissociable subunits of yeast RNA
polymerase II stimulate the initiation of transcription at a promoter in vitro. J Biol Chem.
1991;266(1):71-5. Epub 1991/01/05. PubMed PMID: 1985924.

15.   Schweikhard V, Meng C, Murakami K, Kaplan CD, Kornberg RD, Block SM. Transcription
factors TFIIF and TFIIS promote transcript elongation by RNA polymerase II by synergistic and
independent mechanisms. Proc Natl Acad Sci U S A. 2014;111(18):6642-7. Epub 2014/04/16.
doi: 10.1073/pnas.1405181111. PubMed PMID: 24733897; PubMed Central PMCID:
PMCPMC4020062.
16.   Vos SM, Farnung L, Boehning M, Wigge C, Linden A, Urlaub H, et al. Structure of
activated transcription complex Pol II-DSIF-PAF-SPT6. Nature. 2018;560(7720):607-12. Epub
2018/08/24. doi: 10.1038/s41586-018-0440-4. PubMed PMID: 30135578.
17.   Antosz W, Pfab A, Ehrnsberger HF, Holzinger P, Kollen K, Mortensen SA, et al. The
Composition of the Arabidopsis RNA Polymerase II Transcript Elongation Complex Reveals the
Interplay between Elongation and mRNA Processing Factors. Plant Cell. 2017;29(4):854-70.
Epub 2017/03/30. doi: 10.1105/tpc.16.00735. PubMed PMID: 28351991; PubMed Central
PMCID: PMCPMC5435424.
18.   Dezelee S, Sentenac A, Fromageot P. Role of deoxyribonucleic acid-ribonucleic acid
hybrids in eukaryotes. Synthetic ribo- and deoxyribopolynucleotides as template for yeast
ribonucleic acid polymerase B (or II). J Biol Chem. 1974;249(18):5978-83. Epub 1974/09/25.
PubMed PMID: 4607037.
19.   Flores R, Owens RA, Taylor J. Pathogenesis by subviral agents: viroids and hepatitis delta
virus. Curr Opin Virol. 2016;17:87-94. Epub 2016/02/22. doi: 10.1016/j.coviro.2016.01.022.
PubMed PMID: 26897654.
20.   Wassarman K, Saecker R. Synthesis-mediated release of a small RNA inhibitor of RNA
polymerase. Science. 2006;314(5805):1601-3. PubMed PMID:
2818717177218692492related:jHVpplYYHicJ.
21.   Wagner SD, Yakovchuk P, Gilman B, Ponicsan SL, Drullinger LF, Kugel JF, et al. RNA
polymerase II acts as an RNA-dependent RNA polymerase to extend and destabilize a non-
coding RNA. Embo J. 2013;32(6):781-90. doi: 10.1038/emboj.2013.18. PubMed PMID:
23395899; PubMed Central PMCID: PMCPMC3604716.
22.   Yamaguchi Y, Filipovska J, Yano K, Furuya A, Inukai N, Narita T, et al. Stimulation of RNA
polymerase II elongation by hepatitis delta antigen. Science. 2001;293(5527):124-7. Epub
2001/06/02. doi: 10.1126/science.1057925. PubMed PMID: 11387440.
23.   Wang Y, Qu J, Ji S, Wallace AJ, Wu J, Li Y, et al. A Land Plant-Specific Transcription Factor
Directly Enhances Transcription of a Pathogenic Noncoding RNA Template by DNA-Dependent
RNA Polymerase II. Plant Cell. 2016;28(5):1094-107. Epub 2016/04/27. doi:
10.1105/tpc.16.00100. PubMed PMID: 27113774; PubMed Central PMCID: PMCPMC4904678.
24.   Dissanayaka Mudiyanselage SD, Wang Y. Evidence Supporting That RNA Polymerase II
Catalyzes De Novo Transcription Using Potato Spindle Tuber Viroid Circular RNA Templates.
Viruses. 2020;12(4). Epub 2020/04/02. doi: 10.3390/v12040371. PubMed PMID: 32230827.
25.   Thomas MC, Chiang CM. The general transcription machinery and general cofactors. Crit
Rev Biochem Mol Biol. 2006;41(3):105-78. Epub 2006/07/25. doi:
10.1080/10409230600648736. PubMed PMID: 16858867.
26.   Taylor JM. Chapter 3. Replication of the hepatitis delta virus RNA genome. Adv Virus
Res. 2009;74:103-21. Epub 2009/08/25. doi: 10.1016/S0065-3527(09)74003-5. PubMed PMID:
19698896.

27.   Taylor JM. Hepatitis D Virus Replication. Cold Spring Harb Perspect Med. 2015;5(11).
Epub 2015/11/04. doi: 10.1101/cshperspect.a021568. PubMed PMID: 26525452; PubMed
Central PMCID: PMCPMC4632862.
28.   Qi Y, Ding B. Replication of Potato spindle tuber viroid in cultured cells of tobacco and
Nicotiana benthamiana: the role of specific nucleotides in determining replication levels for
host adaptation. Virology. 2002;302(2):445-56. PubMed PMID: 12441088.
29.   Rackwitz HR, Rohde W, Sänger HL. DNA-dependent RNA polymerase II of plant origin
transcribes viroid RNA into full-length copies. Nature. 1981;291(5813):297-301. PubMed PMID:
7231549.
30.   Ream TS, Haag JR, Wierzbicki AT, Nicora CD, Norbeck AD, Zhu JK, et al. Subunit
compositions of the RNA-silencing enzymes Pol IV and Pol V reveal their origins as specialized
forms of RNA polymerase II. Mol Cell. 2009;33(2):192-203. Epub 2008/12/27. doi:
10.1016/j.molcel.2008.12.015. PubMed PMID: 19110459; PubMed Central PMCID:
PMCPMC2946823.
31.   Farnung L, Ochmann M, Engeholm M, Cramer P. Structural basis of nucleosome
transcription mediated by Chd1 and FACT. Nat Struct Mol Biol. 2021;28(4):382-7. Epub
2021/04/14. doi: 10.1038/s41594-021-00578-6. PubMed PMID: 33846633; PubMed Central
PMCID: PMCPMC8046669.
32.   Francette AM, Tripplehorn SA, Arndt KM. The Paf1 Complex: A Keystone of Nuclear
Regulation Operating at the Interface of Transcription and Chromatin. J Mol Biol.
2021;433(14):166979. Epub 2021/04/04. doi: 10.1016/j.jmb.2021.166979. PubMed PMID:
33811920; PubMed Central PMCID: PMCPMC8184591.
33.   Lee BM, Xu J, Clarkson BK, Martinez-Yamout MA, Dyson HJ, Case DA, et al. Induced fit
and "lock and key" recognition of 5S RNA by zinc fingers of transcription factor IIIA. J Mol Biol.
2006;357(1):275-91. Epub 2006/01/13. doi: 10.1016/j.jmb.2005.12.010. PubMed PMID:
16405997.
34.   Rothfels K, Rowland O, Segall J. Zinc fingers 1 and 7 of yeast TFIIIA are essential for
assembly of a functional transcription complex on the 5 S RNA gene. Nucleic Acids Res.
2007;35(14):4869-81. Epub 2007/07/13. doi: 10.1093/nar/gkm517. PubMed PMID: 17626045;
PubMed Central PMCID: PMCPMC1950542.
35.   Goodman TC, Nagel L, Rappold W, Klotz G, Riesner D. Viroid replication: equilibrium
association constant and comparative activity measurements for the viroid-polymerase
interaction. Nucleic Acids Res. 1984;12(15):6231-46. Epub 1984/08/10. doi:
10.1093/nar/12.15.6231. PubMed PMID: 6473106; PubMed Central PMCID: PMCPMC320069.
36.   Okuda M, Suwa T, Suzuki H, Yamaguchi Y, Nishimura Y. Three human RNA polymerases
interact with TFIIH via a common RPB6 subunit. Nucleic Acids Res. 2022;50(1):1-16. Epub
2021/07/17. doi: 10.1093/nar/gkab612. PubMed PMID: 34268577; PubMed Central PMCID:
PMCPMC8754651.
37.   Ishiguro A, Nogi Y, Hisatake K, Muramatsu M, Ishihama A. The Rpb6 subunit of fission
yeast RNA polymerase II is a contact target of the transcription elongation factor TFIIS. Mol Cell
Biol. 2000;20(4):1263-70. Epub 2000/01/29. doi: 10.1128/MCB.20.4.1263-1270.2000. PubMed
PMID: 10648612; PubMed Central PMCID: PMCPMC85260.
38.   Walmacq C, Kireeva ML, Irvin J, Nedialkov Y, Lubkowska L, Malagon F, et al. Rpb9
subunit controls transcription fidelity by delaying NTP sequestration in RNA polymerase II. J Biol

Chem. 2009;284(29):19601-12. Epub 2009/05/15. doi: 10.1074/jbc.M109.006908. PubMed
PMID: 19439405; PubMed Central PMCID: PMCPMC2740586.

39. Nesser NK, Peterson DO, Hawley DK. RNA polymerase II subunit Rpb9 is important for
transcriptional fidelity in vivo. Proc Natl Acad Sci U S A. 2006;103(9):3268-73. Epub 2006/02/24.
doi: 10.1073/pnas.0511330103. PubMed PMID: 16492753; PubMed Central PMCID:
PMCPMC1413937.

40. Thomas MJ, Platas AA, Hawley DK. Transcriptional fidelity and proofreading by RNA
polymerase II. Cell. 1998;93(4):627-37. Epub 1998/05/30. doi: 10.1016/s0092-8674(00)81191-5.
PubMed PMID: 9604937.

41. Jeon C, Agarwal K. Fidelity of RNA polymerase II transcription controlled by elongation
factor TFIIS. Proc Natl Acad Sci U S A. 1996;93(24):13677-82. Epub 1996/11/26. doi:
10.1073/pnas.93.24.13677. PubMed PMID: 8942993; PubMed Central PMCID: PMCPMC19388.

42. Lopez-Carrasco A, Ballesteros C, Sentandreu V, Delgado S, Gago-Zachert S, Flores R, et
al. Different rates of spontaneous mutation of chloroplastic and nuclear viroids as determined
by high-fidelity ultra-deep sequencing. PLoS Pathog. 2017;13(9):e1006547. Epub 2017/09/15.
doi: 10.1371/journal.ppat.1006547. PubMed PMID: 28910391; PubMed Central PMCID:
PMCPMC5614642.

43. Zaros C, Briand JF, Boulard Y, Labarre-Mariotte S, Garcia-Lopez MC, Thuriaux P, et al.
Functional organization of the Rpb5 subunit shared by the three yeast RNA polymerases.
Nucleic Acids Res. 2007;35(2):634-47. Epub 2006/12/21. doi: 10.1093/nar/gkl686. PubMed
PMID: 17179178; PubMed Central PMCID: PMCPMC1802627.

44. Kim TK, Lagrange T, Wang YH, Griffith JD, Reinberg D, Ebright RH. Trajectory of DNA in
the RNA polymerase II transcription preinitiation complex. Proc Natl Acad Sci U S A.
1997;94(23):12268-73. Epub 1997/11/14. doi: 10.1073/pnas.94.23.12268. PubMed PMID:
9356438; PubMed Central PMCID: PMCPMC24903.

45. Cramer P, Bushnell DA, Kornberg RD. Structural basis of transcription: RNA polymerase II
at 2.8 angstrom resolution. Science. 2001;292(5523):1863-76. Epub 2001/04/21. doi:
10.1126/science.1059493. PubMed PMID: 11313498.

46. Cramer P, Bushnell DA, Fu J, Gnatt AL, Maier-Davis B, Thompson NE, et al. Architecture
of RNA polymerase II and implications for the transcription mechanism. Science.
2000;288(5466):640-9. Epub 2000/04/28. doi: 10.1126/science.288.5466.640. PubMed PMID:
10784442.

47. Chen X, Qi Y, Wu Z, Wang X, Li J, Zhao D, et al. Structural insights into preinitiation
complex assembly on core promoters. Science. 2021;372(6541). Epub 2021/04/03. doi:
10.1126/science.aba8490. PubMed PMID: 33795473.

48. Werner F, Eloranta JJ, Weinzierl RO. Archaeal RNA polymerase subunits F and P are bona
fide homologs of eukaryotic RPB4 and RPB12. Nucleic Acids Res. 2000;28(21):4299-305. Epub
2000/11/01. doi: 10.1093/nar/28.21.4299. PubMed PMID: 11058130; PubMed Central PMCID:
PMCPMC113124.

49. Lehmann E, Brueckner F, Cramer P. Molecular basis of RNA-dependent RNA polymerase
II activity. Nature. 2007;450(7168):445-9. doi: 10.1038/nature06290. PubMed PMID: 18004386.

50. Abrahem A, Pelchat M. Formation of an RNA polymerase II preinitiation complex on an
RNA promoter derived from the hepatitis delta virus RNA genome. Nucleic Acids Res.

2008;36(16):5201-11. Epub 2008/08/07. doi: 10.1093/nar/gkn501. PubMed PMID: 18682525;
PubMed Central PMCID: PMCPMC2532721.
51.   Zhong X, Leontis N, Qian S, Itaya A, Qi Y, Boris-Lawrie K, et al. Tertiary structural and
functional analyses of a viroid RNA motif by isostericity matrix and mutagenesis reveal its
essential role in replication. J Virol. 2006;80(17):8566-81. Epub 2006/08/17. doi:
10.1128/JVI.00837-06. PubMed PMID: 16912306; PubMed Central PMCID: PMCPMC1563885.
52.   Jiang J, Smith HN, Ren D, Dissanayaka Mudiyanselage SD, Dawe AL, Wang L, et al. Potato
Spindle Tuber Viroid Modulates Its Replication through a Direct Interaction with a Splicing
Regulator. J Virol. 2018;92(20). Epub 2018/08/03. doi: 10.1128/JVI.01004-18. PubMed PMID:
30068655; PubMed Central PMCID: PMCPMC6158407.
53.   Takac T, Krenek P, Komis G, Vadovic P, Ovecka M, Ohnoutkova L, et al. TALEN-Based
HvMPK3 Knock-Out Attenuates Proteome and Root Hair Phenotypic Responses to flg22 in
Barley. Front Plant Sci. 2021;12:666229. Epub 2021/05/18. doi: 10.3389/fpls.2021.666229.
PubMed PMID: 33995462; PubMed Central PMCID: PMCPMC8117018.
54.   Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S, Kamatchinathan
S, et al. The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics
evidences. Nucleic Acids Res. 2022;50(D1):D543-D52. Epub 2021/11/02. doi:
10.1093/nar/gkab1038. PubMed PMID: 34723319; PubMed Central PMCID: PMCPMC8728295.
630

Figure Legends

Figure 1. Uncovering the role of TFIIIA-7ZF and Pol II in transcribing (-) PSTVd

oligomers. (A) Reconstituted in vitro transcription (IVT) system using partially purified

Pol II (from wheat germ), (-) PSTVd dimer RNA (7.8 nM), and various amounts of TFIIIA-

7ZF. Sequence-specific riboprobes were used to detect (-) PSTVd templates and (+)

PSTVd products (at the position close to dimer PSTVd). Quantification of product

intensities was performed using ImageJ. The first lane signal was set as 0 and the second

lane signal was set as 1. Data from three replicates were used for graphing the fold

increases induced by various amount of TFIIIA-7ZF. (B) Schematic presentation for RNA-

based affinity purification followed by nLC-MS/MS identifying protein factors in partially

purified Pol II and remodeled Pol II. (C) Peptide counts for each Pol II subunit in all nLC-

MS/MS replicates. The summarized nLC-MS/MS data is listed in S1 Table. The original

data can be found in S1-S6 datasets. (D) IVT assay demonstrating the activity of

remodeled Pol II. The first lane contains free RNA as template, while the second lane

contains mixed free RNA and desthiobiothnylated RNA as template. For the second lane,

labeled RNA was used first to reconstitute the remodeled Pol II, and the free RNA was

then supplied together with NTPs. The reaction condition is described in Methods with

details.

Figure 2. Analyses on transcription elongation factors. (A) Analyses on peptide

counts of the FACT complex (SSRP1-B, SPT16), SPT6, PAF1-C, and TFIIF in partially

purified Pol II and remodeled Pol II. P values were calculated via two-tailed T-test, a built-

in function in Prism. The summarized nLC-MS/MS data are listed in S1 Table. The original

data can be found in S1-S6 datasets. (B) Schematic presentation of Pol II interactions

with SPT6 and PAF1-C during DNA-dependent transcription, based on [16]. P, PAF1. L,

LEO1. R, RTF1. Purple line, CTR9.

Figure 3. Analyses on the role of TFIIIA-7ZF zinc finger domains in aiding Pol II

activity on RNA templates. Reconstituted in vitro transcription (IVT) system using

partially purified Pol II, (-) PSTVd dimer RNA (7.8 nM), and various amounts of TFIIIA-

7ZF mutants. Sequence-specific riboprobes were used to detect (-) PSTVd templates and

(+) PSTVd products (at the position close to dimer PSTVd). Fold changes were analyzed

as described in Figure 1. P values for the most significant fold changes as compared with

Pol II only samples were listed. n.s., no significant comparisons were identified.

Figure 4. Analyses on the binding ability of TFIIIA-7ZF mutants. WT and mutants

TFIIIA-7ZF proteins were used for RNA-based affinity purification in the presence of

partially purified Pol II. (A) Immunoblots for input and RNA-bound TFIIIA-7ZF (anti-TFIIIA)

and the Rpb1 subunit of Pol II (8WG16). None, no TFIIIA-7ZF protein supplied. (B)

Quantification of immunoblotting results in (A). Protein signals in Pull-down blots were

normalized to the corresponding signals in the Input blots. The normalized signals in WT

(TFIIIA and Rpb1) were set as 100. Three replicates were performed to quantification and

statistical analyses. Two tailed T-test was used to calculate P values (listed in tables), by

using the built-in function in Prism.

A

WT

| Pol II (nM) | 0 | 5.4 | 5.4 | 5.4 | 5.4 |
| TFIIIA-7ZF (nM) | 0 | 0 | 18 | 30 | 60 |

Product

(-)-PSTVd template

P=0.02

B

C

	Patially purified Pol II			Pol II on RNA template		
	1	2	3	1	2	3
Rpb1	222	166	175	26	36	46
Rpb2	67	56	63	26	22	22
Rpb3	21	16	15	0	9	8
Rpb4	7	5	5	0	0	0
Rpb5	20	16	18	0	2	0
Rpb6	4	4	4	0	0	0
Rpb7	5	6	5	0	0	0
Rpb8	10	8	8	6	4	4
Rpb9	9	6	8	0	0	0
Rpb10	6	4	6	3	0	2
Rpb11	12	7	8	2	0	2

D

Product

(-)-PSTVd template

Partially purified Pol II   Remodeled Pol II

Figure 1

A

B

Figure 2

Figure 3

A

| | None | WT | zf1 | zf2 | zf3 | zf4 | zf5 | zf6 | zf7 |
Input — TFIIIA-7ZF / Pol II

Pull-down — TFIIIA-7ZF / Pol II

B

TFIIIA-7ZF

	WT	zf4	zf5	zf6	zf7
zf1	<0.0001	0.0053	<0.0001	0.0017	0.0072
zf2	<0.0001	0.0166	0.0002	0.0031	0.0266
zf3	0.0012	0.0358	0.0012	0.0053	0.0585
zf4	0.03			0.0458	
zf5	0.0704			0.9359	
zf6	0.3154				
zf7	0.0119			0.0273	

Pol II

	WT	zf4	zf5	zf6	zf7
None	0.1489				
zf1	<0.0001	<0.0001	0.0002	0.0003	0.0001
zf2	<0.0001	0.0002	0.0008	0.0006	0.0004
zf3	0.0002	0.0013	0.0039	0.0013	0.0015
zf4	<0.0001			0.0201	
zf5	0.0002			0.0114	
zf6	0.303				
zf7	0.0026			0.0848	

Figure 4