

# 1 **Experimental Test of the Contributions of Initial Variation and New Mutations to** 2 **Adaptive Evolution in a Novel Environment**

3 **Minako Izutsu<sup>1,2,3\*</sup>, and Richard E. Lenski<sup>1,2,3\*</sup>**

4 <sup>1</sup>Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,  
5 Michigan, USA

6 <sup>2</sup>BEACON Center for the Study of Evolution in Action, Michigan State University, East Lansing,  
7 Michigan, USA

8 <sup>3</sup>Ecology, Evolutionary Biology and Behavior Program, Michigan State University, East Lansing,  
9 Michigan, USA

10

## 11 **\* Correspondence:**

12 Minako Izutsu

13 [izutsumi@msu.edu](mailto:izutsumi@msu.edu)

14 Richard E. Lenski

15 [lenski@msu.edu](mailto:lenski@msu.edu)

16

17 **Keywords: adaptation, bacteria, convergent evolution, *Escherichia coli*, experimental evolution,**  
18 **fitness, genetic diversity, standing variation**

19

## 20 **ABSTRACT**

21 Experimental evolution is an approach that allows researchers to study organisms as they evolve in  
22 controlled environments. Despite the growing popularity of this approach, there are conceptual gaps  
23 among projects that use different experimental designs. One such gap concerns the contributions to  
24 adaptation of genetic variation present at the start of an experiment and that of new mutations that  
25 arise during an experiment. The primary source of genetic variation has historically depended largely  
26 on the study organisms. In the long-term evolution experiment (LTEE) using *Escherichia coli*, for  
27 example, each population started from a single haploid cell, and therefore adaptation depended  
28 entirely on new mutations. Most other microbial evolution experiments have followed the same  
29 strategy. By contrast, evolution experiments using multicellular, sexually-reproducing organisms  
30 typically start with pre-existing variation that fuels the response to selection. New mutations may  
31 also come into play in later generations of these experiments, but it is generally difficult to quantify  
32 their contribution in these studies. Here, we performed an experiment using *E. coli* to compare the  
33 contributions of initial genetic variation and new mutations to adaptation in a new environment. Our  
34 experiment had four treatments that varied in their starting diversity, with 18 populations in each  
35 treatment. One treatment depended entirely on new mutations, while the other three began with  
36 mixtures of clones, whole-population samples, or mixtures of whole-population samples from the  
37 LTEE. By tracking genetic markers associated with particular founders in two of our treatments, we  
38 could document the impact of the initial variation during the early generations of our experiment.  
39 However, there were no differences in fitness among the treatments after 500 or 2000 generations in

40 the new environment, despite the variation in fitness among the founders. These results indicate that  
41 new mutations quickly overcame, and eventually contributed more to adaptation, than did the initial  
42 variation. Our study thus shows that pre-existing genetic variation can have a strong impact on early  
43 evolution in a new environment, but new beneficial mutations may contribute more to later evolution  
44 and can even drive some initially beneficial variants to extinction.

45

## 46 INTRODUCTION

47 Some basic evolutionary issues can lead to misunderstandings and confusion, even among experts.  
48 One such issue concerns the contributions of standing genetic variation and new mutations to the  
49 process of adaptation by natural selection in a new environment. In this context, standing genetic  
50 variation includes those alleles that existed in a population before it encountered selection in the new  
51 environment, whereas new mutations are those alleles that arose after that selection began. It is a  
52 vexing problem because all genetic variation starts as new mutations and later can become standing  
53 variation, but the timing is important for understanding both the dynamics of evolutionary change  
54 within any single lineage and the repeatability of evolutionary outcomes across multiple lineages.  
55 With respect to the repeatability of evolution, Stern (2013) proposed the new term “collateral  
56 evolution” in juxtaposition with the more familiar idea of “parallel evolution” to emphasize how  
57 these different sources of genetic variation could lead to repeatable outcomes. Collateral evolution  
58 occurs when repeatable phenotypic changes evolve from standing variation in a common ancestral  
59 gene pool (i.e., variation that is identical by descent), whereas parallel evolution occurs when the  
60 similar phenotypes originate from independent mutational events (i.e., new mutations).

61 There is no single “right” answer in terms of the relative importance of standing variation and new  
62 mutations because both can contribute sequentially, simultaneously, and even synergistically to the  
63 process of adaptation by natural selection. But the ways that we do science—both conceptually and  
64 empirically—often lead us to emphasize one or the other source of genetic variation. In the long-term  
65 evolution experiment (LTEE) using *E. coli*, for example, new mutations are emphasized because  
66 each replicate population was founded from a single haploid cell of the ancestral strain in order to  
67 ensure that any repeatable outcomes result from independent mutations and hence parallel, rather  
68 than collateral, evolution (Lenski et al., 1991; Tenaillon et al., 2016; Lenski, 2017a). Hence, there  
69 was no standing variation at the start of the LTEE, and all of the genetic variation was produced by  
70 new mutations after the experiment began. Much of the work in the field of experimental evolution  
71 now follows the same mutation-dependent strategy, including most studies that use microorganisms  
72 (Tenaillon et al., 2012; Johnson et al., 2021). However, that approach is generally not followed in  
73 evolution experiments that use multicellular, sexually-reproducing plants and animals (Scarcelli and  
74 Kover, 2009; Burke et al., 2010; Schulte et al., 2010), for two largely practical reasons. First,  
75 quantitative genetics theory, which was developed for sexual plants and animals, presumes within-  
76 population genetic variation (Roff, 1997). That theory has guided artificial selection experiments to  
77 produce organisms with beneficial phenotypes for agricultural and other human applications (Hill  
78 and Caballero, 1992; Wright et al., 2005; Akey et al., 2010). By starting experiments with large,  
79 outbred populations that harbor abundant standing genetic variation, plant and animal breeders can  
80 improve traits more quickly than with small, inbred populations that lack diversity. Thus, most  
81 quantitative-genetic theories and applications presume that adaptation relies on standing variation,  
82 whereas the input from new mutations is typically ignored or abstracted (Roff, 1997). Second, the  
83 long generation times and small population sizes of larger organisms make evolution experiments  
84 that depend on new mutations (e.g., using near-isogenic inbred lines) impractical in most cases. Some

85 studies using isogenic *Drosophila* populations failed to observe repeatable evolutionary changes  
86 (Harshman and Hoffmann, 2000), and relying on new mutations for adaptation in populations with  
87 long generation times requires experiments that are longer than most researchers are willing to  
88 perform (Izutsu et al., 2012). Therefore, researchers studying animals and plants usually start with  
89 outbred populations that harbor abundant genetic variation, and thus they have largely observed  
90 collateral evolution with respect to the repeatability of changes across replicate populations (Rose,  
91 1984; Hoffmann et al., 2003; Mery and Kawecki, 2002; Barrett et al., 2008; Scarcelli and Kover,  
92 2009; Burke et al., 2010; Schulte et al., 2010; Zhou et al., 2011; Graves et al., 2017).

93 In this study, we directly compare the rates of adaptation based on standing genetic variation versus  
94 new mutations, in order to fill the gap among studies using different model systems. To that end, we  
95 used various sets of bacteria from the LTEE as founders, and we then propagated them in a novel  
96 environment in which D-serine replaced glucose as the limiting resource. We had 18 populations in  
97 each of four treatments (Figure 1). In the Single-Clone (SC) treatment, each population started from a  
98 single clone sampled from one of six LTEE populations. In the Single-Population (SP) treatment,  
99 each population started from an entire LTEE population and all of the genetic variation present in  
100 that population. In the Mixed-Clones (MC) treatment, each population started as an admixture of the  
101 six SC founding clones. Finally, in the Mixed-Populations (MP) treatment, each population started as  
102 an admixture of the six SP founding populations. Thus, the SC populations did not have any initial  
103 within-population genetic variation, and therefore they relied entirely on new mutations for their  
104 evolution. The SP populations began with both the common and rare alleles present at a moment in  
105 time in one of the LTEE populations. The MC populations began with six clones with approximately  
106 equal initial frequencies. The MP populations started with the most diversity, harboring essentially  
107 all of the genetic variation present in the other three treatments at the beginning of the evolution  
108 experiment. All 72 populations evolved for 2,000 generations (300 days) in the novel environment,  
109 with D-serine as their source of carbon and energy. Using stocks that we froze during the evolution  
110 experiment, we subsequently performed competition assays to measure the fitness of the evolved  
111 bacteria relative to common competitors, which allowed us to compare the extent of fitness gains  
112 among the four treatments. We also tracked a genetic marker embedded in our experiment, which  
113 allowed us to observe important dynamics especially during the first 100 generations or so of our  
114 experiment.

115

## 116 MATERIAL AND METHODS

### 117 Evolution Experiment in the D-Serine Environment

118 We used six whole-population samples and six clones from generation 50,000 of the LTEE as the  
119 founders for our new evolution experiment (Supplementary File 1). The populations are those named  
120 Ara-1, Ara-4, Ara-5, Ara-6, Ara+2, and Ara+5, and from those same populations we used the  
121 designated “A” clones that were previously isolated. The whole-population samples and clones were  
122 stored at  $-80^{\circ}\text{C}$ , where they have remained viable and available for future studies. Two of the six  
123 populations (Ara-1 and Ara-4) evolved hypermutability, while the other four retained the low  
124 ancestral mutation rate. Before starting our evolution experiment, we re-isolated clones from the  
125 freezer stocks for the six A clones on Davis minimal (DM) agar plates supplemented with 4 mg/mL  
126 glucose to ensure the genetic homogeneity of the clonal ancestors. Both the re-isolated clones and  
127 120  $\mu\text{L}$  of each whole-population sample were inoculated into 50-mL Erlenmeyer flasks containing  
128 9.9 mL of DM medium supplemented with 1000  $\mu\text{g/mL}$  glucose. These cultures were incubated for  
129 24 h in a shaking incubator at  $37^{\circ}\text{C}$  and 120 rpm. They were then frozen at  $-80^{\circ}\text{C}$  with glycerol as a

130 cryoprotectant, in order to generate and preserve samples of the precise ancestral stocks we used for  
131 our evolution experiment.

132 The new evolution experiment itself was begun as follows. On day  $-2$ , we inoculated 0.1 mL of each  
133 ancestral stock into 9.9 mL of DM medium supplemented with 2000  $\mu\text{g}/\text{mL}$  glucose. We incubated  
134 these cultures for 24 h in the same conditions as described above. On day  $-1$ , we diluted a portion of  
135 each culture 100-fold into saline solution, and then transferred 0.1 mL of the diluted culture into 9.9  
136 mL of DM medium supplemented with 25  $\mu\text{g}/\text{mL}$  glucose (the same medium as used in the LTEE),  
137 and then incubated the cultures for 24 h. On day 0, we took 2 mL from each of the 6 clonal cultures,  
138 mixed them well in a flask, and made a starter mix for the MC treatment. We made a similar mix for  
139 the MP treatment. We then transferred 0.1 mL of each culture into 9.9 mL of DM medium with 150  
140  $\mu\text{g}/\text{mL}$  D-serine (DS150) in an 18 x 150 mm test-tube, vortexed the culture, and then incubated the  
141 cultures for 24 h in a standing incubator at  $37^\circ\text{C}$ . We prepared 3 biological replicates from each of  
142 the 6 clonal and population cultures, making a total of 18 evolving populations in the SC and SP  
143 treatments (Figure 1). In those treatments, six sets of three populations shared the same initial genetic  
144 background (SC treatment) or the same initial genetic diversity (SP treatment). We also started 18  
145 populations from the clonal starter-mix for the MC treatment, and 18 populations from the population  
146 starter-mix for the MP treatment (Figure 1). The 18 populations in the MC treatment share the same  
147 set of initial genetic backgrounds, and the populations in the MP treatment share their initial genetic  
148 diversity, although very rare alleles might have been distributed unevenly, by chance, among the  
149 replicates of these treatments at the start of the evolution experiment.

150 We transferred the 72 populations (18 populations  $\times$  4 treatments) in 9.9 mL of fresh DS150 medium  
151 in test-tubes daily, following the same 100-fold dilution protocol for 300 days. In this environment,  
152 the populations reach a stationary-phase density of  $\sim 5 \times 10^7$  cells/mL and total size of  $\sim 5 \times 10^8$  cells.  
153 The bottleneck population size after the 100-fold dilutions is thus  $\sim 5 \times 10^6$  cells. These values are  
154 essentially the same as those for the glucose-limited LTEE populations. We froze samples of each  
155 population at  $-80^\circ\text{C}$  with glycerol as a cryoprotectant every 15 days through day 165, and then every  
156 15 or 30 days through day 300. We also froze the remaining volume of each culture from day 0.

157 During the evolution experiment, we diluted and spread cells from each population on tetrazolium  
158 arabinose (TA) indicator agar plates every 15 days to check for possible cross-contamination among  
159 the populations in the SC and SP treatments, where each population derived from either an  $\text{Ara}^-$  or  
160  $\text{Ara}^+$  lineage. We did not find any evidence of contamination during the 300 days of our evolution  
161 experiment. The populations in the MC and MP treatments had lineages with both marker states at  
162 the start, and we tracked the marker ratio in those populations for evidence of changing ratios, which  
163 would indicate fitness differences among the heterogenous founders and their descendants in these  
164 populations. To that end, we plated samples from the populations in the MC and MP treatments every  
165 other day until day 15, then every three days until day 45, and finally every five days until day 300.  
166 There was one interruption in the experiment at day 75. When we restarted the populations from the  
167 frozen samples, we plated all of them for the first three days to check whether freezing and reviving  
168 the samples altered the relative abundance of the marker states in the MC and MP populations with  
169 mixed ancestry. We did not see any substantial changes in the marker ratios, indicating that these  
170 steps did not substantially perturb the evolution experiment. Moreover, these procedures were  
171 applied to the populations in all four treatments, and thus they would not systematically bias the  
172 outcome.

## 173 **Fitness Measurements**

174 We isolated clones from each population at generations 500 and 2000 (i.e., days 75 and 300,  
175 Supplementary File 1) on DM agar plates with 900  $\mu\text{g}/\text{mL}$  D-serine, and we re-streaked the clones on  
176 TA plates to confirm their Ara marker state. The clones were chosen at random, except that each  
177 clone had the numerically dominant marker state for its source population at these time points for the  
178 MC and MP treatments. We then isolated Ara<sup>+</sup> mutants of several Ara<sup>-</sup> clones from generation 500 to  
179 identify potential common competitors with intermediate fitness relative to other clones from  
180 generations 0 to 2000. Using a single pair of common competitors (isogenic except for the Ara  
181 marker state) for the fitness assays simplifies procedures and inferences, and having intermediate  
182 fitness allows accurate estimates across a wide range of fitness values. We chose MI2228 and an  
183 Ara<sup>+</sup> revertant MI2339 as the common competitors for the main set of fitness assays (Supplementary  
184 File 1). MI2228 and MI2339 have equal fitness in DS150 medium, which indicates that the Ara<sup>+</sup>  
185 mutation is selectively neutral in that environment.

186 On day -2 of the assays, we transferred 0.1 mL from each competitor's freezer stock into 9.9 mL of  
187 Lysogeny Broth (LB) in a 50-mL Erlenmeyer flask, and we incubated the cultures overnight at 37°C  
188 and 120 rpm. At day -1, we diluted each culture 100-fold in saline solution, transferred 0.1 mL into  
189 9.9 mL of DS150 medium in a test-tube, vortexed it, and then incubated the cultures for 24 h in a  
190 standing incubator at 37°C. This day served as the conditioning step to ensure that competitors were  
191 acclimated to the environment where they would compete, and where the experimental populations  
192 had evolved. The rest of the procedure is the same as described elsewhere for the LTEE (Lenski et  
193 al., 1991; Wisser et al., 2013), except for the medium and culture vessel. In brief, we always competed  
194 the common competitor with the opposite marker state from the clone of interest. We transferred 0.05  
195 mL of each competitor's acclimated culture into 9.9 mL of DS150, and vortexed the new culture to  
196 mix the two competitors. We immediately took a sample, diluted it in saline solution, and plated cells  
197 on TA agar. The cultures were incubated for 24 h, at which time we again sampled the cultures and  
198 plated cells on TA agar. The resulting red (Ara<sup>-</sup>) and white (Ara<sup>+</sup>) colonies were counted after the  
199 plates were incubated for a day at 37°C. We calculated each competitor's realized growth rate as the  
200 log-transformed ratio of its final and initial densities. We then computed the fitness of the clone of  
201 interest relative to the common competitor as the ratio of their growth rates during the competition.

202 We used the generation 0 stocks multiple times for estimating initial fitness levels. We have only 12  
203 generation 0 stocks because we used the same six clones for the three replicates of each clone in the  
204 SC treatment, and the same six whole-population samples for the three replicates of each population  
205 in the SP treatment. The populations in the MC and MP treatments were derived from their  
206 respective starter mixes. We cannot measure the fitness of samples that contain both Ara<sup>-</sup> and Ara<sup>+</sup>  
207 cells using our method, which relies on a common competitor with the opposite marker state.  
208 Therefore, we used the same six clonal stocks at generation 0 for both the SC and MC treatments,  
209 and the same six population stocks at generation 0 for both the SP and MP treatments, and for all  
210 three replicates.

211 We also ran a second set of competition assays using the LTEE ancestors, REL606 and REL607, as  
212 common competitors. For these assays only, we used a 1:4 starting ratio at day 0, instead of the 1:1  
213 starting ratio described above, because of the substantially lower fitness of the LTEE ancestors in  
214 comparison to the common competitors used above. Specifically, we began each competition assay  
215 by mixing 0.08 mL of REL606 or REL607 and 0.02 mL of the strain of interest in the test-tube  
216 containing the DS150 medium. The assay conditions and the calculations of relative fitness were  
217 otherwise the same.

218 **Statistical Analyses**



219 All of our statistical analyses were performed using the referenced tests in R version 4.2.0. The  
220 analysis scripts and underlying data will be deposited in the Dryad Repository upon acceptance of  
221 this paper.

222

## 223 **RESULTS**

### 224 **Effect of Initial Variation on Fitness Improvement**

225 To assess the effect of the initial within-population diversity on adaptation to the new environment,  
226 we measured the relative fitness of the evolved bacteria by competing them against the common  
227 competitor strains. We cannot measure fitness of the entire evolved populations using our method,  
228 however, because that method requires mixing the evolved bacteria with the common competitor  
229 strain bearing the alternative Ara marker, and some populations in the MC and MP treatments had  
230 descendants of lineages with both marker states. Therefore, we isolated random clones at generations  
231 500 and 2000 as representatives of each population, and we measured their fitness. For generation 0,  
232 we used the stocks of the founder clones and populations that we froze immediately after the start of  
233 the evolution experiment. We used the six clone stocks that we had used to found populations in the  
234 SC and MC treatments as the generation 0 samples for those treatments, and we used the six whole-  
235 population stocks used to found populations in the SP and MP treatments as the generation 0 samples  
236 for those treatments. As a consequence, the generation 0 samples for the SC and MC treatments are  
237 technically identical, as are those for the SP and MP treatment.

238 Figure 2 shows the trajectories of the ln-transformed relative fitness values for the four treatments.  
239 As a reminder, the replicate populations in the SC and SP treatments had six different founding  
240 backgrounds. In contrast, the replicate populations in the MC and MP treatments originated from the  
241 same starter mix of six clones or six whole populations, respectively, and thus the replicates in those  
242 treatments shared the same founding backgrounds and diversity. The rate of increase in relative  
243 fitness clearly slowed over time in the D-serine environment (Figure 2). That deceleration is similar  
244 to what was seen during the first 2,000 generations in the glucose-limited environment of the LTEE  
245 (Lenski et al., 1991), and it is indicative of diminishing-returns epistasis (Wiser et al., 2013).

246 Most importantly for our aims and questions, we found no significant difference in fitness among the  
247 four treatments at either generation 500 or 2000 ( $p = 0.2300$  and  $p = 0.7213$ , respectively; one-way  
248 ANOVA, Table S1). The absence of meaningful differences among the treatments in the rate and  
249 extent of their adaptation was surprising to us, given the different levels of within-population genetic  
250 diversity at the beginning of the experiment. One possible explanation for the negative results with  
251 respect to differences in the final fitness values is that the initial variation present in treatments SP,  
252 MC, and MP did not include alleles that were sufficiently beneficial in the novel environment relative  
253 to new mutations. In other words, the populations in all four treatments ultimately depended on new  
254 mutations for adaptation to the novel D-serine medium, regardless of the different levels of initial  
255 genetic diversity. In the sections that follow, we present and examine additional data that helps to  
256 explain this result.

### 257 **Marker Trajectories During the Evolution Experiment**

258 We tracked the relative abundance of the two Ara marker states in all treatments during the evolution  
259 experiment (Figures 3 and S2). The populations in the SC and SP treatments began with a single  
260 marker state; in these populations, checking the marker states allowed us to check for cross-

261 contamination, which we did not see. The populations in the MC and MP treatments began with a  
262 mix of the two marker states. By tracking the relative abundance of the two states in those  
263 populations, we could observe the effects of both initial fitness variation linked to the markers and  
264 later beneficial mutations that gave rise to selective sweeps. The MC and MP treatments started with  
265 equal culture volumes of four Ara<sup>-</sup> lineages and two Ara<sup>+</sup> lineages; therefore, the log-transformed  
266 ratios of Ara<sup>-</sup> to Ara<sup>+</sup> cells were initially > 0 for all of the populations in those treatments (Figures 3  
267 and S2).

268 We observed strikingly similar marker trajectories among the 18 replicate populations in the MC and  
269 MP treatments, especially during the first ~100 generations (Figures 3 and S2). Despite the initially  
270 greater number of Ara<sup>-</sup> lineages, cells derived from one or more Ara<sup>+</sup> lineages increased in relative  
271 abundance in all 36 populations. By 30-50 generations, the Ara<sup>+</sup> cells were numerically dominant in  
272 all 18 MP populations and in most MC populations as well (Figure 3). These initial “bursts” imply  
273 that one or more of the Ara<sup>+</sup> clones and populations initially present in the MC and MP treatments  
274 were substantially more fit than the Ara<sup>-</sup> clones and populations. We will return to this point in the  
275 next section.

276 By generation 100, all 18 populations in the MC treatment, and most of the MP populations, had  
277 reversed course, with descendants of one or more Ara<sup>-</sup> lineages rising sharply in abundance relative  
278 to the Ara<sup>+</sup> descendants (Figure 3). The Ara<sup>-</sup> descendants remained numerically dominant through  
279 the first 500 generations in all 18 MC populations (Figure 3, top), and they evidently fixed in all 18  
280 cases by 2,000 generations (Figure S2, top). By contrast, the later marker-ratio trajectories of the MP  
281 populations were much more variable. Descendants of Ara<sup>-</sup> founders were usually more abundant  
282 through the first 500 generations, but with tremendous dispersion between the trajectories (Figure 3,  
283 bottom). By 2,000 generations, most MP populations had also evidently fixed one of the marker  
284 states, but with several fixations in each direction (Figure S2, bottom).

285 The marker-ratio trajectories also show that bursts leading to the early rise of cells derived from one  
286 or more Ara<sup>+</sup> lineages were much steeper for the populations in the MP treatment than for those in  
287 the MC treatment. While the initial ratios were virtually identical, at generation 47 (day 7) the mean  
288 log<sub>2</sub> ratios were -0.825 and -7.004 for the MC and MP treatments, respectively, even excluding two  
289 MP populations without any Ara<sup>-</sup> cells among the hundreds sampled. In fact, all 18 MP populations  
290 had a much lower ratio than any of the 18 MC populations, a difference that is highly significant ( $p$   
291  $\ll 0.0001$ ; two-tailed Welch’s  $t$ -test). We chose day 7 for this comparison because that is when the  
292 MC treatment showed the lowest average log ratio, although several adjacent days show a similarly  
293 stark difference between these two treatments.

294 In summary, we observed strikingly similar marker trajectories among the replicate populations in  
295 the MC and MP treatments in the early generations of our evolution experiment. Given the inevitable  
296 genetic linkage in asexual populations, this pattern implies that the metagenomes of the populations  
297 also evolved in parallel during this early phase. Moreover, this parallelism indicates that selection  
298 acted on shared genetic variation present in these populations at the start of experiment (i.e., identical  
299 by descent). It is reminiscent of the repeatability observed in previous evolution experiments with  
300 other organisms that were also founded by populations with shared initial variation (Burke et al.,  
301 2014).

## 302 **Fitness Differences Among the Founder Clones and Populations**

303 We examined the relative fitness values of the six founding populations and the six founding clones  
304 to better understand the similar early marker trajectories seen among the replicate populations in the  
305 MP and MC treatments, as well the difference between those treatments in the slope of those early  
306 trajectories (Figure 3). For these analyses, we use the same data as the generation 0 data that  
307 underlies the grand means for each treatment in the fitness trajectories (Figure 2).

308 Given the consistent marker-ratio trajectories towards the Ara<sup>+</sup> marker state, we expect to see that  
309 one or both of the Ara<sup>+</sup> founders had the highest fitness. Also, given that the early trend toward the  
310 Ara<sup>+</sup> state was much faster in the MP treatment than in the MC treatment, we expect that fitness  
311 differential to be greater among the whole-population founders than among the clonal founders.  
312 Figure 4 shows the relative fitness of the founding populations (panel A) and clones (panel B). In  
313 each panel, note that we have arranged the founders from the lowest to highest relative fitness.

314 Focusing first on the whole-population data (Figure 4A), we see that both of the Ara<sup>+</sup> founders have  
315 higher mean fitness than any of the Ara<sup>-</sup> founders in the DS150 environment. An ANOVA confirms  
316 that there is significant variation in fitness among the founders ( $p < 0.0001$ , Table S2, top), and  
317 Tukey's test confirms that the Ara<sup>+</sup> whole-population founders are significantly more fit than any of  
318 the Ara<sup>-</sup> founders. These results thus support our expectation from the marker trajectories that one or  
319 both of the Ara<sup>+</sup> founders had the highest fitness.

320 When we look at the corresponding data for the clonal founders, we see a more ambiguous pattern  
321 (Figure 4B). The relative fitness levels of the clones are more similar; four clones (two Ara<sup>+</sup> and Ara<sup>-</sup>  
322 ) are virtually identical to one another and slightly higher than two others (both Ara<sup>-</sup>). An ANOVA  
323 confirms that there is significant difference in fitness among the clone founders ( $p = 0.0004$ , Table  
324 S2, bottom), while Tukey's test finds no significant difference in fitness among the several most fit  
325 founder clones.

326 Based on the ANOVAs, we estimated the among-founder variance components,  $V_A$ , for fitness in  
327 these two treatments (Sokal and Rohlf, 1995). That founding variation is what would fuel the earliest  
328 response to selection in the evolution experiment before new mutations have had enough time to  
329 become relevant. As expected, the estimated variance in fitness among the whole-population  
330 founders ( $V_A = 0.0052$ ) is much greater than among the clonal founders ( $V_A = 0.0009$ ).

331 We also performed an additional set of competition assays to estimate the fitness of the founders of  
332 our evolution experiment relative to a different pair of common competitors. In this case, we  
333 competed the six founders of whole-populations and clones against the marked ancestors of the  
334 LTEE (Figure S3). The founders generally had higher fitness relative to the LTEE ancestors than  
335 relative to the common competitors used in our other assays. Therefore, we used a 1:4 starting ratio  
336 of the founders relative to the LTEE ancestors, instead of the 1:1 starting ratio used in the other  
337 competitions (Materials and Methods). Otherwise, the assay conditions and calculations of relative  
338 fitness as the ratio of realized growth rates were the same. We also arranged and analyzed these data  
339 as before.

340 These additional data also support one of our two expectations based on the marker trajectories,  
341 namely, that one Ara<sup>+</sup> founder had higher fitness than any of the Ara<sup>-</sup> founders. In this case, we see  
342 that Ara<sup>+</sup> has the highest mean fitness among both the whole-population (Figure S3A) and clonal  
343 (Figure S3B) founders. The results of the Tukey tests confirm that Ara<sup>+</sup> had significantly higher  
344 fitness than all other whole-population founders and higher fitness than all but one clonal founder.  
345 The ANOVAs indicate significant variation in fitness among both the whole-population (Table S3,



346 top) and clonal (Table S3, bottom) founders. However, the variation in fitness is not greater among  
347 the whole-population founders than among the clonal founders. The estimated among-founder  
348 variance component for fitness for the whole-population founders ( $V_A = 0.0274$ ) is essentially  
349 identical to the variance among the clonal founders ( $V_A = 0.0269$ ).

350 Across the four sets of competitions (founder clones and whole populations, against two pairs of  
351 common competitors), we find that the founders derived from LTEE population Ara+5 had the  
352 highest fitness in three of these sets (Figures 4A, S3A, and S3B), while they were tied for the highest  
353 fitness in one set (Figure 4B). These results clearly imply that the early trends toward the Ara<sup>+</sup> state  
354 in the marker-ratio trajectories in the MC (Figure 3, top) and especially the MP (Figure 3, bottom)  
355 treatments were caused by the initial fitness advantage that the Ara+5 founders had in the new DS150  
356 environment. By contrast, the subsequent reversals in most trajectories are presumably associated  
357 with new mutations that arose during our evolution experiment. (In theory, very gradual and uniform  
358 reversals could occur even without new mutations if the single most fit founder had a different  
359 marker state than the maker state with the higher average fitness across its constituent lineages.  
360 However, this hypothetical scenario is clearly not the case for the MP treatment, nor can it explain  
361 the variation in the time and strength of the reversals in the MP and MC treatments shown in Figure  
362 3.) A deeper understanding of the reversals will require future genomic analyses, as we explain in the  
363 Discussion.

364

## 365 DISCUSSION

366 It is generally difficult to disentangle the role of standing genetic variation and new mutations in the  
367 process of adaptation by natural selection. Even with experiments, different study systems tend to  
368 emphasize one source or the other. Selection experiments that use sexually reproducing plants and  
369 animals have typically started from base populations that harbor substantial standing variation, and  
370 they rarely run for more than a few tens of generations owing to the long generation time of these  
371 organisms. As a consequence, these experiments rely largely on variation that was present at the start  
372 of the experiment to fuel the response to selection. The field of experimental evolution with bacteria  
373 and other microorganisms has expanded greatly in recent years (Barrick and Lenski, 2013; Lenski,  
374 2017b; Van den Bergh et al., 2018). These study organisms have rapid generations, and most of them  
375 reproduce asexually during the experiments, even those that may undergo parasexual recombination  
376 (e.g., horizontal gene transfer) in nature. Our experiment was designed to compare the contributions  
377 of initial genetic variation and new mutations during adaptation of strictly asexual populations to a  
378 new environment.

379 To that end, we constructed four treatments with different initial levels of genetic diversity. Each  
380 treatment had 18 populations. In all cases, the founders came from the LTEE, in which *E. coli* have  
381 evolved in and adapted to a glucose-limited medium for 50,000 generations. At one extreme, each  
382 new population was founded by a single genotype, and thus there was no initial within-population  
383 diversity. We call this the Single-Clone (SC) treatment; six different clones, each derived from a  
384 different LTEE lineage, were used to found three replicate populations. At the other extreme, 18  
385 populations derived from an admixture of six whole-population samples that included both common  
386 and rare genotypes from the source populations. We call this the Mixed-Populations (MP) treatment.  
387 We also had two treatments that started with intermediate levels of genetic variation, which we call  
388 the Single-Population (SP) and Mixed-Clones (MC) treatments (Figure 1).

389 We propagated all 72 populations for 2000 generations in a new environment, one in which D-serine  
390 replaced glucose as the source of carbon and energy. We then measured the fitness of evolved strains  
391 from each population at both 500 and 2000 generations. We observed rapid early adaptation to the D-  
392 serine environment in all of the populations, but the rate of further fitness improvement declined over  
393 time, similar to what has been seen in the glucose environment of the LTEE (Wiser et al., 2013) as  
394 well as seen in other microbial evolution experiments (e.g., Johnson et al., 2021; Marad et al., 2018).

395 Most importantly, however, we found no significant differences among the four treatments in their  
396 mean fitness at generations 500 and 2000 (Figure 2), despite their different levels of genetic diversity  
397 at the beginning of the experiment. Thus, the populations in the SC treatment, each of which had no  
398 genetic diversity at the start, achieved the same fitness as the populations in the MP treatment, which  
399 started with all the diversity found in six LTEE populations combined. One possible explanation for  
400 this negative result would be that there were simply no differences in fitness in the D-serine medium  
401 among the founders in the treatments that began the experiment with genetic variation. In that case,  
402 all the populations in all four treatments would have had to depend entirely on new mutations to fuel  
403 adaptation to the new medium. But as we discovered, there was significant initial within-population  
404 variation for fitness in the new environment, at least in the MC and MP treatments.

405 Our first evidence of that initial fitness variation came from tracking the ratio of a neutral genetic  
406 marker that differed among the LTEE-derived founders, and which was therefore polymorphic in  
407 each of the populations in the MC and MP treatments. If there was no initial fitness variation in the  
408 new environment, then that ratio should have remained constant (within sampling error) until such  
409 time as a beneficial mutation occurred and began to sweep through one or the other marked  
410 backgrounds, thereby perturbing that ratio (Barrick et al., 2010; Izutsu et al., 2021). Alternatively, if  
411 the different founding genotypes had unequal fitness, then the marker ratio would systematically and  
412 immediately deviate from its initial value as a result of the inevitable linkage in asexual genomes  
413 between the marker and the alleles responsible for the fitness differences. This alternative outcome is  
414 precisely what we saw. We observed strikingly similar directional shifts in the marker-ratio  
415 trajectories among populations in the MC and MP treatments, especially during the first ~100  
416 generations (Figure 3). These parallel directional trajectories imply the presence of at least one  
417 “preadapted” genotype among the founders in those treatments.

418 We also compared the relative fitness of the founding clones and founding populations used in the  
419 MC and MP treatments, respectively. These comparisons showed that the founders derived from  
420 LTEE Ara+5 lineage had fitness as high as or higher than the other founders in the new D-serine  
421 environment (Figure 4 and S3), consistent with the early and systematic shifts in the marker-ratio  
422 trajectories to the Ara+ marker state. Also, the early marker-ratio trajectories in the MP treatment  
423 were much steeper than in the MC treatment (Figure 3), consistent with greater fitness differentials  
424 favoring the Ara+5 founders in the MP treatment (Figure 4). Thus, the genetic variation initially  
425 present in the MC and MP populations drove adaptation to the new environment during the first 100  
426 generations of our experiment. However, new beneficial mutations soon arose that perturbed and  
427 often reversed those early trends in the marker ratios (Figure 3). By generation 500, the beneficial  
428 effects of these new mutations were sufficiently large that the initial variation no longer mattered,  
429 and all four treatments—including even the SC treatment, in which each population started from a  
430 single clone—had achieved similar average fitness (Figure 2).

431 One might have expected that new beneficial mutations would have arisen randomly with respect to  
432 the marker state of the founders in the MC and MP treatments. Four of the six founders came from  
433 LTEE lineages with the Ara- marker state, and two from lineages with the Ara+ marker state. If the

434 mutations that were beneficial in the D-serine environment arose very early in the new experiment,  
435 then we might expect about two-thirds of the marker trajectories to reverse course and trend toward  
436 the Ara<sup>-</sup> state, after those mutations reached high frequency within the Ara<sup>-</sup> subpopulation. The  
437 expected fraction might be lower than two-thirds, however, because the Ara<sup>+</sup> subpopulation was  
438 increasing in frequency, and would be expected to generate an increasing proportion of the beneficial  
439 mutations, all else being equal. Contrary to this naïve expectation, however, all 18 populations in the  
440 MC treatment and 15 of the 18 populations in the MP treatment ended the experiment with  
441 descendants of the Ara<sup>-</sup> founders being numerically dominant (Figure S2).

442 This bias implies that one or more of the Ara<sup>-</sup> founders had greater potential for future adaptation  
443 than other founders. Of the six LTEE lineages that provided the founders used in our study, two of  
444 them—both with the Ara<sup>-</sup> state—evolved hypermutability during the LTEE (Tenaillon et al., 2016).  
445 The Ara-4 lineage became defective in mismatch repair (Sniegowski et al., 1997), while the Ara-1  
446 lineage acquired mutations in two enzymes that would normally prevent the misincorporation of  
447 oxidized nucleotides into DNA (Wielgoss et al., 2013). It is also possible that epistasis between new  
448 mutations and the various genetic backgrounds has led to differences in evolvability among the  
449 various founders. Background-dependent epistasis leading to differences in evolvability has been  
450 observed in the LTEE using replay experiments (Woods et al., 2011; Blount et al., 2012; Wünsche et  
451 al., 2017). In any case, the populations in the MC and MP treatments had reached similar fitness  
452 levels to those in the SC and SP treatments by generations 500 and 2000. Thus, the effects of both the  
453 initial standing variation and differences among the founders in their genetic potential for adaptation  
454 impacted only the earliest phases of evolution in the new D-serine environment.

455 Genetic variation is essential for populations to adapt to a new environment. We observed that pre-  
456 existing variation was important during the first ~50 generations in the D-serine medium, leading to  
457 substantial changes in the relative abundance of the different founders in the MC and MP treatments  
458 (Figure 3). Those changes depended on the initial genetic variation, which was identical by descent  
459 across the replicate populations in those treatments, and thus they indicate collateral evolution (Stern,  
460 2013; Lenski, 2017a). By contrast, the subsequent reversals in the relative abundance of descendants  
461 of those founders, and the fact that populations in all four treatments eventually achieved similar  
462 fitness levels (Figure 2), resulted from new mutations that arose independently in those populations,  
463 indicating parallel evolution (Stern, 2013; Lenski, 2017a). Thus, we observed both collateral and  
464 parallel evolution in our experiment with bacteria.

465 Two long-term experiments using *Drosophila* also reported collateral evolution, but they were not  
466 followed by parallel evolution (Burke et al., 2010; Graves et al., 2017). The longer generations and  
467 smaller populations of fruit flies probably limited the supply of new beneficial mutations, while  
468 sexual reproduction and the resulting segregation of pre-existing variation may have continued to fuel  
469 the ongoing response to selection. The importance of sexual reproduction with respect to the  
470 contributions of collateral versus parallel evolution was also evident in an evolution experiment  
471 performed using yeast (Burke et al., 2014). That experiment ran for 540 generations with large  
472 populations (10<sup>6</sup> cells during the transfer bottlenecks), and the populations were founded by a diverse  
473 set of diploids obtained by crossing wild strains. Although yeast can reproduce asexually, they  
474 underwent periodic mating and recombination in their experiment. As a consequence, segregating  
475 variation derived from the founders evidently fueled adaptation for the duration of the experiment,  
476 with little or no input from new beneficial mutations (Burke et al., 2014).

477 In any case, our study has shown that strictly asexual populations can also benefit from pre-existing  
478 variation, but the effect is likely to be smaller than in sexual populations. Moreover, any benefit of

479 pre-existing variation in asexual populations may often be short-lived, as we saw in our experiment,  
480 because that variation will be purged when new beneficial mutations sweep to fixation. In particular,  
481 it appears that the pre-existing alleles provided by the founders in our study were not sufficiently  
482 beneficial in the D-serine environment, such that the populations readily produced new mutations  
483 that provided greater benefits and displaced the initial variants. Even the populations in the SC  
484 treatment, which had no genetic diversity at the start of the experiment, achieved fitness levels  
485 comparable to the other treatments (Figure 2).

486 In future work, we intend to sequence the genomes of the founders and evolved samples from several  
487 timepoints. In addition to shedding light on the genetic basis of adaptation to growth on D-serine,  
488 genomic data will enable us to test and refine our inferences based on the fitness measurements and  
489 marker-ratio trajectories. In particular, we make several predictions that can be tested using genomic  
490 data. First, we expect to find an increased frequency of diagnostic alleles from the Ara+5 founders in  
491 the early (~50 generations) metagenomic samples from all of the populations in the MC and MP  
492 treatments. Second, we expect to see the alleles from Ara+5 subsequently disappear in all MC and  
493 most MP populations. Third, we predict that diagnostic alleles from one or more of the Ara<sup>-</sup> founders  
494 will achieve numerical dominance in all of the MC and many MP populations by generation 500 and  
495 remain dominant through generation 2000. In addition, genomic data should clarify whether one or  
496 both of the hypermutable founders (Ara-1 and Ara-4) in the MC and MP treatments dominated over  
497 time in a manner consistent with their having greater evolvability, in the sense of being able to adapt  
498 to the new environment. If so, that raises the interesting question of how the populations derived  
499 from the non-mutator founders in the SC and SP treatments achieved similarly high fitness. Perhaps,  
500 for example, the populations founded by mutators and non-mutators had similar beneficial mutations,  
501 but the hypermutators acquired them slightly earlier in the experiment.

502 In closing, our study contributes to filling the gap between the different experimental designs that are  
503 typically used with different model systems, and to understanding how these differences impact the  
504 dynamics and repeatability of evolution. While it remains difficult to observe adaptation driven by  
505 new mutations using long-standing model systems like *Drosophila*, we demonstrate that one can  
506 disentangle and estimate the contributions of standing variation and new mutations to adaptation in  
507 microbial systems. We also show that these contributions may depend on the particular history of the  
508 founders, and that the relative contributions of pre-existing variation and new mutations are highly  
509 sensitive to when they are measured after the evolving populations encounter a new environment.

510

## 511 **ACKNOWLEDGMENTS**

512 We thank Sydney Anway, Kate Bellgowan, Devin Lake, and Zachary Matson for assistance in the  
513 lab, and Thomas LaBar for valuable discussion. We acknowledge funding from the National Science  
514 Foundation (DEB-1813069), the USDA National Institute of Food and Agriculture (M1CL02253),  
515 the BEACON Center for the Study of Evolution in Action (NSF Cooperative Agreement DBI-  
516 0939454), the Yamada Science Foundation (to MI), and the Japan Society for the Promotion of  
517 Science (to MI).

518

## 519 **FUNDING**



520 This research was partially supported by the National Science Foundation (DEB-1813069), the  
521 USDA National Institute of Food and Agriculture (MICL02253), the BEACON Center for the Study  
522 of Evolution in Action (NSF Cooperative Agreement DBI-0939454), the Yamada Science  
523 Foundation (2016-5042), and the Japan Society for the Promotion of Science (201860528).

524

## 525 REFERENCES

- 526 Akey, J. M., Ruhe, A. L., Akey, D. T., Wong, A. K., Connelly, C. F., Madeoy, J., et al. (2010).  
527 Tracking footprints of artificial selection in the dog genome. *Proc. Natl. Acad. Sci. U.S.A.* 107, 1160–  
528 1165. doi: 10.1073/pnas.0909918107
- 529 Barrett, R. D. H., Rogers, S. M., and Schluter, D. (2008). Natural selection on a major armor gene in  
530 threespine stickleback. *Science* 322, 255–257. doi: 10.1126/science.1159978
- 531 Barrick, J. E., Kauth, M. R., Streliaoff, C. C., and Lenski, R. E. (2010). *Escherichia coli rpoB* mutants  
532 have increased evolvability in proportion to their fitness defects. *Mol. Biol. Evol.* 27, 1338–1347. doi:  
533 10.1093/molbev/msq024
- 534 Barrick, J. E., and Lenski, R. E. (2013). Genome dynamics during experimental evolution. *Nat. Rev.*  
535 *Genet.* 14, 827–839. doi: 10.1038/nrg3564
- 536 Blount, Z. D., Barrick, J. E., Davidson, C. J., and Lenski, R. E. (2012). Genomic analysis of a key  
537 innovation in an experimental *Escherichia coli* population. *Nature* 489, 513–518.  
538 doi:10.1038/nature11514
- 539 Burke, M. K., Dunham, J. P., Shahrestani, P., Thornton, K. R., Rose, M. R., and Long, A. D. (2010).  
540 Genome-wide analysis of a long-term evolution experiment with *Drosophila*. *Nature* 467, 587–590.  
541 doi: 10.1038/nature09352
- 542 Burke, M. K., Liti, G., and Long, A. D. (2014). Standing genetic variation drives repeatable  
543 experimental evolution in outcrossing populations of *Saccharomyces cerevisiae*. *Mol. Biol. Evol.* 31,  
544 3228–3239. doi: 10.1093/molbev/msu256
- 545 Graves, J. L., Hertweck, K. L., Phillips, M. A., Han, M. V., Cabral, L. G., Barter, T. T., et al. (2017).  
546 Genomics of parallel experimental evolution in *Drosophila*. *Mol. Biol. Evol.* 34, 831–842. doi:  
547 10.1093/molbev/msw282
- 548 Harshman, L. G., and Hoffmann, A. A. (2000). Laboratory selection experiments using *Drosophila*:  
549 What do they really tell us? *Trends Ecol. Evol.* 15, 32–36. doi: 10.1016/S0169-5347(99)01756-5
- 550 Hill, W. G., and Caballero, A. (1992). Artificial selection experiments. *Ann. Rev. Ecol. Syst.* 23, 287–  
551 310. doi: 10.1146/annurev.es.23.110192.001443
- 552 Hoffmann, A. A., Hallas, R. J., Dean, J. A., and Schiffer, M. (2003). Low potential for climatic stress  
553 adaptation in a rainforest *Drosophila* species. *Science* 301, 100–102. doi: 10.1126/science.1084296
- 554 Izutsu, M., Lake, D. M., Matson, Z. W. D., Dodson, J. P. and Lenski, R. E. (2022). Effects of  
555 periodic bottlenecks on the dynamics of adaptive evolution in microbial populations. *bioRxiv*  
556 [Preprint] doi:10.1101/2021.12.29.474457v1 (Accessed 27 May 2022)



- 557 Izutsu, M., Zhou, J., Sugiyama, Y., Nishimura, O., Aizu, T., Toyoda, A., et al. (2012). Genome  
558 Features of “Dark-Fly,” a *Drosophila* line reared long-term in a dark environment. *PLoS ONE*  
559 7:e33288. doi: 10.1371/journal.pone.0033288
- 560 Johnson, M. S., Gopalakrishnan, S., Goyal, J., Dillingham, M. E., Bakerlee, C. W., Humphrey, P. T.,  
561 et al. (2021). Phenotypic and molecular evolution across 10,000 generations in laboratory budding  
562 yeast populations. *ELife* 10:e63910. doi: 10.7554/eLife.63910
- 563 Lenski, R. E. (2017a). Convergence and divergence in a long-term experiment with bacteria. *Am.*  
564 *Nat.* 190, S57–S68. doi: 10.1086/691209
- 565 Lenski, R. E. (2017b). Experimental evolution and the dynamics of adaptation and genome evolution  
566 in microbial populations. *ISME J.* 11, 2181–2194. doi: 10.1038/ismej.2017.69
- 567 Lenski, R. E., Rose, M. R., Simpson, S. C., and Tadler, S. C. (1991). Long-term experimental  
568 evolution in *Escherichia coli*. I. Adaptation and divergence during 2000 generations. *Am. Nat.* 138,  
569 1315–1341. doi: 10.1086/285289
- 570 Marad, D. A., Buskirk, S. W., and Lang, G. I. (2018). Altered access to beneficial mutations slows  
571 adaptation and biases fixed mutations in diploids. *Nat Ecol Evol.* 2: 882–889. doi:10.1038/s41559-  
572 018-0503-9
- 573 Mery, F., and Kawecki, T. J. (2002). Experimental evolution of learning ability in fruit flies. *Proc.*  
574 *Natl. Acad. Sci. U.S.A.* 99, 14274–14279. doi: 10.1073/pnas.222371199
- 575 Roff, D. A. (1997). *Evolutionary Quantitative Genetics*. New York: Chapman and Hall.
- 576 Rose M. R. (1984). Laboratory evolution of postponed senescence in *Drosophila melanogaster*.  
577 *Evolution* 38, 1004-1010.
- 578 Scarcelli, N., and Kover, P. X. (2009). Standing genetic variation in FRIGIDA mediates experimental  
579 evolution of flowering time in *Arabidopsis*. *Mol. Ecol.* 18, 2039–2049. doi:10.1111/j.1365-  
580 294X.2009.04145.x
- 581 Schulte, R. D., Makus, C., Hasert, B., Michiels, N. K., and Schulenburg, H. (2010). Multiple  
582 reciprocal adaptations and rapid genetic change upon experimental coevolution of an animal host and  
583 its microbial parasite. *Proc. Natl. Acad. Sci. U.S.A.* 107, 7359–7364. doi: 10.1073/pnas.1003113107
- 584 Sniegowski, P. D., Gerrish, P. J., and Lenski, R. E. (1997). Evolution of high mutation rates in  
585 experimental populations of *E. coli*. *Nature* 387, 703-705. doi: 10.1038/42701
- 586 Sokal, R. R., and Rohlf, F. J. (1995). *Biometry: The Principles and Practice of Statistics in*  
587 *Biological Research, 3rd Edn.* New York: W.H. Freeman.
- 588 Stern, D. L. (2013). The genetic causes of convergent evolution. *Nat. Rev. Genet.* 14, 751–764. doi:  
589 10.1038/nrg3483
- 590 Tenailon, O., Barrick, J. E., Ribeck, N., Deatherage, D. E., Blanchard, J. L., Dasgupta, A., et al.  
591 (2016). Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature* 536, 165–  
592 170. doi: 10.1038/nature18959

- 593 Tenailon, O., Rodriguez-Verdugo, A., Gaut, R. L., McDonald, P., Bennett, A. F., Long, A. D., et al.  
594 (2012). The molecular diversity of adaptive convergence. *Science* 335, 457–461. doi:  
595 10.1126/science.1212986
- 596 Van den Bergh, B., Swings, T., Fauvart, M., and Jan, M. (2018). Experimental design, population  
597 dynamics, and diversity in microbial experimental evolution. *Microbiol. Mol. Biol. Rev.* 82:e00008-  
598 18. doi: 10.1128/MMBR.00008-18
- 599 Wielgoss, S., Barrick, J. E., Tenailon, O., Wisner, M. J., Dittmar, W. J., Cruveiller, S., et al. (2013).  
600 Mutation rate dynamics in a bacterial population reflect tension between adaptation and genetic load.  
601 *Proc. Natl. Acad. Sci. U.S.A.* 110, 222–227. doi: 10.1073/pnas.1219574110
- 602 Wisner, M. J., Ribbeck, N., and Lenski, R. E. (2013). Long-term dynamics of adaptation in asexual  
603 populations. *Science* 342, 1364–1367. doi: 10.1126/science.1243357
- 604 Woods, R. J., Barrick, J. E., Cooper, T. F., Shrestha, U., Kauth, M. R., and Lenski, R. E. (2011).  
605 Second-order selection for evolvability in a large *Escherichia coli* population. *Science* 331, 1433–  
606 1436. doi: 10.1126/science.1198914
- 607 Wright, S. I., Bi, I. V., Schroeder, S. G., Yamasaki, M., Doebley, J. F., McMullen, M. D., et al.  
608 (2005). The effects of artificial selection on the maize genome. *Science* 308, 1310–1314. doi:  
609 10.1126/science.1107891
- 610 Wünsche, A., Dinh, D. M., Satterwhite, R. S., Arenas, C. D., Stoebel, D. M. and Cooper, T. F.,  
611 (2017). Diminishing-returns epistasis decreases adaptability along an evolutionary trajectory. *Nat*  
612 *Ecol Evol.* 1: 0061. doi:10.1038/s41559-016-0061
- 613 Zhou, D., Udpa, N., Gersten, M., Visk, D. W., Bashir, A., Xue, J., et al. (2011). Experimental  
614 selection of hypoxia-tolerant *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U.S.A.* 108, 2349–  
615 2354. doi: 10.1073/pnas.1010643108

1 **Experimental Test of the Contributions of Initial Variation and New Mutations to**  
2 **Adaptive Evolution in a Novel Environment**

3 **Minako Izutsu<sup>1,2,3\*</sup>, and Richard E. Lenski<sup>1,2,3\*</sup>**

4 <sup>1</sup>Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,  
5 Michigan, USA

6 <sup>2</sup>BEACON Center for the Study of Evolution in Action, Michigan State University, East Lansing,  
7 Michigan, USA

8 <sup>3</sup>Ecology, Evolutionary Biology and Behavior Program, Michigan State University, East Lansing,  
9 Michigan, USA

10

11 **\* Correspondence:**

12 Minako Izutsu

13 [izutsumi@msu.edu](mailto:izutsumi@msu.edu)

14 Richard E. Lenski


15 [lenski@msu.edu](mailto:lenski@msu.edu)

16

17


18 **FIGURES**

19



20


21 FIGURE 1 | Experimental design. The colors indicate six different founder lineages. The actual  
22 colors of colonies on TA indicator agar plates are the same, except the cells derived from the four  
23  $Ara^-$  lineages produce red colonies while those derived from the two  $Ara^+$  lineages make pinkish-  
24 white colonies. See Materials and Methods for details of the procedures used.



25


26 FIGURE 2 | Relative fitness of the four treatments at generations 0, 500, and 2000 in the D-serine  
27 environment. Each symbol is the mean of 18 ln-transformed fitness estimates, and error bars show  
28 95% confidence intervals. See Figure S1 for each replicate population.





29

30 FIGURE 3 | Marker trajectories in the Mixed-Clonal (MC) and Mixed-Populations (MP) treatments  
31 during the first 500 generations. The marker ratio indicates the number of cells derived from the Ara<sup>-</sup>  
32 founder lineages divided by the number of cells derived from the Ara<sup>+</sup> founder lineages.



33

34 FIGURE 4 | Relative fitness of founder whole populations (A) and founder clones (B). The founders  
35 in each panel are arranged from lowest to highest fitness. The filled circles show the mean value of  
36 the ln-transformed fitness, based on 18 replicates for each founder. The error bars show 95%  
37 confidence limits, based on the *t*-distribution with 17 degrees of freedom and using the pooled  
38 standard deviation estimated from the corresponding ANOVAs (Table S2). Letters above the error  
39 bars identify sets of founders with values that are not significantly different, based on Tukey's test  
40 for multiple comparisons ( $p > 0.05$ ). For this analysis, we combined data for the SC and MC  
41 treatments, and similarly we combined data for the SP and MP treatments, because we used the same  
42 6 clonal or whole-population samples at generation 0 for those pairs of treatments (see Materials and  
43 Methods).