HSV-1 exploits host heterochromatin for egress

Hannah C Lewis*[1,2], Laurel E Kelnhofer-Millevolte*[1,2,3], Mia R Brinkley[1], Hannah E Arbach[1],

Edward A Arnold[1,4], Srinivas Ramachandran[5, 6, #], and Daphne C Avgousti[1,7,8, #]

[1] Human Biology Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Avenue

North, Seattle, WA 98109

[2] Molecular & Cellular Biology, Graduate Program, University of Washington and Fred

Hutchinson Cancer Research Center, 1959 NE Pacific Street Box 357275 Seattle, WA 98195

[3] UW Medical Scientist Training Program, 1959 NE Pacific Street Box 357470, Seattle, WA

98195

[4] Microbiology Graduate Program, University of Washington, 1705 NE Pacific Street Box

357735, Seattle, WA 98195

[5] RNA Bioscience Initiative, University of Colorado School of Medicine,

[6] Department of Biochemistry and Molecular Genetics, University of Colorado School of

Medicine, 12801 East 17th Avenue Mail Stop 8101, Aurora, CO 80045

[7] Twitter: @avgousti_lab

[8] Lead contact

* Authors contributed equally

Address correspondence to: srinivas.ramachandran@cuanschutz.edu (S.R.),

avgousti@fredhutch.org (D.C.A.)

Abstract

Herpes simplex virus replicates and forms progeny in the nucleus where it must overcome host chromatin to establish a successful infection. During lytic infection, newly formed viral capsids navigate through heterochromatin channels at the nuclear periphery to egress out of the nucleus. In uninfected cells, specific histone modifications such as trimethylation on histone H3 lysine 27 (H3K27me3) and the histone variant macroH2A1 delineate heterochromatin regions, or repressed chromatin, that are predominantly located in the nuclear periphery. We examined these markers during HSV-1 lytic infection in primary cells and discovered a striking increase in the levels of macroH2A1 and H3K27me3. Here, we demonstrate that the loss of macroH2A1 results in significantly lower viral titers but does not impair viral transcription, protein production, or replication. By inhibiting the deposition of H3K27me3 by EZH2, we further show that reduction of H3K27me3 also leads to a significant decrease in viral titers. Through chromatin profiling of macroH2A1 and H3K27me3, we define the specific chromatin regions that change dynamically during HSV-1 lytic infection and show that regions with increased macroH2A1 and H3K27me3 correlate with decreased host transcription as measured by RNA-seq. Furthermore, we find by electron microscopy that loss of macroH2A1 results in reduced heterochromatin at the nuclear periphery and significantly more viral capsids trapped in the nuclear compartment. Using both high and low shedding clinical isolates of HSV-1, we similarly find that HSV-1 titers are significantly reduced in the absence of macroH2A1. Our work demonstrates that heterochromatin imposes an impediment for viral egress, but HSV-1 can exploit its dynamic nature to successfully mediate escape from the nucleus.

INTRODUCTION

Nuclear replicating viruses must contend with host chromatin to establish a successful infection. Like most DNA viruses, herpes simplex virus (HSV-1) takes advantage of host histone factors both by incorporating histones onto its genome to promote gene expression (*1*) and by reorganizing host chromatin during infection (*2–4*). This redistribution of host chromatin results in a global increase in heterochromatin (*5*), or transcriptionally repressed chromatin. Host chromatin that is marginalized to the nuclear periphery during HSV-1 infection creates a potential barrier for capsids to egress from the nucleus. Elegant electron microscopy work showed that viral capsids traverse this dense chromatin in the nuclear periphery through channels of less dense staining, termed heterochromatin channels (*6*). However, the mechanisms by which these channels form are unclear.

In uninfected cells, specific histone modifications such as trimethylation of histone H3 lysine 27 (H3K27me3) and the histone variant macroH2A1 delineate heterochromatin regions that are largely localized to the nuclear periphery. H3K27me3 is deposited by the EZH2 enzyme (*7*), a member of the polycomb repressive complex (PRC2)(*8*). This modification is subsequently bound by PRC2, leading to modification of adjacent nucleosomes, which results in formation of dense heterochromatin and prevents transcription. The deposition of H3K27me3 is often considered to be a consequence of slowed transcription and the regulation of PRC2 is intimately linked to various biological processes including differentiation, proliferation, and maintaining cell identity (*8*). H3K27me3 is heavily deposited on the inactive X chromosome in female mammals, binding broad regions of the genome that overlap with macroH2A binding (*9*, *10*). MacroH2A, the largest of the histone variants, consists of a canonical histone fold domain, a small linker region and a C-terminal 25kDa macro domain that is thought to protrude from the nucleosome (*11*). There are three isoforms referred to collectively as macroH2A: macroH2A1.1, macroH2A1.2 and macroH2A2. MacroH2A1.1 and macroH2A1.2 are splice variants of the same gene that differ by one exon resulting in a 28-amino acid difference in the macro domain. The macro domain of macroH2A1.1 interacts directly with ADP-ribosylation, specifically through the amino acids that differ from macroH2A1.2 (*12*, *13*). MacroH2A1 was also found to be downregulated in melanoma (*14*), suggesting a key role in the maintenance of genome integrity. Importantly, it was reported that loss of macroH2A1 and macroH2A2 results in a significant decrease in heterochromatin in the nuclear periphery observed by electron microscopy (*15*). Furthermore, macroH2A1 also demarcates regions of host chromatin that associate with the nuclear lamina (*16*), termed lamina-associated domains (LADs). Given the complex role of

macroH2A in gene regulation and chromatin structure, studying its exploitation by a virus may

prove an important means to understanding its function.

Herpes simplex virus (HSV) infects an estimated two-thirds of the world's population where it

establishes latency and persists for the lifetime of the host (*17*, *18*). While most infections are

asymptomatic, HSV infection can have severe consequences in neonates and

immunocompromised individuals (*19–21*). During lytic infection, hundreds of host chromatin

factors are dynamically regulated (*3*). We hypothesized that several of these factors are

important for the formation of heterochromatin channels that allow HSV-1 to egress from the

nuclear compartment. We identified upregulation during HSV-1 infection of two key

heterochromatin markers: macroH2A1 and H3K27me3. Here, we use chromatin profiling via

Cleavage Under Targets and Tagmentation (CUT&Tag) of macroH2A1 and H3K27me3 to

define the specific host chromatin regions bound by these markers and how they change during

HSV-1 lytic infection. We show that regions with increased macroH2A1 and H3K27me3

correlate with decreased host transcription as measured by RNA-seq. Next, we demonstrate

that the loss of macroH2A1 results in significantly lower viral titers but does not impair viral

transcription, protein production, or replication. By inhibiting EZH2 deposition of H3K27me3, we

show that reduction of H3K27me3 also leads to a significant decrease in viral titers.

Furthermore, we demonstrate by transmission electron microscopy that loss of macroH2A1

results in reduced heterochromatin at the nuclear periphery and significantly more viral capsids

trapped in the nuclear compartment. Finally, we show that both high- and low-shedding clinical

isolates of HSV-1 also exhibit significantly reduced titers in the absence of macroH2A1. Our

study is the first to demonstrate that HSV-1 infection causes specific changes to

heterochromatin markers that are critical for efficient viral egress.

RESULTS

Heterochromatin markers macroH2A1 and H3K27me3 increase during infection.

Nuclear egress of HSV-1 capsids utilizes heterochromatin channels that form in the nuclear

periphery. We hypothesized that formation of these heterochromatin channels is dependent on

host chromatin markers deposited in specific genomic regions. To test this hypothesis, we first

examined the levels of multiple indicators of heterochromatin as measured by mass

spectrometry during HSV-1 infection in primary human foreskin fibroblast cells (HFFs) (*3*). HFF

cells are commonly used as a model for HSV-1 infection as they represent a common site of

infection in humans (*22*). We discovered striking upregulation of heterochromatin markers

macroH2A1 and H3K27me3 (Fig 1A and Fig S1A). Proteomics data identified the peptide

specific to exon 5 of macroH2A1.2, although other peptides detected as increasing may belong

to both isoforms of macroH2A1. Therefore, using antibodies specific to each isoform, we

examined the levels of macroH2A1.1 and macroH2A1.2 by western blot and found that both

isoforms increase during HSV-1 infection (Fig 1A). We also examined macroH2A2 levels and

found that they did not change during infection, indicating that changes to macroH2A1 levels are

distinct. Since both isoforms of macroH2A1 increase during infection, herein we refer to

macroH2A1 as including both macroH2A1.1 and macroH2A1.2. To determine whether these

results were specific to fibroblasts cells, we also examined macroH2A1 and macroH2A2 levels

during infection in another diploid cell line, retinal pigment epithelial cells (RPEs), which also

represent commonly infected cell types (*23*). We found that macroH2A1 levels, but not

macroH2A2, also increase in RPE cells during infection (Fig S1B). To define the impact of these

heterochromatin markers during HSV-1 infection, we next examined their genomic localization

during infection.

MacroH2A1 and H3K27me3 bind broad regions of the host genome that change

dynamically during HSV-1 infection

We next hypothesized that macroH2A1 is deposited at specific genomic loci on the host

genome during infection to promote the formation of heterochromatin. To test this hypothesis,

we used CUT&Tag to profile the genomic localization of macroH2A1 at 4, 8 and 12 hours post

infection (hpi) in wild-type hTERT-immortalized HFF cells (HFF-T) cells. As a control, we used

CRISPR-Cas9 to knock-out (KO) expression of the *macroH2A1* gene in HFF-T cells, which

resulted in the loss of macroH2A1.1 and macroH2A1.2, termed macroH2A1 KO cells. We also

examined the chromatin profile of H3K27me3 under these conditions to define the genomic

heterochromatin landscape. We used *E. coli* DNA as a spike-in to normalize all datasets. On the

human genome, we observed clear enrichment of H3K27me3 and macroH2A1 compared to IgG

in WT cells (Fig S1C-D). MacroH2A1 CUT&Tag enrichment is below that of IgG in macroH2A1

KO cells, but H3K27me3 enrichment is similar to WT, indicating that globally, CUT&Tag worked

as expected (Fig S1C-D). In contrast, the macroH2A1 CUT&Tag on the viral genome showed

similar enrichments in both WT and macroH2A1 KO cells, indicating that CUT&Tag was not

reporting on the protein enrichments on the viral genomes, and there was a significant

background signal from the viral genome that couldn't be accounted for (Fig S1I-J). Hence, we

focused our analysis on the host genome.

On the human genome, we observed large domains of macroH2A1 and H3K27me3 that were

gained upon viral infection (Fig 1B). Since large regions are not amenable to traditional peak-

based analysis, we instead used domain-based analysis. We partitioned all chromosomes into

domains of macroH2A1: domains had an enrichment that was two times the genome-wide

median and at least four-fold higher than the IgG control. With the minimum domain size of 1 kb,

we observed ~50,000 H3K27me3 domains and ~70,000 macroH2A1 domains genome-wide

(Fig S1E). We observed less than 10 macroH2A1 domains in the datasets generated from

macroH2A1 KO cells, indicating that our algorithm was identifying robust domains. We then

calculated the change in enrichments of macroH2A1 domains during the time course of

infection. We performed k-means clustering of the change in macroH2A1 enrichment in

macroH2A1 during the time course of infection compared to mock to identify patterns of

macroH2A1 gain or loss during infection. With k=6, we observed two clusters to substantially

gain macroH2A1 over the course of infection (Clusters 5 and 6, Fig 1C left, Fig S1F). Clusters 1-

3 had significant decreases in macroH2A1, whereas cluster 4 had a minor increase (Fig 1C left,

Figure S1F). We then asked how these macroH2A1-defined clusters behaved with respect to

H3K27me3. Surprisingly, the overall trends across the clusters were preserved in H3K27me3

(Fig 1C middle, Fig S1G), though the gains were not as high as macroH2A1. This indicated to

us that multiple forms of heterochromatin were gained at specific genomic loci due to viral

infection. We then asked if H3K27me3 increases were driven by macroH2A1 deposition. We

plotted the H3K27me3 changes at the same clusters for macroH2A1 KO cells (Fig 1C right, Fig

S1H). Surprisingly, the trends of H3K27me3 gain/loss were similar between WT and

macroH2A1 cells, pointing to the possibility that an upstream event during infection triggers

formation of new heterochromatin domains that contain both macroH2A1 and H3K27me3.

Furthermore, this upstream event is not affected by loss of macroH2A1. In summary, viral

infection results in formation of new heterochromatin domains that span ~10-100s of kilobases.

Heterochromatin formation is intimately tied to transcription: chromatin states influence

transcription and *vice versa*. Hence, we asked if we could observe changes in transcription that

corresponded to gain/loss of heterochromatin during infection. We performed RNA-seq on WT

and macroH2A1 KO cells over the course of infection and calculated fold changes in RNA levels

at each time point over the mock control. We identified genes that intersected with domains in

each cluster defined in Figure 1C and plotted the distribution of RNA fold changes of the genes

grouped by the clusters to which they belonged. We observed the change in RNA to anti-

correlate with change in macroH2A1: clusters 1-3 had an increase in RNA levels, whereas

clusters 4-6 had a decrease in RNA levels over the course of infection (Fig 2A). To determine

the order of events, that is if the gain in macroH2A1 leads to decreased transcription or if

decreased transcription leads to gain in macroH2A1, we performed the same analysis in

macroH2A1 KO cells. Strikingly, the gene expression changes in macroH2A1 KO cells mirror

that of WT cells, leading us to conclude that transcriptional changes are upstream of

macroH2A1 deposition (Fig 2B).

Steady state RNA changes could occur both due to changes in transcription and changes in

RNA decay. For changes in global RNA levels to cause changes in chromatin state, the most

direct connection would be via changes in transcription: transcription increases nucleosome

turnover (*24*) and counteracts heterochromatin formation and/or maintenance (*25*). Hence, we

asked if changes in RNA reflected changes in transcription. We analyzed published 4sU-RNA-

labeling data over the time course of infection (*26*) and found that the changes at 8hpi largely

followed the same trend as the total RNA: clusters 1-3 featured increased transcription, whereas

clusters 5 and 6 featured decreased transcription during infection (Fig 2C). Interestingly, cluster

4 diverged between total RNA and 4sU-RNA data: 4sU-RNA increased whereas total RNA

decreased. This indication of active transcription explains why cluster 4 features a weak

macroH2A1 gain.

Finally, we asked if the gain or loss of macroH2A1 happened across previously defined genome

compartments (*27*). We determined the distribution of the eigenvector corresponding to A/B

compartments for IMR90 Hi-C data from the 4DN project (*28*). Here, positive values correspond

to the A compartment, which features higher gene density and accessible chromatin, whereas

negative values correspond to the B compartment, which features inactive chromatin. All

clusters except cluster 2 are significantly biased towards one of the compartments: clusters 1

and 3 are significantly biased towards compartment B, whereas clusters 4-6 are significantly

biased towards compartment A (Fig 2D). Strikingly, the median of the clusters increases moving

from Cluster 1 to Cluster 6, correlating with macroH2A1 gain upon infection. Thus, macroH2A1

gains and losses happen in distinct genomic compartments upon viral infection. In summary,

decreased transcription in specific regions in compartment A due to viral infection causes gain

in macroH2A1 at these same regions and the gains in macroH2A1 are mirrored by gains in

H3K27me3. Taken together, these data suggest that macroH2A1 and H3K27me3 function to

form new regions of repressed chromatin on the host genome during HSV-1 infection.

Therefore, we next examined infection progression in the absence of these markers.

Loss of macroH2A1 results in reduced viral progeny but does not affect genome or protein accumulation

To investigate the role of macroH2A1 in HSV-1 infection, we infected WT and macroH2A1 KO HFF-T cells with HSV-1. Our CRISPR-Cas9 disruption of macroH2A1 resulted in loss of both macroH2A1.1 and macroH2A1.2, but not macroH2A2 (Fig 3A). Upon infection, we examined viral protein accumulation and observed similar levels of ICP0 (immediate early (*29*)), VP16 (early (*30*)) and glycoprotein H (gH, late (*31*)) in both wild type (WT) and macroH2A1 KO cells (Fig 3A). We next examined HSV-1 replication and measured viral genome accumulation by droplet digital PCR (ddPCR). We observed no significant difference in viral genome accumulation between WT and macroH2A1 KO cells (Fig 3B), indicating that neither HSV-1 replication nor protein production is affected by macroH2A1 loss. To determine the impact of macroH2A1 loss on viral progeny production, we measured viral genome accumulation in the supernatant. We discovered a 6-fold decrease in the levels of HSV-1 genomes produced in the supernatant of macroH2A1 KO cells compared to those produced from WT cells at 12 hpi (Fig 3C). We further discovered that infectious progeny produced from infected macroH2A1 KO cells were significantly decreased compared to those produced from WT cells at 8 and 12 hpi (Fig 3D). To determine whether these effects held true in multiple cell types, we also infected macroH2A1 KO RPE cells and found similar viral protein accumulation compared to control cells (Fig S2A). Viral titers produced from macroH2A1 KO RPE cells were also significantly lower than those produced from control cells (Fig S2B), consistent with our results in HFF-T cells. Importantly, we also quantified total viral RNA via RNA-seq and determined that there were no significant changes in viral RNA accumulation between WT and macroH2A1 KO cells (Fig S2C), consistent with our measurements of protein and genome accumulation. Taken together, these data indicate that loss of macroH2A1 leads to a significant defect in viral progeny but not protein, RNA, or genome accumulation. Therefore, we conclude that macroH2A1 is important for HSV-1 egress.

Depletion of H3K27me3 also results in reduced viral progeny

Given our observation that both macroH2A1 and H3K27me3 levels increase during HSV-1 infection (Fig S1A), we hypothesized that H3K27me3 is also important for viral egress. Therefore, we investigated how reduction of H3K27me3 would impact HSV-1 infection. To reduce H3K27me3 levels, we targeted the EZH2 enzyme that deposits this mark (*32*) using an inhibitor called Tazemetostat or "Taz" (EPZ-6438) (*33*). We treated cells with 10 µM of Taz or

DMSO control for 3 days to allow for reduction of H3K27me3 and then infected with HSV-1 (Fig

3E). We treated both WT and macroH2A1 KO cells to examine any synergistic effects of these

two heterochromatin markers. Consistent with mass spectrometry measurements, we observe

an increase in H3K27me3 levels as infection progresses (comparing 4, to 8 to 12hpi), although

the mock levels are higher than those at 4 hpi. This increase during infection is also observed in

the macroH2A1 KO cells, suggesting that it is not dependent on the presence of macroH2A1.

We found that accumulation of viral proteins ICP0, VP16 and gH were not affected by the

reduction of H3K27me3 (Fig 3E). Further, we observed that the reduction of H3K27me3 in

macroH2A1 KO cells also did not impact accumulation of these viral proteins at 4, 8 or 12hpi

(Fig 3F). We next examined viral genome accumulation and found that reduction of H3K27me3

resulted in a modest but significant 2-fold decrease in viral genome accumulation within cells

(Fig 3G, grey bars). We observed no additional decrease in viral genomes in macroH2A1 KO

cells treated with the inhibitor, suggesting that regardless of H3K27me3 levels, loss of

macroH2A1 does not impact viral replication as we observed above (Fig 3G, pink bars). We

next measured viral genomes in the supernatant to determine the impact of H3K27me3

reduction on progeny production and found a significant 4-fold decrease at 12 hpi from inhibitor-

treated cells compared to DMSO-treated (Fig 3H). Further, we found infectious progeny

produced from inhibitor-treated cells were significantly reduced at 12 hpi compared to those

produced from control cells (Fig 3I). Interestingly, H3K27me3 reduction in macroH2A1 KO cells

does not result in an additional defect in progeny production, suggesting the two markers

function in the same genetic pathway. Importantly, the modest decrease in viral replication

produced from cells lacking H3K27me3 does not fully explain the greater reduction in progeny,

suggesting that H3K27me3 is also important for HSV-1 egress. These results are consistent

with a previous report that found HSV-1 titers, but not viral replication, were reduced upon EZH2

inhibition (*34*). Furthermore, we did not observe any significant differences in the ratio of

genomes to plaque forming units (pfu) upon loss of macroH2A1, H3K27me3, or both, compared

to control conditions (Fig S2D), indicating that loss of these markers does not result in

significantly more defective particles. Taken together, these results indicate that H3K27me3 is

also important for efficient viral egress.

MacroH2A1-dependent heterochromatin in the nuclear periphery is critical for efficient

HSV-1 nuclear egress

HSV-1 capsids in the nucleus access the nuclear membrane through channels bracketed by

highly dense chromatin at the nuclear periphery (*4, 6, 35*). We found that loss of macroH2A1 or

depletion of H3K27me3 results in decreased viral progeny but has no or minimal impact on viral

replication. Therefore, we hypothesized that efficient HSV-1 egress out of the nuclear

compartment requires macroH2A1- or H3K27me3-dependent heterochromatin in the nuclear

periphery. Because loss of macroH2A1 resulted in no effect on viral replication compared to a

modest effect produced by inhibition of H3K27me3 deposition, we chose to focus on

macroH2A1 KO cells for heterochromatin visualization. We used transmission electron

microscopy (TEM) to image nuclei and examined heterochromatin formation in uninfected cells

(Fig 4A). We found that macroH2A1 KO cells had strikingly less heterochromatin in the nuclear

periphery than WT cells. This is consistent with what has been reported for hepatoma cells upon

loss of macroH2A1 and macroH2A2 (*15*). We quantified the width of heterochromatin at the

nuclear periphery in each cell type through binary thresholding of the intensity of electron-dense

regions and found that loss of macroH2A1 results in a significant reduction in peripheral

heterochromatin (Fig 4B). To test how this loss of macroH2A1-dependent heterochromatin

impacts HSV-1 nuclear egress, we infected these cells and quantified capsids in the nucleus

(Fig 4C-D and S3). Consistent with previous reports, in WT cells we found HSV-1 capsids

egressing through heterochromatin channels (Fig 4C and Fig S3B). We found that infected

macroH2A1 KO nuclei have significantly more HSV-1 capsids than those of WT cells (Fig 4D),

indicating that loss of macroH2A1-dependent heterochromatin is detrimental for efficient nuclear

egress. Herpesvirus infection produces three subtypes of capsids: A capsids which are empty,

B capsids which contain scaffolding proteins, and C capsids which contain viral gDNA and are

considered the precursor to infectious virions (*36*, *37*). To determine whether nuclear egress or

capsid formation is dependent on heterochromatin channels, we quantified the proportion of

each capsid type in WT and macroH2A1 KO cells. We found no difference in the proportion of

each capsid type between the two cell types (Fig S3C-D). Taken together, these data indicate

that macroH2A1-dependent heterochromatin in the nuclear periphery is critical for efficient

nuclear egress.

Clinical isolates of HSV-1 also require macroH2A1 for efficient egress

Our results above were generated using a lab adapted strain of HSV-1 (*syn 17*+), which may

not accurately represent a naturally occurring virus. Therefore, we decided to test whether

clinical isolates of HSV-1 would also be affected by loss of heterochromatin. We infected WT

and macroH2A1 KO cells with HSV-1 categorized as isolated from "low-shedding" or "high-

shedding" patients. We measured protein accumulation at 4, 8, and 12 hpi and viral genome

accumulation in cells and supernatant by ddPCR. We found that there was no significant

change in viral protein or viral genome accumulation for either clinical isolate in macroH2A1 KO

cells (Fig 5A-B). We measured viral genomes in the supernatant and found that there was a

modest reduction in HSV-1 genomes in the supernatant of macroH2A1 KO cells compared to

control cells (Fig 5C). For both clinical isolates, we found that macroH2A1 KO cells produced

significantly fewer progeny than WT cells at 12 hpi (Fig 5D). Taken together, these results

indicate that egress of clinically isolated HSV-1 is also dependent on macroH2A1.

DISCUSSION

Host chromatin is emerging as a critical barrier to viral success. Elegant electron microscopy

work showed that during HSV-1 infection, host chromatin is marginalized to the nuclear

periphery (*2*, *6*). Viral capsids subsequently egress through regions of less dense staining

termed heterochromatin channels (*35*). We found that markers of heterochromatin macroH2A1

and H3K27me3 were upregulated during HSV-1 infection. Therefore, we hypothesized that

these markers are critical for the formation of heterochromatin channels in the nuclear periphery

that allow for efficient viral egress from the nuclear compartment. Here, we showed that

macroH2A1 and H3K27me3 bind broad regions of the host genome which correlate with

decreased transcription, indicative of heterochromatin. Furthermore, we find that these regions

are primarily found in transcriptionally active compartments defined by Hi-C, supporting the idea

that newly formed heterochromatin increases during HSV-1 infection. However, the

mechanisms by which macroH2A1 and H3K27me3 are upregulated are still unclear. One recent

study showed that changes in chromatin accessibility during HSV-1 infection were akin to those

observed during heat shock or salt stress (*38*). Further, H3K27me3 is often deposited as a

consequence of transcriptional changes rather than the cause (*39*). We find that H3K27me3 is

still increased in large genomic regions (clusters 5-6) in the absence of macroH2A1 (Fig 1),

suggesting that it is deposited independently of macroH2A1. Taken together with published

findings, our study suggests that a global host response to the stress of infection results in a

decrease in transcription of many genes and thus increased macroH2A1 and H3K27me3-

dependent heterochromatin. This increased heterochromatin is then exploited by the virus as a

structure through which newly formed capsids can traverse to access the inner nuclear

membrane and egress (Fig 5E).

We found that loss of macroH2A1 or depletion of H3K27me3 resulted in significantly reduced

infectious progeny (Fig 3). Interestingly, we found that depletion of H3K27me3, but not loss of

macroH2A1, results in a modest but significant decrease in viral genome accumulation

compared to control cells. Because we did not detect H3K27me3 on viral genomes, consistent

with a recent study that detected very low levels (*40*), it is possible that inhibiting the deposition

of this mark has other effects in the cell. Indeed, this agrees with the complex effects observed

when multiple EZH1/2 inhibitors were compared in a recent study (*34*). They found treatment

with EZH1/2 inhibitors induced a cellular antiviral state that also suppressed infection with

multiple DNA and RNA viruses. Furthermore, proteomics revealed that other heterochromatin

marks like H3K9me3 are also changing during HSV-1 infection (*3*), suggesting that there may

be other mechanisms of heterochromatin dynamics at play. It will be interesting to consider the

diverse implications of heterochromatin domains not only as a means of transcriptional

regulation but also as a structural component that impacts many aspects of viral infection.

Using transmission electron microscopy, we showed that loss of macroH2A1 results in

decreased heterochromatin in the nuclear periphery. Our results echo those found in Hep2G

cells where loss of macroH2A1 and macroH2A2 was shown to cause significant disruption of

heterochromatin in the nuclear periphery (*15*). It is possible that the presence of macroH2A2 in

our macroH2A1 KO cells compensates to some degree, although we do not see any changes in

macroH2A2 either upon loss of macroH2A1 or during infection suggesting the upregulation

phenotype of macroH2A1 is distinct. It is also possible that different cell types have varying

degrees of dependence on macroH2A1-enriched heterochromatin that we are only beginning to

discern. We used HFFs and RPEs to test our hypothesis, however, in both cases we used

hTERT-immortalized diploid cells. MacroH2A1 was also reported to regulate telomere

maintenance (*41*), suggesting a threshold of macroH2A1 may be required for cell survival. Our

CUT&Tag data indicate that while macroH2A1 is enriched in large regions defined in clusters 5

and 6, we also observe a decrease in clusters 1-3. This suggests that detecting a global

increase in total levels of macroH2A1 may be variable. Indeed, we observed that infected

primary HFFs consistently showed an increase in macroH2A1 while HFF-T cells infected with

clinical isolates of HSV-1 did not. Furthermore, as the heatmaps in Figure 1 indicate, large

regions of the genome are both losing and gaining macroH2A1 and H3K27me3 during infection.

Taken together, it is clear that other factors regulate macroH2A1 levels and future work to

determine the mechanism of upregulation as a response to infection will further define the host

response.

Our electron microscopy revealed greater numbers of capsids trapped in the nucleus of

macroH2A1 KO cells. Our analysis uncovered that the proportion of capsid type was not

different in WT compared to macroH2A1 KO cells (Fig S3C-D), suggesting that heterochromatin

does not play a role in capsid formation but rather acts as a structural highway for formed capsids to reach the inner nuclear membrane. At the inner nuclear membrane, capsids are aided by the nuclear egress complex, comprised of viral proteins UL34 and UL31, to bud into the inner nuclear membrane and fuse with the outer nuclear membrane for further maturation steps in the cytoplasm (*42*). Multiple reports describe that not only does lamina disruption occur independently of capsid formation, but that chromatin is likely the limiting factor in viral egress (*43–48*). Taken together, our results demonstrate that macroH2A1 and H3K27me3- dependent heterochromatin is exploited by HSV-1 to efficiently egress from the nuclear compartment.

AUTHOR CONTRIBUTIONS

Conceptualization, H.C.L, L.K.M., and D.C.A; methodology H.C.L, L.K.M., and D.C.A.; formal analysis, H.C.L, L.K.M., S.R., and D.C.A.; investigation, H.C.L, L.K.M., M.R.B., H.E.A, E.E.A, and D.C.A; resources, H.C.L, L.K.M., M.R.B., H.E.A, E.E.A, S.R., and D.C.A.; data curation, H.C.L, L.K.M., M.R.B., S.R., and D.C.A.; writing – original draft, H.C.L, L.K.M., M.R.B., S.R., and D.C.A.; writing – review & editing, H.C.L, L.K.M., M.R.B., H.E.A, E.E.A, S.R., and D.C.A.; visualization, H.C.L, L.K.M., S.R., and D.C.A.; supervision, H.C.L, L.K.M., and D.C.A.; funding acquisition, H.C.L., E.A.A, S.R., and D.C.A.

ACKNOWLEDGEMENTS

We thank members of the Avgousti lab, M. Emerman, A. Geballe, M. Lagunoff, K. Lynch, S. Parkhurst, and M. Weitzman for insightful comments. We thank D. Janssens and the Henikoff lab for reagents and assistance with CUT&Tag. We thank the Salama and Overbaugh labs for assistance with ddPCR. We thank the Jerome lab for the clinical HSV isolates and the Cellular Imaging, Electron Microscopy (EMSR), F. Wu and Genomics & Bioinformatics Shared Resource Facilities at the Fred Hutchinson Cancer Research Center for help with sequencing and data analysis. We thank J. Dubrulle and the Cellular Imaging Shared Resource (CISR) for help with image analysis. The EMSR, CISR, and Genomics core are supported in part by the Fred Hutch/University of Washington Cancer Consortium (P30 CA015704). This study was supported by start-up funds from the RNA Bioscience Initiative at the University of Colorado School of Medicine (S. R.), the Fred Hutch (D.C.A.), and NIH funding to H.C.L. (AI083203), E.A.A. (AI083203), S.R. (GM133434), and D.C.A. (GM133441).

Figure 1. MacroH2A1 and H3K27me3 increase during lytic HSV-1 infection and form broad chromatin regions.

Figure 1. MacroH2A1 and H3K27me3 increase during lytic HSV-1 infection and form

broad chromatin regions.

A. Western blot of total macroH2A1 (mH2A1), mH2A1.1, mH2A1.2 and mH2A2 with

relative quantification below indicated blots during HSV-1 infection showing mock

415	infected (M), 4-, 8-, and 12-hours post-infection (hpi) in primary HFFs. ICP0, VP16, gH
416	are shown as representative HSV-1 proteins. H4 is a loading control.
417	B. Representative genome browser snapshots of spike-in normalized CUT&Tag enrichment
418	of macroH2A1 and H3K27me3 showing increases at 8hpi of HSV-1 infection as
419	measured by CUT&Tag in WT or macroH2A1 KO as indicated. The region
420	corresponding to chromosome 1, 28,500 kb to 28,660 kb, is shown.
421	C. Changes in log2 enrichment of spike-in normalized CUT&Tag of macroH2A1 and
422	H3K27me3 over IgG compared to mock treatment are shown as a heatmap. Each line in
423	the heatmap represents a domain of macroH2A1.
424	

Figure 2. MacroH2A1 and H3K27me3 presence correlates with decreased transcription.

Figure 2. MacroH2A1 and H3K27me3 presence correlates with decreased transcription.

A. Box plots of total host RNA from RNA-seq for genes overlapping clusters from Figure 1. Kolmogrov-Smirnov test (in R) was performed comparing logCPM values of genes at each time point with logCPM values of genes from mock dataset for each cluster. The p-values were corrected for multiple testing, and the time points and clusters with corrected p-value less than 0.05 are shaded.

B. Same as A for macroH2A1 KO cells.

C. 4sU-RNA counts for genes from Hennig *et al* (*38*) overlapping with clusters from Figure 1 shown as box plots and shading as indicated in (A).

D. Box plots of Hi-C eigenvector scores of regions overlapping with clusters from Figure 1. The Hi-C compartment eigen vector scores were obtained from 4DN project website. Wilcoxon signed rank test with alternate hypothesis that the true location is not equal to 0 was performed on the distribution of eigen vector scores for each cluster. The p-values

440   were corrected for multiple testing, and the clusters with corrected p-value less than 0.05

441   are shaded. Bonferroni correction was used for p-value adjustments.

Figure 3. MacroH2A1 and H3K27me3 are required for efficient HSV-1 egress.

Figure 3. MacroH2A1 and H3K27me3 are required for efficient HSV-1 egress.

A. Western blot of total macroH2A1 (mH2A1), mH2A1.1, mH2A1.2 isoforms and mH2A2 during HSV-1 infection. Mock infected (M), 4-, 8-, and 12-hours post-infection in WT and macroH2A1 knockout (mH2A1 KO) HFF-Ts are shown as indicated. Quantification of relative intensity is shown below indicated blots and H3 is shown as a loading control.

B.   Droplet digital (ddPCR) quantification of HSV-1 genomes extracted from the infected
cells as indicated for each time point. Error bars represent the SEM of three biological
replicates.

C.   ddPCR quantification of HSV-1 genomes released from WT or mH2A1 KO cells and
isolated from supernatants (sups). Error bars represent the SEM of three biological
replicates.

D.   Infectious progeny produced by WT or mH2A1 KO cells quantified by plaque assay. Viral
yield is indicated as the percent yield compared to WT and was calculated by dividing
plaque-forming units (pfu) per mL of the mH2A1 knockout titers by the WT titers of each
respective virus at indicated time points. * $p < 0.05$, **** $p < 0.0001$.

E.   Western blot of H3K27me3 and mH2A1 during HSV-1 infection shown as mock infected
(M), 4-, 8-, and 12-hours post-infection. WT cells were pretreated with DMSO or
tazemetostat inhibitor (Taz) for 3 days prior to infection and remained with inhibitor over
the course of infection. Beta-actin is shown as a loading control. Quantification of relative
intensity over beta-actin normalized to the mock sample on the same blot are shown.

F.   Western blot as in E for macroH2A1 KO cells.

G.   Infection as in E. ddPCR quantification of HSV-1 genome accumulation as in B. Error
bars represent the SEM of three biological replicates. ** $p < 0.01$, *** $p < 0.001$.

H.   Infection as in E and F with quantification as in C comparing HSV-1 genomes released
from Taz or DMSO treated WT or mH2A1 knockout. Error bars represent the SEM of
three biological replicates. ** $p < 0.01$, *** $p < 0.001$.

I.   Infection as in E and F quantifying infectious progeny produced by WT or mH2A1
knockout cells with DSMO or Taz pretreatment by plaque assay. Viral yield is indicated
as the percent yield compared to WT as in D. **** $p < 0.0001$.

Figure 4. HSV-1 requires macroH2A1-dependent hetero-
chromatin for efficient egress.

Figure 4. HSV-1 requires macroH2A1-dependent heterochromatin for efficient egress.

A. Transmission electron microscopy (TEM) images of representative nuclei of uninfected
   WT (left) or macroH2A1 KO (right) cells. Regions outside of the nucleus are colorized
   orange. Dark regions (arrowhead) represent high density heterochromatin. Scale bar is
   1µm.

B. Quantification of peripheral heterochromatin width in uninfected nuclei from A. Width
   was measured in nm from the nuclear periphery via binary thresholding from intensity
   profiles sampled every 10 pixels. Mean width was plotted for each nucleus. $p < 0.05$
   (n=16 WT, n=22 macroH2A1 KO).

C. TEM images of representative WT and macroH2A1 knockout nuclei at 18 hours post-
   infection with HSV-1. Insets show enlarged views of respective boxed areas containing

493   capsids within nuclei with scale bars as indicated. Arrowhead indicates heterochromatin,

494   arrow indicates viral capsid.

495   D. Quantification of all capsids within nuclei from C. Number of capsids was normalized

496   according to nucleus area in μm^2. $p < 0.01$ (n=40 WT, n=55 mH2A1 KO).

Figure 5. MacroH2A1 is required for egress of clinical HSV-1 isolates.

Figure 5. MacroH2A1 is required for egress of clinical HSV-1 isolates.

A. (A-D) WT or macroH2A1 (mH2A1) KO HFF-T were mock-infected (M) or infected with a
low-shedding or high-shedding clinical HSV-1 isolate as indicated for 1, 4, 8 and/or 12

hours. Western blots of proteins as indicate with beta-actin shown as loading control.

Fold-change of mH2A1 protein levels normalized to mH2A1 level in mock-infected cells

are shown below the indicated blots.

B. Infection as in A. ddPCR quantification of HSV-1 genome accumulation within cells.

Error bars represent the SEM of three biological replicates.

C. ddPCR quantification of HSV-1 genomes released from WT or macroH2A1 knockout

cells. Error bars represent the SEM of three biological replicates. * $p < 0.05$.

D. Infectious progeny produced by WT or macroH2A1 knockout cells quantified through

plaque assay. Viral yield is indicated as the percent yield compared to WT. * p. < 0.05, **

$p < 0.01$, **** $p < 0.0001$.

E. Model for heterochromatin support of HSV-1 egress. In uninfected cells (left),

heterochromatin is enriched in the nuclear periphery. Upon infection with HSV-1

(middle), there is an increase in heterochromatin at specific regions in the nuclear

periphery that results in formation of heterochromatin channels that allow capsids to

traverse from the viral replication centers (VRCs) the inner nuclear membrane where

they bud into the perinuclear space and subsequently fuse with the outer nuclear

membrane into the cytoplasm. In infected cells missing heterochromatin markers

macroH2A1 or H3K27me3 (right), these heterochromatin channels cannot form properly

which results in increased capsids trapped in the nuclear compartment.

 SUPPLEMENTAL FIGURES

Supplemental Figure S1. Quantification of macroH2A1 and H3K27me3 enrichment during HSV-1 infection.

Supplemental Figure S1. Quantification of macroH2A1 and H3K27me3 enrichment during

HSV-1 infection.

A. Relative peptide abundance of total macroH2A1 and H3K27me3 by mass spectrometry over the course of HSV-1 infection in primary HFFs from (*3*). Values shown are normalized to mock levels.

B. Western blot of total macroH2A1 (mH2A1), macroH2A1.1, macroH2A1.2, macroH2A2, and H3K27me3 in mock infected (M), and 4-, 8-, and 12-hours post-infection with HSV-1 in RPEs. ICP0, VP16, gH are HSV-1 proteins. H4 is shown as a loading control.

C. Spike-in normalized, genome-wide CUT&Tag read density of macroH2A1 from WT HFF-T and macroH2A1 KO cellular genomes during HSV-1 infection.

D. Spike-in normalized, genome-wide CUT&Tag read density of H3K27me3 on WT HFF-T and macroH2A1 KO cellular genomes during HSV-1 infection.

E. Total number of macroH2A1 and H3K27me3 domains over 1 kb as measured by CUT&Tag of WT HFF-T and macroH2A1 KO cell genomes during HSV-1 infection.

F. Quantification of heatmaps from Figure 1C showing macroH2A1 enrichment across the clusters.

G. Quantification of heatmaps from Figure 1C showing H3K27me3 enrichment in clusters defined by macroH2A1 as in (H).

H. Quantification as in (I) in macroH2A1 KO cells.

I. Spike-in normalized CUT&Tag read density of macroH2A1 on HSV-1 viral genomes during infection of WT HFF-T and macroH2A1 KO cells.

J. Spike-in normalized CUT&Tag read density of H3K27me3 on HSV-1 viral genomes during infection of WT HFF-T and macroH2A1 KO cells.

Supplemental Figure S2. MacroH2A1 is required for egress but does not impact HSV-1 transcription or particle to genome ratios.

Supplemental Figure S2. MacroH2A1 is required for egress but does not impact HSV-1

transcription or particle to genome ratios.

A. Western blot of total macroH2A1 (mH2A1), macroH2A1.1, macroH2A1.2 and

macroH2A2 proteins as indicated in mock infected and 4-, 8-, and 12-hpi with HSV-1 in

hTERT-immortalized RPEs and macroH2A1 knockout hTERT-immortalized RPEs. H3 is

shown as a loading control.

B. Infectious progeny produced by WT or macroH2A1 knockout RPE cells were quantified

by plaque assay. Viral yield is indicated as the percent yield compared to WT and was

calculated by dividing pfu per mL of the macroH2A1 knockout titers by the WT titers of

each respective virus at each indicated time point. *** p. < 0.001.

C. Log2 fold-change of HSV-1 transcripts at 4, 8 and 12 hpi identified by RNA-seq in

macroH2A1 KO compared to WT HFF-T cells. There is no significant difference between

the cell types.

D. Genome to pfu ratio of HSV-1 in WT or mH2A1 knockout HFF-T cells treated with 10 uM

tazemetostat (Taz) or DMSO. Ratio was calculated from the samples used to generate

Figure 3H-I. No significance by ANOVA.

Supplemental Figure S3. HSV-1 capsid formation is independent of macroH2A1.

Supplemental Figure S3. HSV-1 capsid formation is independent of macroH2A1.

A. Transmission electron microscopy (TEM) images of representative nuclei of uninfected WT (left) or macroH2A1 KO (right) HFF-Ts. Regions outside of the nucleus are colorized orange. Insets show enlarged views of respective boxed areas containing peripheral heterochromatin. Arrowhead indicates high density heterochromatin.

B. TEM images of representative WT and macroH2A1 knockout nuclei at 18 hours post-infection with HSV-1. Insets show enlarged views of respective boxed areas containing capsids within nuclei. Arrowhead indicates heterochromatin, arrow indicates viral capsid.

C. TEM images of representative A (empty), B (scaffolding proteins, but no genome), and C (full) HSV-1 capsids. Scale bars as noted for all TEM images.

586  D. Quantification of capsid types within nuclei from B. Values for each capsid type are

587    shown as a proportion of total capsids.

589 **MATERIALS AND METHODS**

590 **Cells and viruses:**

591 Primary human foreskin fibroblasts (HFFs), hTERT-immortalized HFFs, hTERT-immortalized

592 macroH2A1 knockout HFFs were cultured using standard methods with 10% FBS and 1%

593 penicillin-streptomycin as previously described (*49*). Cells were grown at 37 °C with 5% CO_2

594 and tested for mycoplasma contamination approximately once a month.

595 The lab adapted strain of HSV-1, *syn 17+* (*50*) was used for all experiments unless otherwise

596 noted. Monolayers of cells were infected for 1 hour at 37 °C as previously described (*51*). An

597 MOI of 3 was used for all experiments, except for EM samples which used MOI of 10. Cells

598 were collected at mock, 4-, 8-, 12- hpi for western blot. Supernatant was collected at 8- and 12-

599 hpi for plaque assays. Virus stock was grown by infecting Vero cells at an MOI of 0.0001. Virus

600 was harvested ~60-80 hours post-infection and titered on U2OS cells to determine stock plaque

601 forming units per mL (PFU/mL). Experimental plaque assays were set up in Vero cells. Plaque

602 assays were set up on serial 10-fold dilutions in serum-free DMEM. Virus was left on cells for 1

603 hour then aspirated. Cells were washed with 1X PBS (pH 7.46) and 2% methylcellulose overlay

604 in DMEM with 2% FBS and 0.5% penicillin-streptomycin was added to wells. Plaques were fixed

605 with 0.2% crystal violet between 96-100 hours post infection and plaques were counted by

606 hand. All plaque assays were set up with 2 technical replicates.

607 Clinical isolates were acquired in 1994 and 1995 from oral swab collections. Samples were de-

608 identified on collection. Patients provided daily home collections of oral swabs, and qPCR was

609 used to detect how many days HSV was detected in these swabs. Patients that had detectable

610 HSV in only a few swabs were classified as "low-shedders" and patients that had a high

611 percentage of days with positive swabs were classified as "high-shedders." Stocks were a gift

612 from the lab of Keith Jerome.

613 Table 1: Cell line sources.

Cell Line	Source
Human: RPE-1 (hTERT-immortalized)	Gift of E. Hatch Lab (*52*) (Invitrogen)
Human: Primary HFF	Gift of D. Galloway Lab
Human: HFF (hTERT-immortalized)	Gift of J. Kamil Lab (*53*)

| African Green Monkey: Vero | Gift of A. Geballe Lab (*54*) |

Knockout of macroH2A1:

MacroH2A1 guide RNA (gRNA, see Table 3) was cloned into TLCv2 (Addgene plasmid: 87360),

a plasmid encoding doxycycline-inducible Cas9-2A-GFP and gRNA expression. To generate

lentiviral particles, 1.0×10^7 HEK293T cells were transfected with 12 µg TLCV2_sg_mH2A1, 8 µg

pMDL (Addgene plasmid: 12251), 4 µg VSVg, and 2.5 µg pREV (Addgene plasmid: 115989)

using Attractene transfection reagent (Qiagen: 301005). Lentivirus was harvested at 24-, 48-,

and 72-hours post-transfection, filtered through 0.2 µm filters, aliquoted, flash frozen with liquid

nitrogen, and stored at -80C. 2 mL of thawed filtered lentivirus was used to transduce

approximately 2.25×10^6 HFF-T or 3×10^6 RPE cells in 10 cm plates with 8 µg/mL polybrene

(Millipore-Sigma: TR-1003-G). Cells were allowed to reach confluence before selection with 1

µg/mL puromycin for 3 days. Cells were then sorted by GFP positivity or counted by

hemocytometer, plated at 1 cell per well into 96-well plates, and selected through serial

expansion of colonies. Selected cells were screened for macroH2A1 knockout by western blot.

Infections with Tazemetostat pretreatment:

629 HFFs and macroH2A1 KO HFFs were treated with DMSO or 10 µM of Tazemetostat (Taz)

630 (MedChem: HY-13803) in DMSO for 3 days prior to infection. Cells were then infected at MOI of

631 3 and after 1 hour of incubation with virus, fresh media with 10µM Taz was added to previously

632 treated cells. Control samples were treated with equivalent volumes of DMSO. Samples were

633 harvested as above.

Western Blotting:

Western blotting was performed as previously described (*49*). Briefly, cells were counted,

pelleted, resuspended in 1x NuPAGE lithium dodecyl sulfate (LDS) sample buffer (Fisher

Scientific: NP007) + 5% 2-betamercaptoethanol, and boiled for 15 minutes. Protein lysates were

separated by 13.5% SDS-PAGE gels using NuPAGE MOPS buffer (Fisher Scientific: NP0001)

at 75 volts for 30 minutes then 110 volts for 100 minutes, then transferred to a nitrocellulose

membrane (Bio-Rad) at 100 volts for 70 minutes using Transfer Buffer (25 mM Tris Base, 100

mM glycine, 20% methanol). Membranes were ponceau stained and imaged. Membranes were

blocked in 5% milk in Tris-buffered saline with Tween (TBST) for 1 hour, then probed with

primary antibody overnight (see Table 2). Membranes were washed with TBST for thirty

minutes, incubated with secondary antibodies conjugated to horseradish peroxidase (α-mouse

or α-rabbit; 1:5000) at room temperature for 1 hour, washed with TBST for thirty minutes, and

detected using Clarity Western ECL Substrate (Bio-Rad: 1705061) and Chemidoc MP Imaging

System (Bio-Rad). Images were formatted using Adobe Photoshop and Illustrator.

Table 2: Sources and identification of antibodies used.

Antibody	Source	Identifier	Use (Concentration)
Mouse anti-actin	Abcam	Cat: 5441	WB (1:10,000)
Mouse anti-gH	Abcam	Cat: 110227	WB (1:1,000)
Rabbit anti-H3	Abcam	Cat: 1791	WB (1:20,000)
Rabbit anti-H3K27me3	Cell Signal Technologies	Cat: 9733T	WB/CUT&Tag (1:100)
Rabbit anti-H3K9me3	Abcam	Cat: 8898	WB (1:1,000)
Rabbit anti-H4	Abcam	Cat: 10158	WB (1:1,000)
Mouse anti-ICP0	Santa Cruz Biotechnology	Cat: 53070	WB (1:1,000)
Rabbit anti-macroH2A1	Abcam	Cat: 37264	WB (1:1,000) CUT&Tag (1:100)
Rabbit anti-macroH2A1.1	Cell Signaling Technology	Cat: 12455S	WB (1:1,000)
Mouse anti-macroH2A1.2	Cell Signaling Technology	Cat: 4827S	WB (1:1,000)
Rabbit anti-macroH2A2	Active Motif	Cat: 39874	WB (1:1,000)
HRP-conjugated anti-VP16	Santa Cruz Biotechnology	Cat. 7546	WB (1:1,000)
Peroxidase-AffiniPure Goat Anti-Mouse	Jackson Immunoresearch Laboratories	Cat: 115-035-003	WB (1:5,000)
Peroxidase-AffiniPure Goat Anti-Rabbit	Jackson Immunoresearch Laboratories	Cat: 111-035-045	WB (1:5,000)
Rabbit anti-mouse IgG	Abcam	Cat: 46540	CUT&Tag (1:100)
Guinea pig anti-rabbit IgG	Antibodies Online	Cat: AB1N101961	CUT&Tag (1:100)

Quantification of HSV-1 genomes by ddPCR:

Cells were harvested at the indicated times post-infection by trypsinization, washed with 1X

PBS and centrifuged at 5000 xg for 2 minutes. Pellets were flash-frozen in liquid nitrogen and

stored at -80 °C until processed. HSV-1 DNA within cells was isolated from frozen pellets using

QIAamp DNAMini Kit (Qiagen: 51304).

Supernatants were harvested at the indicated times post-infection, centrifuged at > 3500 xg and

filtered through 40 micron sterile syringe filters. DNA on the exterior of filtered capsids was

digested for 1 hour at 25 °C with 20.3 units DNase (Qiagen: 79254) supplemented with 10 mM

$MgCl_2$. DNase was inactivated at 75 °C for 10 minutes followed by vortexing. Capsids were

digested with 3 mg/mL proteinase K (Fisher Scientific: BP1700) in 100 mM KCl, 25 mM EDTA,

10 mM Tris-HCl pH 7.4 and 1% Igepal for 1 hour at 50 °C. HSV-1 genomes from digested

capsids were isolated using QIAamp DNAMini Kit.

A duplexed digital droplet PCR was performed to measure the levels of cellular or supernatant

HSV-1 genomes on the QX100 droplet digital PCR system (Bio-Rad Laboratories, Hercules,

CA) using a primer/probe set specific to HSV-1 gB. Cell numbers were determined using a

primer/probe set specific to human Beta-globin, a reference gene that exists at two copies per

cell. See Table 3 for oligonucleotide sequences, as have been previously established for HSV-1

(*55*). The digital droplet PCR (ddPCR) reaction mixture consisted of 12.5 µL of a 2X ddPCR

Supermix for Probes no dUTP (Bio-Rad: 1863024), 1.25 µL of each 20X primer-probe mix, and

10 µL of template DNA. 20 µL of each reaction mixture was loaded onto a disposable plastic

cartridge (Bio-Rad: 1864008) with 70 µL of droplet generation oil (Bio-Rad: 1863005) and

placed in the droplet generator (Bio-Rad). Droplets generated were transferred to a 96-well PCR

plate (Bio-Rad 12001925), and PCR amplification was performed on a Bio-Rad C1000 Touch

Thermal Cycler with the following conditions: 95 °C for 10 minutes, 40 cycles of 94 °C for 30

seconds, and 60 °C for 1 minute, followed by 98 °C for 10 minutes and ending at 4 °C. After

amplification, the plate was loaded onto the droplet reader (Bio-Rad: QX200), and the droplets

from each well of the plate were automatically read with droplet reader oil (Bio-Rad: 186-3004)

at a rate of 32 wells per hour. Data were analyzed with QuantaSoft analysis software and

quantitation of target molecules presented as copies per µL of PCR reaction. HSV-1 genome

values were standardized to cellular β-globin levels. Experiments were completed in biological

triplicate and statistical analysis was performed using Prism v10 (GraphPad Software).

Table 3: Primers used for knockout and ddPCR

CRISPR MacroH2A1 gRNA	CCTCAATAGCAAGCCATCCTGT
HSV1_gB_qPCR_F1	CCGTCAGCACCTTCATCGA
HSV1_gB_qPCR_R1	CGCTGGACCTCCGTGTAGTC
HSV1_gB_probe1	6FAM-CCACGAGATCAAGGACAGCGGCC-BHQ1
beta-globin_qPCR_F1	TGAAGGCTCATGGCAAGAAA
beta-globin_qPCR_R1	GCTCACTCAGTGTGGCAAAGG
beta-globin_probe2_hex	/5HEX/TC AGG TGA GCA GCC ATC ACT A/3BHQ_1/

Cleavage Under Targets & Tagmentation (CUT&Tag):

Two biological replicates per time point were obtained from independent infections. Protocol was adapted from the established CUT&Tag methods (*56*). Cells were harvested using trypsin, washed three times with ice-cold phosphate-buffered saline (PBS) via centrifugation at 600 xg for 3 minutes and counted using a hemocytometer. Nuclei from 600,000 cells were isolated by hypotonic lysis in 1 mL buffer NE1 (20 mM HEPES-KOH pH 7.9; 10 mM KCl; 0.5 mM spermidine; 0.1% Triton X-100; 20% Glycerol; Roche EDTA-free protease inhibitor) for 10 minutes on ice followed by centrifugation at 1300 xg for 4 minutes. Nuclei were resuspended in Wash buffer (20 mM HEPES-NaOH pH 7.5; 150 mM NaCl; 0.5 mM spermidine; Roche EDTA-free protease inhibitor) and counted using a hemocytometer. BioMag Plus Concanavalin A coated beads (Polysciences: 86057-3) were equilibrated with Binding buffer (20 mM HEPES-KOH pH 7.9; 10 mM KCL; 1mM $CaCl_2$; 1 mM $MnCl_2$). Beads (5 µL) were mixed with aliquots of 75,000 nuclei and incubated at 25 °C for 10 minutes followed by magnetic separation of beads. Beads were resuspended in 50 µL primary antibody [anti-mH2A1 (Abcam: ab37624), anti-H3K27me3 (Cell Signaling:9733) or anti-mouse IgG (Abcam: ab46540) in Wash buffer supplemented with 2 mM EDTA and 0.1% bovine serum albumin (BSA) and incubated on a nutator at 4 °C overnight. The beads were decanted on a magnet stand then resuspended in 50 µL secondary antibody [Guinea pig anti-rabbit IgG (Antibodies-Online: ABIN101961) 1:100] in Wash buffer supplemented with 2 mM EDTA and 0.1% BSA and incubated on a nutator at room temperature for 1 hour. The beads were decanted on a magnet stand and washed with 200 µL Wash buffer, then were resuspended in 50 µL pA-Tn5 (1:200) in 300-Wash buffer (Wash buffer containing 300 mM NaCl) and incubated on a nutator at room temperature for 1 hour. The beads were washed twice with 200 µL 300-Wash buffer, then resuspended in 50 µL Tagmentation buffer (300-Wash buffer supplemented with 10 mM $MgCl_2$) and incubated at 37 °C for 1 hour. Beads were then washed with 50 µL TAPS wash buffer (10 mM TAPS pH 8.5, 0.2 mM EDTA), then resuspended in 5 µL TAPS wash buffer supplemented with 0.1% SDS and incubated at 58 °C for 1 hour. SDS was neutralized on ice with 15 µL 0.67% Triton-X100. 2 µL of 10 mM indexed P5 and P7 primer solutions and 25 µL NEBnext High-Fidelity 2X Master Mix (New England BioLabs: ME541L) were added. Gap-filling and 15 cycles of PCR were performed using an MJ PTC-200 Thermocycler. Library clean-up was performed by incubating beads with 65 µL SPRI bead slurry for 5-10 minutes, then magnetization and two washes with 200 µL 80% ethanol. Libraries were eluted with 22 µL Tris-HCl pH 8.0 and 2 µL was used for Agilent 4200 Tapestation analysis. The barcoded libraries were mixed to achieve equimolar representation as

desired aiming for a final concentration as recommended by the manufacturer for sequencing

on an Illumina HiSeq 2500 2-lane Turbo flow cell.

CUT&Tag data processing:

CUT&Tag raw sequencing data were aligned to a custom genome made by concatenating

human (hg38), HSV (JN555585.1), and E. coli (U00096.3 Escherichia coli str. K-12 substr.

MG1655, complete genome). We performed alignments using bowtie2.

Domain calling:

Coverage at 100 bp windows of the human hg38 reference genome was calculated as number

of reads of given length (120-1000 bp for the analyses presented here) that mapped at that

window normalized by the factor N:

N= 10,000/(Total number of reads that mapped to E. coli genome)

Here 10,000 is an arbitrarily chosen number. The normalized coverage at each base-pair from

each replicate was averaged when combining multiple replicates. The normalized read density

in 100 bp bins were then smoothed with a running average over the bins spanning +/- 1000 bp

around each bin. We then calculated the genome-wide distribution of normalized read density

and medians that were plotted in Figure S1. We averaged medians across WT macroH2A1

datasets (mock, 4hpi, 8hpi, and 12hpi), and multiplied the average by 2, to set the cutoff for

domain definition for macroH2A1 datasets. We defined a similar cutoff for H3K27me3 datasets

using H3K27me3 WT datasets.

Identifying domain level dynamics of macroH2A1 over time course of infection:

To measure changes in macroH2A1 across time points where the domain boundaries are not

the same, we first concatenated domain definitions from all macroH2A1 datasets and then

defined a set of non-overlapping intervals using the "disjoin" method of GenomicRanges R

package (*57*). We then calculated the log2 ratio of macroH2A1 enrichment over IgG for the non-

overlapping regions for the mock, 4, 8, and 12hpi. The 4, 8, and 12hpi enrichments were then

divided by the enrichment of the mock dataset to obtain change in macroH2A1 over time course

at the non-overlapping regions. k-means clustering (k=6) using R (*58*) was performed on the

matrix where the rows are the non-overlapping regions and the columns are change in

macroH2A1 over mock at 4, 8, and 12hpi. We extended the non-overlapping regions by 5 bp on

each end and then merged regions within each cluster using "reduce" method of

GenomicRanges, to obtain domains in each cluster. We recalculated change in macroH2A1 at

4, 8, and 12hpi over mock, which was used to plot heatmaps and boxplots shown in Figure 1

and Figure S1. H3K27me3 enrichment for WT and macroH2A1 KO cells was calculated at

clustered macroH2A1 domains.

RNA sequencing:

Three biological replicates per time point were obtained from independent infections. Cells were

harvested at the indicated times post-infection by trypsinization, washed with PBS and

centrifuged at 5000 xg for 2 minutes. Pellets were lysed with TRIzol (Thermo Fisher: 15-596-

026) and total RNA was harvested according to manufacturer's instructions. RNAs were then

treated with DNase (Qiagen: 79254) on RNeasy columns (Qiagen: 74104) per manufacturer's

instructions. RNA was precipitated with 3 volumes ice-cold 96% ethanol, 1 volume 3 M sodium

acetate pH 5.5, and 1 µL glycogen (Thermo Fischer: R055) overnight at -80 °C. Precipitated

RNAs were pelleted at 15,000 xg and 4 °C for 30 minutes, washed with ice-cold 75% ethanol

and spun as above for 10 min. RNA was resuspended in nuclease-free water.

RNA was quantified by Nanodrop and integrity analyzed with the 4200 Tapestation Bioanalyzer

system (Agilent). 500 ng of total RNA with an RNA Integrity Number (RIN) greater than 9.5 were

used to prepare sequencing libraries with the TruSeq® Stranded mRNA Library Prep Kit

(Illumina: 20020594). Library concentrations were measured with Qubit dsDNA HS Assay Kit

(Thermo Fisher: Q32854) then analyzed with Agilent High Sensitivity D5000 ScreenTape

System and pooled. Libraries were sequenced with 100-bp paired-end reads on an Illumina

NovaSeq 6000 SP sequencer at the Fred Hutch Genomics Core Facility

RNA/4sU-RNA/Hi-C data processing:

Fastq files were filtered to exclude reads that didn't pass Illumina's base call quality threshold.

STAR v2.7.1 (*59*) with 2-pass mapping was used to align paired-end reads to a combined

reference of human genome build hg38 and HSV1 genome JN555585.1

(https://www.ncbi.nlm.nih.gov/nuccore/JN555585.1/). FastQC 0.11.9

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and RSeQC 4.0.0 (*60*) were used

for QC including insert fragment size, read quality, read duplication rates, gene body coverage

and read distribution in different genomic regions. FeatureCounts (*61*) in Subread 1.6.5 was

used to quantify gene-level expression by strand-specific paired-end read counting.

Gene annotation were based on GENCODE V31 (https://www.gencodegenes.org/human/) and

GCA_000859985.2_ViralProj15217 (https://www.ncbi.nlm.nih.gov/data-

hub/genome/GCF_000859985.2/). For HSV1, annotated genes were collapsed into non-

overlapping transcribed regions, e.g. X indicating a transcribed region unique to gene X, X:Y

indicating an overlapping transcribed region for genes X and Y, and so on.

Bioconductor package edgeR 3.26.8 (*62*) was used to detect differential gene expression

between sample groups. Genes with low expression were excluded by requiring at least one

count per million in at least N samples (N is equal to the number of samples in the smaller

group). The filtered expression matrix was normalized by TMM method

(https://genomebiology.biomedcentral.com/articles/10.1186/gb-2010-11-3-r25) and subject to

significance testing using quasi-likelihood pipeline implemented in edgeR. Genes were deemed

differentially expressed if fold changes were greater than 2 in either direction and Benjamini-

Hochberg adjusted p-values were less than 0.01.

The intervals representing start and end of each gene were intersected with clustered

macroH2A1 domains to obtain cluster assignments for genes. Intersection was performed using

"intersect" function of bedtools (*63*). Genes that did not uniquely intersect with domains in a

single cluster were discarded. The cluster assignments for genes were used for plotting total

RNA fold changes in Figure 2. Reads per kilobase of transcript per million reads mapped

(RPKM) values for 4sU-RNA was obtained from GEO (GSE59717) and converted to transcripts

per million (TPM). The TPM values across two 4sU-RNA replicates for each condition was

averaged and box plots were generated similarly to total RNA.

For calculating eigenvector distributions, compartment wig file was downloaded from the

following link:

https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/b543cbf4-ce54-

4d2d-8960-281528ff18a6/4DNFI342UZP1.bw

Regions from compartment file intersecting with each domain cluster were extracted and the

values were used for generating box plots shown in Figure 2D.

Electron Microscopy:

Cells were fixed in 2% paraformaldehyde and 2.5% glutaraldehyde in 0.1M sodium cacodylate

buffer (pH 7.3) at 4 °C. Fixed cells were rinsed briefly in 1% sucrose in 50mM cacodylate (pH

7.2), then postfixed on ice for 30 mins in a solution of 1% osmium (EM Sciences: RT19152) and

0.8% potassium ferricyanide in 50mM cacodylate (pH 7.2). Cell pellets were washed twice

briefly at 25 °C in 1% sucrose in 50mM cacodylate (pH 7.2), then washed in three changes of

50mM cacodylate (pH 7.2) for 5 min each. Cell pellets were treated with 0.2% tannic acid

(Sigma-Aldrich: 1401-55-4) in 50 mM cacodylate (pH 7.2) for 15 minutes at 25 °C, then rinsed

several times in water. Cells were dehydrated through a graded ethanol series and embedded

in Epon 12 resin (Ted Pella: 18010). 70nm thin sections were cut using an Ultracut UC7

ultramicrotome (Leica Mikrosysteme) and collected on 200 mesh formvar/carbon copper grids

(Ted Pella: 01800). Sections were stained with 2% aqueous uranyl acetate and Reynolds lead

citrate. Cell pellet sections were imaged using a Talos L120C microscope operated at 120kV

with a Ceta-16M (4096 x 4096) camera (ThermoFisher Scientific).

All data were collected at spot size 5 with a 100um C2 aperture and 70um objective aperture.

For quantification, nuclei were targeted at 1250x and manually circled. Autofocus was set to -

2.0um. Nuclei were then imaged at 11000x as a montage and stitched together automatically

using SerialEM (Nexperion Inc). Stitched maps were exported as uncompressed 16-bit tif files

for further analysis. For qualitative analysis, images were manually focused to -2.0um defocus

and then 20 1-second frames were collected and drift-corrected using Velox software (Thermo

Fisher Scientific, Eindhoven, NL). Final summed images were exported as 16-bit tif files and

cropped using ImageJ.

Image analysis pipelines for counting capsids and for measuring chromatin density at the

nuclear envelope were deployed in MATLAB R2020b. Scripts are available on the Fred Hutch

GitHub repository. Capsid pipeline follows three steps: 1- nuclear boundaries identification, 2-

capsid detection, 3- capsid classification. Nuclear boundaries were identified with user input, by

outlining a freehand contour of the nucleus of interest, from which a binary mask is extracted.

Capsid detection was performed on the median-filtered complement of the original image, using

a Circular Hough Transform based algorithm, with phase coding for radii estimation (*64*), and

search radius ranging from 10.6 to 31.7 nm. Capsids residing within the nuclear mask were then

counted and classified. Detected capsids were classified in three categories (Empty,

Intermediate and Full), depending on the distribution of the pixel grayscale intensities within the

capsids relative to a normal distribution.

Width of heterochromatin abutting the nuclear envelope was quantified by measuring the length

of the binarized chromatin from 1D intensity profiles along the normal of the nuclear perimeter,

sampled at every 10 perimeter pixels. Dense heterochromatin was binarized using global Otsu's

thresholding method applied to the background-corrected complement of the contrast-adjusted

original image. Noise from the binarized image was further reduced by applying a 2D order

statistic filter using the minimum value of a varying domain interactively defined by the user, with

default value of 8-by-8 pixels. The resulting heterochromatin density distribution was normalized

to the total length of the nucleus' perimeter.

REFERENCES CITED

1. D. M. Knipe, A. Cliffe, Chromatin control of herpes simplex virus lytic and latent infection. *Nat Rev Microbiol*. 6, 211–221 (2008).

2. K. Monier, J. C. G. Armas, S. Etteldorf, P. Ghazal, K. F. Sullivan, Annexation of the interchromosomal space during viral infection. *Nat Cell Biol*. 2, 661–665 (2000).

3. K. Kulej, D. C. Avgousti, S. Sidoli, C. Herrmann, A. N. D. Fera, E. T. Kim, B. A. Garcia, M. D. Weitzman, Time-resolved Global and Chromatin Proteomics during Herpes Simplex Virus Type 1 (HSV-1) Infection. *Mol Cell Proteomics*. 16, S92–S107 (2017).

4. V. Aho, E. Mäntylä, A. Ekman, S. Hakanen, S. Mattola, J.-H. Chen, V. Weinhardt, V. Ruokolainen, B. Sodeik, C. Larabell, M. Vihinen-Ranta, Quantitative Microscopy Reveals Stepwise Alteration of Chromatin Structure during Herpesvirus Infection. *Viruses*. 11, 935 (2019).

5. V. Aho, S. Salminen, S. Mattola, A. Gupta, F. Flomm, B. Sodeik, J. B. Bosse, M. Vihinen-Ranta, Infection-induced chromatin modifications facilitate translocation of herpes simplex virus capsids to the inner nuclear membrane. *Plos Pathog*. 17, e1010132 (2021).

6. M. Myllys, V. Ruokolainen, V. Aho, E. A. Smith, S. Hakanen, P. Peri, A. Salvetti, J. Timonen, V. Hukkanen, C. A. Larabell, M. Vihinen-Ranta, Herpes simplex virus 1 induces egress channels through marginalized host chromatin. *Sci Rep-uk*. 6, 28844 (2016).

7. P. Chi, C. D. Allis, G. G. Wang, Covalent histone modifications — miswritten, misinterpreted and mis-erased in human cancers. *Nat Rev Cancer*. 10, 457–469 (2010).

8. R. Margueron, D. Reinberg, The Polycomb complex PRC2 and its mark in life. *Nature*. 469, 343–349 (2011).

9. V. Pasque, A. Gillich, N. Garrett, J. B. Gurdon, Histone variant macroH2A confers resistance to nuclear reprogramming. *Embo J*. 30, 2373–2387 (2011).

10. F. G. Ghiraldini, D. Filipescu, E. Bernstein, Solid tumours hijack the histone variant network. *Nat Rev Cancer*. 21, 257–275 (2021).

11. M. J. Gamble, W. L. Kraus, Multiple facets of the unique histone variant macroH2A: From genomics to cell biology. *Cell Cycle*. 9, 2568–2574 (2010).

12. G. Timinszky, S. Till, P. O. Hassa, M. Hothorn, G. Kustatscher, B. Nijmeijer, J. Colombelli, M. Altmeyer, E. H. K. Stelzer, K. Scheffzek, M. O. Hottiger, A. G. Ladurner, A macrodomain-containing histone rearranges chromatin upon sensing PARP1 activation. *Nat Struct Mol Biol*. 16, 923–929 (2009).

13. H. Chen, P. D. Ruiz, L. Novikov, A. D. Casill, J. W. Park, M. J. Gamble, MacroH2A1.1 and PARP-1 cooperate to regulate transcription by promoting CBP-mediated H2B acetylation. *Nat Struct Mol Biol*. 21, 981–989 (2014).

14. A. Kapoor, M. S. Goldberg, L. K. Cumberland, K. Ratnakumar, M. F. Segura, P. O. Emanuel, S.
Menendez, C. Vardabasso, G. LeRoy, C. I. Vidal, D. Polsky, I. Osman, B. A. Garcia, E. Hernando, E.
Bernstein, The histone variant macroH2A suppresses melanoma progression through regulation of CDK8.
Nature. 468, 1105–1109 (2010).

15. J. Douet, D. Corujo, R. Malinverni, J. Renauld, V. Sansoni, M. P. Marjanović, N. Cantariño, V.
Valero, F. Mongelard, P. Bouvet, A. Imhof, M. Thiry, M. Buschbeck, MacroH2A histone variants
maintain nuclear organization and heterochromatin architecture. *J Cell Sci*. 130, 1570–1582 (2017).

16. Y. Fu, P. Lv, G. Yan, H. Fan, L. Cheng, F. Zhang, Y. Dang, H. Wu, B. Wen, MacroH2A1 associates
with nuclear lamina and maintains chromatin architecture in mouse liver cells. *Sci Rep-uk*. 5, 17186
(2015).

17. C. James, M. Harfouche, N. J. Welton, K. M. Turner, L. J. Abu-Raddad, S. L. Gottlieb, K. J. Looker,
Herpes simplex virus: global infection prevalence and incidence estimates, 2016. *B World Health Organ*.
98, 315–329 (2020).

18. K. J. Looker, A. S. Magaret, M. T. May, K. M. E. Turner, P. Vickerman, L. M. Newman, S. L.
Gottlieb, First estimates of the global and regional incidence of neonatal herpes infection. *Lancet Global
Heal*. 5, e300–e309 (2017).

19. L. A. Sadowski, R. Upadhyay, Z. W. Greeley, B. J. Margulies, Current Drugs to Treat Infections with
Herpes Simplex Viruses-1 and -2. *Viruses*. 13, 1228 (2021).

20. J. Matthias, S. du Bernard, J. A. Schillinger, J. Hong, V. Pearson, T. A. Peterman, Estimating neonatal
herpes simplex virus incidence and mortality using capture-recapture, Florida. *Clin Infect Dis*. 73,
ciaa727- (2020).

21. A. Berkhout, V. Kapoor, C. Heney, C. A. Jones, J. E. Clark, V. L. Vaska, M. Lai, C. Nourse, Herpes
Simplex Virus Infection in Infants: 13 Year Evaluation (2005–2017) of Laboratory Confirmed Cases in
Queensland, Australia. *Pediatr Infect Dis J*. Publish Ahead of Print (2020),
doi:10.1097/inf.0000000000002970.

22. P. G. Spear, Herpes simplex virus: receptors and ligands for cell entry. *Cell Microbiol*. 6, 401–410
(2004).

23. V. Tiwari, M. Oh, M. Kovacs, S. Y. Shukla, T. Valyi-Nagy, D. Shukla, Role for nectin-1 in herpes
simplex virus 1 entry and spread in human retinal pigment epithelial cells. *Febs J*. 275, 5272–5285
(2008).

24. R. B. Deal, J. G. Henikoff, S. Henikoff, Genome-Wide Kinetics of Nucleosome Turnover Determined
by Metabolic Labeling of Histones. *Science*. 328, 1161–1164 (2010).

25. E. M. Riising, I. Comet, B. Leblanc, X. Wu, J. V. Johansen, K. Helin, Gene Silencing Triggers
Polycomb Repressive Complex 2 Recruitment to CpG Islands Genome Wide. *Mol Cell*. 55, 347–360
(2014).

26. E. Lieberman-Aiden, N. L. van Berkum, L. Williams, M. Imakaev, T. Ragoczy, A. Telling, I. Amit,
B. R. Lajoie, P. J. Sabo, M. O. Dorschner, R. Sandstrom, B. Bernstein, M. A. Bender, M. Groudine, A.

Gnirke, J. Stamatoyannopoulos, L. A. Mirny, E. S. Lander, J. Dekker, Comprehensive Mapping of Long-
Range Interactions Reveals Folding Principles of the Human Genome. *Science*. 326, 289–293 (2009).

27. S. S. P. Rao, M. H. Huntley, N. C. Durand, E. K. Stamenova, I. D. Bochkov, J. T. Robinson, A. L.
Sanborn, I. Machol, A. D. Omer, E. S. Lander, E. L. Aiden, A 3D Map of the Human Genome at
Kilobase Resolution Reveals Principles of Chromatin Looping. *Cell*. 159, 1665–1680 (2014).

28. W. Cai, P. A. Schaffer, Herpes simplex virus type 1 ICP0 regulates expression of immediate-early,
early, and late genes in productively infected cells. *J Virol*. 66, 2904–2915 (1992).

29. R. Naldinho-Souto, H. Browne, T. Minson, Herpes Simplex Virus Tegument Protein VP16 Is a
Component of Primary Enveloped Virions. *J Virol*. 80, 2582–2584 (2006).

30. E. U. Lorentzen, B. R. Eing, W. Hafezi, R. Manservigi, J. E. Kühn, Replication-Competent Herpes
simplex Virus Type 1 Mutant Expressing an Autofluorescent Glycoprotein H Fusion Protein.
Intervirology. 44, 232–242 (2001).

31. R. Rickels, D. Hu, C. K. Collings, A. R. Woodfin, A. Piunti, M. Mohan, H.-M. Herz, E. Kvon, A.
Shilatifard, An Evolutionary Conserved Epigenetic Mark of Polycomb Response Elements Implemented
by Trx/MLL/COMPASS. *Mol Cell*. 63, 318–328 (2016).

32. J. K. Lue, J. E. Amengual, Emerging EZH2 Inhibitors and Their Application in Lymphoma. *Curr
Hematol Malig R*. 13, 369–382 (2018).

33. J. H. Arbuckle, P. J. Gardina, D. N. Gordon, H. D. Hickman, J. W. Yewdell, T. C. Pierson, T. G.
Myers, T. M. Kristie, Inhibitors of the Histone Methyltransferases EZH2/1 Induce a Potent Antiviral State
and Suppress Infection by Diverse Viral Pathogens. *Mbio*. 8, e01141-17 (2017).

34. V. Aho, M. Myllys, V. Ruokolainen, S. Hakanen, E. Mäntylä, J. Virtanen, V. Hukkanen, T. Kühn, J.
Timonen, K. Mattila, C. A. Larabell, M. Vihinen-Ranta, Chromatin organization regulates viral egress
dynamics. *Sci Rep-uk*. 7, 3692 (2017).

35. W. Gibson, B. Roizman, Proteins Specified by Herpes Simplex Virus VIII. Characterization and
Composition of Multiple Capsid Forms of Subtypes 1 and 2. *J Virol*. 10, 1044–1052 (1972).

36. X.-K. Yu, C. M. O'Connor, I. Atanasov, B. Damania, D. H. Kedes, Z. H. Zhou, Three-Dimensional
Structures of the A, B, and CCapsids of Rhesus Monkey Rhadinovirus: Insights into
GammaherpesvirusCapsid Assembly, Maturation, and DNAPackaging. *J Virol*. 77, 13182–13193 (2003).

37. T. Hennig, M. Michalski, A. J. Rutkowski, L. Djakovic, A. W. Whisnant, M.-S. Friedl, B. A. Jha, M.
A. P. Baptista, A. L'Hernault, F. Erhard, L. Dölken, C. C. Friedel, HSV-1-induced disruption of
transcription termination resembles a cellular stress response but selectively increases chromatin
accessibility downstream of genes. *Plos Pathog*. 14, e1006954 (2018).

38. Y. Cai, Y. Zhang, Y. P. Loh, J. Q. Tng, M. C. Lim, Z. Cao, A. Raju, E. L. Aiden, S. Li, L.
Manikandan, V. Tergaonkar, G. Tucker-Kellogg, M. J. Fullwood, H3K27me3-rich genomic regions can
function as silencers to repress gene expression via chromatin interactions. *Nat Commun*. 12, 719 (2021).

39. C. Gao, L. Chen, S.-B. Tang, Q.-Y. Long, J.-L. He, N.-A. Zhang, H.-B. Shu, Z.-X. Chen, M. Wu, L.-
Y. Li, The epigenetic landscapes of histone modifications on HSV-1 genome in human THP-1 cells.
Antivir Res. 176, 104730 (2020).

40. J. Kim, C. Sun, A. D. Tran, P.-J. Chin, P. D. Ruiz, K. Wang, R. J. Gibbons, M. J. Gamble, Y. Liu, P.
Oberdoerffer, The macroH2A1.2 histone variant links ATRX loss to alternative telomere lengthening. *Nat*
Struct Mol Biol. 26, 213–219 (2019).

41. J. Arii, Host and Viral Factors Involved in Nuclear Egress of Herpes Simplex Virus 1. *Viruses*. 13,
754 (2021).

42. S. L. Bjerke, R. J. Roller, Roles for herpes simplex virus type 1 UL34 and US3 proteins in disrupting
the nuclear lamina during herpes simplex virus type 1 egress. *Virology*. 347, 261–276 (2006).

43. M. Simpson-Holley, J. Baines, R. Roller, D. M. Knipe, Herpes Simplex Virus 1 UL31 and UL34
Gene Products Promote the Late Maturation of Viral Replication Compartments to the Nuclear Periphery.
J Virol. 78, 5591–5600 (2004).

44. N. R. Leach, R. J. Roller, Significance of host cell kinases in herpes simplex virus type 1 egress and
lamin-associated protein disassembly from the nuclear lamina. *Virology*. 406, 127–137 (2010).

45. M. M. Bahnamiri, R. J. Roller, Mechanism of Nuclear Lamina Disruption and the Role of pUS3 in
Herpes Simplex Virus 1 Nuclear Egress. *J Virol*. 95 (2021), doi:10.1128/jvi.02432-20.

46. J. M. Bigalke, T. Heuser, D. Nicastro, E. E. Heldwein, Membrane deformation and scission by the
HSV-1 nuclear egress complex. *Nat Commun*. 5, 4131–4131 (2014).

47. E. B. Draganova, M. K. Thorsen, E. E. Heldwein, Nuclear Egress of Herpesviruses. *Curr Issues Mol*
Biol. 41, 125–170 (2021).

48. K. L. Lynch, M. R. Dillon, M. Bat-Erdene, H. C. Lewis, R. J. Kaai, E. A. Arnold, D. C. Avgousti, A
viral histone-like protein exploits antagonism between linker histones and HMGB proteins to obstruct the
cell cycle. *Curr Biol* (2021), doi:10.1016/j.cub.2021.09.050.

49. S. M. Brown, D. A. Ritchie, J. H. Subak-Sharpe, Genetic Studies with Herpes Simplex Virus Type 1.
The Isolation of Temperature-sensitive Mutants, their Arrangement into Complementation Groups and
Recombination Analysis Leading to a Linkage Map. *J Gen Virol*. 18, 329–346 (1973).

50. C. E. Lilley, C. T. Carson, A. R. Muotri, F. H. Gage, M. D. Weitzman, DNA repair proteins affect the
lifecycle of herpes simplex virus 1. *P Natl Acad Sci Usa*. 102, 5844–5849 (2005).

51. E. M. Hatch, A. Kulukian, A. J. Holland, D. W. Cleveland, T. Stearns, Cep152 interacts with Plk4 and
is required for centriole duplication. *J Cell Biol*. 191, 721–729 (2010).

52. G. Li, C. C. Nguyen, B. J. Ryckman, W. J. Britt, J. P. Kamil, A viral regulator of glycoprotein
complexes contributes to human cytomegalovirus cell tropism. *Proc National Acad Sci*. 112, 4471–4476
(2015).

53. S. J. Child, A. L. Greninger, A. P. Geballe, Rapid adaptation to human protein kinase R by a unique
genomic rearrangement in rhesus cytomegalovirus. *Plos Pathog*. 17, e1009088 (2021).

54. M. Aubert, N. M. Boyle, D. Stone, L. Stensland, M.-L. Huang, A. S. Magaret, R. Galetto, D. J.
Rawlings, A. M. Scharenberg, K. R. Jerome, In vitro Inactivation of Latent HSV by Targeted
Mutagenesis Using an HSV-specific Homing Endonuclease. *Mol Ther Nucleic Acids*. 3, e146 (2014).

55. H. S. Kaya-Okur, S. J. Wu, C. A. Codomo, E. S. Pledger, T. D. Bryson, J. G. Henikoff, K. Ahmad, S.
Henikoff, CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat Commun*.
10, 1930 (2019).

56. M. Lawrence, W. Huber, H. Pagès, P. Aboyoun, M. Carlson, R. Gentleman, M. T. Morgan, V. J.
Carey, Software for Computing and Annotating Genomic Ranges. *Plos Comput Biol*. 9, e1003118 (2013).

57. R. C. Team, R: A Language and Environment for Statistical Computing. *R Foundation for Statistical*
Computing (2019) (available at https://www.R-project.org).

58. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R.
Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 29, 15–21 (2013).

59. L. Wang, S. Wang, W. Li, RSeQC: quality control of RNA-seq experiments. *Bioinformatics*. 28,
2184–2185 (2012).

60. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics*. 30, 923–930 (2014).

61. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics*. 26, 139–140 (2010).

62. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 26, 841–842 (2010).

63. T. J. Atherton, D. J. Kerbyson, Size invariant circle detection. *Image Vision Comput*. 17, 795–803
(1999).

1009