The conserved AAA-ATPase PCH-2 distributes its regulation of meiotic prophase events

by remodeling multiple meiotic HORMADs in *C. elegans*

Anna E. Russo*, Stefani Giacopazzi*, Alison Deshong*, Malaika Menon*, Valery Ortiz*, Kaori M.

Ego#, Kevin D. Corbett# and Needhi Bhalla*[1]

* Department of Molecular, Cell and Developmental Biology, University of California, Santa

Cruz, Santa Cruz, CA 95064

Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla,

CA 92093

[1] corresponding author: nbhalla@ucsc.edu

Department of Molecular, Cell and Developmental Biology

225 Sinsheimer Labs

University of California, Santa Cruz

Santa Cruz, CA 95064

phone: (831) 459-1319

fax: (831) 459-3139

Abstract

During meiotic prophase, the essential events of homolog pairing, synapsis, and recombination are coordinated with meiotic progression to promote fidelity and prevent aneuploidy. The conserved AAA+ ATPase PCH-2 coordinates these events to guarantee crossover assurance and accurate chromosome segregation. How PCH-2 accomplishes this coordination is poorly understood. Here, we provide evidence that PCH-2 decelerates pairing, synapsis and recombination in *C. elegans* by remodeling meiotic HORMADs. We propose that PCH-2 converts the closed versions of these proteins, which drive these meiotic prophase events, to unbuckled conformations, destabilizing interhomolog interactions and delaying meiotic progression. Further, we find that PCH-2 distributes this regulation among three essential meiotic HORMADs in *C. elegans*: PCH-2 acts through HTP-3 to regulate pairing and synapsis, HIM-3 to promote crossover assurance, and HTP-1 to control meiotic progression. In addition to identifying a molecular mechanism for how PCH-2 regulates interhomolog interactions, our results provide a possible explanation for the expansion of the meiotic HORMAD family as a conserved evolutionary feature of meiosis. Taken together, our work demonstrates that PCH-2's remodeling of meiotic HORMADs has functional consequences for the rate and fidelity of homolog pairing, synapsis, recombination and meiotic progression, ensuring accurate meiotic chromosome segregation.

<u>Introduction</u>

All sexually reproducing organisms depend on the specialized cell division called meiosis, which

produces haploid gametes such as sperm, eggs, and pollen. During meiosis, cells combine one

round of DNA replication with two rounds of chromosome segregation, all of which is tightly

regulated to ensure that gametes inherit the correct number of chromosomes. Errors in meiosis

can lead to aneuploid gametes that cause infertility, birth defects and miscarriages [1].

Therefore, understanding how meiotic events are coordinated has important implications for

reproductive health.

During prophase I of meiosis, chromosome pairs or homologs undergo a series of events known

as pairing, synapsis, and crossover recombination. First, homologs find each other and pair

within the nucleus. Next, a molecular structure called the synaptonemal complex (SC)

assembles between homologs and stabilizes homolog pairing in a process called synapsis.

Finally, homologs undergo crossover recombination, which serves two important purposes.

First, it shuffles alleles to create new genetic combinations that can be acted upon by natural

selection. Second, it physically links homolog pairs, which is required for homologs to properly

remodel, orient on the spindle, and segregate correctly during the first meiotic division.

Despite being described as a linear pathway, the processes of pairing, synapsis and

recombination between homologous chromosomes are highly dynamic and can occur

simultaneously, requiring active coordination. Further, these events can also initiate

inappropriately between non-homologous chromosomes and therefore need to be corrected.

We have shown that the highly conserved AAA+ ATPase, Pachytene checkpoint 2 (PCH-2),

contributes to both the coordination and correction of meiotic events [2, 3]. However, how it

carries out these roles is currently unclear.

PCH-2, and its mammalian ortholog TRIP13, has been shown to act through a family of proteins

that contain a HORMA domain (HORMAD proteins) to mediate a variety of chromosome

behaviors, including meiotic interhomolog interactions. HORMADs were named after the first

proteins in this family to be discovered in budding yeast: Hop1, Rev7, and Mad2 [4] and meiotic

HORMADs are essential to organize meiotic chromosomes for the meiosis-specific events of

homolog pairing, synapsis, and recombination [5]. Despite having essential roles in meiotic

prophase, meiotic HORMADs differ greatly across eukaryotes in the number of family members

they contain. For example, *S. cerevisiae* and *S. pombe* each have one meiotic HORMAD [6, 7],

mice and *Arabidopsis* have two [8-10] and *C. elegans* has four [11-14]. In both mice and plants,

one meiotic HORMAD plays a more integral role in the events of pairing, synapsis, and

recombination than the other, indicating some specialization of function [9, 10, 15-18]. This

specialization is also observed in *C. elegans* among the three essential HORMADs [19]. HTP-3

is responsible for DNA double-strand break formation [12] and REC-8-mediated sister chromatid

cohesion [20]. HIM-3 promotes synapsis [19] and homolog access to ensure repair off the

homolog versus the sister chromatid during crossover recombination [21]. HTP-1, in addition to

contributing to homolog access along with HIM-3, also controls the timely initiation of SC

assembly once homologs are paired properly [11, 13] and the protection of sister chromatid

cohesion during the first meiotic division [22]. The fourth meiotic HORMAD, HTP-2, is not

essential and is largely redundant with HTP-1 [13, 22].

Meiotic HORMAD proteins have a characteristic N-terminal HORMA domain that allows

HORMADs to adopt two different conformations: a closed conformation and an open/unlocked

conformation (reviewed in [23]). In the closed conformation, a peptide on a binding partner

called the "closure motif" binds the core of the HORMAD's HORMA domain. This interaction is

stabilized by a C-terminal region of the HORMA domain called the safety belt, which wraps

around the HORMAD to lock the binding partner's closure motif against the HORMA domain

core. Conversion of the HORMAD from the closed to open conformation occurs when the PCH-

2/TRIP13 binds and partially unfolds the extreme N-terminus of the HORMA domain, resulting in

a structural rearrangement that releases the bound closure motif and causes the safety belt to

occupy the closure motif interaction site. Biochemical and structural studies show that PCH-

2/TRIP13 is directly responsible for the conversion of closed conformers to open conformers for

the mitotic HORMADs Mad2 [24] and Rev7 [25], where this remodeling relies on a direct

interaction between PCH-2/TRIP13 and the N-terminus of the HORMAD [26, 27].

Conformational switching has also been shown to be essential for meiotic HORMAD function.

Closed versions of meiotic HORMADs assemble on meiotic chromosomes to drive pairing,

synapsis, recombination, and meiotic progression [19]. While stable, open conformers of meiotic

HORMADs have not been observed *in vitro* or *in vivo*, the budding yeast meiotic HORMAD,

Hop1, exists in solution as two distinct conformations that are in equilibrium: the closed

conformation and an intermediate "extended" state that has been referred to as unbuckled [28].

The physiological relevance of unbuckled meiotic HORMADs *in vivo* is unclear but work in

plants and budding yeast indicates that their unbuckling is required for nuclear import and

association with meiotic chromosomes [29, 30].

While studies with other HORMADs, such as Mad2 and Rev7, provide a powerful framework to

conceptualize PCH-2's molecular mechanism during meiotic prophase, identifying PCH-2's

specific role during meiosis has been challenging. In contrast to Mad2 and Rev7, attempts to

undertake similar structural and biochemical analysis of PCH-2/TRIP13 and meiotic HORMADs

have been less informative. However, prior evidence from budding yeast, plants, and mammals

indicates that PCH-2/TRIP13 binds and acts directly on meiotic HORMADs. In budding yeast,

Pch2 has been shown to bind to and remodel meiotic HORMADs *in vitro* [31]. Additionally,

PCH-2's orthologs in other systems have been shown to promote meiotic progression [32] by

depleting or redistributing meiotic HORMADs from chromosome axes upon synapsis, including

budding yeast [33, 34], *Arabidopsis* [35] and mice [10]. In mice, the removal of the meiotic

HORMAD, HORMAD1, at the onset of synapsis is abolished in both TRIP13 mutants [10] or

when the N-terminus of HORMAD1 is deleted [27], indicating that PCH-2 and its orthologs'

remodeling of meiotic HORMADs is similar to its remodeling of Mad2 and Rev7. However, in

some systems, such as *C. elegans*, meiotic HORMADs do not rely on PCH-2 for nuclear import

[2] nor are they removed from chromosomes upon synapsis [21, 22, 36]. Further, PCH-2 and its

orthologs localize to meiotic chromosomes as foci prior to synapsis in budding yeast, plants and

C. elegans [2, 34, 35, 37, 38] and meiotic HORMADs are not fully removed during synapsis in

budding yeast, plants or mice [10, 33, 35]. These data raise the important question of whether

remodeling of meiotic HORMADs plays additional roles during meiotic prophase.

Here, we use a combination of genetic, cytological and biochemical data to demonstrate that

PCH-2 remodels meiotic HORMADs to regulate and coordinate pairing, synapsis, crossover

recombination, and meiotic progression in *C. elegans*. We show that a missense mutation in the

HORMA domain of HTP-3 suppresses pairing and synapsis defects in *pch-2* mutants but

exacerbates its recombination defects, indicating these two proteins cooperate to promote

pairing and synapsis but regulate recombination independently. We also show that a similar

mutation in *him-3* exacerbates synapsis defects in *pch-2* mutants but suppresses its

recombination defects, suggesting that PCH-2 acts on HIM-3 specifically to promote crossover

recombination. Finally, we show that corresponding mutations in the meiotic HORMAD HTP-1

produce a delay in meiotic progression, with consequences for synapsis and recombination, and

can be partially suppressed by a mutation in *pch-2*, suggesting that PCH-2 cooperates with

HTP-1 to regulate meiotic progression. Biochemical analysis demonstrates that these mutations

limit the ability of meiotic HORMADs to adopt their closed conformations. Taken together, our

work strongly supports a model in which PCH-2's remodeling of meiotic HORMADs from closed

to unbuckled conformations decelerates pairing, synapsis, recombination, and meiotic

progression, to proofread homolog interactions and maintain meiotic fidelity. Further, PCH-2

cleanly delegates its regulation of different meiotic prophase events to different meiotic

HORMADs, offering new insight into how PCH-2 mechanistically coordinates the events of

pairing, synapsis, and recombination. Lastly, it provides a potential explanation for why this

family of meiotic HORMADs has dramatically expanded in *C. elegans* and provides a possible

explanation about why some model systems have multiple meiotic HORMADs.

Results

PCH-2's localization is consistent with remodeling HORMADs on the axial elements of meiotic

chromosomes

Meiotic HORMADs collaborate with cohesins and other axial proteins to organize meiotic

chromosomes into linear axial elements functionally capable of undergoing interhomolog

pairing, synapsis and recombination [5]. While PCH-2/TRIP13 has been implicated in

remodeling meiotic HORMADs away from meiotic chromosomes to ensure their availability [29,

39] and entry into the nucleus [30], its localization to meiotic chromosomes before and after

synapsis [2, 34, 35, 37, 38] suggests a role remodeling meiotic HORMADs on chromosomes to

regulate pairing, synapsis and recombination as well as meiotic progression. To test this, we

used structured illumination microscopy to more closely investigate whether PCH-2's

localization was consistent with remodeling meiotic HORMADs on axial elements.

Before synapsis and during axis assembly, when meiotic HORMADs are diffuse throughout

nuclei, PCH-2 localizes to foci on meiotic chromosomes [2]. After synapsis, PCH-2 localizes

along the synaptonemal complex (SC) and this localization depends on the central element

component SYP-1 [2, 40], which assembles between axial elements during synapsis (Figure

1A). We performed indirect immunofluorescence on wildtype meiotic nuclei against both PCH-2

and SYP-1 and performed structured illumination microscopy to resolve the distance between

axial elements (Figure 1B) [41]. While SYP-1 appeared as a line of continuous width in all

meiotic nuclei, PCH-2 staining often appeared wider than SYP-1, sometimes splitting as parallel

tracks with SYP-1 staining between PCH-2 tracks (inset and arrows in Figure 1B, Figure 1C).

Indeed, when we quantified the signal intensities for both PCH-2 and SYP-1, we found that the

signal for PCH-2 spanned a wider distance compared to SYP-1 (Figure 1D), indicating that

PCH-2 can reside on or close to axial elements of synapsed chromosomes and remodel meiotic

HORMADs on chromosomes.

htp-3 ^{H96Y} suppresses the acceleration of pairing and synapsis in pch-2 mutants

Having established that PCH-2's localization is consistent with the ability to remodel meiotic

HORMADs on meiotic chromosomes, we looked for genetic interactions that might support this

possibility. *htp-3(vc75)* is a missense mutation that replaces histidine 96 with a tyrosine in the

HORMA domain (H96Y) (Figure 2A). HTP-3^{H96Y} properly localizes to meiotic chromosomes [42]

and does not affect pairing, synapsis and crossover recombination [42, 43]. However, we and

others previously showed that this mutation in *htp-3* abolished meiotic checkpoint responses in

C. elegans [42, 43]. Because of this phenotype, we analyzed HTP-3's relationship with PCH-2 in

regulating pairing, synapsis, recombination, and meiotic progression.

To test whether PCH-2 genetically interacts with HTP-3 to regulate pairing, we generated *pch-*

2;syp-1 double mutants, *htp-3^{H96Y};syp-1* double mutants*, and *pch-2;htp-3^{H96Y};syp-1* triple

mutants and assessed pairing when compared with *syp-1* single mutants. *syp-1* null mutants

successfully undergo pairing despite the failure to load SC components, particularly at specific

loci called Pairing Centers (PCs) [40]. Therefore, these experiments allow us to analyze pairing

independent of synapsis. We performed immunofluorescence against the HIM-8 protein, which

localizes to PCs on the X chromosome [44], to assess pairing at this locus throughout the

germline: two HIM-8 foci indicate the X-chromosomes are unpaired while a single HIM-8 focus

indicates paired X-chromosomes. Because meiotic nuclei are arranged in spatiotemporal

gradient in germlines, we divided the germline into six equivalently sized zones to perform a

meiotic time-course and determined the percentage of nuclei with a single HIM-8 focus in each

zone (Figure 3A and 3B).

Similar to previous work [2, 3], we found that there is an increase in percentage of nuclei with

paired X chromosomes in zone 2 for *pch-2;syp-1* double mutants (76%, Figure 3C) compared to

syp-1 single mutant controls (49%, Figure 3C), showing that the rate of pairing is accelerated in

the absence of *pch-2* (p value <0.0001 by two-tailed Fisher's exact test). *htp-3^{H96Y};syp-1* double

mutants have a similar rate of pairing compared to *syp-1* single mutant controls (46%, Figure

3C, p value = 0.4001 by two-tailed Fisher's exact test), demonstrating that this mutation does

not affect the rate of pairing. Interestingly, we found that *pch-2;htp-3^{H96Y};syp-1* triple mutants

have a similar rate of pairing compared to *syp-1* single mutants and *htp-3^{H96Y};syp-1* double

mutants (49%, Figure 3C), indicating that *htp-3^{H96Y}* suppresses the acceleration of pairing

observed in *pch-2* mutants.

Next, we tested for a genetic interaction between *pch-2* and *htp-3* during synapsis. We analyzed

the rate of synapsis in germlines from control, *pch-2* single mutants, *htp-3^{H96Y}* single mutants,

and *pch-2;htp-3^{H96Y}* double mutant worms by performing immunofluorescence against the axial

element HTP-3 [12, 36] and the central element SYP-1 [40]. We then determined the fraction of

nuclei with complete synapsis in each zone. Complete synapsis is visible as full co-localization

of HTP-3 and SYP-1, whereas nuclei with incomplete synapsis have stretches of HTP-3 without

SYP-1 signal (arrowheads, Figure 3D). Similar to previous data, we report that *pch-2* null

mutants have an accelerated rate of synapsis (61%, Figure 3E) compared to wildtype controls

(32%, Figure 3E, p value <0.0001 by two-tailed Fisher's exact test) in zone 2 and a delay in

desynapsis in zone 6 [2, 3] (Figure 3E). htp-3^{H96Y} single mutants have a similar rate of synapsis

compared to wildtype controls (36%, Figure 3E, p value = 0.228 by two-tailed Fisher's exact

test), suggesting that the rate of synapsis is unaffected in this background, and accelerated

desynapsis, as previously reported [42]. Analogous to our pairing results, we found that *pch-*

2;htp-3^{H96Y} double mutants have a similar rate of synapsis compared to wildtype controls and

htp-3^{H96Y} single mutants (34%, Figure 3E), indicating that *htp-3^{H96Y}* suppresses the acceleration

of synapsis in *pch-2* mutants.

To further solidify that PCH-2 acts on HTP-3 to regulate synapsis, we analyzed synapsis in

meDf2 heterozygotes (*meDf2/+*). *meDf2* is a mutation in which PC regions are deleted from the

X chromosome [45]. Since PCs are essential for pairing and synapsis [36], *meDf2* homozygotes

fail to pair and synapse X chromosomes while *meDf2* heterozygotes behave like a partial loss of

function mutation: ~50% of nuclei complete synapsis of X chromosomes while the remaining

50% have unsynapsed X chromosomes [36]. In *pch-2;meDf2/+* worms, ~85% of nuclei complete

synapsis, demonstrating that PCH-2 normally inhibits synapsis when PC function is

compromised [2].

To test whether *htp-3^{H96Y}* also suppressed the synapsis in *pch-2;meDf2/+* worms, we generated

pch-2;htp-3^{H96Y};meDf2/+ triple mutants and analyzed the rate of synapsis, comparing it to

controls. Similar to previous results, we found that 87% of nuclei in *pch-2;meDf2/+* double

mutants achieve complete synapsis in zone 5 compared to 48% complete synapsis for *meDf2/+*

single mutant worms (Figure 3F). *htp-3^{H96Y};meDf2/+* worms have 52% of nuclei with complete

synapsis in zone 5, similar to *meDf2/+* single mutants (Figure 3F). *pch-2;htp-3^{H96Y};meDf2/+*

triple mutants show maximal levels of synapsis in zone 4 (36%, Figure 3F) and more closely

resemble the levels of synapsis observed in to *htp-3^{H96Y};meDf2/+* double and *meDf2/+* single

mutants than *pch-2;meDf2/+* double mutants. These data indicate that PCH-2's ability to inhibit

synapsis when PC function is compromised depends on the function of HTP-3, and perhaps

more specifically, on the HORMA domain of HTP-3, similar to what we observed when PCs are

fully functional (Figure 3E). To determine whether these defects in synapsis produce

chromosome segregation errors, we scored the percent of male progeny in these strains. Male

progeny, which are genotypically XO in contrast to XX hermaphrodites, are diagnostic of

misegregation of X chromosomes in meiosis. The ability of *htp-3^{H96Y}* to suppress the synapsis of

X chromosomes in *pch-2;meDf2/+* was further supported by the increase in male progeny

observed in *pch-2;htp-3^{H96Y};meDf2/+* triple mutants (10.84%, Table 1) in comparison to *pch-*

2;meDf2/+ double mutants (0.83%, Table 1).

htp-3^{H96Y} exacerbates defects in DNA repair and crossover formation in *pch-2* mutants

We next tested whether PCH-2 regulates recombination through HTP-3 by assaying DNA repair

and crossover formation in *pch-2;htp-3^{H96Y}* double mutants. The rate of DNA repair is assessed

by quantifying the number of RAD-51 foci per nucleus throughout the germline (Figure 3G). The

appearance of RAD-51 foci on meiotic chromosomes is associated with the introduction of DNA

double-strand breaks and their disappearance is associated with the entry of DNA double-

strand breaks into repair pathways [46]. RAD-51 numbers peak in zone 4 in wildtype animals,

increasing to an average of 2 per nuclei, and decrease as RAD-51 is removed during DNA

repair. *pch-2* single mutants have a lower average of RAD-51 foci per nucleus in zones 4-6

(Figure 3H), consistent with previous findings that that *pch-2* mutants exhibit a faster rate of

DNA repair compared to wildtype controls [2]. *htp-3^{H96Y}* single mutants show a similar rate of

DNA repair as wildtype germlines (Figure 3H), although the average number of RAD-51 foci is

slightly lower than wildtype [42]. By contrast, *pch-2;htp-3^{H96Y}* double mutants not only show an

accelerated rate of DNA repair, similar to *pch-2* single mutants, they have a reduced average

number of RAD-51 foci in zone 5 (0.75 foci, Figure 3H), showing that mutation of *htp-3*

exacerbates the acceleration of DNA repair in *pch-2* mutants. Because *htp-3^{H96Y}* and *pch-2*

mutants do not have reduced numbers of DNA double-stranded breaks [2, 42], these results

reflect changes in the rate of DNA repair. Further, this indicates that, in contrast to our findings

for pairing and synapsis, PCH-2 does not act through HTP-3 to regulate the rate of DNA repair.

Instead, both proteins regulate DNA repair independent of each other.

We previously reported that *pch-2;htp-3^{H96Y}* double mutants exhibit more achiasmate

chromosomes than either single mutant [2], indicating that the double mutant has a more severe

defect in meiotic crossover recombination than either single mutant and consistent with our

analysis of DNA repair (Figure 3H). To further verify this result on crossover formation, we

quantified the number of GFP::COSA-1 foci for each meiotic nucleus. COSA-1 is a cyclin-like

protein that localizes specifically to presumptive crossovers, serving as a cytological marker for

crossover formation [47]. Because *C. elegans* contain 6 pairs of chromosomes and exhibit

strong crossover interference, most wildtype nuclei have 6 GFP::COSA-1 foci per nucleus

(Figure 3I) [47]. Our findings also reflect this, where we report only 3% of nuclei have less than

6 GFP::COSA-1 foci (Figure 3J). We also found that the incidence of nuclei with less than 6

GFP::COSA-1 foci increases to 9% of nuclei in *pch-2* single mutants (Figure 3J, p value <

0.0001 by Fisher's exact test), recapitulating previous findings that *pch-2* mutants exhibit a loss

of crossover assurance [2, 3]. *htp-3^{H96Y}* single mutants also show an increase in the number of

nuclei with less than 6 GFP::COSA-1 crossovers compared to wildtype (7%, Figure 3J, p value

= 0.003 by Fisher's exact test), consistent with their checkpoint role [42]. Similar to our analysis

of DNA repair, the number of nuclei with less than 6 GFP::COSA-1 foci in *pch-2;htp-3^{H96Y}*

double mutants increases to 11% percent (Figure 3J, p value <0.0001 by Fisher's exact test),

showing an additive effect for crossover defects. These data illustrate that PCH-2 and HTP-3

work independently to regulate crossover formation.

Altogether, our analysis of pairing, synapsis and crossover recombination indicate that PCH-2

and HTP-3 act together to regulate pairing and synapsis but independently to regulate DNA

repair and crossover formation. Since htp-3^{H96Y} single mutants also show defects in the DNA

damage response [43] while pch-2 mutants do not [48], this separation of function is not

necessarily surprising. However, more importantly these data strongly support a model in which

PCH-2 acts through HTP-3's HORMA domain to control the rate of pairing and synapsis but not

the process of meiotic recombination.

htp-3^{H96Y} does not affect meiotic progression

In budding yeast, Pch2's remodeling of the meiotic HORMAD, Hop1, contributes to meiotic

progression [32]. Since htp-3^{H96Y} mutants genetically interact with pch-2 mutants in _C. elegans_,

we also tested whether htp-3^{H96Y} mutants affected meiotic progression. We assessed meiotic

progression by performing immunofluorescence against a factor required for DNA double-strand

break formation, DSB-1 (Figure S1A). DSB-1 localizes to meiotic chromosomes to facilitate

DNA double-strand break formation [49]. In response to meiotic defects, DSB-1 persists on

chromosomes to maintain a period of competency for DNA double-strand break formation to

promote crossover assurance [49] and this persistence correlates with other markers of delays

in meiotic progression [50, 51]. We quantified the fraction of meiotic nuclei that were positive for

DSB-1 and could not detect any difference between control and htp-3^{H96Y} mutant germlines

(Figure S1B, p value = 0.8269 by Fisher's exact test), illustrating that this mutation does not

affect meiotic progression.

A null mutation in htp-3 abolishes the delay in meiotic progression observed when there are

defects in synapsis, such as in syp-1 mutants [50]. To determine whether htp-3^{H96Y} affected

meiotic progression in syp-1 mutants, we quantified the fraction of DSB-1 positive meiotic nuclei

in both syp-1 single mutants and syp-$1;htp$-3^{H96Y} double mutants and could not observe any

difference between these genotypes (Figure S1C, p value = 1.0 by Fisher's exact test),

indicating that *htp-3*^{H96Y} does not affect the extension of meiotic progression observed when

there are meiotic defects.

him-3^{R93Y} does not suppress the acceleration of pairing or synapsis in *pch-2* mutants

Our findings that PCH-2 acts on HTP-3 to promote pairing and synapsis, but not recombination,

indicates that PCH-2 regulates recombination through an HTP-3 independent mechanism.

Further, it is unclear whether PCH-2 regulates pairing and synapsis only through its interaction

with HTP-3 or whether other meiotic HORMADs also contribute to this regulation. To address

these possibilities, we next tested whether *pch-2* genetically interacts with similar mutations in

him-3 or *htp-1*, genes that encode the two other HORMADs essential for meiosis in *C. elegans*.

We performed a Clustal Omega alignment of HTP-3 with the other three meiotic HORMADs [52]

and observed that the mutation in *htp-3*^{H96Y} was not of a conserved residue but was adjacent to

two invariant residues in all four meiotic HORMADs (red box in Figure 2B). Further, based on

this alignment, the arginine at position 93 of HIM-3 should be positioned similarly within the

HORMA domain (Figures 2C and D). Using CRISPR/Cas9 gene editing, we made a

corresponding missense mutation to the *htp-3*^{H96Y} allele in *him-3*, where arginine 93 was

replaced with a tyrosine (*him-3*^{R93Y}). Indirect immunofluorescence verified that this mutation did

not affect HIM-3's localization to chromosomes (Figure S2), progeny viability or the production

of male progeny (Table 1).

First we analyzed pairing in *syp-1*, *pch-2;syp-1*, *him-3*^{R93Y}*;syp-1*, and *pch-2;him-3*^{R93Y}*;syp-1*

mutants by performing immunofluorescence against HIM-8 (Figure 4A and 4B). *him-3*^{R93Y}*;syp-1*

double mutants show 44% of nuclei completed pairing in zone 2, similar to *syp-1* single mutants

(p value =1.00 by two-tailed Fisher's exact test), demonstrating that this mutation does not

affect pairing (Figure 4B). In contrast to our analysis of the *htp-3^{H96Y}* allele, we found that *pch-*

2;him-3^{R93Y};syp-1 triple mutants also show an accelerated rate of pairing phenotype (63%,

Figure 4B) similar to that of *pch-2;syp-1* double mutants (p value = 0.52 by two-tailed Fisher's

exact test). Therefore, we detect no genetic interaction between *pch-2* and *him-3^{R93Y}* in affecting

the rate of pairing.

Next, we assessed the rate of synapsis in wildtype, *pch-2*, *him-3^{R93Y}*, and *pch-2;him-3^{R93Y}*

germlines by performing immunofluorescence against SYP-1 and HTP-3 (Figure 4C and 4D). In

both *pch-2* and *him-3^{R93Y}* single mutants, we found that there was an increase in the number of

nuclei that completed synapsis in zone 2 (13% for both, compared to the wildtype control of 3%,

Figure 4D), suggesting that synapsis is accelerated in both mutants (p values <0.0001 for both

comparisons by two-tailed Fisher's exact test). However, we found that *pch-2;him-3^{R93Y}* double

mutants display more severe synapsis defects than either single mutant: the rate of synapsis is

delayed throughout zones 2-5 and these double mutants reach only 87% complete synapsis

throughout the germline (Figure 4D). These results demonstrate that PCH-2 and HIM-3 regulate

the progression of synapsis independent of each other.

Finally, we further assessed synapsis in *him-3^{R93Y} and pch-2;him-3^{R93Y}* mutants by scoring

synapsis in *meDf2* heterozygotes (*meDf2/+*). Consistent with a role in regulating synapsis, *him-*

3^{R93Y};meDf2/+ double mutants show a reduced rate of synapsis (Figure 4E) compared to

meDf2/+ controls (see zones 4 and 5). This effect was intermediate in *pch-2;him-3^{R93Y};meDf2/+*

triple mutant worms, providing further support that PCH-2 and HIM-3 both regulate synapsis but

accomplish this through independent mechanisms.

him-3^{R93Y} does not suppress DNA repair defects in pch-2 mutants

We next tested whether *pch-2* genetically interacts with *him-3^R93Y* during the regulation of DNA

repair by scoring RAD-51 foci in N2, *pch-2*, *him-3^R93Y* and *pch-2;him-3^R93Y* mutants (Figure 4F).

In *him-3^R93Y* single mutants, the average number of RAD-51 foci was higher in zone 3 than both

pch-2 and control germlines, suggesting a more rapid installation of RAD-51 and potentially

more DNA double-strand breaks. However, the rate of DNA repair was similar to wildtype

(Figure 4G). *pch-2*;*him-3^R93Y* worms more closely resembled *him-3^R93Y* single mutants than *pch-*

2 single mutants in zone 3 but was intermediate between *him-3^R93Y* and *pch-2* single mutants in

zone 4 (Figure 4G), indicating that the accelerated rate of DNA repair observed in *pch-2*

mutants is not fully suppressed by the *him-3^R93Y* mutation and that these two factors do not act

in the same pathway to regulate DNA repair.

<u>*him-3^R93Y* suppresses crossover recombination defects in *pch-2* mutants</u>

To analyze the effect of *him-3^R93Y* on crossover recombination, and test for a genetic interaction

with *pch-2* mutants, we scored GFP::COSA-1 foci in wildtype, *pch-2, him-3^R93Y* and *pch-2;him-*

3^R93Y mutant strains (Figure 4H). Similar to *pch-2* single mutants, we found that there was an

increase of nuclei with less than 6 GFP::COSA-1 foci in *him-3^R93Y* mutants compared to wildtype

controls (10%, Figure 4I, p value = 0.0004 by two-tailed Fisher's exact test), showing that

mutation of this residue in HIM-3 causes a loss of crossover assurance. In contrast to *pch-2*

mutants and *him-3^R93Y* single mutants, *pch-2;him-3^R93Y* double mutants have a reduced number

of nuclei with less than 6 foci compared to either single mutant and more closely resembles the

frequency observed in wildtype nuclei (5%, Figure 4I, p values = 0.016 and 0.0214 by two-tailed

Fisher's exact test, respectively). Therefore, *him-3^R93Y* suppresses the defect on crossover

assurance observed in *pch-2* mutants.

We next visualized whether we could observe this genetic suppression later in meiotic

prophase, when chromosome pairs are linked by chiasmata. Wildtype *C. elegans* oocytes

contain 6 pairs of chromosomes or bivalents, and these can be visualized as 6 "DAPI stained

bodies". Defects in recombination produce homologs without chiasmata, or univalents, in

oocytes (arrows, Figure 4J). In our analysis of wildtype oocytes, we did not observe any

univalents (Figure 4K). Similar to our previous findings [2], we found that there was an increase

in the percent of nuclei with univalents in *pch-2* single mutants compared to wildtype controls

(0.8%, Figure 4K), further demonstrating a loss of crossover assurance that leads to achiasmate

homolog pairs. Correlating with our GFP::COSA-1 analysis, *him-3^{R93Y}* single mutants also have

an increase in the number of oocytes containing univalents (1.3%, Figure 4K), demonstrating

that *him-3^{R93Y}* mutants also observe a loss of crossover assurance. In *pch-2;him-3^{R93Y}* double

mutants, we found no instances of univalents (Figure 4K), further indicating that these two

mutant alleles genetically suppress each other in the context of crossover recombination.

Taken together, our results are consistent with a model in which PCH-2 acts through HIM-3 to

promote crossover recombination but not pairing or synapsis. Further, the roles of the two

meiotic HORMADs appear to be cleanly delegated, with PCH-2 controlling pairing and synapsis

through HTP-3 and crossover recombination through HIM-3. Since we do not observe a similar

genetic interaction when we analyze DNA repair in *pch-2;him-3^{R93Y}* double mutants, this

regulation of crossover recombination occurs later than the installation and removal of RAD-51.

him-3^{R93Y} does not affect meiotic progression

We next tested whether *him-3^{R93Y}* mutants affect meiotic progression by performing

immunofluorescence against DSB-1 and quantifying the fraction that were DSB-1 positive in

wildtype and *him-3^{R93Y}*mutant strains (Figure S3A). We found that there was not a significant

difference between the number of DSB-1 positive nuclei between wildtype and *him-3^{R93Y}* mutant

strains (Figure S3B, p value =0.645 blank by two-tailed Fisher's exact test), suggesting that this

mutation does not affect meiotic progression.

We further tested whether the *him-3^{R93Y}* mutation affected meiotic progression by quantifying

the fraction of DSB-1 positive nuclei in *syp-1* and *him-3^{R93Y};syp-1* mutant strains. Similar to our

findings for *htp-3^{H96Y}*, we found that there was not a significant difference in the number of DSB-

1 positive nuclei between *syp-1* single mutant*s* and *him-3^{R93Y};syp-1* double mutants (Figure

S3C, p value =1.00 by two-tailed Fisher's exact test). This result further demonstrates that *him-*

3^{R93Y} does not affect meiotic progression.

htp-1^{G97T} does not suppress the accelerated rate of pairing in *pch-2* mutants

Our findings suggest that PCH-2 acts through HTP-3 to promote pairing and synapsis, and HIM-

3 to promote crossover formation. We next wanted to test whether *pch-2* genetically interacts

with a similar mutation in *htp-1*, the gene that encodes the final essential meiotic HORMAD in *C.*

elegans.

Preliminary structural analysis of the HORMA domain of HTP-1 indicates that the glycine

residue that is analogous to the histidine and arginine residues of HTP-3 and HIM-3 (Figure 2B),

respectively, is positioned more internally within the HORMA domain (Figure 2E) and therefore

potentially more constrained in the mutations we can introduce at this site without affecting

HTP-1's stability. Indeed, when we replaced this glycine with a tyrosine by CRISPR/Cas9 gene

editing (*htp-1^{G97Y}*), this mutant phenotypically resembled the *htp-1*(*me84)* null allele [11] in which

this glycine residue is mutated to a glutamate, producing very few viable progeny and a high

incidence of male (HIM) phenotype. Given that the severity of the *htp-1^{G97Y}* mutation would

preclude our analysis of meiotic prophase events, we sought to create a less severe

hypomorphic allele of HTP-1 to test whether it genetically interacts with the *pch-2* null mutation.

To identify a mutation that would allow analysis of pairing, synapsis, recombination, and meiotic

progression, we used CRISPR/Cas9 to substitute glycine 97 with amino acids that progressively

add bulkier side chains at this position (*htp-1^{G97A}*, *htp-1^{G97S}, and htp-1^{G97T}*). Upon initial

examination of these strains, we observed a HIM phenotype associated with the *htp-1^{G97T}*

mutant strain (Table 1), suggesting that this mutation causes meiotic chromosome

misegregation. This HIM phenotype was accompanied by a reduction in embryonic viability

(Table 1). Therefore, we limited our analysis of testing genetic interactions with *pch-2* for

pairing, synapsis, and recombination to *htp-1^{G97T}* mutants. Indirect immunofluorescence showed

no difference in the loading of HTP-1 to meiotic chromosomes in *htp-1^{G97T}* mutants or in *htp-*

1^{G97T} htp-2 double mutants (Figure S4).

We first tested whether pairing is disrupted in this strain by performing immunofluorescence

against HIM-8 in *syp-1*, *pch-2;syp-1*, *htp-1^{G97T};syp-1* and *pch-2;htp-1^{G97T};syp-1* mutant germlines

(Figure 5A and 5B). In *htp-1^{G97T};syp-1* double mutants, the percent of nuclei that have

completed pairing in zone 2 is similar compared to *syp-1* single mutants (Figure 5B, p value

=0.553 by two-tailed Fisher's exact test), indicating that *htp-1^{G97T};syp-1* double mutants do not

have a defect in the progression of pairing. *pch-2;htp-1^{G97T};syp-1* triple mutants have a similar

rate of pairing in zone 2 compared to *pch-2;syp-1* double mutants (Figure 5B, p value =0.311 by

two-tailed Fisher's exact test), showing that the *htp-1^{G97T}* mutation does not suppress the

acceleration of pairing observed in *pch-2* mutants.

htp-1^{G97T} causes severe synapsis defects that are not suppressed by mutation of *pch-2*

Next, we tested whether synapsis is perturbed in *htp-1^{G97T}* mutants and whether this allele

genetically interacts with the *pch-2* null allele (Figure 5C and 5D). Synapsis is severely

disrupted in *htp-1^{G97T}* mutants: despite some chromosomes completing synapsis (Figure 5D),

almost all meiotic nuclei have some number of unsynapsed chromosomes throughout the

germline, where only 1.3% of nuclei reach complete synapsis (Figure 5D). In *pch-2;htp-1^{G97T}*

double mutants, we find that synapsis is disrupted as severely as *htp-1^{G97T}* single mutants,

where 1.3% of nuclei also reach complete synapsis (Figure 5D, p value =0.717 by two-tailed

Fisher's exact test), indicating that these alleles do not genetically interact and regulate

synapsis independent of each other.

htp-1^{G97T} leads to a delay in RAD-51 removal that is not suppressed by mutation of *pch-2*

Next, we assessed the progress of DNA repair in *htp-1^{G97T}* and *pch-2;htp-1^{G97T}* mutants by

scoring RAD-51 foci throughout the germline (Figure 5E and 5F). In comparison to control

germlines, *htp-1^{G97T}* single mutants exhibit a delay in the installation of RAD-51 foci (Figure 5F).

However, as meiotic prophase continues, meiotic nuclei in *htp-1^{G97T}* single mutants have more

RAD-51 foci and substantially delay their removal, possibly as an indirect consequence of the

defects in synapsis we observe in this mutant (Figure 5D). *pch-2;htp-1^{G97T}* double mutants no

longer exhibit this delay in RAD-51 installation, suggesting a partial suppression of this

particular phenotype, but display more RAD-51 foci, particularly in zones 4 and 5, than wildtype

meiotic nuclei and *pch-2* single mutants, similar to *htp-1^{G97T}* single mutants. Thus, PCH-2 and

HTP-1 do not appear to genetically interact to regulate the rate of DNA repair.

htp-1^{G97T} causes defects in crossover recombination that are not suppressed by *pch-2*

mutations

To continue our analysis, we analyzed crossover recombination by scoring the number of

GFP::COSA-1 foci per nucleus in wildtype, *pch-2*, *htp-1^{G97T}*, and *pch-2;htp-1^{G97T}* mutants (Figure

5G and 5H). Surprisingly, we found that in *htp-1^{G97T}* single mutants, the number of COSA-1 foci

varies greatly across nuclei, ranging from 4 to 9, with only 39% of nuclei have 6 GFP::COSA-1

foci (Figure 5H), suggesting that the *htp-1*^{G97T} mutation causes misregulation of crossover

formation. In addition to the loss of crossover assurance (nuclei with less than 6 GFP::COSA-1

foci), we also observe a loss of crossover interference (nuclei with greater than 6 GFP::COSA-1

foci), indicating a general loss of crossover control in *htp-1*^{G97T} mutants. In *pch-2*;*htp-1*^{G97T}

double mutants, we observe a similar phenotype where the number of GFP::COSA-1 varies

across nuclei, ranging from 4 to 9, with only 45% having 6 COSA-1 foci (Figure 5H, p value

=0.1725 by two-tailed Fisher's exact test). Overall, these data indicate that PCH-2 and HTP-1

do not genetically interact to regulate crossover recombination.

We further analyzed crossover formation by looking directly at bivalents in oocytes and

quantified the frequency of univalents in wildtype, *pch-2*, *htp-1*^{G97T}, and *pch-2*;*htp-1*^{G97T} mutants

(Figure 5I and 5J). In *htp-1*^{G97T} single mutants, we found that the frequency of univalents

significantly increases to 41% (Figure 5J, p value < 0.0001 by two-tailed Fisher's exact test),

further verifying the defect in crossover formation in *htp-1*^{G97T} single mutants. For *pch-2*;*htp-*

1^{G97T} double mutants, we also found that the frequency of univalents is statistically higher

compared to wildtype oocytes (31%,Figure 5J, p value < 0.0001 by two-tailed Fisher's exact

test) but not significantly different when compared to *htp-1*^{G97T} single mutants (31%, Figure 5J, p

value =0.182 by two-tailed Fisher's exact test). Taken together, this suggests that both *pch-2*

and *htp-1* both promote crossover assurance but regulate it through different mechanisms.

htp-1^{G97T} delays meiotic progression and this delay is partially suppressed by mutation of *pch-2*

The absence of HTP-1 accelerates meiotic progression, producing non-homologous synapsis

and more rapid DNA repair [11]. In contrast, mutations within the HORMA domain of HTP-1 that

prevent its loading on meiotic chromosomes cause a delay in meiotic progression, producing

asynapsis [53]. We tested whether htp-1^{G97T} mutations similarly affect meiotic progression by

performing immunostaining against DSB-1.

We quantified the fraction of DSB-1 positive meiotic nuclei in wildtype and htp-1^{G97T} mutant

germlines (Figure 6A). We found that htp-1^{G97T} mutants display a statistically significant increase

in the fraction of DSB-1 positive nuclei (81%, p value < 0.0001 by two-tailed Fisher's exact test

Figure 6B), indicating that this mutation produces a delay in meiotic progression. Since DSB-1 is

required for DNA double-strand break formation, the extension of DSB-1 likely explains the

increase in RAD-51 foci (Figure 5F) and the attenuation of crossover interference (Figure 5H) in

htp-1^{G97T} mutants. Moreover, since the increase in RAD-51 foci formation (Figure 5F) is

accompanied by an increase in GFP::COSA-1 foci (Figure 5H), htp-1^{G97T} mutants also appear to

have defects in crossover homeostasis, a mechanism that ensures that the number of

crossovers remain invariant even when the number of crossover precursors, such as DNA

double-strand breaks, vary [54] and which is particularly robust in $C.$ $elegans$ [47].

Given the severity of this phenotype, we wondered if the less severe mutations we generated,

htp-1^{G97A} and htp-1^{G97S} mutants, might also affect meiotic progression, albeit more weakly

(Figure 6A). While htp-1^{G97A} mutants did not affect the fraction of DSB-1 positive nuclei, htp-

1^{G97S} mutants produced a small but statistically significant delay in meiotic progression (63%

DSB-1 positive nuclei, Figure 6B, p value =0.0006 by two-tailed Fisher's exact test), indicating

that the introduction of bulkier amino acids at this residue produces an allelic series with respect

to defects in meiotic progression.

We next tested whether this defect in meiotic progression in htp-1^{G97T} single mutants is

suppressed by the pch-2 mutation. We found that the percent of DSB-1 stained nuclei is slightly

but significantly lower in pch-2;htp-1^{G97T} double mutants (77%, Figure 6B) compared to htp-1^{G97T}

single mutants (Figure 6B, p value = 0.0047 by two-tailed Fisher's exact test), indicating that the

meiotic progression defect in *htp-1^{G97T}* mutants is partially suppressed by mutations in *pch-2*

and that these two proteins cooperate to regulate meiotic progression.

him-3^{R93Y} and *htp-1^{G97T}* mutants have functional meiotic checkpoints

We previously showed that *htp-3^{H96Y}* mutants abolishes both the DNA damage response and

the synapsis checkpoint [43], similar to null mutations in *htp-3* and *him-3* [55]. A null mutation in

htp-1 only affects the synapsis checkpoint [55]. To determine whether *him-3^{R93Y}* or *htp-1^{G97T}*

mutants affect meiotic checkpoint activation, we introduced each mutation into the *syp-1* mutant

background and evaluated the levels of germline apoptosis. *syp-1* mutants activate both the

DNA damage response and the synapsis checkpoint, producing high levels of germline

apoptosis [48]. *him-3^{R93Y}* mutants showed wildtype levels of apoptosis while *htp-1^{G97T}* mutants

exhibited germline apoptosis similar to that of *syp-1* single mutants (Figure S6, p value < 0.0001

by student t-test), likely as a consequence of the asynapsis and defects in recombination we

observe (Figure 5). Both *syp-1;him-3^{R93Y}* and *syp-1;htp-1^{G97T}* double mutants showed similar

levels of germline apoptosis as *syp-1* single mutants (Figure S6), indicating that both *him-3^{R93Y}*

or *htp-1^{G97T}* mutants are competent for meiotic checkpoint activation, unlike *htp-3^{H96Y}* mutants

[43].

him-3^{R93Y} mutants maintain the structure of the HORMA domain but have reduced affinity for

closure motifs

Our data thus far demonstrates that *pch-2* genetically interacts with specific alleles of *htp-3*,

him-3 and *htp-1*, with varying effects on pairing, synapsis, recombination and meiotic

progression. However, how these genetic interactions translate into a molecular understanding

of the functional relationship between PCH-2 and these meiotic HORMADs is unclear. To

address this, we tested what effect these mutations had on the ability of meiotic HORMADs to

bind their closure motifs and adopt the closed conformation.

While the closure motif for HTP-3 is unknown, HIM-3 is known to bind closure motifs found

within the C-terminal tail of HTP-3 [19]. HIM-3 also contains a closure motif in its own C-

terminus but it preferentially binds closure motifs in HTP-3 to contribute to a hierarchical

HORMAD assembly [19]. We cloned and expressed wildtype HIM-3 and HIM-3^{R93Y} in *E. coli*,

purified each protein, and performed a fluorescence polarization peptide binding assay to test

how well HIM-3^{R93Y} binds a peptide encoding HTP-3 closure motif #4 *in vitro*. Consistent with

prior work, we found that wildtype HIM-3 binds the HTP-3 closure motif with a K_d of 220 +/- 20

nM and found that it does not bind to a control peptide with a scrambled sequence (Figure 7A).

HIM-3^{R93Y} binds to the same closure motif with an increased K_d of 690 +/- 130 nM, indicating

that HIM-3^{R93Y} binds the HTP-3 closure motif with reduced affinity compared to wildtype HIM-3

(Figure 7A). The thermal stability of HIM-3^{R93Y} is similar to that of wild-type HIM-3 (Figure S5),

and a crystal structure of HIM-3^{R93Y} in the closed conformation is equivalent to that of wildtype

HIM-3 (Figure 2D), suggesting that reduced affinity for the closure motif for HIM-3^{R93Y} is not due

to instability of the protein or an inability to adopt the closed conformation. These results

demonstrate that HIM-3^{R93Y} binds the HTP-3 closure motif less readily than wildtype HIM-3,

which would potentially limit its ability to adopt the closed conformation *in vivo* (Figure 7B).

Because this residue is positioned similarly within all three-essential meiotic HORMADs (Figure

2B), our biochemical analysis of HIM-3^{R93Y} may be generalizable to the mutant versions of HTP-

3 and HTP-1: HTP-3^{H96Y} and HTP-1^{G97T} could bind their respective closure motifs with less

affinity, affecting their ability to adopt the closed conformation *in vivo*.

Unlike the budding yeast meiotic HORMAD, Hop1 [56], HIM-3 is difficult to purify in any

conformation aside from its closed conformation [19], preventing us from directly assessing

whether the HIM-3^{R93Y} mutation affects the protein's closed-versus-unbuckled conformational

equilibrium. However, the reduced affinity of HIM-3^{R93Y} for the HTP-3 closure motif, when

combined with the established conformational dynamics of Hop1 [56], the localization of PCH-2

by structured illumination microscopy (Figure 1) and our genetic and cytological analysis

(Figures 3-6) of how these mutant alleles suppress some defects observed in *pch-2* mutants,

presents a model consistent with PCH-2 remodeling meiotic HORMADs from a closed to

unbuckled conformation on meiotic chromosomes to control the progress of pairing, synapsis,

meiotic recombination and cell cycle progression (Figure 8).

Discussion

Since the discovery of Pch2 in budding yeast more than 20 years ago, the particular roles of

Pch2/PCH-2/TRIP13 in meiosis have been enigmatic. More recently, biochemical and structural

analysis of how mammalian TRIP13 contributes to spindle checkpoint progression by

remodeling the HORMA domain protein MAD2 have provided a useful framework to establish

Pch2/PCH-2/TRIP13's diverse roles in structuring meiotic chromosomes for homolog pairing,

synapsis, and recombination, and signaling meiotic progression. In the spindle checkpoint,

PCH-2/TRIP13 promotes checkpoint activation by maintaining a pool of soluble "open" MAD2,

which is used to build the mitotic checkpoint complex at unattached kinetochores and delay

anaphase onset [57-59]. After kinetochore attachment, TRIP13 plays an additional role in

human cells, disassembling the mitotic checkpoint complex by remodeling MAD2, promoting

checkpoint inactivation and anaphase progression [60, 61]. This activity may not necessarily be

conserved, given that a similar role in silencing the checkpoint is not observed in *C. elegans* [59,

62], and Pch2 is not expressed outside meiosis in *S. cerevisiae* [38]. Nonetheless, in some

systems, TRIP13 appears to play dual roles in spindle checkpoint activation and inactivation,

through a single biochemical activity of converting "closed" MAD2 to the "open" conformer.

Recently, studies of Pch2 in budding yeast, and by extension plants and mammals, have

suggested that this enzyme plays a similar dual role in meiotic chromosome organization,

homologous recombination, and meiotic progression. Meiotic HORMADs in most organisms

adopt a "self-closed" conformation with their HORMA domains bound to a closure motif in their

own C-terminus [19, 28, 30]. In budding yeast and plants, Pch2/TRIP13's remodeling activity is

needed to convert these "self-closed" HORMADs into an "open" or "unbuckled" conformation to

promote their nuclear import and association with chromosomes [29, 30]. After assembly into

axial elements of meiotic chromosomes, HORMADs promote meiotic recombination and

homolog pairing [6, 63], likely through direct protein-protein interactions with partner proteins

[23]. Once homologs have recombined and synapsed, Pch2/TRIP13 is recruited to the

synaptonemal complex [34, 35, 38], where it is thought to disassemble HORMAD-axis protein

complexes to deplete them [10, 33, 35]. HORMAD depletion down-regulates recombination,

promotes rapid repair of remaining DNA breaks, and facilitates meiotic progression [64, 65].

In *C. elegans*, several key differences in the timing of meiotic events relative to yeast, plants,

and mammals necessitate a rethinking of PCH-2's roles in meiotic progression. First, homolog

pairing and synapsis occur independently of recombination in *C. elegans* [66], such that

synapsis cannot serve as a signal for successful interhomolog recombination as in other

organisms. Second, meiotic HORMADs localize to meiotic chromosomes independently of

PCH-2 [2] and are not visibly depleted in response to synapsis [21, 22, 36], indicating the dual

role for Pch2/TRIP13 proposed in other systems is not relevant to understanding PCH-2's

function in *C. elegans* meiosis. Finally, a null mutation of *pch-2* or mutations that eliminate its

ATP hydrolysis activity result in accelerated homolog pairing, synapsis, and meiotic

recombination, leading to a loss of meiotic fidelity [2, 3]. Because these events are likely

mediated by direct HORMAD-partner protein interactions on meiotic chromosomes, and

because PCH-2 is localized to chromosomes during the relevant stages of meiotic prophase

(Figure 1) [2], these phenotypes suggest that PCH-2 either reduces overall levels of HORMADs

or increases their association-dissociation dynamics to modulate these key events.

Our experiments from this study shed further light on this model. We have identified mutations

in the HORMA domains of both HTP-3 and HIM-3 (Figure 2A) that appear to load normally on

chromosomes [42] (Figures S2 and S4), reduce the affinity of closure motif interactions in vitro

(Figure 7) and suppress specific defects of a *pch-2* null mutation in homolog pairing/synapsis

and recombination, respectively (Figures 3 and 4). These data lead us to propose a model in

which PCH-2 remodels the HORMA domains of chromosome-associated HORMADs from the

"closed" conformation to the "open/unbuckled" conformation to temporarily reduce HORMAD

occupancy on chromosomes, decelerating HORMAD-dependent homolog pairing, synapsis,

and recombination to promote meiotic fidelity (Figure 8). An attractive but still speculative

extension to this model is that PCH-2's remodeling of meiotic HORMADs destabilizes

HORMAD-partner protein interactions essential for homolog pairing, synapsis, and

recombination, deliberately slowing their progress.

Our model has important implications for the established roles of Pch2/TRIP13 in other

organisms. First, Pch2/PCH2 in both yeast and plants localize specifically to sites associated

with synapsis initiation and recombination [34, 35], suggesting a similar role temporarily

reducing HORMAD occupancy at these sites prior to synapsis. Further, despite the observation

that Pch2/TRIP13 strongly depletes meiotic HORMADs from chromosomes upon synapsis,

these proteins are not completely removed from chromosomes in any organism [10, 33, 35],

suggesting roles for both meiotic HORMADs and Pch2/TRIP13 even after chromosome

synapsis. Given that the gradual implementation of crossover formation partially overlaps when

chromosomes are synapsed in budding yeast, plants, and mice [67-70], we propose that the

presence of Pch2/TRIP13 on synapsed chromosomes remodels meiotic HORMADS to regulate

the number and location of crossovers, similar to one role of PCH-2 in *C. elegans* [2, 71].

The relationship between PCH-2 and HTP-1 is less clear. Despite the small structural

differences between HIM-3 and HTP-1 (Figure 2E) that required some adjustments in how we

designed our mutational analysis of HTP-1, we predict that HTP-1^{G97T} also shows a reduced

affinity for its closure motifs in both HTP-3 and HIM-3, potentially shifting its equilibrium toward

the unbuckled conformation. The consequences of this difference in protein function are a

significant delay in meiotic progression (Figure 6B) and defects in synapsis, DNA repair and

crossover recombination (Figure 5D, 5F, & 5H). Since previous data has implicated HTP-1 and

its regulation in controlling meiotic progression, with indirect effects on synapsis, DNA repair

and crossover recombination [11, 53, 72] , we favor the interpretation that the primary defect in

htp-1^{G97T} mutants is the delay in meiotic progression, with secondary, indirect consequences for

synapsis and recombination. Mutation of *pch-2* partially suppresses this defect in meiotic

progression (Figure 6B), with little effect on synapsis or recombination (Figure 5D & 5H),

supporting our interpretation. How might a delay in meiotic progression prevent complete

synapsis or the normal completion of recombination? This meiotic delay might maintain a cell

cycle state in which the synaptonemal complex and/or recombination intermediates are more

labile and subject to turnover, through mechanisms such as post-translational modifications,

high chromosome mobility, some other unknown condition or some combination of these

conditions. Finally, if we are correct and an increase in unbuckled HTP-1^{G97T} is responsible for a

meiotic delay, with indirect consequences on synapsis and recombination, our data also

suggest that, unlike open MAD2, unbuckled versions of meiotic HORMADs may have additional

functions aside from being an important structural intermediate poised for binding to closure

motifs in partner proteins.

That mutation of *pch-2* only weakly suppresses the delay in meiotic progression we observe in

htp-1^{G97T} mutants raises two possibilities: 1) the *htp-1^{G97T}* mutation is too severe to allow us to

observe this genetic interaction; or 2) PCH-2 does not directly act on HTP-1 to regulate meiotic

progression. How might PCH-2 contribute to the regulation of meiotic progression without

directly remodeling HTP-1? Since both HTP-3 and HIM-3 have HTP-1 closure motifs in their C-

termini [19], PCH-2's remodeling of HTP-3 and HIM-3 may indirectly result in the removal, and

remodeling, of HTP-1, contributing to the regulation of meiotic progression. This possibility is

particularly attractive since it explicitly coordinates meiotic progression with the rate of pairing,

synapsis and recombination.

Finally, our work shows that PCH-2's regulation of pairing, synapsis, recombination and meiotic

progression are cleanly delegated to the remodeling of different meiotic HORMADs: PCH-2

remodels HTP-3 from a closed to an unbluckled conformation to decelerate pairing and

synapsis and contribute to checkpoint activation, HIM-3 to decelerate crossover recombination

and HTP-1, either directly or indirectly, to control meiotic progression. This deceleration allows

for proofreading of the homolog interactions that underlie these events and ensure fidelity [3].

Similar separation of function has been observed in plants and mice: each have two meiotic

HORMADs, only one of which (HORMAD1 in mice, ASY1 in plants), appears to be essential for

pairing, synapsis and recombination [9, 15-17, 73]. In the case of mice, a second HORMAD,

HORMAD2, is not required for pairing, synapsis or recombination but is essential for efficient

meiotic checkpoint responses [18, 74]. In plants, no role for ASY2 has been reported yet. The

separation of function we observe in *C. elegans* is slightly different: The three essential meiotic

HORMADs have overlapping but distinct roles that nonetheless significantly impact pairing,

synapsis and recombination [11-13, 19, 21, 22]. However, it is through their regulation by PCH-2

that their distinct roles become more clear, allowing for the coordination of pairing, synapsis,

recombination and meiotic progression. This delegation of tasks, as revealed by their regulation

by PCH-2, may also explain why this family has expanded so dramatically in *C. elegans*. Since

multiple systems, including mammals, also have more than one meiotic HORMAD, this

delegation of tasks may be a conserved evolutionary feature of meiosis.

Another possible contributing factor to the expansion of this family in *C. elegans* is the

mechanistic uncoupling of meiotic recombination with synapsis. Mice, budding yeast and plants

all rely on the initiation of meiotic recombination [75-78] to promote accurate synapsis. Similarly,

all of these organisms have a single essential meiotic HORMAD controlling pairing, synapsis

and recombination [6, 9, 15], suggesting a possible connection between these observations. In

contrast to these systems, *C. elegans* does not rely on meiotic recombination for accurate

chromosome synapsis [66], suggesting that the family of meiotic HORMADs may have

expanded to accommodate the fact that synapsis and recombination are regulated by two

different mechanisms in this system.

While *C. elegans* is unique in that it has four meiotic HORMADs which assemble into a

hierarchical complex, it is formally possible that other systems also have different functional

classes of meiotic HORMADs, even when they have a single meiotic HORMAD in their genome.

For example, in budding yeast, Hop1 can bind directly to nucleosomal DNA [79], a closure motif

in the axis protein Red1 and closure motifs in Hop1 once it has bound to Red1, producing

oligomerization [28]. These three different classes of Hop1, potentially defined by the molecular

nature of their binding to meiotic chromosomes, may also assemble hierarchically and have

different effects on the events of meiotic prophase. These differences could affect their

regulation by Pch2, producing a similar delegation of functions as we observe with HTP-3, HIM-

3 and HTP-1. For example, it has already been shown in budding yeast that in regions adjacent

to telomeres, Hop1 is resistant to Pch2-mediated redistribution [80], consistent with this

possibility of differential regulation by Pch2.

Our data show that the remodeling of meiotic HORMADs by PCH-2 have consequences on the

progress, coordination, and fidelity of meiotic prophase events. Major questions that arise from

this work are 1) how is PCH-2 activity limited to meiotic HORMADs in some contexts but not

others and 2) how do these meiotic HORMADs contribute to the events of pairing, synapsis and

crossover recombination? How does HTP-3 molecularly participate in pairing and synapsis?

How does HIM-3 promote crossover recombination? And how does HTP-1 control meiotic

progression? Future studies focused on understanding how meiotic HORMADs molecularly

interact with each other, other axis proteins, and/or additional binding partners will shed new

light on how meiotic HORMADs, in collaboration with PCH-2, drive the coordination of meiotic

prophase events to promote fidelity and prevent aneuploidy.

Materials and Methods

Genetics, Worm strains

The C. *elegans* Bristol N2 strain was used as the wildtype strain throughout this study [81]. All

strains were grown at 20°C under standard conditions unless otherwise stated. Mutations and

rearrangements used were as follows:

LG I: *htp-3(vc75), mnDp66*

LG II: *pch-2(tm1458), meIs8 [Ppie-1::GFP::cosa-1 + unc-119(+)]*

LG IV: *him-3(blt9), htp-1(blt10), htp-1(blt11), htp-1(blt12), htp-2(tm2543), nT1[unc-?(n754let-*

?(m435)]

LG V: *syp-1(me17), bcIs39[Plim-7::ced-1::gfp; lin-15(+)]*

LG X: *meDf2*

The *him-3^{R93Y}* (*him-3[blt9]*), *htp-1^{G97A}* (*htp-1[blt10]*), *htp-1^{G97S}* (*htp-1[blt11]*), and *htp-1^{G97T}* (*htp-*

1[blt12]) were generated by CRISPR-Cas9 genome editing supplemented with the *dpy-10* co-

conversion method [82].

The following guide RNA sequences were used:

dpy-10 sgRNA: 5'GCU ACC AUA GGC ACC ACG AGG UUU UAG AGC UAU GCU 3'

him-3 sgRNA: 5'CGU GUG UCU UCA ACA UUC GAG UUU UAG AGC UAU GCU 3'

htp-1 sgRNA: 5' UCA ACU ACU UCG AAA UGC UGG UUU UAG AGC UAU GCU 3'

The following DNA repair oligonucleotide sequences were used:

dpy-10: 5' CAC TTG AAC TTC AAT ACG GCA AGA TGA GAA TGA CTG GAA ACC GTA CCG

CAT GCG GTG CCT ATG GTA GCG GAG CTT CAC ATG GCT TCA GAC CAA CAG CCT AT

3'

him-3^{R93Y}: 5' GCA TTC CAG AGC AGT GTT CCT GCC GCA AAG CGT GTG TCT TCA ACA

TTT GAC GGA TTG TAC GAT GCG ATT CAA CAA GGC TAT TTG CGA GAG TTC GCA ATC

GTG TTC TAC AAG 3'

htp-1^{G97A}: 5' GCT TCA GAA ACC CAC GGT CTA ACC AAA TCG CTC AAC TAC TTC GAA

ATG CTG CTG ATG CAA CAA AGG ATG GGT TTC TGA AAG AAG TCT CCC TCG TGA TCA

CAA ATA ATG 3'

htp-1^G97S: 5' GCT TCA GAA ACC CAC GGT CTA ACC AAA TCG CTC AAC TAC TTC GAA

ATG CTT CAG ATG CAA CAA AGG ATG GGT TTC TGA AAG AAG TCT CCC TCG TGA TCA

CAA ATA ATG 3'

htp-1 ^G97T: 5' 5' GCT TCA GAA ACC CAC GGT CTA ACC AAA TCG CTC AAC TAC TTC GAA

ATG CTA CAG ATG CAA CAA AGG ATG GGT TTC TGA AAG AAG TCT CCC TCG TGA TCA

CAA ATA ATG 3'

For CRISPR/Cas9 gene editing, all mutations were made in the N2 wildtype strain, with the

exception of the *htp-1^G97T htp-2* double mutant strain, in which the *htp-1^G97T* mutation was

introduced in the *htp-2(tm2543)* mutant background. sgRNAs and oligo repair templates were

synthesized by Integrated DNA technologies (IDT, Coraville, IA). Injection mixes contained

either *htp-1* or *him-3* sgRNA (100 μM final), *dpy-10* sgRNA (100 μM final, [82]), tracrRNA (100

μM final, IDT), purified Cas9 protein (40 μM final), *htp-1* or *him-3* DNA repair oligo (50 μM final,

IDT) and *dpy-10* repair oligo (50 μM final, IDT, [82]). Young adults were injected, recovered and

stored at either 15 or 20 degrees. Roller or *dpy* F1's were singled to individual plates seeded

with OP50 and screened for presence of the mutant allele by PCR and BspHI restriction digest

(*htp-1)* or RsaI restriction digest (*him-3*). Individual F2's were then isolated from F1 plates that

contained the allele to identify a homozygous strain.

All mutations were verified by sequencing and strains were backcrossed against wildtype worms

three times.

Immunofluorescence, Antibodies and Microscopy

Day 1 adult hermaphrodites were dissected 24-28 hours post-late L4 stage and immunostaining

and DAPI staining were performed similarly to [48]. For analyzing bivalents, hermaphrodites

were dissected and DAPI stained 48 hours post late L4 stage.

The following primary antibodies were used for this study: rat anti-HIM-8 1:500 [44], guinea pig

anti-HTP-3 1:250 [36], chicken anti-HTP-3 1:250 [36], rabbit anti-SYP-1 1:500 [40], guinea pig

anti-SYP-1 1:500 [40], rabbit anti-HIM-3 1:500 [14], rabbit anti-HTP-1 1:400 [22], rabbit anti-

RAD-51 1:250 (Novus Biologicals, Littleton, CO), guinea pig anti-DSB-1 1:250 [49], mouse anti-

GFP 1:100 (Invitrogen, Waltham, MA) and rabbit anti-PCH-2 1:500 [2].

The following secondaries were used for this study: Alexa 488 anti-guinea pig (Invitrogen),

Alexa 488 anti-rabbit (Invitrogen), Alexa 488 anti-mouse (Invitrogen), Cy3 anti-rabbit (Jackson

Immunochemicals, West Grove, PA), Cy3 anti-rat (Jackson Immunochemicals), Cy3 anti-guinea

pig (Jackson Immunochemicals), and Cy5 anti-chicken (Jackson Immunochemicals). All

secondary antibodies were used at a 1:500 dilution.

To visualize meiotic nuclei and bivalents, DAPI was used at a dilution of 1:10,000.

For analysis of the rate of paining, "paired" nuclei were scored as a single focus, whether it was

elongated or circular. "Unpaired" nuclei were scored as two completely separate foci that each

had an individual border.

For analysis of the rate of synapsis, nuclei were considered "completely synapsed" when

complete colocalization of HTP-3 and SYP-1 was observed. "Unsynapsed" nuclei were

assigned when nuclei had stretched of HTP-3 with missing SYP-1 signal.

Scoring of germline apoptosis was performed as previously described in [48] in strains

containing *bcIs39 [lim-7p::ced-1::GFP + lin-15(+)]* with the following exceptions: L4

hermaphrodites were allowed to age for 22-24 hours. They were then mounted under coverslips

on 2% agarose pads containing 0.2mM levamisole and scored. A minimum of twenty-five

germlines was analyzed for each genotype.

Line scans measuring pixel intensity for Figure 1D were performed in ImageJ. Channels were

split and the pixel intensities for each fluorophore was measured across the distance of the

region of interest (Figure 1C, yellow line). Distance in pixels and normalized intensity were

plotted using R (R core team) with the ggplot2 package [83].

All images except for those in Figure 1 were acquired using a DeltaVision Personal DV system

(Applied Precision) equipped with a 100X N.A. 1.40 oil-immersion objective (Olympus), resulting

in an effective XY pixel spacing of 0.064 or 0.040 μm. Three-dimensional image stacks were

collected at 0.2-μm Z-spacing and processed by constrained, iterative deconvolution. For

structured Illumination microscopy, images were obtained as 125 nm spaced Z stacks, using a

100x NA 1.40 objective on a DeltaVison OMX Blaze microscopy system, 3D-reconstructed and

corrected for registration using SoftWoRx. Image scaling and analysis were performed using

functions in the softWoRx software package. Projections were calculated by a maximum

intensity algorithm. Composite images were assembled and some false coloring was performed

with ImageJ.

Whole germline images were assembled in ImageJ as in [84] using pairwise stitching and/or

grid collection stitching [85]. Quantification of pairing, synapsis, RAD-51 foci, GFP::COSA-1 foci,

and DSB-1 positive nuclei was performed as in [3] with a minimum of three germlines per

genotype.

Protein Expression and Purification

The full-length HIM-3 gene (residues 2-291) was PCR-amplified from a *C. elegans* cDNA library

and cloned into UC Berkeley Macrolab vector 2-CT (Addgene #29706), which encodes a TEV

protease-cleavable His_6-MBP (maltose binding protein) tag. PCR-based mutagenesis was used

to generate the R93Y mutant construct. Proteins were expressed in *E. coli* strain Rosetta2

pLysS (EMD Millipore) at 20°C for 16 hours, then harvested by centrifugation. Proteins were

purified using Ni^{2+} affinity chromatography (HisTrap HP, Cytiva), followed by TEV protease

treatment [86] for 48 hours. The cleaved mixture was passed through Ni^{2+} affinity resin a second

time to remove uncleaved protein, free His_6-MBP tags, and His_6-tagged TEV protease, and the

flow-through was concentrated and further purified by size exclusion chromatography (Superdex

200; Cytiva). Fractions were concentrated by ultrafiltration (Amicon Ultra, EMD Millipore),

aliquoted, and stored at -80°C.

Fluorescence polarization binding assays

N-terminal FITC-Ahx labeled peptides were synthesized (BioMatik), resuspended in DMSO,

then diluted into binding buffer (20 mM Tris pH 7.5, 300 mM NaCl, 10% glycerol, 1 mM DTT,

0.1% NP-40). Fifty μL reactions containing 50 nM peptide plus indicated amounts of HIM-3 (WT

or R93Y) protein were incubated 60 minutes at room temperature, then fluorescence

polarization was read in 384-well plates using a TECAN Infinite M1000 PRO fluorescence plate

reader. All binding curves were done in triplicate. Binding data were analyzed with Graphpad

Prism v. 8 using a single-site binding model.

ThermoFluor assays

For protein stability measurement by ThermoFluor assay, purified proteins were diluted to 0.1

mg/mL in a buffer containing 20 mM Tris-HCl pH 7.5, 300 mM NaCl, 10% glycerol, and 1 mM

DTT. 0.5 μL of SPYRO Orange dye (5,000X concentrate; Thermo Fisher Scientific #S6650) was

mixed with 49.5 μL protein solution, and the mixture was sealed in an optically clear PCR plate.

In a Bio-Rad CFX Connect real-time PCR system, fluorescence was measured in scan mode

"FRET" during a 25°-99° temperature ramp (0.5° intervals, 30 second hold per interval). For

analysis, the change in fluorescence per step was plotted against temperature, and the peak of

this plot (the representing the inflection point in the upward slope of the raw fluorescence curve)

was taken as the melting temperature (Tm).

Protein crystallization and structure determination

For crystallization, HIM-3(R93Y) was concentrated to 15-20 mg/mL in crystallization buffer (20

mM HEPES pH 7.5, 100 mM NaCl, 1 mM DTT) and mixed 1:1 with well solution containing 1.4-

1.6 M sodium malonate pH 6.5-7.0. Large crystals grown over the course of two weeks were

cryoprotected in 2.5 M (total) sodium malonate and flash-frozen in liquid nitrogen. Diffraction

data were collected at the Advanced Light source beamline 8.3.1 (support statement below).

Data were indexed and reduced with XDS [87], scaled with AIMLESS [88], and converted to

structure factors with TRUNCATE [89] (**Table S1**). The structure was determined by molecular

replacement in PHASER using the structure of HIM-3 (PDB ID 4TRK) [19] as a search model.

An initial model was rebuilt in COOT [90] and refined in phenix.refine [91] using positional and

individual B-factor (anisotropic for protein atoms, isotropic for water molecules) refinement with

riding hydrogens. Structure figures were generated in PyMOL version 2.0 (Schrödinger, LLC).

ALS Beamline 8.3.1 support statement

The Berkeley Center for Structural Biology is supported in part by the Howard Hughes Medical

Institute. The Advanced Light Source is a Department of Energy Office of Science User Facility

under Contract No. DE-AC02-05CH11231. The Pilatus detector on 5.0.1. was funded under NIH

grant S10OD021832. The ALS-ENABLE beamlines are supported in part by the National

Institutes of Health, National Institute of General Medical Sciences, grant P30 GM124169.

Author contributions:

A.E.R., S.G., A.D., K.D.C., and N.B. designed the experiments. A.E.R., S.G., A.D., M.M., V.O.,

and K.E. performed the experiments. A.E.R., S.G., A.D., M.M., V.O., K.E., K.D.C., and N.B.

analyzed the data. A.E.R and N.B. wrote the initial draft of the manuscript and A.E.R., S.G.,

A.D., M.M., V.O., K.E., K.D.C., and N.B. revised the manuscript. K.D.C., and N.B. acquired

funding.

Acknowledgements:

We would like to thank Abby Dernburg, Monique Zetka and Anne Villeneuve for valuable strains

and reagents. We would also like to thank the members of the Bhalla lab as well as Brandt

Warecki and Alice Devigne for careful review of the manuscript. This work was supported by the

NIH (grant number R35GM141835 [N.B.], T32GM008646 [A.E.R] and R35GM144121 [K.D.C.]).

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440).

References:

1.   Hassold T, Hunt P. To err (meiotically) is human: the genesis of human aneuploidy. Nat Rev Genet. 2001;2(4):280-91. doi: 10.1038/35066065. PubMed PMID: 11283700.

2.   Deshong AJ, Ye AL, Lamelza P, Bhalla N. A quality control mechanism coordinates meiotic prophase events to promote crossover assurance. PLoS Genet. 2014;10(4):e1004291. doi: 10.1371/journal.pgen.1004291. PubMed PMID: 24762417; PubMed Central PMCID: PMCPMC3998905.

3.   Giacopazzi S, Vong D, Devigne A, Bhalla N. PCH-2 collaborates with CMT-1 to proofread meiotic homolog interactions. PLoS Genet. 2020;16(7):e1008904. doi: 10.1371/journal.pgen.1008904. PubMed PMID: 32730253; PubMed Central PMCID: PMCPMC7433886.

4.   Aravind L, Koonin EV. The HORMA domain: a common structural denominator in mitotic checkpoints, chromosome synapsis and DNA repair. Trends Biochem Sci. 1998;23(8):284-6. PubMed PMID: 9757827.

5.   Ur SN, Corbett KD. Architecture and Dynamics of Meiotic Chromosomes. Annu Rev

Genet. 2021;55:497-526. Epub 20210916. doi: 10.1146/annurev-genet-071719-020235.

PubMed PMID: 34530636.

6.   Hollingsworth NM, Goetsch L, Byers B. The HOP1 gene encodes a meiosis-specific

component of yeast chromosomes. Cell. 1990;61(1):73-84. PubMed PMID: 2107981.

7.   Lorenz A, Wells JL, Pryce DW, Novatchkova M, Eisenhaber F, McFarlane RJ, et al. S.

pombe meiotic linear elements contain proteins related to synaptonemal complex components.

J Cell Sci. 2004;117(Pt 15):3343-51. doi: 10.1242/jcs.01203. PubMed PMID: 15226405.

8.   Fukuda T, Daniel K, Wojtasz L, Toth A, Hoog C. A novel mammalian HORMA domain-

containing protein, HORMAD1, preferentially associates with unsynapsed meiotic

chromosomes. Exp Cell Res. 2010;316(2):158-71. doi: 10.1016/j.yexcr.2009.08.007. PubMed

PMID: 19686734.

9.   Caryl AP, Armstrong SJ, Jones GH, Franklin FC. A homologue of the yeast HOP1 gene

is inactivated in the Arabidopsis meiotic mutant asy1. Chromosoma. 2000;109(1-2):62-71. doi:

10.1007/s004120050413. PubMed PMID: 10855496.

10.   Wojtasz L, Daniel K, Roig I, Bolcun-Filas E, Xu H, Boonsanay V, et al. Mouse

HORMAD1 and HORMAD2, two conserved meiotic chromosomal proteins, are depleted from

synapsed chromosome axes with the help of TRIP13 AAA-ATPase. PLoS Genet.

2009;5(10):e1000702. doi: 10.1371/journal.pgen.1000702. PubMed PMID: 19851446; PubMed

Central PMCID: PMCPMC2758600.

11.   Martinez-Perez E, Villeneuve AM. HTP-1-dependent constraints coordinate homolog

pairing and synapsis and promote chiasma formation during C. elegans meiosis. Genes Dev.

2005;19(22):2727-43. doi: 10.1101/gad.1338505. PubMed PMID: 16291646; PubMed Central

PMCID: PMCPMC1283965.

12.   Goodyer W, Kaitna S, Couteau F, Ward JD, Boulton SJ, Zetka M. HTP-3 links DSB

formation with homolog pairing and crossing over during C. elegans meiosis. Dev Cell.

2008;14(2):263-74. doi: 10.1016/j.devcel.2007.11.016. PubMed PMID: 18267094.

13.   Couteau F, Zetka M. HTP-1 coordinates synaptonemal complex assembly with homolog

alignment during meiosis in C. elegans. Genes Dev. 2005;19(22):2744-56. doi:

10.1101/gad.1348205. PubMed PMID: 16291647; PubMed Central PMCID: PMCPMC1283966.

14.   Zetka MC, Kawasaki I, Strome S, Muller F. Synapsis and chiasma formation in

Caenorhabditis elegans require HIM-3, a meiotic chromosome core component that functions in

chromosome segregation. Genes Dev. 1999;13(17):2258-70. PubMed PMID: 10485848;

PubMed Central PMCID: PMC317003.

15.   Daniel K, Lange J, Hached K, Fu J, Anastassiadis K, Roig I, et al. Meiotic homologue

alignment and its quality surveillance are controlled by mouse HORMAD1. Nat Cell Biol.

2011;13(5):599-610. doi: 10.1038/ncb2213. PubMed PMID: 21478856; PubMed Central

PMCID: PMCPMC3087846.

16.   Kogo H, Tsutsumi M, Ohye T, Inagaki H, Abe T, Kurahashi H. HORMAD1-dependent

checkpoint/surveillance mechanism eliminates asynaptic oocytes. Genes Cells. 2012;17(6):439-

54. doi: 10.1111/j.1365-2443.2012.01600.x. PubMed PMID: 22530760.

17.   Shin YH, Choi Y, Erdin SU, Yatsenko SA, Kloc M, Yang F, et al. Hormad1 mutation

disrupts synaptonemal complex formation, recombination, and chromosome segregation in

mammalian meiosis. PLoS Genet. 2010;6(11):e1001190. doi: 10.1371/journal.pgen.1001190.

PubMed PMID: 21079677; PubMed Central PMCID: PMCPMC2973818.

18.   Wojtasz L, Cloutier JM, Baumann M, Daniel K, Varga J, Fu J, et al. Meiotic DNA double-

strand breaks and chromosome asynapsis in mice are monitored by distinct HORMAD2-

independent and -dependent mechanisms. Genes Dev. 2012;26(9):958-73. doi:

10.1101/gad.187559.112. PubMed PMID: 22549958; PubMed Central PMCID:

PMCPMC3347793.

19.   Kim Y, Rosenberg SC, Kugel CL, Kostow N, Rog O, Davydov V, et al. The chromosome

axis controls meiotic events through a hierarchical assembly of HORMA domain proteins. Dev

Cell. 2014;31(4):487-502. doi: 10.1016/j.devcel.2014.09.013. PubMed PMID: 25446517;

PubMed Central PMCID: PMCPMC4254552.

20.   Severson AF, Ling L, van Zuylen V, Meyer BJ. The axial element protein HTP-3

promotes cohesin loading and meiotic axis assembly in C. elegans to implement the meiotic

program of chromosome segregation. Genes Dev. 2009;23(15):1763-78. doi:

10.1101/gad.1808809. PubMed PMID: 19574299; PubMed Central PMCID: PMCPMC2720254.

21.   Couteau F, Nabeshima K, Villeneuve A, Zetka M. A component of C. elegans meiotic

chromosome axes at the interface of homolog alignment, synapsis, nuclear reorganization, and

recombination. Curr Biol. 2004;14(7):585-92. doi: 10.1016/j.cub.2004.03.033. PubMed PMID:

15062099.

22.   Martinez-Perez E, Schvarzstein M, Barroso C, Lightfoot J, Dernburg AF, Villeneuve AM.

Crossovers trigger a remodeling of meiotic chromosome axis composition that is linked to two-

step loss of sister chromatid cohesion. Genes Dev. 2008;22(20):2886-901. doi:

10.1101/gad.1694108. PubMed PMID: 18923085; PubMed Central PMCID: PMCPMC2569886.

23.   Rosenberg SC, Corbett KD. The multifaceted roles of the HORMA domain in cellular

signaling. J Cell Biol. 2015;211(4):745-55. doi: 10.1083/jcb.201509076. PubMed PMID:

26598612; PubMed Central PMCID: PMCPMC4657174.

24.   Ye Q, Rosenberg SC, Moeller A, Speir JA, Su TY, Corbett KD. TRIP13 is a protein-

remodeling AAA+ ATPase that catalyzes MAD2 conformation switching. eLife. 2015;4. Epub

2015/04/29. doi: 10.7554/eLife.07367. PubMed PMID: 25918846; PubMed Central PMCID:

PMC4439613.

25.   Clairmont CS, Sarangi P, Ponnienselvan K, Galli LD, Csete I, Moreau L, et al. TRIP13

regulates DNA repair pathway choice through REV7 conformational change. Nat Cell Biol.

2020;22(1):87-96. doi: 10.1038/s41556-019-0442-y. PubMed PMID: 31915374; PubMed Central

PMCID: PMCPMC7336368.

26.   Alfieri C, Chang L, Barford D. Mechanism for remodelling of the cell cycle checkpoint

protein MAD2 by the ATPase TRIP13. Nature. 2018;559(7713):274-8. doi: 10.1038/s41586-

018-0281-1. PubMed PMID: 29973720; PubMed Central PMCID: PMCPMC6057611.

27.   Ye Q, Kim DH, Dereli I, Rosenberg SC, Hagemann G, Herzog F, et al. The AAA+

ATPase TRIP13 remodels HORMA domains through N-terminal engagement and unfolding.

EMBO J. 2017. doi: 10.15252/embj.201797291. PubMed PMID: 28659378.

28.   West AMV, Komives EA, Corbett KD. Conformational dynamics of the Hop1 HORMA

domain reveal a common mechanism with the spindle checkpoint protein Mad2. Nucleic Acids

Res. 2017. doi: 10.1093/nar/gkx1196. PubMed PMID: 29186573.

29.   Herruzo E, Lago-Maciel A, Baztan S, Santos B, Carballo JA, San-Segundo PA. Pch2

orchestrates the meiotic recombination checkpoint from the cytoplasm. PLoS Genet.

2021;17(7):e1009560. doi: 10.1371/journal.pgen.1009560. PubMed PMID: 34260586; PubMed

Central PMCID: PMCPMC8312941.

30.   Yang C, Hu B, Portheine SM, Chuenban P, Schnittger A. State changes of the HORMA

protein ASY1 are mediated by an interplay between its closure motif and PCH2. Nucleic Acids

Res. 2020;48(20):11521-35. doi: 10.1093/nar/gkaa527. PubMed PMID: 32558910; PubMed

Central PMCID: PMCPMC7672429.

31.   Chen C, Jomaa A, Ortega J, Alani EE. Pch2 is a hexameric ring ATPase that remodels

the chromosome axis protein Hop1. Proc Natl Acad Sci U S A. 2014;111(1):E44-53. doi:

10.1073/pnas.1310755111. PubMed PMID: 24367111; PubMed Central PMCID:

PMCPMC3890899.

32.   Raina VB, Vader G. Homeostatic Control of Meiotic Prophase Checkpoint Function by

Pch2 and Hop1. Curr Biol. 2020;30(22):4413-24.e5. Epub 20200910. doi:

10.1016/j.cub.2020.08.064. PubMed PMID: 32916108.

33.   Borner GV, Barot A, Kleckner N. Yeast Pch2 promotes domainal axis organization,

timely recombination progression, and arrest of defective recombinosomes during meiosis. Proc

Natl Acad Sci U S A. 2008;105(9):3327-32. doi: 10.1073/pnas.0711864105. PubMed PMID:

18305165; PubMed Central PMCID: PMCPMC2265181.

34.   Joshi N, Barot A, Jamison C, Borner GV. Pch2 links chromosome axis remodeling at

future crossover sites and crossover distribution during yeast meiosis. PLoS Genet.

2009;5(7):e1000557. doi: 10.1371/journal.pgen.1000557. PubMed PMID: 19629172; PubMed

Central PMCID: PMCPMC2708914.

35.   Lambing C, Osman K, Nuntasoontorn K, West A, Higgins JD, Copenhaver GP, et al.

Arabidopsis PCH2 Mediates Meiotic Chromosome Remodeling and Maturation of Crossovers.

PLoS Genet. 2015;11(7):e1005372. doi: 10.1371/journal.pgen.1005372. PubMed PMID:

26182244; PubMed Central PMCID: PMCPMC4504720.

36.   MacQueen AJ, Phillips CM, Bhalla N, Weiser P, Villeneuve AM, Dernburg AF.

Chromosome sites play dual roles to establish homologous synapsis during meiosis in C.

elegans. Cell. 2005;123(6):1037-50. doi: 10.1016/j.cell.2005.09.034. PubMed PMID: 16360034;

PubMed Central PMCID: PMCPMC4435800.

37.   Cuacos M, Lambing C, Pachon-Penalba M, Osman K, Armstrong SJ, Henderson IR, et

al. Meiotic chromosome axis remodelling is critical for meiotic recombination in Brassica rapa. J

Exp Bot. 2021;72(8):3012-27. doi: 10.1093/jxb/erab035. PubMed PMID: 33502451; PubMed

Central PMCID: PMCPMC8023211.

38.   San-Segundo PA, Roeder GS. Pch2 links chromatin silencing to meiotic checkpoint

control. Cell. 1999;97(3):313-24. PubMed PMID: 10319812.

39.   Herruzo E, Ontoso D, Gonzalez-Arranz S, Cavero S, Lechuga A, San-Segundo PA. The

Pch2 AAA+ ATPase promotes phosphorylation of the Hop1 meiotic checkpoint adaptor in

response to synaptonemal complex defects. Nucleic Acids Res. 2016;44(16):7722-41. doi:

10.1093/nar/gkw506. PubMed PMID: 27257060; PubMed Central PMCID: PMCPMC5027488.

40.   MacQueen AJ, Colaiacovo MP, McDonald K, Villeneuve AM. Synapsis-dependent and -

independent mechanisms stabilize homolog pairing during meiotic prophase in C. elegans.

Genes Dev. 2002;16(18):2428-42. PubMed PMID: 12231631.

41.   Carlton PM. Three-dimensional structured illumination microscopy and its application to

chromosome structure. Chromosome Res. 2008;16(3):351-65. doi: 10.1007/s10577-008-1231-

9. PubMed PMID: 18461477.

42.   Couteau F, Zetka M. DNA damage during meiosis induces chromatin remodeling and

synaptonemal complex disassembly. Dev Cell. 2011;20(3):353-63. doi:

10.1016/j.devcel.2011.01.015. PubMed PMID: 21397846.

43.   Bohr T, Nelson CR, Giacopazzi S, Lamelza P, Bhalla N. Shugoshin Is Essential for

Meiotic Prophase Checkpoints in C. elegans. Curr Biol. 2018;28(20):3199-211.e3. Epub

20181004. doi: 10.1016/j.cub.2018.08.026. PubMed PMID: 30293721; PubMed Central PMCID:

PMCPMC6200582.

44.   Phillips CM, Wong C, Bhalla N, Carlton PM, Weiser P, Meneely PM, et al. HIM-8 binds

to the X chromosome pairing center and mediates chromosome-specific meiotic synapsis. Cell.

2005;123(6):1051-63. doi: 10.1016/j.cell.2005.09.035. PubMed PMID: 16360035; PubMed

Central PMCID: PMCPMC4435792.

45.   Villeneuve AM. A cis-acting locus that promotes crossing over between X chromosomes

in Caenorhabditis elegans. Genetics. 1994;136(3):887-902. PubMed PMID: 8005443; PubMed

Central PMCID: PMCPMC1205894.

46.   Colaiacovo MP, MacQueen AJ, Martinez-Perez E, McDonald K, Adamo A, La Volpe A,

et al. Synaptonemal complex assembly in C. elegans is dispensable for loading strand-

exchange proteins but critical for proper completion of recombination. Dev Cell. 2003;5(3):463-

74. PubMed PMID: 12967565.

47.   Yokoo R, Zawadzki KA, Nabeshima K, Drake M, Arur S, Villeneuve AM. COSA-1 reveals

robust homeostasis and separable licensing and reinforcement steps governing meiotic

crossovers. Cell. 2012;149(1):75-87. doi: 10.1016/j.cell.2012.01.052. PubMed PMID: 22464324;

PubMed Central PMCID: PMCPMC3339199.

48.   Bhalla N, Dernburg AF. A conserved checkpoint monitors meiotic chromosome synapsis

in Caenorhabditis elegans. Science. 2005;310(5754):1683-6. doi: 10.1126/science.1117468.

PubMed PMID: 16339446.

49.   Stamper EL, Rodenbusch SE, Rosu S, Ahringer J, Villeneuve AM, Dernburg AF.

Identification of DSB-1, a protein required for initiation of meiotic recombination in

Caenorhabditis elegans, illuminates a crossover assurance checkpoint. PLoS Genet.

2013;9(8):e1003679. doi: 10.1371/journal.pgen.1003679. PubMed PMID: 23990794; PubMed

Central PMCID: PMCPMC3749324.

50.   Kim Y, Kostow N, Dernburg AF. The Chromosome Axis Mediates Feedback Control of

CHK-2 to Ensure Crossover Formation in C. elegans. Dev Cell. 2015;35(2):247-61. doi:

10.1016/j.devcel.2015.09.021. PubMed PMID: 26506311; PubMed Central PMCID:

PMCPMC4624198.

51.   Rosu S, Zawadzki KA, Stamper EL, Libuda DE, Reese AL, Dernburg AF, et al. The C.

elegans DSB-2 protein reveals a regulatory network that controls competence for meiotic DSB

formation and promotes crossover assurance. PLoS Genet. 2013;9(8):e1003674. doi:

10.1371/journal.pgen.1003674. PubMed PMID: 23950729; PubMed Central PMCID:

PMCPMC3738457.

52.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable

generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst

Biol. 2011;7:539. Epub 20111011. doi: 10.1038/msb.2011.75. PubMed PMID: 21988835;

PubMed Central PMCID: PMCPMC3261699.

53.   Silva N, Ferrandiz N, Barroso C, Tognetti S, Lightfoot J, Telecan O, et al. The fidelity of

synaptonemal complex assembly is regulated by a signaling mechanism that controls early

meiotic progression. Dev Cell. 2014;31(4):503-11. doi: 10.1016/j.devcel.2014.10.001. PubMed

PMID: 25455309.

54.   Martini E, Diaz RL, Hunter N, Keeney S. Crossover homeostasis in yeast meiosis. Cell.

2006;126(2):285-95. doi: 10.1016/j.cell.2006.05.044. PubMed PMID: 16873061; PubMed

Central PMCID: PMCPMC1949389.

55.   Bohr T, Ashley G, Eggleston E, Firestone K, Bhalla N. Synaptonemal Complex

Components Are Required for Meiotic Checkpoint Function in Caenorhabditis elegans.

Genetics. 2016;204(3):987-97. doi: 10.1534/genetics.116.191494. PubMed PMID: 27605049;

PubMed Central PMCID: PMCPMC5105873.

56.   West AMV, Komives EA, Corbett KD. Conformational dynamics of the Hop1 HORMA

domain reveal a common mechanism with the spindle checkpoint protein Mad2. Nucleic Acids

Res. 2018;46(1):279-92. doi: 10.1093/nar/gkx1196. PubMed PMID: 29186573; PubMed Central

PMCID: PMCPMC5758881.

57.   Ma HT, Poon RY. TRIP13 Regulates Both the Activation and Inactivation of the Spindle-

Assembly Checkpoint. Cell Rep. 2016;14(5):1086-99. doi: 10.1016/j.celrep.2016.01.001.

PubMed PMID: 26832417.

58.   Ma HT, Poon RYC. TRIP13 Functions in the Establishment of the Spindle Assembly

Checkpoint by Replenishing O-MAD2. Cell Rep. 2018;22(6):1439-50. doi:

10.1016/j.celrep.2018.01.027. PubMed PMID: 29425500.

59.   Nelson CR, Hwang T, Chen PH, Bhalla N. TRIP13PCH-2 promotes Mad2 localization to

unattached kinetochores in the spindle checkpoint response. J Cell Biol. 2015;211(3):503-16.

doi: 10.1083/jcb.201505114. PubMed PMID: 26527744; PubMed Central PMCID:

PMCPMC4639874.

60.   Eytan E, Wang K, Miniowitz-Shemtov S, Sitry-Shevah D, Kaisari S, Yen TJ, et al.

Disassembly of mitotic checkpoint complexes by the joint action of the AAA-ATPase TRIP13

and p31(comet). Proc Natl Acad Sci U S A. 2014;111(33):12019-24. doi:

10.1073/pnas.1412901111. PubMed PMID: 25092294; PubMed Central PMCID:

PMCPMC4142996.

61.   Wang K, Sturt-Gillespie B, Hittle JC, Macdonald D, Chan GK, Yen TJ, et al. Thyroid

hormone receptor interacting protein 13 (TRIP13) AAA-ATPase is a novel mitotic checkpoint-

silencing protein. J Biol Chem. 2014;289(34):23928-37. doi: 10.1074/jbc.M114.585315. PubMed

PMID: 25012665; PubMed Central PMCID: PMCPMC4156041.

62.   Defachelles L, Russo AE, Nelson CR, Bhalla N. The conserved AAA-ATPase PCH-2

(TRIP13) regulates spindle checkpoint strength. Mol Biol Cell. 2020;31(20):2219-33. doi:

10.1091/mbc.E20-05-0310. PubMed PMID: 32697629; PubMed Central PMCID:

PMCPMC7550697.

63.   Sanchez-Moran E, Santos JL, Jones GH, Franklin FC. ASY1 mediates AtDMC1-

dependent interhomolog recombination during meiosis in Arabidopsis. Genes Dev.

2007;21(17):2220-33. doi: 10.1101/gad.439007. PubMed PMID: 17785529; PubMed Central

PMCID: PMCPMC1950860.

64.   Raina VB, Vader G. Homeostatic Control of Meiotic Prophase Checkpoint Function by

Pch2 and Hop1. Curr Biol. 2020. doi: 10.1016/j.cub.2020.08.064. PubMed PMID: 32916108.

65.   Subramanian VV, MacQueen AJ, Vader G, Shinohara M, Sanchez A, Borde V, et al.

Chromosome Synapsis Alleviates Mek1-Dependent Suppression of Meiotic DNA Repair. PLoS

Biol. 2016;14(2):e1002369. doi: 10.1371/journal.pbio.1002369. PubMed PMID: 26870961;

PubMed Central PMCID: PMCPMC4752329.

66.   Dernburg AF, McDonald K, Moulder G, Barstead R, Dresser M, Villeneuve AM. Meiotic

recombination in C. elegans initiates by a conserved mechanism and is dispensable for

homologous chromosome synapsis. Cell. 1998;94(3):387-98. PubMed PMID: 9708740.

67.   Capilla-Perez L, Durand S, Hurel A, Lian Q, Chambon A, Taochy C, et al. The

synaptonemal complex imposes crossover interference and heterochiasmy in Arabidopsis. Proc

Natl Acad Sci U S A. 2021;118(12). doi: 10.1073/pnas.2023613118. PubMed PMID: 33723072;

PubMed Central PMCID: PMCPMC8000504.

68.   Cole F, Kauppi L, Lange J, Roig I, Wang R, Keeney S, et al. Homeostatic control of

recombination is implemented progressively in mouse meiosis. Nat Cell Biol. 2012;14(4):424-

30. doi: 10.1038/ncb2451. PubMed PMID: 22388890; PubMed Central PMCID:

PMCPMC3319518.

69.   Joshi N, Brown MS, Bishop DK, Borner GV. Gradual implementation of the meiotic

recombination program via checkpoint pathways controlled by global DSB levels. Mol Cell.

2015;57(5):797-811. doi: 10.1016/j.molcel.2014.12.027. PubMed PMID: 25661491; PubMed

Central PMCID: PMCPMC4392720.

70.   Morgan C, Fozard JA, Hartley M, Henderson IR, Bomblies K, Howard M. Diffusion-

mediated HEI10 coarsening can explain meiotic crossover positioning in Arabidopsis. Nat

Commun. 2021;12(1):4674. doi: 10.1038/s41467-021-24827-w. PubMed PMID: 34344879;

PubMed Central PMCID: PMCPMC8333306.

71.   Bhalla N. PCH-2 and meiotic HORMADs: A module for evolutionary innovation in

meiosis? Current Topics in Developmental Biology. 2022;in press. doi:

https://doi.org/10.1016/bs.ctdb.2022.07.001.

72.   Das D, Chen SY, Arur S. ERK phosphorylates chromosomal axis component HORMA

domain protein HTP-1 to regulate oocyte numbers. Sci Adv. 2020;6(44). Epub 20201030. doi:

10.1126/sciadv.abc5580. PubMed PMID: 33127680; PubMed Central PMCID:

PMCPMC7608811.

73.   Shin YH, McGuire MM, Rajkovic A. Mouse HORMAD1 is a meiosis i checkpoint protein

that modulates DNA double- strand break repair during female meiosis. Biol Reprod.

2013;89(2):29. doi: 10.1095/biolreprod.112.106773. PubMed PMID: 23759310; PubMed Central

PMCID: PMCPMC4076362.

74.   Kogo H, Tsutsumi M, Inagaki H, Ohye T, Kiyonari H, Kurahashi H. HORMAD2 is

essential for synapsis surveillance during meiotic prophase via the recruitment of ATR activity.

Genes Cells. 2012;17(11):897-912. doi: 10.1111/gtc.12005. PubMed PMID: 23039116.

75.   Baudat F, Manova K, Yuen JP, Jasin M, Keeney S. Chromosome synapsis defects and

sexually dimorphic meiotic progression in mice lacking Spo11. Mol Cell. 2000;6(5):989-98.

PubMed PMID: 11106739.

76.   Giroux CN, Dresser ME, Tiano HF. Genetic control of chromosome synapsis in yeast

meiosis. Genome. 1989;31(1):88-94. PubMed PMID: 2687110.

77.   Grelon M, Vezon D, Gendrot G, Pelletier G. AtSPO11-1 is necessary for efficient meiotic

recombination in plants. EMBO J. 2001;20(3):589-600. doi: 10.1093/emboj/20.3.589. PubMed

PMID: 11157765; PubMed Central PMCID: PMCPMC133473.

78.   Romanienko PJ, Camerini-Otero RD. The mouse Spo11 gene is required for meiotic

chromosome synapsis. Mol Cell. 2000;6(5):975-87. PubMed PMID: 11106738.

79.   Heldrich J, Milano CR, Markowitz TE, Ur SN, Vale-Silva LA, Corbett KD, et al. Two

pathways drive meiotic chromosome axis assembly in Saccharomyces cerevisiae. Nucleic Acids

Res. 2022;50(8):4545-56. doi: 10.1093/nar/gkac227. PubMed PMID: 35412621; PubMed

Central PMCID: PMCPMC9071447.

80.   Subramanian VV, Zhu X, Markowitz TE, Vale-Silva LA, San-Segundo PA, Hollingsworth

NM, et al. Persistent DNA-break potential near telomeres increases initiation of meiotic

recombination on short chromosomes. Nat Commun. 2019;10(1):970. doi: 10.1038/s41467-019-

08875-x. PubMed PMID: 30814509; PubMed Central PMCID: PMCPMC6393486.

81.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94.

PubMed PMID: 4366476; PubMed Central PMCID: PMCPMC1213120.

82.   Arribere JA, Bell RT, Fu BX, Artiles KL, Hartman PS, Fire AZ. Efficient marker-free

recovery of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans.

Genetics. 2014;198(3):837-46. doi: 10.1534/genetics.114.169730. PubMed PMID: 25161212;

PubMed Central PMCID: PMCPMC4224173.

83.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag;

2016.

84.   Toraason E, Adler VL, Kurhanewicz NA, DiNardo A, Saunders AM, Cahoon CK, et al.

Automated and customizable quantitative image analysis of whole Caenorhabditis elegans

germlines. Genetics. 2021;217(3). doi: 10.1093/genetics/iyab010. PubMed PMID: 33772283;

PubMed Central PMCID: PMCPMC8045727.

85. Preibisch S, Saalfeld S, Tomancak P. Globally optimal stitching of tiled 3D microscopic

image acquisitions. Bioinformatics. 2009;25(11):1463-5. Epub 20090403. doi:

10.1093/bioinformatics/btp184. PubMed PMID: 19346324; PubMed Central PMCID:

PMCPMC2682522.

86. Tropea JE, Cherry S, Waugh DS. Expression and purification of soluble His(6)-tagged

TEV protease. Methods in Molecular Biology. 498. Totowa, NJ: Humana Press; 2009. p. 297-

307.

87. Kabsch W. XDS. Acta crystallographica Section D, Biological crystallography.

2010;66:125-32. doi: 10.1107/S0907444909047337. PubMed PMID: 20124692.

88. Evans PR, Murshudov GN. How good are my data and what is the resolution? Acta

crystallographica Section D, Biological crystallography. 2013;69:1204-14. doi:

10.1107/S0907444913000061. PubMed PMID: 23793146.

89. Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, et al. Overview of

the CCP4 suite and current developments. Acta crystallographica Section D, Biological

crystallography. 2011;67:235-42. doi: 10.1107/S0907444910045749. PubMed PMID: 21460441.

90. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta

crystallographica Section D, Biological crystallography. 2010;66:486-501. doi:

10.1107/S0907444910007493. PubMed PMID: 20383002.

91. Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, et al. PHENIX: a

comprehensive Python-based system for macromolecular structure solution. Acta

crystallographica Section D, Biological crystallography. 2010;66:213-21. doi:

10.1107/S0907444909052925.

92.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold

Protein Structure Database: massively expanding the structural coverage of protein-sequence

space with high-accuracy models. Nucleic Acids Res. 2022;50(D1):D439-D44. doi:

10.1093/nar/gkab1061. PubMed PMID: 34791371; PubMed Central PMCID: PMCPMC8728224.

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317   **<u>Figure Legends</u>**

1318

Figure 1. PCH-2's localization is consistent with remodeling HORMADs on meiotic axes.

A. Cartoon of synaptonemal complex with axial elements made up of meiotic HORMADs and the central element component, SYP-1. **B.** Structured illumination microscopy (SIM) images of meiotic nuclei stained with antibodies against PCH-2 (green) and SYP-1 (magenta). Arrows represent regions of PCH-2 potentially associated with meiotic axes. Scalebar indicates 5 microns. **C.** Cropped insets of PCH-2 and SYP-1 signal. Yellow line represents area of pixel intensities measured. **D**. Line scans of PCH-2 (green) and SYP-1 (magenta) pixel intensities plotted against distance in pixels.

Figure 2. HORMA domain structure of meiotic HORMADs in *C. elegans.* A. Cartoon depicting protein domains of HTP-3, HIM-3, HTP-1, and HTP-2. HORMA domains and closure motifs are shown as colored boxes and labeled. Mutations generated by CRISPR/Cas9 are shown. **B.** Clustal Omega alignment of the HORMA domains of HTP-3, HIM-3, HTP-1, and HTP-2. Asterisks represent conserved residues, : represents conservative changes and . represents semi-conservative changes. Red box indicates targeted reside for mutation for each respective HORMAD. **C.** Structure of the *C. elegans* HTP-3 HORMA domain (pink) bound to a closure motif (gray), predicted by AlphaFold 2 [92]. Right panels show the position of H96. **D.** Overlay of wild-type (gray; PDB ID 4TRK) and R93Y mutant (green/pink) of *C. elegans* HIM-3. Right panels show the position of residue 93 and show that the overall HORMA domain structure is not impacted upon mutation of arginine 93 to tyrosine. **E.** View of the *C. elegans* HTP-1 HORMA domain (PDB ID 4TZQ) equivalent to the close-up panel in (D), showing the position of residue G97 (yellow spheres), which is partially buried against a neighboring alpha-helix.

Figure 3. *htp-3*^{H96Y} suppresses the acceleration of pairing and synapsis but exacerbates

defects in DNA repair and crossover formation in *pch-2* mutants. A. Cartoon depicting

zones for quantification in the *C. elegans* germline. **B.** Meiotic nuclei stained with DAPI

(magenta) and antibody against HIM-8 (green). **C.** Quantification of percent nuclei with paired

HIM-8 signals in zones 1-6 for *syp-1* (blue), *pch-2;syp-1* (red), *htp-3*^{H96Y}*;syp-1* (green) and *pch-*

2;htp-3^{H96Y}*;syp-*1 (purple) mutant strains. Error bars represent 95% confidence intervals. **D.**

Meiotic nuclei stained with antibodies against SYP-1(magenta) and HTP-3 (green). Arrows

indicates unsynapsed chromosomes. **E.** Quantification of percent nuclei with complete synapsis

in zones 1-6 for wildtype (blue), *pch-2* (red), *htp-3*^{H96Y} (green) and *pch-2;htp-3*^{H96Y} (purple)

mutant strains. Error bars represent 95% confidence intervals. **F.** Quantification of percent

nuclei with complete synapsis in zones 1-6 for *meDf2/+* (blue), *pch-2;meDf2/+* (red), *htp-*

3^{H96Y}*;meDf2/+* (green) and *pch-2;htp-3*^{H96Y}*;meDf2/+* (purple) mutant strains. Error bars represent

95% confidence intervals. **G.** Meiotic nuclei stained with DAPI (magenta) and antibody against

RAD-51 (green). **H.** Quantification of average number of RAD-51 foci in zones 1-6 for wildtype

(blue), *pch-2* (red), *htp-3*^{H96Y} (green) and *pch-2;htp-3*^{H96Y} (purple) mutant strains. Error bars

represent standard error of the mean (SEM). **I.** Meiotic nuclei stained with DNA (magenta) and

antibody against GFP::COSA-1 (green). Left indicates representative nuclei with 6 GFP::COSA-

1 foci. Right indicates representative nuclei with 5 GFP::COSA-1 foci. **J.** Quantification of

percent nuclei with 5 GFP::COSA-1 foci for wildtype, *pch-2*, *htp-3*^{H96Y}, and *pch-2;htp-3*^{H96Y}

mutant strains. Error bars represent 95% confidence intervals. In all images, scalebar indicates

5 microns. In all graphs, ** indicates p-value < 0.01 and **** indicated p-value <0.0001,

Significance was assessed using two-tailed Fisher's exact-tests.

1365

Figure 4. *him-3*^{R93Y} does not suppress the acceleration of pairing, synapsis, or DNA

repair, but does suppress defects in crossover formation in *pch-2* mutants. A. Meiotic

nuclei stained with DAPI (magenta) and antibody against HIM-8 (green). **B.** Quantification of

percent nuclei with paired HIM-8 signals in zones 1-6 for *syp-1* (blue), *pch-2;syp-1* (red), *him-*

3^{R93Y};syp-1 (green) and *pch-2;him-3^{R93Y};syp*-1 (purple) mutant strains. Error bars represent 95%

confidence intervals. **C.** Meiotic nuclei stained with antibodies against SYP-1(magenta) and

HTP-3 (green). Arrow indicates unsynapsed chromosomes. **D.** Quantification of percent nuclei

with complete synapsis in zones 1-6 for wildtype (blue), *pch-2* (red), *him-3^{R93Y}* (green) and *pch-*

2;him-3^{R93Y} (purple) mutant strains. Error bars represent 95% confidence intervals. **E.**

Quantification of percent nuclei with complete synapsis in zones 1-6 for *meDf2/+* (blue), *pch-*

2;meDf2/+ (red), *him-3^{R93Y};meDf2/+* (green) and *pch-2;him-3^{R93Y};meDf2/+* (purple) mutant

strains. Error bars represent 95% confidence intervals. **F.** Meiotic nuclei stained with DAPI

(magenta) and antibody against RAD-51 (green). **G.** Quantification of average number of RAD-

51 foci in zones 1-6 for wildtype (blue), *pch-2* (red), *him-3^{R93Y}* (green) and *pch-2;him-3^{R93Y}*

(purple) mutant strains. Error bars represent standard error of the mean (SEM). **H.** Meiotic

nuclei stained with DNA (magenta) and antibody against GFP::COSA-1 (green). Left indicates

representative nuclei with 6 GFP::COSA-1 foci. Right indicates representative nuclei with 5

GFP::COSA-1 foci. **I.** Quantification of percent nuclei with 5 GFP::COSA-1 foci for wildtype, *pch-*

2, *him-3^{R93Y}*, and *pch-2;him-3^{R93Y}* mutant strains. Error bars represent 95% confidence intervals.

J. Oocytes stained with DAPI. Left indicates representative oocyte with zero univalents. Right

indicates representative oocyte with univalents (arrows). Scalebar indicates 5 microns. **K.**

Quantification of percent oocytes with univalents for wildtype, *pch-2*, *him-3^{R93Y}*, and *pch-2;him-*

3^{R93Y} mutant strains. Error bars represent 95% confidence intervals. In all graphs, NS indicates

non-significant, * indicates p-value < 0.05, ** indicates p-value < 0.01, *** indicates p-value <

0.001, and **** indicated p-value <0.0001. Significance was assessed using two-tailed Fisher's

exact-tests. In all images, scalebar indicates 5 microns.

1392

Figure 5. *htp-1^{G97T}* does not genetically interact with *pch-2* in the context of pairing,

synapsis, DNA repair, or crossover formation. A. Meiotic nuclei stained with DAPI (magenta)

and antibody against HIM-8 (green). **B.** Quantification of percent nuclei with paired HIM-8

signals in zones 1-6 for *syp-1* (blue), *pch-2;syp-1* (red), *htp-1^{G97T};syp-1* (green) and *pch-2;htp-*

1^{G97T};syp-1 (purple) mutant strains. Error bars represent 95% confidence intervals. **C.** Meiotic

nuclei stained with antibodies against SYP-1(magenta) and HTP-3 (green). Arrow indicates

unsynapsed chromosomes. **D.** Quantification of percent nuclei with complete synapsis in zones

1-6 for wildtype (blue), *pch-2* (red), *htp-1^{G97T}* (green) and *pch-2;htp-1^{G97T}* (purple) mutant strains.

Error bars represent 95% confidence intervals. **E.** Meiotic nuclei stained with DAPI (magenta)

and antibody against RAD-51 (green). **F.** Quantification of average number of RAD-51 foci in

zones 1-6 for wildtype (blue), *pch-2* (red), *htp-1^{G97T}* (green) and *pch-2;htp-1^{G97T}* (purple) mutant

strains. Error bars represent standard error of the mean (SEM). **G.** Meiotic nuclei stained with

DNA (magenta) and antibodies against GFP::COSA-1 (green). Left indicates representative

nuclei with 6 GFP::COSA-1 foci. Right indicates representative nuclei with 5 GFP::COSA-1 foci.

H. Quantification of distribution of GFP::COSA-1 foci per nucleus for wildtype (blue), *pch-2* (red),

htp-1^{G97T} (green) and *pch-2;htp-1^{G97T}* (purple) mutant strains. **E.** Oocytes stained with DAPI. Left

indicates representative oocyte with zero univalents. Right indicates representative oocyte with

univalents (arrows). **F.** Quantification of percent oocytes with univalents for wildtype, *pch-2*, *htp-*

1^{G97Y}, and *pch-2;htp-1^{G97Y}* mutant strains. In all graphs, NS indicates non-significant and ****

indicates p-value < 0.0001. Significance was assessed using two-tailed Fisher's exact tests. In

all images, scalebar indicates 5 microns.

1414

Figure 6. *htp-1^{G97T}* delays meiotic progression and this delay is partially suppressed by

mutation of *pch-2*. A. Full length representative germlines stained with DAPI (magenta) and

antibody against DSB-1 (green). Yellow dashed line represents region of DSB-1 positive nuclei.

Scalebar indicates 20 microns. **B.** Quantification of DSB-1 positive meiotic nuclei for wildtype,

htp-1^{G97A}, htp-1^{G97S}, htp-1^{G97T} and pch-2;htp-1^{G97T} mutant strains. NS indicates non-significant. **

indicates p-value <0.01. *** indicates p-value < 0.001. **** indicates p-value <0.0001. Statistical

significance was assessed using two-tailed Fisher's exact tests. Error bars represent 95%

confidence intervals.

1423

1424   **Figure 7. *him-3^{R93Y}* mutants have reduced affinity for closure motifs. A.** Fluorescence

1425   polarization peptide binding assay of HIM-3 (blue) and HIM-3^{R93Y} (green) and measured Kd's. **B.**

1426   Cartoon depicting proposed model for open and closed conformation switching of wildtype HIM-

1427   3 and HIM-3^{R93Y}.

1428

1429   **Figure 8. PCH-2 distributes its regulation of meiotic prophase events by remodeling**

1430   **different meiotic HORMADs.** Cartoon depicting proposed model for PCH-2's regulation of

1431   meiotic prophase events. PCH-2 regulates the progression of pairing and by acting through

1432   HTP-3, crossover recombination through HIM-3, and may regulate meiotic progression through

1433   HTP-1.

1434

Figure S1. *htp-3^{H96Y}* does not affect meiotic progression. A. Full length representative

germlines stained with DAPI (magenta) and DSB-1 (green). Yellow dashed line represents

region of DSB-1 positive nuclei. Scalebar indicates 20 microns. **B.** Quantification of DSB-1

positive meiotic nuclei for wildtype and *htp-3 ^{H96Y}* mutant strains. NS indicates non-significant. **C.**

Quantification of DSB-1 positive meiotic nuclei for *syp-1* and *htp-3 ^{H96Y};syp-1* mutants. NS

indicates non-significant. Statistical significance was assessed using two-tailed Fisher's exact

tests. Error bars represent 95% confidence intervals.

1442

1443   **Figure S2. *him-3^{R93Y}* localizes to meiotic chromosomes. Top:** Mid-pachytene nuclei from

1444   wildtype and *him-3^{R93Y}* germlines stained with DAPI (magenta) and antibody against HIM-3

1445   (green). **Bottom:** Mid-pachytene nuclei from wildtype and *him-3^{R93Y}* germlines stained with HIM-

1446   3 (white). Scalebar indicates 5 microns.

1447

Figure S3. *him-3^{R93Y}* does not affect meiotic progression. A. Full length representative

germlines stained with DAPI (magenta) and antibody against DSB-1 (green). Yellow dashed line

represents region of DSB-1 positive nuclei. Scalebar indicates 20 microns. **B.** Quantification of

DSB-1 positive meiotic nuclei for wildtype and *him-3^{R93Y}* mutant strains. NS indicates non-

significant. **C.** Quantification of DSB-1 positive meiotic nuclei for *syp-1* and *him-3^{R93Y}*;*syp-1*

mutants. NS indicates non-significant. Statistical significance was assessed using two-tailed

Fisher's exact tests.

1455

1456 **Figure S4. *htp-1^{G97T}* localizes to meiotic chromosomes. Top:** Mid-pachytene nuclei from

1457 wildtype, *htp-1^{G97T}* and *htp-1^{G97T} htp-2* germlines stained with DAPI (magenta) and antibody

1458 against HTP-1/2 (green). **Bottom:** Mid-pachytene nuclei from wildtype, *htp-1^{G97T}* and *htp-1^{G97T}*

1459 *htp-2* germlines stained with HTP-1/2 (white). Scalebar indicates 5 microns.

1460

1461 **Figure S5. HIM3^{R93Y} does not affect HIM-3's protein stability.** Stability curve for purified

1462 wildtype HIM-3 (blue) and HIM-3^{R93Y} (green).

1463

1464 **Figure S6. *him-3^{R93Y}* and *htp-1^{G97T}* mutants have functional meiotic checkpoints.**

1465 Quantification of the average number of apoptotic nuclei per germline. NS indicates non-

1466 significant and **** indicates p-value <0.0001. Statistical significance was assessed using two-

1467 tailed Student t-tests. Error bars indicate 2X standard error of the mean (SEM).

1468

1469 **Table 1. Viability assay and male counts for meiotic HORMAD mutants.** Table with percent

1470 viable progeny and males. Total number of progeny listed in parentheses.

1471

1472 **Table S1. Data collection and refinement statistics.** Table with data collection and

1473 refinement statistics for Figure 7. Explanations of equations listed below the table.

Table 1: Viability and percent male-self-progeny

genotype	percent viability (total no. of animals)	percent male self-progeny
wildtype	100% (2489)	0.22%
meDf2/+	97% (2286)	10.91%
pch-2	100% (2388)	0.71%
pch-2;meDf2/+	98.15% (1808)	0.83%
htp-3^{H96Y}	100% (2586)	0.12%
htp-3^{H96Y};meDf2/+	97.03% (1957)	11.34%
pch-2;htp-3^{H96Y}	81.89% (1067)	1.12%
pch-2;htp-3^{H96Y};meDf2/+	83.90% (1652)	10.84%
him-3^{R93Y}	100% (3304)	0.15%
him-3^{R93Y};meDf2/+	97.69% (3170)	11.30%
pch-2;him-3^{R93Y}	100% (3027)	0.03%
pch-2;him-3^{R93Y};meDf2/+	98.43% (3140)	10.34%
htp-1^{G97T}	67.29% (1984)	2.25%
pch-2;htp-1^{G97T}	58.44% (2175)	1.97%

A

HTP-3 — H96Y — 739

HIM-3 — R93Y — 291

HTP-1 — G97T — 352

HTP-2 — 352

HORMA domain   **closure motifs**

B

```
HTP-3 (57-116) VIPAEYFQDFQVYGDVSGYTLRQDIPEGRSVSSKLLASHDAIRQKLLQKLAIVVEESLN-
HIM-3 (55-114) ILDEKCFKTRTIDKLL-VTAFQSSVPAAKRVSSTFDGLRDAIQQGYLREFAIVFYKKPNE
HTP-1 (59-118) ILGQEYFTKNYITEKLKCMTLCFRNPRSNQIAQLLRNAGDATKDGFLKEVSLVITNNEGD
HTP-2 (59-118) ILEKEYFSKNYITENLSCMTLSFKNLRAHQIAQLLRAAGDATKDGFLKEISLVVTEHDGD
               ::   :  *    :    :   ::    ..::. :   ** ::  *::.::*. :   .
```

C

HTP-3 (AlphaFold)

closure motif

180°

H96

D

⬤ HIM-3 WT (4TRK)
⬤ HIM-3 R93Y

closure motif

180°

Y93
R93

E

HTP-1 (4TZQ)

G97 Cα

Figure 4

Figure 5

A

Wildtype

htp1^G97A

htp1^G97S

htp1^G97T

pch-2;htp1^G97T

DNA
DSB-1

B

% DSB-1 positive meiotic nuceli

wildtype   *htp-1^G97A*   *htp-1^G97S*   *htp-1^G97T*   *pch-2;*
                                                          htp-1^G97T

NS   ***   ****   ****   **

A

HIM-3 WT and R93Y binding HTP-3 motif #4

Protein	Peptide	Kd
HIM-3^{WT}	HTP-3 motif #4	220 +/- 20 nm
HIM-3^{WT}	Scrambled	*no binding*
HIM-3^{R93Y}	HTP-3 motif #4	690 +/- 130 nm
HIM-3^{R93Y}	Scrambled	*no binding*

B

HIM-3
wildtype closure motif

HIM-3^{R93Y}
wildtype closure motif

Table S1. Data collection and refinement statistics

	HIM-3 R93Y
Data collection	
Synchrotron/Beamline	ALS 8.3.1
Date collected	9/14/2018
Resolution (Å)	63.4-1.3
Wavelength (Å)	1.11583
Space Group	$P6_1$
Unit Cell Dimensions (a, b, c) Å	73.23, 73.23, 110.82
Unit cell Angles (α,β,γ) °	90, 90, 120
I/σ (last shell)	27.4
[1]R_{sym} (last shell)	0.055 (2.245)
[2]R_{meas} (last shell)	0.056 (2.081)
[3]$CC_{1/2}$ (last shell)	1 (0.677)
Completeness (last shell) %	100 (100)
Number of reflections	1,632,957
unique	82,280
Multiplicity (last shell)	19.8 (19.7)
Refinement	
Resolution (Å)	63.4-1.3
No. of reflections	82274
working	4255
free	78019
[4]R_{work} (last shell) (%)	14.39 (31.35)
[4]R_{free} (last shell) (%)	16.39 (32.51)
Structure/Stereochemistry	
No. of atoms	4093
solvent	252
hydrogen	1907
r.m.s.d. bond lengths (Å)	0.021
r.m.s.d. bond angles (°)	1.58
Ramachandran favored/allowed (%)	97.03%/100%
Molprobity score	1.55
[5]SBGrid Data Bank ID	8CXK
[6]Protein Data Bank ID	910

[1] $R_{sym} = \sum\sum_j |I_j - \langle I \rangle | / \sum I_j$, where I_j is the intensity measurement for reflection j and $\langle I \rangle$ is the mean intensity for multiply recorded reflections.

[2] $R_{meas} = \sum_h [\sqrt{(n/(n-1))} \sum_j [I_{hj} - \langle I_h \rangle] / \sum_{hj} \langle I_h \rangle$, where I_{hj} is a single intensity measurement for reflection h, $\langle I_h \rangle$ is the average intensity measurement for multiply recorded reflections, and n is the number of observations of reflection h.

[3] $CC_{1/2}$ is the Pearson correlation coefficient between the average measured intensities of two randomly-assigned half-sets of the measurements of each unique reflection.

[4] R_{work}, $R_{free} = \sum ||F_{obs}| - |F_{calc}|| / |F_{obs}|$, where the working and free R-factors are calculated using the working and free reflection sets, respectively.

A

Wildtype

htp-3^{H96Y}

syp-1

htp-3^{H96Y};syp-1

DNA
DSB-1

B

C

wildtype   *him-3*^{R93Y}

MERGE

HIM-3

DNA
HIM-3

A

wildtype

him-3^{R93Y}

syp-1

him-3^{R93Y};syp-1

DNA
DSB-1

B

C

wildtype   *htp-1^{G97T}*   *htp-1^{G97T} htp-2*

MERGE

HTP-1/2

DNA
HTP-1/2

A