**A Parabrachial to Hypothalamic Neurocircuit Mediates Defensive Behavior**

Fan Wang[1,2†], Yuge Chen[1,2†], Yuxin Lin[1,2$], Xuze Wang[1,2$], Kaiyuan Li[1,2], Yong Han[1,2],

Jintao Wu[1,2], Xingyi Shi[1,2], Zhenggang Zhu[1,2], Chaoying Long[1,2], Xiaojun Hu[1,3],

Shumin Duan[1,2,3,4,5,6]* and Zhihua Gao[1,2,3]*

[1]Department of Neurobiology and Department of Neurosurgery of Second Affiliated

Hospital, Zhejiang University School of Medicine, Hangzhou, P.R. China;

[2]Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, P.R. China.

[3]NHC and CAMS Key Laboratory of Medical Neurobiology, MOE Frontier Science

Center for Brain Science and Brain-Machine Integration, School of Brain Science and

Brain Medicine, Zhejiang University, Hangzhou, P.R. China;

[4]Institute for Translational Brain Research, MOE Frontiers Center for Brain Science,

Fudan University, Shanghai, P.R. China;

[5]The Institute of Brain and Cognitive Sciences, Zhejiang University City College,

Hangzhou, P.R. China;

[6]Chuanqi Research and Development Center of Zhejiang University, Hangzhou, P.R.

China

[†]These authors contributed equally to this work

[$] These authors contributed equally to this work

*For correspondence: duanshumin@zju.edu.cn (S.D.); zhihuagao@zju.edu.cn (Z.G.)

**Abstract**

Defensive behaviors are critical for animal's survival. The parabrachial nucleus (PBN) is critical in transmitting danger stimuli to trigger behavioral responses. Here, by viral tracing and electrophysiological recordings, we uncover that cholecystokinin (CCK) neurons in the lateral PB (LPB$^{CCK}$ neurons) are monosynaptically connected with the paraventricular nucleus (PVN) neurons. Selective photoactivation of LPB$^{CCK}$ neurons, their axon terminals within the PVN and glutamatergic neurons in the PVN, trigger defensive-like flight-to-nest behavior. Conversely, photoinhibition of LPB$^{CCK}$ neurons suppresses defensive responses. We further show that LPB$^{CCK}$ neurons provide excitatory transmission to the PVN neurons, and blocking glutamatergic transmission inhibits LPB$^{CCK}$-PVN pathway activation-induced flight behaviors. Also, chemogenetic inhibition of PVN neurons prevents photoactivation of LPB$^{CCK}$-PVN pathway-driven flight responses. These data suggest that LPB$^{CCK}$ neurons recruit downstream PVN neurons to actively engage in flight responses. Our study identifies a novel circuit from LPB$^{CCK}$ to PVN and dissects its role in defensive behaviors.

**Introduction**

Living in an environment full of diverse threats, animals develop a variety of defensive

behaviors for survival during evolution (Anderson & Adolphs, 2014; Blanchard et al.,

2001; Crawford & Masterson, 1982; Fanselow & Bolles, 1979; LeDoux, 2012; Yilmaz

& Meister, 2013). From the perspective of predator and prey interactions, defensive

responses are generally divided into two categories: defensive behaviors avoiding being

discovered (for example, freezing or hiding behaviors) and defensive behaviors

avoiding being caught (for example, running or attacking behaviors) (Crawford &

Masterson, 1982; Fanselow & Bolles, 1979; Yilmaz & Meister, 2013). Which type of

defensive behavior is selected depends on many factors, such as the distance from threat

and access to shelters. In general, escape is triggered when animals are faced with an

imminent threat with a shelter nearby (Eilam, 2005; Perusini & Fanselow, 2015; L. Sun

et al., 2020; Zhang et al., 2018).

Several brain regions, including the periaqueductal gray (PAG), amygdala and

medial hypothalamic zone (MHZ), have been implicated in mediating escape behaviors

(Evans et al., 2018; Shang et al., 2018; Shang et al., 2015; Silva et al., 2013; Tovote et

al., 2016; Wang et al., 2015; Wang, Schuette, Nagai, et al., 2021; Wei et al., 2015).

Recent studies have highlighted a role for MHZ, including the anterior hypothalamic

nucleus (AHN), the dorsomedial part of ventromedial hypothalamic nucleus (VMHdm)

and the dorsal pre-mammillary nucleus (PMd), in mediating escape behaviors. For

example, projections from PMd to the dorsolateral periaqueductal gray region (dlPAG)

and those from PVN to the anteromedial ventral thalamic region (AMV) have been

shown to control escape behaviors (Wang, Schuette, Nagai, et al., 2021). Another study

showed that a subset of VMH neurons collaterally project to the AHN and PAG to

promote escape and avoidance behaviors (Wang et al., 2015). In addition, PVN, an area

enriched with endocrine neurons, has also been associated with escape behaviors.

Optogenetic activation of $Sim1^+$ neurons in the PVN triggers escape behavior (Mangieri

et al., 2019b). Projections from PVN to the ventral midbrain region (vMB) mediate

profound defensive-like behaviors, including hiding, escape jumping, hyperlocomotion

(Mangieri et al., 2019b), whereas projections from PVN to the ventral lateral septum

(LSv) drive fear-related escape jumping (Xu et al., 2019). Despite a growing number

of studies illustrating the importance of PVN in mediating defensive behaviors,

upstream input that transmit the danger signals to PVN remains unclear.

The parabrachial nucleus located in the dorsolateral pons, serves as a relay station

for sensory transmission during the aversive sensory processes (Fulwiler & Saper,

1984). PBN bears association with various sensory processes including the pain, itch,

hunger, and thermoregulation to protect the body from noxious stimuli (Alhadeff et al.,

2018; Campos et al., 2017; Campos et al., 2018; Carter et al., 2013; Chen et al., 2018;

Geerling et al., 2016; Kobayashi & Osaka, 2003; Mu et al., 2017; Nakamura &

Morrison, 2008, 2010; Uddin et al., 2018). In addition, PBN contains several subregions

with diverse peptide-expressing neuronal subtypes. The external lateral nuclei of PBN

(PBel), rich in calcitonin gene-related peptide (CGRP) neurons, plays an important role

in affective-behavior states to limit the harm caused by potential threats, such as painful

stimuli, itch-related stimuli, novel foods (Campos et al., 2018; Carter et al., 2013;

Gauriau & Bernard, 2002; Han et al., 2015; Kobayashi & Osaka, 2003). The dorsal

lateral area of PB (PBdl) contains dynorphin neurons and is crucial for thermoregulation

(Norris et al., 2021). The superior lateral area of PB (PBsl) contains a large number of

CCK neurons, which has been linked to glucose regulation via projections to VMH

(Garfield et al., 2014) and heat defense via projections to preoptic area (POA) (Yang et

al., 2020). Among these peptide-expressing neurons, CGRP neurons in the LPB are

implicated in defensive behavior under stressful or threatening circumstances. $LPB^{CGRP}$

neurons have been implicated in encoding alarm response by projecting to a number of

limbic regions including the amygdala and bed nucleus of stria terminalis (BNST). It

remains unclear whether neurons in the LPB are directly connected to the hypothalamus

to mediate defensive behaviors.

CCK and CCK receptor-expressing neurons have been associated with defensive

behaviors (Bertoglio et al., 2007; Chen et al., 2022; Wang, Schuette, La-Vu, et al.,

2021). PVN neurons also express CCK receptors and are involved in defensive

behaviours (Mangieri et al., 2019a; O'Shea & Gundlach, 1993; Xu et al., 2019), raising

a possibility that upstream CCK input to the PVN may be implicated in defensive

behaviours. In attempting to answer these questions, we firstly injected the Cre-

dependent retrograde tracer (AAV2-Retro-DIO-EYFP) into the PVN of CCK-Cre mice

and found that LPB was an important upstream area transmission to the PVN. By

electrophysiological analyses, we verified the direct connections between $LPB^{CCK}$

neurons and PVN neurons. Using optogenetic tools, we showed that activation of

$LPB^{CCK}$ neurons drove rapid flight-to-nest behaviors, whereas inhibition of $LPB^{CCK}$

neurons suppressed heat-evoked defensive behaviors, suggesting that LPB^CCK neurons

were sufficient and required for defensive behaviors. Interestingly, photoactivation of

terminals from LPB^CCK to PVN or activation of PVN^Vglut2 neurons alone also elicited

flight-to-nest behavior. Both chemogenetic inhibition of PVN neurons and

pharmacological blockade of glutamatergic transmission prevented axonal termianl

activation of LPB^CCK within the PVN-induced flight-to-nest behavior, suggesting that

PVN was an important target mediating LPB^CCK-driven defensive responses. Our data

suggest a novel neurocircuit that mediates defensive behavior, which may shed light on

the mechanism of fear-related psychiatric diseases.

**Results**

**CCK$^+$ neurons in the LPB provides monosynaptic glutamatergic projections to the**

**PVN**

To identify the upstream CCKergic inputs to the PVN, we injected the retrograde viral

tracer (AAV2-Retro-DIO-EYFP, hereafter referred to as AAV2-Retro) into the PVN

region of a knock-in mouse line expressing Cre recombinase at the CCK Locus (CCK-

ires-Cre) **(Figure 1A)**. We observed abundant EYFP$^+$ neurons in the medial orbital

cortex (MO), cingulate cortex (CC), LPB and scattered GFP$^+$ cells in the PAG and

dysgranular insular cortex (DI) **(Figure 1B-C，Supplementary Fig.1).** Among these

nuclei, we mainly focused on the LPB, a region critical for sensory signal processing

(Fulwiler, C.E., and Saper, C.B.,1985; Garfield AS et al., 2014). To further verify the

anatomical connections between the LPB and PVN, we injected the anterograde viral

tracer (AAV-hSyn-FLEx-mGFP-2A-Synaptophysin-mRuby) into the LPB of CCK-Cre

mice **(Figure 1D)**. We found that prominent GFP$^+$ fibers and mRuby$^+$ bouton-like

structures, reminiscent axonal terminals, within the PVN, indicating that LPB$^{CCK}$

neurons directly project to the PVN **(Figure 1E-F, Supplementary Fig 2).**

CCK$^+$ neurons have been shown to primarily located within PBsl (Garfield et al.,

2014). To further visualize and characterize LPB$^{CCK}$ neurons, we crossed the CCK-ires-

Cre mice with the Cre-dependent tdTomato (tdT) reporter line (Ai14) and found that

the expression of CCK was abundantly expressed in the rostral and mid LPB **(Figure**

**2A-C)**. The majority of LPB neurons have been shown to be glutamatergic neurons

with a small population being GABAergic (Geerling et al., 2016; Kaur et al., 2013). To

dissect the identity of CCK$^+$ neurons in the LPB, we performed *in situ* hybridization in

CCK-ires-Cre::Ai14 mice using probes for *Vglut2* and *Vgat*, markers specific to

glutamatergic and GABAergic neurons. We found that LPB$^{CCK}$ neurons (84.0% ± 1.7%)

predominantly expressed *Vglut2*, with a tiny subpopulation (2.0% ± 1.2%) expressed

*Vgat*, suggesting that LPB$^{CCK}$ neurons were mostly glutamatergic neurons. In addition,

co-labeling with another neuropeptide, CGRP, in these mice revealed a slight co-

localization (2.2% ± 0.37%) between CCK and CGRP **(Figure 2D-G)**, suggesting a

separation between these two subpopulations of neurons within different areas of LPB.

Collectively, these data indicate that CCK neurons comprise a subpopulation of

glutamatergic neurons and barely overlap with CGRP neurons within the LPB.

Next, we used electrophysiological recording to investigate the functional

connections between the LPB and PVN by injecting the Cre-dependent adeno-

associated virus (AAV) expressing channelrhodopsin-2 virus (AAV2/9-DIO-ChR2-EYFP) into the LPB of CCK-Cre mice **(Figure 2H)**. By optic stimulating the axonal terminals of LPB$^{CCK}$ neurons within the PVN, we successfully recorded excitatory postsynaptic currents (EPSCs), rather than inhibitory postsynaptic currents (IPSCs). These EPSCs were blocked in the presence of a sodium channel antagonist TTX and rescued by the potassium channel antagonist 4-AP, suggesting monosynaptic connections between these neurons. Further, the EPSCs were blocked by NMDA receptor antagonist AP5 **(Figure 2I)**, and AMPA receptor antagonist CNQX **(Figure 2J-K)**. These data suggest that neurons in the PVN receive monosynaptic excitatory innervations from LPB$^{CCK}$ neurons.

**Photostimulation of LPB$^{CCK}$ neurons triggers defensive-like flight-to-nest behavior**

Previous studies have shown that LPB$^{CCK}$ neurons participate in the regulation of glucose homeostasis through SF1 neurons in the VMH (Garfield et al., 2014) and thermoregulation via connections with the POA (Yang et al., 2020). It remains unclear what effects direct optogenetic activation of LPB$^{CCK}$ neurons may induce. We firstly tested the efficacy of ChR2-mediated activation by injecting the AAV2/9-DIO-ChR2-EYFP virus into the LPB of CCK-Cre mice and carried out whole-cell recordings from brain slices for viral-infected neurons. As shown in **Figure 3A**, different frequencies of blue laser pulses (from 5-20 Hz) induced time-locked action potential firing in LPB$^{CCK}$ neurons expressing ChR2, with 20 Hz inducing maximal firing capacities, verifying the

effectiveness of optic tools.

PBN is an important relay nucleus that transmits peripheral sensations to the brain

to limit harm (Campos et al., 2018). Recent studies have found that LPB[CGRP] neurons

play an important role in mediating fear-related behaviors (Campos et al., 2018; Han et

al., 2015). It remains unclear whether LPB[CCK] neurons also participate in fear-related

behaviors. To determine whether LPB[CCK] neurons mediate fear-related behavior, we

optically activated these neurons as described above **(Figure 3B-3C)**, and investigated

fear-related defensive-like behaviors in mice using a flight-to-nest test experimental

paradigm previously used (Zhou et al., 2019). We placed a nest in the corner to test the

ability of mice to actively search for hiding. Optogenetic activation of ChR2-expressing

CCK neurons induced defensive-like flight-to-nest behavior in the experimental

apparatus, as mice escaped towards the nest (latency: EYFP $87.71 \pm 22.38$ s versus

ChR2 $7.571 \pm 0.8123$ s; speed: EYFP $184.4\% \pm 30.4\%$ versus ChR2 $361.7\% \pm 31.5\%$)

followed by long-lasting time stay in the nest (EYFP $63.906\% \pm 3.645\%$ versus ChR2

$97.26\% \pm 2.737\%$) **(Figure 3D-J)**. Our data suggest that activation of LPB[CCK] neurons

is sufficient to drive defensive-like flight-to-nest behavior.

**Photostimulation of LPB[CCK] neurons induces anxiety-like behavior and**

**autonomic responses**

Since activation of LPB[CCK] neurons induced fear-related flight-to-nest behavior, and

long-term fear may trigger an anxiety-like state in animals (Tovote et al., 2005; Yang

et al., 2016), we next determined whether mice exhibit anxiety-like behaviors after

prolonged activation of LPB$^{CCK}$ neurons. We gave mice 10-min of light stimulation and

tested their responses in the open field and elevated plus maze 20-min later **(Figure**

**4A)**. We found that optogenetic stimulation of LPB$^{CCK}$ neurons expressing ChR2-

EYFP, but not those expressing EYFP, significantly reduced the total distance, center

time and center entries in the open field tests, as well as the open arm time and open

arm entries in the elevated plus maze tests **(Figure 4B-E)**. Our results indicate that

prolonged optogenetic manipulation of LPB$^{CCK}$ neurons induces anxiety-like behaviors.

When animals and humans face danger, the sympathetic nervous system is usually

activated, as characterized by increased heart rates and dilated pupil size (Dong et al.,

2019; Gross & Canteras, 2012; Wang et al., 2015). We found that photostimulation of

LPB$^{CCK}$ neurons significantly increased heart rate and enlarged pupil diameter,

suggesting a fast arousal state (Salay et al., 2018) **(Figure 4F-J)**. Moreover, plasma

levels of corticosterone were also increased upon photostimulation of LPB$^{CCK}$ neurons

**(Figure 4K)**, indicating a stress-related response in mice. In summary, photoactivation

of LPB$^{CCK}$ neurons induces accompanying changes in sympathetic nervous system and

stress responses in mice.

**Inhibition of LPB$^{CCK}$ neurons suppressed heat-evoked defensive behavior**

To examine whether LPB$^{CCK}$ neurons are required for defensive behavior, we then

inhibited LPB$^{CCK}$ neurons using optogenetic tools. By injecting the Cre-dependent

adeno-associated virus (AAV) expressing the guillardia theta anion channel

rhodopsins-1 (GtACR1) into the LPB of CCK-Cre mice, followed by optic stimulation,

we were able to effectively inhibit the firing of LPB$^{CCK}$ neurons **(Figure 5A)**.

Mammals are sensitive to temperature changes and develop multiple mechanisms to avoid excessive high or low temperature. When confronted extreme heat or cold, animals would turn to an area with comfortable temperature. Since LPB$^{CCK}$ neurons have been recently shown to be implicated in thermosensation (Yang et al., 2020), we speculated that they may mediated heat defense. To determine whether LPB$^{CCK}$ neurons might be involved in heat defense, we designed an apparatus, which containing two zones of different temperatures, 25 degree, the comfortable zone at room temperature, and 43 degree, the uncomfortable zone at a higher temperature which induces uncomfortable feeling but not pain (Wang, Schuette, Nagai, et al., 2021). To test whether inhibition of LPB$^{CCK}$ neurons suppressed heat-evoked defensive behaviors, we bilaterally injected Cre-dependent virus (AAV-CAG-DIO-GtACR1-EYFP or AAV-CAG-DIO-EYFP) into LPB and positioned optical fibers above the LPB of CCK-Cre mice **(Figure 5B and 5C).** After habituation in the customized apparatus (Figure 5D), control mice exhibited quick heat defense by running to the comfortable zone when placed in the hot zone edge (43˚C) **(Figure 5D)**. However, optic inhibition of LPB$^{CCK}$ neurons led to a significant delay in the heat defense behavior, with a substantially increased time in the uncomfortable zone, compared to the controls (EYFP 14.66%±4.119% versus GtACR1 57.3% ± 14.33%), accompanied by a much longer latency returning to the comfortable zone (EYFP 22.23 ± 13.06 s versus GtACR1 163.5 ± 45.75 s), suggesting a delayed response to the hot stimuli **(Figure 5E-G)**. Our data suggest that LPB$^{CCK}$ neurons are required to induce heat defensive behavior.

**Photostimulation of LPB$^{CCK}$-PVN neurons triggers defensive-like flight-to-nest**

**behavior**

To further investigate whether activation of the LPB$^{CCK}$-PVN pathway induces

defensive-like flight-to-nest behavior, we unilaterally injected Cre-dependent adeno-

associated virus (AAV-CAG-DIO-ChR2-EYFP or AAV-CAG-DIO-EYFP) into the

LPB of CCK-Cre mice and positioned an optical fiber above the PVN **(Figure 6A and**

**6B)**. By photostimulating the axonal terminals of LPB$^{CCK}$ neurons in the PVN, we also

observed apparent defensive-like flight-to-nest behavior (latency: EYFP 73.43 ± 21.08

s versus ChR2 8 ± 1.047 s; speed: EYFP 122.3% ± 17.81% versus ChR2 277.9% ±

32.85%; time in the nest: EYFP 3.751% ± 1.23% versus ChR2 89.29% ± 10.71%)

**(Figure 6C-H)**, similar to soma activation. Immunostaining of c-fos further verified the

activation of PVN neurons upon optic stimulation of axonal terminals of LPB$^{CCK}$

neurons **(Figure 6I-J)**. As well, terminal activation also increased heart rate and plasma

corticosterone levels **(Figure 6K-L)**, suggesting that PVN is an important downstream

target region mediating defensive-like flight-to-nest behavior and accompanying

activation of autonomic nervous system.

**PVN is required for defensive response to threatening situations**

Since LPB$^{CCK}$ neurons project to multiple regions **(Supplementary figure 1)**, thus the

effects observed above may arise from other downstream targets. To test the possibility,

we investigated whether inhibition of the PVN is required for the LPB$^{CCK}$ neurons-

induced defensive-like flight-to-nest behavior. To achieve this aim, we used a dual-

virus mediated optogenetic and chemogenetic strategy, by injecting the photoactivable

virus AAV2/9-DIO-ChR2-EYFP into the LPB, and a chemogenetic-inhibitory virus

AAV2/9-hsyn-hM4Di-mCherry into the PVN of CCK-Cre mice. By putting an optical

fiber above the PVN, we were able to activate the LPB$^{CCK}$ terminals and induce

defensive-like flight-to-nest behavior as shown above **(Figure 7A and 7B)**. Upon CNO

administration to inhibit PVN neurons, however, we observed a much longer latency,

lower speed of the mice in returning to the nest, and lower percentage of hiding time

spent in the nest after photostimulation of LPB$^{CCK}$-PVN pathway (latency: saline 8.286

± 1.475 s versus CNO 27.86 ± 6.978 s; speed: saline 289.3% ± 25.22% versus CNO

111.7% ± 17.38%; time in the nest: saline 81.55% ± 11.16% versus CNO 25.6% ±

10.33%) **(Figure 7C-F)**. These data suggest that PVN is essential for the LPB$^{CCK}$

neurons activation-induced defensive-like flight-to-nest behavior.

LPB$^{CCK}$ neurons may release either glutamate or CCK to induce the defensive

behavior. To determine which neurotransmitter or modulator is engaged in the above

responses, we combined pharmacology with optogenetic manipulation. Two types of

CCK receptors are present in the central nervous system and CCKaR is widely

distributed throughout the PVN, whereas the expression of CCKbR receptors in PVN

is low **(Figure 7G-H)**. We infused glutamate receptor antagonists (CNQX + AP5),

CCK receptor antagonists (Devazepide + L-365,260), or artificial cerebrospinal fluid

(aCSF) through an implanted cannula 30-min before optogenetic activation of LPB$^{CCK}$

neurons **(Fig 7I-J)**. We found that blocking glutamate receptors induced a longer

latency, slower speed returning to the nest and lower percentage of hiding time in the nest, compared with ACSF-treated mice. However, blocking CCK receptors showed no significant difference compared to the control **(Figure 7K-N).** These results suggest that glutamatergic transmission from LPB[CCK] neurons to PVN regulates defensive-like flight-to-nest behavior.

**Photostimulation of PVN[Vglut2] neurons rather than PVN[CRH] neurons triggers defensive-like flight-to-nest behavior**

Of note, PVN neurons are mostly glutamatergic (Vong et al., 2011; Xu et al., 2013). We further assessed whether optogenetic activation of PVN[Vglut2] neurons elicit defensive-like behavior. We injected a Cre-dependent adeno-associated virus (AAV-CAG-DIO-ChR2-EYFP or AAV-CAG-DIO-EYFP) into the PVN and positioned an optical fiber above the PVN of Vglut2-Cre mice **(Figure 8A-B)**. We found that optogenetic activation of ChR2-expressing Vglut2 neurons in the PVN also induced defensive-like flight-to-nest behavior. Mice escaped to the nest (latency: EYFP $127.5 \pm 22.5$ s versus ChR2 $12.83 \pm 2.937$ s; speed: EYFP $142.4 \pm 32.07\%$ versus ChR2 $384.4\% \pm 83.52\%$), followed by longer time stay in the nest (EYFP $0.2783\% \pm 0.2783\%$ versus ChR2 $61.11\% \pm 17.58\%$) **(Figure 8C-E)**, compared to the control. These data suggest that optogenetic activation of PVN[Vglut2] neurons was sufficient to evoke defensive-like flight-to-nest behavior.

Recently, a study showed PVN[CRH] neurons could predict the occurrence of defensive behavior in mice (Daviu et al., 2020). To determine whether activation of the

PVN<sup>CRH</sup> neurons induced defensive-like flight-to-nest behavior, we injected a Cre-dependent adeno-associated virus (AAV-CAG-DIO-ChR2-EYFP or AAV-CAG-DIO-EYFP) into the PVN and positioned an optical fiber above the PVN of CRH-Cre mice **(Figure 8F-G)**. However, we found that optogenetic activation of ChR2-expressing CRH neurons in the PVN did not elicit defensive-like flight-to-nest behavior **(Figure 8H-J)** (latency: EYFP $77 \pm 20.69$ s versus ChR2 $90.17 \pm 22.03$ s; speed: EYFP $156.6 \pm 22.59\%$ versus ChR2 $217.2\% \pm 61.54\%$; time in the nest: EYFP $20.95\% \pm 13.89\%$ versus ChR2 $35.28\% \pm 20.48\%$).

**Discussion**

In the present study, we found that LPB$^{CCK}$ neurons regulate defensive-like flight-to-nest behavior through a glutmatergic excitatory projections to PVN. Morover, photoactivation of PVN$^{Vglut2}$ neurons, rather than PVN$^{CRH}$ neurons, elicits flight-to-nest behavior. Taken together, these results revealed that a novel circuit from the brainstem to hypothalamus regulates defensive behaviors **(Figure 9)**.

CCK-expressing neurons are widely distributed throughout the brain. Previous studies showed that CCK neurons in the cortex and hippocampus are mainly GABAergic (Liu et al., 2020; X. C. Sun et al., 2020; Whissell et al., 2015), while LPB$^{CCK}$ neurons are mainly glutamatergic neurons. Furthermore, LPB$^{CCK}$ neurons are anatomically separated from LPB$^{CGRP}$ neurons, which have been associated with learned fear responses and freezing behaviors, a passive coping strategy, when encountered with threats (Han et al., 2015; Keay & Bandler, 2001). By contrast, we show that selective activation of LPB$^{CCK}$ neurons triggered defensive-like flight-to-nest behavior (Keay & Bandler, 2001), an active coping strategy. Our data suggest that different neuronal subpopulations in spatially segregated regions within the LPB may encode distinct threat signals and drive distinct defensive responses when animals encounter danger. Whether LPB$^{CCK}$ neurons are further related to conditioned fear, needs further exploration.

LPB neurons are involved in regulating a series of physiological activities to protect the body from threat, including extremely heat and cold (Geerling et al., 2016; Kobayashi & Osaka, 2003; Nakamura & Morrison, 2008, 2010). Studies have demonstrated that LPB contains thermosensory neurons that facilitate various

homeostatic functions (Chiang et al., 2019). Recently, it has been shown that several neuropeptide-containing neurons in the LPB including cholecystokinin, prodynorphin and proenkephalin neurons, are activated by warm temperature (Norris et al., 2021; Yang et al., 2020). Their projections to the POA are involved in maintaining temperature homeostasis by driving physiological or behavioral heat defenses. In this study, we show that LPB$^{CCK}$ neurons are required for the heat-induced defensive behavior, suggesting that they are likely to encode heat-related threat signals and drive defensive responses.

LPB receives diverse sources of inputs and sends projections to many forebrain regions (cortex, central amygdala, BNST), hypothalamus (PVN, POA, dorsomedial hypothalamic nuclei, lateral hypothalamic area), hindbrain regions (nucleus of the solitary tract and spinal dorsal horn) (Chiang et al., 2019; Cooper Jr & Blumstein, 2015; Ellard & Eller, 2009; Han et al., 2015; Krout & Loewy, 2000; L. Sun et al., 2020). Despite many early studies mapping downstream nuclei for LPB neurons, our study is the first to demonstrate that LPB$^{CCK}$ neurons possess direct anatomical and functional connections with PVN neurons. We further show that LPB$^{CCK}$ neurons convey threatening signals to the PVN that drives the flight-to-nest behavior. Interestingly, blocking glutamate receptors rather than CCK receptors in the PVN, paired with photoactivation of LPB$^{CCK}$-PVN pathway, suppressed the flight-to-nest behavior, demonstrating the necessity and sufficiency of LPB$^{CCK}$-PVN glutamatergic transmission for defensive responses. Besides the PVN, we also observed that LPB$^{CCK}$ neurons project to the paraventricular thalamic nucleus (PVT), VMH and PAG, all of

which have been associated with aversion and/or defensive behaviors (LeDoux, 2012;

Vianna et al., 2001; Wang et al., 2015). It would be interesting to further investigate

how these different downstream areas contribute to defensive responses.

PVN plays an essential role in modulating neuroendocrine and autonomous

functions (Canteras et al., 2001; Coote, 2005; Ferguson et al., 2008; Sutton et al., 2016).

Although earlier literatures suggest that PVN neurons are activated in response to

different threat stimuli by c-fos staining (Canteras et al., 2001; Faturi et al., 2014;

Martinez et al., 2008; Staples et al., 2008), it was unclear whether these neurons are

only involved in neuroendocrine responses to stress or directly regulate defensive

behaviors. Recently, increasing evidence demonstrated that PVN neurons play

important roles in defensive behaviors independent of hormonal actions (Daviu et al.,

2020; Mangieri et al., 2019b; Xu et al., 2019). Our results also support a role of PVN

neurons in mediating behavioral response of defense, as selective photostimulation of

$PVN^{Vglut2}$ neurons drives a rapid flight-to-nest behavior. While studies showed that

$PVN^{CRH}$ neurons participate in the initiation of escape behaviors upon a visual threat

and encode stress signals (Daviu et al., 2020), we found that photoactivation of $PVN^{CRH}$

neurons was insufficient to induce the flight-to-nest behavior. The following factors

may contribute to these discrepancies. First, despite that $PVN^{CRH}$ neurons receive inputs

from the LPB (Zhang et al., 2019), these neurons may not necessarily be $LPB^{CCK}$

neurons. Second, $PVN^{CRH}$ neurons receive inputs from multiple upstream nuclei and

activation of $PVN^{CRH}$ neuronal ensemble may not fully reflect the function of CRH

neurons in input-specific manner. Third, CRH neurons may only be involved in

initiation of escape but may not necessarily involved in the flight-to-nest behavior.

Future studies using additional projection-specific approaches and more genetically

defined cell tools may help resolve this problem. In addition, since PVN contains

diverse neuronal types, it is likely that different subpopulations of neurons may

coordinatively mediate defensive responses. It is worthwhile to further investigate

which type of neurons in the PVN are involved in these responses.

Our data show that activation of LPB$^{CCK}$ neurons drive defensive and anxiety-like

behaviors, eliciting an arousal and stressed state. Fear, anxiety and stress responses are

closely associated with mental illnesses involving dysregulated neural circuits

(Deisseroth, 2014; Jamalamadaka et al., 2020; Luthi & Luscher, 2014; Pitman et al.,

2012). It is worth further investigation to examine whether alterations of the LPB$^{CCK}$-

PVN pathway are implicated in these diseases, in which manipulation of this pathway

might be an effective target for potential therapeutic strategy.

**Materials and Methods**

**Key resources table**

| Reagent type (species) or resource | Designation | Source or reference | Identifiers | Additional information |
|---|---|---|---|---|
| Strain, strain background (Mus musculus) | CCK-ires-Cre(Cck$^{tm1.1(cre)Zjh}$/J) | The Jackson Laboratory | JAX012706 | |
| Strain, strain background (Mus musculus) | CRH-ires-Cre (B6(Cg)-Crh$^{tm1(cre)Zjh}$/J | The Jackson Laboratory | JAX012704 | |
| Strain, strain background (Mus musculus) | Vglut2-ires-Cre (Slc17a6$^{tm2(cre)Lowl}$/J) | The Jackson Laboratory | JAX 016963 | |
| Strain, strain background (Mus musculus) | Ai14 (B6;129S6-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J) | The Jackson Laboratory | JAX007908 | |
| Strain, strain background (Mus musculus) | Ai3; (B6.Cg-Gt (ROSA)26Sor$^{tm3(CAG-EYFP)Hze}$/J) | The Jackson Laboratory | JAX007903 | |
| Strain, strain background (Mus musculus) | C57BL6/J | Shanghai SLAC Laboratory Animal Co. Ltd | http://www.slaccas.com/ | |
| Antibody | Anti-Dsred, rabbit polyclonal | Takara | Cat# 632496 RRID: AB_10013483 | (1:800) |
| Antibody | Anti-CGRP, mouse monoclonal | Abcam | Cat# 81887 RRID: AB_1658411 | (1:800) |
| Antibody | Anti-GFP, goat polyclonal | Abcam | Cat# 5450 RRID: AB_304896 | (1:500) |
| Antibody | Anti-CRH, rabbit polyclona | Phoenix Biotech | Cat# H-019-06 | (1:500) |
| Antibody | Anti-c-fos, Guinea pig polyclonal | Synaptic Systems | Cat# 226004 RRID: AB_2619946 | (1:10000) |
| Antibody | Alexa Fluor 488 donkey anti-guinea pig IgG (H+L) | Jackson | Cat#112-486-068 RRID: AB_2617153 | (1:1000) |
| Antibody | Alexa Fluor 555 donkey anti-rabbit IgG (H+L) | Invitrogen | Cat#A31572 RRID: AB_162543 | (1:1000) |
| Antibody | Alexa Fluor 488 donkey anti-mouse | Invitrogen | Cat# R37114 RRID: AB_2556542 | (1:1000) |

| | IgG (H+L) | | | |
|---|---|---|---|---|
| Antibody | Alexa Fluor 488 donkey anti-goat IgG (H+L) | Invitrogen | Cat# A11055 RRID: AB_2534102 | (1:1000) |
| Antibody | Alexa Fluor 488 donkey anti-rabbit IgG (H+L) | Jackson | Cat# R37118 RRID: AB_2556546 | (1:1000) |
| Commercial assay or kit | RNAscope Multiplex Fluorescent Reagent Kit v2 | Advanced Cell Diagnostics | Cat# 323100 | |
| Sequence-based reagent | RNAscope probe *Slc17a6* | Advanced Cell Diagnostics | accession number NM_080853.3 | probe region 1986-2998 |
| Sequence-based reagent | RNAscope probe *Slc32a1* | Advanced Cell Diagnostics | Accession number NM_009508.2 | probe region 894-2037 |
| Sequence-based reagent | RNAscope probe *CckaR* | Advanced Cell Diagnostics | accession number NM_009827.2 | probe region 328-1434 |
| Sequence-based reagent | RNAscope probe *CckbR* | Advanced Cell Diagnostics | accession number NM_007627.4 | probe region 136-1164 |
| Chemical compound, drug | CNQX disodium salt hydrate | Sigma-Aldrich | Cat#1045 | |
| Chemical compound, drug | Devazepide | Sigma-Aldrich | Cat#2304 | |
| Chemical compound, drug | Clozapine N-oxide | Sigma-Aldrich | Cat#C0832 | |
| Chemical compound, drug | L-365,260 | Sigma-Aldrich | Cat#143626 | |
| Chemical compound, drug | DAPI | Sigma-Aldrich | N/A | |
| Chemical compound, drug | Tween-20 | Sigma-Aldrich | N/A | |
| Strain, strain background (AAV2/9) | AAV2/9-hEF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA | Shanghai Taitool Bioscience Co. | Cat# S0199-9 | Viral titers: 2.95x1013 particles/ml |
| Strain, strain background (AAV2/9) | AAV2/9-hEF1a-DIO-EYFP-WPRE-pA | Shanghai Taitool Bioscience Co. | Cat#S0196-9 | Viral titers:$1.0x10^{12}$ particles/ml |
| Strain, strain background | AAV2/9-CAG-DIO-hGtACR1-P2A- | Shanghai Taitool Bioscience Co. | Cat# S0311-9 | Viral titers:$5 \times 10^{13}$ particles/ml |

| (AAV2/9) | EGFP-WPRE-pA | | | |
|---|---|---|---|---|
| Strain, strain background (AAV2/9) | rAAV2/9-hSyn-DIO-mGFP-2A-Synaptophysin-mRuby | Shanghai Taitool Bioscience Co. | Cat# S0250-9 | Viral titers :$1.55 \times 10^{13}$ particles/ml |
| Strain, strain background (AAV2/2) | rAAV2/2-Retro-hEF1a-DIO-EYFP-WPRE-pA | Shanghai Taitool Bioscience Co. | Cat# S0196-2R | Viral titers: $2.52 \times 10^{13}$ particles/ml |
| Strain, strain background (AAV2/9) | AAV2/9-hSyn-mCherry-WPRE-pA | Shanghai Taitool Bioscience Co. | Cat# S0238-9 | Viral titers: $\geq 1.0 \times 10^{13}$ particles/ml |
| Strain, strain background (AAV2/9) | AAV2/9-hSyn-hM4D(Gi)-mCherry-WPRE-pA | Shanghai Taitool Bioscience Co. | Cat# S0279-9 | Viral titers: $\geq 1.0 \times 10^{13}$ particles/ml |
| Software, algorithm | ANY-Maze software 5.3 | Global Biotech Inc. | http://www.anymaze.co.uk/ | |
| Software, algorithm | Image J | NIH | https://imagej.nih.gov/ij/index.html;%20 RRID:SCR_003070 | |
| Software, algorithm | GraphPad Prism 6 | GraphPad Software | https://www.graphpad.com/scientificsoftware/prism/; RRID: SCR_002798 | |
| Software, algorithm | MatLab R2016a | MathWorks | https://www.mathworks.com/products.html; RRID:SCR_001622 | |

**Animals**

CCK-ires-Cre (Cck[tm1.1(cre)Zjh]/J; Stock No. 012706), CRH-ires-Cre (B6(Cg)-

Crh[tm1(cre)Zjh]/J; Stock No. 012704), Vglut2-ires-Cre (Slc17a6[tm2(cre)Lowl]/J; Stock No.

016963), Ai14 (B6;129S6-Gt(ROSA)26Sor[tm14(CAG-tdTomato)Hze]/J; Stock No. 007908),

C57BL/6 mice were obtained from the Shanghai Laboratory Animal Center. Adult male

mice were used in our study. Mice housed at $22 \pm 1$ °C and $55 \pm 5\%$ humidity on a 12-

hr light/12-hr dark cycle (light on from 07:00 to 19:00) with food and water ad libitum.

All experimental procedures were approved by the Animal Advisory Committee at

Zhejiang University and were performed in strict accordance with the National

Institutes of Health Guidelines for the Care and Use of Laboratory Animals. All

surgeries were performed under sodium pentobarbital anesthesia, and every effort was

made to minimize suffering.

**Immunohistochemistry**

Animals were transcardially perfused with saline and 4% PFA. Brains were post-fixed

overnight in 4% PFA at 4 °C, followed by immersed in 30% sucrose. Coronal sections

(40 μm) were cut by a CM1950 Microtome (Leica). Immunostaining was performed as

previously described (Zhang et al., 2018). The brain slices were permeabilized in 0.5%

Triton X-100 in Tris-buffered saline, blocked with 100 mM glycine and 5% bovine

serum albumin (BSA) containing 5% normal donkey serum. Tissue sections were

subsequently incubated with diluted primary and secondary antibodies as indicated,

nuclei stained with 6-diamidino2-phenylindole (DAPI), and slides mounted with

antifade reagents. The primary antibodies used were: Guinea pig anti-c-fos (Synaptic

Systems, Cat# 226004), Rabbit anti-Dsred (Takara, Cat# 632496), Goat anti-GFP

(Abcam, Cat# 5450), Mouse anti-CGRP (Abcam, Cat# 81887). Slides were imaged

with a confocal microscope (Olympus FluoView FV1200).

**RNAscope in situ hybridization**

We used RNAscope multiplex fluorescent reagent kit and designed probes (ACDBio

Inc.) to perform fluorescence in situ hybridization. Mouse brain tissue was sectioned

into 20 μm sections by cryostat (Leica CM 1950). Then sections were mounted on slides

and air-dried at room temperature. Subsequently, the sections were dehydrated in 50%

EtOH, 70% EtOH, and 100% EtOH for 5-10 min each time and air-dried at room

temperature again. Thereafter, protease digestion was performed in a 40 °C HybEZ

oven for 30 min pretreatment, slides were hybridized with pre-warmed probe in a 40 °C

HybEZ oven for 2 hours. Probes used in our paper were: *Slc17a6* probe (*VgluT2,*

accession number NM_080853.3, probe region 1986-2998), *Slc32a1* probe (*VGAT,*

accession number NM_009508.2, probe region 894-2037), *CCKaR* probe (accession

number NM_009827.2, probe region 328-1434), *CCKbR* probe (accession number

NM_007627.4, probe region 136-1164). After hybridization, the brain sections went

through 4 steps of signal amplification fluorescent label. Anti-DsRed or GFP staining

was performed after the RNAscope staining. Slides were imaged with a confocal

microscope (Olympus FluoView FV1200).

**Stereotaxic injections and optical fiber/cannula implantation**

For surgical procedures, mice were anaesthetized with sodium pentobarbital (0.1 g/kg)

and placed in a stereotaxic apparatus (RWD). Stereotaxic surgery was performed as

described (Li Sun et al., 2020). Briefly, holes were made into the skull over the target

areas to inject the virus with glass pipettes (diameter,10-15 mm) or to implant optical

fibers [outer diameter (o.d.): 200 μm; length: 6.0 mm, 0.37 NA; Inper] or to implant of

guide cannula [outer diameter (o.d.): 0.41 mm; RWD]. The coordinates relative to

bregma were as follows according to the Paxinos and Franklin (2001) atlas. For all

experiments, mice with incorrect injection sites were excluded from further analysis.

For optical activation of LPB$^{CCK}$ neurons, AAV2/9-hEF1a-DIO-hChR2(H134R)-

EYFP-WPRE-pA(viral titers: $2.95 \times 10^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-

hEF1a-DIO-EYFP-WPRE-pA (viral titers: $1.0 \times 10^{12}$ particles/ml; Taitool Bioscience)

virus was unilaterally microinjected into the LPB (AP:-4.8; ML:-1.35; DV: -3.4; mm

relative to bregma) of CCK-Cre mice. The virus was diluted into $5.9 \times 10^{12}$ genomic

copies per ml with phosphate-buffered saline (PBS) before use and injected with 65 nl

into the LPB. Virus was delivered at a flow rate of 10 nl/min. The glass capillary was

left in place for an additional 10 min after injection to allow diffusion of the virus. The

cannulas were held in place with dental cement above the LPB (AP: -4.8; ML: -1.35;

DV: -3.2; mm relative to bregma).

For optical activation of PVN$^{Vglut2}$ neurons, AAV2/9-EF1a-DIO-hChR2-EYFP

(viral titers: $2.95 \times 10^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-EF1α-DIO-EYFP

(viral titers: $1.0 \times 10^{12}$ particles/ml; Taitool Bioscience) virus was unilaterally

microinjected into the PVN (AP: -0.4; ML: -0.15; DV: -4.95; mm relative to bregma)

of Vglut2-Cre mice. The virus was diluted into $2.0 \times 10^{12}$ genomic copies per ml with

PBS before use and injected with 70 nl into the PVN. Virus was delivered at a flow rate

of 15 nl/min. The glass capillary was left in place for an additional 10 min after injection

to allow diffusion of the virus. The cannulas were held in place with dental cement

above the PVN (AP: -0.4; ML: -0.15; DV: -4.75; mm relative to bregma).

For optical activation of PVN$^{CRH}$ neurons, AAV2/9-EF1a-DIO-hChR2-EYFP

(viral titers: $2.95 \times 10^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-EF1α-DIO-EYFP

(viral titers: $1.0 \times 10^{12}$ particles/ml ; Taitool Bioscience) was unilaterally microinjected

into the PVN (AP: -0.4; ML: -0.15; DV: -4.90 mm relative to bregma) of CRH-Cre

mice. The virus was diluted into $4.0 \times 10^{12}$ genomic copies per ml with PBS before use

and injected with 75 nl into the PVN. Virus was delivered at a flow rate of 15 nl/min.

The glass capillary was left in place for an additional 10 min after injection to allow

diffusion of the virus. The cannulas were held in place with dental cement above the

PVN (AP: -0.4; ML: -0.15; DV: -4.7; mm relative to bregma).

For optical inhibition of LPB$^{CCK}$ neurons, AAV2/9-EF1α-DIO-hGtACR1-P2A-

eYFP-WPRE ($5 \times 10^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-hEF1a-DIO-EYFP-

WPRE-pA (viral titers: $1.0 \times 10^{12}$ particles/ml; Taitool Bioscience) virus was bilaterally

infused to the LPB (AP: -4.8; ML: -1.35; DV: -3.4; mm relative to bregma) of CCK-

Cre mice. The virus was diluted into $1.67 \times 10^{12}$ genomic copies per ml with PBS before

use and injected with 70 nl into the LPB. Optical fibers were bilaterally implanted at an

angle of 5° above the LPB (AP: -4.8; ML: -1.72; DV: -2.94; mm relative to bregma).

In order to study the downstream of LPB$^{CCK}$ neurons, AAV2/9-hSyn-DIO-mGFP-

2A-Synaptophysin-mRuby (viral titers: $1.55 \times 10^{13}$ particles/ml; Taitool Bioscience)

virus was unilaterally injected into the LPB (AP: -4.8 mm; ML: -1.35 mm; DV: -3.4

mm) of CCK-Cre mice; in order to study the source of CCKergic upstream of PVN,

AAV2/2-Retro-hEF1a-DIO-EYFP-WPRE-pA (viral titers: $2.52 \times 10^{13}$ particles/ml;

Taitool Bioscience) virus was unilaterally injected into the PVN (AP, -0.4 mm; ML, -

0.2mm; DV, -4.85mm) of CCK-Cre mice.

For LPB$^{CCK}$-PVN axon terminal stimulation, AAV2/9-hEF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA (viral titers: 2.95×10$^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-hEF1a-DIO-EYFP-WPRE-pA (viral titers: 1.0×10$^{12}$ particles/ml; Taitool Bioscience) virus was unilaterally microinjected into the LPB (AP: -4.8; ML: -1.35; DV: -3.4; mm relative to bregma) of CCK-Cre mice. The virus was diluted into 5.9×10$^{12}$ genomic copies per ml with PBS before use and injected with 65 nl into the LPB. After the virus was expressed for 2 weeks, for LPB$^{CCK}$-PVN axon terminals stimulation, optical fibers were unilaterally implanted above the PVN (AP: -0.4; ML: -0.2; DV: -4.7; mm relative to bregma).

For pharmacological experiments (procedures of optical activation of LPB$^{CCK}$ neurons have been mentioned above), drug cannulas were ipsilaterally implanted into the PVN (AP, -0.4 mm; ML, -0.2mm; DV, -4.7mm).

For prolonged inhibition of PVN neurons, AAV2/9-hSyn-mCherry-WPRE-pA (viral titers: ≥1.0×10$^{13}$ particles/ml; Taitool Bioscience) or AAV2/9-hSyn-hM4D(Gi)-mCherry-WPRE-pA(viral titers: ≥1.0×10$^{13}$ particles/ml; Taitool Bioscience) virus was microinjected into the PVN (AP, -0.4 mm; ML, -0.2mm; DV, -4.85mm) of CCK-Cre mice.

**In vivo optogenetic manipulation**

For optogenetic manipulation experiments, an implanted fiber was connected to a 473 nm laser power source (Newdoon Inc., Hangzhou, China). The power of the blue (473 nm) (Newdoon Inc., Hangzhou, China) was 0.83-3.33 mW mm$^2$ as measured at the tip

of the fiber. 473 nm laser (ChR2: power 5-15 mW, frequency 20 Hz, pulse width 5 ms;

GtACR1: power 15 mW, direct current) was supplied to activate or inhibit neurons,

respectively.

**Pharmacological antagonism**

Antagonism was delivered 30 min before optical activation of LPB$^{CCK}$ neurons. For

blocking glutamatergic neurotransmission, we firstly connected guiding cannula with a

Hamilton syringe via a polyethylene tube. Then we infused 0.25 μl mixed working

solution containing CNQX disodium salt hydrate (0.015 μg; Sigma-Aldrich), a

glutamate AMPA receptor antagonist, and AP5 (0.03 μg; Sigma-Aldrich), a NMDA

receptor antagonist, into the PVN with a manual microinfusion pump (RWD, 68606)

over 5 min. For blocking CCKergic neurotransmission, 0.25 μl mixed solution

containing Devazepide (0.0625 μg; Sigma-Aldrich), a CCKaR receptor antagonist, and

L-365,260(0.0625 μg; Sigma-Aldrich), a CCKbR receptor antagonist, was infused into

the PVN over a period of 5 min. To prevent backflow of fluid, the fluid-delivery cannula

was left for 10 min after infusion.

**Chemogenetic manipulation**

CNO (Sigma) was dissolved in saline (5 mg in 10 mL DMSO & 190 mL NaCl). CNO

was injected intraperitoneally at 0.3 mg per kg of body weight for chemogenetic

manipulation.

**Fos induction and analysis**

For c-Fos quantification, mice were perfused 1.5 h after 10 min blue photostimulation illumination, and sections were cut. The boundaries of the nuclei were defined according to brain atlases Mouse Brain Atlas (Franklin and Paxinos, 2008). Cell counting was carried out manually using Image J-win64.

**Behavioral task**

For all behavioral tests, experimenters were blinded to genotypes and treatments. Mice were handled daily at least seven days before performing behavioral tests. All the apparatuses and cages were sequentially wiped with 70% ethanol and ddH$_2$O then air-dried between stages. At the end of behavioral tests, mice were perfused with 4% PFA followed by post hoc analysis to confirm the viral injection sites, optic fiber and cannula locations. Mice with incorrect viral injection sites, incorrect positioning of optical fibers or cannulae were excluded.

**Flight-to-nest test**

Flight-to-nest test was performed using previously described methods (Zhou, Z., et al, 2019). Flight-to-nest test was performed in a 40×40×30 cm closed box with a 27-inch LED monitor stationed on top to display the stimulus. A nest in the shape of a 20 cm wide×12 cm high triangular prism was in the corner of the closed Plexiglas box. There are two cameras, one from the top, one from the side (Logitech) that record the mouse's activity simultaneously. Briefly, on day 1, the mice were habituated to the box

conditions for 15 min. On day 2, the mice were first allowed to explore the box for 5-

10 min. When the mice were in the corner furthest from the nest and within in a body-

length distance from the wall, they were given optical stimulus. For optogenetic

activation of LPB$^{CCK}$ neurons or LPB$^{CCK}$-PVN experiments, mice received a 20 s 473

nm blue laser (frequency 20 Hz, pulse width 5 ms) with 15-20 mW (terminal) or 5-10

mW (soma) light power at the fiber tips. For prolonged inhibition of PVN neurons,

Clozapine Noxide (CNO, Sigma-Aldrich) was dissolved in saline to a concentration of

3 mg/ml. The flight-to-nest test was performed 1.5 h after CNO intraperitoneally

injection.

**Flight-to-nest behavioral analysis**

The behaviors of the mice were recorded and analysed automatically with Anymaze

software (Global Biotech Inc). Behavioral analysis was performed as previously

described (Zhou, Z., et al, 2019). Flight-to-nest behavior was characterized on the basis

of the three aspects: latency to return nest, speed (% of baseline speed), time spent in

nest (% of 2 min). Latency to return nest refers to the moment from optical stimulus

onset to moment when the mouse first went into the nest. Speed (% of baseline speed)

refers to the ratio of the post-stimulation speed to the baseline speed. We recorded the

speed of the mice in the 50 seconds before photostimulation presentation as the baseline

speed. The post-stimulation speed was record from the time of stimulation to 15 s after.

It was averaged over a 1 s time window centered on the maximum speed. Time spent

in nest (% of 2 min) refers to the time from mouse's body was first completely under the shelter after photostimulation to the time when the mouse left the nest.

**Heat defense test**

For optogenetic inhibition tests exposing mice to a heat defense environment. We used a translucent plastic box (38×25 cm) divided into two parts, the smaller part is the unheated comfortable zone (5×25 cm), and the larger part with rubber heating pad is the hot uncomfortable zone (33×25 cm). The rubber heating pad was heated at 43℃, and the heat insulated pad in the comfortable zone was placed to avoid the heat. The test involves two stages, the first stage (habituation), the mouse was placed in the hot uncomfortable zone and allowed to explore the whole apparatus freely for 5 min to acclimate. The second stage (test) , the mouse was placed on the edge of the hot uncomfortable zone, while exposed to light（473 nm, 15 mw, 20 s）and allowed to explore for 5 min. In the second stage, the time spent in the hot uncomfortable zone and the latency time from the hot uncomfortable zone to the comfortable zone was calculated by Anymaze software. The heating temperature 43℃ was chosen because it would cause heat escape but not heat pain.

**Open field test**

The open-field chamber was made of plastic (50 × 50 × 50 cm). At the start of the test, mice were placed in the periphery of the open-field chamber. The open field test lasted 5 min.

**Elevated plus-maze test**

The elevated plus maze was made of plastic with two open arms (30×5 cm), two closed

arms (30×5×30 cm) and a central platform (5×5×5 cm). At the beginning of the

experiments, mice were placed in the centeral platform facing a closed arm. The

elevated plus-maze test lasted 5 min.

**Heart rate measurements**

Heart rate was measured via a pulse oximeter (MouseOx Plus; Starr Life Sciences).

Mice were placed in the home cage with a detector fixed around the neck. After

habituation for 10 min, heart rate was simultaneously measured for 10 min while

intermittent 2 min period blue light was applied. Each mouse was tested 3 times, and

the mean heart rate was calculated. Heart rate data was analyzed with the MouseOx

Plus Conscious Applications Software.

**Pupil size measurements**

Mice were adapted to constant room light (100 lx) for 1 h before testing. Mice were

kept unanaesthetized and restrained in a stereotaxic apparatus during the experiment.

The pupil size was recorded using Macro module camera under constant light

conditions before stimulation, during stimulation, and after stimulation. The test lasted

90 s, consisting of 30 s light off-on-off epochs. The pupil size was later measured by

Matlab software.

**Measurements of corticosterone**

Blood samples were collected to determine the hormone levels. Taking Blood was performed in the morning by rapidly collecting heart blood after anesthetization. We immediately collected blood after 10 min of light stimulation (10 mw, 20 Hz, 5 ms pulse width, 15 s per min, 473 nm). Blood samples were temporarily placed in iced plastic tubes coated with heparin. All serum was prepared after every blood sample was centrifuged at 2000 g for 2.5 min at 4°C. Supernatant was collected and plasma corticosterone concentration was measured using commercially-available ELISA kits (Enzo ADI-900-097).

**In vitro electrophysiology**

Each mouse was anesthetized with pentobarbital sodium (100 mg/kg, i.p.) and decapitated. Then the whole brain was quickly dissected into ice-cold oxygenated (95% $O_2$ and 5% $CO_2$) artificial cerebrospinal fluid (aCSF) (93 mM N-methyl-D-glucamine, 2.5 mM KCl, 1.2 mM $NaH_2PO_4$, 20 mM HEPES, 25 mM D-glucose, 30 mM $NaHCO_3$, 10 mM $MgSO_4$, 0.5 mM $CaCl_2$, 5 mM sodium ascorbate, 3 mM sodium pyruvate, and 1 mM kynurenic acid), followed by cutting coronally into 300 μm slices on a microtome (VTA-1200S; Leica). Slices containing the LPB were transferred to a similar solution (93 mM NaCl, 2.5 mM KCl, 1.2 mM $NaH_2PO_4$, 20 mM HEPES, 25 mM D-glucose, 30 mM $NaHCO_3$, 2 mM $MgSO_4$, 2 mM $CaCl_2$, 5 mM sodium ascorbate, 3 mM sodium pyruvate, and 1 mM kynurenic acid), and incubated for at lowest 1 h at room

temperature (24-26°C). Then the brain slices were transferred to a recording chamber

attached to the fixed stage of an BX51WI microscope (Olympus) (solution containing:

125 mM NaCl, 3 mM KCl, 1.25 mM $NaH_2PO_4$, HEPES, 10 mM D-glucose, 26 mM

$NaHCO_3$, 2 mM $MgSO_4$ and 2 mM $CaCl_2$). Patch glass electrodes were pulled from

borosilicate capillaries (BF150-86-10; Sutter Instrument Co, Novato, CA, USA) and

filled with artificial intracellular fluid following component: 135 mM $CsMeSO_3$, 10

mM HEPS, 0.5 mM EGTA, 3.3 mM QX-314, 4 mM Mg-ATP, 0.3 mM $Na_2$-GTP, 8 mM

$Na_2$-Phosphocreatine. Whole-cell voltage-clamp recordings were made with a

MultiClamp 700B amplifier (Molecular Devices). To tested the efficacy of ChR2-

mediated activation, LED-generated blue light pulses were applied to recorded neurons

using 4 different frequencies (5,10, 20, and 40 Hz). To test the effects of photoactivation

of $LPB^{CCK}$ projection terminals within PVN, blue light pulses were applied to the

recorded PVN neurons. To confirm that postsynaptic currents were monosynaptic, the

blue light-evoked currents were recorded in the presence of ACSF (Ctrl), TTX (1 μM)

and 4-AP (100 μM). To confirm that postsynaptic currents were monosynaptic. CNQX

(20 μM) and AP5 (50 μM) were perfused with ACSF to examine the neurotransmitter

type used on $LPB^{CCK}$-PVN projection. Signals were low-pass filtered at 10 kHz and

digitized at 10 kHz (MICRO3 1401, Cambridge Electronic Design). Data were acquired

and analyzed using Spike2 7.04 software (Cambridge Electronic Design).

**Quantification and statistical analysis**

All data analyses were conducted blinded. All statistical analyses were performed with

GraphPad Prism (version 7.0) and analyzed by Unpaired Student's t-tests, one-way ANOVA, two-way ANOVA according to the form of the data. Nonparametric tests were used if the data did not match assumed Gaussian distribution. Animals were randomly assigned to treatment groups. All data were presented as Mean ± SEM, with statistical significance taken as $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$.

**Acknowledgements**

We thank Weisheng Wang, Xiaoming Li, Bin Zhang, Zhi Chen, Kai Liu, Yu-dong Zhou, Huifang Lou for technical support. This work was supported by the National Key Research and Development Program (2021ZD0202700 and 2021ZD0202703), the National Natural Science Foundation of China (82090033) the NSFC-Guangdong Joint Fund (U20A6005), the Key Realm R&D Program of Guangdong province (2019B030335001), the Key Research and Development Program of Zhejiang province (2020C03009 and 2020R01001), the Research Units for Emotion and Emotion Disorders, Chinese Acedemy of Medical Sciences (2019-I2M-5-057).

**Author contributions**

Fan Wang, conceptualization, data curation, investigation, writing original draft, writing-review, revising and editing, designed the experiments, performed the experiments including immunohistochemistry, optogenetics, chemogenetics, behavioral tests, neuropharmacology, collected and analyzed the data, project administration; Yuge Chen, investigation, designed the experiments, performed the experiments including immunohistochemistry, optogenetics, behavioral tests, collected

and analyzed the data, writing-review, revising and editing; Yuxin Lin, Xuze Wang, Performed the experiments including immunohistochemistry, optogenetics, chemogenetics, behavioral tests, neuropharmacology, collected and analyzed the data, writing-review and editing; Kaiyuan Li, Jintao Wu, performed electrophysiology experiments; Yong Han, Zhenggang Zhu writing-review, and editing; Xingyi Shi, Chaoying Long, analyzed the data; Xiaojun Hu, technical support; Shumin Duan, Zhihua Gao, conceptualization, project administration，supervision, conceived and designed the experiments, funding acquisition, writing the manuscript, writing-review and editing.

**Declaration of interests**

The authors declare no competing interests.

**References**

Alhadeff, A. L., Su, Z., Hernandez, E., Klima, M. L., Phillips, S. Z., Holland, R. A., Guo, C., Hantman, A. W., De Jonghe, B. C., & Betley, J. N. (2018, Mar 22). A Neural Circuit for the Suppression of Pain by a Competing Need State. *Cell, 173*(1), 140-152 e115. DOI:10.1016/j.cell.2018.02.057

Anderson, D. J., & Adolphs, R. (2014, Mar 27). A framework for studying emotions across species. *Cell, 157*(1), 187-200. DOI:10.1016/j.cell.2014.03.003

Bertoglio, L. J., de Bortoli, V. C., & Zangrossi, H., Jr. (2007, Jul 2). Cholecystokinin-2

receptors modulate freezing and escape behaviors evoked by the electrical

stimulation of the rat dorsolateral periaqueductal gray. *Brain Res, 1156*, 133-

138. DOI:10.1016/j.brainres.2007.04.038

Blanchard, D. C., Griebel, G., & Blanchard, R. J. (2001, May). Mouse defensive

behaviors: pharmacological and behavioral assays for anxiety and panic.

*Neurosci Biobehav Rev, 25*(3), 205-218. DOI:10.1016/s0149-7634(01)00009-4

Campos, C. A., Bowen, A. J., Han, S., Wisse, B. E., Palmiter, R. D., & Schwartz, M. W.

(2017, Jul). Cancer-induced anorexia and malaise are mediated by CGRP

neurons in the parabrachial nucleus. *Nat Neurosci, 20*(7), 934-942.

DOI:10.1038/nn.4574

Campos, C. A., Bowen, A. J., Roman, C. W., & Palmiter, R. D. (2018, Mar 29).

Encoding of danger by parabrachial CGRP neurons. *Nature, 555*(7698), 617-

622. DOI:10.1038/nature25511

Canteras, N. S., Ribeiro-Barbosa, E. R., & Comoli, E. (2001, Dec). Tracing from the

dorsal premammillary nucleus prosencephalic systems involved in the

organization of innate fear responses. *Neurosci Biobehav Rev, 25*(7-8), 661-668.

DOI:10.1016/s0149-7634(01)00048-3

Carter, M. E., Soden, M. E., Zweifel, L. S., & Palmiter, R. D. (2013, Nov 7). Genetic

identification of a neural circuit that suppresses appetite. *Nature, 503*(7474),

111-114. DOI:10.1038/nature12596

Chen, J. Y., Campos, C. A., Jarvie, B. C., & Palmiter, R. D. (2018, Nov 21). Parabrachial

CGRP Neurons Establish and Sustain Aversive Taste Memories. *Neuron, 100*(4),

891-899 e895. DOI:10.1016/j.neuron.2018.09.032

Chen, L., Liu, Y., Su, P., Hung, W., Li, H., Wang, Y., Yue, Z., Ge, M. H., Wu, Z. X., Zhang, Y., Fei, P., Chen, L. M., Tao, L., Mao, H., Zhen, M., & Gao, S. (2022, Feb 8). Escape steering by cholecystokinin peptidergic signaling. *Cell Rep, 38*(6), 110330. DOI:10.1016/j.celrep.2022.110330

Chiang, M. C., Bowen, A., Schier, L. A., Tupone, D., Uddin, O., & Heinricher, M. M. (2019, Oct 16). Parabrachial Complex: A Hub for Pain and Aversion. *J Neurosci, 39*(42), 8225-8230. DOI:10.1523/JNEUROSCI.1162-19.2019

Cooper Jr, W. E., & Blumstein, D. T. (2015). *Escaping From Predators: Escape decisions prior to pursuit*. DOI:10.1017/cbo9781107447189

Coote, J. H. (2005, Mar). A role for the paraventricular nucleus of the hypothalamus in the autonomic control of heart and kidney. *Exp Physiol, 90*(2), 169-173. DOI:10.1113/expphysiol.2004.029041

Crawford, M., & Masterson, F. A. (1982, Oct-Dec). Species-specific defense reactions and avoidance learning. An evaluative review. *Pavlov J Biol Sci, 17*(4), 204-214. DOI:10.1007/BF03001275

Daviu, N., Fuzesi, T., Rosenegger, D. G., Rasiah, N. P., Sterley, T. L., Peringod, G., & Bains, J. S. (2020, Mar). Paraventricular nucleus CRH neurons encode stress controllability and regulate defensive behavior selection. *Nat Neurosci, 23*(3), 398-410. DOI:10.1038/s41593-020-0591-0

Deisseroth, K. (2014, Jan 16). Circuit dynamics of adaptive and maladaptive behaviour. *Nature, 505*(7483), 309-317. DOI:10.1038/nature12982

Dong, P., Wang, H., Shen, X. F., Jiang, P., Zhu, X. T., Li, Y., Gao, J. H., Lin, S., Huang, Y., He, X. B., Xu, F. Q., Duan, S., Lian, H., Wang, H., Chen, J., & Li, X. M. (2019, Jun). A novel cortico-intrathalamic circuit for flight behavior. *Nat Neurosci, 22*(6), 941-949. DOI:10.1038/s41593-019-0391-6

Eilam, D. (2005). Die hard: a blend of freezing and fleeing as a dynamic defense-- implications for the control of defensive behavior. *Neurosci Biobehav Rev, 29*(8), 1181-1191. DOI:10.1016/j.neubiorev.2005.03.027

Ellard, C. G., & Eller, M. C. (2009, Mar). Spatial cognition in the gerbil: computing optimal escape routes from visual threats. *Anim Cogn, 12*(2), 333-345. DOI:10.1007/s10071-008-0193-9

Evans, D. A., Stempel, A. V., Vale, R., Ruehle, S., Lefler, Y., & Branco, T. (2018, Jun). A synaptic threshold mechanism for computing escape decisions. *Nature, 558*(7711), 590-594. DOI:10.1038/s41586-018-0244-6

Fanselow, M. S., & Bolles, R. C. (1979, Aug). Naloxone and shock-elicited freezing in the rat. *J Comp Physiol Psychol, 93*(4), 736-744. DOI:10.1037/h0077609

Faturi, C. B., Rangel, M. J., Jr., Baldo, M. V., & Canteras, N. S. (2014, May). Functional mapping of the circuits involved in the expression of contextual fear responses in socially defeated animals. *Brain Struct Funct, 219*(3), 931-946. DOI:10.1007/s00429-013-0544-4

Ferguson, A. V., Latchford, K. J., & Samson, W. K. (2008, Jun). The paraventricular nucleus of the hypothalamus - a potential target for integrative treatment of autonomic dysfunction. *Expert Opin Ther Targets, 12*(6), 717-727.

DOI:10.1517/14728222.12.6.717

Fulwiler, C. E., & Saper, C. B. (1984, Aug). Subnuclear organization of the efferent connections of the parabrachial nucleus in the rat. *Brain Res, 319*(3), 229-259. DOI:10.1016/0165-0173(84)90012-2

Garfield, A. S., Shah, B. P., Madara, J. C., Burke, L. K., Patterson, C. M., Flak, J., Neve, R. L., Evans, M. L., Lowell, B. B., Myers, M. G., Jr., & Heisler, L. K. (2014, Dec 2). A parabrachial-hypothalamic cholecystokinin neurocircuit controls counterregulatory responses to hypoglycemia. *Cell Metab, 20*(6), 1030-1037. DOI:10.1016/j.cmet.2014.11.006

Gauriau, C., & Bernard, J. F. (2002, Mar). Pain pathways and parabrachial circuits in the rat. *Exp Physiol, 87*(2), 251-258. DOI:10.1113/eph8702357

Geerling, J. C., Kim, M., Mahoney, C. E., Abbott, S. B., Agostinelli, L. J., Garfield, A. S., Krashes, M. J., Lowell, B. B., & Scammell, T. E. (2016, Jan 1). Genetic identity of thermosensory relay neurons in the lateral parabrachial nucleus. *Am J Physiol Regul Integr Comp Physiol, 310*(1), R41-54. DOI:10.1152/ajpregu.00094.2015

Gross, C. T., & Canteras, N. S. (2012, Sep). The many paths to fear. *Nat Rev Neurosci, 13*(9), 651-658. DOI:10.1038/nrn3301

Han, S., Soleiman, M. T., Soden, M. E., Zweifel, L. S., & Palmiter, R. D. (2015, Jul 16). Elucidating an Affective Pain Circuit that Creates a Threat Memory. *Cell, 162*(2), 363-374. DOI:10.1016/j.cell.2015.05.057

Jamalamadaka, T., Griffith, E., Steer, H., & Salkovskis, P. (2020, Sep). Fear of illness

recurrence and mental health anxiety in people recovering from psychosis and common mental health problems. *Br J Clin Psychol, 59*(3), 403-423. DOI:10.1111/bjc.12253

Kaur, S., Pedersen, N. P., Yokota, S., Hur, E. E., Fuller, P. M., Lazarus, M., Chamberlin, N. L., & Saper, C. B. (2013, May 1). Glutamatergic signaling from the parabrachial nucleus plays a critical role in hypercapnic arousal. *J Neurosci, 33*(18), 7627-7640. DOI:10.1523/JNEUROSCI.0173-13.2013

Keay, K. A., & Bandler, R. (2001, Dec). Parallel circuits mediating distinct emotional coping reactions to different types of stress. *Neurosci Biobehav Rev, 25*(7-8), 669-678. DOI:10.1016/s0149-7634(01)00049-5

Kobayashi, A., & Osaka, T. (2003, Sep). Involvement of the parabrachial nucleus in thermogenesis induced by environmental cooling in the rat. *Pflugers Arch, 446*(6), 760-765. DOI:10.1007/s00424-003-1119-7

Krout, K. E., & Loewy, A. D. (2000, Dec 18). Parabrachial nucleus projections to midline and intralaminar thalamic nuclei of the rat. *J Comp Neurol, 428*(3), 475-494. DOI:10.1002/1096-9861(20001218)428:3<475::aid-cne6>3.0.co;2-9

LeDoux, J. (2012, Mar 8). Rethinking the Emotional Brain (vol 73, pg 653, 2012). *Neuron, 73*(5), 1052-1052. DOI:10.1016/j.neuron.2012.02.018

Liu, X., Dimidschstein, J., Fishell, G., & Carter, A. G. (2020, Oct 9). Hippocampal inputs engage CCK+ interneurons to mediate endocannabinoid-modulated feed-forward inhibition in the prefrontal cortex. *Elife, 9*. DOI:10.7554/eLife.55267

Luthi, A., & Luscher, C. (2014, Dec). Pathological circuit function underlying addiction

and anxiety disorders. *Nat Neurosci, 17*(12), 1635-1643. DOI:10.1038/nn.3849

Mangieri, L. R., Jiang, Z., Lu, Y., Xu, Y., Cassidy, R. M., Justice, N., Xu, Y., Arenkiel, B. R., & Tong, Q. (2019a, Jul/Aug). Defensive Behaviors Driven by a Hypothalamic-Ventral Midbrain Circuit. *eNeuro, 6*(4). DOI:10.1523/ENEURO.0156-19.2019

Mangieri, L. R., Jiang, Z., Lu, Y., Xu, Y., Cassidy, R. M., Justice, N., Xu, Y., Arenkiel, B. R., & Tong, Q. (2019b, Jul/Aug). Defensive Behaviors Driven by a Hypothalamic-Ventral Midbrain Circuit. *eNeuro, 6*(4), ENEURO.0156-0119.2019. DOI:10.1523/ENEURO.0156-19.2019

Martinez, R. C., Carvalho-Netto, E. F., Amaral, V. C., Nunes-de-Souza, R. L., & Canteras, N. S. (2008, Oct 10). Investigation of the hypothalamic defensive system in the mouse. *Behav Brain Res, 192*(2), 185-190. DOI:10.1016/j.bbr.2008.03.042

Mu, D., Deng, J., Liu, K. F., Wu, Z. Y., Shi, Y. F., Guo, W. M., Mao, Q. Q., Liu, X. J., Li, H., & Sun, Y. G. (2017, Aug 18). A central neural circuit for itch sensation. *Science, 357*(6352), 695-699. DOI:10.1126/science.aaf4918

Nakamura, K., & Morrison, S. F. (2008, Jan). A thermosensory pathway that controls body temperature. *Nat Neurosci, 11*(1), 62-71. DOI:10.1038/nn2027

Nakamura, K., & Morrison, S. F. (2010, May 11). A thermosensory pathway mediating heat-defense responses. *Proc Natl Acad Sci U S A, 107*(19), 8848-8853. DOI:10.1073/pnas.0913358107

Norris, A. J., Shaker, J. R., Cone, A. L., Ndiokho, I. B., & Bruchas, M. R. (2021, Mar

5). Parabrachial opioidergic projections to preoptic hypothalamus mediate behavioral and physiological thermal defenses. *Elife, 10*. DOI:10.7554/eLife.60779

O'Shea, R. D., & Gundlach, A. L. (1993, Dec). Regulation of cholecystokinin receptors in the hypothalamus of the rat: reciprocal changes in magnocellular nuclei induced by food deprivation and dehydration. *J Neuroendocrinol, 5*(6), 697-704. DOI:10.1111/j.1365-2826.1993.tb00542.x

Perusini, J. N., & Fanselow, M. S. (2015, Sep). Neurobehavioral perspectives on the distinction between fear and anxiety. *Learn Mem, 22*(9), 417-425. DOI:10.1101/lm.039180.115

Pitman, R. K., Rasmusson, A. M., Koenen, K. C., Shin, L. M., Orr, S. P., Gilbertson, M. W., Milad, M. R., & Liberzon, I. (2012, Nov). Biological studies of post-traumatic stress disorder. *Nat Rev Neurosci, 13*(11), 769-787. DOI:10.1038/nrn3339

Salay, L. D., Ishiko, N., & Huberman, A. D. (2018, May). A midline thalamic circuit determines reactions to visual threat. *Nature, 557*(7704), 183-189. DOI:10.1038/s41586-018-0078-2

Shang, C., Chen, Z., Liu, A., Li, Y., Zhang, J., Qu, B., Yan, F., Zhang, Y., Liu, W., Liu, Z., Guo, X., Li, D., Wang, Y., & Cao, P. (2018, Mar 26). Divergent midbrain circuits orchestrate escape and freezing responses to looming stimuli in mice. *Nat Commun, 9*(1), 1232. DOI:10.1038/s41467-018-03580-7

Shang, C., Liu, Z., Chen, Z., Shi, Y., Wang, Q., Liu, S., Li, D., & Cao, P. (2015, Jun 26).

BRAIN CIRCUITS. A parvalbumin-positive excitatory visual pathway to trigger fear responses in mice. *Science, 348*(6242), 1472-1477. DOI:10.1126/science.aaa8694

Silva, B. A., Mattucci, C., Krzywkowski, P., Murana, E., Illarionova, A., Grinevich, V., Canteras, N. S., Ragozzino, D., & Gross, C. T. (2013, Dec). Independent hypothalamic circuits for social and predator fear. *Nat Neurosci, 16*(12), 1731-1733. DOI:10.1038/nn.3573

Staples, L. G., McGregor, I. S., Apfelbach, R., & Hunt, G. E. (2008, Feb 19). Cat odor, but not trimethylthiazoline (fox odor), activates accessory olfactory and defense-related brain regions in rats. *Neuroscience, 151*(4), 937-947. DOI:10.1016/j.neuroscience.2007.11.039

Sun, L., Liu, R., Guo, F., Wen, M. Q., Ma, X. L., Li, K. Y., Sun, H., Xu, C. L., Li, Y. Y., Wu, M. Y., Zhu, Z. G., Li, X. J., Yu, Y. Q., Chen, Z., Li, X. Y., & Duan, S. (2020, Nov 25). Parabrachial nucleus circuit governs neuropathic pain-like behavior. *Nat Commun, 11*(1), 5974. DOI:10.1038/s41467-020-19767-w

Sun, X. C., Bernstein, M. J., Meng, M. Z., Rao, S. Y., Sorensen, A. T., Yao, L., Zhang, X. H., Anikeeva, P. O., & Lin, Y. X. (2020, Apr 16). Functionally Distinct Neuronal Ensembles within the Memory Engram. *Cell, 181*(2), 410-+. DOI:10.1016/j.cell.2020.02.055

Sutton, A. K., Myers, M. G., Jr., & Olson, D. P. (2016). The Role of PVH Circuits in Leptin Action and Energy Balance. *Annu Rev Physiol, 78*(1), 207-221. DOI:10.1146/annurev-physiol-021115-105347

Tovote, P., Esposito, M. S., Botta, P., Chaudun, F., Fadok, J. P., Markovic, M., Wolff, S. B., Ramakrishnan, C., Fenno, L., Deisseroth, K., Herry, C., Arber, S., & Luthi, A. (2016, Jun 9). Midbrain circuits for defensive behaviour. *Nature, 534*(7606), 206-212. DOI:10.1038/nature17996

Tovote, P., Meyer, M., Ronnenberg, A., Ogren, S. O., Spiess, J., & Stiedl, O. (2005). Heart rate dynamics and behavioral responses during acute emotional challenge in corticotropin-releasing factor receptor 1-deficient and corticotropin-releasing factor-overexpressing mice. *Neuroscience, 134*(4), 1113-1122. DOI:10.1016/j.neuroscience.2005.05.027

Uddin, O., Studlack, P., Akintola, T., Raver, C., Castro, A., Masri, R., & Keller, A. (2018, Jan-Jul). Amplified parabrachial nucleus activity in a rat model of trigeminal neuropathic pain. *Neurobiol Pain, 3*, 22-30. DOI:10.1016/j.ynpai.2018.02.002

Vianna, D. M., Landeira-Fernandez, J., & Brandão, M. L. (2001, Dec). Dorsolateral and ventral regions of the periaqueductal gray matter are involved in distinct types of fear. *Neurosci Biobehav Rev, 25*(7-8), 711-719. DOI:10.1016/s0149-7634(01)00052-5

Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., Jr., & Lowell, B. B. (2011, Jul 14). Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron, 71*(1), 142-154. DOI:10.1016/j.neuron.2011.05.028

Wang, L., Chen, I. Z., & Lin, D. (2015, Mar 18). Collateral pathways from the ventromedial hypothalamus mediate defensive behaviors. *Neuron, 85*(6), 1344-1358. DOI:10.1016/j.neuron.2014.12.025

Wang, W., Schuette, P. J., La-Vu, M. Q., Torossian, A., Tobias, B. C., Ceko, M., Kragel,

P. A., Reis, F. M., Ji, S., Sehgal, M., Maesta-Pereira, S., Chakerian, M., Silva,

A. J., Canteras, N. S., Wager, T., Kao, J. C., & Adhikari, A. (2021, Sep 1). Dorsal

premammillary projection to periaqueductal gray controls escape vigor from

innate and conditioned threats. *Elife, 10*. DOI:10.7554/eLife.69178

Wang, W., Schuette, P. J., Nagai, J., Tobias, B. C., Cuccovia, V. R. F. M., Ji, S., de Lima,

M. A. X., La-Vu, M. Q., Maesta-Pereira, S., Chakerian, M., Leonard, S. J., Lin,

L., Severino, A. L., Cahill, C. M., Canteras, N. S., Khakh, B. S., Kao, J. C., &

Adhikari, A. (2021, Jun 2). Coordination of escape and spatial navigation

circuits orchestrates versatile flight from threats. *Neuron, 109*(11), 1848-1860

e1848. DOI:10.1016/j.neuron.2021.03.033

Wei, P., Liu, N., Zhang, Z., Liu, X., Tang, Y., He, X., Wu, B., Zhou, Z., Liu, Y., Li, J.,

Zhang, Y., Zhou, X., Xu, L., Chen, L., Bi, G., Hu, X., Xu, F., & Wang, L. (2015,

Apr 9). Processing of visually evoked innate fear by a non-canonical thalamic

pathway. *Nat Commun, 6*, 6756. DOI:10.1038/ncomms7756

Whissell, P. D., Cajanding, J. D., Fogel, N., & Kim, J. C. (2015, Sep 23). Comparative

density of CCK- and PV-GABA cells within the cortex and hippocampus.

*Frontiers in Neuroanatomy, 9*, 124. DOI:ARTN 124

10.3389/fnana.2015.00124

Xu, Y., Lu, Y., Cassidy, R. M., Mangieri, L. R., Zhu, C., Huang, X., Jiang, Z., Justice,

N. J., Xu, Y., Arenkiel, B. R., & Tong, Q. (2019, Aug 1). Identification of a

neurocircuit underlying regulation of feeding by stress-related emotional

responses. *Nat Commun, 10*(1), 3446. DOI:10.1038/s41467-019-11399-z

Xu, Y., Wu, Z., Sun, H., Zhu, Y., Kim, E. R., Lowell, B. B., Arenkiel, B. R., Xu, Y., & Tong, Q. (2013, Dec 3). Glutamate mediates the function of melanocortin receptor 4 on Sim1 neurons in body weight regulation. *Cell Metab, 18*(6), 860-870. DOI:10.1016/j.cmet.2013.11.003

Yang, H., Yang, J., Xi, W., Hao, S., Luo, B., He, X., Zhu, L., Lou, H., Yu, Y. Q., Xu, F., Duan, S., & Wang, H. (2016, Feb). Laterodorsal tegmentum interneuron subtypes oppositely regulate olfactory cue-induced innate fear. *Nat Neurosci, 19*(2), 283-289. DOI:10.1038/nn.4208

Yang, W. Z., Du, X., Zhang, W., Gao, C., Xie, H., Xiao, Y., Jia, X., Liu, J., Xu, J., Fu, X., Tu, H., Fu, X., Ni, X., He, M., Yang, J., Wang, H., Yang, H., Xu, X. H., & Shen, W. L. (2020, Sep). Parabrachial neuron types categorically encode thermoregulation variables during heat defense. *Sci Adv, 6*(36). DOI:10.1126/sciadv.abb9414

Yilmaz, M., & Meister, M. (2013, Oct 21). Rapid innate defensive responses of mice to looming visual stimuli. *Curr Biol, 23*(20), 2011-2015. DOI:10.1016/j.cub.2013.08.015

Zhang, B., Wang, W., Zhang, Z., Hu, Y., Meng, F., Wang, F., Lou, H., Zhu, L., Godbout, R., Duan, S., & Gao, Z. (2018, Oct 1). Alternative Splicing of Disabled-1 Controls Multipolar-to-Bipolar Transition of Migrating Neurons in the Neocortex. *Cereb Cortex, 28*(10), 3457-3467. DOI:10.1093/cercor/bhx212

Zhang, S., Lv, F., Yuan, Y., Fan, C., Li, J., Sun, W., & Hu, J. (2019). Whole-Brain

Mapping of Monosynaptic Afferent Inputs to Cortical CRH Neurons. *Front*

*Neurosci, 13*, 565. DOI:10.3389/fnins.2019.00565–

Zhou, Z., Liu, X., Chen, S., Zhang, Z., Liu, Y., Montardy, Q., Tang, Y., Wei, P., Liu, N.,

Li, L., Song, R., Lai, J., He, X., Chen, C., Bi, G., Feng, G., Xu, F., & Wang, L.

(2019, Aug 7). A VTA GABAergic Neural Circuit Mediates Visually Evoked

Innate    Defensive    Responses.    *Neuron,    103*(3),    473-488    e476.

DOI:10.1016/j.neuron.2019.05.027

**Figure Legends**

**Figure 1.** CCK$^+$ neurons in the LPB project to the PVN. **(A)** Scheme of viral strategy for retrograde tracing from the PVN in CCK-Cre mice using AAV2-Retro virus. **(B)** CCK-Cre mice were injected with AAV2-Retro-EYFP virus at the PVN. And traced neurons (EYFP$^+$) in the LPB, PAG; scale bar, 100 μm; n=3 mice. **(C)** A heatmap demonstrating the distribution of EYFP$^+$ neurons in the MO, CC, DI, PAG. The number of cells in individual areas are color coded as indicated; n = 3 mice. **(D)** Scheme of viral strategy for anterograde tracing from LPB$^{CCK}$ neurons using AAV-hSyn-FLEX-mGFP-2A-Synaptophysin-mRuby virus. **(E)** Representative histological images of AAV-hSyn-FLEx-mGFP-2A-Synaptophysin-mRuby virus expression in the CCK neurons in the LPB; scale bar, 100 μm. **(F)** LPB$^{CCK}$ neurons have projections to the PVN; scale bar, 100 μm. The right panel shows a magnified view of rectangular part; scale bar, 100 μm; n=3 mice.

**Figure 2.** CCK$^+$ neurons in the LPB are mostly glutamatergic neurons and send glutamatergic projections to the PVN. **(A)** Schematic of mice crossing to generate a CCK-ires-Cre::Ai14 mouse line. **(B)** Representative coronal section images illustrating the expression of CCK-tdT in the LPB; scale bar, 100 μm. **(C)** Quantification of CCK-tdT cells across the rostral-to-caudal extent of the LPB; n=3 mice. **(D)** Representative images showing co-labeling (arrowheads) of Vglut2-positive (Green) and CCK-tdT expressing cells in the LPB. Rightmost panel, magnification of the box-shaped area in the left panel (white dashed lines). Blue, DAPI staining. Scale bar, 100 μm. **(E)**

Example images showing Vgat-positive (Green) and CCK-tdT-expressing cells in the

LPB. Rightmost panel, magnification of the boxed area in the left panel. Blue, DAPI

staining. Scale bar, 100 μm. **(F)** Representative image from a CCK-ires-Cre::Ai14

mouse in the LPB region stained with antibodies against CGRP. Rightmost panels,

magnification of the boxed areas in the left panels. Blue, DAPI staining. Scale bar, 100

μm. **(G)** Percentage of CCK-positive neurons that co-localized with markers; n=4 mice.

**(H)** Schematic of recording from PVN cells following optogenetic activation of

LPB$^{CCK}$ axonal terminals. **(I-J)** Representative traces of light-evoked EPSCs recorded

from PVN neurons following light stimulation of LPB$^{CCK}$ axonal terminals in the

presence of ACSF(Ctrl), TTX (1 μM) and 4- AP (100 μM). Representative traces of

light-evoked EPSCs recorded from PVN neurons following light stimulation of LPB$^{CCK}$

axonal terminals in the presence of CNQX (20 μM) and AP5 (50 μM) (n = 9 neurons).

**(K)** Quantification of excitatory postsynaptic currents (EPSCs) and inhibitory

postsynaptic currents (IPSCs) from identified PVN neurons receiving inputs from the

LPB$^{CCK}$ neurons.

1024

**Figure 3.** Activation of LPB$^{CCK}$ neurons triggers defensive-like flight-to-nest behavior.

**(A)** Left, schematic of light stimulation of and patch-clamp recording from ChR2-EYFP

expressing CCK neurons in the LPB. Right, example of action potentials evoked by

optogenetic stimulation LPB$^{CCK}$ neurons from using whole cell patch-clamp slice

recording. **(B)** Schematic diagram of optogenetic activation of LPB$^{CCK}$ neurons. **(C)**

Representative image showing the ChR2-EYFP expression and optical fiber tip

locations in LPB of a CCK-Cre mouse. Scale bar, 100 μm. **(D)** Schematic of the timing

and behavioral paradigm of flight-to-nest behavior test. **(E)** Experimental apparatus:

Open field with a nest in the corner. **(F)** Schematic of the timing and behavioral

paradigm with optical activation of LPB$^{CCK}$ neurons. **(G)** Optogenetic activation of

LPB$^{CCK}$ neurons induced defensive-like flight-to-nest behavior. Mice movements were

tracked in the experimental apparatus. **(H-J)** Compared with control group, ChR2 group

showed shorter latency, higher speed back into the nest and higher percentage of hiding

time spent in the nest (EYFP: n =7 mice, ChR2: n =7 mice; for latency, p = 0.0006, U

= 0; Mann-Whitney test; for speed, p = 0.0016, t = 4.052, df = 12; unpaired t test; for

time in the nest, p < 0.0001, t = 19.82, df = 12; unpaired t test).

**Figure 4.** Photostimulation of LPB$^{CCK}$ neurons induces anxiety-like behavior and

autonomic responses. **(A)** Schematic of behavioral test protocol. OFT, open field test;

EPM, elevated plus maze test. **(B)** Movement traces of a control group mouse (AAV2/9-

EF1a-DIO-EYFP, left) and an experimental group mouse (AAV2/9-EF1a-DIO-ChR2-

EYFP, right) in the open field. **(C)** Quantification of the total distance moved, time

spent in center and total number of center entries in the open field tests before (Pre) and

after (Post) photostimulation of LPB$^{CCK}$ neurons in CCK-Cre mice.(EYFP: n = 7 mice,

ChR2: n = 7 mice; for total distance, Pre-light EYFP vs. Post-light EYFP ns p > 0.9999,

Post-light EYFP vs. Post-light ChR2 ns p = 0.2236, Pre-light ChR2 vs. Post-light ChR2

ns p = 0.9026; two-way ANOVA, Bonferroni's multiple comparisons test; for center

time, Pre-light EYFP vs. Post-light EYFP ns p > 0.9999, Post-light EYFP vs. Post-light

ChR2 **p < 0.01 p=0.005, Pre-light ChR2 vs. Post-light ChR2 **p < 0.01 p = 0.005;

two-way ANOVA, Bonferroni's multiple comparisons test; for center entries, Pre-light

EYFP vs. Post-light EYFP ns p > 0.9999, Post-light EYFP vs. Post-light ChR2 ***p <

0.0005 p = 0.0004, Pre-light ChR2 vs. Post-light ChR2 **p < 0.01 p = 0.0042; two-way

ANOVA, Bonferroni's multiple comparisons test). **(D)** Example traces of a control

group mouse (AAV2/9-EF1a-DIO-EYFP, left) and an experimental group mouse

(AAV2/9-EF1a-DIO-EYFP, right) in the elevated plus maze. **(E)** Quantification of the

open arm time and open arm entries in the elevated plus maze tests before (Pre) and

after (Post) photostimulation of LPB$^{CCK}$ neurons in CCK-Cre mice. (EYFP: n = 7 mice,

ChR2: n = 7 mice; for open arm time, Pre-light EYFP vs. Post-light EYFP ns p > 0.9999,

Post-light EYFP vs. Post-light ChR2 **p < 0.01 p = 0.0053, Pre-light ChR2 vs. Post-

light ChR2 ***p<0.0005 p = 0.0005; two-way ANOVA, Bonferroni's multiple

comparisons test; for open arm entries, Pre-light EYFP vs. Post-light EYFP ns p >

0.9999, Post-light EYFP vs. Post-light ChR2 **p < 0.01 p = 0.0159, Pre-light ChR2 vs.

Post-light ChR2 ****p < 0.0001; two-way ANOVA, Bonferroni's multiple

comparisons test). **(F)** Relative heart rate changes induced by photostimulation of

LPB$^{CCK}$ neurons. (EYFP: n = 7 mice, ChR2: n = 7 mice; p = 0.0006, t = 4.603, df = 12;

unpaired t test). **(G)** Schematic of detection of pupil size. **(H)** Example image of

computer-detected pupil before, during and after photo stimulation of LPB$^{CCK}$ neurons.

**(I)** Relative pupil size (during photostimulation/before photostimulation) of LPB$^{CCK}$

neurons. (EYFP: n = 6 mice, ChR2: n = 6 mice; p = 0.0022, U = 0; Mann-Whitney test).

**(J)** Cartoon of the arousal state during the activation of LPB$^{CCK}$ neurons. **(K)** Serum

1075 corticosterone levels in EYFP and ChR2 groups. (EYFP: n = 5 mice, ChR2: n = 5 mice;

1076 p = 0.0121, t = 3.23, df = 8; unpaired t test).

1077

**Figure 5.** Optogenetic inhibition of LPB$^{CCK}$ neurons suppressed heat-evoked defensive

behavior. **(A)** Left, schematic of light stimulation of and patch-clamp recording from

GtACR1-expressing CCK neurons in the LPB. Right, example of action potentials

evoked by optogenetic inhibition of LPB$^{CCK}$ neurons from using whole cell patch-

clamp slice recording. **(B)** Schematic diagram of optogenetic inhibition of LPB$^{CCK}$

neurons. **(C)** Representative image showing the GtACR1-EYFP expression in the LPB

and optical fiber tip locations above the LPB of a CCK-Cre mouse. **(D)** Heat defense

apparatus: a translucent plastic box divided into two parts, the smaller part is the

unheated comfortable zone, and the larger part with rubber heating pad is the hot

uncomfortable zone. **(E)** Schematic of the timing and behavioral paradigm of heat-

evoked defensive behavioral test. **(F-G)** Photoinhibition of LPB$^{CCK}$ neurons suppressed

heat-evoked defensive behavior. Compared with control group, GtACR1 group showed

longer latency and lower percentage of time spent in the hot zone (EYFP: n = 8 mice,

GtACR1: n = 7 mice; for latency, p = 0.0059, U = 2; Mann-Whitney test; for time in

the nest, p = 0.02, U = 8; Mann-Whitney test).

1093

**Figure 6.** Photostimulation of LPB$^{CCK}$-PVN pathway induces defensive-like flight-to-

nest behavior. **(A)** Schematic diagram of optogenetic activation of LPB$^{CCK}$-PVN

neurons. **(B)** Left, Representative image showing the ChR2-EYFP expression in the

LPB of a CCK-Cre mouse. Right, Optical fiber tip locations in the PVN from a ChR2-

EYFP expressing CCK-Cre mouse. **(C)** Experimental apparatus: Open field with nest

in the corner. **(D)** Schematic of the timing and behavioral paradigm with optical

activation of LPB$^{CCK}$-PVN. **(E)** Optogenetic activation of LPB$^{CCK}$-PVN induced

defensive-like flight-to-nest behavior. Mice movements were tracked in the

experimental apparatus. **(F-H)** Optogenetic activation of LPB$^{CCK}$-PVN induced

defensive-like flight-to-nest behavior. Compared with control group, ChR2 group

showed shorter latency, higher speed back into the nest and higher percentage of hiding

time spent in the nest (EYFP: n = 7 mice, ChR2: n = 7 mice; for latency, p = 0.0012, U

= 0; Mann-Whitney test; for speed, p = 0.0013, t = 4.163, df = 12; unpaired t test; for

time in the nest, p = 0.0006, U = 0; Mann-Whitney test). **(I)** C-fos expression in the

PVN with blue light activation of LPB-originating dio-EYFP (left) or ChR2-EYFP

(right) expressing terminals in the PVN. Scale bar: 100 μm. **(J)** Quantification of c-fos

positive cells in the PVN following ChR2 activation (EYFP: n = 3 mice, ChR2: n = 3

mice; p = 0.0033, t = 6.253, df = 4; unpaired t test). **(K)** Relative heart rate in mice with

and without LPB$^{CCK}$-PVN projection stimulation (EYFP: n = 7 mice, ChR2: n = 7 mice;

P = 0.0002, t = 5.249, df = 12; unpaired t test). **(L)** Serum corticosterone levels in EYFP

and ChR2 groups. (EYFP: n = 5 mice, ChR2: n = 5 mice; P = 0.0079, U = 0; Mann-

Whitney test).

1116

1117 **Figure 7.** PVN, as a downstream nuclei of LPB$^{CCK}$ neurons, is involved in defensive-

1118 like flight-to-nest behavior. **(A)** Schematics of experimental setup. CCK-Cre mice with

AAV2/9-DIO-ChR2-EYFP virus delivery to the LPB, AAV2/9-hsyn-hM4Di-mCherry virus delivery to the PVN, and optic fiber implantation targeting PVN were used for flight- to-nest behavior test. **(B)** Left, representative image showing the ChR2-EYFP expression in the LPB of a CCK-Cre mouse. Scale bar, 100 μm. Right, representative image showing the hM4Di-mCherry expression in the PVN and optic fiber placement in the PVN from a ChR2-EYFP expressing mouse. **(C)** Schematic of behavioral test protocol. **(D-F)** Chemogenetic inhibition of PVN neurons before optical activation of LPB$^{CCK}$-PVN showed longer latency and lower speed returning to the nest and lower percentage of hiding time spent in the nest (saline: n = 7 mice, CNO: n = 7 mice; for latency, p = 0.0012, U = 1; Mann-Whitney test; for speed, p < 0.0001, t = 5.798, df = 12; unpaired t test; for time in the nest, p = 0.0031, t = 3.68, df = 12; unpaired t test). **(G-H)** Expression of CCKaR and CCKbR throughout the PVN. Representative histological images showing CCKaR (left) and CCKbR (right) in the PVN. **(I)** Schematics of experimental setup. Optogenetic activation of LPB$^{CCK}$-PVN with a cannula implanted in the PVN for aCSF or CNQX + AP5 or Devazepide + L-365,260 delivery. Scale bar, 100 μm. **(J)** Left, representative image showing the ChR2-EYFP expression in the LPB of a CCK-Cre mouse. Scale bar, 100 μm. Right, representative image showing the cannula placement in the PVN from a ChR2-EYFP expressing CCK-Cre mouse. Scale bar, 100 μm. **(K)** Schematic of behavioral test protocol. **(L-N)** Compared with the ACSF group, microinjection of the glutamatergic receptor antagonist rather than CCK receptor antagonist into the PVN showed longer latency returning to the nest and no change in the speed back into the nest and percentage of

1141    hiding time spent in the nest (ACSF: n = 6 mice, CNQX + AP5: n = 6 mice, Devazepide

1142    + L-365260: n = 5 mice; for latency, $F_{(2,14)}$ = 7.658, p=0.057; One-way ANOVA; for

1143    speed, $F_{(2,14)}$ = 11.49, p = 0.0011; for time in the nest, One-way ANOVA; $F_{(2,14)}$

1144    =16.44, p = 0.0002; One-way ANOVA).

1145

**Figure 8.** Photoactivation of $PVN^{Vglut2}$ neurons rather than $PVN^{CRH}$ neurons triggers

defensive-like flight-to-nest behavior. **(A)** Schematic diagram of optogenetic activation

of $PVN^{Vglut2}$ neurons. **(B)** Representative image showing the ChR2-EYFP expression

and optical fiber tip locations in the PVN of a Vglut2-Cre mouse. Scale bar, 100 μm.

**(C-E)** Optogenetic activation of $PVN^{Vglut2}$ neurons induced defensive-like flight-to-nest

behavior. Compared with control group, ChR2 group showed shorter latency, higher

speed back into the nest and higher percentage of hiding time spent in the nest. (EYFP:

n = 6 mice, ChR2: n = 6 mice; for latency, p = 0.0065, u = 2; Mann-Whitney test; for

speed, p = 0.0221, t = 2.705, df = 10; unpaired t test; for time in the nest, p = 0.022, u

= 0; Mann-Whitney test). **(F)** Schematic diagram of optogenetic activation of $PVN^{CRH}$

neurons. **(G)** Representative image showing the ChR2-EYFP expression and optical

fiber tip locations in the PVN of a CRH-Cre mouse. Scale bar, 100 μm. **(H-J)**

Optogenetic activation of $PVN^{CRH}$ neurons did not induce flight-to-nest behavior.

Compared with control group, ChR2 group showed no changes in latency, speed back

into the nest and total percentage of hiding time spent in the nest (EYFP: n = 5 mice,

ChR2: n = 6 mice; for latency, p = 0.6664, t = 0.4456, df = 9; unpaired t test; for speed,

p = 0.0221, t = 2.705, df = 10; unpaired t test; for time in the nest, p = 0.2088, t = 1.367,

df = 8; unpaired t test).

1164

1165    **Figure 9.** Schematic diagram of the working model. Graphical summary. These studies

1166    suggest a novel circuit that mediates defensive-like flight-to-nest behavior.

1167

1168

1169    **Supplementary figure legends**

1170    **Figure 1-figure supplement 1.** Upstream nuclei of LPB$^{CCK}$ neurons in the whole brain.

1171    **(A-D)** CCK-Cre mice with PVN delivery AAV2-Retro-EYFP virus and EYFP$^{+}$ neurons

1172    in the MO, CC, DI, PAG; scale bar, 100 μm; n = 3 mice.

1173

**Figure1-figure supplement 2. (A-I)** Projection mapping from LPB$^{CCK}$ neurons.

Representative images showing projections from LPB$^{CCK}$ neurons in the median

preoptic nucleus (MnPO), lateral preoptic area (LPO), paraventricular thalamic nucleus

(PVT), ventromedial hypothalamic nucleus (VMH), ventral posteromedial thalamic

nucleus (VPM), lateral posterior thalamic nucleus, mediorostral part (LPMR),

parasubthalamic nucleus (PSTh), PAG, deep gray layer of the superior colliculus (DpG）

(n = 3 mice; green, GFP; red, mrubby; blue, DAPI). Scale bar, 100 μm; n = 3 mice.

1181

1182    **Source data files**

1183    **Figure 1-source data 1.** Data for Figure 1. The number of EYFP$^{+}$ neurons in the MO,

1184    CC, DI, PAG, LPB.

**Figure 2-source data 1.** Data for Figure 2. Quantification of CCK-tdT cells across the rostral-to-caudal extent of the LPB; Percentage of CCK-positive neurons that co-localized with markers; Quantification of excitatory postsynaptic currents (EPSCs) and inhibitory postsynaptic currents (IPSCs) from identified PVN neurons.

**Figure 3-source data 1.** Data for Figure 3. Values for latency, speed back into the nest and percentage of hiding time spent in the nest during flight-to-nest behavior test.

**Figure 4-source data 1.** Data for Figure 4. Quantification of the total distance moved, time spent in center and total number of center entries in the open field tests; Quantification of the open arm time and open arm entries in the elevated plus maze tests; Values for relative heart rate changes; Values for relative pupil size; Values for serum corticosterone levels in EYFP and ChR2 groups.

**Figure 5-source data 1.** Data for Figure 5. Values for suppressed heat-evoked defensive behavior.

**Figure 6-source data 1.** Data for Figure 6. Values for latency, speed back into the nest and percentage of hiding time spent in the nest during flight-to-nest behavior test; Quantification of c-fos positive cells in the PVN; Values for relative heart rate; Values for serum corticosterone levels in EYFP and ChR2 groups.

**Figure 7-source data 1.** Data for Figure 7. Values for latency, speed back into the nest and percentage of hiding time spent in the nest during flight-to-nest behavior test;

**Figure 8-source data 1.** Data for Figure 8. Values for latency, speed back into the nest and percentage of hiding time spent in the nest during flight-to-nest behavior test.

**Figure 1**

1223 **Figure 2**

1224
1225

1226

1227

1228    **Figure 3**

1229

1230

1231

1232    **Figure 4**

1233

1234

1235

1236 **Figure 5**

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251    **Figure 6**

1252

1253

1254

1255    **Figure 7**

1256

1257

1258

 **Figure 8**

1263 **Figure 9**

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277 **Figure 1–figure supplement 1**

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

 **Figure 1–figure supplement 2**