Effective conservation decisions require models designed for purpose: a case study for

boreal caribou in Ontario's Ring of Fire

Matt Dyson[1], Sarah Endicott[2], Craig Simpkins[1], Julie W. Turner[3], Stephanie Avery-Gomm[2],

Cheryl A. Johnson[2], Mathieu Leblond[2], Eric W. Neilson[4], Rob Rempel[5], Philip A. Wiebe[6],

Jennifer L. Baltzer[1], Josie Hughes[2*], Frances E.C. Stewart[1*]

[1]Biology Department, Wilfrid Laurier University, Waterloo, ON, Canada.

matt.e.dyson@gmail.com; simpkinscraig063@gmail.com; jbaltzer@wlu.ca; fstewart@wlu.ca

[2]National Wildlife Research Centre, Environment and Climate Change Canada, Ottawa, ON,

Canada. sarah.endicott@ec.gc.ca; Stephanie.Avery-Gomm@ec.gc.ca; Cheryl-

Ann.Johnson@ec.gc.ca; Mathieu.Leblond@ec.gc.ca; josie.hughes@ec.gc.ca

[3]University of British Columbia, Vancouver, BC, Canada. julwturner@gmail.com

[4]Northern Forestry Centre, Canadian Forest Service, Natural Resources Canada, Edmonton,

AB, Canada. Eric.W.Neilson@NRCan-RNCan.gc.ca

[5]FERIT Environmental Consulting, Thunder Bay, ON, Canada. northernbio@gmail.com

[6]Great Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, Sault Ste.

Marie, ON, Canada. philip.wiebe@NRCan-RNCan.gc.ca

*shared senior authorship as Co-PIs

ABSTRACT

Decision making in conservation science often relies on the best available information. This may include using models that were not designed for purpose and are not accompanied by an assessment of limitations. To begin addressing these issues, we sought to reproduce, and evaluate the suitability of, the best available models for predicting impacts of proposed mining on boreal woodland caribou (*Rangifer tarandus caribou)* resource selection and demography in northern Ontario. We then evaluated their suitability for projecting the impacts of development in the Ring of Fire region. To aid in accessibility, we developed an R package for data preparation, analyses of resource selection, and demographic parameters. We found existing models were either ill suited, or lacking, for ongoing regional planning. The specificity of the regional resource selection model limited its usefulness for predicting impacts of development, and the high variability across caribou ranges limited the usefulness of a national aspatial demographic model for predicting range-specific impacts. Variability in model coefficients across caribou ranges suggests selection responses vary with habitat availability (i.e. a functional response) while demographic responses continue to decline with increasing disturbance. Models designed for forecasting that are continuously updated by range-specific demographic and habitat information, are required to better inform conservation decisions and ongoing policy and planning practices in the Ring of Fire region.

Keywords: *Rangifer tarandus caribou*; Open science; Decision-support; Resource selection function; Demographic model; environmental impact assessment; caribouMetrics

1 INTRODUCTION

Decisions about natural resource development should be informed by the best available information (Fuller et al. 2020). Models that quantitatively link wildlife species responses to development, such as wildlife-habitat relationships and demography (Beyer et al., 2010; Matthiopoulos et al., 2020), can inform these decisions and associated policies (Wilson et al., 2021). However, these tools are generally not designed for decision support, and decision makers may be unaware of their limitations. For example, models are rarely reproducible; data and modelling processes are rarely reported nor accessible without author engagement, resulting in limited applicability outside of the specific context for which models were built. These problems can be addressed by couching models within an open source, and easily reproduced, decision-support tool (e.g., Eacker et al., 2019; Nagy-Reis et al., 2020; Nowak et al., 2018; McIntire et al. 2022) that can inform impact assessment, particularly for Species at Risk (Roche et al., 2020).

As a country supporting some of the largest intact landscapes globally, Canada has a heightened responsibility to conduct natural resource development decisions in the most sustainable way possible. For example, the Hudson Bay Lowlands of Northern Ontario is one of the largest intact wetlands and peatland carbon stores on the planet (Ibisch et al., 2016; Poley et al., 2022; Sothe et al., 2022; Tootchi et al., 2019). The region is home to many wildlife species and is largely inaccessible by road. At the same time, it is rich in valuable minerals and is a target for resource extraction and economic development aimed at a green energy transition (Carlson and Chetkiewicz, 2013; Far North Science Advisory Panel (Ont.) et al., 2011; IAAC, 2021). With an anticipated Regional Assessment in the region (IAAC, 2023), and much interest in critical minerals for batteries and other purposes (NRCan 2022), there is a need for modelling tools to assess the potential environmental impacts of the proposed mining of 'Ring of Fire' (RoF) mineral deposits and associated development on wildlife in this globally significant ecosystem (Far North Science Advisory Panel (Ont.) et al., 2011; IAAC, 2021; Fig. 1).

Quantitative knowledge of wildlife-habitat relationships are commonly represented by

resource selection functions (RSFs), which can produce maps of habitat suitability through

space and time, and can infer consequences of habitat loss (Boyce et al., 2002; Harju et al.,

2011; Johnson et al., 2004; Matthiopoulos et al., 2019). However, RSFs represent descriptive

associations between the behaviour of a sample of animals and habitat under current, scale-

specific conditions, limiting their predictive capacity and transferability to novel conditions

(Boyce, 2006; Decesare et al., 2012b; Kunegel-Lion et al., 2022; Yates et al., 2018). Aspatial

wildlife demographic models are another tool and provide insight into drivers of population

growth, such as survival and recruitment, and are also used for characterizing demographic

responses to landscape alteration (Johnson et al., 2020; Sorensen et al., 2008; Stewart et al.,

2020). However, high uncertainty about demographic parameters, drivers of change, and vital

rates can limit the utility of models for decision making and regional application (Chaudhary and

Oli, 2020; Sleep and Loehle, 2010; but see Stewart et al. 2020).

Boreal populations of woodland caribou (*Rangifer tarandus caribou*; hereafter 'boreal

caribou') inhabit large portions of Canada's boreal forest (ECCC, 2011, 2019; Festa-Bianchet et

al., 2011), where anthropogenic disturbance from natural resource extraction sector threatens

population persistence (Fryxell et al., 2020; Hebblewhite, 2017; Stewart et al. 2020; Johnson et

al., 2020). Nationally, only 29% of population ranges (15 of 51) are considered self-sustaining

and boreal caribou are listed as threatened under the federal Species at Risk Act (ECCC,

2024a). In the province of Ontario, boreal caribou are listed as threatened due to development

from forestry and mineral exploration, including their road networks, which together increase

predation and reduces adult female and calf survival (MNRF, 2014a) Ongoing landscape

disturbances are predicted to cause continued population declines (Fryxell et al., 2020).

Effective conservation of caribou in Ontario requires spatial and demographic tools that can

project the effects of future landscape and climatic changes on caribou habitats and population

performance. An existing Ontario RSF (Hornseth and Rempel, 2016) and national demographic

model (Johnson et al. 2020) represent the best available regional prediction tools for this

species but have many potential shortcomings (ECCC 2024b): notably, they rely on a limited

amount of data from over a decade ago, and their utility for transferability to other contexts and

projecting anticipated anthropogenic impacts has not been evaluated.

To advance the practice of conservation science decision making for a species at risk in

a changing landscape, we sought to evaluate the suitability of the best available models for

predicting impacts of proposed mining in the RoF on Ontario's northern boreal caribou. To

assess the potential impacts, we reproduced existing caribou RSF and demographic models

and evaluated their suitability for spatial transferability and prediction. Species at risk models

can be difficult to fully reproduce as information is rarely made publicly available. In this context,

location data for the RSF was not available and error estimates for model coefficients were not

published (Hornseth and Rempel 2016). For the national demographic model, code and

regression model parameters were made available upon request from the authors (Johnson et

al., 2020), but data owned by provinces was not accessible. We demonstrate and discuss the

challenges of this exercise and suggest open source tools and reproducible workflows as a

solution by reproducing the existing models within an R package framework (Wickham, 2015).

To aid this effort, we created an R package (caribouMetrics) that can be integrated into

predictive frameworks to support resource use decisions or modified and advanced by other

practitioners for other uses (Bodner et al., 2021a; McIntire et al., 2022; Micheletti et al., 2021;

Miller and Frid, 2022).

2 METHODS

2.1 Study Area

We focused on four caribou ranges, as delineated by Ontario's provincial government,

that are adjacent to or overlap with the RoF mining claims and had published RSF coefficients:

Missisa, James Bay, Pagwachuan, and Nipigon. The Missisa range contains most of the RoF

mineral claims (Far North Science Advisory Panel (Ont.) et al., 2011; IAAC, 2021; Fig. 1) and is

the focus of our study. Missisa and James Bay ranges occur in Ontario's Far North (Far North

Science Advisory Panel (Ont.) et al., 2011), where anthropogenic disturbance has so far been

limited to small communities and exploration camps, winter roads, and the now inactive Victor

Diamond Mine. Nipigon and Pagwachuan ranges are almost entirely within the Managed Forest

Zone (MFZ) where industrial forestry is common (Fig. 1).

These four ranges straddle the Ontario Shield and Hudson Bay Lowlands ecozones (Fig

1). The Shield consists of mixed and coniferous forests dominated by black spruce (*Picea*

mariana), and containing jack pine (*Pinus banksiana*), balsam fir (*Abies balsamea*), white

spruce (*Picea glauca*), trembling aspen (*Populus tremuloides*), and balsam poplar (*Populus*

balsamifera; Crins et al., 2009). The Missisa and Pagwachuan ranges include the transition

between the Shield and Lowlands, which is mainly composed of peatland complexes with poor

drainage, and trees including stunted black spruce and tamarack (*Larix laricina*; Crins et al.,

2009). The James Bay range is almost entirely within the Lowlands and the Nipigon range is

entirely within the Shield (Fig 1).

2.2 RSF Model Reproduction & Validation

To reproduce existing RSFs for caribou in northern Ontario we used tables of published

boreal caribou RSF coefficients for the same areas as Hornseth and Rempel (2016) (hereafter

referred to as the 'original RSFs'). Hornseth and Rempel referred to these models as resource

selection probability functions (RSPFs), but their approach differs from what is commonly

understood as an RSPF (Johnson et al., 2006); as such we use the term 'RSF' throughout. The

data used to produce the original RSFs were collected between 2009 and 2013 from GPS-

collared caribou across a variety of companion studies within the ranges of interest; however,

these data are not publicly available (Hornseth and Rempel, 2016; MNRF, 2014b, 2014b;

Rempel and Hornseth, 2018). Instead, we obtained the published coefficients for range-specific

seasonal RSFs of each reported top model (Table S1.1) for the Nipigon, Pagwachuan, Missisa,

and James Bay ranges. However, associated error or uncertainty information for individual

coefficients was not reported in Hornseth and Rempel (2016) and model performance was only

reported at the model level (e.g., sensitivity, specificity, and area under the receiver operating [

ROC] characteristic curve).

The predictors used in the original RSFs were developed from land cover, fire and

harvest disturbance history, linear features, and esker datasets (Tables S1.2, S1.3). To

reproduce the original RSFs, we acquired data sets for the predictor variables that were publicly

available and matched the data sources described by Hornseth and Rempel (2016) as closely

as possible. Harvest, fire, and linear features data from the present (2020) were available and

we used information on the timing of the event or the feature's construction to recreate the data

that was available when the original models were produced (Table S1.2). We used current

versions of these same datasets to project the RSFs under different development scenarios

(Table S1.2).

To assess our ability to reproduce the original models, we acquired the original projected

surfaces for comparison from the authors (R. Rempel pers. comm, 2021). We validated our

models visually and quantitatively. The original models used a nested hexagon grid, common in

Ontario (e.g. Poley et al. 2014; Ray et al. 2018), to generate predictions. For ease of future

implementation using raster tools, we approximated this approach with distance-weighted

moving windows on a rectangular grid, and compared the two approaches by extracting the

values from our rectangular grid to the 592-hectare hexagonal grid provided by the authors

using the weighted mean. We mapped the difference in predictions and explanatory variables

between the models and used scatterplots to compare the values of the model responses for

each grid cell produced by both methods. A perfect reproduction would produce a Pearson

correlation coefficient of 1, and any deviation from the original prediction would reduce this

value. We expected some differences in the predictions to result from the type of grid and

predictor data sets used.

2.3 Disturbance Scenarios and RSF Model Projection

To assess the suitability of existing RSFs for projecting the impacts of disturbance on boreal

caribou habitat selection we considered three simple disturbance scenarios. We created a

'base' scenario with no change in disturbance footprint (0.41% anthropogenic disturbance), a

'roads-only' scenario that included proposed RoF access roads (1.11 % anthropogenic

disturbance; MNDMNRF, 2022), and a 'roads-and-mines' scenario that included the proposed

access roads and mining claims associated with the RoF (6.95% anthropogenic disturbance;

Fig. 2; Table S1.2). The original RSF included road density as a predictor and required

additional information on roads within mining areas that we were not able to obtain; we therefore

only compared the base and road-only scenarios for the RSF but included all three scenarios for

the demographic model (see below).

To understand how the 'roads-only' scenario might affect resource selection, we

projected the original RSFs across updated landscape conditions as represented by i) temporal

changes in forest structure between 2010 and 2020 (e.g., including simulated fires; Table S1.2)

and ii) the new proposed roads-only scenario (Fig. 2). We also assessed the potential for

borrowing RSF information from other ranges by transferring coefficients from the top original

RSFs from adjacent ranges (James Bay, Nipigon, and Pagwachuan) to Missisa. We visualized

the spatial transferability of these models, and then examined their sensitivity to changing

habitat availability using scatterplots where we compared the value of the model response for

each grid cell produced by the Missisa model and each respective adjacent range.

2.4 Demographic Model Reproduction, Validation, and Projection

Canada's national demographic boreal caribou model was developed from adult female

survival (S), calf recruitment (R), and landscape data across 58 boreal caribou study areas,

including 13 study areas in Ontario (Johnson et al., 2020; Table S1). We used models (Johnson

et al., 2020) to predict changes in S and R within the roads-only and roads-and-mines

disturbance scenarios. The national demographic model is aspatial and all types of

anthropogenic disturbances are combined into a single measure of disturbed area within a

range; using the 'roads-and-mines' scenario is sufficient and there is no need to specify the

location of roads within mining claims. We calculated the relevant predictor variables for the

Missisa range based on Johnson et al. (2020;i.e., % anthropogenic disturbance buffered by 500

m; % wildfire within the last 40 years; Table 1), and calculated expected R and S as a function

of disturbance according to the beta regression models with highest support (M4 and M1

respectively from Johnson et al., 2020):

$$R \sim Beta(\mu^R, \phi^R); log(\mu^R) = \beta_0^R + \beta_a^R\ anthro\ + \beta_f^R\ fireExclAnthro, \qquad \text{(eq 1a)}$$

$$S \sim (46 \times Beta(\mu^S, \phi^S) - 0.5)/45; log(\mu^S) = \beta_0^S + \beta_a^S\ anthro, \qquad \text{(eq 1b)}$$

Where $\Box^R$ ~ *Normal(19.862,2.229)* and $\Box^S$ ~ *Normal(63.733,8.311)* are precisions of the Beta

distributed errors (Ferrari and Cribari-Neto, 2004), and survival rates are back transformed as in

Johnson et al. 2020. Table 3 of Johnson et al. (2020) provides the expected values and 95%

confidence intervals of all regression coefficients ($\beta_0{}^R$, $\beta_a{}^R$, $\beta_f{}^R$, $\beta_0{}^S$, $\beta_a{}^S$), which are assumed to

be Gaussian distributed. To evaluate these recruitment (eq 1a) and survival (eq 1b) models, we

sampled expected demographic rates across a range of anthropogenic and fire disturbance (0-

100%) to reproduce expected values and 95% predictive intervals from Fig. 3 and Fig. 5 of

Johnson et al. (2020).

In areas of low anthropogenic disturbance, such as the Missisa and James Bay ranges,

there is substantial among-population variability in recruitment (Fig. 3 in Johnson et al., 2020).

To model this variation, we selected regression model parameter values for each study area

(i.e. a sample population) at the beginning of simulations and assigned to quantiles of the error

distributions for survival and recruitment. Sample populations remained in their quantiles as the

landscape changed, allowing us to distinguish the effects of changing disturbance from variation

in initial population status. To show these effects, we projected population growth for 35 sample

populations across a wide range of anthropogenic disturbance levels (0-90%). We slightly

altered Johnson et al's (2020) demographic model by including demographic stochasticity

(Supplementary Materials Part 2; see Hughes et al. *in review* for details).

We characterized the distribution of outcomes for our three disturbance scenarios (Fig.

2, Table 1) more thoroughly by projecting the dynamics of 500 sample populations. Population

growth rate each year is given by $\lambda_t = N_{t+1}/N_t$ when $N_t > 0$ and $\lambda_t = 0$ when $N_t = 0$. For each sample

population, population growth was projected for 20 years from an assumed initial population

size of 373 females, which we derived from the minimum animal count of 745 caribou between

2009 and 2011, assuming 50% were female (MNRF, 2014b). This estimate is conservative

because adult male:female ratios are typically lower in caribou (ECCC, 2011; Espinoza and

Weckerly, 2021). We report the average growth rate λ over that time. To verify our reproduction

of the demographic model used by Johnson et al. (2020), we compared our outputs to those

from model code supplied by the authors.

2.5 caribouMetrics: an R Package for Model Reproduction, Validation, and Projection

We incorporated RSF and demographic model components into the *caribouMetrics* R

package with documentation and vignettes explaining their use and used GitHub to promote

version control and transparency of the development process. This package can be accessed

on GitHub: https://github.com/LandSciTech/caribouMetrics.

The standardized nature of packages allows them to easily integrate into the larger R

ecosystem, facilitating ease of extension and adaptation for tasks beyond their original design

(e.g. prediction or forecasting; McIntire et al., 2022). *caribouMetrics* also contains functions that

automate the geospatial data preparation process to facilitate application to new landscapes.

The original RSFs were developed using a closed scripting language called Landscape

Scripting Language (LSL; Kushneriuk and Rempel, 2011), which we re-implemented in R. For

the demographic model, we began by borrowing demographic rate sampling code from a

SpaDES (Chubaty and McIntire, 2022) module (*caribouPopGrowth*; https://github.com/tati-

micheletti/caribouPopGrowthModel; Stewart et al., 2023), and advanced the method to include

precision and quantiles. Together, this makes the national caribou demographic model

(Johnson et al. 2020) more transparent, reproducible, and adaptable to local population

estimates.

3 RESULTS

3.1 RSF Model Reproduction, Validation, and Projection

We were able to produce a reasonably accurate representation of the original RSFs for

the Missisa range, as evidenced by high Pearson's r values across all seasons (r > 0.935; Fig.

S1.1). However, some locations showed different predictions depending on the season (Fig.

S1.2). This was expected given the input data was not exactly the same between the two

methods (Fig. S1.3). In the Missisa range, the model predicted the highest relative use

probabilities in the northwest of the study area during winter (Fig. 3), consistent with the original

RSFs. During the summer and spring, the eastern portion of the range was used more than the

northwest (Fig. 3).

We observed a lower relative probability of use in areas associated with proposed roads

(Fig. 4). Lower relative probability of use along the road corridors was strongest in the spring

and summer, consistent with seasonal changes in the response to roads described in the

original RSFs (Fig. 4). There was high variability in the estimated response to roads among

ranges (Fig. 4). The James Bay range prediction appeared the most similar to the Missisa range

prediction (Fig. 4); however it varied by season (Fig S1.4). The Nipigon and Pagwachuan range

projections, which visually differed substantially from the Missisa projection, did not show a

strong response to proposed roads (Fig. 4; Fig S1.4).

3.2 Demographic Model Reproduction, Validation, and Projection

Our regression models for survival and recruitment closely matched those of Johnson et

al. 2020 (Fig. 5). Anthropogenic disturbance remained low in all our disturbance scenarios

(Table 1), and the corresponding range of variability in demographic rates among sample

populations was high (Fig. 5). The model predicted that increasing anthropogenic disturbance

would decrease both survival and recruitment (Fig. 5), but the importance of that decrease for

self-sustainability of the population was highly uncertain and depended on initial population

status (Fig. 5).

4 DISCUSSION

To evaluate existing models and assess their limitations for projecting anticipated landscape

impacts we examined the applicability of two kinds of existing models; resource selection

functions (RSF) and demographic models, for boreal caribou. We found that the original Missisa

range RSF was poorly suited for projecting the impacts of disturbance because it was specific to

conditions at the time of model development in an area that contained limited anthropogenic

disturbance. This led to high uncertainty in the estimated impacts of linear features, with a low

standardized effect size, high variability among seasons, and the possibility that the model

understinated future impacts of linear features. In contrast, the existing national demographic

model was too general to project the impacts of disturbance in this range. Additional data

collection is necessary but not sufficient to address these limitations (see also ECCC 2024b);

there is also a need for new models that are better suited to forecasting the impacts of

disturbance. These models should be designed for purpose. They should identify and account

for uncertainty, be updateable with new information (e.g., local data), be transparent, simple,

and reproducible (Bodner et al., 2021b).

4.1 Lessons learnt from using non-predictive models to forecast impacts

Across caribou ranges where anthropogenic disturbance is low, the national

demographic model predicted high variation in recruitment and survival among populations (Fig.

3 and Fig. 5 of Johnson et al., 2020), leading to high variability in projected impact of increasing

disturbance on population growth rate in the RoF (Fig. 5 here). This variability highlights the

need for local information about the current (i.e., baseline) demographic parameters of caribou

populations in the RoF and ongoing monitoring of disturbance impacts in the area. An

assessment of the status of the Missisa population based on 2008 – 2012 data (MNRF, 2014b)

suggested survival, reproduction, and population growth rate in this range were lower than

predicted by the national demographic model of Johnson et al. 2020 (see black diamond in Fig.

5). The national demographic model (Johnson et al., 2020) predicts that increasing

anthropogenic disturbance will decrease both survival and recruitment, but the importance of

that decrease for population viability depends on baseline demographic parameters. Even small

changes in adult female survival can affect the sustainability of a population in ranges with low

adult survival (Johnson et al., 2020). We a method for integrating local survival, recruitment, and

status information into range-specific demographic projections informed by national

demographic-disturbance relationships, but see Hughes et al. *in review* for recent

advancements. Without adequate recent information on population size and demographic

parameters, it is difficult to predict the impacts of proposed RoF disturbance on the viability of

this particular population with any degree of confidence.

4.2 Modelling challenges and opportunities

The original RSF models (Hornseth and Rempel, 2016) were fit independently with data

from each range. This is a reasonable approach when the objective is to characterize current

habitat use, but not for projecting responses to changing landscape conditions. In the original

RSFs there was high variability in the effect of linear features among range-specific RSFs,

suggesting that the behavioural response of caribou to linear features may vary with the amount

or type of disturbance in a range (i.e., a functional response; Mysterud and Ims, 1998), or that

effects of linear features were confounded with other correlated predictors (e.g., forest harvest;

Hornseth and Rempel, 2016). Hierarchical regional models that include functional responses to

disturbance (Matthiopoulos et al., 2011; Muff et al., 2020; Olson et al., 2021; Teitelbaum et al.,

2021) could yield models that are better suited for projection.

Our method of modelling variation in survival and recruitment among populations (Fig. 5)

adequately reproduced the observed variation among populations (Fig. 3 and Fig. 5 in Johnson

et al., 2020), though it differs from the range-specific scenario approach taken by Johnson et al.

(2020). We assumed demographic parameters (i.e., recruitment and survival) vary with

disturbance according to the best supported national models, but we note that other competing

models with comparable support (i.e., Table 2 of Johnson et al., 2020) are also plausible and

might yield somewhat different projections when applied to northern ranges with low

anthropogenic disturbance (*sensu* Stewart et al. 2023). This simple population model assumes

no variation in recruitment or survival with age, sex, or other parameters (Supplementary

Material Part 2). The simple demographic model also ignores uncertainty associated with

imperfect survey methods (DeCesare et al., 2012a; Ellington et al., 2020). A more thorough

investigation of the sensitivity of caribou demographic projections to variation in model form and

measurement errors is warranted but beyond the scope of this paper.

4.3 Next steps for supporting actionable science in the Ring of Fire region

In the Ring of Fire region, existing data and models are not sufficient for forecasting

changes in caribou habitat selection and demography. Recent caribou movement and

demographic data is lacking, and there are also important deficits in our understanding of

disturbance history, impacts, and recovery in the low productivity peatlands of the Hudson Bay

Lowlands. The region is incredibly remote, making data collection expensive logistically difficult.

There are various efforts underway to collect data needed to inform resource development and

wildlife management decisions in the RoF (e.g., Government of Canada, 2022), and we hope

these efforts also lead to opportunities for collaborative development of region-specific open-

source forecasting models. We hope our analysis of existing models helps clarify that new data

and models are needed to inform decisions in this region; existing models are not enough to

provide spatially explicit information to minimize detrimental effects of human development on

caribou recovery in northern Ontario

Although we believe that more data is needed in the Ring of Fire region, it does not

follow that more data is needed in all regions and circumstances. We advocate for an open,

transparent, iterative modelling approach that allows for integration of data and models as they

become available, *sensu* caribou metrics and Hughes et al. (*in review*). This approach allows

ongoing assessment of the implications of additional data for model uncertainty and can

address important questions about what additional data is required to inform conservation

decisions. A well-designed, open, transparent, iterative modelling framework can help support

and inform important, nuanced, and region-specific decision making about exactly what data is

needed and why.

5 CONCLUSIONS

We reiterate a call for improvements in future models, and their workflows, used for

conservation decision-making. To assess disturbance impacts on wildlife, models need to be

designed for purpose, identify and account for uncertainty, be updateable as new predictor

information is available (e.g., local data), and be reproducible (Bodner et al., 2021b). We

encourage developers of wildlife response models and collectors of relevant wildlife data to

work together toward this goal (Davidson et al., 2021; Russell et al., 2021).

Developing usable decision-support tools demands time and specialized skills that may

not be accessible to everyone engaged in open conservation science. One solution is to work in

multi-disciplinary teams (Bodner et al., 2021b). Another is to recognize that steps and training

toward transparency and reproducibility are valuable, even if the result is not always an easily

usable tool. In this project, we were able to reproduce existing models only because the

developers of those models were willing to share their code and discuss their model generation

procedures. Code does not have to be flawless to enable others to build on previous work, and

a shift to more open workflows will reduce the chance of errors, increase efficiency, and

advance our ability to generalize across studies and systems (Alston and Rick, 2021; Lewis et

al., 2018). We hope that improving the transparency, reproducibility, and decision-relevance of

wildlife response models will strengthen our collective ability to make informed decisions and

advance conservation science and practice.

AUTHOR CONTRIBUTIONS MD led the writing and co-led the analysis; SE and CS co-led

package development and provided support for analysis and writing; JT provided contributions

to writing and evaluation of the RSF; SAG, CAJ, ML, EN, RR, PW (listed in alphabetical order)

provided advice on package development, analysis, and writing; RR and CAJ also provided

code and insight into the RSFs and demographic models. FS, JB, and JH provided supervision

and oversaw the writing and analysis of this piece.

DATA AVAILABILITY STATEMENT The caribouMetrics R package is available at

https://github.com/LandSciTech/caribouMetrics, and analysis code is available at

https://github.com/LandSciTech/MissisaBooPaper. Data required for analysis is available in this

OSF repository: https://osf.io/r9mkp/?view_only=fb71321265d14dbeb3d932e4de66be0c

Contributed paper

TARGET AUDIENCE We target conservation scientists and decision makers addressing

Environmental Impact and wildlife policy.

Supplementary Material

Supplementary material is available in this OSF repository:

https://osf.io/r9mkp/?view_only=fb71321265d14dbeb3d932e4de66be0c

WORD COUNT 4377

CORRESPONDENCE

Frances Stewart, fstewart@wlu.ca

Wilfrid Laurier University, Biology Department

75 University Ave., Waterloo, Ontario, N2L 3C5

COMPETING INTERESTS STATEMENT: RR is principal of FERIT Consulting. FS and JB were

project leads at Wilfrid Laurier University and JH was project lead at ECCC.

FUNDING STATEMENT: Funding for this work was provided by the Canadian Wildlife Service

and the Wildlife and Landscape Science Directorate of Environment and Climate Change

Canada. F. Stewart and J. Baltzer were supported by the NSERC Canada Research Chairs

Program.

ACKNOWLEDGEMENTS

We thank J. Girard, S. McFarlane, R. Bhatti, S. Hayne, M. Hornseth, and G.D. Sutherland for

discussion and suggestions that helped improve this manuscript. We also thank E. McIntire, T.

Micheletti, A. Chubaty, and the Predictive Ecology group for their support and contributions to R

package development.

References

Alston, J.M., Rick, J.A., 2021. A beginner's guide to conducting reproducible research. Bull Ecol Soc Am. https://doi.org/10.1002/bes2.1801

Avgar, T., Potts, J.R., Lewis, M.A., Boyce, M.S., 2016. Integrated step selection analysis: Bridging the gap between resource selection and animal movement. Methods in Ecology and Evolution 7, 619–630. https://doi.org/10.1111/2041-210X.12528

Beyer, H.L., Haydon, D.T., Morales, J.M., Frair, J.L., Hebblewhite, M., Mitchell, M., Matthiopoulos, J., 2010. The interpretation of habitat preference metrics under use-availability designs. Philisophical Transactions of The Royal Society of Biological Sciences 365, 2245–2254.

Bodner, K., Rauen Firkowski, C., Bennett, J.R., Brookson, C., Dietze, M., Green, S., Hughes, J., Kerr, J., Kunegel-Lion, M., Leroux, S.J., McIntire, E., Molnár, P.K., Simpkins, C., Tekwa, E., Watts, A., Fortin, M.-J., 2021a. Bridging the divide between ecological forecasts and environmental decision making. Ecosphere 12, e03869. https://doi.org/10.1002/ecs2.3869

Bodner, K., Brimacombe, C., Chenery, E.S., Greiner, A., McLeod, A.M., Penk, S.R., Vargas Soto, J.S., 2021b. Ten simple rules for tackling your first mathematical models: A guide for graduate students by graduate students. PLoS Comput Biol 17, e1008539. https://doi.org/10.1371/journal.pcbi.1008539

Boyce, M.S., 2006. Scale for resource selection functions. Diversity and Distributions 12, 269–276. https://doi.org/10.1111/j.1366-9516.2006.00243.x

Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions. Ecological Modelling 157, 281–300.

Carlson, M., Chetkiewicz, C., 2013. A fork in the road: future development in Ontario's Far North. Canadian Boreal Initiative, Ottawa, ON.

Chaudhary, V., Oli, M.K., 2020. A critical appraisal of population viability analysis. Conservation Biology 34, 26–40. https://doi.org/10.1111/cobi.13414

Chubaty, A.M., McIntire, E.J.B., 2022. SpaDES: Develop and Run Spatially Explicit Discrete Event Simulation Models.

Crins, W.J., Gray, P.A., Uhlig, P.W.C., Wester, M.C., 2009. Ecosystems of Ontario. Part 1, Part 1, (No. SIB TER IMA TR-01). Ontario, Ministry of Natural Resources, Inventory, Monitoring and Assessment Section, Toronto, Ont.

Davidson, S., Bohrer, G., Kölzsch, A., Vinciguerra, C., Kays, R., 2021. Mobilizing Animal Movement Data: API use and the Movebank platform. BISS 5, e74312. https://doi.org/10.3897/biss.5.74312

DeCesare, N.J., Hebblewhite, M., Bradley, M., Smith, K.G., Hervieux, D. and Neufeld, L., 2012a. Estimating ungulate recruitment and growth rates using age ratios. The Journal of Wildlife Management, 76(1), pp.144-153.

Decesare, N.J., Hebblewhite, M., Schmiegelow, F., Hervieux, D., McDermid, G.J., Neufeld, L., Bradley, M., Whittington, J., Smith, K.G., Morgantini, L.E., Wheatley, M., Musiani, M., 2012b. Transcending scale dependence in identifying habitat with resource selection functions. Ecological Applications 22, 1068–1083.

Eacker, D.R., Hebblewhite, M., Steenweg, R., Russell, M., Flasko, A., Hervieux, D., 2019. Web-based application for threatened woodland caribou population modeling. Wildl. Soc. Bull. 43, 167–177. https://doi.org/10.1002/wsb.950

ECCC, 2011. Scientific assessment to inform the identification of critical habitat for woodland caribou (Rangifer tarandus caribou), boreal population, in Canada. Canadian Wildlife Service, Ottawa.

ECCC, 2019. Amended Recovery Strategy for the Woodland Caribou (Rangifer tarandus
caribou), Boreal population, in Canada [Proposed], Species at Risk Act Recovery
Statagy Series. Environment and Climate Change Canada, Ottawa.
ECCC, 2024a. Report on the Progress of the Recovery Strategy Implementation (Period 2017 –
2022) and the Action Plan Implementation (Period 2018 - 2023) for Caribou (Rangifer
tarandus), Boreal Population, in Canada. Species at Risk Act Recovery Strategy
Report Series. Environment and Climate Change Canada, Ottawa.
ECCC, 2024b. Scientific Assessment of Federal and Provincial Frameworks for the
Conservation of Boreal Caribou in Ontario. Scientific Assessment Report. Environment
and Climate Change Canada, Ottawa. https://wildlife-species.az.ec.gc.ca/species-risk
registry/virtual_sara/files//Sa-BorealCaribouBoreale-v00-2024Ma-eng.pdf
Ellington, E.H., Lewis, K.P., Koen, E.L., Vander Wal, E., 2020. Divergent estimates of herd-wide
caribou calf survival: Ecological factors and methodological biases. Ecology and
Evolution 10, 8476–8505. https://doi.org/10.1002/ece3.6553.
Espinoza, Z.S., Weckerly, F.W., 2021. A Comparative Analysis of Adult Sex Ratios in
Polygynous and Monogamous Mammal Populations. amid 186, 299–308.
https://doi.org/10.1674/0003-0031-186.2.299
Far North Science Advisory Panel (Ont.), Ontario, Ministry of Natural Resources, 2011. Science
for a changing far north: the report of the Far North Science Advisory Panel. Queen's
Printer for Ontario, Toronto, Ont.
Ferrari, S., Cribari-Neto, F., 2004. Beta Regression for Modelling Rates and Proportions.
Journal of Applied Statistics 31, 799–815.
https://doi.org/10.1080/0266476042000214501
Festa-Bianchet, M., Ray, J.C., Boutin, S., Côté, S.D., Gunn, A., 2011. Conservation of caribou (
Rangifer tarandus) in Canada: an uncertain future. Can. J. Zool. 89, 419–434.
https://doi.org/10.1139/z11-025
Fryxell, J.M., Avgar, T., Liu, B., Baker, J.A., Rodgers, A.R., Shuter, J., Thompson, I.D., Reid,
D.E.B., Kittle, A.M., Mosser, A., Newmaster, S.G., Nudds, T.D., Street, G.M., Brown,
G.S., Patterson, B., 2020. Anthropogenic disturbance and population viability of
woodland caribou in Ontario. Jour. Wild. Mgmt. 84, 636–650.
https://doi.org/10.1002/jwmg.21829
Government of Canada, 2022. Agreement for the Conservation of Caribou, Boreal population in
Ontario. Ottawa, Ontario. April 22, 2022. https://sararegistry.gc.ca/virtual_sara/files/Ca-
CaribouOntario-v00-2022April-Eng.pdf
Harju, S.M., Dzialak, M.R., Osborn, R.G., Hayden-Wing, L.D., Winstead, J.B., 2011.
Conservation planning using resource selection models: altered selection in the
presence of human activity changes spatial prediction of resource use: Human activity
alters spatial use predictions. Anim Conserv 14, 502–511. https://doi.org/10.1111/j.1469-
1795.2011.00456.x
Hebblewhite, M., 2017. Billion dollar boreal woodland caribou and the biodiversity impacts of the
global oil and gas industry. Biological Conservation 206, 102–111.
https://doi.org/10.1016/j.biocon.2016.12.014
Hornseth, M.L., Rempel, R.S., 2016. Seasonal resource selection of woodland caribou
(*Rangifer tarandus caribou*) across a gradient of anthropogenic disturbance. Can. J.
Zool. 94, 79–93. https://doi.org/10.1139/cjz-2015-0101
Hughes, J., Endicott, S., Calvert, A.M., and Johnson, C.A. 2024. Integration of national
demographic-disturbance relationships into local caribou population viability projections
can reduce uncertainty and inform monitoring decisions. In review at Ecological
Informatics. ECOINF-D-24-01983.
IAAC, 2021. Draft Agreement to Conduct a Regional Assessment in the Ring of Fire Area. (last
accessed June 2024) https://iaac-aeic.gc.ca/050/documents/p80468/142280E.pdf

IAAC, 2023. Regional Assessment in the Ring of Fire Area. Latest Update (last accessed June
2024). https://iaac-aeic.gc.ca/050/evaluations/proj/80468
Ibisch, P.L., Hoffmann, M.T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L., DellaSala,
D.A., Vale, M.M., Hobson, P.R., Selva, N., 2016. A global map of roadless areas and
their conservation status. Science 354, 1423–1427.
https://doi.org/10.1126/science.aaf7166
Johnson, C.A., Sutherland, G.D., Neave, E., Leblond, M., Kirby, P., Superbie, C., McLoughlin,
P.D., 2020. Science to inform policy: Linking population dynamics to habitat for a
threatened species in Canada. J Appl Ecol 57, 1314–1327. https://doi.org/10.1111/1365-
2664.13637
Johnson, C.J., Nielsen, S.E., Merrill, E.H., Trent, L., Boyce, M.S., Science, E., Program, M.,
British, N., George, P., 2006. Resource Selection Functions Based on Use – Availability
Data: Theoretical Motivation and Evaluation Methods. Journal of Wildlife Management
70, 347–357. https://doi.org/10.2193/0022-541X(2006)70
Johnson, C.J., Seip, D.R., Boyce, M.S., 2004. A quantitative approach to conservation planning:
using resource selection functions to map the distribution of mountain caribou at multiple
spatial scales. J Appl Ecology 41, 238–251. https://doi.org/10.1111/j.0021-
8901.2004.00899.x
Kunegel-Lion, M., Neilson, E.W., Mansuy, N., Goodsman, D.W., 2022. Habitat quality does not
predict animal population abundance on frequently disturbed landscapes. Ecological
Modelling 469, 109943. https://doi.org/10.1016/j.ecolmodel.2022.109943
Kushneriuk, R.S., Rempel, R.S., 2011. LSL - Landscape Scripting Language. Ontario Ministry of
Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, ON.
Lewis, K.P., Vander Wal, E., Fifield, D.A., 2018. Wildlife biology, big data, and reproducible
research. Wildlife Society Bulletin. https://doi.org/10.1002/wsb.847
Matthiopoulos, J., Fieberg, J., Aarts, G., 2020. Species-Habitat Associations: Spatial data,
predictive models, and ecological insights, 1st ed. University of Minnesota Libraries.
https://doi.org/10.24926/2020.081320
Matthiopoulos, J., Field, C., MacLeod, R., 2019. Predicting population change from models
based on habitat availability and utilization. Proceedings of the Royal Society B:
Biological Sciences 286. https://doi.org/10.1098/rspb.2018.2911
Matthiopoulos, J., Hebblewhite, M., Aarts, G., 2011. Generalized functional responses for
species distributions. Ecology 92, 583–589.
McIntire, E.J.B., Chubaty, A.M., Cumming, S.G., Andison, D., Barros, C., Boisvenue, C., Haché,
S., Luo, Y., Micheletti, T., Stewart, F.E.C., 2022. PERFICT A Re-imagined foundation
for predictive ecology.pdf. Ecology Letters Online, 1–7.
Micheletti, T., Stewart, F.E.C., Cumming, S.G., Haché, S., Stralberg, D., Tremblay, J.A., Barros,
C., Eddy, I.M.S., Chubaty, A.M., Leblond, M., Pankratz, R.F., Mahon, C.L., Van
Wilgenburg, S.L., Bayne, E.M., Schmiegelow, F., McIntire, E.J.B., 2021. Assessing
Pathways of Climate Change Effects in SpaDES: An Application to Boreal Landbirds of
Northwest Territories Canada. Front. Ecol. Evol. 9, 679673.
https://doi.org/10.3389/fevo.2021.679673
Miller, B.W., Frid, L., 2022. A new approach for representing agent-environment feedbacks:
coupled agent-based and state-and-transition simulation models. Landsc Ecol 37, 43–
58. https://doi.org/10.1007/s10980-021-01282-y
MNDMNRF, 2022. Mining Lands Administration System. Ministry of Northern Development,
Mines, Natural Resources, and Forestry, Peterborough, ON.
MNRF, 2014a. State of the Woodland Caribou Resource Report. Ministry of Natural Resources
and Forestry - Species at Risk Branch, Thunder Bay, ON.

MNRF, 2014b. Integrated Range Assessment for Woodland Caribou and their Habitat - The Far North of Ontario 2013 (No. Version 1.1). Ministry of Natural Resources and Forestry - Species at Risk Branch, Thunder Bay, ON.

Moeller, A.K., Nowak, J.J., Neufeld, L., Bradley, M., Manseau, M., Wilson, P., McFarlane, S., Lukacs, P.M., Hebblewhite, M., 2021. Integrating counts, telemetry, and non-invasive DNA data to improve demographic monitoring of an endangered species. Ecosphere 12. https://doi.org/10.1002/ecs2.3443

Muff, S., Signer, J., Fieberg, J., 2020. Accounting for individual-specific variation in habitat-selection studies: Efficient estimation of mixed-effects models using Bayesian or frequentist computation. J Anim Ecol 89, 80–92. https://doi.org/10.1111/1365-2656.13087

Mysterud, A., Ims, R.A., 1998. Functional responses in habitat use: availability influences relative use intrade-off situations. Ecology 79, 1435–1441.

Nagy-Reis, M., Dickie, M., Sólymos, P., Gilbert, S.L., DeMars, C.A., Serrouya, R., Boutin, S., 2020. 'WildLift': An Open-Source Tool to Guide Decisions for Wildlife Conservation. Frontiers in Ecology and Evolution 8.

Nowak, J.J., Lukacs, P.M., Hurley, M.A., Lindbloom, A.J., Robling, K.A., Gude, J.A., Robinson, H., 2018. Customized software to streamline routine analyses for wildlife management: Apps for Wildlife Management. Wildl. Soc. Bull. 42, 144–149. https://doi.org/10.1002/wsb.841

NRCan, 2022. The Canadian Critical Minerals Strategy: From Exploration to Recycling: Powering the Green and Digital Economy for Canada and the World. Natural Resources Canada. Ottawa.

Olson, L.E., Bjornlie, N., Hanvey, G., Holbrook, J.D., Ivan, J.S., Jackson, S., Kertson, B., King, T., Lucid, M., Murray, D., Naney, R., Rohrer, J., Scully, A., Thornton, D., Walker, Z., Squires, J.R., 2021. Improved prediction of Canada lynx distribution through regional model transferability and data efficiency. Ecol. Evol. 11, 1667–1690. https://doi.org/10.1002/ece3.7157

Poley, L.G., Pond, B.A., Schaefer, J.A., Brown, G.S., Ray, J.C. and Johnson, D.S., 2014. Occupancy patterns of large mammals in the Far North of Ontario under imperfect detection and spatial autocorrelation. *Journal of Biogeography*, *41*(1), pp.122-132.

Poley, L.G., Schuster, R., Smith, W., Ray, J.C., 2022. Identifying differences in roadless areas in Canada based on global, national, and regional road datasets. Conservation Science and Practice n/a, e12656. https://doi.org/10.1111/csp2.12656

Prokopenko, C.M., Boyce, M.S., Avgar, T., 2017. Extent-dependent habitat selection in a migratory large herbivore: road avoidance across scales. Landscape Ecology 32, 313–325. https://doi.org/10.1007/s10980-016-0451-1

Ray, J.C., Poley, L.G., Magoun, A.J., Chetkiewicz, C.L.B., Meg Southee, F., Neil Dawson, F. and Chenier, C., 2018. Modelling broad☐scale wolverine occupancy in a remote boreal region using multi☐year aerial survey data. *Journal of biogeography*, *45*(7), pp.1478-1489.

Rempel, R.S., Carlson, M., Rodgers, A.R., Shuter, J.L., Farrell, C.E., Cairns, D., Stelfox, B., Hunt, L.M., Mackereth, R.W., Jackson, J.M., 2021. Modeling Cumulative Effects of Climate and Development on Moose, Wolf, and Caribou Populations. Jour. Wild. Mgmt. 85, 1355–1376. https://doi.org/10.1002/jwmg.22094

Rempel, R.S., Hornseth, M.L., 2018. Range-specific seasonal resource selection probability functions for 13 caribou ranges in Northern Ontario (No. IFR-01), Science and Research Internal File Report. Ministry of Natural Resources and Forestry, Peterborough, ON.

Roche, D.G., Granados, M., Austin, C.C., Wilson, S., Mitchell, G.M., Smith, P.A., Cooke, S.J., Bennett, J.R., 2020. Open government data and environmental science: a federal Canadian perspective. FACETS 5, 942–962. https://doi.org/10.1139/facets-2020-0008

611   Russell, D., Gunn, A., White, R., 2021. A decision support tool for assessing cumulative effects
612   on an Arctic migratory tundra caribou population. E&S 26, art4.
613   https://doi.org/10.5751/ES-12105-260104

614   Sleep, D.J.H., Loehle, C., 2010. Validation of a Demographic Model for Woodland Caribou. The
615   Journal of Wildlife Management 74, 1508–1512. https://doi.org/10.1111/j.1937-
616   2817.2010.tb01278.x

617   Sorensen, T., McLoughlin, P.D., Hervieux, D., Dzus, E., Nolan, J., Wynes, B., Boutin, S., 2008.
618   Determining Sustainable Levels of Cumulative Effects for Boreal Caribou. Journal of
619   Wildlife Management 72, 900–905. https://doi.org/10.2193/2007-079

620   Sothe, C., Gonsamo, A., Arabian, J., Kurz, W.A., Finkelstein, S.A., Snider, J., 2022. Large Soil
621   Carbon Storage in Terrestrial Ecosystems of Canada. Global Biogeochemical Cycles 36.
622   https://doi.org/10.1029/2021GB007213

623   Stewart, F.E.C., Micheletti, T., Cumming, S.G., Barros, C., Chubaty, A.M., Dookie, A.E., Duclos,
624   I., Eddy, I., Hache, S., Hodson, J., Hughes, J., Johnson, C.A., Leblond, M.,
625   Schmiegelow, F.K.A., Tremblay, J.A., McIntire, E.J.B. 2023. Climate-informed forecasts
626   reveal dramatic local habitat change and population uncertainty for threatened caribou in
627   Canada's north. Ecological Applications. 33(3) p.e2816

628   Stewart, F.E.C., Nowak, J.J., Micheletti, T., McIntire, E.J.B., Schmiegelow, F.K.A., Cumming,
629   S.G., 2020. Boreal Caribou Can Coexist with Natural but Not Industrial Disturbances.
630   The Journal of Wildlife Management 84, 1435–1444.

631   Teitelbaum, C.S., Sirén, A.P.K., Coffel, E., Foster, J.R., Frair, J.L., Hinton, J.W., Horton, R.M.,
632   Kramer, D.W., Lesk, C., Raymond, C., Wattles, D.W., Zeller, K.A., Morelli, T.L., 2021.
633   Habitat use as indicator of adaptive capacity to climate change. Divers Distrib ddi.13223.
634   https://doi.org/10.1111/ddi.13223

635   Tootchi, A., Jost, A., Ducharne, A., 2019. Multi-source global wetland maps combining surface
636   water imagery and groundwater constraints. Earth System Science Data 11, 189–220.
637   https://doi.org/10.5194/essd-11-189-2019

638   Travers, H., Selinske, M., Nuno, A., Serban, A., Mancini, F., Barychka, T., Bush, E.,
639   Rasolofoson, R.A., Watson, J.E.M., Milner-Gulland, E.J., 2019. A manifesto for
640   predictive conservation. Biological Conservation 237, 12–18.
641   https://doi.org/10.1016/j.biocon.2019.05.059

642   Wickham, H., 2015. R Packages: Organize, test, document, and share your code. O'Reilly.

643   Wilson, S.F., Nudds, T.D., Vries, A., 2021. A causal modelling approach to informing woodland
644   caribou conservation policy from observational studies. Biological Conservation 264,
645   109370. https://doi.org/10.1016/j.biocon.2021.109370

646   Yates, K.L., Bouchet, P.J., Caley, M.J., Mengersen, K., Randin, C.F., Parnell, S., Fielding, A.H.,
647   Bamford, A.J., Ban, S., Barbosa, A.M., Dormann, C.F., Elith, J., Embling, C.B., Ervin,
648   G.N., Fisher, R., Gould, S., Graf, R.F., Gregr, E.J., Halpin, P.N., Heikkinen, R.K.,
649   Heinänen, S., Jones, A.R., Krishnakumar, P.K., Lauria, V., Lozano-Montes, H.,
650   Mannocci, L., Mellin, C., Mesgaran, M.B., Moreno-Amat, E., Mormede, S., Novaczek, E.,
651   Oppel, S., Ortuño Crespo, G., Peterson, A.T., Rapacciuolo, G., Roberts, J.J., Ross,
652   R.E., Scales, K.L., Schoeman, D., Snelgrove, P., Sundblad, G., Thuiller, W., Torres,
653   L.G., Verbruggen, H., Wang, L., Wenger, S., Whittingham, M.J., Zharikov, Y., Zurell, D.,
654   Sequeira, A.M.M., 2018. Outstanding Challenges in the Transferability of Ecological
655   Models. Trends in Ecology & Evolution 33, 790–802.
656   https://doi.org/10.1016/j.tree.2018.08.001

Table 1. Summary of changes in buffered anthropogenic disturbance, and fire excluding

anthropogenic disturbance (as defined in Johnson et al. 2020), in three scenarios: the Base

Scenario without additional development, Roads Only scenario, and Roads-and-Mines scenario

that included the proposed roads and mining claims within the Ring of Fire area.

Figure 1. The Missisa boreal caribou range (black outline), which includes the proposed Ring of

Fire (RoF) mining claims (dark red), is the focus of this study. The blue dashed line

distinguishes the Managed Forest Zone (MFZ) from the Far North, and shading distinguishes

the Ontario Shield ecozone (pale red) from the Hudson Bay Lowlands(blue). Inlay map (top-

right) indicates the location of the Ontario boreal caribou ranges (grey) relative to Canada.

Figure 2. Maps of the Missisa range with the extent of linear features included in (a) the original

model developed by Hornseth and Rempel (base), (b) the roads-only scenario used to project

the RSF model, and (c) the roads-and-mines scenario used for demographic modelling. Existing

roads are represented as green lines, proposed roads as orange lines, and mines are coloured

in purple.

Figure 3. Reproduction of the seasonal RSF for the Missisa range from Hornseth and Rempel

(2016) using *caribouMetrics* and the published coefficients to reproduce the relative probability

of use (0-1) by boreal caribou during spring, summer, fall, and winter. The predictor variables

used are approximations of those used by Hornseth and Rempel (2016) based on currently

available data. Scale ranges from dark blue to yellow with yellow representing a higher relative

probability of use.

Figure 4. Seasonal RSF predictions from *caribouMetrics* in the Missisa range under the roads

only scenario using the coefficients from Hornseth and Rempel (2016) from the Missisa, James

Bay, Nipigon, and Pagwachuan range to estimate the relative probability of use (0-1) by boreal

caribou during spring, summer, fall, and winter. Scale ranges from dark blue to yellow with

yellow representing a higher relative probability of use.

Figure 5. Demographic rate simulations derived from regression models in Johnson et al.

(2020) for (a) Adult female survival (S), (b) Recruitment (calves per 100 cows), and (c) Average

population rate (λ). Overlap of grey and black dotted lines in (a) and (b) indicates a good match

between expected values from our model (grey) and Johnson et al. 2020 (black). Coloured lines

show effects of changing anthropogenic disturbance on demographic rates in 35 sample

populations, assuming sample populations are randomly distributed among quantiles of the beta

distribution, and each population remains in the same quantile of the beta distribution as

disturbance changes. Alignment of these sample trajectories with 95% predictive intervals from

Johnson et al. 2020 (pale grey bands in panels a and b) indicates that we have adequately

reproduced variability in that model. Bars show the 2.5th and 97.5th percentiles for 500 sample

populations under the three disturbance scenarios for the Missisa range (from left to right, Base,

Road Only, and Roads and Mines). The black diamond indicates the demographic rates for the

Missisa range according to a 2014 assessment (MNRF, 2014a). Populations with an average

population trend of less than 0.99 are considered not self-sustaining.

704 **Table 1**

Scenario	Anthropogenic (%)	Fire (%)	Total Disturbance (%)	Fire Excluding Anthropogenic (%)
Base	0.41	4.27	4.58	4.17
Roads Only	1.11	4.27	5.22	4.12
Roads-and-Mines	6.95	4.27	11.02	4.71

Ecozone
- Hudson's Bay Lowlands
- Ontario Shield
- Managed Forest Boundary
- Mining Claims

Swan

Spirit

Ozhiski

Missisa

James Bay

Berens

Kinloch

Sydney

Churchill

Brightsand

Nipigon

Pagwachuan

Kesagami

Discontinuous Distribution

Lake Superior Coast

0   100   200   300 km

N

Missisa

James Bay

Nipigon

Pagwachuan