Existing caribou habitat and demographic models are poorly suited for Ring of Fire impact

assessment: A roadmap for improving the usefulness, transparency, and availability of models

for conservation

Matt Dyson[1], Sarah Endicott[2], Craig Simpkins[1], Julie W. Turner[3], Stephanie Avery-Gomm[2],

Cheryl A. Johnson[2], Mathieu Leblond[2], Eric W. Neilson[4], Rob Rempel[5], Philip A. Wiebe[6],

Jennifer L. Baltzer[1], Frances E.C. Stewart[1,*], Josie Hughes[2,7,*]

[1]Biology Department, Wilfrid Laurier University, Waterloo, ON, Canada.

matt.e.dyson@gmail.com; simpkinscraig063@gmail.com; jbaltzer@wlu.ca; fstewart@wlu.ca

[2]National Wildlife Research Centre, Environment and Climate Change Canada, Ottawa, ON,

Canada. sarah.endicott@ec.gc.ca; Stephanie.Avery-Gomm@ec.gc.ca; Cheryl-

Ann.Johnson@ec.gc.ca; Mathieu.Leblond@ec.gc.ca; josie.hughes@ec.gc.ca

[3]University of British Columbia, Vancouver, BC, Canada. julwturner@gmail.com

[4]Northern Forestry Centre, Canadian Forest Service, Natural Resources Canada, Edmonton,

AB, Canada. Eric.W.Neilson@NRCan-RNCan.gc.ca

[5]FERIT Environmental Consulting, Thunder Bay, ON, Canada. northernbio@gmail.com

[6]Great Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, Sault Ste.

Marie, ON, Canada. philip.wiebe@NRCan-RNCan.gc.ca

[7]Corresponding Author (josie.hughes@ec.gc.ca)

* Co-PIs

ABSTRACT

Environmental impact assessments often rely on best available information, which may include models that were not designed for purpose and are not accompanied by an assessment of limitations. We reproduced available models of boreal woodland caribou resource selection and demography and evaluated their suitability for projecting impacts of development in the Ring of Fire (RoF) on boreal caribou in the Missisa range (Ontario, Canada). The specificity of the resource selection model limited usefulness for predicting impacts, and high variability in model coefficients among ranges suggests responses vary with habitat availability. The aspatial demographic model projects decreasing survival and recruitment with increasing disturbance, but high variability among populations implies the importance of these impacts depends on the status of the Missisa population, and there no current status estimate. New models that are designed for forecasting the cumulative effects of development and climate change are required to better inform RoF decisions. To demonstrate how open-source tools and reproducible workflows can improve the transparency and reusability of models we developed an R package for data preparation, resource selection, and demographic calculations. Open-source tools, reproducible workflows, and reuseable forecasting models can improve our collective ability to inform wildlife management decisions in a timely manner.

Keywords: *Rangifer tarandus caribou*; Open science; Decision-support; Resource selection function; Demographic model; environmental impact assessment, cumulative effects

INTRODUCTION

Decisions about natural resource development should be informed by sound

environmental impact assessment (Fortin et al., 2020; Hebblewhite, 2017; Johnson et al., 2019;

Johnson and St. Laurent, 2011). Increasingly, these assessments mandate quantitative

appraisals of cumulative effects (e.g. ECCC, 2021). Models that quantitatively link wildlife

species responses to project impacts can influence policy and natural resource development

(Wilson et al., 2021). However, information is not always available or accessible to support

decisions. Practitioners are often required to use tools and models that were not designed to

support impact assessment and may be unaware of their limitations. There is a need to develop

open source, and easily reproduced, wildlife management decision-support tools (Eacker et al.,

2019; Nagy-Reis et al., 2020; Nowak et al., 2018; Stewart et al., 2020) that increase the speed,

scope, and quality of science that informs impact assessment, particularly for Species at Risk

(McIntire et al., 2022; Roche et al., 2020).

Boreal woodland caribou (*Rangifer tarandus caribou*; hereafter 'boreal caribou') inhabit

large portions of Canada's boreal forest (ECCC, 2011; Festa-Bianchet et al., 2011), where

anthropogenic disturbance from the natural resource sector threatens population persistence

(Fortin et al., 2020; Fryxell et al., 2020; Hebblewhite, 2017; Johnson et al., 2020). Nationally,

only 29% of population ranges are considered self-sustaining and boreal caribou are listed as

threatened under the federal Species at Risk Act. In the province of Ontario, boreal caribou are

also listed as threatened due to development from forestry and mineral exploration (ECCC,

2011; MNRF, 2014a). Commercial logging and associated road development have contributed

to landscape disturbance and are predicted to cause continued population declines (Fryxell et

al., 2020).

The Hudson Plains of Northern Ontario, Canada, is one of the largest intact wetlands

and peatland carbon stores on the planet (Ibisch et al., 2016; Poley et al., 2022; Sothe et al.,

2022; Tootchi et al., 2019). This region is rich in valuable minerals and is a target for resource

extraction and economic development (Carlson and Chetkiewicz, 2013; Far North Science

Advisory Panel (Ont.) et al., 2011; IAAC, 2021). The region is home to Indigenous people and

wildlife, including threatened boreal caribou, and is largely inaccessible by road. There is a need

to assess the potential environmental impacts of proposed mining of 'Ring of Fire' (RoF) mineral

deposits and associated development in this globally significant ecosystem. Development

decisions in the RoF will impact caribou and tools are required to quantify this impact (Far North

Science Advisory Panel (Ont.) et al., 2011; IAAC, 2021; Fig. 1).

Information to support impact assessments comes in many forms, including models of

wildlife-habitat relationships and demography (Beyer et al., 2010; Matthiopoulos et al., 2020).

Wildlife-habitat relationships are commonly represented by resource selection functions (RSFs),

which can produce maps to predict habitat suitability and infer the consequences of habitat loss

(Boyce et al., 2002; Harju et al., 2011; Johnson et al., 2004; Matthiopoulos et al., 2019).

However, RSFs represent descriptive associations between animal behaviour and habitat under

current, scale-specific conditions, limiting their predictive capacity and transferability to novel

conditions (Boyce, 2006; Decesare et al., 2012; Harju et al., 2011; Kunegel-Lion et al., 2022;

Mysterud and Ims, 1998; Yates et al., 2018). Aspatial demographic models that provide insight

into drivers of population growth, such as survival and recruitment, are also used for

characterizing wildlife responses to landscape alteration (Johnson et al., 2020; Sorensen et al.,

2008; Stewart et al., 2020). However, high uncertainty about demographic parameters, drivers

of change, and vital rates can limit the utility of these models for impact assessment (Chaudhary

and Oli, 2020; Sleep and Loehle, 2010). Regardless of their potential limitations, RSFs and

demographic models often represent the best available information to be used to support

conservation decision making.

In Ontario, Canada, existing RSF (Hornseth and Rempel, 2016) and demographic

models (Johnson et al., 2020) are used to support decisions provincially (Rempel et al., 2021)

and federally (ECCC, 2019). These models represent the best available information for the RoF,

but their utility for projecting impacts has not been formally evaluated, nor are they easy to

scrutinize or apply. We reproduced the available RSF and demographic models and evaluated

their suitability to predict some of the anticipated impacts of proposed mining in the RoF. To

demonstrate the utility of open source tools and reproducible workflows in this context, we used

an R package framework (Wickham, 2015) to create an R package that reproduces the existing

RSF and demographic models. Our package can be integrated into predictive frameworks to

support resource use decisions or modified and advanced by other practitioners for other uses

(Bodner et al., 2021b; McIntire et al., 2022; Micheletti et al., 2021; Miller and Frid, 2022). We

identify and discuss challenges with using existing models for decision-support and

opportunities for improving the usefulness, transparency, and availability of environmental

impact assessment tools for SAR.

Methods

Study Area

Our study area included caribou ranges as delineated by Ontario's provincial

government that contain and surround the RoF mining claims: Missisa, James Bay,

Pagwachuan, and Nipigon, with most mineral claims being situated in the Missisa range (Far

North Science Advisory Panel (Ont.) et al., 2011; IAAC, 2021; Fig. 1). The Missisa and James

Bay ranges occur in Ontario's Far North (Far North Science Advisory Panel (Ont.) et al., 2011),

where anthropogenic disturbance has thus far been limited to communities and exploration

camps, winter roads, and the now inactive Victor Diamond Mine. The Nipigon and Pagwachuan

ranges are almost entirely within the more productive southern portion of the Boreal Shield

ecozone (Crins et al., 2009) where industrial forestry is common (the Managed Forest Zone or

MFZ; Fig. 1). The Boreal Shield consists of mixed and coniferous forests dominated by black

spruce (*Picea mariana*), and containing jack pine (*Pinus banksiana*), balsam fir (*Abies*

balsamea), white spruce (*Picea glauca*), trembling aspen (*Populus tremuloides*), and balsam

poplar (*Populus balsamifera*) (Crins et al., 2009). The Pagwachuan range also includes a

portion of the Boreal Plains and the Great Clay Belt. The Missisa range includes the transition

between the Boreal Shield and Hudson Plains, which is mainly composed of peatland

complexes with poor drainage, with trees including stunted black spruce and tamarack (*Larix*

laricina) (Crins et al., 2009). The James Bay range is almost entirely within the Hudson Plains.

We compare the Missisa range to adjacent ranges to investigate the transferability of range-

specific RSFs to different landscape conditions.

Ring of Fire Development Scenarios

We considered three simple development scenarios (Fig 2); a 'base' scenario with no

change in development footprint, a 'roads-only' scenario that includes proposed RoF access

roads (MNDMNRF, 2022), and a 'roads-and-mines' scenario which includes the proposed

access roads and mining claims associated with the RoF (Table S1.2) . Our intention is not to

speculate about the scale of future developments, but to assess the suitability of existing

models for projecting impacts of development on boreal caribou.

RSF Model Reproduction

We used tables of published boreal caribou range- and season-specific RSF coefficients

for Ontario (Hornseth and Rempel, 2016; Rempel et al., 2021), hereafter referred to as the

original RSFs. The authors referred to these models as resource selection probability functions

(RSPFs), but their approach differs from what is commonly understood as an RSPF (Johnson et

al., 2006) so we use the term RSF. The data to produce the original RSFs were collected from

GPS collared caribou across a variety of companion studies within the ranges of interest

occurring between 2009 and 2013 (Hornseth and Rempel, 2016; MNRF, 2014b, 2014b; Rempel

and Hornseth, 2018). We obtained the published coefficients for range-specific seasonal RSFs

of each reported top model (Table S1.1) for the Nipigon, Pagwachuan, Missisa, and James Bay

range, but information on the uncertainty associated with these coefficient estimates is not

available. The predictors used in the original RSFs were developed from land cover, forest fire

and harvest disturbance history, linear features, and esker datasets but the original data was

not available (Table S1.2, S1.3). To reproduce the original RSFs, we acquired data sets for the

predictor variables that were publicly available and, where possible, used information on the

timing of the event or the feature's construction to recreate the data that was available when the

original model was produced (Table S1.2). We used current versions of these same datasets to

project the RSFs on the landscape.

RSF Model Validation

To assess our ability to reproduce the original model, we acquired the original projected

surfaces for comparison from the authors (R. Rempel pers. comm, 2021). We validated our

models visually and quantitatively. The original model used a nested hexagon grid to generate

predictions. We opted to approximate this approach with distance-weighted moving windows on

a rectangular grid, which is easier to implement using standard raster-processing tools. To

compare these approaches, we extracted the values from our rectangular grid to the hexagonal

grid using the weighted mean. We mapped the difference in predictions and explanatory

variables between the models and used scatterplots to compare the value of the model

response for each grid cell produced by the two models. A perfect reproduction would produce a

Pearson correlation coefficient of 1, and any deviation from the original prediction would reduce

this coefficient. We expected some differences in the predictions as a result of the type of grid

and predictor data sets used.

RSF Model Projection Under Disturbance Scenarios

The RSF includes road density as a predictor and would require additional information

on roads within mining areas that we do not have; for the RSF we only compare the base and

road-only scenarios. To understand how the 'roads only' scenario would affect the caribou RSF,

we projected the original RSFs across updated landscape conditions as represented by i)

temporal changes in forest structure between 2010 and 2020 (e.g., fires; Table S1.2) and ii) the

new proposed roads only scenario (Fig. 2). We also assessed the potential for borrowing

information from other ranges by using the coefficients from the top original RSFs from the

adjacent James Bay, Nipigon, and Pagwachuan ranges transferred to Missisa. We visualized

the spatial transferability of the models and examined their sensitivity to changing habitat

availability using scatterplots to compare the value of the model response for each grid cell

produced by the Missisa model and each respective adjacent range.

Demographic Model

Canada's national demographic boreal caribou models were developed from adult

female survival (*S*), calf recruitment (*R*), and landscape data across 58 boreal caribou study

areas, including 13 study areas in Ontario (Johnson et al., 2020; Table S1). We used these

models (Johnson et al., 2020) to predict changes in *S* and *R*; in the roads-only and roads-and-

mines development scenarios. The demographic model is aspatial and all types of

anthropogenic disturbances are combined into a single measure of disturbed area, so the

'roads-and-mines' footprint is sufficient and there is no need to specify the location of roads

within mining claim areas. We calculated the relevant predictor variables for the Missisa range

(e.g., % anthropogenic disturbance buffered by 500 m; % wildfire within the last 40 years; Table

1), and calculated expected *R* and *S* as a function of disturbance according to the beta

regression models with highest support (M4 and M1 respectively from Johnson et al., 2020):

$$R \sim Beta(\mu^R, \phi^R); log(\mu^R) = \beta_0^R + \beta_a^R\ anthro\ + \beta_f^R\ fireExclAnthro, \quad \text{(eq 1a)}$$

$$S \sim (46 \times Beta(\mu^S, \phi^S) - 0.5)/45; log(\mu^S) = \beta_0^S + \beta_a^S\ anthro, \quad \text{(eq 1b)}$$

Where $\square^R$ ~ *Normal(19.862,2.229)* and $\square^S$ ~ *Normal(63.733,8.311)* are precisions of the Beta

distributed errors (Ferrari and Cribari-Neto, 2004), and survival rates are back transformed as in

Johnson et al. 2020. Table 3 of Johnson et al. (2020) provides the expected values and 95%

confidence intervals of all regression coefficients (β_0^R, β_a^R, β_f^R, β_0^S, β_a^S), which are assumed to

be Gaussian distributed. To evaluate our recruitment and survival models, we sampled

expected demographic rates across a range of anthropogenic and fire disturbance (0-100%) to

reproduce expected values and 95% predictive intervals from Fig. 3 and Fig. 5 of Johnson et al.

(2020).

In areas of low anthropogenic disturbance, substantial among-population variability in

recruitment existed (Fig. 3 in Johnson et al., 2020). To model this variation, regression model

parameter values for each sample population were selected at the beginning of simulations and

each population was assigned to quantiles of the error distributions for survival and recruitment.

Sample populations remained in their quantiles as the landscape changed, allowing us to

distinguish the effects of changing disturbance from variation in initial population status. To

show these effects, we projected population growth for 35 sample populations across a wide

range of anthropogenic disturbance levels (0-90%) using a two-stage demographic model with

density dependence and interannual variability that differed from the model used by Johnson et

al. (2020) in the method used to ensure whole numbers of animals (Supplementary Material

Part 2). We characterized the distribution of outcomes for our three disturbance scenarios (Fig.

2, Table 1) more thoroughly by projecting the dynamics of 500 sample populations. Population

growth rate each year is given by $\lambda_t = N_{t+1}/N_t$ when $N_t > 0$ and $\lambda_t = 0$ when $N_t = 0$. For each sample

population, population growth was projected for 20 years from an assumed initial population

size of 373 females, which we derived from the minimum animal count of 745 caribou between

2009 and 2011 and assumed 50% were female (MNRF, 2014b), and we report the average

growth rate λ over that time. This estimate is conservative since adult male:female ratios are

typically lower (ECCC, 2011; Espinoza and Weckerly, 2021). To verify our reproduction of the

demographic model used by Johnson et al. (2020), we compared our outputs to those from

model code supplied by the authors.

R Package Framework: caribouMetrics

We incorporated RSF and demographic model components into an R package with

documentation and vignettes explaining their use, and used GitHub (github, 2020) to promote

version control and transparency of the development process:

https://github.com/LandSciTech/caristMetrics. The standardized nature of packages allows

them to easily integrate into the larger R ecosystem, allowing them to be extended and adapted

towards tasks beyond their original design (e.g. prediction or forecasting; McIntire et al., 2022).

cariboMetrics also contains functions that automate the geospatial data preparation process to

facilitate application to new landscapes. The original RSFs were developed using a closed

scripting language called Landscape Scripting Language (LSL; Kushneriuk and Rempel, 2011),

which we re-implemented in R. For the aspatial demographic model, we began by borrowing

demographic rate sampling code from a SpaDES (Chubaty and McIntire, 2022) module

(*cariboPopGrowth*; https://github.com/tati-micheletti/cariboPopGrowthModel; Stewart et al., in

review), and modified the method to include precision and quantiles. We extracted the two-

stage demographic model with density dependence and interannual variability from unpublished

code (Johnson et al., 2020), and modified the code to improve transparency and ensure whole

numbers of animals (see Supplementary Material Part 2).

Results

RSF Model Reproduction and Validation

We were able to produce a reasonably accurate representation of the original RSFs for

the Missisa range, as evidenced by high Pearson's r values across all seasons (r > 0.935; Fig.

S1.1). Maps highlighting the differences between predictions and some variation in the input

data layers are provided in Supplementary Material (Fig. S1.2, S1.3). In the Missisa range, the

model predicted the highest relative use probabilities in the northwest of the study area during

winter (Fig. 3), consistent with the original RSFs. During the summer and spring, the eastern

portion of the range was used more than the northwest (Fig. 3).

RSF Model Projection

We observed a lower relative probability of use in areas associated with proposed roads

(Fig. 4). Lower relative probability of use along the road corridors was strongest in the spring

and summer, consistent with seasonal changes in the response to roads described in the

original RSFs (Fig. 4). There was high variability in the estimated response to roads among

ranges (Fig. 4). The James Bay range prediction appeared the most similar to the Missisa range

prediction (Fig. 4); however it varied by season (Fig S1.4). The Nipigon and Pagwachuan range

projections, which visually differed substantially from the Missisa projection, did not show a

strong response to proposed roads (Fig. 4; Fig S1.4).

Demographic model

Comparison with Johnson et al. (2020) indicates a good reproduction of the regression

models for survival and recruitment (Fig. 5). Anthropogenic disturbance remains relatively low in

all our development scenarios (Table 1), and the corresponding range of variability in

demographic rates among sample populations is high (Fig. 5). The model predicts increasing

anthropogenic disturbance will decrease both survival and recruitment (Fig. 5), but the

importance of that decrease for self-sustainability of the population is highly uncertain and

depends on initial population status (Fig. 5). A 2014 assessment informed by data from 2008-

2012 indicated lower than expected recruitment, survival, and population growth rate in the

Missisa range (diamonds in Fig. 5; MNRF, 2014a); however, we lack recent information on the

survival, recruitment and status of this population to project the impacts of disturbance.

Discussion

We examined the applicability of two existing caribou models for projecting impacts of

development in the RoF on boreal caribou in the Missisa range. We highlight limitations of

existing tools and point out possible solutions. We found that the original RSFs are poorly suited

for projecting the impacts of development in the RoF because they are specific to current

conditions. In contrast to the RSFs, the existing aspatial demographic model is too general to

project the impacts of development on this particular population. Ideally, forecasting models

used to inform environmental decisions should be designed for the purpose, identify and

account for uncertainty, be updateable with new information, and be transparent and

reproducible (Bodner et al., 2021a). However, in practice, environmental impact assessments

often rely on best available information, which may include tools and models that were not

designed for the purpose. We highlight limitations of existing tools and point out possible

solutions.

The RSF models for each caribou range were fit independently using data from that

range. This is a reasonable approach when the objective is to characterize current habitat use,

but not for projecting responses to changing landscape conditions. There has been very little

anthropogenic disturbance in the Missisa range, so projecting impacts of disturbance on boreal

caribou requires use of information from more disturbed areas. However, as formulated, the

RSFs do not allow borrowing of information. There was high variability in the effect of linear

features among range-specific RSFs, suggesting either that the behavioral response of caribou

to linear features may vary with the amount or type of disturbance in a range (i.e. functional

response; Mysterud and Ims, 1998), or that effects of linear features were confounded with

other correlated predictors (Hornseth and Rempel, 2016). Hierarchical regional models that

include functional responses to disturbance (Matthiopoulos et al., 2011; Muff et al., 2020; Olson

et al., 2021; Teitelbaum et al., 2021) could help distinguish between these possibilities, and

integrated step-selection analysis approaches could yield models that are better suited for

projection (Avgar et al., 2016; Passoni et al., 2021; Prokopenko et al., 2017). The original RSFs

used road density as a proxy for other anthropogenic disturbances (e.g., harvest) and did not

distinguish among road types (Hornseth and Rempel, 2016). Evidence from other locations and

contexts suggests that road type and traffic volume is important for wildlife, and metrics that

include this information may be more informative (D'Amico et al., 2016; Jaeger et al., 2005;

Leblond et al., 2013; Loosen et al., 2021). Although projecting impacts in a unique and

previously undisturbed region is inherently challenging, we are optimistic that more informative

models can be developed. In the meantime, these limitations should be considered when using

or interpreting the original RSFs.

Among caribou ranges where anthropogenic disturbance was low, we saw that the

demographic model predicted high variation in recruitment and survival among populations (Fig.

3 and Fig. 5 of Johnson et al., 2020), leading to high variability in projections of the impact of

increasing disturbance on population growth rate (Fig. 5). This variability highlights the need for

local information about the current status of caribou populations in the RoF and ongoing

monitoring of development impacts in the area. An assessment of the status of the Missisa

population based on 2008 – 2012 data (MNRF, 2014b) suggested survival, reproduction, and

growth rate in this range was lower than predicted by the aspatial national demographic model

of Johnson et al. 2020 (see black diamond in Fig. 5). More frequent and consistent data to

estimate survival, reproduction, or population size are required to understand population trends

and status. The national demographic model (Johnson et al., 2020) predicts that increasing

anthropogenic disturbance will decrease both survival and recruitment, but the importance of

that decrease for population viability depends on population status. Even small changes in adult

female survival can affect the sustainability of a population in ranges with low adult survival

(Johnson et al., 2020). Without information on the current status of the population, it is difficult to

predict impacts of disturbance associated with RoF development on the viability of this particular

population with any degree of confidence.

Our method of modelling variation in survival and recruitment among populations (Fig. 5)

adequately reproduced the observed variation among populations (Fig. 3 and Fig. 5 in Johnson

et al., 2020), but differed from the range-specific scenario approach taken by Johnson et al.

(2020). We also opted to use a different method for ensuring whole numbers of animals, which

lead to different estimates of population growth rate for small populations (see Supplementary

Material Part 2). We assumed that demographic rates (i.e., recruitment and survival) vary with

disturbance according to the best supported national models (Table 2 of Johnson et al., 2020)

but note that other competing models in that candidate set were nearly as well supported by the

data. This simple population model also assumed no variation in recruitment or survival with age

or other parameters (Supplementary Material Part 2). Most female caribou reproduced at 2-3

years of age rather than at 1 year, as assumed in the model. This likely results in overestimating

demographic parameter values under changing conditions. A more thorough investigation of the

sensitivity of results, parameters, and representation of uncertainty and stochasticity in aspatial

caribou demographic models is beyond the scope of this paper but seems warranted.

We lack region-specific information on disturbance history, vegetation, and vegetation

recovery trajectories post-disturbance that are required for assessing impacts of climate change

and anthropogenic disturbance in the RoF (McLaughlin and Webster, 2014; McLaughlin and

Packalen, 2021). There may also be variability in caribou responses to fire (Konkolics et al.,

2021; Palm et al., 2022) and interactions with predators (Bergerud, 1974) as has been found in

other ranges. Disturbance-mediated apparent competition is generally the accepted mechanism

for boreal caribou declines across Canada (Festa-Bianchet et al., 2011; Serrouya et al., 2016;

Wittmer et al., 2013), but recent studies have noted that apparent competition may be less

important in environments where above ground productivity is low because of the absence or

lower abundance of other ungulate competitors (Superbie et al., 2022). The James Bay

Lowlands, which make up a considerable portion of the Missisa range (Fig. 1), are a unique low

productivity peatland ecosystem that currently experiences little fire (Abraham and Keddy,

2005). Climate change is predicted to alter fire regimes, but even if fire risk increases, this area

will likely remain relatively wet and resistant to fire (Balshi et al., 2009; Wang et al., 2020). Given

the uniqueness of this area, area-specific information is required; we demonstrate that data and

models from other regions is not sufficient for understanding the effects of disturbance on

wildlife-habitat relationships and boreal caribou demography in the RoF (MNRF, 2014a). Even

where relevant data exists, political and social barriers to data access make it difficult or

impossible to do synthetic work that would yield better insight into consistencies and differences

among regions (Rutz et al., 2020; Tucker et al., 2018).

There are various efforts underway to collect data needed to inform resource

development and wildlife management decisions in the RoF, yielding opportunities for

collaborative development of region-specific open source forecasting models. We hope that our

investigation of the limitations of existing models, and demonstration of the usefulness of

implementing models an R package such as *caribouMetrics*, can help inform the development

of new models and tools. Hierarchical regional models that include functional responses to

disturbance (Matthiopoulos et al., 2011; Muff et al., 2020; Olson et al., 2021; Teitelbaum et al.,

2021), integrated step-selection analysis (Avgar et al., 2016; Prokopenko et al., 2017), and

hierarchical integrated population models (Moeller et al., 2021; Nowak et al., 2018) would yield

more reliable projections of impacts in this previously undisturbed region. Open, transparent,

reproducible workflows should be designed to enable ongoing incorporation of new data into

models (Dietze, 2017; McIntire et al., 2022; Micheletti et al., 2021). Models should also be

designed for flexibility and nimbleness, allowing decision makers to quickly adapt to specific

contexts without duplicating efforts (Bodner et al., 2021a; McIntire et al., 2022; Travers et al.,

2019); an R package framework provides a practical option for achieving these goals.

We reiterate a call for transparent, quantitative decision-support tools to assess

industrial development impacts on wildlife, encourage developers of wildlife response models

and collectors of relevant wildlife data to work together toward this goal (Davidson et al., 2021;

Russell et al., 2021), and are encouraged by progress toward knowledge sharing and

collaboration (e.g. CCLM, 2019). We also recognize that the development of usable decision-

support tools requires time and skills that are not possessed by everyone interested in doing

open science. One solution is to work in multi-disciplinary teams (Bodner et al., 2021a). Another

is to recognize that steps toward transparency and reproducibility are valuable, even if the result

is not always an easily usable tool. In this project, we were able to reproduce existing models

because the developers of those models were willing to share their code. Code does not have

to be flawless to enable others to build on previous work, and there are many ways that

researchers can shift to more open and reproducible workflows that reduce the chance of

errors, increase efficiency, improve reproducibility, and increase our ability to generalize across

studies (Alston and Rick, 2021; Lewis et al., 2018). We are hopeful that improving the

transparency, reproducibility and decision-relevance of wildlife response models will improve

our collective ability to inform decisions for SAR and improve conservation outcomes.

Acknowledgements

We thank Judith Girard, Samantha McFarlane, Raj Bhatti, Shari Hayne, Megan Hornseth, and Glenn D. Sutherland for discussion and suggestions that helped improve this manuscript. We also thank Eliot McIntire, Tati Micheletti, Alex Chubaty, and the Predictive Ecology group for their support and contributions to R package development.

Competing Interests Statement: Rob Rempel is principal of FERIT Consulting.

Author Contribution Statement: Matt Dyson led the writing and co-led the analysis; Sarah Endicott co-led package development and provided support for analysis and writing; Craig Simpkins co-led package development and contributed to writing; Julie Turner provided contributions to writing and evaluation of the RSF; Stephanie Avery-Gomm, Cheryl A. Johnson, Mathieu Leblond, Eric Neilson, Rob Rempel, Philip Wiebe (listed in alphabetical order) provided advice on package development, analysis, and writing; Rob Rempel and Cheryl A. Johnson also provided code and insight into the RSFs and demographic models, respectively. Frances Stewart and Jennifer Baltzer were project leads at Wilfrid Laurier University and Josie Hughes was project lead at ECCC.

Funding Statement: Funding for this work was provided by the Canadian Wildlife Service and the Wildlife and Landscape Science Directorate of Environment and Climate Change Canada. FEC Stewart was supported by a CRC in Northern Wildlife Biology and J. Baltzer was supported by a CRC in Forest and Global Change.

Data availability statement: The caribouMetrics R package is available at https://github.com/LandSciTech/caribouMetrics, and analysis code is available at https://github.com/LandSciTech/MissisaBooPaper. Data required for analysis is available in this OSF repository: https://osf.io/r9mkp/?view_only=fb71321265d14dbeb3d932e4de66be0c

References

Abraham, K.F., Keddy, C.J., 2005. The Hudson Bay Lowland, in: Fraser, L.H., Keddy, P.A. (Eds.), The World's Largest Wetlands: Ecology and Conservation. Cambridge University Press, Cambridge, UK, pp. 118–148.

Alston, J.M., Rick, J.A., 2021. A beginner's guide to conducting reproducible research. Bull Ecol Soc Am. https://doi.org/10.1002/bes2.1801

Avgar, T., Potts, J.R., Lewis, M.A., Boyce, M.S., 2016. Integrated step selection analysis: Bridging the gap between resource selection and animal movement. Methods in Ecology and Evolution 7, 619–630. https://doi.org/10.1111/2041-210X.12528

Balshi, M.S., McGuire, A.D., Duffy, P., Flannigan, M., Kicklighter, D.W., Melillo, J., 2009. Vulnerability of carbon storage in North American boreal forests to wildfires during the 21st century. Global Change Biology. 15: 1491-1510 15, 1491–1510. https://doi.org/10.1111/j.1365-2486.2009.01877.x

Bergerud, A.T., 1974. Decline of Caribou in North America Following Settlement. The Journal of Wildlife Management 38, 757. https://doi.org/10.2307/3800042

Beyer, H.L., Haydon, D.T., Morales, J.M., Frair, J.L., Hebblewhite, M., Mitchell, M., Matthiopoulos, J., 2010. The interpretation of habitat preference metrics under use-availability designs. Philisophical Transactions of The Royal Society of Biological Sciences 365, 2245–2254.

Bodner, K., Brimacombe, C., Chenery, E.S., Greiner, A., McLeod, A.M., Penk, S.R., Vargas Soto, J.S., 2021a. Ten simple rules for tackling your first mathematical models: A guide for graduate students by graduate students. PLoS Comput Biol 17, e1008539. https://doi.org/10.1371/journal.pcbi.1008539

Bodner, K., Rauen Firkowski, C., Bennett, J.R., Brookson, C., Dietze, M., Green, S., Hughes, J., Kerr, J., Kunegel-Lion, M., Leroux, S.J., McIntire, E., Molnár, P.K., Simpkins, C., Tekwa, E., Watts, A., Fortin, M.-J., 2021b. Bridging the divide between ecological forecasts and environmental decision making. Ecosphere 12, e03869. https://doi.org/10.1002/ecs2.3869

Boyce, M.S., 2006. Scale for resource selection functions. Diversity and Distributions 12, 269–276. https://doi.org/10.1111/j.1366-9516.2006.00243.x

Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions. Ecological Modelling 157, 281–300.

Carlson, M., Chetkiewicz, C., 2013. A fork in the road: future development in Ontario's Far North. Canadian Boreal Initiative, Ottawa, ON.

CCLM, 2019. Canadian Conservation and Land Management Knowledge Network [WWW Document]. Canadian Conservation and Land Management Knowledge Network. URL https://www.cclmportal.ca/home (accessed 6.8.22).

Chaudhary, V., Oli, M.K., 2020. A critical appraisal of population viability analysis. Conservation Biology 34, 26–40. https://doi.org/10.1111/cobi.13414

Chubaty, A.M., McIntire, E.J.B., 2022. SpaDES: Develop and Run Spatially Explicit Discrete Event Simulation Models.

Crins, W.J., Gray, P.A., Uhlig, P.W.C., Wester, M.C., 2009. Ecosystems of Ontario. Part 1, Part 1, (No. SIB TER IMA TR-01). Ontario, Ministry of Natural Resources, Inventory, Monitoring and Assessement Section, Toronto, Ont.

D'Amico, M., Périquet, S., Román, J., Revilla, E., 2016. Road avoidance responses determine the impact of heterogeneous road networks at a regional scale. Journal of Applied Ecology 53, 181–190. https://doi.org/10.1111/1365-2664.12572

Davidson, S., Bohrer, G., Kölzsch, A., Vinciguerra, C., Kays, R., 2021. Mobilizing Animal Movement Data: API use and the Movebank platform. BISS 5, e74312. https://doi.org/10.3897/biss.5.74312

Decesare, N.J., Hebblewhite, M., Schmiegelow, F., Hervieux, D., McDermid, G.J., Neufeld, L.,
Bradley, M., Whittington, J., Smith, K.G., Morgantini, L.E., Wheatley, M., Musiani, M.,
2012. Transcending scale dependence in identifying habitat with resource selection
functions. Ecological Applications 22, 1068–1083.
Dietze, M., 2017. Ecological Forecasting. Princeton University Press, New Jersey.
Eacker, D.R., Hebblewhite, M., Steenweg, R., Russell, M., Flasko, A., Hervieux, D., 2019.
Web-based application for threatened woodland caribou population modeling. Wildl.
Soc. Bull. 43, 167–177. https://doi.org/10.1002/wsb.950
ECCC, 2021. Regional Assessment under the Impact Assessment Act.
ECCC, 2019. Amended Recovery Stategy for the Woodland Caribou (Rangifer tarandus
caribou), Boreal population, in Canada [Proposed], Species at Risk Act Recovery
Stategy Series. Environment and Climate Change Canada, Ottawa.
ECCC, 2011. Scientific assessment to inform the identification of critical habitat for woodland
caribou (Rangifer tarandus caribou), boreal population, in Canada. Canadian Wildlife
Service, Ottawa.
Espinoza, Z.S., Weckerly, F.W., 2021. A Comparative Analysis of Adult Sex Ratios in
Polygynous and Monogamous Mammal Populations. amid 186, 299–308.
https://doi.org/10.1674/0003-0031-186.2.299
Far North Science Advisory Panel (Ont.), Ontario, Ministry of Natural Resources, 2011. Science
for a changing far north: the report of the Far North Science Advisory Panel. Queen's
Printer for Ontario, Toronto, Ont.
Ferrari, S., Cribari-Neto, F., 2004. Beta Regression for Modelling Rates and Proportions.
Journal of Applied Statistics 31, 799–815.
https://doi.org/10.1080/0266476042000214501
Festa-Bianchet, M., Ray, J.C., Boutin, S., Côté, S.D., Gunn, A., 2011. Conservation of caribou (
Rangifer tarandus) in Canada: an uncertain future. Can. J. Zool. 89, 419–434.
https://doi.org/10.1139/z11-025
Fortin, D., Mcloughlin, P.D., Hebblewhite, M., 2020. When the protection of a threatened
species depends on the economy of a foreign nation 1–16.
https://doi.org/10.1371/journal.pone.0229555
Fryxell, J.M., Avgar, T., Liu, B., Baker, J.A., Rodgers, A.R., Shuter, J., Thompson, I.D., Reid,
D.E.B., Kittle, A.M., Mosser, A., Newmaster, S.G., Nudds, T.D., Street, G.M., Brown,
G.S., Patterson, B., 2020. Anthropogenic disturbance and population viability of
woodland caribou in Ontario. Jour. Wild. Mgmt. 84, 636–650.
https://doi.org/10.1002/jwmg.21829
github, 2020. GitHub.
Harju, S.M., Dzialak, M.R., Osborn, R.G., Hayden-Wing, L.D., Winstead, J.B., 2011.
Conservation planning using resource selection models: altered selection in the
presence of human activity changes spatial prediction of resource use: Human activity
alters spatial use predictions. Anim Conserv 14, 502–511. https://doi.org/10.1111/j.1469-
1795.2011.00456.x
Hebblewhite, M., 2017. Billion dollar boreal woodland caribou and the biodiversity impacts of the
global oil and gas industry. Biological Conservation 206, 102–111.
https://doi.org/10.1016/j.biocon.2016.12.014
Hornseth, M.L., Rempel, R.S., 2016. Seasonal resource selection of woodland caribou (
Rangifer tarandus caribou) across a gradient of anthropogenic disturbance. Can. J.
Zool. 94, 79–93. https://doi.org/10.1139/cjz-2015-0101
IAAC, 2021. Draft Agreement to Conduct a Regional Assessment in the Ring of Fire Area.
Ibisch, P.L., Hoffmann, M.T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L., DellaSala,
D.A., Vale, M.M., Hobson, P.R., Selva, N., 2016. A global map of roadless areas and

their conservation status. Science 354, 1423–1427. https://doi.org/10.1126/science.aaf7166

Jaeger, J.A.G., Bowman, J., Brennan, J., Fahrig, L., Bert, D., Bouchard, J., Charbonneau, N., Frank, K., Gruber, B., von Toschanowitz, K.T., 2005. Predicting when animal populations are at risk from roads: an interactive model of road avoidance behavior. Ecological Modelling 185, 329–348. https://doi.org/10.1016/j.ecolmodel.2004.12.015

Johnson, C.A., Sutherland, G.D., Neave, E., Leblond, M., Kirby, P., Superbie, C., McLoughlin, P.D., 2020. Science to inform policy: Linking population dynamics to habitat for a threatened species in Canada. J Appl Ecol 57, 1314–1327. https://doi.org/10.1111/1365-2664.13637

Johnson, C.J., Nielsen, S.E., Merrill, E.H., Trent, L., Boyce, M.S., Science, E., Program, M., British, N., George, P., 2006. Resource Selection Functions Based on Use – Availability Data□: Theoretical Motivation and Evaluation Methods. Journal of Wildlife Management 70, 347–357. https://doi.org/10.2193/0022-541X(2006)70

Johnson, C.J., Seip, D.R., Boyce, M.S., 2004. A quantitative approach to conservation planning: using resource selection functions to map the distribution of mountain caribou at multiple spatial scales. J Appl Ecology 41, 238–251. https://doi.org/10.1111/j.0021-8901.2004.00899.x

Johnson, C.J., St. Laurent, M.-H., 2011. Unifying framework for understanding impacts of human developments on wildlife, in: Naugle, D.E. (Ed.), Energy Development and Wildlife Conservation in Western North America. Island Press, Washington, D.C., p. 305.

Johnson, C.J., Venter, O., Ray, J.C., Watson, J.E.M., 2019. Growth-inducing infrastructure represents transformative yet ignored keystone environmental decisions. Conservation Letters 1–7. https://doi.org/10.1111/conl.12696

Konkolics, S., Dickie, M., Serrouya, R., Hervieux, D., Boutin, S., 2021. A Burning Question: What are the Implications of Forest Fires for Woodland Caribou? Jour. Wild. Mgmt. jwmg.22111. https://doi.org/10.1002/jwmg.22111

Kunegel-Lion, M., Neilson, E.W., Mansuy, N., Goodsman, D.W., 2022. Habitat quality does not predict animal population abundance on frequently disturbed landscapes. Ecological Modelling 469, 109943. https://doi.org/10.1016/j.ecolmodel.2022.109943

Kushneriuk, R.S., Rempel, R.S., 2011. LSL - Landscape Scripting Language. Ontario Ministry of Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, ON.

Leblond, M., Dussault, C., Ouellet, J.-P., 2013. Avoidance of roads by large herbivores and its relation to disturbance intensity: Avoidance of roads and disturbance intensity. J Zool 289, 32–40. https://doi.org/10.1111/j.1469-7998.2012.00959.x

Lewis, K.P., Vander Wal, E., Fifield, D.A., 2018. Wildlife biology, big data, and reproducible research. Wildlife Society Bulletin. https://doi.org/10.1002/wsb.847

Loosen, A.E., Devineau, O., Zimmermann, B., Cromsigt, J.P.G.M., Pfeffer, S.E., Skarpe, C., Marie Mathisen, K., 2021. Roads, forestry, and wolves interact to drive moose browsing behavior in Scandinavia. Ecosphere 12. https://doi.org/10.1002/ecs2.3358

Matthiopoulos, J., Fieberg, J., Aarts, G., 2020. Species-Habitat Associations: Spatial data, predictive models, and ecological insights, 1st ed. University of Minnesota Libraries. https://doi.org/10.24926/2020.081320

Matthiopoulos, J., Field, C., MacLeod, R., 2019. Predicting population change from models based on habitat availability and utilization. Proceedings of the Royal Society B: Biological Sciences 286. https://doi.org/10.1098/rspb.2018.2911

Matthiopoulos, J., Hebblewhite, M., Aarts, G., 2011. Generalized functional responses for species distributions. Ecology 92, 583–589.

McIntire, E.J.B., Chubaty, A.M., Cumming, S.G., Andison, D., Barros, C., Boisvenue, C., Haché, S., Luo, Y., Micheletti, T., Stewart, F.E.C., 2022. PERFICT A Re-imagined foundation for predictive ecology.pdf. Ecology Letters Online, 1–7.

McLaughlin, J., Webster, K., 2014. Effects of Climate Change on Peatlands in the Far North of Ontario, Canada: A Synthesis. Arctic, Antarctic, and Alpine Research 46, 84–102. https://doi.org/10.1657/1938-4246-46.1.84

McLaughlin, J.W., Packalen, M.S., 2021. Peat Carbon Vulnerability to Projected Climate Warming in the Hudson Bay Lowlands, Canada: A Decision Support Tool for Land Use Planning in Peatland Dominated Landscapes. Front. Earth Sci. 9, 650662. https://doi.org/10.3389/feart.2021.650662

Micheletti, T., Stewart, F.E.C., Cumming, S.G., Haché, S., Stralberg, D., Tremblay, J.A., Barros, C., Eddy, I.M.S., Chubaty, A.M., Leblond, M., Pankratz, R.F., Mahon, C.L., Van Wilgenburg, S.L., Bayne, E.M., Schmiegelow, F., McIntire, E.J.B., 2021. Assessing Pathways of Climate Change Effects in SpaDES: An Application to Boreal Landbirds of Northwest Territories Canada. Front. Ecol. Evol. 9, 679673. https://doi.org/10.3389/fevo.2021.679673

Miller, B.W., Frid, L., 2022. A new approach for representing agent-environment feedbacks: coupled agent-based and state-and-transition simulation models. Landsc Ecol 37, 43–58. https://doi.org/10.1007/s10980-021-01282-y

MNDMNRF, 2022. Mining Lands Administration System. Ministry of Northern Development, Mines, Natural Resoruces, and Forestry, Peterborough, ON.

MNRF, 2014a. State of the Woodland Caribou Resource Report. Ministry of Natural Resources and Forestry - Species at Risk Branch, Thunder Bay, ON.

MNRF, 2014b. Integrated Range Assessment for Woodland Caribou and their Habitat - The Far North of Ontario 2013 (No. Version 1.1). Ministry of Natural Resources and Forestry - Species at Risk Branch, Thunder Bay, ON.

Moeller, A.K., Nowak, J.J., Neufeld, L., Bradley, M., Manseau, M., Wilson, P., McFarlane, S., Lukacs, P.M., Hebblewhite, M., 2021. Integrating counts, telemetry, and non-invasive DNA data to improve demographic monitoring of an endangered species. Ecosphere 12. https://doi.org/10.1002/ecs2.3443

Muff, S., Signer, J., Fieberg, J., 2020. Accounting for individual‐specific variation in habitat‐selection studies: Efficient estimation of mixed‐effects models using Bayesian or frequentist computation. J Anim Ecol 89, 80–92. https://doi.org/10.1111/1365-2656.13087

Mysterud, A., Ims, R.A., 1998. Functional responses in habitat use: availability influences relative use intrade-off situations. Ecology 79, 1435–1441.

Nagy-Reis, M., Dickie, M., Sólymos, P., Gilbert, S.L., DeMars, C.A., Serrouya, R., Boutin, S., 2020. 'WildLift': An Open-Source Tool to Guide Decisions for Wildlife Conservation. Frontiers in Ecology and Evolution 8.

Nowak, J.J., Lukacs, P.M., Hurley, M.A., Lindbloom, A.J., Robling, K.A., Gude, J.A., Robinson, H., 2018. Customized software to streamline routine analyses for wildlife management: Apps for Wildlife Management. Wildl. Soc. Bull. 42, 144–149. https://doi.org/10.1002/wsb.841

Olson, L.E., Bjornlie, N., Hanvey, G., Holbrook, J.D., Ivan, J.S., Jackson, S., Kertson, B., King, T., Lucid, M., Murray, D., Naney, R., Rohrer, J., Scully, A., Thornton, D., Walker, Z., Squires, J.R., 2021. Improved prediction of Canada lynx distribution through regional model transferability and data efficiency. Ecol. Evol. 11, 1667–1690. https://doi.org/10.1002/ece3.7157

Palm, E.C., Suitor, M.J., Joly, K., Herriges, J.D., Kelly, A.P., Hervieux, D., Russell, K.L.M., Bentzen, T.W., Larter, N.C., Hebblewhite, M., 2022. Increasing fire frequency and severity will increase habitat loss for a boreal forest indicator species. Ecological Applications 32. https://doi.org/10.1002/eap.2549

Passoni, G., Coulson, T., Ranc, N., Corradini, A., Hewison, A.J.M., Ciuti, S., Gehr, B., Heurich, M., Brieger, F., Sandfort, R., Mysterud, A., Balkenhol, N., Cagnacci, F., 2021. Roads

constrain movement across behavioural processes in a partially migratory ungulate.
Movement Ecology 9, 57. https://doi.org/10.1186/s40462-021-00292-4

Poley, L.G., Schuster, R., Smith, W., Ray, J.C., 2022. Identifying differences in roadless areas
in Canada based on global, national, and regional road datasets. Conservation Science
and Practice n/a, e12656. https://doi.org/10.1111/csp2.12656

Prokopenko, C.M., Boyce, M.S., Avgar, T., 2017. Extent-dependent habitat selection in a
migratory large herbivore: road avoidance across scales. Landscape Ecology 32, 313–
325. https://doi.org/10.1007/s10980-016-0451-1

Rempel, R.S., Carlson, M., Rodgers, A.R., Shuter, J.L., Farrell, C.E., Cairns, D., Stelfox, B.,
Hunt, L.M., Mackereth, R.W., Jackson, J.M., 2021. Modeling Cumulative Effects of
Climate and Development on Moose, Wolf, and Caribou Populations. Jour. Wild. Mgmt.
85, 1355–1376. https://doi.org/10.1002/jwmg.22094

Rempel, R.S., Hornseth, M.L., 2018. Range-specific seasonal resource selection probability
functions for 13 caribou ranges in Northern Ontario (No. IFR-01), Science and Research
Internal File Report. Ministry of Natural Resources and Forestry, Peterborough, ON.

Roche, D.G., Granados, M., Austin, C.C., Wilson, S., Mitchell, G.M., Smith, P.A., Cooke, S.J.,
Bennett, J.R., 2020. Open government data and environmental science: a federal
Canadian perspective. FACETS 5, 942–962. https://doi.org/10.1139/facets-2020-0008

Russell, D., Gunn, A., White, R., 2021. A decision support tool for assessing cumulative effects
on an Arctic migratory tundra caribou population. E&S 26, art4.
https://doi.org/10.5751/ES-12105-260104

Rutz, C., Loretto, M.-C., Bates, A.E., Davidson, S.C., Duarte, C.M., Jetz, W., Johnson, M., Kato,
A., Kays, R., Mueller, T., Primack, R.B., Ropert-Coudert, Y., Tucker, M.A., Wikelski, M.,
Cagnacci, F., 2020. COVID-19 lockdown allows researchers to quantify the effects of
human activity on wildlife. Nat Ecol Evol 4, 1156–1159. https://doi.org/10.1038/s41559-
020-1237-z

Serrouya, R., Oort, H., DeMars, C., Boutin, S., 2016. Human footprint, habitat, wolves and
boreal caribou population growth rates. Alberta Biovdiversity Monitoring Institute.

Sleep, D.J.H., Loehle, C., 2010. Validation of a Demographic Model for Woodland Caribou. The
Journal of Wildlife Management 74, 1508–1512. https://doi.org/10.1111/j.1937-
2817.2010.tb01278.x

Sorensen, T., McLoughlin, P.D., Hervieux, D., Dzus, E., Nolan, J., Wynes, B., Boutin, S., 2008.
Determining Sustainable Levels of Cumulative Effects for Boreal Caribou. Journal of
Wildlife Management 72, 900–905. https://doi.org/10.2193/2007-079

Sothe, C., Gonsamo, A., Arabian, J., Kurz, W.A., Finkelstein, S.A., Snider, J., 2022. Large Soil
Carbon Storage in Terrestrial Ecosystems of Canada. Global Biogeochemical Cycles 36.
https://doi.org/10.1029/2021GB007213

Stewart, F.E.C., Micheletti, T., Cumming, S.G., Barros, C., Chubaty, A.M., Dookie, A.E., Duclos,
I., Eddy, I., Hache, S., Hodson, J., Hughes, J., Johnson, C.A., Leblond, M.,
Schmiegelow, F.K.A., Tremblay, J.A., McIntire, E.J.B., in review. Climate-informed
forecasts reveal dramatic local habitat change and population uncertainty for threatened
caribou in Canada's north. In review Journal of Applied Ecology.

Stewart, F.E.C., Nowak, J.J., Micheletti, T., McIntire, E.J.B., Schmiegelow, F.K.A., Cumming,
S.G., 2020. Boreal Caribou Can Coexist with Natural but Not Industrial Disturbances.
The Journal of Wildlife Management 84, 1435–1444.

Superbie, C., Stewart, K.M., Regan, C.E., Johnstone, J.F., McLoughlin, P.D., 2022. Northern
boreal caribou conservation should focus on anthropogenic disturbance, not
disturbance-mediated apparent competition. Biological Conservation 265, 109426.
https://doi.org/10.1016/j.biocon.2021.109426

Teitelbaum, C.S., Sirén, A.P.K., Coffel, E., Foster, J.R., Frair, J.L., Hinton, J.W., Horton, R.M.,
Kramer, D.W., Lesk, C., Raymond, C., Wattles, D.W., Zeller, K.A., Morelli, T.L., 2021.

Habitat use as indicator of adaptive capacity to climate change. Divers Distrib ddi.13223. https://doi.org/10.1111/ddi.13223

Tootchi, A., Jost, A., Ducharne, A., 2019. Multi-source global wetland maps combining surface water imagery and groundwater constraints. Earth System Science Data 11, 189–220. https://doi.org/10.5194/essd-11-189-2019

Travers, H., Selinske, M., Nuno, A., Serban, A., Mancini, F., Barychka, T., Bush, E., Rasolofoson, R.A., Watson, J.E.M., Milner-Gulland, E.J., 2019. A manifesto for predictive conservation. Biological Conservation 237, 12–18. https://doi.org/10.1016/j.biocon.2019.05.059

Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., Fryxell, J.M., Moorter, B.V., Alberts, S.C., Ali, A.H., Allen, A.M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar, B., Belant, J.L., Bertassoni, A., Beyer, D., Bidner, L., Beest, F.M. van, Blake, S., Blaum, N., Bracis, C., Brown, D., Bruyn, P.J.N. de, Cagnacci, F., Calabrese, J.M., Camilo-Alves, C., Chamaillé-Jammes, S., Chiaradia, A., Davidson, S.C., Dennis, T., DeStefano, S., Diefenbach, D., Douglas-Hamilton, I., Fennessy, J., Fichtel, C., Fiedler, W., Fischer, C., Fischhoff, I., Fleming, C.H., Ford, A.T., Fritz, S.A., Gehr, B., Goheen, J.R., Gurarie, E., Hebblewhite, M., Heurich, M., Hewison, A.J.M., Hof, C., Hurme, E., Isbell, L.A., Janssen, R., Jeltsch, F., Kaczensky, P., Kane, A., Kappeler, P.M., Kauffman, M., Kays, R., Kimuyu, D., Koch, F., Kranstauber, B., LaPoint, S., Leimgruber, P., Linnell, J.D.C., López-López, P., Markham, A.C., Mattisson, J., Medici, E.P., Mellone, U., Merrill, E., Mourão, G. de M., Morato, R.G., Morellet, N., Morrison, T.A., Díaz-Muñoz, S.L., Mysterud, A., Nandintsetseg, D., Nathan, R., Niamir, A., Odden, J., O'Hara, R.B., Oliveira-Santos, L.G.R., Olson, K.A., Patterson, B.D., Paula, R.C. de, Pedrotti, L., Reineking, B., Rimmler, M., Rogers, T.L., Rolandsen, C.M., Rosenberry, C.S., Rubenstein, D.I., Safi, K., Saïd, S., Sapir, N., Sawyer, H., Schmidt, N.M., Selva, N., Sergiel, A., Shiilegdamba, E., Silva, J.P., Singh, N., Solberg, E.J., Spiegel, O., Strand, O., Sundaresan, S., Ullmann, W., Voigt, U., Wall, J., Wattles, D., Wikelski, M., Wilmers, C.C., Wilson, J.W., Wittemyer, G., Zięba, F., Zwijacz-Kozica, T., Mueller, T., 2018. Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. Science 359, 466–469. https://doi.org/10.1126/science.aam9712

Wang, X., Studens, K., Parisien, M.-A., Taylor, S.W., Candau, J.-N., Boulanger, Y., Flannigan, M.D., 2020. Projected changes in fire size from daily spread potential in Canada over the 21st century. Environ. Res. Lett. 15, 104048. https://doi.org/10.1088/1748-9326/aba101

Wickham, H., 2015. R Packages: Organize, test, document, and share your code. O'Reilly.

Wilson, S.F., Nudds, T.D., Vries, A., 2021. A causal modelling approach to informing woodland caribou conservation policy from observational studies. Biological Conservation 264, 109370. https://doi.org/10.1016/j.biocon.2021.109370

Wittmer, H.U., Serrouya, R., Elbroch, L.M., Marshall, A.J., 2013. Conservation Strategies for Species Affected by Apparent Competition: Conservation Biology. Conservation Biology 27, 254–260. https://doi.org/10.1111/cobi.12005

Yates, K.L., Bouchet, P.J., Caley, M.J., Mengersen, K., Randin, C.F., Parnell, S., Fielding, A.H., Bamford, A.J., Ban, S., Barbosa, A.M., Dormann, C.F., Elith, J., Embling, C.B., Ervin, G.N., Fisher, R., Gould, S., Graf, R.F., Gregr, E.J., Halpin, P.N., Heikkinen, R.K., Heinänen, S., Jones, A.R., Krishnakumar, P.K., Lauria, V., Lozano-Montes, H., Mannocci, L., Mellin, C., Mesgaran, M.B., Moreno-Amat, E., Mormede, S., Novaczek, E., Oppel, S., Ortuño Crespo, G., Peterson, A.T., Rapacciuolo, G., Roberts, J.J., Ross, R.E., Scales, K.L., Schoeman, D., Snelgrove, P., Sundblad, G., Thuiller, W., Torres, L.G., Verbruggen, H., Wang, L., Wenger, S., Whittingham, M.J., Zharikov, Y., Zurell, D., Sequeira, A.M.M., 2018. Outstanding Challenges in the Transferability of Ecological Models. Trends in Ecology & Evolution 33, 790–802. https://doi.org/10.1016/j.tree.2018.08.001

714 **Table 1.** Summary of changes in buffered anthropogenic disturbance and fire excluding

715 anthropogenic disturbance (as defined in Johnson et al. 2020) in the Base Scenario without

716 additional development, Roads Only scenario, and a Roads-and-Mines scenario that included

717 the proposed roads and mining claims within the Ring of Fire area.

Scenario	Anthropogenic (%)	Fire (%)	Total Disturbance (%)	Fire Excluding Anthropogenic (%)
Base	0.41	4.27	4.58	4.17
Roads Only	1.11	4.27	5.22	4.12
Roads-and-Mines	6.95	4.27	11.02	4.71

Figure 1. The Missisa boreal caribou range (black outline), which includes the proposed Ring of

Fire (RoF) mining claims (dark red), is the focus of this study. The blue dashed line

distinguishes the Managed Forest Zone (MFZ) from the Far North, and shading distinguishes

the Boreal Shield ecozone (pale red) from the Hudson Plains (blue). Inlay map (top-right)

indicates the location of the Ontario boreal caribou ranges (grey) relative to Canada.

Figure 2. Maps of the Missisa range with the extent of linear features included in (a) the original

model developed by Hornseth and Rempel (Base), (b) the roads only scenario used to project

the RSF model, and (c) the roads-and-mines scenario used for demographic modelling. Existing

roads are represented as green lines, proposed roads as orange lines, and mines are coloured

in purple.

Figure 3. Reproduction of the seasonal RSF for the Missisa range from Hornseth and Rempel

(2016) using caribouMetrics and the published coefficients to reproduce the relative probability

of use (0-1) by boreal caribou during spring, summer, fall, and winter. The predictor variables

used are approximations of those used by Hornseth and Rempel (2016) based on currently

available data. Scale ranges from dark blue to yellow with yellow representing a higher relative

probability of use.

Figure 4. Seasonal RSF predictions from *caribouMetrics* in the Missisa range under the roads

only scenario using the coefficients from Hornseth and Rempel (2016) from the Missisa, James

Bay, Nipigon, and Pagwachuan range to estimate the relative probability of use (0-1) by boreal

caribou during spring, summer, fall, and winter. Scale ranges from dark blue to yellow with

yellow representing a higher relative probability of use.

Figure 5. Demographic rate simulations derived from regression models in Johnson et al.

(2020) for (a) Adult female survival (S), (b) Recruitment (calves per 100 cows), and (c) Average

population rate (λ). Overlap of grey and black dotted lines in (a) and (b) indicates a good match

between expected values from our model (grey) and Johnson et al. 2020 (black). Coloured lines

show effects of changing anthropogenic disturbance on demographic rates in 35 sample

populations, assuming sample populations are randomly distributed among quantiles of the beta

distribution, and each population remains in the same quantile of the beta distribution as

disturbance changes. Alignment of these sample trajectories with 95% predictive intervals from

Johnson et al. 2020 (pale grey bands in panels a and b) indicates that we have adequately

reproduced variability in that model. Bars show the 2.5th and 97.5th percentiles for 500 sample

populations under the three disturbance scenarios for the Missisa range (from left to right, Base,

Road Only, and Roads and Mines). The black diamond indicates the demographic rates for the

Missisa range according to a 2014 assessment (MNRF, 2014a). Populations with an average

population trend of less than 0.99 are considered not self-sustaining.

767 **Supplementary Material**

768 Supplementary material is available in this OSF repository:

769 https://osf.io/r9mkp/?view_only=fb71321265d14dbeb3d932e4de66be0c

Ecozone
- Boreal Shield
- Hudson Plain
- Managed Forest Boundary
- Mining Claims

Swan

Spirit

Ozhiski

Missisa

James Bay

Berens

Kinloch

Sydney

Churchill

Nipigon

Pagwachuan

Brightsand

Kesagami

Discontinuous Distribution

Lake Superior Coast

N

0   100   200   300 km

Missisa

Spring	Summer	Fall	Winter

 |

 |

 |

 |

RSF Value

0.00
0.25
0.50
0.75
1.00

James Bay

Spring	Summer	Fall	Winter

 |

 |

 |

 |

Nipigon

Spring	Summer	Fall	Winter

 |

 |

 |

 |

Pagwachuan

Spring	Summer	Fall	Winter

 |

 |

 |

 |