Title: Interacting abiotic and biotic drivers shape woody invasions across Hawaiian forests

Running head: Alien woody invasions across Hawaii

Authors: Dylan Craven [1,2,3*], Jonathan M. Chase [2,4], Tiffany M. Knight [2,3,5]

Institutional affiliations:

1 Centro de Modelación y Monitoreo de Ecosistemas, Facultad de Ciencias, Universidad Mayor,

Santiago, Chile

2 German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig,

Germany

3 Department of Community Ecology, Helmholtz Centre for Environmental Research–UFZ, 06120 Halle

(Saale), Germany

4 Department of Computer Science, Martin Luther University, Halle-Wittenberg, 06108 Halle (Saale),

Germany

5 Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108 Halle (Saale),

Germany

Contact information:

* Telephone: +56 973887180

E-mail: dylan.craven@aya.yale.edu

Abstract

The same features that generate biodiversity patterns across and within oceanic islands over evolutionary

time – interactions between isolation, area, and heterogeneity – also influence their vulnerability to

biological invasions. Here, we identify the factors that shape the richness and abundance of woody aliens

in forest communities across the Hawaiian archipelago, and assess the relative importance of abiotic,

biotic, and anthropogenic factors and their interactions on the establishment and dominance of woody

alien species. Using a database of 460 forest plots distributed across the six major Hawaiian islands, we

examine variation in *i*) relative alien species richness and abundance as a function of multiple factors

(e.g., temperature, aridity, soil age, and the human influence index) and *ii*) establishment and dominance

of alien species as a function of abiotic and anthropogenic factors, as well as phylogenetic and trait

distinctiveness. We found that relative alien species richness and abundance were higher in areas where

temperature was high and aridity low. Gradients in temperature, aridity, soil age, and human influence

also modulated the importance of biotic factors in determining establishment of alien species. In contrast,

whether these alien species could become locally dominant was not strongly influenced by abiotic or

biotic factors, or their interactions. Our results suggest that environmental filtering mediates the strength

of biotic filtering in determining where woody aliens are able to colonize and establish on these oceanic

islands, but not whether they become dominant. The strong context dependence of multi-species

invasions highlights the complexity of developing management strategies to mitigate the biodiversity and

ecosystem impacts of biological invasions.

Keywords: biological invasions, context dependency, oceanic islands, Darwin's Naturalisation

Hypothesis, functional traits

62
63
64
65

Introduction

On just 3 to 5% of the world's area, islands harbor a disproportionate amount (15% to 20%) of its known plant diversity (Tershy *et al.*, 2015; Whittaker *et al.*, 2017). Especially on oceanic islands, this biodiversity has emerged as a result of isolation and dynamic geophysical processes that influence island area and environmental heterogeneity (Gillespie, 2016; Schluter & Pennell, 2017; Whittaker *et al.*, 2017). However, these isolated islands are also highly susceptible to invasions by alien species (Moser *et al.*, 2018), which can alter community structure, displace endemic species, and threaten the provisioning of vital ecosystem services (Vitousek *et al.*, 1987; Sax & Gaines, 2008; Vilà *et al.*, 2011; Simberloff *et al.*, 2013; Castro-Díez *et al.*, 2019). There is no sign that the number of alien species that are successfully establishing is reaching equilibrium (Seebens *et al.*, 2017), suggesting that without prevention and management, new alien species will continue to arrive and disrupt island ecosystems. It is therefore critical to deepen current understanding of the factors that allow alien species to successfully establish and come to dominate in island ecosystems (Mack *et al.*, 2000; Pyšek *et al.*, 2010; Blackburn *et al.*, 2011; Dawson *et al.*, 2017).

Across and within oceanic islands, abiotic, biogeographical, and anthropogenic factors jointly determine where biological invasions occur. Hot spots of alien plant species are typically found on islands that are isolated, large, and topographically complex, as well as on islands with high human impacts (Sax & Gaines, 2008; Denslow *et al.*, 2009; Dawson *et al.*, 2017; Moser *et al.*, 2018; Wohlwend *et al.*, 2021). Within islands, richness and abundance of alien plants are greater at lower elevations in locations with high precipitation (Ibanez *et al.*, 2019) and where human population density is higher and land use is more intense (Denslow, 2003). However, community-level patterns of biological invasions often fail to provide mechanistic insights, in part because they overlook functional and phylogenetic differences between native and alien species (biotic filtering). A growing body of literature has emerged to fill this gap, which also highlights that biological invasions are multi-faceted processes most appropriately addressed by jointly considering multiple potential mechanisms, including environmental and biotic

filtering, propagule pressure, and human impacts (Catford *et al.*, 2009; Gallien & Carboni, 2017; Pearson

et al., 2018; Bennett, 2019).

Environmental filtering structures communities by influencing the establishment of species in a

given place (Kraft *et al.*, 2015), via their traits that determine tolerance of abiotic factors (e.g. water

availability, soil pH). This environmental filtering can result in the co-occurrence of species with similar

traits. In the context of biological invasions, environmental filtering is frequently framed in terms of the

pre-adaptation hypothesis (Cadotte *et al.*, 2018; Bennett, 2019), which posits that alien species should be

sufficiently similar to native species in their traits or evolutionary history to allow them to establish viable

populations. Support for this hypothesis is particularly expected in environments with high abiotic stress

such as those with high aridity, cold temperatures, or young soils (Cornwell & Ackerly, 2009). Yet empty

niche space on islands, created by the rarity and non-randomness of colonizations (Losos & Ricklefs,

2009; Carvajal-Endara *et al.*, 2017; Taylor *et al.*, 2019), may occur even in environments with high

abiotic stress. For example, alien plant species were more different from native plant species in warm,

arid environments on the lower slopes of Mauna Loa, Hawai'i than in cooler, wet environments,

suggesting that more unfilled niche space was available in environments with greater abiotic stress (Henn

et al., 2019). Empirical evidence in support of the pre-adaptation hypothesis on island ecosystems (and in

general) is therefore quite mixed (Schaefer *et al.*, 2011; Ma *et al.*, 2016; Marx *et al.*, 2016; Cadotte *et al.*,

2018).

On the other hand, biotic filtering can promote the co-occurrence of ecologically dissimilar

species by minimizing competition (Kraft *et al.*, 2008; Kraft & Ackerly, 2010). However, these species

may still not completely fill available niche space (Ashby *et al.*, 2017). For example, dispersal limitation

might result in the absence of particular trait syndromes on oceanic islands (Taylor *et al.*, 2019; König *et*

al., 2021). This leaves niche space available for colonization by radiating lineages, such as the Hawaiian

silversword alliance (Givnish *et al.*, 2009) and the *Periploca laevigata* complex in the Canary Islands

(García-Verdugo *et al.*, 2020), or by alien species. Assuming that alien species can persist in the new

environment, Darwin's naturalisation conundrum (Darwin, 1859) predicts that alien species are more

likely to establish in native communities in which they are sufficiently distinct from native species, either

phylogenetically or functionally, to avoid competition by exploiting unused niche space (Cadotte *et al.*,

2018; Bennett, 2019). A meta-analysis using phylogenetic relatedness as a surrogate for niche differences

found that biotic filtering influences the establishment of alien species at local spatial scales in a manner

consistent with Darwin's naturalisation hypothesis (Ma et al., 2016). While phylogenetic dissimilarity is a

reliable proxy for detecting differences in how species occupy multidimensional niche space (Tucker *et*

al., 2018), differences in traits associated with competition and resource use (Kunstler *et al.,* 2012; Díaz

et al., 2016) provide a more direct means for understanding the relationship between ecological similarity

and invasion (Marx *et al.,* 2016; Carboni *et al.,* 2018; Levin *et al.,* 2020; van der Sande *et al.,* 2020).

There is a growing appreciation that biological invasions are context dependent, as they are

frequently the result of interactions among abiotic, biotic, and anthropogenic factors (Ricciardi *et al.,*

2013; Pearson *et al.,* 2018; Sapsford *et al.*, 2020; Catford *et al.*, 2022). Along abiotic gradients, native

communities are expected to be less invaded and alien plant species more similar to native communities

where abiotic stress is high, while native communities are expected to be more invaded and alien species

less similar to native communities where abiotic stress is low (Perelman *et al.*, 2007; Tecco *et al.*, 2010;

Gross *et al.,* 2013; González-Moreno *et al.*, 2014; Sapsford *et al.*, 2020). Anthropogenic disturbances (or

habitats) may also generate context dependent invasion outcomes (González-Moreno *et al.*, 2014; Jauni *et*

al., 2015) by favoring non-native species with disturbance-tolerant trait values (Richardson & Rejmánek,

2011; Ni *et al.,* 2021; Sperry *et al.,* 2021). Particularly on oceanic islands, island age (or soil age) - which

typically has been used to address the macroevolutionary processes underlying native biodiversity

patterns (Emerson, 2002; Gillespie, 2016; Whittaker *et al.*, 2017) - may further strengthen the context

dependency of biological invasions by differentially shaping the biodiversity patterns for native

(Whittaker *et al.*, 2008; Craven *et al.*, 2019) and alien species (Leihy *et al.*, 2018). Older islands with high

native biodiversity may be more resistant to invasion ("Eltonian biotic resistance"; Fridley *et al.*, 2007),

yet there is growing evidence that more diverse native plant communities can also be more vulnerable to

invasion (Stohlgren *et al.,* 2003; Qian & Sandel, 2022), possibly due to empty niche space, high habitat

heterogeneity or greater resource availability. Absent disturbances, alien species are more apt to establish

on older soils that usually have greater resource availability than on younger soils, which tend to be more

nutrient poor (Aplet *et al.,* 1998; Hughes & Denslow, 2005; Zimmerman *et al.,* 2008). This suggests that

older soils may favor stronger biotic filtering, while younger soils may act as an environmental filter of

biological invasions. However, the extent to which soil age and anthropogenic and abiotic drivers

underpin context-dependent biological invasions across oceanic islands remains uncertain.

Oceanic archipelagos, such as the Hawaiian islands, are frequently used as natural laboratories to

disentangle ecological and evolutionary drivers of native biodiversity patterns (Emerson, 2002; Gillespie,

2016; Craven *et al.,* 2019), yet also provide a compelling template for understanding biological invasions

because of their recency and severity (Barton *et al.,* 2021; Wohlwend *et al.,* 2021). Here, we examine

patterns of invasions of alien woody species, which represent approximately 60% of alien species on

oceanic islands (Kueffer *et al.,* 2010), and the relative importance of abiotic, biotic and anthropogenic

drivers and their interactions. We first identify the abiotic and anthropogenic drivers of biological

invasions (e.g., temperature, aridity, soil age, and the human influence index) in terms of alien species

richness and abundance relative to native communities. We then evaluate the relative importance of

phylogenetic distinctiveness and differences in traits of alien woody species to the native community

along environmental and anthropogenic gradients for the establishment and dominance of alien woody

species. Lastly, we test the sensitivity of the interactive effects of abiotic, biotic, and anthropogenic

factors to the presence of the widespread and abundant native woody species, *Metrosideros polymorpha*

(local name, Ohi'a). *Metrosideros polymorpha* is closely related to several alien and native woody species

(Fig. S1; Craven *et al.,* 2018) yet is functionally unique, exhibiting greater adult plant size, heavier wood

density, and lighter seeds than most native and alien woody species in Hawaiian forests. Determining if

our results depend on the presence of this one native species is particularly urgent because an emerging

soil fungal pathogen (*Ceratocystis fimbriata*), also known as rapid Ohi'a death (ROD; Keith *et al.,* 2015;

Fortini *et al.,* 2019), kills large numbers of *M. polymorpha* (Mortenson *et al.,* 2016), and may impact

future biological invasions of native forests across Hawai'i.

Methods

Plot data selection and preparation

To examine patterns of woody plant invasion of Hawaiian forests, we used a forest plot database that

compiled data from multiple sources (Zimmerman *et al.,* 2007; Ainsworth *et al.,* 2011; Gillespie Thomas

W. *et al.,* 2013; Ostertag *et al.,* 2014; US Forest Service, 2016) to include species identity and size from

across the Hawaiian archipelago (Craven et al. 2018). For our analyses, we used a total of 460 plots that

range in area from 100 to 1,018 m^2 (median plot area = 1,000 m^2) with a minimum tree size threshold of 5

cm diameter at 1.3 m height. These plots are distributed across the six major islands in the archipelago:

Hawai'i, Maui, Lana'i, Moloka'i, O'ahu, and Kaua'i (Fig. 1).

Fig. 1. Map of alien species richness in forests across the Hawaiian archipelago n= 460 plots). Color corresponds to
species richness of alien woody species rarefied to 100 stems; grey circles indicate plots that have not been invaded.
Inset shows the number of plots invaded by zero, one, and more than one alien woody species.

Species selection

Species names were standardized using The Plant List v 1.1 (www.theplantlist.org) with 'Taxonstand'
(Cayuela *et al.*, 2017) and alien status was obtained from the flora of the Hawaiian Islands (Wagner *et al.*,
2005). While there are few non-angiosperm alien woody species in Hawai'i that have naturalized (*Pinus*
caribea and *Cryptomeria japonica* (Wagner *et al.*, 2005)), their branches in a phylogeny may be longer
than those of angiosperms and may result in unusual phylogenetic measures (Qian *et al.*, 2017). Similarly,
monocot species likely have different ecological strategies than woody angiosperms (Spicer & Groover,
2010; Pierce *et al.*, 2013). We therefore excluded 14 non-angiosperm and monocot species from our
dataset. We further excluded alien woody species that have not been reported as naturalised or whose
populations are planted and have not spread widely, such as *Eucalyptus robusta* and *E. grandis* (Wagner
et al., 2005; Imada, 2012). In total, we included data for 129 woody species, of which 41 are alien and 88
are native (Table S1).

Levels of biological invasion

We calculated commonly used metrics that are proxies for the establishment and dominance of alien
species (Catford *et al.*, 2012; Leung *et al.*, 2012). At the community level, we calculated relative alien
species richness as the proportion of alien woody species to all woody species, and relative alien species
abundance as the proportion of alien woody stems to all woody stems and thus indicate the extent of
invasion. At the species level, we calculated presence/absence and relative abundances of each alien
woody species. We classified absences as true absences if a species was reported as naturalized on an
island by Wagner et al. (2005) but was not present in a plot. From here on, we refer to presence/absence
as establishment and relative abundance (at the species level) as dominance.

Phylogenetic distinctiveness of alien woody species

We used an updated version of the molecular phylogeny from Zanne et al. (Jiang & Jiang 2016; Zanne et
al. 2013) to build phylogenies of all species (native and alien) and only alien species. We conservatively

bound species to the backbone using dating information from congeners with the 'congeneric.merge'

function in the 'pez' package (Pearse *et al.,* 2015). In total, we placed all 129 woody species in the plot

database on the phylogeny (Fig. S1).

Following Thuiller et al. (2010), we calculated phylogenetic distinctiveness of alien woody

species to the community of woody native species using two metrics: (i) phylogenetic distance to the

nearest native species (DNNS) and (ii) mean weighted distance to native community (MWDNC). Both

metrics were calculated for each alien woody species in all communities, including those in which they

were not present. Because phylogenetic distinctiveness may be confounded with species richness

(Kembel, 2009), both DNNS and MWDNC are likely to be smaller in more diverse communities. To

correct for this, we calculated a standardized effect size (SES) by shuffling the tips of the phylogeny

1,000 times:

$SES = (observed - mean_{null}) / standard\ deviation_{null}$.

As DNNS is more sensitive to topological uncertainty in the phylogeny than MWDNC (Park Daniel S. *et*

al., 2020), we use MWDNC in our analyses. Lower (more negative) SES values indicate that alien woody

species are more closely related to the native plant community, while higher SES values indicate that

alien woody species are less closely related to the native plant community.

Functional traits

We selected three functional traits related to competitive ability to explore variation in establishment and

dominance of alien woody species and that are associated with key axes of ecological variation (Díaz *et*

al., 2016): wood density, adult plant size, and seed mass. Wood density is related to interspecific variation

in shade tolerance (Muller-Landau, 2004; Chave *et al.*, 2009) and growth and mortality rates (Chave *et*

al., 2009), adult plant size is associated with competition for light (King *et al.*, 2006) and seed dispersal

distance (Thomson *et al.,* 2011) variation in seed mass drives the tolerance-fecundity trade-off (Muller-

Landau Helene C., 2010). Data for wood density were obtained from Chave *et al.* (2009) and from the

BAAD, BIEN, and TRY databases (Falster *et al.*, 2015; Burrill *et al.*, 2018; Kattge *et al.*, 2020; Maitner,

2020). For seed mass, we obtained data from Shiels (2011) and the BIEN and TRY databases (Kattge *et*

al., 2020; Maitner, 2020). Because wood density and seed mass are strongly conserved phylogenetically

(Chave *et al.*, 2009; Igea *et al.*, 2017), we calculated genus-level means when species-level data were not

available. We estimated adult plant size from the plot data as the 95^{th} percentile of diameter of all

diameters ≥ 0.1 x maximum observed diameter ($D95_{0.1}$) for species with 20 or more stems or as the 95^{th}

percentile of diameter of all diameters for species with fewer than 20 stems (D95) following King et al.

(2006). Both measures are independent of abundance, strongly correlated with maximum height, and are

robust to outliers (King *et al.*, 2006).

Trait imputation

Because data were not available at the species level for all species in our data set (wood density: 5.4%

unavailable; seed mass: <1% unavailable; adult plant size: 13.2% unavailable) and these gaps did not

exceed 30% of our dataset (Penone *et al.*, 2014), we used phylogenetic trait imputation to fill gaps in our

data set. Including imputed trait data also has been shown to reduce bias, as compared to removing

species with missing trait data (Penone *et al.*, 2014). We used a random forest imputation algorithm in

combination with phylogenetic information, which we optimized for each trait to minimize imputation

error. For each trait, we assessed the imputation error for a range of phylogenetic eigenvectors (1 - 30)

and used the imputed trait values for the number of phylogenetic eigenvectors that minimized imputation

error. Prior to imputation, we log-transformed seed mass and adult plant size and back-transformed

imputed values and imputation errors. We calculated imputation errors as the percent normalized root

mean squared error, which were 1.95% for wood density, 13.6% for adult plant size, and 23.8% for seed

mass. Our gap-filled trait data had similar distributions and mean values as the observed trait data (Fig.

S2).

Trait differences

To measure trait differences among native and alien species, we calculated differences between trait

values of alien species and the abundance-weighted trait values of native species for each trait (Fig. S3;

van der Sande *et al.,* 2020). We standardized trait differences by calculating the log-response ratio of trait

values of alien species to native species. Positive trait differences indicate that alien species had higher

trait values than the native plant community and negative trait differences indicate the opposite.

Abiotic conditions and anthropogenic impacts

To understand the role of abiotic conditions in shaping woody invasions, we extracted mean annual

temperature (MAT), precipitation (MAP) and potential evapotranspiration (PET; calculated using the

Penman-Monteith equation) for each plot from locally interpolated climate data (Giambelluca *et al.,*

2013, 2014). We then calculated aridity, a widely used measure to assess the impacts of water availability

on plant communities (e.g., Berdugo *et al.,* 2020), as the ratio of MAP to PET (Zomer *et al.,* 2008). We

multiplied aridity by -1, such that low values of this index correspond to low aridity and high values to

high aridity. We also extracted elevation (Jarvis *et al.,* 2008) and soil substrate age for each plot (hereon

referred to as soil age; Sherrod *et al.,* 2007). On the Hawaiian archipelago, soil age is strongly related to

the timing of volcanic activity over a volcanic hotspot, such that the western-most large islands (Kauai

and O'ahu) are several millions of years older than the younger islands of Maui and Hawaii (Clague &

Sherrod, 2014).

Because the establishment of alien woody species is facilitated by roads, and large populations of

alien species can be maintained in areas where anthropogenic disturbance is high, we also included an

index of human impacts in our analyses. Specifically, we extracted the human influence index (HII), a

composite of human population density, land use, and infrastructure (HII ; Wildlife Conservation Society

- WCS & Center for International Earth Science Information Network - CIESIN - Columbia University,

2005). .

Data analysis

Prior to model fitting, we tested for collinearity among explanatory variables using pairwise Pearson

correlations. We selected variables that had a pairwise correlation below 0.7 (Dormann *et al.*, 2013).

Based on this criteria, we retained mean annual temperature (MAT), aridity, soil age, and human

influence index (HII) in our models examining relative alien species richness and abundance and MAT,

aridity, soil age, HII, phylogenetic distinctiveness (in terms of mean weighted distance to native

community (MWDNC)) and trait differences for wood density, adult plant size, and seed mass in our

models examining variation in establishment and dominance.

We examined the relative importance of aridity, MAT, HII, and soil age, in determining variation

in relative alien species richness and abundance using hierarchical Bayesian models with Beta and log-

normal distributions, respectively. Relative alien species richness and abundance were modeled as the

proportion of alien woody species or individuals in a plot. In both models we used random intercept terms

for island and plot size.

Because phylogenetically closely related species are likely to share similar trait values (e.g.,

Felsenstein, 1985), not accounting for phylogenetic relationships may reduce estimation accuracy and

increase type I error rates (Li & Ives, 2017). We therefore fit phylogenetic hierarchical Bayesian models

to examine variation in establishment and dominance of alien woody species as a function of abiotic,

biogeographical, and anthropogenic factors (aridity, MAT, HII, soil age), phylogenetic distinctiveness,

competitive trait differences, and two-way interactions of phylogenetic distinctiveness or trait differences

with abiotic conditions. We fit the model for probability of occurrence (i.e. establishment) using a

Bernoulli distribution and the model for the relative abundance of alien woody species using a lognormal

distribution, which fit the data better than a Beta distribution. For both models, we used a crossed random

effects structure with random intercept terms for island, species and plot, as not all alien species occurred

in each plot or island. In the random intercept term for plot, we nested plot identity within a group factor

of plot area to account for the increased probability of occurrence in larger plots. To account for

phylogenetic correlations among species, we included two random intercept terms for species; one term

models phylogenetic covariance and another term accounts for repeated measurements and other effects

that may be independent of phylogenetic relationships among species. We first fit a full model with all

two-way interactions and then simplified it by removing two-way interactions whose 95% credible

intervals overlapped with zero.

We fit all models using weakly informative priors, four chains, and 2,500 burn-in samples per

chain, after which 2,000 samples per chain were used to calculate posterior distributions of model

parameters (total post-warmup samples= 8,000). Priors were flat for population-level effects. We

increased the '*adapt_delta*' parameter in the 'brm' function until there were no divergent transitions.

Model convergence was evaluated visually with trace plots and by estimating '*Rhat*' using the 'rhat'

function, where values greater than 1.01 indicate that models have failed to converge (Tables S1 and S2).

Model fit was assessed visually by comparing observed data to simulated data from the posterior

predictive distribution (Fig. S4 and S5). Additionally, we estimated a Bayesian r^2 for each model to

represent an estimate of the proportion of variation explained for new data (Tables S1 and S2). Model fit

for establishment was also assessed by calculating ROC and AUC (Fig. S6). We evaluated spatial

autocorrelation of model residuals using Moran's I. All models showed minimal evidence of spatial

autocorrelation (Figs. S7 and S8). We also tested all models for scale dependence by evaluating the

precision and accuracy of model estimates per plot size (Spake *et al.*, 2021); random effect estimates and

their accuracy and precision did not vary systematically with plot size (Fig. S9 and S10). We also tested

the sensitivity of our results to the presence of the widespread and abundant native woody species, *M.*

polymorpha, by fitting the same set of models described above, but after removing *M. polymorpha* from

our data set and phylogeny. All models were fitted using 'brms' (Bürkner, 2017).

For all analyses, environmental and functional trait data were visually inspected for normality;

soil age was natural-log transformed. All variables were standardized using a *z*-transformation to enable

comparisons across models. Analyses were performed using R v. 3.6.1 (R Core Team, 2020).

Results

Woody invasions were widespread across Hawaiian forests on all six islands (Fig. 1). Approximately 36% of forest plots were invaded, 19.6% being invaded by multiple alien woody species and 16.7% being invaded by one alien woody species. Across all plots, the most frequent woody invaders were *Psidium cattleianum, Cecropia obtusifolia, Psidium guajava*, and *Melastoma malabathricum* (Fig. S1).

Abiotic and anthropogenic drivers of woody invasions

Relative alien species richness and abundance were moderately positively correlated ($r = 0.39$, P < 0.001). At the community level, invasion of forests by alien woody species varied with abiotic conditions (Fig. 2). Relative species richness and abundance of alien woody species increased with MAT, and relative abundance of alien woody species decreased with increasing aridity. Relative species richness and abundance of alien woody species also showed a tendency to increase with HII, as 80% credible intervals did not overlap with zero. We found no evidence for an effect of soil age on the richness or abundance of alien woody species.

Fig. 2. Coefficient estimates of main effects used to examine **a)** relative alien woody species richness and **b)** relative alien woody species abundance across the Hawaiian archipelago. MAT is mean annual temperature, aridity is the ratio of mean annual potential evapotranspiration (PET) to mean annual precipitation, soil age is the estimated soil age, and HII is the human influence index. Medians, 80%, and 95% credible intervals of model coefficients were estimated using a phylogenetic hierarchical Bayesian with a beta distribution.

Establishment of alien woody species

Our analysis showed strong interactive effects among biotic and abiotic factors on the establishment of

alien woody species across Hawaiian forests. Establishment of alien woody species increased with

increasing temperature, soil age, and Δ adult plant size (Fig. 4). Yet, all biotic factors exhibited

differential responses along abiotic and anthropogenic gradients (Fig. 5). At higher temperatures, alien

woody species more closely related to the native plant community were more likely to occur than those

distantly related to the native plant community (Fig. 5a). Likewise, alien woody species with lighter seeds

and smaller adult plant size than the native plant community were more likely to establish at higher

temperatures (Fig. 5b and c). In areas with high aridity, alien woody species distantly related to the native

plant community were more likely to occur than less distantly related ones, whose probability of

establishment increased as aridity decreased (Fig. 5d). Alien woody species with heavier seeds and that

are taller than the native plant community were more likely to occur where aridity was high, a pattern that

switched sharply where aridity was low (Fig. 5e and f). Alien woody species with larger seeds and denser

wood than the native plant community were more likely to occur on older soils than on younger soils

(Fig. 5h and i). In areas with low human impacts, alien woody species with denser wood than the native

plant community were more likely to establish than those with lighter wood than the native plant

community (Fig. 5g). In contrast, alien woody species with lighter wood than the native plant community

were more likely to establish in areas with higher human impacts (Fig. 5g).

365
Fig. 3. Coefficient estimates of main effects used to evaluate the establishment of alien woody species across the
Hawaiian archipelago. Aridity is the ratio of mean annual potential evapotranspiration (PET) to mean annual
precipitation, MAT is mean annual temperature, soil age is the estimated soil age, HII is the human influence index,
MWDNC is mean weighted phylogenetic distance to the native tree community, and ΔWD, Δ seed mass, and Δ
adult plant size are the standardised differences in wood density, seed mass, and adult plant size, respectively,
between alien woody species and the native tree community species. Medians, 80%, and 95% credible intervals of
model coefficients were estimated using a phylogenetic hierarchical Bayesian model with a Bernoulli distribution.
373

374
Fig. 4. The estimated effects of phylogenetic distinctiveness (MDWNC) and trait differences among alien and native
species in terms of seed mass, adult plant size, and wood density along gradients of (**a,b,c**) mean annual temperature
(MAT), (**d, e, f**), aridity, (**g**) human influence index (HII), and (**h & i**) soil age on the probability of establishment of
alien woody species across Hawaiian forests. MDWNC is the standardized mean weighted phylogenetic distance
between each alien species and the native community; positive values indicate that alien species are phylogenetically
distinct from the native plant community and negative values indicate that alien species are phylogenetically similar
to the native plant community. Seed mass, adult plant size, and wood density are standardized trait differences
between alien woody species and the native community; positive values indicate that trait values of alien species are
larger than those of the native community and negative values indicate that trait values of the alien woody species
are smaller than those of the native community. Black and colored lines are marginal predictions evaluated at the
50th and 5th to the 95th quantiles of the moderator, respectively, and estimated with a phylogenetic hierarchical
Bayesian model with a Bernoulli distribution. All x axes are scaled to unit variance.

Dominance of alien woody species

Abiotic and biotic factors and their interactions did not have marked impacts on dominance patterns of

alien woody species (Fig. 6). Only phylogenetic distinctiveness had a moderate, negative effect on the

relative abundance of alien woody species (80% credible intervals did not overlap zero), and no

interactions between abiotic and biotic factors were included in the final model as the 95% credible

intervals of all two-way interactions overlapped with zero.

Fig. 5. Coefficient estimates of main effects used to examine dominance of alien woody species across Hawaiian
forests. Aridity is the ratio of mean annual potential evapotranspiration (PET) to mean annual precipitation, MAT is
mean annual temperature, soil age is the estimated soil age, HII is the human influence index, c is mean weighted
phylogenetic distance to the native tree community, and ΔWD, Δ seed mass, and Δ adult plant size are the
standardised differences in wood density, seed mass, and adult plant size, respectively, between alien woody species
and the native tree community species. Medians and 80% and 95% credible intervals of model coefficients were
estimated using a phylogenetic hierarchical Bayesian model with a log-normal distribution.
401

Sensitivity of context-dependent drivers of establishment and dominance

Further analyses provided added insight to the role played by *M. polymorpha* in mediating the context

dependency of drivers affecting the establishment and dominance of alien woody species. When we

removed *M. polymorpha* from the dataset, the influence of many interactive effects identified in the main

analysis on the establishment of alien woody species was reduced considerably (Fig. S9). Indeed, only the

interactions between soil age and Δ seed mass, MAT and Δ seed mass, and HII and Δ wood density

retained their influence on the probability of establishment of alien woody species. Independent of the

presence of *M. polymorpha,* temperature, soil age, and Δ adult plant size exhibited strong, positive effects

on the probability of establishment of alien woody species, while Δ seed mass had strong, negative effects

on the probability of establishment of alien woody species.

In contrast, the removal of *M. polymorpha* from the data set resulted in minimal changes to the

impacts of abiotic and biotic drivers on the dominance of alien woody species (Fig. S10). The exception

to this was Δ seed mass, which had a strong, positive effect on dominance. As found for the analysis of

the full data set, no interactive effects were detected and phylogenetic distinctiveness had a marginally

negative effect on the dominance of alien woody species.

Discussion

Our results highlight the context dependence of biological invasions (e.g., Catford et al. 2021) in forests

across the Hawaiian archipelago. Abiotic factors determine the richness and abundance of alien species in

communities in the expected direction, with more woody aliens in hotter and wetter locations.

Interactions among biotic, abiotic, and anthropogenic factors jointly determine where alien woody species

establish, but not where they become dominant. We found that alien woody species establish more readily

in areas with high aridity and in older soils, which they most likely colonize successfully by exploiting

niche space not occupied by native woody species. In areas with harsh abiotic conditions, such as those

with younger soils, environmental filtering was usually - but not always - the likely dominant ecological

mechanism operating on the establishment of alien woody species. Our results further reveal how the

impacts of soil age, a biogeographical factor often used to examine macroevolutionary processes that

generate native biodiversity across and within oceanic islands (Emerson, 2002; Whittaker *et al.*, 2008;

Gillespie, 2016), extend to shaping local coexistence of alien species. Together, our results suggest that

the ecological mechanisms underlying both the pre-adaptation hypothesis (environmental filtering) and

Darwin's naturalisation conundrum (biotic filtering) are likely to interactively shape the assembly of alien

woody communities across the Hawaiian archipelago.

Abiotic drivers of multi-species woody invasions

We found that local communities are generally more invaded in forests with higher temperatures and low

aridity. These findings indicate that biological invasions are more severe in lowland, moist tropical forests

across the Hawaiian archipelago (Ibanez *et al.*, 2019), which have been heavily impacted by agriculture

and land-use change (Jacobi *et al.*, 2017), as temperature is strongly negatively correlated with elevation

($r = - 0.99$). While our results show certain similarities with macroecological patterns of alien species

richness and composition on oceanic islands in terms of abiotic factors, we detected weaker human

impacts on the severity of biological invasions at local scales than have been reported at larger spatial

scales (Moser *et al.*, 2018; Wohlwend *et al.*, 2021). Yet, the low explanatory power of our models for

relative alien species richness ($R^2 = 0.25$, 95% credible interval: 0.15-0.33) and abundance ($R^2 = 0.12$,

95% credible interval: 0.06-0.20) highlights the need to include additional drivers, such as biotic

interactions, historical legacies, and propagule pressure, to enhance the robustness of predictions of the

severity of biological invasions at local spatial scales.

The weak, positive correlation between the two metrics of invasion impacts suggests that while in

general there are more alien species when there are more alien stems, there is also a range of invaded

forests across Hawai'i, including those that are heavily invaded by a small number of dominant alien

woody species such as *P. cattleianum* and those dominated by native woody species, but contain several

alien woody species. This finding also suggests that invasion by one species may not facilitate invasion

by other alien species within the same trophic group (e.g., Zarnetske *et al.*, 2013), possibly due to niche

preemption or niche modification (Fukami, 2015). As such priority effects are often contingent upon

abiotic factors (Grainger *et al.,* 2019), it is possible that high temperatures facilitate coexistence among

alien woody species, while low aridity may strengthen priority effects (Fig. 2). Further studies would be

needed to determine whether and how priority effects of dominant alien species such as *P. cattleianum*

are influenced by their demographic traits (Fukami *et al.,* 2016) to better understand the mechanisms that

underpin multi-species biological invasions.

Context dependency of alien species establishment and dominance

Our analysis suggests that the establishment of alien woody species was strongly context dependent

(sensu Catford *et al.,* 2022), meaning that the impacts of phylogenetic and trait differences between alien

and native species on establishment were not consistent across Hawaiian forests (Tecco *et al.,* 2010;

Gross *et al.,* 2013; González-Moreno *et al.,* 2014; Sapsford *et al.,* 2020). The strength of biotic filtering is

expected to shift along environmental gradients (e.g., Cornwell & Ackerly, 2009; Kraft & Ackerly, 2010),

with the expectation that biotic filtering should be stronger in more favorable environments compared to

harsher environments (Bertness & Callaway, 1994). In contrast, we found that environmental filtering

strengthened with increasing temperature (for MWDNC, Δ seed mass, and Δ adult plant size), meaning

that alien woody species that are more similar to native woody species are more likely to establish where

environmental filters are weaker. In line with expectations, we found that environmental filtering

strengthened with increasing human impacts (for Δ wood density) and decreasing soil age (for Δ seed

mass and Δ wood density); alien woody species that are more similar to native woody species were more

likely to establish in areas with stronger human impacts or with younger soils. Yet, we also found

evidence that biotic filtering could be stronger in harsh environments, particularly where aridity was high

(for MWDNC, Δ adult plant size, and Δ seed mass), which is consistent with results reported by Henn et

al. (2019). This finding suggests that native woody communities do not fully occupy niche space in harsh

environments, thereby facilitating the establishment of phylogenetically and functionally distinct alien

woody species. It could also suggest that alien woody species may exhibit greater tolerance to water- or

nutrient-limited conditions than phylogenetically similar native woody species, which could be facilitated

by greater intraspecific trait variation (Westerband *et al.*, 2021). Together, our results reveal that alien

woody species are likely to be able to colonize native forests by competing less strongly with native

woody species (Levin *et al.*, 2020), either because conditions are too harsh (i.e. where environmental

filtering is strong) or empty niche space is available in places where biotic filtering is strong.

Our analysis revealed that soil age is an important underlying factor driving context dependent

biological invasions across Hawaiian forests. Despite its importance for native biodiversity patterns,

empirical evidence showing that soil age shapes the spatial distribution of alien species within and across

islands has been limited (Aplet *et al.*, 1998; Hughes & Denslow, 2005), particularly at macroecological

scales where greater emphasis is usually given to anthropogenic drivers (e.g., Dawson *et al.*, 2017; Moser

et al., 2018; Wohlwend *et al.*, 2021). One explanation for the positive effects of soil age on the

probability of establishment of alien woody species observed in this study is that woody communities

with high native plant diversity on older soils are less resistant to invasion ("the rich get richer";

Stohlgren *et al.*, 2003; Fridley *et al.*, 2007). However, once accounting for soil age, native and alien

species richness exhibited a significantly negative correlation (Pearson's partial correlation = -0.11, *P*

=0.01), indicating that more diverse woody communities are moderately more resistant to invasion than

less diverse plant communities. This suggests that it not soil age per se, but rather the macroevolutionary

processes associated with it, such as niche partitioning, that have generated local scale patterns of native

biodiversity over millions of years (Craven *et al.*, 2019), and that this may confer resistance to biological

invasions (Fridley *et al.*, 2007). Our results are also consistent with the idea that soil age directly affects

biological invasions by filtering alien woody species on younger soils that have low soil nutrient content

(Aplet *et al.*, 1998). Moreover, our results suggest that alien woody species successfully establish in areas

with older soils by occupying different regeneration niches or acquiring soil nutrients differently than

native woody species (Fig. 5h and i), strategies that may covary with either wood density or seed mass

(but see Carmona *et al.*, 2021). More broadly, the importance of gradients in soil age mediating biological

invasions reveals that biogeographical factors may play an underappreciated role in shaping seemingly

idiosyncratic patterns of biological invasions within and across oceanic islands.

We found little evidence that abiotic or biotic filtering consistently influenced the dominance of

alien woody species across Hawaiian forests. Our results indicate that competitive outcomes between

alien and native woody species are largely independent of environmental conditions or functional or

phylogenetic differences. Thus, spatial variation in dominance of the alien woody community could be

driven principally by dispersal, particularly if propagule pressure is high (Lockwood *et al.*, 2005;

Richardson & Pyšek, 2012; Cassey *et al.*, 2018). We detected a moderate negative effect of phylogenetic

distinctiveness on dominance (80% confidence intervals do not overlap with zero), which is consistent

with the idea that environmental filtering is the dominant process structuring invaded forest ecosystems

(Lemoine *et al.*, 2015). Nevertheless, our ability to detect the influence of abiotic and biotic filtering on

dominance may have been limited by the spatial grain of our analysis (median plot area = 1,000 m^2; range

of plot area: 100 to 1,018 m^2), which may be too large to capture the effects of neighborhood competition

or crowding (e.g., Uriarte *et al.*, 2010). Further, we used phylogenetic distinctiveness and trait differences

as proxies for competitive ability, which is rarely quantified directly (but see Levin *et al.*, 2020), and may

explain further variation in dominance among alien woody species.

Presence of M. polymorpha
We found that the patterns of alien establishment critically depend on the presence of *M. polymorpha.*

When we excluded *M. polymorpha* from the analysis, the factors that shape the establishment of alien

woody species were markedly less context dependent; only three interactions (soil age and Δ seed mass,

MAT and Δ adult plant size, and HII and Δ wood density) were retained. This result suggests that where

temperature and human impacts are high, alien woody species that are phylogenetically and functionally

similar to native species are more likely to establish; this is likely driven by the pervasiveness of *P.*

cattleianum across Hawaiian forests and its functional similarity to *M. polymorpha* (Fig. S11, Table S6).

Also, it suggests that independent of the presence of *M. polymorpha*, soil age mediates the strength of

biotic filtering. Thus, the apparent context dependency of biological invasions appears to have been

driven at least in part by the broad environmental gradients across which *M. polymorpha* occurs, whose

light seeds allow it to disperse widely and high wood density confers greater drought tolerance (Cordell *et*

al., 1998), and functional similarities between alien woody species and *M. polymorpha*.

Our sensitivity analysis indicates that unoccupied niche space persists even in the absence of *M.*

polymorpha, as evidenced by retained interactions (soil age *x* Δ seed mass, MAT *x* and Δ adult plant size,

and HII *x* Δ wood density). These results suggest that being taller, having lighter seeds or denser wood

than the native community in areas with older soils or lower human impacts may confer a competitive

advantage to alien woody species invading forests, regardless of whether they are dominated by *M.*

polymorpha. Consequently, the growing impacts of the fungal pathogen that causes Rapid O'hia Death on

M. polymorpha could shift alien woody invasions in Hawaiian forests (Gómez-Aparicio *et al.*, 2012;

Anderegg *et al.*, 2015), potentially favoring alien woody species with lighter seeds and taller stature than

the native woody community, such as *Aleurites moluccanus, Melia azedarach,* and *Syzygium cumini.*

Conclusions

Our results highlight that alien woody species likely establish in island ecosystems by occupying empty

niche space and/or habitats that are too harsh to compete directly with native woody species. While these

alien species are clearly changing the composition and dynamics of Hawaiian forests (Craven *et al.*, 2019;

Barton *et al.*, 2021), our results show that biological invasions are shaped in part by the same

environmental and biogeographical factors, such as climate and soil age, that generate and maintain

patterns of native biodiversity. Our sensitivity analysis provides evidence that the context dependency of

alien woody invasions across Hawai'i arises due to ecological interactions between abiotic,

anthropogenic, and biotic factors (Catford *et al.*, 2022), particularly along gradients in soil age, human

impacts, and temperature. While this context dependency makes our results difficult to generalize to other

archipelagos, our results emphasize that future efforts should account for how soil age mediates biotic

filtering. This strong context dependence of multi-species invasions also highlights the complexity of

developing management strategies to mitigate the biodiversity and ecosystem impacts of biological

invasions.

Acknowledgments

DC, TMK and JMC acknowledge funding by German Centre for Integrative Biodiversity Research (iDiv)

Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118). DC also received funding

from the Agencia Nacional de Investigación y Desarrollo (Chile; FONDECYT Regular No. 1201347).

TMK also acknowledges support from the Alexander von Humboldt Foundation, and the National

Geographic Society. We thank Valentin Stephan for helping to prepare the environmental data, Nathaly

Guerrero-Ramírez, Patrick Weigelt, Gunnar Keppel, Thomas Ibanez, and Holger Kreft for their input to

conceptual development, and to data providers for collecting the field data. We also thank Rebecca Spake

for her feedback on data analysis. The authors certify that they do not have any conflict of interest.

Data availability statement

The code and data that support the findings of this study are available in Github:

https://github.com/dylancraven/Hawaii_BiologicalInvasions. These data were derived from the following

resources available in Dryad: https://doi.org/10.5061/dryad.1kk02qr.

References

Ainsworth, A., Berkowitz, P, Jacobi, J.D., Loh, R.K. & Kozar, K. (2011) Focal Terrestrial Plant
571   Communities Monitoring Protocol. *Focal Terrestrial Plant Communities Monitoring Pacific*
572   *Island Network NPS/PACN/NRR–2011/410 National Park Service, Fort Collins, Colorado*.
573   Anderegg, W.R.L., Hicke, J.A., Fisher, R.A., Allen, C.D., Aukema, J., Bentz, B., Hood, S., Lichstein,
574   J.W., Macalady, A.K., McDowell, N., Pan, Y., Raffa, K., Sala, A., Shaw, J.D., Stephenson, N.L.,
575   Tague, C. & Zeppel, M. (2015) Tree mortality from drought, insects, and their interactions in a
576   changing climate. *New Phytologist*, **208**, 674–683.
577   Aplet, G.H., Hughes, R.F. & Vitousek, P.M. (1998) Ecosystem development on Hawaiian lava flows:
578   biomass and species composition. *Journal of Vegetation Science*, **9**, 17–26.
579   Ashby, B., Watkins, E., Lourenço, J., Gupta, S. & Foster, K.R. (2017) Competing species leave many
580   potential niches unfilled. *Nature Ecology & Evolution*, **1**, 1495–1501.
581   Barton, K.E., Westerband, A., Ostertag, R., Stacy, E., Winter, K., Drake, D.R., Fortini, L.B., Litton, C.M.,
582   Cordell, S., Krushelnycky, P., Kawelo, K., Feliciano, K., Bennett, G. & Knight, T. (2021)
583   Hawai'i Forest Review: Synthesizing the Ecology, Evolution, and Conservation of a Model

System. *Perspectives in Plant Ecology, Evolution and Systematics*, 125631.

Bennett, J.A. (2019) Similarities between invaders and native species: Moving past Darwin's naturalization conundrum. *Journal of Vegetation Science*, **30**, 1027–1034.

Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J.J., Gross, N., Saiz, H., Maire, V., Lehman, A., Rillig, M.C., Solé, R.V. & Maestre, F.T. (2020) Global ecosystem thresholds driven by aridity. *Science*, **367**, 787–790.

Bertness, M.D. & Callaway, R. (1994) Positive interactions in communities. *Trends in Ecology & Evolution*, **9**, 191–193.

Blackburn, T.M., Pyšek, P., Bacher, S., Carlton, J.T., Duncan, R.P., Jarošík, V., Wilson, J.R.U. & Richardson, D.M. (2011) A proposed unified framework for biological invasions. *Trends in Ecology & Evolution*, **26**, 333–339.

Bürkner, P.-C. (2017) brms: An R package for Bayesian multilevel models Using Stan. *Journal of Statistical Software*, **80**, 1–28.

Burrill, E.A., Wilson, A.M., Turner, J.A., Pugh, S.A., Menlove, J., Christiansen, G., Conkling, B.L. & David, W. (2018) *The Forest Inventory and Analysis Database: database description and user guide version 8.0 for Phase 2*, U.S. Department of Agriculture, Forest Service.

Cadotte, M.W., Campbell, S.E., Li, S., Sodhi, D.S. & Mandrak, N.E. (2018) Preadaptation and Naturalization of Nonnative Species: Darwin's Two Fundamental Insights into Species Invasion. *Annual Review of Plant Biology*, **69**, 661–684.

Carboni, M., Calderon-Sanou, I., Pollock, L., Violle, C. & Thuiller, W. (2018) Functional traits modulate the response of alien plants along abiotic and biotic gradients. *Global Ecology and Biogeography*, **27**, 1173–1185.

Carmona, C.P., Bueno, C.G., Toussaint, A., Träger, S., Díaz, S., Moora, M., Munson, A.D., Pärtel, M., Zobel, M. & Tamme, R. (2021) Fine-root traits in the global spectrum of plant form and function. *Nature*, **597**, 683–687.

Carvajal-Endara, S., Hendry, A.P., Emery, N.C. & Davies, T.J. (2017) Habitat filtering not dispersal limitation shapes oceanic island floras: species assembly of the Galápagos archipelago. *Ecology Letters*, **20**, 495–504.

Cassey, P., Delean, S., Lockwood, J.L., Sadowski, J. & Blackburn, T.M. (2018) Dissecting the null model for biological invasions: A meta-analysis of the propagule pressure effect. *PLOS Biology*, **16**, 1–16.

Castro-Díez, P., Vaz, A.S., Silva, J.S., Loo, M. van, Alonso, Á., Aponte, C., Bayón, Á., Bellingham, P.J., Chiuffo, M.C., DiManno, N., Julian, K., Kandert, S., Porta, N.L., Marchante, H., Maule, H.G., Mayfield, M.M., Metcalfe, D., Monteverdi, M.C., Núñez, M.A., Ostertag, R., Parker, I.M., Peltzer, D.A., Potgieter, L.J., Raymundo, M., Rayome, D., Reisman-Berman, O., Richardson, D.M., Roos, R.E., Saldaña, A., Shackleton, R.T., Torres, A., Trudgen, M., Urban, J., Vicente, J.R., Vilà, M., Ylioja, T., Zenni, R.D. & Godoy, O. (2019) Global effects of non-native tree species on multiple ecosystem services. *Biological Reviews*, **94**, 1477–1501.

Catford, J.A., Jansson, R. & Nilsson, C. (2009) Reducing redundancy in invasion ecology by integrating hypotheses into a single theoretical framework. *Diversity and Distributions*, **15**, 22–40.

Catford, J.A., Vesk, P.A., Richardson, D.M. & Pyšek, P. (2012) Quantifying levels of biological invasion: towards the objective classification of invaded and invasible ecosystems. *Global Change Biology*, **18**, 44–62.

Catford, J.A., Wilson, J.R.U., Pyšek, P., Hulme, P.E. & Duncan, R.P. (2022) Addressing context dependence in ecology. *Trends in Ecology & Evolution*, **37**, 158–170.

Cayuela, L., Stein, A. & Oksanen, J. (2017) Taxonstand: taxonomic standardization of plant species names. *R package version 2*.

Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009) Towards a worldwide wood economics spectrum. *Ecol Lett*, **12**, 351–66.

Clague, D.A. & Sherrod, D.R. (2014) *Growth and degradation of Hawaiian volcanoes*. *Characteristics of Hawaiian Volcanoes*, pp. 97–146. US Geological Survey Reston, VA.

Cordell, S., Goldstein, G., Mueller-Dombois, D., Webb, D. & Vitousek, P.M. (1998) Physiological and
morphological variation in Metrosideros polymorpha, a dominant Hawaiian tree species, along an
altitudinal gradient: the role of phenotypic plasticity. *Oecologia*, **113**, 188–196.
Cornwell, W.K. & Ackerly, D.D. (2009) Community assembly and shifts in plant trait distributions across
an environmental gradient in coastal California. *Ecological Monographs*, **79**, 109–126.
Craven, D., Knight, T.M., Barton, K.E., Bialic-Murphy, L. & Chase, J.M. (2019) Dissecting
macroecological and macroevolutionary patterns of forest biodiversity across the Hawaiian
archipelago. *Proceedings of the National Academy of Sciences*, **116**, 16436–16441.
Craven, D., Knight, T.M., Barton, K.E., Bialic-Murphy, L., Cordell, S., Giardina, C.P., Gillespie, T.W.,
Ostertag, R., Sack, L. & Chase, J.M. (2018) OpenNahele: the open Hawaiian forest plot database.
Biodiversity Data Journal, **6**, e28406.
Darwin, C. (1859) *On the origins of species by means of natural selection*, John Murray, United
Kingdom.
Dawson, W., Moser, D., van Kleunen, M., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Winter, M.,
Lenzner, B., Blackburn, T.M., Dyer, E.E., Cassey, P., Scrivens, S.L., Economo, E.P., Guénard,
B., Capinha, C., Seebens, H., García-Díaz, P., Nentwig, W., García-Berthou, E., Casal, C.,
Mandrak, N.E., Fuller, P., Meyer, C. & Essl, F. (2017) Global hotspots and correlates of alien
species richness across taxonomic groups. *Nature Ecology & Evolution*, **1**, 0186.
Denslow, J.S. (2003) Weeds in Paradise: Thoughts on the Invasibility of Tropical Islands. *Annals of the
Missouri Botanical Garden*, **90**, 119–127.
Denslow, J.S., Space, J.C. & Thomas, P.A. (2009) Invasive Exotic Plants in the Tropical Pacific Islands:
Patterns of Diversity. *Biotropica*, **41**, 162–170.
Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C.,
Colin Prentice, I., Garnier, E., Bönisch, G., Westoby, M., Poorter, H., Reich, P.B., Moles, A.T.,
Dickie, J., Gillison, A.N., Zanne, A.E., Chave, J., Joseph Wright, S., Sheremet'ev, S.N., Jactel,
H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J.S.,
Günther, A., Falczuk, V., Rüger, N., Mahecha, M.D. & Gorné, L.D. (2016) The global spectrum
of plant form and function. *Nature*, **529**, 167–171.
Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B.,
Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking, B.,
Schröder, B., Skidmore, A.K., Zurell, D. & Lautenbach, S. (2013) Collinearity: a review of
methods to deal with it and a simulation study evaluating their performance. *Ecography*, **36**, 27–
46.
Emerson, B. (2002) Evolution on oceanic islands: molecular phylogenetic approaches to understanding
pattern and process. *Molecular ecology*, **11**, 951–966.
Falster, D.S., Duursma, R.A., Ishihara, M.I., Barneche, D.R., FitzJohn, R.G., Vårhammar, A., Aiba, M.,
Ando, M., Anten, N., Aspinwall, M.J., Baltzer, J.L., Baraloto, C., Battaglia, M., Battles, J.J.,
Bond-Lamberty, B., Breugel, M. van, Camac, J., Claveau, Y., Coll, L., Dannoura, M.,
Delagrange, S., Domec, J.-C., Fatemi, F., Feng, W., Gargaglione, V., Goto, Y., Hagihara, A.,
Hall, J.S., Hamilton, S., Harja, D., Hiura, T., Holdaway, R., Hutley, L.S., Ichie, T., Jokela, E.J.,
Kantola, A., Kelly, J.W.G., Kenzo, T., King, D., Kloeppel, B.D., Kohyama, T., Komiyama, A.,
Laclau, J.-P., Lusk, C.H., Maguire, D.A., Maire, G. le, Mäkelä, A., Markesteijn, L., Marshall, J.,
McCulloh, K., Miyata, I., Mokany, K., Mori, S., Myster, R.W., Nagano, M., Naidu, S.L.,
Nouvellon, Y., O'Grady, A.P., O'Hara, K.L., Ohtsuka, T., Osada, N., Osunkoya, O.O., Peri, P.L.,
Petritan, A.M., Poorter, L., Portsmuth, A., Potvin, C., Ransijn, J., Reid, D., Ribeiro, S.C.,
Roberts, S.D., Rodríguez, R., Saldaña-Acosta, A., Santa-Regina, I., Sasa, K., Selaya, N.G., Sillett,
S.C., Sterck, F., Takagi, K., Tange, T., Tanouchi, H., Tissue, D., Umehara, T., Utsugi, H.,
Vadeboncoeur, M.A., Valladares, F., Vanninen, P., Wang, J.R., Wenk, E., Williams, R.,
Ximenes, F. de A., Yamaba, A., Yamada, T., Yamakura, T., Yanai, R.D. & York, R.A. (2015)
BAAD: a Biomass And Allometry Database for woody plants. *Ecology*, **96**, 1445–1445.
Felsenstein, J. (1985) Phylogenies and the Comparative Method. *The American Naturalist*, **125**, 1–15.

Fortini, L.B., Kaiser, L.R., Keith, L.M., Price, J., Hughes, R.F., Jacobi, J.D. & Friday, J.B. (2019) The
evolving threat of Rapid ʻŌhiʻa Death (ROD) to Hawaiʻi's native ecosystems and rare plant
species. *Forest Ecology and Management*, **448**, 376–385.
Fridley, J.D., Stachowicz, J.J., Naeem, S., Sax, D.F., Seabloom, E.W., Smith, M.D., Stohlgren, T.J.,
Tilman, D. & Holle, B.V. (2007) The invasion paradox: reconciling pattern and process in species
invasions. *Ecology*, **88**, 3–17.
Fukami, T. (2015) Historical Contingency in Community Assembly: Integrating Niches, Species Pools,
and Priority Effects. *Annual Review of Ecology, Evolution, and Systematics*, **46**, 1–23.
Fukami, T., Mordecai, E.A. & Ostling, A. (2016) A framework for priority effects. *Journal of Vegetation
Science*, **27**, 655–657.
Gallien, L. & Carboni, M. (2017) The community ecology of invasive species: where are we and what's
next? *Ecography*, **40**.
García-Verdugo, C., Monroy, P., Pugnaire, F.I., Jura-Morawiec, J., Moreira, X. & Flexas, J. (2020) Leaf
functional traits and insular colonization: Subtropical islands as a melting pot of trait diversity in
a widespread plant lineage. *Journal of Biogeography*, **47**, 2362– 2376.
Giambelluca, T., Shuai, X., Barnes, M., Alliss, R., Longman, R., Miura, T., Chen, Q., Frazier, A., Mudd,
R., Cuo, L. & others (2014) Evapotranspiration of Hawaiʻi. *Final report submitted to the US
Army Corps of Engineers—Honolulu District, and the Commission on Water Resource
Management, State of Hawaiʻi.*
Giambelluca, T.W., Chen, Q., Frazier, A.G., Price, J.P., Chen, Y.-L., Chu, P.-S., Eischeid, J.K. &
Delparte, D.M. (2013) Online rainfall atlas of Hawaiʻi. *Bulletin of the American Meteorological
Society*, **94**, 313–316.
Gillespie, R.G. (2016) Island time and the interplay between ecology and evolution in species
diversification. *Evolutionary Applications*, **9**, 53–73.
Gillespie Thomas W., Keppel Gunnar, Pau Stephanie, Price Jonathan P., Jaffré Tanguy & O'Neill Kristin
(2013) Scaling species richness and endemism of tropical dry forests on oceanic islands.
Diversity and Distributions, **19**, 896–906.
Givnish, T.J., Millam, K.C., Mast, A.R., Paterson, T.B., Theim, T.J., Hipp, A.L., Henss, J.M., Smith, J.F.,
Wood, K.R. & Sytsma, K.J. (2009) Origin, adaptive radiation and diversification of the Hawaiian
lobeliads (Asterales: Campanulaceae). *Proceedings of the Royal Society of London B: Biological
Sciences*, **276**, 407–416.
Gómez-Aparicio, L., Ibáñez, B., Serrano, M.S., Vita, P.D., Ávila, J.M., Pérez-Ramos, I.M., García, L.V.,
Sánchez, M.E. & Marañón, T. (2012) Spatial patterns of soil pathogens in declining
Mediterranean forests: implications for tree species regeneration. *New Phytologist*, **194**, 1014–
1024.
González-Moreno, P., Diez, J.M., Ibáñez, I., Font, X. & Vilà, M. (2014) Plant invasions are context-
dependent: multiscale effects of climate, human activity and habitat. *Diversity and Distributions*,
20, 720–731.
Grainger, T.N., Letten, A.D., Gilbert, B. & Fukami, T. (2019) Applying modern coexistence theory to
priority effects. *Proceedings of the National Academy of Sciences*, **116**, 6205–6210.
Gross, N., Börger, L., Duncan, R.P. & Hulme, P.E. (2013) Functional differences between alien and
native species: do biotic interactions determine the functional structure of highly invaded
grasslands? *Functional Ecology*, **27**, 1262–1272.
Henn, J.J., Yelenik, S. & Damschen, E.I. (2019) Environmental gradients influence differences in leaf
functional traits between native and non-native plants. *Oecologia*, **191**, 397–409.
Hughes, R.F. & Denslow, J.S. (2005) Invasion by a N2-Fixing Tree Alters Function and Structure in Wet
Lowland Forests of Hawaii. *Ecological Applications*, **15**, 1615–1628.
Ibanez, T., Hart, P., Ainsworth, A., Gross, J. & Monello, R. (2019) Factors associated with alien plant
richness, cover and composition differ in tropical island forests. *Diversity and Distributions*, **25**,
1910–1923.
Igea, J., Miller, E.F., Papadopulos, A.S.T. & Tanentzap, A.J. (2017) Seed size and its rate of evolution

correlate with species diversification across angiosperms. *PLOS Biology*, **15**, 1–16.

Imada, C.T. (2012) Hawaiian native and naturalized vascular plants checklist. *Honolulu: Hawaii Biological Survey, Bishop Museum*.

Jacobi, J.D., Price, J.P., Fortini, L.B., Gon, S.M. & Berkowitz, P. (2017) Baseline land cover. *Baseline and projected future carbon storage and carbon fluxes in ecosystems of Hawai'i. US Geological Survey Professional Paper 1834. Reston, VA: US Department of the Interior, US Geological Survey*, 9–20.

Jarvis, A., Reuter, H., Nelson, A. & Guevara, E. (2008) *SRTM 90m Digital Elevation Database v4. 1*, CGIAR-CSI.

Jauni, M., Gripenberg, S. & Ramula, S. (2015) Non-native plant species benefit from disturbance: a meta-analysis. *Oikos*, **124**, 122–129.

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Tautenhahn, S., Werner, G.D.A., Aakala, T., Abedi, M., Acosta, A.T.R., Adamidis, G.C., Adamson, K., Aiba, M., Albert, C.H., Alcántara, J.M., C, C.A., Aleixo, I., Ali, H., Amiaud, B., Ammer, C., Amoroso, M.M., Anand, M., Anderson, C., Anten, N., Antos, J., Apgaua, D.M.G., Ashman, T.-L., Asmara, D.H., Asner, G.P., Aspinwall, M., Atkin, O., Aubin, I., Baastrup-Spohr, L., Bahalkeh, K., Bahn, M., Baker, T., Baker, W.J., Bakker, J.P., Baldocchi, D., Baltzer, J., Banerjee, A., Baranger, A., Barlow, J., Barneche, D.R., Baruch, Z., Bastianelli, D., Battles, J., Bauerle, W., Bauters, M., Bazzato, E., Beckmann, M., Beeckman, H., Beierkuhnlein, C., Bekker, R., Belfry, G., Belluau, M., Beloiu, M., Benavides, R., Benomar, L., Berdugo-Lattke, M.L., Berenguer, E., Bergamin, R., Bergmann, J., Carlucci, M.B., Berner, L., Bernhardt-Römermann, M., Bigler, C., Bjorkman, A.D., Blackman, C., Blanco, C., Blonder, B., Blumenthal, D., Bocanegra-González, K.T., Boeckx, P., Bohlman, S., Böhning-Gaese, K., Boisvert-Marsh, L., Bond, W., Bond-Lamberty, B., Boom, A., Boonman, C.C.F., Bordin, K., Boughton, E.H., Boukili, V., Bowman, D.M.J.S., Bravo, S., Brendel, M.R., Broadley, M.R., Brown, K.A., Bruelheide, H., Brumnich, F., Bruun, H.H., Bruy, D., Buchanan, S.W., Bucher, S.F., Buchmann, N., Buitenwerf, R., Bunker, D.E., Bürger, J., Burrascano, S., Burslem, D.F.R.P., Butterfield, B.J., Byun, C., Marques, M., Scalon, M.C., Caccianiga, M., Cadotte, M., Cailleret, M., Camac, J., Camarero, J.J., Campany, C., Campetella, G., Campos, J.A., Cano-Arboleda, L., Canullo, R., Carbognani, M., Carvalho, F., Casanoves, F., Castagneyrol, B., Catford, J.A., Cavender-Bares, J., Cerabolini, B.E.L., Cervellini, M., Chacón-Madrigal, E., Chapin, K., Chapin, F.S., Chelli, S., Chen, S.-C., Chen, A., Cherubini, P., Chianucci, F., Choat, B., Chung, K.-S., Chytrý, M., Ciccarelli, D., Coll, L., Collins, C.G., Conti, L., Coomes, D., Cornelissen, J.H.C., Cornwell, W.K., Corona, P., Coyea, M., Craine, J., Craven, D., Cromsigt, J.P.G.M., Csecserits, A., Cufar, K., Cuntz, M., Silva, A.C. da, Dahlin, K.M., Dainese, M., Dalke, I., Fratte, M.D., Dang-Le, A.T., Danihelka, J., Dannoura, M., Dawson, S., Beer, A.J. de, Frutos, A.D., Long, J.R.D., Dechant, B., Delagrange, S., Delpierre, N., Derroire, G., Dias, A.S., Diaz-Toribio, M.H., Dimitrakopoulos, P.G., Dobrowolski, M., Doktor, D., Dřevojan, P., Dong, N., Dransfield, J., Dressler, S., Duarte, L., Ducouret, E., Dullinger, S., Durka, W., Duursma, R., Dymova, O., E-Vojtkó, A., Eckstein, R.L., Ejtehadi, H., Elser, J., Emilio, T., Engemann, K., Erfanian, M.B., Erfmeier, A., Esquivel-Muelbert, A., Esser, G., Estiarte, M., Domingues, T.F., Fagan, W.F., Fagúndez, J., Falster, D.S., Fan, Y., Fang, J., Farris, E., Fazlioglu, F., Feng, Y., Fernandez-Mendez, F., Ferrara, C., Ferreira, J., Fidelis, A., Finegan, B., Firn, J., Flowers, T.J., Flynn, D.F.B., Fontana, V., Forey, E., Forgiarini, C., François, L., Frangipani, M., Frank, D., Frenette-Dussault, C., Freschet, G.T., Fry, E.L., Fyllas, N.M., Mazzochini, G.G., Gachet, S., Gallagher, R., Ganade, G., Ganga, F., García-Palacios, P., Gargaglione, V., Garnier, E., Garrido, J.L., Gasper, A.L. de, Gea-Izquierdo, G., Gibson, D., Gillison, A.N., Giroldo, A., Glasenhardt, M.-C., Gleason, S., Gliesch, M., Goldberg, E., Göldel, B., Gonzalez-Akre, E., Gonzalez-Andujar, J.L., González-Melo, A., González-Robles, A., Graae, B.J., Granda, E., Graves, S., Green, W.A., Gregor, T., Gross, N., Guerin, G.R., Günther, A., Gutiérrez, A.G., Haddock, L., Haines, A., Hall, J., Hambuckers, A., Han, W., Harrison, S.P., Hattingh, W., Hawes, J.E., He, T., He, P., Heberling, J.M., Helm, A., Hempel, S., Hentschel, J., Hérault, B., Hereş, A.-M., Herz, K.,

Heuertz, M., Hickler, T., Hietz, P., Higuchi, P., Hipp, A.L., Hirons, A., Hock, M., Hogan, J.A.,
Holl, K., Honnay, O., Hornstein, D., Hou, E., Hough-Snee, N., Hovstad, K.A., Ichie, T., Igić, B.,
Illa, E., Isaac, M., Ishihara, M., Ivanov, L., Ivanova, L., Iversen, C.M., Izquierdo, J., Jackson,
R.B., Jackson, B., Jactel, H., Jagodzinski, A.M., Jandt, U., Jansen, S., Jenkins, T., Jentsch, A.,
Jespersen, J.R.P., Jiang, G.-F., Johansen, J.L., Johnson, D., Jokela, E.J., Joly, C.A., Jordan, G.J.,
Joseph, G.S., Junaedi, D., Junker, R.R., Justes, E., Kabzems, R., Kane, J., Kaplan, Z., Kattenborn,
T., Kavelenova, L., Kearsley, E., Kempel, A., Kenzo, T., Kerkhoff, A., Khalil, M.I., Kinlock,
N.L., Kissling, W.D., Kitajima, K., Kitzberger, T., Kjøller, R., Klein, T., Kleyer, M., Klimešová,
J., Klipel, J., Kloeppel, B., Klotz, S., Knops, J.M.H., Kohyama, T., Koike, F., Kollmann, J.,
Komac, B., Komatsu, K., König, C., Kraft, N.J.B., Kramer, K., Kreft, H., Kühn, I.,
Kumarathunge, D., Kuppler, J., Kurokawa, H., Kurosawa, Y., Kuyah, S., Laclau, J.-P., Lafleur,
B., Lallai, E., Lamb, E., Lamprecht, A., Larkin, D.J., Laughlin, D., Bagousse-Pinguet, Y.L.,
Maire, G. le, Roux, P.C. le, Roux, E. le, Lee, T., Lens, F., Lewis, S.L., Lhotsky, B., Li, Y., Li, X.,
Lichstein, J.W., Liebergesell, M., Lim, J.Y., Lin, Y.-S., Linares, J.C., Liu, C., Liu, D., Liu, U.,
Livingstone, S., Llusià, J., Lohbeck, M., López-García, Á., Lopez-Gonzalez, G., Lososová, Z.,
Louault, F., Lukács, B.A., Lukeš, P., Luo, Y., Lussu, M., Ma, S., Pereira, C.M.R., Mack, M.,
Maire, V., Mäkelä, A., Mäkinen, H., Malhado, A.C.M., Mallik, A., Manning, P., Manzoni, S.,
Marchetti, Z., Marchino, L., Marcilio-Silva, V., Marcon, E., Marignani, M., Markesteijn, L.,
Martin, A., Martínez-Garza, C., Martínez-Vilalta, J., Mašková, T., Mason, K., Mason, N.,
Massad, T.J., Masse, J., Mayrose, I., McCarthy, J., McCormack, M.L., McCulloh, K., McFadden,
I.R., McGill, B.J., McPartland, M.Y., Medeiros, J.S., Medlyn, B., Meerts, P., Mehrabi, Z., Meir,
P., Melo, F.P.L., Mencuccini, M., Meredieu, C., Messier, J., Mészáros, I., Metsaranta, J.,
Michaletz, S.T., Michelaki, C., Migalina, S., Milla, R., Miller, J.E.D., Minden, V., Ming, R.,
Mokany, K., Moles, A.T., Molnár, A., Molofsky, J., Molz, M., Montgomery, R.A., Monty, A.,
Moravcová, L., Moreno-Martínez, A., Moretti, M., Mori, A.S., Mori, S., Morris, D., Morrison, J.,
Mucina, L., Mueller, S., Muir, C.D., Müller, S.C., Munoz, F., Myers-Smith, I.H., Myster, R.W.,
Nagano, M., Naidu, S., Narayanan, A., Natesan, B., Negoita, L., Nelson, A.S., Neuschulz, E.L.,
Ni, J., Niedrist, G., Nieto, J., Niinemets, Ü., Nolan, R., Nottebrock, H., Nouvellon, Y.,
Novakovskiy, A., Nystuen, K.O., O'Grady, A., O'Hara, K., O'Reilly-Nugent, A., Oakley, S.,
Oberhuber, W., Ohtsuka, T., Oliveira, R., Öllerer, K., Olson, M.E., Onipchenko, V., Onoda, Y.,
Onstein, R.E., Ordonez, J.C., Osada, N., Ostonen, I., Ottaviani, G., Otto, S., Overbeck, G.E.,
Ozinga, W.A., Pahl, A.T., Paine, C.E.T., Pakeman, R.J., Papageorgiou, A.C., Parfionova, E.,
Pärtel, M., Patacca, M., Paula, S., Paule, J., Pauli, H., Pausas, J.G., Peco, B., Penuelas, J., Perea,
A., Peri, P.L., Petisco-Souza, A.C., Petraglia, A., Petritan, A.M., Phillips, O.L., Pierce, S., Pillar,
V.D., Pisek, J., Pomogaybin, A., Poorter, H., Portsmuth, A., Poschlod, P., Potvin, C., Pounds, D.,
Powell, A.S., Power, S.A., Prinzing, A., Puglielli, G., Pyšek, P., Raevel, V., Rammig, A.,
Ransijn, J., Ray, C.A., Reich, P.B., Reichstein, M., Reid, D.E.B., Réjou-Méchain, M., Dios, V.R.
de, Ribeiro, S., Richardson, S., Riibak, K., Rillig, M.C., Riviera, F., Robert, E.M.R., Roberts, S.,
Robroek, B., Roddy, A., Rodrigues, A.V., Rogers, A., Rollinson, E., Rolo, V., Römermann, C.,
Ronzhina, D., Roscher, C., Rosell, J.A., Rosenfield, M.F., Rossi, C., Roy, D.B., Royer-Tardif, S.,
Rüger, N., Ruiz-Peinado, R., Rumpf, S.B., Rusch, G.M., Ryo, M., Sack, L., Saldaña, A., Salgado-
Negret, B., Salguero-Gomez, R., Santa-Regina, I., Santacruz-García, A.C., Santos, J., Sardans, J.,
Schamp, B., Scherer-Lorenzen, M., Schleuning, M., Schmid, B., Schmidt, M., Schmitt, S.,
Schneider, J.V., Schowanek, S.D., Schrader, J., Schrodt, F., Schuldt, B., Schurr, F., Garvizu,
G.S., Semchenko, M., Seymour, C., Sfair, J.C., Sharpe, J.M., Sheppard, C.S., Sheremetiev, S.,
Shiodera, S., Shipley, B., Shovon, T.A., Siebenkäs, A., Sierra, C., Silva, V., Silva, M., Sitzia, T.,
Sjöman, H., Slot, M., Smith, N.G., Sodhi, D., Soltis, P., Soltis, D., Somers, B., Sonnier, G.,
Sørensen, M.V., Sosinski, E.E., Soudzilovskaia, N.A., Souza, A.F., Spasojevic, M., Sperandii,
M.G., Stan, A.B., Stegen, J., Steinbauer, K., Stephan, J.G., Sterck, F., Stojanovic, D.B., Strydom,
T., Suarez, M.L., Svenning, J.-C., Svitková, I., Svitok, M., Svoboda, M., Swaine, E., Swenson,
N., Tabarelli, M., Takagi, K., Tappeiner, U., Tarifa, R., Tauugourdeau, S., Tavsanoglu, C., Beest,

M. te, Tedersoo, L., Thiffault, N., Thom, D., Thomas, E., Thompson, K., Thornton, P.E.,
Thuiller, W., Tichý, L., Tissue, D., Tjoelker, M.G., Tng, D.Y.P., Tobias, J., Török, P., Tarin, T.,
Torres-Ruiz, J.M., Tóthmérész, B., Treurnicht, M., Trivellone, V., Trolliet, F., Trotsiuk, V.,
Tsakalos, J.L., Tsiripidis, I., Tysklind, N., Umehara, T., Usoltsev, V., Vadeboncoeur, M., Vaezi,
J., Valladares, F., Vamosi, J., Bodegom, P.M. van, Breugel, M. van, Cleemput, E.V., Weg, M.
van de, Merwe, S. van der, Plas, F. van der, Sande, M.T. van der, Kleunen, M. van, Meerbeek,
K.V., Vanderwel, M., Vanselow, K.A., Vårhammar, A., Varone, L., Valderrama, M.Y.V.,
Vassilev, K., Vellend, M., Veneklaas, E.J., Verbeeck, H., Verheyen, K., Vibrans, A., Vieira, I.,
Villacís, J., Violle, C., Vivek, P., Wagner, K., Waldram, M., Waldron, A., Walker, A.P., Waller,
M., Walther, G., Wang, H., Wang, F., Wang, W., Watkins, H., Watkins, J., Weber, U., Weedon,
J.T., Wei, L., Weigelt, P., Weiher, E., Wells, A.W., Wellstein, C., Wenk, E., Westoby, M.,
Westwood, A., White, P.J., Whitten, M., Williams, M., Winkler, D.E., Winter, K., Womack, C.,
Wright, I.J., Wright, S.J., Wright, J., Pinho, B.X., Ximenes, F., Yamada, T., Yamaji, K., Yanai,
R., Yankov, N., Yguel, B., Zanini, K.J., Zanne, A.E., Zelený, D., Zhao, Y.-P., Zheng, J., Zheng,
J., Ziemińska, K., Zirbel, C.R., Zizka, G., Zo-Bi, I.C., Zotz, G. & Wirth, C. (2020) TRY plant
trait database – enhanced coverage and open access. *Global Change Biology*, **26**, 119–188.
Keith, L.M., Hughes, R.F., Sugiyama, L.S., Heller, W.P., Bushe, B.C. & Friday, J.B. (2015) First Report
of Ceratocystis Wilt on `Ōhi`a (Metrosideros polymorpha). *Plant Disease*, **99**, 1276.
Kembel, S.W. (2009) Disentangling niche and neutral influences on community assembly: assessing the
performance of community phylogenetic structure tests. *Ecology Letters*, **12**, 949–960.
King, D.A., Davies, S.J. & Noor, N.S.M. (2006) Growth and mortality are related to adult tree size in a
Malaysian mixed dipterocarp forest. *Forest Ecology and Management*, **223**, 152–158.
König, C., Weigelt, P., Taylor, A., Stein, A., Dawson, W., Essl, F., Pergl, J., Pyšek, P., van Kleunen, M.,
Winter, M., Chatelain, C., Wieringa, J.J., Krestov, P. & Kreft, H. (2021) Source pools and
disharmony of the world's island floras. *Ecography*, **44**, 44–55.
Kraft, N.J.B. & Ackerly, D.D. (2010) Functional trait and phylogenetic tests of community assembly
across spatial scales in an Amazonian forest. *Ecological Monographs*, **80**, 401–422.
Kraft, N.J.B., Adler, P.B., Godoy, O., James, E.C., Fuller, S. & Levine, J.M. (2015) Community
assembly, coexistence and the environmental filtering metaphor. *Functional Ecology*, **29**, 592–
599.
Kraft, N.J.B., Valencia, R. & Ackerly, D.D. (2008) Functional Traits and Niche-Based Tree Community
Assembly in an Amazonian Forest. *Science*, **322**, 580–582.
Kueffer, C., Daehler, C.C., Torres-Santana, C.W., Lavergne, C., Meyer, J.-Y., Otto, R. & Silva, L. (2010)
A global comparison of plant invasions on oceanic islands. *Perspectives in Plant Ecology,*
Evolution and Systematics, **12**, 145–161.
Kunstler, G., Lavergne, S., Courbaud, B., Thuiller, W., Vieilledent, G., Zimmermann, N.E., Kattge, J. &
Coomes, D.A. (2012) Competitive interactions between forest trees are driven by species' trait
hierarchy, not phylogenetic or functional similarity: implications for forest community assembly.
Ecology Letters, **15**, 831–840.
Leihy, R.I., Duffy, G.A. & Chown, S.L. (2018) Species richness and turnover among indigenous and
introduced plants and insects of the Southern Ocean Islands. *Ecosphere*, **9**, e02358.
Lemoine, N.P., Shue, J., Verrico, B., Erickson, D., Kress, W.J. & Parker, J.D. (2015) Phylogenetic
relatedness and leaf functional traits, not introduced status, influence community assembly.
Ecology, **96**, 2605–2612.
Leung, B., Roura-Pascual, N., Bacher, S., Heikkilä, J., Brotons, L., Burgman, M.A., Dehnen-Schmutz, K.,
Essl, F., Hulme, P.E., Richardson, D.M., Sol, D. & Vilà, M. (2012) TEASIng apart alien species
risk assessments: a framework for best practices. *Ecology Letters*, **15**, 1475–1493.
Levin, S.C., Crandall, R.M., Pokoski, T., Stein, C. & Knight, T.M. (2020) Phylogenetic and functional
distinctiveness explain alien plant population responses to competition. *Proceedings of the Royal*
Society B: Biological Sciences, **287**, 20201070.
Li, D. & Ives, A.R. (2017) The statistical need to include phylogeny in trait-based analyses of community

composition. *Methods in Ecology and Evolution*, **8**, 1192–1199.

Lockwood, J.L., Cassey, P. & Blackburn, T. (2005) The role of propagule pressure in explaining species
invasions. *Trends in Ecology & Evolution*, **20**, 223–228.

Losos, J.B. & Ricklefs, R.E. (2009) Adaptation and diversification on islands. *Nature*, **457**, 830.

Ma, C., Li, S., Pu, Z., Tan, J., Liu, M., Zhou, J., Li, H. & Jiang, L. (2016) Different effects of invader–
native phylogenetic relatedness on invasion success and impact: a meta-analysis of Darwin's
naturalization hypothesis. *Proceedings of the Royal Society B: Biological Sciences*, **283**,
20160663.

Mack, R.N., Simberloff, D., Lonsdale, W.M., Evans, H., Clout, M. & Bazzaz, F.A. (2000) Biotic
Invasions: Causes, Epidemiology, Global Consequences, and Control. *Ecological Applications*,
10, 689–710.

Maitner, B. (2020) BIEN: tools for accessing the botanical information and ecology network database. *R*
package version 1.2.5.

Marx, H.E., Giblin, D.E., Dunwiddie, P.W. & Tank, D.C. (2016) Deconstructing darwin's naturalization
conundrum in the san juan islands using community phylogenetics and functional traits. *Diversity*
and Distributions, **22**, 318–331.

Mortenson, L.A., Flint Hughes, R., Friday, J.B., Keith, L.M., Barbosa, J.M., Friday, N.J., Liu, Z. &
Sowards, T.G. (2016) Assessing spatial distribution, stand impacts and rate of Ceratocystis
fimbriata induced 'ōhi'a (Metrosideros polymorpha) mortality in a tropical wet forest, Hawai'i
Island, USA. *Forest Ecology and Management*, **377**, 83–92.

Moser, D., Lenzner, B., Weigelt, P., Dawson, W., Kreft, H., Pergl, J., Pyšek, P., Kleunen, M. van, Winter,
M., Capinha, C., Cassey, P., Dullinger, S., Economo, E.P., García-Díaz, P., Guénard, B.,
Hofhansl, F., Mang, T., Seebens, H. & Essl, F. (2018) Remoteness promotes biological invasions
on islands worldwide. *Proceedings of the National Academy of Sciences*, **115**, 9270–9275.

Muller-Landau, H.C. (2004) Interspecific and Inter-site Variation in Wood Specific Gravity of Tropical
Trees. *Biotropica*, **36**, 20–32.

Muller-Landau Helene C. (2010) The tolerance–fecundity trade-off and the maintenance of diversity in
seed size. *Proceedings of the National Academy of Sciences*, **107**, 4242–4247.

Ni, M., Deane, D.C., Li, S., Wu, Y., Sui, X., Xu, H., Chu, C., He, F. & Fang, S. (2021) Invasion success
and impacts depend on different characteristics in non-native plants. *Diversity and Distributions*,
27, 1194–1207.

Ostertag, R., Inman-Narahari, F., Cordell, S., Giardina, C.P. & Sack, L. (2014) Forest structure in low-
diversity tropical forests: a study of Hawaiian wet and dry forests. *PLOS ONE*, **9**, 1–18.

Park Daniel S., Feng Xiao, Maitner Brian S., Ernst Kacey C. & Enquist Brian J. (2020) Darwin's
naturalization conundrum can be explained by spatial scale. *Proceedings of the National*
Academy of Sciences, **117**, 10904–10910.

Pearse, W.D., Cadotte, M.W., Cavender-Bares, J., Ives, A.R., Tucker, C.M., Walker, S.C. & Helmus,
M.R. (2015) pez: phylogenetics for the environmental sciences. *Bioinformatics*, **31**, 2888–2890.

Pearson, D.E., Ortega, Y.K., Eren, Ö. & Hierro, J.L. (2018) Community Assembly Theory as a
Framework for Biological Invasions. *Trends in Ecology & Evolution*, **33**, 313–325.

Penone, C., Davidson, A.D., Shoemaker, K.T., Marco, M.D., Rondinini, C., Brooks, T.M., Young, B.E.,
Graham, C.H. & Costa, G.C. (2014) Imputation of missing data in life-history trait datasets:
which approach performs the best? *Methods in Ecology and Evolution*, **5**, 961–970.

Perelman, S.B., Chaneton, E.J., Batista, W.B., Burkart, S.E. & León, R.J.C. (2007) Habitat stress, species
pool size and biotic resistance influence exotic plant richness in the Flooding Pampa grasslands.
Journal of Ecology, **95**, 662–673.

Pierce, S., Brusa, G., Vagge, I. & Cerabolini, B.E.L. (2013) Allocating CSR plant functional types: the
use of leaf economics and size traits to classify woody and herbaceous vascular plants.
Functional Ecology, **27**, 1002–1010.

Pyšek, P., Jarošík, V., Hulme, P.E., Kühn, I., Wild, J., Arianoutsou, M., Bacher, S., Chiron, F., Didžiulis,
V., Essl, F., Genovesi, P., Gherardi, F., Hejda, M., Kark, S., Lambdon, P.W., Desprez-Loustau,

941   M.-L., Nentwig, W., Pergl, J., Poboljšaj, K., Rabitsch, W., Roques, A., Roy, D.B., Shirley, S.,
942   Solarz, W., Vilà, M. & Winter, M. (2010) Disentangling the role of environmental and human
943   pressures on biological invasions across Europe. *Proceedings of the National Academy of*
944   *Sciences*, **107**, 12157–12162.
945 Qian, H., Jin, Y. & Ricklefs, R.E. (2017) Patterns of phylogenetic relatedness of angiosperm woody
946   plants across biomes and life-history stages. *Journal of Biogeography*, **44**, 1383–1392.
947 Qian, H. & Sandel, B. (2022) Darwin's preadaptation hypothesis and the phylogenetic structure of native
948   and alien regional plant assemblages across North America. *Global Ecology and Biogeography*,
949   **31**, 531–545.
950 R Core Team (2020) *R: A Language and Environment for Statistical Computing*, R Foundation for
951   Statistical Computing, Vienna, Austria.
952 Ricciardi, A., Hoopes, M.F., Marchetti, M.P. & Lockwood, J.L. (2013) Progress toward understanding
953   the ecological impacts of nonnative species. *Ecological Monographs*, **83**, 263–282.
954 Richardson, D.M. & Pyšek, P. (2012) Naturalization of introduced plants: ecological drivers of
955   biogeographical patterns. *New Phytologist*, **196**, 383–396.
956 Richardson, D.M. & Rejmánek, M. (2011) Trees and shrubs as invasive alien species – a global review.
957   *Diversity and Distributions*, **17**, 788–809.
958 van der Sande, M.T., Bruelheide, H., Dawson, W., Dengler, J., Essl, F., Field, R., Haider, S., van
959   Kleunen, M., Kreft, H., Pagel, J., Pergl, J., Purschke, O., Pyšek, P., Weigelt, P., Winter, M.,
960   Attorre, F., Aubin, I., Bergmeier, E., Chytrý, M., Dainese, M., De Sanctis, M., Fagundez, J.,
961   Golub, V., Guerin, G.R., Gutiérrez, A.G., Jandt, U., Jansen, F., Jiménez-Alfaro, B., Kattge, J.,
962   Kearsley, E., Klotz, S., Kramer, K., Moretti, M., Niinemets, Ü., Peet, R.K., Penuelas, J., Petřík,
963   P., Reich, P.B., Sandel, B., Schmidt, M., Sibikova, M., Violle, C., Whitfeld, T.J.S., Wohlgemuth,
964   T. & Knight, T.M. (2020) Similar factors underlie tree abundance in forests in native and alien
965   ranges. *Global Ecology and Biogeography*, **29**, 281–294.
966 Sapsford, S.J., Brandt, A.J., Davis, K.T., Peralta, G., Dickie, I.A., Gibson, R.D., Green, J.L., Hulme, P.E.,
967   Nuñez, M.A., Orwin, K.H., Pauchard, A., Wardle, D.A. & Peltzer, D.A. (2020) Towards a
968   framework for understanding the context dependence of impacts of non-native tree species.
969   *Functional Ecology*, **34**, 944–955.
970 Sax, D.F. & Gaines, S.D. (2008) Species invasions and extinction: the future of native biodiversity on
971   islands. *Proceedings of the National Academy of Sciences*, **105**, 11490–11497.
972 Schaefer, H., Hardy, O.J., Silva, L., Barraclough, T.G. & Savolainen, V. (2011) Testing Darwin's
973   naturalization hypothesis in the Azores. *Ecology letters*, **14**, 389–396.
974 Schluter, D. & Pennell, M.W. (2017) Speciation gradients and the distribution of biodiversity. *Nature*,
975   **546**, 48–55.
976 Seebens, H., Blackburn, T.M., Dyer, E.E., Genovesi, P., Hulme, P.E., Jeschke, J.M., Pagad, S., Pyšek, P.,
977   Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C., Celesti-Grapow,
978   L., Dawson, W., Dullinger, S., Fuentes, N., Jäger, H., Kartesz, J., Kenis, M., Kreft, H., Kühn, I.,
979   Lenzner, B., Liebhold, A., Mosena, A., Moser, D., Nishino, M., Pearman, D., Pergl, J., Rabitsch,
980   W., Rojas-Sandoval, J., Roques, A., Rorke, S., Rossinelli, S., Roy, H.E., Scalera, R., Schindler,
981   S., Štajerová, K., Tokarska-Guzik, B., van Kleunen, M., Walker, K., Weigelt, P., Yamanaka, T. &
982   Essl, F. (2017) No saturation in the accumulation of alien species worldwide. *Nature*
983   *Communications*, **8**, 14435.
984 Sherrod, D.R., Sinton, J.M., Watkins, S.E. & Brunt, K.M. (2007) *Geologic map of the State of Hawaii*,
985   Geological Survey (US).
986 Shiels, A.B. (2011) Frugivory by introduced black rats (Rattus rattus) promotes dispersal of invasive
987   plant seeds. *Biological Invasions*, **13**, 781–792.
988 Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp, F., Galil,
989   B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E. & Vilà, M. (2013) Impacts
990   of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution*, **28**,
991   58–66.

Spake, R., Mori, A.S., Beckmann, M., Martin, P.A., Christie, A.P., Duguid, M.C. & Doncaster, C.P.
(2021) Implications of scale dependence for cross-study syntheses of biodiversity differences.
Ecology Letters, **24**, 374–390.
Sperry, J.H., O'Hearn, D., Drake, D.R., Hruska, A.M., Case, S.B., Vizentin-Bugoni, J., Arnett, C.,
Chambers, T. & Tarwater, C.E. (2021) Fruit and seed traits of native and invasive plant species in
Hawai'i: implications for seed dispersal by non-native birds. *Biological Invasions*, **23**, 1819–
1835.
Spicer, R. & Groover, A. (2010) Evolution of development of vascular cambia and secondary growth.
New Phytologist, **186**, 577–592.
Stohlgren, T.J., Barnett, D.T. & Kartesz, J.T. (2003) The rich get richer: patterns of plant invasions in the
United States. *Frontiers in Ecology and the Environment*, **1**, 11–14.
Taylor, A., Weigelt, P., König, C., Zotz, G. & Kreft, H. (2019) Island disharmony revisited using orchids
as a model group. *New Phytologist*, **223**, 597–606.
Tecco, P.A., Díaz, S., Cabido, M. & Urcelay, C. (2010) Functional traits of alien plants across contrasting
climatic and land-use regimes: do aliens join the locals or try harder than them? *Journal of
Ecology*, **98**, 17–27.
Tershy, B.R., Shen, K.-W., Newton, K.M., Holmes, N.D. & Croll, D.A. (2015) The importance of islands
for the protection of biological and linguistic diversity. *BioScience*, **65**, 592–597.
Thomson, F.J., Moles, A.T., Auld, T.D. & Kingsford, R.T. (2011) Seed dispersal distance is more
strongly correlated with plant height than with seed mass. *Journal of Ecology*, **99**, 1299–1307.
Thuiller, W., Gallien, L., Boulangeat, I., Bello, F.D., Münkemüller, T., Roquet, C. & Lavergne, S. (2010)
Resolving Darwin's naturalization conundrum: a quest for evidence. *Diversity and Distributions*,
16, 461–475.
Tucker, C.M., Davies, T.J., Cadotte, M.W. & Pearse, W.D. (2018) On the relationship between
phylogenetic diversity and trait diversity. *Ecology*, **99**, 1473–1479.
Uriarte, M., Swenson, N.G., Chazdon, R.L., Comita, L.S., Kress, W.J., Erickson, D., Forero-Montaña, J.,
Zimmerman, J.K. & Thompson, J. (2010) Trait similarity, shared ancestry and the structure of
neighbourhood interactions in a subtropical wet forest: implications for community assembly.
Ecology Letters, **13**, 1503–1514.
US Forest Service (2016) Forest inventory and analysis national core field guide, Volume 1: Field data
collection procedures for Phase 2 Plots, version 7.1.
Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U., Sun, Y.
& Pyšek, P. (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on
species, communities and ecosystems. *Ecology Letters*, **14**, 702–708.
Vitousek, P.M., Walker, L.R., Whiteaker, L.D., Mueller-Dombois, D. & Matson, P.A. (1987) Biological
invasion by Myrica faya alters ecosystem development in Hawaii. *Science*, **238**, 802–804.
Wagner, W., Herbst, D. & Lorence, D. (2005) Flora of the Hawaiian Islands website.
(https://naturalhistory2.si.edu/botany/hawaiianflora/).
Westerband, A.C., Knight, T.M. & Barton, K.E. (2021) Intraspecific trait variation and reversals of trait
strategies across key climate gradients in native Hawaiian plants and non-native invaders. *Annals
of Botany*, **127**, 553–564.
Whittaker, R.J., Fernández-Palacios, J.M., Matthews, T.J., Borregaard, M.K. & Triantis, K.A. (2017)
Island biogeography: Taking the long view of nature's laboratories. *Science*, **357**.
Whittaker, R.J., Triantis, K.A. & Ladle, R.J. (2008) A general dynamic theory of oceanic island
biogeography. *Journal of Biogeography*, **35**, 977–994.
Wildlife Conservation Society - WCS & Center for International Earth Science Information Network -
CIESIN - Columbia University (2005) Last of the Wild Project, Version 2, 2005 (LWP-2):
Global Human Influence Index (HII) Dataset (Geographic).
Wohlwend, M.R., Craven, D., Weigelt, P., Seebens, H., Winter, M., Kreft, H., Zurell, D., Cabral, J.S.,
Essl, F., Kleunen, M. van, Pergl, J., Pyšek, P. & Knight, T.M. (2021) Anthropogenic and
environmental drivers shape diversity of naturalized plants across the Pacific. *Diversity and*

Distributions, **27**, 1120– 1133.

Zarnetske, P.L., Gouhier, T.C., Hacker, S.D., Seabloom, E.W. & Bokil, V.A. (2013) Indirect effects and

facilitation among native and non-native species promote invasion success along an

environmental stress gradient. *Journal of Ecology*, **101**, 905–915.

Zimmerman, N., Flint Hughes, R., Cordell, S., Hart, P., Chang, H.K., Perez, D., Like, R.K. & Ostertag, R.

(2008) Patterns of primary succession of native and introduced plants in lowland wet forests in

eastern Hawai 'i. *Biotropica*, **40**, 277–284.

Zimmerman, N., Hughes, R., Cordell, S., Hart, P., Chang, H., Perez, D., Like, R. & Ostertag, R. (2007)

Data from: Patterns of primary succession of native and introduced plants in lowland wet forests

in eastern Hawai'i, Dryad Digital Repository.

Zomer, R.J., Trabucco, A., Bossio, D.A. & Verchot, L.V. (2008) Climate change mitigation: A spatial

analysis of global land suitability for clean development mechanism afforestation and

reforestation. *Agriculture, Ecosystems & Environment*, **126**, 67–80.