TDP 2 modulates the expression of estrogen-responsive oncogenes

Nicholas Manguso[1], Minhyung Kim[1], Neeraj Joshi[1], Rasel Al Mahmud[2, 6], Juan Aldaco[1],

Ryusuke Suzuki[1], Felipe Cortes-Ledesma[3,8], Xiaojiang Cui[1,4], Shintaro Yamada[2], Shunichi

Takeda[2,7], Armando Giuliano[1,4], Sungyong You[1,4,5,] and Hisashi Tanaka[1,4,5]

[1]Department of Surgery, [4]Biomedical Sciences, and [5]Cedars-Sinai Cancer, Cedars-Sinai Medical

Center, West Hollywood, CA, 90048, USA

[2]Department of Radiation Genetics, Graduate School of Medicine, Kyoto University, 606-8501

Kyoto, Japan

[3]Centro Andaluz de Biologia Moleculary Medicina Regenerativa (CAMBIER), CSIC-

Universidad de Sevilla (Departmento de genetica) Sevilla, Spain

Corresponding author: Hisashi Tanaka (Hisashi.Tanaka@cshs.org)

Present addresses

[6]Center for Cancer Research, National Cancer Institute, Bethesda, MD, USA

[7]Shenzhen University School of Medicine, Shenzhen, Guangdong, China

[8]Topology and DNA breaks Group, Spanish National Cancer Centre (CNIO), Madrid, Spain

N.M and M.K. contributed equally to this work

Conflict of interest: The authors have no conflict of interest to declare regarding this study.

Abstract

With its ligand estrogen, the estrogen receptor (ER) stimulates tumor cell growth by activating a global transcriptional program. This activation involves topoisomerase 2 (TOP2), which resolves topological problems by transiently creating and re-ligating DNA double-strand breaks (DSBs). Recent studies have uncovered the involvement of DNA repair proteins in the repair of TOP2-induced DSBs. These noncanonical repair pathways may serve as backup processes when TOP2 is halted and fails to re-ligate DSBs, but their impact on transcription remains elusive. In this study, we investigated the role of tyrosyl-DNA phosphodiesterase 2 (TDP2), an enzyme that acts for the removal of halted TOP2 from the 5'-end of the DNA, in the estrogen-induced transcriptome. Using TDP2-deficient ER-positive cells and mice, we showed that TDP2 regulates the expression of oncogene *MYC*. *MYC* induction by estrogen was a very early event (1 hour) and TOP2-dependent. In TDP2-deficient cells, the induction of *MYC* by estrogen became prolonged and volatile. Bulk and single-cell RNA-seq identified the oncogenes *MYC* and *CCND1* as genes whose estrogen response was regulated by TDP2. These results suggest that TDP2 may play a role in the repair of TOP2-induced DSBs in specific genomic loci and tightly regulates the expression of oncogenes.

Introduction

With a lifetime risk of one in eight women, breast cancer stands as the most prevalent female cancer, remaining the most frequent cause of cancer-related death in women. Approximately two-thirds of breast tumors express estrogen receptor alpha (ERα), and their growth depends on the hormone estrogen [1]. Estrogen plays a crucial role in the development and regulation of female reproductive systems and secondary sex characteristics. 17beta-

estradiol (E2) is the predominant estrogen during reproductive years both in terms of absolute

serum levels and activity. As with all steroid hormones, secreted E2 diffuses into the cell nucleus,

where it binds to a receptor ERα, forming E2-ERα complex [2]. The E2-ERα complex

subsequently binds to DNA, acting as potent regulators of gene expression. Transcriptional

activation *in vitro* occurs within an hour after E2 stimulation and involves Type II topoisomerase

(TOP2), specifically, TOP2 beta (TOP2β) [3, 4].

TOP2 activities are crucial in managing torsional stress during various processes of DNA

metabolism, including replication, chromosome segregation, and transcription [5, 6]. It catalyzes

to induce a transient DNA double-stranded break (DSB) in a DNA duplex, with passing another

intact duplex through the broken DNA. This process involves a transient intermediate with a

covalent bond between the catalytic tyrosine in TOP2 and the 5' phosphate groups of the DNA

ends (TOP2 cleavage complex, TOP2cc), followed by the re-ligation of the DSB ends. In the

context of general transcription, advancing RNA polymerase II (PolII) on the DNA templates

creates over-wound DNA in front and leaves under-wound DNA behind. By relieving torsional

stress in front, TOP2 promotes the release of paused PolII for transcriptional elongation [7-9].

Because of the important and broad roles of TOP2 in nucleotide metabolisms, small molecule

inhibitors that intervene in each step of TOP2 activity have been developed. TOP2 catalytic

inhibitors prevent the induction of DSBs, while TOP2-poisons, by stabilizing TOP2cc, prevent

the re-ligation of TOP2-induced DSBs [10].

In addition to relieving torsional stress during general transcription, TOP2 plays a crucial

role in the induction of a specific set of genes in response to exogenous stimuli, including E2,

heat shock, serum starvation, and neuronal activity [7, 9, 11, 12]. On some of these gene loci,

TOP2β was shown to induce DSBs at transcriptional units, including promoters, in response to

these stimuli, and following the DSBs were the recruitment of DSB repair proteins, DNAPK and

Ku70 [9, 11]. This recruitment is intriguing, considering that TOP2 is established as a self-

sufficient enzyme, and DSBs induced by TOP2 are transient and usually re-ligated by TOP2 [5,

13]. TOP2β occupies over 2,000 genomic sites upon E2 treatment in MCF7 breast cancer cells

[14], suggesting a widespread involvement in transcriptional regulation. A fraction of these sites

would possibly recruit DSB repair proteins, as noncanonical repair processes for Top2-induced

DSBs. Two classes of repair proteins have been identified; (1) proteins involved in processing

(removing) TOP2cc that remains covalently attached to the 5' DSB ends and (2) those repairing

the resulting clean DSB ends. The first process was executed either by 5'-tyrosyl-DNA

phosphodiesterase (TDP2) or the Mre11-RAD50-Nbs1 (MRN) complex [15-17]. The resulting

clean ends are then repaired by non-homologous end-joining (NHEJ) [18]. BRCA1 is also

implicated in the repair of DSBs with TOP2cc [17, 19]. Although these noncanonical processes

are currently viewed as backup processes, for instance where TOP2 is trapped at the DNA ends

and requires assistance, it remains elusive whether noncanonical processes are inherently

programmed to contribute to the repair of TOP2-induced DSBs in specific gene loci.

Intriguingly, when left unrepaired, TOP2-induced DSBs can activate a specific set of

genes. For example, inhibiting the re-ligation of TOP2-induced DSBs by TOP2-poison etoposide

resulted in the upregulation of early response genes to neuronal activity in cultured primary

neurons [11]. Understanding the mechanism behind the upregulation is crucial. It is also essential

to identify which genes are upregulated in response to external stimuli when the repair of TOP2-

induced DSBs is hindered. The upregulation of target genes can pose a serious problem,

particularly when external stimuli, such as E2, promote cell proliferation. Such activation could

potentially lead to abnormal cell proliferation and, ultimately, contribute to tumorigenesis.

However, it remains unknown whether such upregulation can occur in a more natural setting of

DSB repair deficiency, for example, in cells lacking a DSB repair protein, compared to etoposide

treatment. Stalled TOP2cc by etoposide occurs very frequently in the entire cell population,

enabling the detection of transcriptional upregulation using techniques for measuring the average

gene expression levels in a cell population, such as qPCR, digital PCR, and bulk RNA-seq. In

contrast, in cells deficient in a DSB repair protein, stalled TOP2cc may not occur throughout the

entire cell population. In such a case, the aforementioned techniques might not detect

transcriptional upregulation. However, upregulation could still be observed in DSB repair-

deficient cells if TOP2-induced DSB is programmed to involve the DSB repair protein at a

certain gene locus. This possibility was shown at the protein level in mice lacking NHEJ repair;

E2 stimulation induced MYC protein in mammary epithelial cells of mice lacking DNA-PKcs

[20].

We aimed to identify a set of E2-induced genes that become upregulated in cells deficient

in TOP2-induced DSB repair. This gene set could also inform a potentially novel process by

elucidating gene loci where noncanonical repair of TOP2-induced DSBs is programmed. To

achieve this aim, we established ER-positive breast cancer cell lines lacking TDP2 and

investigated their global transcriptional program. We chose TDP2 because TDP2 knockout

cells/mice are viable [18] and amenable to E2 stimulation experiments. Using digital PCR and

bulk- and single-cell RNA-seq, we identified that *MYC* was among several genes that exhibited

enhanced transcription in response to E2 in TDP2-deficient cells compared to TDP2-proficient

cells. We also found that another breast cancer oncogene, *CCND1*, showed increased

transcription in TDP2-deficient cells. These results suggest that noncanonical repair processes of

TOP2-induced DSBs may tightly regulate oncogenic gene expression. The results may also

explain why deficiency of DSB repair proteins is often associated with the susceptibility of

breast tumors.

Results

MYC **induction by 17beta-estradiol (E2) depends on TOP2**

MYC is a well-studied oncogene and known to transcriptionally respond to E2 [21]. To

assess this response, we collected RNA from two ER-positive breast cancer cell lines MCF7 and

T47D, before and after treatment with E2 (**Fig. 1A**). We employed digital PCR assays to

measure the expression of *MYC* mRNA. In this assay, a *MYC* Taqman probe labeled with a dye

FAM and a control probe, either *TBP* or *TFRC,* labeled with a dye VIC were mixed with cDNA

from cell lines and PCR-amplified in approximately 20,000 x 1 nL reaction (**Fig. 1B**). *MYC*

expression levels (the number of *MYC*-positive 1 nL reactions) were normalized by the internal

control (the number of internal control-positive reactions). We added 10 nM E2 and incubated

cells for two hours before extracting RNA. The fold induction of *MYC* was calculated as the ratio

of normalized *MYC* expression with E2 treatment to normalized *MYC* expression without E2

treatment.

ER-positive (ER+) cell lines, MCF7 and T47D, showed an increase in *MYC* expression

after E2 treatment (2.59-fold and 2.17-fold, respectively). As expected, the ER-negative (ER-)

cell line MDA-MB-231 did not show the induction (0.93-fold change) (**Fig. 1C**). To understand

the temporal regulation of *MYC* expression, we measured *MYC* mRNA at 1, 2, and 6 hours after

the addition of 10 nM E2. In both the MCF7 and T47D cell lines, *MYC* expression peaked at 1

hour (MCF7 2.89-fold, T47D 2.79-fold), followed by gradual decreases until 6 hours (**Fig. 1D**).

As previous studies reported that TOP2 mediates the induction of E2-responsive genes [3], we

then tested whether *MYC* induction is also TOP2-dependent. We used a TOP2 catalytic inhibitor,

ICRF-193, which prevents TOP2 from creating DSBs [22, 23]. We added ICRF-193 to the cell

culture media 1 hour before adding 10 nM E2 and incubated cells for 2 hours after E2 treatment.

We observed a dose-dependent inhibition of *MYC* induction in both MCF7 and T47D (Figure

1E), strongly suggesting that E2 stimulation of *MYC* expression in ER-positive cells is TOP2-

dependent.

MYC transcriptional activation in TDP2 knockout cells

Since ICRF-193 inhibits DSB induction, DSBs may play a pivotal role in the

transcriptional activation of *MYC*. When left unrepaired, TOP2-induced DSBs promote

transcription for a specific set of genes [11]. Therefore, our next objective was to determine

whether unrepaired DSBs could further enhance *MYC* induction by E2. To test the idea, we

knocked out *TDP2* in ER-positive cell lines. When TOP2 fails to re-ligate DSBs and is aborted,

TOP2cc is processed by a ubiquitin-proteasome pathway or, alternatively, a proteosome-

independent pathway [24, 25]. TDP2 removes the remnant aborted TOP2 and creates a clean

DSB for NHEJ machinery to repair [18]. We used CRISPR-Cas9 to target *TDP2* in both MCF7

(MCF7 TDP2KO-1) and T47D cell lines. Additionally, we employed an independently created

TDP2 knockout MCF7 cell line (MCF7 TDP2KO-2)[17]. We confirmed the lack of TDP2

protein in knockout cells by western blotting (**Fig. 2A**). The functional TDP2 deficiency was also

confirmed using the TOP2 poison, etoposide. Etoposide traps TOP2cc at DSBs, which would

require TDP2 to repair. Both the MCF7 and T47D cells deficient in TDP2 were significantly

more sensitive to etoposide than parental cell lines (**Fig. 2B**).

We monitored the dynamics of *MYC* induction using digital PCR, as illustrated in **Figure**

1C. The highest level of induction at 1 hour after the addition of E2 (10nM) was similar between

MCF7 and MCF7 TDP2KO-1 cells (**Fig. 2C**). However, *MYC* expression continued to be high in

MCF7 TDP2KO cells, whereas it declined quickly in MCF7 cells. We further measured the

induction of *MYC* at 2 hours after the addition of 10 nM E2 and found the increased fold

induction of *MYC* in all three TDP2 knockout cell lines relative to their parental TDP2 wild-type

counterparts (**Fig. 2D**). Because the expression was measured by total RNA of entire cell

populations, these results suggest that *MYC* induction by E2 was enhanced in a significant

proportion of the TDP2KO cell population. The prevailing view is that alternative repair proteins,

including TDP2, only repair abortive TOP2-induced DSBs that could occur in a very small

fraction of cells. However, these results suggest that a significant fraction of cells require TDP2

to regulate *MYC* induction. The delayed repair of TOP2-induced DSBs could promote the *MYC*

response to E2 at 2 hours, as previously described for early response genes to neuronal activity

[11]. Consistently, we observed that etoposide promotes the induction of *MYC* in MCF7 cells

(**Fig. 2E**, left). Inhibiting NHEJ could also promote *MYC* response, as previously shown *in vivo*

in DNA-PKcs deficient mice [20]. To explore this possibility, we inhibited NHEJ by DNA-PKcs

inhibitor NU7441 [26]. Indeed, *MYC* induction at 2 hours after the addition of E2 was increased

by NU7441 (5 μM) (**Fig. 2E**, right).

Induction of MYC by estrogen *in vivo* in mice mammary glands

To assess the relevance of the above observation *in vivo*, we examined *MYC* response to

E2 at the protein level in mice mammary glands. We administered E2 (300 μg/kg body weight)

or PBS (control) intraperitoneally into three *Tdp2* wild-type and *Tdp2*$^{-/-}$ mice [18]. Mice were

examined before injection (0 hours) and 6 and 20 hours after injection for MYC protein

expression in CK8/18-positive mammary epithelial cells by immunostaining (**Fig. 3A**). We

quantified MYC-positive epithelial cells in 3 to 12 mammary ducts at each timepoint and scored

the fraction of MYC-positive cells in each mammary duct (**Fig. 3B**). MYC-positive cells

increased in both wild-type and *Tdp2*$^{-/-}$ mice at 6 hours. MYC-positive cells become less frequent

at 20 hours, showing transient induction of MYC in mammary epithelial cells. MYC-positive

cells in *Tdp2*$^{-/-}$ mice at 20 hours were significantly more abundant than in *Tdp2* wild-type mice

(p<0.01) at 20 hours or in *Tdp2*$^{-/-}$ mice without E2 at 0 hours (p<0.01). The prolonged induction

of MYC in *Tdp2*$^{-/-}$ mice parallels the observation in transcription in TDP2KO MCF7 cells (**Fig.**

2C) and aligned with our *in vitro* findings.

MYC and MYC downstream targets induced by E2

MYC is a beta Helix-Loop-Helix transcription factor that, in conjunction with a partner

protein Max, binds to DNA at the CACGTG E-Box sequence, thereby modulating global gene

expression [27]. To examine the expression of MYC-regulated genes in MCF7 and MCF7

TDP2KO cells, we conducted RNA-seq analysis using total RNA collected before and 2 hours

after the addition of 10nM E2 (bulk RNA-seq). We found a large number of differentially

expressed genes in both MCF7 (2,021 genes) and MCF7 TDP2 KO cells (1,034 genes) (**Tables**

S1 and **S2**; fold change ≥ 1.5 and combined P-value < 0.05). Notably, *MYC* showed upregulation

in both cell types, with a more pronounced increase in MCF7 TDP2KO cells (3.16 fold)

compared to parental MCF7 cells (1.94 fold) by E2 stimulation (**Fig. 4A**), validating the

prolonged response of *MYC* to estrogen in the absence of TDP2. *MYC* transcriptional response

was slightly more pronounced in MCF7 cells treated with etoposide (2.17 fold, **Table S3**),

suggesting the unrepaired TOP2-induced DSBs contributed to the enhanced response.

To assess the outcomes of *MYC* activation following E2 treatment in MCF7 TDP2KO

and parental MCF7 cells, we performed Gene Set Enrichment Analysis (GSEA) [28] with

predefined MYC target genes. GSEA determines whether a defined set of genes shows

statistically significant, concordant differences between two biological states. Specifically, we

examined the significant changes in the expression of genes whose promoters harbor MYC

binding sites within the 4 kb of the transcription start sites (MYC_Q2), potentially representing

direct targets of MYC (**Fig. S1, A**, top). Upon treating cells with E2 for 2 hours, the majority of

genes in MYC_Q2 exhibited nonrandom distribution, predominantly showing upregulation in

both MCF7 and MCF7 TDP2KO cells. We determined the normalized enrichment score (NES)

(**Fig. S1, B**), which reports the degree of overrepresentation of genes at the extremes of all the

ranked genes. We found that MYC_Q2 genes were slightly more clustered in highly upregulated

genes in MCF7 TDP2KO cells than in MCF7 cells. Furthermore, we examined gene sets whose

expression was enhanced by overexpressing MYC in human primary mammary epithelial cells

(Hallmark MYC target v2 and MYC_UP.V1_UP, **Fig. S1A** and **B**) [29]. Both sets demonstrated

he enrichment of MYC targets in upregulated genes, with comparable NES values between

MCF7 and MCF7 TDP2KO. These results suggest that the upregulation of MYC by E2

modulated global transcriptional programs in both cell types.

MYC induction at the single-cell level

Bulk RNA-seq offers insights into the average RNA expression across an entire cell

population. The observed upregulation of *MYC* in MCF7 TDP2KO cells could stem from either

(1) a scenario in which only a subset of cells robustly upregulates *MYC,* creating a bimodal

distribution that elevates the population average or (2) a situation where *MYC* is induced more

robustly in the vast majority of MCF7 TDP2KO cell population than in MCF7 cell population,

resulting in the normal distribution. The first scenario suggests that, for those DSBs that are

involved in *MYC* regulation after E2 treatments, TOP2-induced DSBs are primarily re-ligated by

TOP2 itself, and TDP2 plays a role in repair only in a subset of cells with abortive TOP2ccs. The

second scenario implies that TDP2 is programmed in the repair of TOP2-induced DSBs that

regulate *MYC* expression. These possibilities can be distinguished by single-cell RNA-seq (sc

RNA-seq) experiments.

Four groups of cells were subject to scRNA-seq: MCF7 parental cells, MCF7 parental

cells treated with E2 for 2 hours, MCF7 TDP2KO cells, and MCF7 TDP2KO cells treated with

E2 for 2 hours. Quality control and normalization of scRNA-seq data were executed using the

Seurat pipeline [30]. Following the exclusion of poor-quality cells and genes, the transcriptome

data comprised 354 genes in E2-treated MCF7, 585 genes in untreated MCF7, 708 genes in E2-

treated TDP2KO, and 399 genes in untreated TDP2KO. For the filtered and normalized data, we

conducted two independent comparisons: E2-treated MCF7 versus untreated MCF7 and E2-

treated TDP2KO versus untreated TDP2KO. Applying an adjusted p-value <0.05 and log fold

change ≥ 0.1 (1.11 fold), we identified a total of 111 genes that were differentially expressed

between untreated and E2-treated MCF7; 100 upregulated and 11 down-regulated genes,

respectively (**Table S4**). For TDP2-KO cells, E2 treatment resulted in gene expression changes

in 327 genes: 138 up- and 189 down-regulated genes (**Table S5**).

Consistent with the findings from digital PCR and bulk RNA-seq, *MYC* expression was

induced by E2 treatment in both MCF7 TDP2KO and MCF7 cells, with log2-fold changes of

0.43 (1.54 fold) and 0.17 (1.19 fold), respectively (**Fig. 4A**). Notably, the *MYC* induction by E2

was more pronounced in MCF7 TDP2KO cells than in MCF7. Normalized expression data

revealed that MCF7 TDP2KO cell populations displayed continuous distributions without

distinct subpopulations and uniquely included cells with very high *MYC* expression (normalized

expression >1.5) after E2 treatment (**Fig. 4B**). Histograms of raw read count were consistent with

the skewed unimodal distribution across all four populations (**Fig. 4C**). The normal distribution

of expression values at a single cell level suggests that most cells in the population contributed

equally to the mean increase, providing supports for the scenario (2).

We next examined whether the enhanced transcriptional response of the *MYC* in MCF7

TDP2KO cells is associated with cellular processes, such as cell cycle phases. Cell cycle phases

are considered confounding factors in single-cell studies [31]. To address this concern, we

allocated individual cells to G1, S, and G2/M phases using the cell cycle prediction tool based on

single transcriptome data [32]. We then examined whether cell cycle distribution might confound

the observed difference in *MYC* transcriptional responses by collecting *MYC* expression profiles

with and without E2 in each cell cycle phase (Fig. 4D). Based on the cell cycle phase prediction,

we found that MCF7 TDP2KO cells exhibited modestly higher enrichment in G1 cell

populations than MCF7 cells; >60% of TDP2 KO cells were in G1, while 50% of MCF7 cells

were in G1 (**Fig. S2**). However, the difference of *MYC* transcriptional induction by E2 was less

pronounced between MCF7 and MCF7 TDP2KO cells in G1 cells (1.26-fold with $p<0.0001$ in

MCF7 vs. 1.47-fold with $p<0.0001$ in MCF7 TDP2KO, Wilcoxon rank-sum test) than in S (1.08-

fold with $p=0.0332$ vs. 1.52-fold with $p<0.0001$) and G2M cells (1.24-fold with $p=0.0014$ vs.

1.88-fold with $p<0.0001$) (**Fig. 4C**). Therefore, the difference in the distribution of cell cycles

was not a major contributor to increased *MYC* induction in MCF7 TDP2KO cells. Instead, cell

cycle profiling revealed that the increased response to E2 would likely stem from cells in S/G2M

phases.

The dominance of upregulated genes by the short period of E2 treatment

The global transcriptional responses to E2 have been extensively studied in the MCF7

cell line, and both upregulated and downregulated genes have been documented. However,

variations in the amounts of E2 and treatment durations across experiments, as well as the

differences in genomic platforms (e.g., microarray, RNA-seq, and scRNA-seq), could influence

the repertoire of E2-responsive genes, as previously discussed [33, 34]. To establish a reliable

set of E2 (10nM)--responsive genes at an early time point (2 hours), we sought the intersections

between different genomic platforms, bulk RNA-seq, and scRNA-seq.

Bulk RNA-seq analysis identified a large number of differentially expressed genes,

including 2021 for MCF7 and 1034 for MCF7 TDP2KO, between untreated and E2-treated cells.

In contrast, the numbers of differentially expressed genes in our scRNA-seq data were modest:

111 genes for MCF7 and 327 genes for MCF7 TDP2KO (**Fig. 5A**). Thirty-two genes were

common between bulk and single-cell RNA-seq in MCF7, while 62 genes were shared in MCF7

TDP2KO cells. The majority of these common genes were upregulated genes in both cell lines

(**Fig. 5B; Tables S6** and **S7**). In MCF7, 28 genes were upregulated, and four genes were

repressed, with one gene, GPRC5A, in opposite directions between bulk- and sc-RNA-seq. In

MCF7 TDP2KO, 58 genes were upregulated, and four genes were repressed. There were 8

common differentially expressed genes between MCF7 and MCF7 TDP2KO, including *MYC*,

CCND1, *NRIP*, *SH3BP5*, *IGFBP4*, *RET*, *NBPF1* and *CA12* (**Fig, 5C**). Notably, *MYC* and

CCND1 are well-established oncogenes in breast cancer [35-38]. In summary, a reliable set of

estrogen-responsive genes at an early time point (2h) were predominantly upregulated genes,

with TDP2 potentially regulating responses to E2 for both *MYC* and *CCND1*.

Discussion

In this study, we assessed the transcriptional responses to 17beta-estradiol (E2), an

abundant form of estrogen, both *in vitro* and *in vivo*, in ER-positive breast cancer cells and breast

epithelial cells in mice. The transcriptional response to estrogen is known to involve

topoisomerase 2 (TOP2) [3], which creates transient DNA double-strand breaks, relieves

torsional stress, and reseal the ends. Recent studies have highlighted the roles of DNA repair

proteins, including TDP2, Mre11, BRCA1, and NHEJ proteins, in the repair of TOP2-induced

DSBs [15-18]. We aimed to gain insights into how these proteins contribute to the biological

processes involving TOP2-induced DSBs. To do so, we monitored transcriptional response to E2

in TDP2 wild-type and TDP2KO MCF7 and T47D cell line. Defects in the repair of TOP2-

induced DSB have been shown to upregulate transcriptional response to external stimuli [11].

Therefore, we anticipated that the extent of changes in the E2 response would be proportional to

the fraction of the population experiencing repair defects due to the absence of TDP2.

We confirmed that transcription of an oncogene *MYC* responded to E2 (10 nM) very

early, as early as 1 hour after the addition of E2 (**Fig. 1D**). The early response was inhibited

when the catalytic activity of TOP2 was blocked by ICRF-193 in a dose-dependent manner in

both MCF7 and T47D cell lines (**Fig. 1D**), strongly suggesting that TOP2 plays a crucial role in

the early transcriptional response of *MYC*. The transcriptional response is most likely through

TOP2-induced DSBs, as the TOP2 poison etoposide enhanced the transcriptional response (**Fig.**

2E). This notion was further supported by the *in vitro* experiments with cell lines deficient in

TDP2 using various platforms for gene expression measurements (**Fig. 2, 3** and **4**). The enhanced

MYC response was also observed in cells treated with an NHEJ inhibitor, indicating that

unrepaired TOP2-induced DSBs were likely the cause of upregulation (**Fig. 2E**). Importantly, we

detected the upregulation by the techniques that measured the average gene expression level in

the entire cell population, such as digital PCR and bulk mRNA-seq. Therefore, TDP2 seems to

be involved in the transcriptional regulation of *MYC* in a substantial fraction of the cell

population.

Using scRNA-seq, we further revealed that the observed enhanced response of *MYC* in

MCF7 TDP2KO cells, compared to MCF7 cells, was driven by the population-wide upregulation

of *MYC* by E2. This was evidenced by the unimodal distribution of *MYC* mRNA in MCF7

TDP2KO cells (**Fig. 4C**). The unimodality of gene expression was seen for *MYC* and other

commonly upregulated genes. In a previous study of scRNA-seq for the E2-stimulated

transcriptional response in MCF7 cells, a large proportion of genes showed a bimodal

distribution of gene expression [39]. However, the smaller numbers of cells captured for scRNA-

seq in that study (87 MCF7 cells) compared to our study (354 cells from MCF7 with E2, and 585

cells from MCF7 without E2) may contribute to the difference. With these results along with the

TOP2-dependent E2 response of *MYC* (**Fig. 1E**), we speculate that TDP2-dependent repair of

TOP2-induced DSBs is programmed to regulate *MYC* expression when stimulated by E2.

Unrepaired DSBs could potentially suppress the transcription of nearby genes [40-42].

Conversely, the accumulation of DNA damages at stimuli-induced genes would promote

transcription by releasing PolII from pausing [11]. Also, unrepaired DSBs far from the target

genes, such as enhancer elements, would not directly disrupt the elongation of target gene

mRNA. This scenario could apply to *MYC,* whose enhancer elements are dispersed within a few

Mb gene deserts surrounding *MYC* [43, 44]. Notably, an enhancer located at 135kb distal to *MYC*

(Myc+135kb enhancer) has an ER binding site and responds to E2 treatments [45]. The estrogen-

induced expression of enhancer RNA and the physical interaction between enhancer and *MYC*

promoter was greatly promoted when the repair of TOP2-induced DSBs was inhibited by an

ATM inhibitor (KU-55933). Although ATM and TDP2 function independently to repair

etoposide-induced DSBs [46], and spatial and temporal regulations of these two repair pathways

remain unclear, TDP2 may also regulate the estrogen-induced activation of *MYC* enhancers

through the repair of TOP2-induced DSBs.

scRNA-seq also enabled us to investigate the responses to E2 within distinct cell cycle

phases. We found that TDP2 KO cells responded to E2 more robustly in the S and G2/M phases

than in the G1 phase. There are two pathways for the removal of abortive TOP2ccs from the

ends: TDP2 and Mre11/BRCA1. It is currently unclear how these two pathways delineate their

tasks in removal. BRCA1 supports the nuclease activity of Mre11 to remove abortive TOP2ccs

[17], which was shown to be active in G1 cells. Considering the robust upregulation of *MYC* in S

and G2/M phases in TDP2-KO cells, we speculate that these two pathways repair DSBs with

abortive TOP2cc in different cell cycle phases. A recent study showed that TDP2 binds to K63

and K27 poly-ubiquitinated cellular proteins, and this binding stimulates TDP2-catalyzed

removal of TOP2 from DNA [47]. The identity of the K27 and/or K63 poly-ubiquitinated

proteins bound to TDP2 is not known. However, it is conceivable that such a protein could

regulate TDP2 activity in a cell-cycle-dependent manner.

We utilized both bulk and scRNA-seq for evaluating the responses of E2-responsive

genes at a 2-hour time point. The ER-positive MCF7 cell line has been employed to study E2-

responsive genes [33, 34]. However, these studies varied in their experimental conditions,

including measurement platforms, E2 concentrations, and treatment durations, resulting in

substantial variations in the numbers and spectra of E2-responsive genes. Additionally, genetic

variations among MCF7 sublines [48] may contribute to the variability in results. To address this,

we employed the same clones and maintained identical experimental conditions across two

RNA-seq platforms to minimize variabilities associated with the experimental design. The genes

called as common genes between the two platforms were strongly restricted by scRNA-seq data,

as only small fractions of differentially expressed genes detected by bulk RNA-seq were also

called by scRNA-seq. This discrepancy is likely attributed to the inherent heterogeneity of

scRNA-seq data, which often includes numerous zero values for a substantial fraction of genes

[49]. Nevertheless, these common differentially expressed genes showed a pronounced bias

toward upregulated genes. Notably, these genes included oncogenes *MYC* and *CCND1,*

frequently upregulated in ER-positive breast tumors. Estrogen induces mammary ductal

hyperplasia through MYC in mice deficient in NHEJ, which is downstream of TDP2-mediated

repair of TOP2cc. CCND1 is more frequently upregulated in ER-positive tumors than other

subtypes and promotes cell cycle progression [50, 51]. By participating in the regulation of these

oncogenes, TDP2 would emerge as a potential suppressor for the most common subtype of

breast tumors.

Materials and Methods

Cell culture and digital PCR   MCF7 and MCF7 TDP2KO cells were maintained with

MEM with 10% FBS. T47D, T47D TDP2KO, and MDA-MB-231 cells were maintained in

DMEM with 10% FBS. TDP2KO cells were created by targeting exon 2 of the *TDP2* gene (5'-

GGCTCAGAGATGGTTTCAGGT-3'). The targeting vector was constructed using

pSpCas9(BB)-2A-GFP(PX458), a gift from Feng Zhang (Addgene plasmid #48138).

Etoposide sensitivity was assessed through the colony formation assay. In brief, 3000

cells were plated in each well of a 6-well plate, and cells were cultured for 8 days with or without

etoposide before staining with crystal violet. Experiments were conducted in triplicate. Colonies

were counted using the Hybrid Cell Function of the Keyence BZ-X700 microscopy (Keyence).

RNA was extracted using the RNeasy mini kit, and first-strand cDNA synthesis was

performed using Superscript III (Invitrogen). Cells were cultured in phenol red-free medium

supplemented with 5% FBS-charcoal stripped for 24 hours then treated with 10 nM 17β-estradiol

for either 1, 2 or 6 hours before RNA extraction. NU7441 was added to the media 3 hours before

adding 10 nM E2. Etoposide was added to the media 1 hour before adding E2. Digital PCR was

conducted using the QuantStudio 3D Digital PCR System. A TaqMan probe labeled with FAM

(MYC-Hs00153408, IL20-Hs00218888 or OLFML3-Hs01113293) and VIC (TBP-Hs00427620

or TFRC-Hs00951083) were mixed in a reaction, and the amount of transcript (copies/microliter)

was determined using the Analysis Suite (Thermo Fisher Scientific).

Mouse experiments

Tdp2 Wild-type and *Tdp2$^{-/-}$* strains originated from the C57BL6/J strain. E2 (300 μg/kg

body weight) was administered via intraperitoneal injection (IP) into three independent two-

month-old female mice at a volume of 100 μL. As a control, an equivalent volume of a solvent

(PBS) is injected.

The mice were sacrificed to harvest mammary gland tissues and frozen blocks were

prepared. After a brief wash of the isolated tissues with cold PBS, the tissues were fixed with

paraformaldehyde (4%, Cat# 163-20145, Wako, Japan) for 15 min at 4° C and then washed three

times with 1x PBS. To preserve tissue integrity, incubation was performed in 30% sucrose (in

PBS) for 30 min to 3 hours. Tissues were then embedded in optimal cutting temperature (OCT)

compound (Cat# 4583, Sakura, Japan) into Cryomolds (Cat# 4565, Sakura, Japan). For cryo-

sectioning of frozen blocks, a cryostat (CM1850, Leica, Germany) was set at -25° C and

sectioned into 10 μm slices. The slides were then heated at 55° C for 10 to 30 min to dry. Dried

slides were washed three times with Tween-20 in PBS (0.05 %, PBS-T). For blocking, 1 % BSA

and 5 % goat serum in PBS-T were used for 3 hours at room temperature, followed by washing

once with PBS-T. Slides were then incubated overnight at 4° C with both α-Cytokeratin-8/18

(1/10, rat monoclonal, University of Iowa, US) and α- Myc antibody (1/200 [Y69], ab32072,

rabbit monoclonal, Abcam). After washing with PBS-T, slides were incubated with both α-rabbit

(Alexa Fluor 546) and α-rat (Alexa Fluor 488) secondary antibodies (Molecular probe, US). Myc

and Cytokeratin-8/18 signals were detected using LEICA SP8.

Data preprocessing and normalization of bulk RNA-seq data

MCF7 cells, MCF7 cells treated with 10 nM E2 for 2 hours, MCF7 TDP2KO cells,

MCF7 TDP2KO cells treated with 10 nM E2 for 2 hours, and etoposide-treated MCF7 with or

without 10 nM E2 were subject to RNA-seq at the Cedars-Sinai Genomics Core. Biological

duplicates were obtained for all of the experimental groups. Our quality control pipeline

employed Illumina standards to assess base call quality and truncated low-quality reads. The

quality of sequencing data was evaluated using MultiQC software [52]. Sequence alignment and

quantification were performed using the STAR-RSEM pipeline [53, 54]. Reads overlapping

exons in the annotation of Genome Reference Consortium Human Build 38 (GRCh38) will be

identified. Genes failing to achieve raw read counts of at least two across all the libraries were

filtered out and excluded from downstream analysis. Batch effects were corrected by ComBat

[55]. The trimmed mean of the M-values normalization method (TMM) [56] (version 1.6.1) was

used for calculating normalized count data.

Differential expression analysis of bulk RNA-seq data

For the two comparisons - 1) E2 treated versus untreated in MCF7 cells and 2) E2 treated

versus untreated in MCF7 TDP2KO cells - we employed the integrated hypothesis testing

method, as previously reported [57]. Briefly, for each gene, two P-values were computed by

conducting a two-tailed Student's T-test and a two-tailed median different test using the

empirical distributions that were estimated by random permutations of the samples. We then

combined two P-values into a single combined P-value using Stouffer's method. Multiple testing

correction was done by Storey's correction method [58]. Finally, the differentially expressed

genes were selected with the ones having a combined p-value < 0.05 and a fold-change ≥ 1.5.

The Gene Set Enrichment Analysis (GSEA) was employed to assess the enrichment of gene sets

[28].

Single-cell RNA-seq data preprocessing and differential expression analysis

Single-cell RNA-seq was performed using the 10X Genomics scRNA-seq approach at the

Cedars-Sinai Genomics Core. The Cell Ranger Single-Cell Software Suite was used for

demultiplexing, barcode assignment, and quantification of unique molecular identifiers (UMIs).

Reads were aligned to the hg38 reference genome using a pre-built annotation package obtained

from the 10X Genomics website. All lanes per sample were processed using the 'cellranger

count' function. The output from different lanes was then aggregated using 'cellranger aggr' with

–normalise set to 'none.' Quality control and analysis were conducted using the Seurat packages

in R (version 3.2.2) [30]. Poor-quality cells, defined by the number of detected genes less than

2,000, were filtered out. In addition, all cells with 11.89% (three standard deviations) or more of

UMIs mapping to mitochondrial genes were considered non-viable or apoptotic and excluded

from the analysis. Genes detected in at least three cells were retained for further analysis,

resulting in a total of 19,043 genes for the analysis. To ensure no index swapping occurred, cells

with unique barcodes that appeared in more than one sample (non-unique barcodes) as 'NULL'

were excluded for further analysis. Gene expression values were normalized through log

normalization using the "NormalizeData" function in Seurat. We performed two comparisons: 1)

E2 treated versus untreated in MCF7 cells and 2) E2 treated versus untreated in MCF7 TDP2KO

cells. The "FindMarkers" function in Seurat was used with filtered data to identify differentially

expressed genes between cell groups. We called differentially expressed genes with adjusted p-

value <0.05 and log fold change ≥ 0.1 (1.11 fold). The fold change threshold was determined at

the 95 percentile value of the empirical null distribution of fold changes, which was generated

through 1,000 times random permutation of the cells.

Statistical Analysis

We conducted Principal Component Analysis (PCA) to visualize the samples and assessed the

distribution of the samples based on their gene expression profile. We used MATLAB (v.9.0;

Mathworks, Natick, MA, USA), R (v.3.5), and Python (v.3.7) for bioinformatic analysis.

Acknowledgment

We thank Cedars-Sinai Cancer, Applied Genomics, Computation and Translational Core and

Flow Core for technical support. This work is supported by Cedars-Sinai CTSI Clinical Scholar

Award (to N.M.), NIHR01CA149385, Cedars-Sinai Cancer Biology Development Fund and

Institutional Support (to H.T.), Leukemia & Lymphoma Society (to N.J.) and Margie and Robert

E. Petersen Foundation, The Fashion Footwear Charitable Foundation of New York, Inc.,

Avon Foundation and Associates for Breast and Prostate Cancer Studies (to A.G.).

Data availability

Bulk and single-cell RNA-seq data are available in Gene Expression Omnibus (accession #

GSE225423).

Author Contributions

Conceptualization, N.M., M.K., S.T., S.Y.Y., and H.T.; Data collection, N.M., N.J., R.A.M., J.A,

R.S., and H.T.; Data analysis and interpretation, N.M., M.K., R.A.M., S.Y.Y., and H.T.;

Materials, F.C-L., S.Y., X.C.; Writing the manuscript, N.M., M.K., S.T., S.Y.Y., and H.T.;

Funding Acquisition, N.M., A.E.G., and H.T.

References

462 1   Katzenellenbogen BS, Katzenellenbogen JA. Estrogen receptor transcription and
463   transactivation: Estrogen receptor alpha and estrogen receptor beta: regulation by
464   selective estrogen receptor modulators and importance in breast cancer. *Breast Cancer*
465   *Res* 2000; 2: 335-344.

466
467 2   Fuentes N, Silveyra P. Estrogen receptor signaling mechanisms. *Adv Protein Chem Struct*
468   *Biol* 2019; 116: 135-170.

469
470 3   Ju BG, Lunyak VV, Perissi V, Garcia-Bassets I, Rose DW, Glass CK *et al*. A
471   topoisomerase IIbeta-mediated dsDNA break required for regulated transcription. *Science*
472   2006; 312: 1798-1802.

473
474 4   Hah N, Danko CG, Core L, Waterfall JJ, Siepel A, Lis JT *et al*. A rapid, extensive, and
475   transient transcriptional response to estrogen signaling in breast cancer cells. *Cell* 2011;
476   145: 622-634.

477
478 5   Pommier Y, Sun Y, Huang SN, Nitiss JL. Roles of eukaryotic topoisomerases in
479   transcription, replication and genomic stability. *Nat Rev Mol Cell Biol* 2016; 17: 703-721.

480
481 6   Morimoto S, Tsuda M, Bunch H, Sasanuma H, Austin C, Takeda S. Type II DNA
482   Topoisomerases Cause Spontaneous Double-Strand Breaks in Genomic DNA. *Genes*
483   *(Basel)* 2019; 10.

484
485 7   Tubbs A, Nussenzweig A. Endogenous DNA Damage as a Source of Genomic Instability
486   in Cancer. *Cell* 2017; 168: 644-656.

487
8   King IF, Yandava CN, Mabb AM, Hsiao JS, Huang H-S, Pearson BL *et al*.
Topoisomerases facilitate transcription of long genes linked to autism. *Nature* 2013; 501:
58-62.

9   Bunch H, Lawney BP, Lin Y-F, Asaithamby A, Murshid A, Wang YE *et al*.
Transcriptional elongation requires DNA break-induced signalling. *Nature*
Communications 2015; 6: 10191.

10   Nitiss JL. Targeting DNA topoisomerase II in cancer chemotherapy. *Nat Rev Cancer*
2009; 9: 338-350.

11   Madabhushi R, Gao F, Pfenning AR, Pan L, Yamakawa S, Seo J *et al*. Activity-Induced
DNA Breaks Govern the Expression of Neuronal Early-Response Genes. *Cell* 2015; 161:
1592-1605.

12   Calderwood SK. A critical role for topoisomerase IIb and DNA double strand breaks in
transcription. *Transcription* 2016; 7: 75-83.

13   Nitiss JL. DNA topoisomerase II and its growing repertoire of biological functions.
Nature Reviews Cancer 2009; 9: 327-337.

14   Manville CM, Smith K, Sondka Z, Rance H, Cockell S, Cowell IG *et al*. Genome-wide
ChIP-seq analysis of human TOP2B occupancy in MCF7 breast cancer epithelial cells.
Biol Open 2015; 4: 1436-1447.

15   Ledesma FC, El Khamisy SF, Zuma MC, Osborn K, Caldecott KW. A human 5′-tyrosyl
DNA phosphodiesterase that repairs topoisomerase-mediated DNA damage. *Nature*
2009; 461: 674-678.

16   Hoa Nguyen N, Shimizu T, Zhou Zhong W, Wang Z-Q, Deshpande Rajashree A, Paull
Tanya T *et al*. Mre11 Is Essential for the Removal of Lethal Topoisomerase 2 Covalent
Cleavage Complexes. *Molecular Cell* 2016; 64: 580-592.

17   Sasanuma H, Tsuda M, Morimoto S, Saha LK, Rahman MM, Kiyooka Y *et al*. BRCA1
ensures genome integrity by eliminating estrogen-induced pathological topoisomerase II–
DNA complexes. *Proceedings of the National Academy of Sciences* 2018; 115: E10642-
E10651.

18   Gómez-Herreros F, Romero-Granados R, Zeng Z, Alvarez-Quilón A, Quintero C, Ju L *et*
al. TDP2-dependent non-homologous end-joining protects against topoisomerase II-
induced DNA breaks and genome instability in cells and in vivo. *PLoS Genet* 2013; 9:
e1003226.

19   Aparicio T, Baer R, Gottesman M, Gautier J. MRN, CtIP, and BRCA1 mediate repair of
topoisomerase II–DNA adducts. *Journal of Cell Biology* 2016; 212: 399-408.

533
534   20   Itou J, Takahashi R, Sasanuma H, Tsuda M, Morimoto S, Matsumoto Y *et al*. Estrogen
535   Induces Mammary Ductal Dysplasia via the Upregulation of Myc Expression in a DNA-
536   Repair-Deficient Condition. *iScience* 2020; 23: 100821-100821.

537
538   21   Shang Y, Hu X, DiRenzo J, Lazar MA, Brown M. Cofactor dynamics and sufficiency in
539   estrogen receptor-regulated transcription. *Cell* 2000; 103: 843-852.

540
541   22   Ishida R, Miki T, Narita T, Yui R, Sato M, Utsumi KR *et al*. Inhibition of intracellular
542   topoisomerase II by antitumor bis(2,6-dioxopiperazine) derivatives: mode of cell growth
543   inhibition distinct from that of cleavable complex-forming type inhibitors. *Cancer Res*
544   1991; 51: 4909-4916.

545
546   23   Skoufias DA, Lacroix FB, Andreassen PR, Wilson L, Margolis RL. Inhibition of DNA
547   decatenation, but not DNA damage, arrests cells at metaphase. *Mol Cell* 2004; 15: 977-
548   990.

549
550   24   Mao Y, Desai SD, Ting CY, Hwang J, Liu LF. 26 S proteasome-mediated degradation of
551   topoisomerase II cleavable complexes. *J Biol Chem* 2001; 276: 40652-40658.

552
553   25   Schellenberg MJ, Lieberman JA, Herrero-Ruiz A, Butler LR, Williams JG, Munoz-
554   Cabello AM *et al*. ZATT (ZNF451)-mediated resolution of topoisomerase 2 DNA-
555   protein cross-links. *Science* 2017; 357: 1412-1416.

556
557   26   Zhao Y, Thomas HD, Batey MA, Cowell IG, Richardson CJ, Griffin RJ *et al*. Preclinical
558   evaluation of a potent novel DNA-dependent protein kinase inhibitor NU7441. *Cancer*
559   *Res* 2006; 66: 5354-5362.

560
561   27   Grandori C, Cowley SM, James LP, Eisenman RN. The Myc/Max/Mad network and the
562   transcriptional control of cell behavior. *Annu Rev Cell Dev Biol* 2000; 16: 653-699.

563
564   28   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA *et al*. Gene
565   set enrichment analysis: A knowledge-based approach for interpreting genome-wide
566   expression profiles. *Proceedings of the National Academy of Sciences* 2005; 102: 15545-
567   15550.

568
29   Bild AH, Yao G, Chang JT, Wang Q, Potti A, Chasse D *et al*. Oncogenic pathway
signatures in human cancers as a guide to targeted therapies. *Nature* 2006; 439: 353-357.

30   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd *et al*.
Comprehensive Integration of Single-Cell Data. *Cell* 2019; 177: 1888-1902 e1821.

31   Hsiao CJ, Tung P, Blischak JD, Burnett JE, Barr KA, Dey KK *et al*. Characterizing and
inferring quantitative cell cycle phase in single-cell RNA-seq data analysis. *Genome Res*
2020; 30: 611-621.

32   Nestorowa S, Hamey FK, Pijuan Sala B, Diamanti E, Shepherd M, Laurenti E *et al*. A
single-cell resolution map of mouse hematopoietic stem and progenitor cell
differentiation. *Blood* 2016; 128: e20-31.

33   Cheung E, Kraus WL. Genomic Analyses of Hormone Signaling and Gene Regulation.
Annual Review of Physiology 2010; 72: 191-218.

34   Jagannathan V, Robinson-Rechavi M. Meta-analysis of estrogen response in MCF-7
distinguishes early target genes involved in signaling and cell proliferation from later
target genes involved in cell cycle and DNA repair. *BMC Syst Biol* 2011; 5: 138.

35   Escot C, Theillet C, Lidereau R, Spyratos F, Champeme MH, Gest J *et al*. Genetic
alteration of the c-myc protooncogene (MYC) in human primary breast carcinomas. *Proc*
Natl Acad Sci U S A 1986; 83: 4834-4838.

36   Sicinski P, Donaher JL, Parker SB, Li T, Fazeli A, Gardner H *et al*. Cyclin D1 provides a
link between development and oncogenesis in the retina and breast. *Cell* 1995; 82: 621-
630.

37   Weinstat-Saslow D, Merino MJ, Manrow RE, Lawrence JA, Bluth RF, Wittenbel KD *et*
al. Overexpression of cyclin D mRNA distinguishes invasive and in situ breast
carcinomas from non-malignant lesions. *Nat Med* 1995; 1: 1257-1260.

38   D'Cruz CM, Gunther EJ, Boxer RB, Hartman JL, Sintasath L, Moody SE *et al*. c-MYC
induces mammary tumorigenesis by means of a preferred pathway involving spontaneous
Kras2 mutations. *Nature Medicine* 2001; 7: 235-239.

39   Zhu D, Zhao Z, Cui G, Chang S, Hu L, See YX *et al*. Single-Cell Transcriptome Analysis
Reveals Estrogen Signaling Coordinately Augments One-Carbon, Polyamine, and Purine
Synthesis in Breast Cancer. *Cell Rep* 2018; 25: 2285-2298 e2284.

40   Harding SM, Boiarsky JA, Greenberg RA. ATM Dependent Silencing Links Nucleolar
Chromatin Reorganization to DNA Damage Recognition. *Cell Rep* 2015; 13: 251-259.

41   Larsen DH, Hari F, Clapperton JA, Gwerder M, Gutsche K, Altmeyer M *et al*. The
NBS1-Treacle complex controls ribosomal RNA transcription in response to DNA
damage. *Nat Cell Biol* 2014; 16: 792-803.

42   Shanbhag NM, Rafalska-Metcalf IU, Balane-Bolivar C, Janicki SM, Greenberg RA.
ATM-dependent chromatin changes silence transcription in cis to DNA double-strand
breaks. *Cell* 2010; 141: 970-981.

43   Bahr C, von Paleske L, Uslu VV, Remeseiro S, Takayama N, Ng SW *et al*. A Myc
enhancer cluster regulates normal and leukaemic haematopoietic stem cell hierarchies.
Nature 2018; 553: 515-520.

44   Ahmadiyeh N, Pomerantz MM, Grisanzio C, Herman P, Jia L, Almendro V *et al*. 8q24
prostate, breast, and colon cancer risk loci show tissue-specific long-range interaction
with MYC. *Proc Natl Acad Sci U S A* 2010; 107: 9742-9746.

45   Najnin RA, Al Mahmud MR, Rahman MM, Takeda S, Sasanuma H, Tanaka H *et al*.
ATM suppresses c-Myc overexpression in the mammary epithelium in response to
estrogen. *Cell Reports* 2023; 42: 111909.

46   Álvarez-Quilón A, Serrano-Benítez A, Ariel Lieberman J, Quintero C, Sánchez-Gutiérrez
D, Escudero LM *et al*. ATM specifically mediates repair of double-strand breaks with
blocked DNA ends. *Nature Communications* 2014; 5: 3347.

47   Schellenberg MJ, Appel CD, Riccio AA, Butler LR, Krahn JM, Liebermann JA *et al*.
Ubiquitin stimulated reversal of topoisomerase 2 DNA-protein crosslinks by TDP2.
Nucleic Acids Res 2020; 48: 6310-6325.

48   Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K *et al*. Genetic and
transcriptional evolution alters cancer cell line drug response. *Nature* 2018; 560: 325-330.

49   Risso D, Perraudeau F, Gribkova S, Dudoit S, Vert J-P. A general and flexible method
for signal extraction from single-cell RNA-seq data. *Nature Communications* 2018; 9:
284.

50   Roy PG, Pratt N, Purdie CA, Baker L, Ashfield A, Quinlan P *et al*. High CCND1
amplification identifies a group of poor prognosis women with estrogen receptor positive
breast cancer. *International Journal of Cancer* 2010; 127: 355-360.

51   Ortiz AB, Garcia D, Vicente Y, Palka M, Bellas C, Martin P. Prognostic significance of
cyclin D1 protein expression and gene amplification in invasive breast carcinoma. *PLOS*
ONE 2017; 12: e0188068.

52   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* 2011; 12: 323.

53   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S *et al*. STAR: ultrafast
universal RNA-seq aligner. *Bioinformatics* 2012; 29: 15-21.

54   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* 2016; 32: 3047-3048.

55   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
using empirical Bayes methods. *Biostatistics* 2006; 8: 118-127.

56   Robinson MD, Oshlack A. A scaling normalization method for differential expression
analysis of RNA-seq data. *Genome Biology* 2010; 11: R25.

57   You S, Yoo S-A, Choi S, Kim J-Y, Park S-J, Ji JD *et al*. Identification of key regulators
for the migration and invasion of rheumatoid synoviocytes through a systems approach.
Proceedings of the National Academy of Sciences 2014; 111: 550-555.

58   Storey JD, Tibshirani R. Statistical significance for genomewide studies. *Proceedings of*
the National Academy of Sciences 2003; 100: 9440-9445.

Figure Legends

Figure 1   Estrogen and TOP2-dependent induction of *MYC*

(A) Experiment design: MCF7 and T47D cells were treated with 10 nM 17β-estradiol for 1, 2,

and 6 hours. RNA was extracted at each time point for 3D PCR analysis, and bulk- and single-

cell RNA-seq was performed at the 2-hour time point.

(B) 3D PCR output from Analysis Suites. Blue dots represent wells only with amplification of

genes detected by FAM (*MYC*) probe, red dots represent wells only with the amplification

detected by VIC (*TBP* or *TFRC*) probe (internal control), yellow dots represent wells without

amplification, and green dots represent wells with the amplification of both probes. Normalized

MYC expression was calculated by the ratio of FAM (*MYC*) to VIC (*TBP* or *TFRC*).

(C) *MYC* induction in cells after 2 hours of treatment with 10 nM E2 relative to cells with no E2

in two ER-positive breast cancer cell lines (MCF7 and T47D) and an ER-negative cell line (MD-

MBA-231). Averages of three independent experiments are shown.

(D) Dynamic responses of *MYC* induction: Cells were treated with 10 nM E2 for 0, 1, 2, and 6

hours. Averages of two independent experiments are shown.

(E) A TOP2 catalytic inhibitor ICRF-193 Effects on *MYC* mRNA induction: MCF7 and T47D

cells were treated with 2 and 8 μM ICRF-193 for 1 hour, followed by the treatment with 10 nM

E2 for 2 hours. *MYC* induction decreased by ICRF-193 in both cell lines. Averages of three

independent experiments are shown. * $p<0.05$, ** $p<0.01$, Student's t-test.

Figure 2   TDP2 suppresses the volatile and prolonged induction of *MYC* by estrogen

(A) Western Blot Analysis displaying MCF7-1, T47D and MCF7-2 cell lines alongside TDP2

KO cells for each cell line.

(B) Etoposide sensitivity measured by colony formation assay for MCF7 and MCF7

TDP2KO(top) and T47D and T47D TDP2KO (bottom). Results from three independent

experiments. * $p<0.05$, ** $p<0.01$, Student's t-test

(C) Persistent *MYC* transcriptional responses: MCF7 and MCF7 TDP2 KO cells were treated

with 10 nM E2 for 1, 2, and 6 hours. Averages from two independent experiments.

(D) *MYC* transcriptional responses at 2 hours: MCF7-1, MCF7-2 and T47D, and their TDP2 KO

counterparts were treated with 10 nM E2 for 2 hours, and RNA was extracted for digital PCR

assays. Results from two independent experiments.

(E) NU7441 and etoposide increased the *MYC* transcriptional response. MCF7 cells, cultured

with or without NU7441 or etoposide for three hours were then treated with 10 nM E2 for 2

hours. Results from two independent experiments.

Figure 3   **The induction of MYC protein by E2** *in vivo*

(A) Representative images of mammary ducts: Shown are images of mammary ducts from wild-

type (left) and TDP2KO (right) mice, immunostained for MYC (red) and CK-8/18 (green)

proteins. The images were captured before injection (0 hours) and 6 and 20 hours after the

injection of E2.

(B) Box plots illustrating the fraction of MYC-positive cells: Each plot represents Each plot

represents the distribution of the fraction of MYC-positive cells among CK8/18-positive

mammary epithelial cells. MYC-positive epithelial cells were scored in 3 to 12 mammary ducts

in each mouse at each timepoint. p values were determined by t-test.

Figure 4   **Bulk and Single-cell RNA sequencing analysis of** *MYC* **transcriptional**

response by E2

(A) *MYC* transcriptional response to E2 (2 hours) in MCF7 and MCF7 TDP2KO cells, measured

by bulk RNA-seq and single-cell RNA-seq.

(B) Violin plots displaying *MYC* expression: Normalized *MYC* expression distributions in MCF7

and MCF7 TDP2KO cells with or without E2 (2 hours) from single-cell RNA-seq data.

(C) Histograms illustrating *MYC* expression distribution: Showing the skewed unimodal

distribution of MYC expression in MCF7 and MCF7 TDP2KO cells with or without E2 (2

hours), obtained from single-cell RNA-seq data.

(D) Violin plots of *MYC* expression at single-cell level: Depicting the normalized expression of

MYC in G1, S, and G2/M cell populations.

Figure 5   **Commonly upregulated genes between bulk- and single-cell RNA-seq**

(A) Venn diagrams showing the differentially expressed genes: The differentially expressed

genes by E2 treatment were presented for MCF7 and MCF7 TDP2KO cells. 32 genes in MCF7

and 62 genes in MCF7 TDP2KO are common between the two platforms. Significances of

overlap were assessed using Fisher's exact test.

(B) Bar plots showing the number of common differentially regulated genes s by E2 (2 hours):

28 and 4 genes were commonly up- and down-regulated in E2-treated MCF7 compared to

untreated MCF7. Similarly, 58 and 4 genes were commonly up- and downregulated in E2 treated

MCF7 TDP2KO compared to untreated MCF7 TDP2KO.

(C) Venn-diagram and a Table of commonly differentially expressed genes: Depicting 32 and 62

commonly differentially regulated genes from bulk and single cell RNA-seq data, along with the

table highlighting 8 overlaping genes. Significance of overlap was determined using Fisher's

exact test.

Figure 1 Manguso et al,

Figure 2 Manguso et al,

Figure 3, Manguso et al,

A

Wild-type   TDP2⁻/⁻

0 h

6 h

20 h

DNA/CK-8/18/MYC

B

Figure 4, Manguso et al,

A

Comparison	Bulk RNA-seq		Single-cell RNA-seq	
	log2-FC	P-value	log-FC	P-value
MCF7 with E2 vs without E2	0.9760	0.0194	0.1722	<0.0001
MCF7 TDP2KO with E2 vs without E2	1.6595	0.0001	0.4267	<0.0001

Figure 5, Manguso et al,

A

Bulk RNA-seq
MCF7 treated vs. untreated
(2021)

scRNA-seq
MCF7 treated vs. untreated
(111)

1989   32   79

P<0.0001

Bulk RNA-seq
TDP2KO treated vs. untreated
(1034)

scRNA-seq
TDP2KO treated vs. untreated
(327)

972   62   265

P<0.0001

B

■ up   ■ down

number of DEGs

MCF7   MCF7 TDP2KO

C

MCF7 common
treated vs. untreated
(32)

24   8   54

P<0.0001   TDP2KO common
treated vs. untreated
(62)

| | **Bulk RNA-seq** | | | | **scRNA-seq** | | | |
| | MCF7 | | MCF7 TDP2KO | | MCF7 | | MCF7 TDP2KO | |
	log FC	p value	log FC	p value	log FC	p value	log FC	p value
MYC	0.98	0.02	1.66	<0.01	0.17	3.8E-06	0.43	1.3E-41
CCND1	0.61	0.01	0.81	0.01	0.27	4E-15	0.35	1.2E-35
NRIP1	1.61	<0.01	1.97	<0.01	0.15	9E-06	0.19	2.7E-24
SH3BP5	0.64	0.01	0.61	0.01	0.14	0.01	0.25	5.6E-24
IGFBP4	1.27	<0.01	0.81	0.04	0.14	<0.01	0.39	3.5E-39
RET	1.45	0.01	0.99	0.03	0.18	5.4E-16	0.16	2.4E-13
NBPF1	0.86	0.02	0.90	0.02	0.15	3E-07	0.14	4.1E-08
CA12	1.21	0.01	0.62	0.05	0.24	2.3E-15	0.58	2.9E-77