The reef-building coral *Galaxea fascicularis*: a new model system for coral

symbiosis

Giulia Puntin[1], Jamie Craggs[2,3], Róisín Hayden[4], Kara E. Engelhardt[1], Shelby McIlroy[4], Michael

Sweet[3,5], David M. Baker[4], Maren Ziegler[1,]*

[1]Marine Holobiomics Lab, Department of Animal Ecology & Systematics, Justus Liebig

University Giessen, Heinrich-Buff-Ring 26-32 IFZ, 35392 Giessen, Germany

[2]Horniman Museum and Gardens, Forest Hill, London, SE23 3PQ, United Kingdom.

[3]Coral Spawning Lab, Unit 6, Midas Metro Centre, 193 Garth Rd, Morden, SM4 4NE, United

Kingdom

[4]The Swire Institute of Marine Science, University of Hong Kong, Cape D'Aguilar, Hong Kong

[5]Aquatic Research Facility, Environmental Sustainability Research Centre, University of Derby,

DE22 1GB, United Kingdom

*Corresponding author: maren.ziegler@bio.uni-giessen.de

Abstract

Reef-building corals owe their evolutionary success to their symbiosis with unicellular algae (Symbiodiniaceae). However, increasingly frequent heat waves lead to coral mass-bleaching events and pose a serious threat to the survival of reef ecosystems. Despite significant efforts, a mechanistic understanding of coral-algal symbiosis functioning, what leads to its breakdown and what can prevent it, remains incomplete. The main obstacles are low amenability of corals to experimental handling and, owing to its obligatory nature, the difficulties of manipulating the coral-algal association. Indeed, many studies on the symbiotic partnership are conducted on other cnidarian model organisms and their results may therefore not be fully transferable to tropical reef-building corals. Here, we identify the tropical stony coral species *Galaxea fascicularis* as a novel candidate coral model system. Individual polyps of this species can be separated, enabling highly replicated genotype studies, and are well suited to experimental investigation of the symbiosis as they can be easily and effectively rid of their algal symbionts (bleached). We show that bleached adult individuals can reestablish symbiosis with homologous and heterologous symbionts. We also report the completion of the gametogenic cycle *ex-situ* and the successful spawning in aquaria over multiple years. These achievements help overcome several of the major limitations to direct research on corals and highlight the potential of *G. fascicularis* as an important new model system for investigations of symbiosis functioning and manipulation.

Keywords: Model organisms; reef-building corals; menthol bleaching; symbiosis manipulation; *ex situ* sexual reproduction

Introduction

Reef-building corals form obligatory endosymbiotic association with unicellular algae of the family Symbiodiniaceae. This association is key to their evolutionary success, but it is also at the heart of corals' susceptibility to global climate change, which manifests in coral bleaching - the breakdown of the coral-algal symbiosis (Dietzel et al. 2020). Bleaching is mostly driven by marine heatwaves which are predicted to worsen, causing the loss of virtually all coral reefs by the end of this century (van Hooidonk et al. 2016).

The interaction between heat stress and bleaching has been studied for more than 30 years (Gates et al. 1992; McLachlan et al. 2020), but a detailed understanding of the mechanisms underlying coral bleaching is still missing. Progress in coral symbiosis research is hampered by two main aspects. First, corals are challenging to maintain in aquarium settings as they tolerate only a narrow range of environmental conditions. Second, the obligatory nature of the coral-algal association makes it particularly challenging to physically and functionally separate the partners, an approach often necessary to unravel the mechanisms underlying symbiosis breakdown and what could prevent it (Weis et al. 2008; Voolstra 2013).

Model organisms are integral for understanding fundamental biological principles and symbiotic cnidarians have successfully been used as "coral models", advancing our understanding of coral bleaching and holobiont functioning (Weis et al. 2008). These model organisms share important traits with corals, such as being cnidarians that associate with microalgae, yet they also lack other features that normally represent obstacles to research work such as the calcium carbonate skeleton. In addition, the ability to study the animal host and its algal symbiont in isolation is more easily achieved with facultatively symbiotic organisms such as *Hydra* spp., Aiptasia (*Exaiptasia diaphana*), and *Astrangia* spp. (Dimond and Carrington 2007; Weis et al. 2008; Galliot 2012).

Cnidarian model organisms allow the study of basal or shared traits or phenomena at a speed that

would not be possible with reef-building corals, but testing on the latter remains necessary for the

understanding of coral-specific or ecologically relevant aspects. The obligatory symbiosis with

Symbiodiniaceae in reef-building corals has important biological implications (Falkowski et al.

1984; Hoegh-Guldberg 1999), similar to the calcification process that is deeply intertwined with

coral physiology and ecology (Gattuso et al. 1999). It is therefore important to establish a "true"

tropical reef-building coral model species.

The richness of the scleractinian taxon offers a vast array of species to choose from in the quest

for a suitable candidate coral model species. These can be evaluated for their tractability (as

amenability to experimental work) considering several aspects which, as proposed by Puntin et al.

(2022) should include: 1) pre-existing knowledge: baseline information of the organism's biology

is necessary to interpret and contextualize results; 2) compatibility with aquarium rearing: the

possibility to maintain the organism and preferably to complete its life cycle in the lab is essential

to increase its availability, reduce confounding effects such as unknown life history, improve

reproducibility, and lower the pressure on threatened wild populations; 3) amenability to symbiosis

manipulations is necessary to help unravel complex coral-algal functional interactions, and can be

broken down into the organism's suitability to be rendered aposymbiotic (bleached), to be

maintained aposymbiotic for a certain time, and to then be re-infected with Symbiodiniaceae.

We have identified the coral species *Galaxea fascicularis* (Linnaeus, 1767) (clade: Complexa,

family: Euphyllidae) as a promising candidate coral model system. The species is relatively

common across the Indo-Pacific region (Veron et al. 2016), where the phylogeny of the genus

Galaxea is geographically well resolved (Wepfer et al. 2020b). It is also one of the more commonly

used species in heat-stress experiments (McLachlan et al. 2020) and in studies on coral

calcification (e.g., Marshall and Clode 2004; Al-Horani 2005). Importantly, other community resources such as an annotated draft genome are available (Liew et al. 2016; Niu et al. 2016, http://www.gfas.reefgenomics.org/). In addition, *G. fascicularis* is known for its tolerance to stressors as it is "largely unaffected by bleaching" in the field (Marshall and Baird 2000) and for being easy to grow and propagate in aquaria (Pavia and Estacion 2019). Further, its morphology provides additional practical advantages with relatively big polyps, which can be easily isolated (Al-Horani 2005). This conveniently allows reduction of the complexity of the organism from colonial to individual, enables high clonal replication, and facilitates visualization (Al-Horani et al. 2003; Marshall and Clode 2004). Its association with Symbiodiniaceae is relatively well characterized (Huang et al. 2011; Wepfer et al. 2020a). Furthermore, the species' reproductive mode and spawning patterns in the wild are well documented (Babcock et al. 1986; Harrison 1988; Keshavmurthy et al. 2012). However, its amenability to symbiosis manipulation, to experimental handling of bleached individuals, and the ability to complete gametogenic cycles and spawning in closed aquaria has not yet been described.

The overarching aim of this study was to explore the potential of *Galaxea fascicularis* as a coral model. Specifically, we investigated: 1) the feasibility to render polyps aposymbiotic by eliminating Symbiodiniaceae (bleaching), 2) the amenability to experimental manipulation of symbiotic and aposymbiotic polyps in a simplified system, 3) the re-establishment of the symbiosis with various species of Symbiodiniaceae, and 4) the possibility to induce full gametogenic cycles *ex situ*, with subsequent spawning under aquarium conditions. These aspects are central to overcoming the two main obstacles to coral research work: low tractability of corals and their complex nature.

Materials and methods

Coral collection and long-term aquarium rearing

Colonies of *Galaxea fascicularis* were collected from three geographical locations: the Red Sea, Hong Kong, and the Great Barrier Reef. Red Sea colonies were collected at 9-13 m depth at the North-Eastern protected end of the reef "Al Fahal" in the central Saudi Arabian Red Sea (22.3054°, 38.9655°) in March 2019, and transported to the Ocean2100 aquarium facility at Justus Liebig University Giessen (Germany) where they are part of the live collection (CITES permit 19-SA-000096-PD). In the aquarium system, light was provided by white and blue fluorescent lamps with a light:dark cycle of 12:12 h at 130 -160 µmol photons m^{-2} s^{-1}, which approximates light condition at the collection site (Ziegler et al. 2015). Salinity was maintained around 35 and temperature at 26 °C. Colonies were fed daily with a combination of frozen copepods, *Artemia*, krill, and *Mysis*. Hong Kong colonies were collected from 5 m (max. depth) from Crescent Island (22.5308°, 114.3150°) in November 2020 and transported to the University of Hong Kong, where they were fragmented and acclimated to aquarium conditions for one month. The aquarium system consisted of 6-L acrylic tanks placed within a Plant Growth Chamber (Panasonic MLR-352H-PA). Each tank held eight ~3 cm-nubbins and was fitted with a submersible pump (Atman AT301) and a small water filter (Shiruba PF120). The fragments used in symbiosis reestablishment were fed twice-weekly with a powdered blend of marine plankton (ReefRoids, Polyp Lab) followed by a partial water change. Light, salinity, and temperature conditions were consistent with those maintained in the Ocean2100 facility.

For *ex situ* spawning, 35 colonies were collected from Arlington Reef, Great Barrier Reef, Australia (-16.7000°, 146.0500°) in September 2019, and transported to the Horniman Museum and Gardens (United Kingdom; CITES permit 585319/01). Colonies were distributed into four

coral spawning systems that replicated the natural environmental parameters (seasonal

temperature, photoperiod, solar irradiance, and lunar cycle) required to stimulate reproduction.

Aquarium design and coral husbandry protocol was based on the mesocosms described by Craggs

et al. (2017), and the seasonal temperature profile were based on a non-sequential eight year

average (1998 - 2017) from Moore Reef (-16.8667°, 146.2334°')

(http://data.aims.gov.au/aimsrtds/datatool.xhtml?site=931¶m=water%20temperature).

Preparation of single polyps for experimentation

135 For the menthol bleaching and thermal performance experiment, we used single polyps from three

136 Red Sea colonies each showing distinct coloration (Fig. S1). Replicate clonal polyps were isolated

137 from each colony using an electric rotary cutter, mounted on coral glue (Grotech, CoraFix

138 SuperFast), and allowed to heal for two weeks in the Ocean2100 aquarium facility (Schubert and

139 Wilke 2018).

Menthol bleaching to remove algal symbionts

An equal number of polyps from each colony were randomly assigned to the control group

(maintained in a symbiotic state), and the bleached group (chemically bleached with menthol).

Menthol bleaching followed a protocol modified from Wang et al. (2012) and consisted of three

days of treatment in 0.38 mM menthol solution in seawater, followed by a day of rest and a fourth

day of menthol treatment. Each day the polyps were incubated in menthol for 8 h under light and

stirring (Fig. S2). After, the polys were rinsed and kept in clean containers with air bubbling and

stirring. During menthol treatment, the treated polyps were only exposed to filtered (1.2 μm)

artificial seawater (FASW) to prevent exposure to Symbiodiniaceae. Following, all polyps

(including, for consistency, the symbiotic group) were maintained in FASW.

To confirm bleaching, half of the polyps from each group (bleached and symbiotic) and each

colony were visually assessed with a fluorescence stereomicroscope (Leica MZ16 F) 10 days after

the termination of the menthol treatment. Representative pictures were taken under natural light

(brightfield, light source: 2950 K) and under UV light with GF2 filters allowing visualization of

the green fluorescent proteins and chlorophyll (Fig. 1a). Following the same bleaching protocol,

menthol-bleached polyps were also documented using a compound epifluorescence microscope to

reach higher resolution (Leica DM 5500B, TX2 filter) (Fig. 1b).

Thermal performance experimental design

Symbiotic and bleached polyps were moved to the experimental tanks on the day after the last

menthol exposure and allowed to recover and acclimate for 10 days. The experimental system

consisted of eight 5-L glass tanks (20 cm × 30 cm) equipped with a small pump (Resun SP-500)

in a temperature-controlled water bath (26 ˚C). White fluorescent lamps were used to maintain the

light cycle and intensity consistent with long-term rearing conditions. Bleached and symbiotic

polyps were kept in separate tanks (four per group) filled with FASW, with three to six polyps

each (Fig. S3). Polyps were fed each day after the end of the dark incubation, followed by 10 %

water change after 2-3 h. Each polyp was fed one small frozen adult Artemia shrimp pipetted

directly on top of the oral opening. The polyps were cleaned every two to three days after the

incubations to remove fouling that would interfere with respirometry measurements.

To test the amenability to experimental handling of single polyps and to compare metabolic rates

of bleached and symbiotic polyps, we conducted a thermal performance curve experiment.

Photosynthesis and respiration of *G. fascicularis* polyps were measured across 12 ˚C temperature

over 10 days. For this, the polyps were acclimated to a different temperature each day, in the

following order: 20, 22, 23, 24, 25, 26, 27, 28, 30 to 32 °C. The temperature of the water bath was

changed overnight, and incubations took place the subsequent day at the respective temperature.

Photosynthesis and respiration of symbiotic and bleached polyps

We measured oxygen evolution of individual *G. fascicularis* polyps in light and dark incubations

to calculate net photosynthesis (PN) and dark respiration (R) respectively. For these measurements

four bleached and four symbiotic polyps from three colonies were used (n = 12). Every day, each

polyp was placed in individual 160 mL glass incubation jars (Weck, Germany), equipped with a

magnetic stirrer (~5 cm s^{-1}). Three additional jars containing only seawater and stirring bars were

used to control background biological activity. Light intensity of 230 - 280 µmol photons m^{-2} s^{-1}

during light incubations (ATI 80W Aquablue Special) and temperatures were maintained through

thermostat-controlled water baths placed on a custom-made multiplate stirring system (Rades et

al. 2022).

Dissolved oxygen (DO) was measured at the start and at the end of each incubation using a

handheld multiparameter probe (WTW Multi 3620 IDS set). Light incubations lasted 2 h around

mid-day followed by 1.5 h of dark incubation. Net photosynthesis and respiration were calculated

using the following formula (Schneider and Erez 2006):

$$PN \text{ or } R[mgO_2 \times cm^{-2} \times h^{-1}] = \frac{\Delta O_2[mg \times L^{-1}] \times V_{incubation}[L]}{T[h] \times SA[cm^2]}$$

189   Where ΔO_2 is the difference in dissolved oxygen between the end and the start of the incubation

190   (DO_{end} - DO_{start}) corrected for the controls' ΔO_2. This is multiplied by the volume of the incubation

191   chambers in L ($V_{incubation}$), and divided by incubation time (T) in hours and polyp surface area (SA)

192   in cm^2.

Surface area measurements

Polyp surface area was measured through photogrammetry. Between 40 to 50 pictures of each polyp were taken with a phone camera two days before the beginning of the incubations. These were used to create 3D models with 3DF Zephyr Free (v.4.523), cleaned on Artec3D (Studio 11 Professional v.11.2.2.16), and loaded on MeshLab (v.2016.12, Cignoni et al. 2008) for size scaling and calculation of live coral surface area.

Thermal performance data analysis

All analyses were conducted in the R statistical environment (v.4.1.0, R Core Team 2021) and plotted with ggplot2 (v.3.3.5, Wickham 2016). The effects of symbiotic state ('symbiotic' vs. 'bleached') or colony identity on net photosynthesis (PN) and respiration (R) were analyzed using linear mixed-effect models. The effect of symbiotic state on R was tested by setting 'symbiotic state' as a fixed factor and 'polyp identity', 'colony', and 'temperature' as random factors. The effect of colony identity on PN and R was tested considering light and dark incubations separately and setting 'colony identity' as fixed factor and 'polyp identity', 'temperature', and 'symbiotic state' as random factors. The package 'lmerTest' (v.3.1-3, Kuznetsova et al. 2017) was used to construct the models and calculate p-values. The package 'performance' was used to check the residuals of fixed and random factors and to compare alternative models (v.0.7.3, Lüdecke et al. 2021). Differences between colonies were tested as pairwise comparisons of estimated marginal means (Searle et al. 1980) with Bonferroni correction with the 'emmeans' package (v.1.6.2-1).

Symbiosis reestablishment after bleaching

We explored the possibility of returning polyps to the symbiotic state after menthol bleaching using three approaches. The first and simplest approach consisted of testing natural symbiont acquisition. One month after the menthol treatment, bleached polyps were placed next to symbiotic

colonies of *G. fascicularis* to expose them to Symbiodiniaceae naturally released from conspecifics

into the water column. The second approach consisted of inoculating bleached polyps with freshly

isolated Symbiodiniaceae from symbiotic *G. fascicularis*. For this, a donor polyp from each colony

was placed upside-down inside a 1.5 mL tube and centrifuged (Heraeus Biofuge Pico, England) at

8,300 *g* for 1 min to detach the tissue from the skeleton (Wilkinson et al. 2016). The slurry was

homogenized with 600 µl of FASW, centrifuged at 3,400 *g* for 1 min, and the resulting pellet

resuspended twice to clean the algal fraction. This Symbiodiniaceae suspension was used to

inoculate two to three polyps each with Symbiodiniaceae from the same colony or from one

respective other colony. For the inoculation, bleached polyps were placed in individual beakers,

the Symbiodiniaceae inoculum was pipetted on the oral opening together with one Artemia shrimp,

and allowed to feed for 3 h. Subsequently, inoculated polyps were transferred to the long-term

rearing aquarium facility. Symbiosis reestablishment was visually assessed after one month

(Fig. 3).

The third approach consisted of inoculating menthol-bleached coral nubbins with cultured

Symbiodiniaceae. Two species of symbionts, *Cladocopium goreaui* and *Durusdinium trenchii*

were chosen for targeted reinfection of the bleached fragments, as species from both these genera

are known to form stable symbiosis with *G. fascicularis* across the South China Sea (Tong et al.

2017). Batch cultures of *C. goreaui* (AIMS-SCF-055, isolated from *Acropora tenuis*) and *D.*

trenchii (AIMS-SCF-088, isolated from *A. muricata*) were obtained from the Symbiont Culture

Facility at the Australian Institute of Marine Science and maintained in f/2 media (Guillard 1975)

under the same conditions as the corals. A total of 16 fragments were inoculated with a

concentration of 200 symbiont cells ml^{-1} in each 6-L aquarium. The concentration of symbionts

was divided in a 90:10 ratio (180 cells ml^{-1} *D. trenchii* : 20 cells ml^{-1} *C. goreaui*). This ratio was

chosen because *G. fascicularis* is more commonly associated with symbionts from the genus

Cladocopium under the temperature conditions of this experiment (Dong et al. 2009; Zhou et al.

2017b). The symbiont aliquots were centrifuged to remove the culture media, washed once with

FASW before final resuspension in 10 mL FASW. The bleached fragments were first fed with

ReefRoids before being exposed to the chosen symbionts. For this, a suspension of ReefRoids in

ASW was pipetted directly into the oral opening of each polyp. The same process was then

immediately repeated with the suspended symbionts. Aquarium filters were turned off for 1 h to

allow the corals to feed and to prevent removal of symbiont cells from the water column.

Fragments were inoculated twice a week for 6 weeks, until symbiosis reestablishment was visually

confirmed. Symbiodiniaceae populations *in hospite* were then characterized with fluorescent in-

situ hybridization (FISH) and flow cytometry (McIlroy et al. 2020).

Fluorescent In-Situ Hybridisation (FISH)

After 6 weeks of inoculation, a single polyp was removed from each fragment with a chisel. Tissue

was removed from the skeleton with an airbrush containing deionized water. The resulting slurry

was homogenized using a 20-µl pipette tip. 1.5 mL of homogenate was transferred to a 2-mL

Eppendorf tube and centrifuged at 500 rpm for seven minutes, after which the supernatant was

discarded. The remaining pellet was washed once before being preserved in 5X SET buffer and

stored at -80 °C for further analysis.

Samples were prepared following an adapted protocol from McIlroy et al. (2020). Briefly, after

washing with 5X SET in IGEPAL at 0.4 and 0.1 % vol/vol samples were split equally between

three 1.5-mL black Eppendorf tubes. Each sample had one tube treated with a SymC probe, one

tube with a SymD probe and one without any probe as a control. Both SymC and SymD probes

were designed to have genus-level specificity for *Cladocopium* and *Durusdinium*, respectively.

After overnight hybridization at 45 °C (5X SET, 0.1 % vol/vol IGE- PAL, 10 % vol/vol formamide

and 100 pmol of the relevant probe), samples were washed once in warm 1X SET, vortexed and

resuspended in 500 µl of 1X SET for same-day cytometric analysis on a BD FACSAria Fusion

flow cytometer (BD Biosciences, CA). FlowJo™ Software (v.10.8.1, BD Life Sciences) was used

to determine the percentage of probe-positive cells (either *Cladocopium* or *Durusdinium*), to

identify a) the presence and b) the relative abundance of each algal genus within the sample.

Completing gametogenic cycles and *ex-situ* spawning

Galaxea fascicularis in the GBR spawns between October and December, 0.05-200 minutes after

sunset (MAS) and 1 – 8 nights after full moon (NAFM) (Babcock et al. 1986; Baird et al. 2021).

To determine if the *ex situ* reproductive pattern remained in synchrony with the wild we followed

our colonies over three consecutive reproductive seasons (2019 – 2021).

To ascertain gamete development in each colony, two months prior to predicted spawning, and

two to four days prior to the full moon, whole polyps were removed from each colony and

longitudinal sections imaged (Canon 5d MKIII, Fig. 4a,d). Based on the stage of gamete

development in these sections *ex situ* spawning activity was predicted.

To record spawning observations each year, colonies were observed with red head torches during

the predicted spawning months. Observations commenced two NAFM and 30 mins prior to the

predicted spawning time, during which time the broodstock aquariums were isolated from the

filtration system. In addition, all internal water pumps were also turned off leaving the aquariums'

water static. Observations continued over multiple consecutive nights until all colonies that had

developed gametes, during that reproductive season, spawned. Spawning date and time of each

colony was recorded, with onset of spawning being denoted as the time of first gamete release.

Following release, gametes were pooled to allow fertilization to occur, and subsequent embryos

were reared and larvae settled following the method described by Craggs et al. (2019).

Results

Production of single polyps and long-term rearing of bleached polyps

We produced clonal replicates of *G. fascicularis* colonies from the Red Sea and Hong Kong and

bleached them with menthol. All polyps (n = 24, including additional polyps not used in this

experiment) and all coral fragments (n = 16) survived the menthol treatment and appeared visually

completely white. Visual inspection under a fluorescent stereomicroscope revealed no detectable

algal cells in the bleached polyps (Red Sea colonies, Fig. 1a), but inspection at higher

magnification (compound fluorescent microscope, ×400) revealed the presence of few scattered

algal cells, limited to the base of the tentacles (Fig. 1b).

All polyps that were employed in the thermal performance experiment (n = 12) remained visually

completely bleached and viable throughout the 10 days of temperature treatment, with the

exception of two polyps that appeared dead on the 7^{th} and 8^{th} day of incubation, respectively. These

polyps showed atrophic and unresponsive tentacles, and overall thinning of the soft tissue until it

was not recognizable. Overall, we successfully maintained the bleached polyps for three weeks

from the termination of the menthol treatment.

Fig. 1 Microscopic comparison between symbiotic and menthol-bleached polyps. Representative stereo and compound micrographs in bright field and UV light, with filters for chlorophyll (red) and coral tissue (green) autofluorescence. **a** Symbiodiniaceae cells are abundant in symbiotic polyps while not detectable in bleached polyps. **b** At higher magnifications, few algal cells are still detectable in bleached polyps only at the base of the tentacles (arrows)

Amenability to experimental manipulation and broad thermal tolerance

Symbiotic (untreated) polyps had positive net photosynthesis across all temperatures from 21 to 32 °C (Fig. 2). Net photosynthesis peaked at 23 °C (mean 13.05 µg O_2 cm^{-2} h^{-1}) and reached the minimum at 32 °C (mean 3.26 µg O_2 cm^{-2} h^{-1}). In contrast, respiration overall increased with temperature with minimum values at 21 °C (mean 5.77 µg O_2 cm^{-2} h^{-1}) and maximum at 32 °C (mean 25.70 µg O_2 cm^{-2} h^{-1}; Fig. 2). Bleached polyps, as expected by their lack of photosymbionts, exhibited a net oxygen depletion. Similar to symbiotic polyps, respiration in bleached polyps increased with temperature, both in light and dark incubations (Fig. 2). Overall, respiration in symbiotic polyps was higher than in bleached polyps ($P_{lmer} < 0.001$).

316　While net photosynthesis was similar between colonies ($P_{lmer} > 0.05$), respiration was significantly

317　higher in one colony ($P_{lmer} < 0.001$) compared to the other two (RS2-RS1: $P_{Bonf} < 0.01$; RS2-RS3:

318　$P_{Bonf} < 0.001$; RS1-RS3: $P_{Bonf} > 0.05$; Fig. S4).

Fig. 2 Net photosynthesis and respiration rate of symbiotic states in *Galaxea fascicularis.*

321　Symbiotic (brown symbols) and Bleached (white symbols) polyps in light and dark incubations.

322　Noise added (jittering) for ease of visualization, means per group depicted as diamonds. n = 12

Symbiosis re-establishment after bleaching

Natural symbiont uptake by placing menthol-bleached polyps next to symbiotic colonies resulted

in the re-establishment of the coral algal symbiosis. In the long-term, these polyps grew into small

colonies that were visually indistinguishable from untreated corals (Fig. S5a). The degree and

speed of repopulation differed between polyps from different colonies. One month after natural

uptake, polyps from one colony appeared fully symbiotic while polyps from another colony were

only intermediately populated (Fig. 3a).

Inoculation with freshly isolated Symbiodiniaceae was followed by symbiont repopulation in three

out of seven bleached polyps. These belonged to the same colony and repopulation was

independent of the source inoculum (i.e. symbionts isolated from the same or from a different

colony). Similar to the successfully repopulated polyps from natural uptake, successfully

inoculated polyps grew into small colonies visually indistinguishable from untreated corals one

year later (Fig. S5b).

Inoculation with cultured symbionts resulted in successful symbiosis re-establishment for 100 %

of the samples (n = 16), which were macroscopically visibly symbiotic. Of these, nine samples

were successfully FISH hybridized, with the genus-specific probes identifying the presence of both

target genera in each sample (Fig. 3b). Individual variation in the relative ratio of each genus was

evident among the fragments, with *Durusdinium* contributing to 4 – 63 % of cells and

Cladocopium to 37 – 82 % (Fig. 3b). A small, but significant proportion of symbionts remained

unlabeled in some samples. These may represent additional symbiont genera or be the result of

variable probe efficiency.

Fig. 3 Symbiosis reestablishment of menthol-bleached polyps following inoculation with Symbiodiniaceae. **a** Representative stereomicrographs of untreated ('Symbiotic') polyp and fully aposymbiotic ('Bleached') polyp 10 days after menthol treatment. One month after inoculation with freshly isolated Symbiodiniaceae, polyps originating from different colonies show different degrees of repopulation. UV light visualizes chlorophyll (red) and host tissue (green) fluorescence. Bright field: natural light (2950 K); UV light + GF2 filter; (Leica, MZ16 F). Scale bar = 2 mm. **b** Relative abundance (%) of *Cladocopium* (gray), *Durusdinium* (black), and unlabeled cells (white) following menthol bleaching and inoculation with cultured algae of these genera. The presence of both genera is confirmed, with individual variation in relative ratios across fragments (n = 9).

Ex-situ spawning

Spawning was observed each year over the three-year observation period for all 35 colonies. Annual fecundity, as a percentage of colonies completing a full gametogenic cycle, was 43.97 % (3.02 sd) (2019: 40.48 %, 2020 and 2021: 45.71 %) during these three spawning cycles. Spawning occurred in synchrony with wild predictions, with gametes being released in November and December each year, between 0.26 - 259 MAS and 5 - 9 NAFM (Fig. 4b-c & e-f). Following embryo rearing, Symbiodiniaceae were observed within primary polyps 14 days post settlement (Fig. 4g). Polyps were fully developed 40 days post settlement (Fig. 4h) and grew into small colonies within the first six months (Fig. 4i).

Fig. 4 *Galaxea fascicularis* spawning *ex situ*. Representative images of gamete development and spawning in female and male colonies. **a** Longitudinal section of female polyp with pinkish-red pigmented oocytes. **b** Female colony releasing oocyte bundle *ex situ*. **c** Close-up of female bundle post release. **d** Longitudinal section of male polyp with white oocytes. **e** Male colony releasing oocytes/sperm bundle *ex situ*. **f** Close-up of male gamete bundles post release undergoing disassociation. **g** Newly settled *G. fascicularis* primary polyps showing initial Symbiodiniaceae (arrow) uptake 14 days post settlement. **h** Primary polyp 40 days post settlement. **i** Multiple colonies six months post settlement. Scale bar: a-h = 1 mm, i = 1 cm

Discussion

Here, we explored the potential of *G. fascicularis* as a novel coral model by assessing its amenability to aquarium rearing, experimental handling, and symbiosis manipulation.

Successful bleaching with menthol

We showed that *G. fascicularis* can be readily and reliably rendered aposymbiotic through menthol bleaching, similar to other cnidarian model systems (e.g., Matthews et al. 2015; Röthig et al. 2021). We lowered the menthol concentration in the protocol developed by Wang et al. (2012) to further limit stress on the host, and indeed all coral polyps survived this bleaching procedure on three independently replicated occasions. Overall, menthol treatment was very effective, with only a few algal cells remaining at the base of the tentacles. From a photophysiological perspective, these scant remnants can be considered negligible, as illustrated by the light and dark oxygen production data. Furthermore, bleached polyps remained aposymbiotic for weeks and we therefore considered this bleaching protocol suitable for experimental investigation of aposymbiotic *G. fascicularis* in physiological studies and symbiosis manipulation. However, as menthol has photo-inhibitory (Wang et al. 2012; Clowez et al. 2021) and antimicrobial properties (İşcan et al. 2002), further studies should investigate the effects of menthol bleaching on the remaining microbiome.

Tractability of symbiotic and aposymbiotic *G. fascicularis*

Tractability, as amenability to experimental work and laboratory conditions, is one of the most important requirements for model organisms. The *G. fascicularis* polyps and fragments could be maintained for three and ten weeks respectively in simple systems consisting of small independent tanks (5-6 L) with regular feeding and basic water quality care. These time frames match the requirements for most experimental designs and we believe could be further extended. Indeed, symbiotic individuals looked healthy and responsive throughout the whole period, testifying to a

comparatively hardy coral with only low demands on rearing conditions. Additionally, the positive

net photosynthetic rate across a large temperature range confirmed the broad thermal tolerance of

this species (Al-Horani 2005; McIlroy et al. 2019), which was retained in the simplified single

polyp application.

Moreover, bleached polyps could be maintained in the simple experimental system for three weeks

even with the additional stress of the thermal experiment. However, they showed signs of

deteriorating condition over time indicating that not all nutritional needs were met. Thus one

priority is to develop bespoke feed for long-term rearing of bleached coral polyps (Wang et al.

2012).

Thermal performance of symbiotic and aposymbiotic *G. fascicularis*

Studies that compare symbiotic and aposymbiotic coral holobionts are particularly informative for

understanding coral response to climate change, and therefore we investigated the comparability

between symbiotic and aposymbiotic *G. fascicularis* polyps. As expected, these differed in their

photosynthetic rates, where the latter were net oxygen consumers even under illumination. Overall,

symbiotic polyps had higher respiration rates than aposymbiotic polyps, which could be ascribed

to the additional metabolic burden of the associated Symbiodiniaceae, as well as a greater

metabolic capacity derived from the availability of photosynthates (Muscatine et al. 1981; Al-

Horani et al. 2003). For all polyps, respiration rates overall increased with temperature in line with

optimum thermal performance of corals and macroalgae from the central Red Sea (27-33 ˚C)

(Anton et al. 2020).

Interestingly, we observed intraspecific variability in physiological performance. One colony had

significantly higher respiration rates than the others (Fig. S4). This manifested exclusively under

dark conditions and was therefore more likely linked to host rather than symbiont identity. This

variability warrants further investigations to better characterize host genotypes, as in the long term

it will be necessary to establish representative host clonal lines.

Symbiosis reestablishment in adult corals

We showed that bleached *G. fascicularis* polyps successfully reestablished symbiosis with

Symbiodiniaceae using qualitative and quantitative approaches. The simpler approaches included

exposure to symbionts through the water medium (by proximity with symbiotic colonies), and a

one-time raw inoculum of Symbiodiniaceae. In the short-term, bleached hosts retained the ability

to feed, which facilitates symbiont uptake (Fitt and Trench 1983) and likely contributes to

symbiosis re-establishment. In the long-term, these polyps continued to grow into small colonies

and became visually indistinguishable from the untreated controls. Therefore, these trials allowed

us to qualitatively demonstrate that symbiosis reestablishment of aposymbiotic adults after

menthol treatment is possible.

Further, the quantitative approach also demonstrated that it is possible to reestablish symbiosis in

adult *G. fascicularis* with cultured Symbiodiniaceae. Remarkably, these were likely heterologous

strains (different from the natively associated ones) as they were initially isolated from *Acropora*

spp. from the GBR (Australia) and inoculated to *G. fascicularis* from Hong Kong. The

experimental repopulation of adult corals with cultured heterologous symbionts has only recently

been demonstrated (Scharfenstein et al. 2022), and it is particularly relevant in the context of

symbiosis manipulations to enhance heat tolerance in reef-building corals (van Oppen et al. 2015).

Further assessments of Symbiodiniaceae identity from algal cultures, symbiotic corals, and

inoculated corals are however necessary to confirm that this heterologous symbiosis

reestablishment occurred in our experiment.

Nevertheless, our targeted inoculation highlighted other interesting prospects for symbiosis

manipulations. The high thermal tolerance of *G. fascicularis* (Marshall and Baird 2000) has been

hypothesized to be linked to its flexible host-algal association, where *Cladocopium*, *Durusdinium*,

and their simultaneous co-occurrence have been documented, and with an equatorial-ward shift

from *Cladocopium* dominance to *Durusdinium* dominance (Huang et al. 2011, and references

therein). At Hong Kong's latitude, and at our rearing temperature, *G. fascicularis* is expected to

be exclusively associated with *Cladocopium* (Huang et al. 2011; Zhou et al. 2017a), yet the

targeted inoculation produced mixed-genus associations. This shows that targeted inoculation can

successfully shift Symbiodiniaceae community composition towards a higher proportion of heat

tolerant strains (Silverstein et al. 2015) without thermal treatment. Although the longevity of these

partnerships is still unclear, these are promising results for future development of *in hospite* studies

on the role of temperature on inter-partner symbiosis dynamics.

***Ex situ* spawning**

Galaxea fascicularis is a good model species candidate for sexual reproductive work due its high

survivorship in aquarium conditions. As demonstrated herein, a combination of seasonal

programming and husbandry enabled the species to complete a full gametogenic cycle *ex situ* over

multiple years, which closely mimics that observed in the wild (Babcock et al. 1986; Baird et al.

2021). Furthermore, we showed *in vitro* fertilization, embryological development, and *ex situ*

larval settlement resulting in spat of known ages. As the age at which the species reaches sexual

maturity is currently unknown, future studies should focus on this aspect, which is also important

for producing an F2 generation and for closing the species life cycle *ex situ* (Craggs et al. 2020).

Access to a F2 generation of known parental lineage will provide a platform to conduct

experiments in areas such as mapping quantitative trait loci (Zhang 2012) or phenotypic traits such

as growth, disease resistance, or thermal tolerance (van Oppen et al. 2015). It also opens

possibilities for studies on heritability in symbiont selection, and for rearing of axenic or

gnotobiotic coral lineages, paralleling procedures established for *Hydra* (Fraune et al. 2015).

Conclusions

The study of coral-algal symbiosis has been constrained by the difficulties of maintaining corals

ex situ and of manipulating the coral-algal association to unravel symbiosis functioning. Here, we

showed that adopting *G. fascicularis* as a model system can alleviate these limitations. Its

demonstrated compatibility to rearing in simplified systems and experimental applications,

together with the repeated spawning in aquaria, confirm the high tractability of this coral species.

Further, the possibility to readily obtain aposymbiotic individuals suitable for detailed study and

symbiosis re-establishment set the stage for exciting future developments in coral symbiosis

research which is necessary to understand corals' potential to cope with changing oceans.

Besides these strengths, our study also highlights areas worth future efforts on the path to

developing the *G. fascicularis* model system further. These include: i) better characterization of

the effect of menthol on the rest of the coral microbiome (e.g., bacteria); ii) development of custom

feed for long-term maintenance of bleached individuals; iii) assessment of long-term persistence

of heterologous Symbiodiniaceae, and testing of a variety of symbiont strains to understand the

breadth of symbiont diversity that can be accommodated by the *G. fascicularis* host; iv) further

characterization of host colonies and genotypes with the aim of establishing clonal lines, mirroring

for example the development of the Aiptasia model; and v) development of collaborative platforms

and open-access community resources around this emerging model system to accelerate research

and discovery.

Acknowledgements

We thank Dr. Jafargholi Imani and Dr. Kathrin Ehlers (both JLU) for access and guidance in the

fluorescence microscope facilities, André Dietzmann and Catarina P. P. Martins (both JLU) for

assistance during experiments, and the Li Ka Shing Faculty of Medicine Core Facility (HKU) for

the flow cytometry equipment.

Data and code availability

Data and R scripts used for this study are available at:

https://github.com/sPuntinG/Galaxea_Coral_Model/

Author contributions

GP and MZ conceived and designed the study. GP, JC, RH, KEE, SM produced and analyzed data.

MS, DMB, MZ contributed reagents/materials/analysis tools. GP and MZ wrote the first draft with

contributions from all authors.

Conflict of Interest

The authors declare no conflict of interest for this submission.

References

Al-Horani FA (2005) Effects of changing seawater temperature on photosynthesis and

calcification in the scleractinian coral *Galaxea fascicularis*, measured with O_2, Ca^{2+} and pH

microsensors. Sci Mar 69:347–354

http://scientiamarina.revistas.csic.es/index.php/scientiamarina/article/view/263/261

Al-Horani FA, Al-Moghrabi SM, de Beer D (2003) The mechanism of calcification and its

relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*.

Mar Biol 142:419–426 http://link.springer.com/10.1007/s00227-002-0981-8

Anton A, Randle JL, Garcia FC, Rossbach S, Ellis JI, Weinzierl M, Duarte CM (2020)

Differential thermal tolerance between algae and corals may trigger the proliferation of

algae in coral reefs. Glob Chang Biol 26:4316–4327

https://onlinelibrary.wiley.com/doi/10.1111/gcb.15141

Babcock RC, Bull GD, Harrison PL, Heyward AJ, Oliver JK, Wallace CC, Willis BL (1986)

Synchronous spawnings of 105 scleractinian coral species on the Great Barrier Reef. Mar

Biol 90:379–394 http://link.springer.com/10.1007/BF00428562

Baird AH, Guest JR, Edwards AJ, et al. (2021) An Indo-Pacific coral spawning database. Sci

Data 8:35 http://www.nature.com/articles/s41597-020-00793-8

Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G (2008) MeshLab: an

Open-Source Mesh Processing Tool.

Clowez S, Renicke C, Pringle JR, Grossman AR (2021) Impact of menthol on growth and

photosynthetic function of *Breviolum minutum* (Dinoflagellata, Dinophyceae,

Symbiodiniaceae) and interactions with its *Aiptasia* host. J Phycol 57:245–257

https://onlinelibrary.wiley.com/doi/10.1111/jpy.13081

Craggs J, Guest JR, Bulling M, Sweet M (2019) Ex situ co culturing of the sea urchin, *Mespilia*

globulus and the coral *Acropora millepora* enhances early post-settlement survivorship. Sci

Rep 9:12984 http://www.nature.com/articles/s41598-019-49447-9

Craggs J, Guest JR, Davis M, Simmons J, Dashti E, Sweet M (2017) Inducing broadcast coral

spawning ex situ: closed system mesocosm design and husbandry protocol. Ecol Evol

7:11066 https://onlinelibrary.wiley.com/doi/10.1002/ece3.3538

Craggs J, Guest JR, Davis M, Sweet M (2020) Completing the life cycle of a broadcast spawning
coral in a closed mesocosm. Invertebr Reprod Dev 64:244–247
https://doi.org/10.1080/07924259.2020.1759704

Dietzel A, Bode M, Connolly SR, Hughes TP (2020) Long-term shifts in the colony size
structure of coral populations along the Great Barrier Reef. Proc R Soc B Biol Sci
287:20201432 https://royalsocietypublishing.org/doi/10.1098/rspb.2020.1432

Dimond J, Carrington E (2007) Temporal variation in the symbiosis and growth of the temperate
scleractinian coral *Astrangia poculata*. Mar Ecol Prog Ser 348:161–172 http://www.int-
res.com/abstracts/meps/v348/p161-172/

Dong Z-J, Huang H, Huang L-M, Li Y-C (2009) Diversity of symbiotic algae of the genus
Symbiodinium in scleractinian corals of the Xisha Islands in the South China Sea. J Syst
Evol 47:321–326 https://onlinelibrary.wiley.com/doi/10.1111/j.1759-6831.2009.00034.x

Falkowski PG, Dubinsky Z, Muscatine L, Porter JW (1984) Light and the bioenergetics of a
symbiotic coral. Bioscience 34:705–709 https://academic.oup.com/bioscience/article-
lookup/doi/10.2307/1309663

Fitt WK, Trench RK (1983) Endocytosis of the symbiotic dinoflagellate *Symbiodinium*
microadriaticum Freudenthal by endodermal cells of the scyphistomae of *Cassiopeia*
xamachana and resistance of the algae to host digestion. J Cell Sci 64:195–212

Fraune S, Anton-Erxleben F, Augustin R, Franzenburg S, Knop M, Schröder K, Willoweit-Ohl
D, Bosch TCG (2015) Bacteria–bacteria interactions within the microbiota of the ancestral
metazoan *Hydra* contribute to fungal resistance. ISME J 9:1543–1556
http://www.nature.com/articles/ismej2014239

Galliot B (2012) *Hydra*, a fruitful model system for 270 years. Int J Dev Biol 56:411–423
http://www.intjdevbiol.com/paper.php?doi=120086bg

Gates RD, Baghdasarian G, Muscatine L (1992) Temperature stress causes host cell detachment
in symbiotic cnidarians: implications for coral bleaching. Biol Bull 182:324–332
https://www.journals.uchicago.edu/doi/10.2307/1542252

Gattuso J-P, Allemand D, Frankignoulle M (1999) Photosynthesis and Calcification at Cellular,

Organismal and Community Levels in Coral Reefs: A Review on Interactions and Control

by Carbonate Chemistry. Am Zool 39:160–183 https://academic.oup.com/icb/article-

lookup/doi/10.1093/icb/39.1.160

Guillard RRL (1975) Culture of Phytoplankton for Feeding Marine Invertebrates. In: Smith,

W.L., Chanley M.H. (eds) Culture of Marine Invertebrate Animals. Springer US, Boston,

MA, pp 29–60 http://link.springer.com/10.1007/978-1-4615-8714-9_3

Harrison PL (1988) Pseudo-gynodioecy: an unusual breeding system in the scleractinian coral

Galaxea fascicularis. 699–705

https://researchportal.scu.edu.au/esploro/outputs/conferencePaper/Pseudo-gynodioecy-an-

unusual-breeding-system-in-the-scleractinian-coral-Galaxea-

fascicularia/991012821519702368

Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral

reefs. Mar Freshw Res 50:839–866 http://www.publish.csiro.au/?paper=MF99078

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadia G, Raymundo L, Williams G,

Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of

the Paris Agreement. Sci Rep 6:39666 http://www.nature.com/articles/srep39666

Huang H, Dong Z, Huang L, Yang J, Di B, Li Y, Zhou G, Zhang C (2011) Latitudinal variation

in algal symbionts within the scleractinian coral *Galaxea fascicularis* in the South China

Sea. Mar Biol Res 7:208–211

http://www.tandfonline.com/doi/abs/10.1080/17451000.2010.489616

İşcan G, Kirimer N, Kürkcüoğlu M, Hüsnü Can Başer, Demirci F (2002) Antimicrobial

screening of *Mentha piperita* essential oils. J Agric Food Chem 50:3943–3946

https://doi.org/10.1021/jf011476k

Keshavmurthy S, Hsu CM, Kuo CY, Denis V, Leung JKL, Fontana S, Hsieh HJ, Tsai W Sen, Su

WC, Chen CA (2012) Larval development of fertilized "pseudo-gynodioecious" eggs

suggests a sexual pattern of gynodioecy in *Galaxea fascicularis* (Scleractinia: Euphyllidae).

Zool Stud 51:143–149

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. J Stat Softw 82:1–26

https://www.jstatsoft.org/index.php/jss/article/view/v082i13

Liew YJ, Aranda M, Voolstra CR (2016) Reefgenomics.Org - a repository for marine genomics

data. Database 2016:baw152 https://academic.oup.com/database/article-

lookup/doi/10.1093/database/baw152

Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D (2021) performance: An R

Package for Assessment, Comparison and Testing of Statistical Models. J Open Source

Softw 6:3139 https://joss.theoj.org/papers/10.21105/joss.03139

Marshall AT, Clode P (2004) Calcification rate and the effect of temperature in a zooxanthellate

and an azooxanthellate scleractinian reef coral. Coral Reefs 23:218–224

Marshall PA, Baird AH (2000) Bleaching of corals on the Great Barrier Reef: differential

susceptibilities among taxa. Coral Reefs 19:155–163

http://link.springer.com/10.1007/s003380000086

Matthews JL, Sproles AE, Oakley CA, Grossman AR, Weis VM, Davy SK (2015) Menthol-

induced bleaching rapidly and effectively provides experimental aposymbiotic sea

anemones (*Aiptasia* sp.) for symbiosis investigations. J Exp Biol 219:306–310

http://jeb.biologists.org/cgi/doi/10.1242/jeb.128934

McIlroy SE, Thompson PD, Yuan FL, Bonebrake TC, Baker DM (2019) Subtropical thermal

variation supports persistence of corals but limits productivity of coral reefs. Proc R Soc B

Biol Sci 286:20190882 https://royalsocietypublishing.org/doi/10.1098/rspb.2019.0882

McIlroy SE, Wong JCY, Baker DM (2020) Competitive traits of coral symbionts may alter the

structure and function of the microbiome. ISME J 14:2424–2432

https://www.nature.com/articles/s41396-020-0697-0

McLachlan RH, Price JT, Solomon SL, Grottoli AG (2020) Thirty years of coral heat-stress

experiments: a review of methods. Coral Reefs 39:885–902

https://link.springer.com/10.1007/s00338-020-01931-9

Muscatine L, R. McCloskey L, E. Marian R (1981) Estimating the daily contribution of carbon

from zooxanthellae to coral animal respiration. Limnol Oceanogr 26:601–611

http://doi.wiley.com/10.4319/lo.1981.26.4.0601

Niu W, Huang H, Lin R, Chen C-H, Shen K-N, Hsiao C-D (2016) The complete mitogenome of

the Galaxy Coral, *Galaxea fascicularis* (Cnidaria: Oculinidae). Mitochondrial DNA Part B

1:10–11 https://www.tandfonline.com/doi/full/10.1080/23802359.2015.1137796

van Oppen MJH, Oliver JK, Putnam HM, Gates RD (2015) Building coral reef resilience

through assisted evolution. Proc Natl Acad Sci 112:2307–2313

http://www.pnas.org/lookup/doi/10.1073/pnas.1422301112

Pavia RTB, Estacion JS (2019) Survival and growth of isolated polyps of *Galaxea fascicularis*

(Linnaeus 1767) on six kinds of culture substrates: implications for mariculture, aquarium

culture, and conservation. J World Aquac Soc 50:219–230

https://onlinelibrary.wiley.com/doi/10.1111/jwas.12538

Puntin G, Sweet M, Fraune, Sebastian Medina M, Sharp K, Weis VM, Ziegler M (2022)

Harnessing the power of model organisms to unravel microbial functions in the coral

holobiont. bioRxiv

R Core Team (2021) R: A Language and Environment for Statistical Computing. https://www.r-

project.org/ https://www.r-project.org/

Rades M, Schubert P, Ziegler M, Kröckel M, Reichert J (2022) Building plan for a temperature-

controlled multi-point stirring incubator. https://www.protocols.io/view/building-plan-for-a-

temperature-controlled-multi-p-dm6gpb34dlzp/v1

Röthig T, Puntin G, Wong JCY, Burian A, McLeod W, Baker DM (2021) Holobiont nitrogen

control and its potential for eutrophication resistance in an obligate photosymbiotic

jellyfish. Microbiome 9:127

https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-021-01075-0

Scharfenstein HJ, Chan WY, Buerger P, Humphrey C, van Oppen MJH (2022) Evidence for de

novo acquisition of microalgal symbionts by bleached adult corals. ISME J 16:1676–1679

https://www.nature.com/articles/s41396-022-01203-0

Schneider K, Erez J (2006) The effect of carbonate chemistry on calcification and photosynthesis

in the hermatypic coral *Acropora eurystoma*. Limnol Oceanogr 51:1284–1293

https://doi.org/10.4319/lo.2006.51.3.1284

Schubert P, Wilke T (2018) Coral Microcosms: Challenges and Opportunities for Global Change

Biology. Corals in a Changing World. InTech, http://www.intechopen.com/books/corals-in-

a-changing-world/coral-microcosms-challenges-and-opportunities-for-global-change-

biology

Searle SR, Speed FM, Milliken GA (1980) Population Marginal Means in the Linear Model: An

Alternative to Least Squares Means. Am Stat 34:216–221

http://www.tandfonline.com/doi/abs/10.1080/00031305.1980.10483031

Silverstein RN, Cunning R, Baker AC (2015) Change in algal symbiont communities after

bleaching, not prior heat exposure, increases heat tolerance of reef corals. Glob Chang Biol

21:236–249 https://onlinelibrary.wiley.com/doi/10.1111/gcb.12706

Tong H, Cai L, Zhou G, Yuan T, Zhang W, Tian R, Huang H, Qian P-Y (2017) Temperature

shapes coral-algal symbiosis in the South China Sea. Sci Rep 7:1–12

http://dx.doi.org/10.1038/srep40118

Veron JEN, Stafford-Smith MG, Turak E, DeVantier LM (2016) Corals of the World.

http://www.coralsoftheworld.org/species_factsheets/species_factsheet_summary/galaxea-

fascicularis/

Voolstra CR (2013) A journey into the wild of the cnidarian model system *Aiptasia* and its

symbionts. Mol Ecol 22:4366–4368

Wang J-T, Chen Y-Y, Tew KS, Meng P-J, Chen CA (2012) Physiological and biochemical

performances of menthol-induced aposymbiotic corals. PLoS One 7:e46406

https://dx.plos.org/10.1371/journal.pone.0046406

Weis VM, Davy SK, Hoegh-Guldberg O, Rodriguez-Lanetty M, Pringle JR (2008) Cell biology

in model systems as the key to understanding corals. Trends Ecol Evol 23:369–376

https://linkinghub.elsevier.com/retrieve/pii/S0169534708001614

Wepfer P, Nakajima Y, Hui F, Mitarai S, Economo E (2020a) Metacommunity ecology of

Symbiodiniaceae hosted by the coral *Galaxea fascicularis*. Mar Ecol Prog Ser 633:71–87

https://www.int-res.com/abstracts/meps/v633/p71-87/

Wepfer PH, Nakajima Y, Sutthacheep M, Radice VZ, Richards Z, Ang P, Terraneo T, Sudek M,

Fujimura A, Toonen RJ, Mikheyev AS, Economo EP, Mitarai S (2020b) Evolutionary

biogeography of the reef-building coral genus *Galaxea* across the Indo-Pacific ocean. Mol

Phylogenet Evol 151:106905 https://doi.org/10.1016/j.ympev.2020.106905

Wickham H (2016) Ggplot2: Elegant graphics for data analysis. Springer International

Publishing, Cham, Switzerland https://books.google.de/books?id=RTMFswEACAAJ

Wilkinson SP, Pontasch S, Fisher PL, Davy SK (2016) The distribution of intra-genomically

variable dinoflagellate symbionts at Lord Howe Island, Australia. Coral Reefs 35:565–576

http://link.springer.com/10.1007/s00338-015-1357-0

Zhang Y (2012) Quantitative Trait Loci (QTL). Humana Press, Totowa, NJ

http://link.springer.com/10.1007/978-1-61779-785-9

Zhou G, Cai L, Li Y, Tong H, Jiang L, Zhang Y, Lei X, Guo M, Liu S, Qian P-Y, Huang H

(2017a) Temperature-driven local acclimatization of *Symbiodnium* hosted by the coral

Galaxea fascicularis at Hainan Island, China. Front Microbiol 8:1–9

http://journal.frontiersin.org/article/10.3389/fmicb.2017.02487/full

Zhou J, Huang H, Beardall J, Gao K (2017b) Effect of UV radiation on the expulsion of

Symbiodinium from the coral *Pocillopora damicornis*. J Photochem Photobiol B Biol

166:12–17 http://dx.doi.org/10.1016/j.jphotobiol.2016.11.003

Ziegler M, Roder C, Büchel C, Voolstra C (2015) Niche acclimatization in Red Sea corals is

dependent on flexibility of host-symbiont association. Mar Ecol Prog Ser 533:149–161

http://www.int-res.com/abstracts/meps/v533/p149-161/