1 **The reef-building coral *Galaxea fascicularis*: a new model system for coral**

2 **symbiosis research**

Giulia Puntin[1], Jamie Craggs[2,3], Róisín Hayden[4], Kara E. Engelhardt[1], Shelby McIlroy[4], Michael

Sweet[3,5], David M. Baker[4], Maren Ziegler[1,]*

[1]Marine Holobiomics Lab, Department of Animal Ecology & Systematics, Justus Liebig

University Giessen, Heinrich-Buff-Ring 26-32 IFZ, 35392 Giessen, Germany

[2]Horniman Museum and Gardens, Forest Hill, London, SE23 3PQ, United Kingdom

[3]Coral Spawning Lab, Unit 6, Midas Metro Centre, 193 Garth Rd, Morden, SM4 4NE, United

Kingdom

[4]The Swire Institute of Marine Science, University of Hong Kong, Cape D'Aguilar, Hong Kong

[5]Aquatic Research Facility, Environmental Sustainability Research Centre, University of Derby,

DE22 1GB, United Kingdom

ORCiD:
GP: 0000-0001-9099-1730
JC: 0000-0002-3787-2203
SM: 0000-0003-2482-8817
MS: 0000-0003-4983-8333
DMB: 0000-0002-0308-4954
MZ: 0000-0003-2237-9261
20
21   *Corresponding author: maren.ziegler@bio.uni-giessen.de

Abstract

Reef-building corals owe their evolutionary success to their symbiosis with unicellular algae (Symbiodiniaceae). However, increasingly frequent heat waves lead to coral mass-bleaching events and pose a serious threat to the survival of reef ecosystems. Despite significant efforts, a mechanistic understanding of coral-algal symbiosis functioning, what leads to its breakdown and what can prevent it, remains incomplete. The main obstacles are low amenability of corals to experimental handling and, owing to its obligatory nature, the difficulties of manipulating the coral-algal association. Indeed, many studies on the symbiotic partnership are conducted on other cnidarian model organisms and their results may therefore not be fully transferable to tropical reef-building corals. Here, we identify the tropical stony coral species *Galaxea fascicularis* as a novel candidate coral model system. Individual polyps of this species can be separated, enabling highly replicated genotype studies, and are well suited to experimental investigation of the symbiosis as they can be easily and effectively rid of their algal symbionts (bleached). We show that bleached adult individuals can reestablish symbiosis with non-native symbionts, and we report the completion of the gametogenic cycle *ex-situ*, with the successful spawning in aquaria over multiple years. These achievements help overcome several of the major limitations to direct research on corals and highlight the potential of *G. fascicularis* as an important new model system for investigations of symbiosis functioning and manipulation.

Keywords: Model organisms; reef-building corals; menthol bleaching; symbiosis manipulation; *ex situ* sexual reproduction

Introduction

Reef-building corals form obligatory endosymbiotic association with unicellular algae of the family Symbiodiniaceae. This association is key to their evolutionary success, but it is also at the heart of corals' susceptibility to global climate change, which manifests in coral bleaching - the breakdown of the coral-algal symbiosis (Hoegh-Guldberg 1999). Bleaching is mostly driven by marine heatwaves which are predicted to worsen, causing the loss of virtually all coral reefs by the end of this century (van Hooidonk et al. 2016).

The interaction between heat stress and bleaching has been studied for more than 30 years (Gates et al. 1992; McLachlan et al. 2020), but a detailed understanding of the mechanisms underlying coral bleaching is still missing. Progress in coral symbiosis research is hampered by two main aspects. First, corals are challenging to maintain in aquarium settings as they tolerate only a narrow range of environmental conditions, i.e., bleaching can be triggered by just 1-2 °C increases above summer mean temperatures (Glynn and D'Croz 1990). Second, the obligatory nature of the coral-algal association makes it particularly challenging to physically and functionally separate the partners, an approach often necessary to unravel the mechanisms underlying symbiosis breakdown and what could prevent it (Weis et al. 2008; Voolstra 2013).

Model organisms are integral for understanding fundamental biological principles and symbiotic cnidarians have successfully been used as "coral models", advancing our understanding of coral bleaching and holobiont functioning (Weis et al. 2008). These model organisms share important traits with corals, such as being cnidarians that associate with microalgae, yet they also lack other features that normally represent obstacles to research work such as the calcium carbonate skeleton. In addition, the ability to study the animal host and its algal symbiont in isolation is more easily achieved with facultatively symbiotic organisms such as *Hydra* spp., Aiptasia (*Exaiptasia*

diaphana), and *Astrangia poculata* (Dimond and Carrington 2007; Weis et al. 2008; Galliot 2012).

Cnidarian model organisms allow the study of basal or shared traits or phenomena at a speed that would not be possible with reef-building corals, but testing on the latter remains necessary for the understanding of coral-specific or ecologically relevant aspects. The obligatory symbiosis with Symbiodiniaceae in reef-building corals has important biological implications (Falkowski et al. 1984; Hoegh-Guldberg 1999), similar to the calcification process that is deeply intertwined with coral physiology and ecology (Gattuso et al. 1999). It is therefore important to establish a "true" tropical reef-building coral model species.

The richness of the scleractinian taxon offers a vast array of species to choose from in the quest for a suitable candidate coral model species. These can be evaluated for their tractability (as amenability to experimental work) considering several aspects which, as proposed by Puntin et al. (2022) should include: 1) pre-existing knowledge: baseline information of the organism's biology is necessary to interpret and contextualize results; 2) compatibility with aquarium rearing: the possibility to maintain the organism and preferably to complete its life cycle in the lab is essential to increase its availability, reduce confounding effects such as unknown life history, improve reproducibility, and lower the pressure on threatened wild populations; 3) amenability to symbiosis manipulations is necessary to help unravel complex coral-algal functional interactions, and can be broken down into the organism's suitability to be rendered aposymbiotic (bleached), to be maintained aposymbiotic for a certain time, and to then be re-infected with Symbiodiniaceae.

We have identified the coral species *Galaxea fascicularis* (Linnaeus, 1767) (clade: Complexa, family: Euphyllidae) as a promising candidate coral model system. The species is relatively common across the Indo-Pacific region (Veron et al. 2016), where the phylogeny of the genus *Galaxea* is geographically well resolved (Wepfer et al. 2020b). It is also one of the more commonly

used species in heat-stress experiments (McLachlan et al. 2020) and in studies on coral

calcification (e.g., Marshall and Clode 2004; Al-Horani 2005). Importantly, other community

resources such as an annotated draft genome are available (Liew et al. 2016; Niu et al. 2016,

http://www.gfas.reefgenomics.org/). In addition, *G. fascicularis* is known for its tolerance to

stressors as it is "largely unaffected by bleaching" in the field (Marshall and Baird 2000) and for

being easy to grow and propagate in aquaria (Pavia and Estacion 2019). Further, its morphology

provides additional practical advantages with relatively big polyps, which can be easily isolated

(Pavia and Estacion 2019). This conveniently allows reduction of the complexity of the organism

from colonial to individual, enables high clonal replication, and facilitates visualization (Al-Horani

et al. 2003; Marshall and Clode 2004). Its association with Symbiodiniaceae is relatively well

characterized (Huang et al. 2011; Wepfer et al. 2020a). Furthermore, the species' reproductive

mode and spawning patterns in the wild are well documented (Babcock et al. 1986; Harrison 1988;

Keshavmurthy et al. 2012). However, its amenability to symbiosis manipulation, to experimental

handling of bleached individuals, and the ability to complete gametogenic cycles and spawning in

closed aquaria have not yet been described.

The overarching aim of this study was to explore the potential of *Galaxea fascicularis* as a coral

laboratory model. Specifically, we investigated: 1) the feasibility to render polyps aposymbiotic

by eliminating Symbiodiniaceae (bleaching), 2) the amenability to experimental manipulation of

symbiotic and aposymbiotic polyps in a simplified system, 3) the re-establishment of the symbiosis

with different species of Symbiodiniaceae, and 4) the possibility to induce full gametogenic cycles

ex situ, with subsequent spawning under aquarium conditions. These aspects are central to

overcoming the two main obstacles to coral research work: low tractability of corals and their

complex nature.

Materials and methods

Coral collection and long-term aquarium rearing

Colonies of *Galaxea fascicularis* were collected from three geographical locations: the Red Sea, Hong Kong, and the Great Barrier Reef (Fig. 1). To increase comparability, rearing conditions were adjusted to be similar between labs, with few exceptions pertaining to the particular experimental needs or feeding regime, which was based on the long-term rearing arrangements in each facility (Tab. S1). Red Sea colonies were collected at 9-13 m depth at the North-Eastern protected end of the reef "Al Fahal" in the central Saudi Arabian Red Sea (22.3054°, 38.9655°) in March 2019, and transported to the Ocean2100 aquarium facility at Justus Liebig University Giessen (Germany) where they are part of the live collection (CITES permit 19-SA-000096-PD). In the aquarium system, light was provided by white and blue fluorescent lamps with a light:dark cycle of 12:12 h at 130 -160 µmol photons m^{-2} s^{-1}, which approximates light condition at the collection site (Ziegler et al. 2015). Salinity was maintained around 35 and temperature at 26 °C. Colonies were fed daily with a combination of frozen copepods, *Artemia*, krill, and *Mysis*.

Hong Kong colonies were collected from 5 m (max. depth) from Crescent Island (22.5308°, 114.3150°) in November 2020 and transported to the University of Hong Kong, where they were fragmented and acclimated to aquarium conditions for one month. The aquarium system consisted of 6-L acrylic tanks placed within a Plant Growth Chamber (Panasonic MLR-352H-PA). Each tank held eight ~3 cm-fragments and was fitted with a submersible pump (Atman AT301) and a small water filter (Shiruba PF120). The fragments used in symbiosis reestablishment were fed twice-weekly with a powdered blend of marine plankton (ReefRoids, Polyp Lab) followed by a partial water change. Light, salinity, and temperature conditions were consistent with those maintained in the Ocean2100 facility.

For *ex situ* spawning, 35 colonies (10.1 cm ± 2.5 (mean ± sd)) were collected from Arlington Reef,

Great Barrier Reef, Australia (-16.7000°, 146.0500°) in September 2019, and transported to the

Horniman Museum and Gardens (United Kingdom; CITES permit 585319/01) with the 'inverted

submersion method' as described by Craggs et al. (2018). Colonies were distributed into four coral

spawning systems (www.coralspawninglab.org) that replicated the natural environmental

parameters (seasonal temperature, photoperiod, solar irradiance, and lunar cycle) required to

stimulate reproduction (Tab. S1). Aquarium design and coral husbandry protocol was based on the

mesocosms described by Craggs et al. (2017), and the seasonal temperature profile were based on

a non-sequential eight year average (1998 - 2017) from Moore Reef (-16.8667°, 146.2334°')

(http://data.aims.gov.au/aimsrtds/datatool.xhtml?site=931¶m=water%20temperature) (Fig.

S7). Colonies were fed daily on a mix of phytoplankton (*Tisochrysis lutea*, *Chaetoceros calitrans*

and *Rhodomonas salina*) and zooplankton (newly hatched *Artemia salina* nauplii, frozen red

plankton, rotifers, and lobster / fish eggs) (Tab. S1). During feeding the broodstock tank was

isolated from the filtration for two hours to aid prey capture and uptake.

Fig. 1 Summary of *G. fascicularis* coloniy origins, participating labs, and experiments described in this study. Colonies were collected from three distant locations (Red Sea, Hong Kong, and the Great Barrier Reef) and separately employed in different experiments (menthol bleaching, thermal performance, symbiosis reestablishment, and *ex situ* spawning). Map modified from Corals of the World (www.coralsoftheworld.org, Veron et al. (2016)).

Thermal performance experiment on single polyps

Preparation of single polyps

For the menthol bleaching and thermal performance experiment, we used single polyps from three Red Sea colonies each showing distinct coloration (Fig. S1). Replicate clonal polyps were isolated from each colony using an electric rotary cutter, mounted on coral glue (Grotech, CoraFix SuperFast), and allowed to heal for two weeks in the Ocean2100 aquarium facility (Schubert and Wilke 2018).

Menthol bleaching to remove algal symbionts

An equal number of polyps from each colony were randomly assigned to the control group

(maintained in a symbiotic state), and the bleached group (chemically bleached with menthol).

Menthol bleaching followed a protocol modified from Wang et al. (2012) and consisted of three

days of treatment in 0.38 mM menthol solution in seawater, followed by a day of rest and a fourth

day of menthol treatment. Each day the polyps were incubated in menthol for 8 h under light and

stirring (Fig. S2). After, the polys were rinsed and kept in clean containers with air bubbling and

stirring. During menthol treatment, the treated polyps were only exposed to filtered (1.2 μm)

artificial seawater (FASW) to prevent exposure to Symbiodiniaceae. Following, all polyps

(including, for consistency, the symbiotic group) were maintained in FASW.

To confirm bleaching, half of the polyps from each group (bleached and symbiotic) and each

colony were visually assessed with a fluorescence stereomicroscope (Leica MZ16 F) 10 days after

the termination of the menthol treatment. Representative pictures were taken under natural light

(brightfield, light source: 2950 K) and under UV light with GF2 filters allowing visualization of

the green fluorescent proteins and chlorophyll (Fig. 2a). Following the same bleaching protocol,

menthol-bleached polyps were also documented using a compound epifluorescence microscope to

reach higher resolution (Leica DM 5500B, TX2 filter) (Fig. 2b).

Thermal performance experimental design

Symbiotic and bleached polyps were moved to the experimental tanks on the day after the last

menthol exposure and allowed to recover and acclimate for 10 days. The experimental system

consisted of eight 5-L glass tanks (20 cm × 30 cm) equipped with a small pump (Resun SP-500)

in a common temperature-controlled water bath (26 ˚C). White fluorescent lamps were used to

maintain the light cycle and intensity consistent with long-term rearing conditions. Bleached and

symbiotic polyps were kept in separate tanks, with four replicate tanks per treatment filled with

FASW. Each tank contained one polyp from each colony (n = 3) that were used for the thermal

performance experiment. Additional polyps were kept in each tank as back-up (Fig. S3). Polyps

were fed each day after the end of the dark incubation, followed by 10 % water change after 2-3

h. Each polyp was fed one small frozen adult Artemia pipetted directly on top of the oral opening.

The polyps were cleaned every two to three days after the incubations to remove fouling that would

interfere with respirometry measurements.

To test the amenability to experimental handling of single polyps and to compare metabolic rates

of bleached and symbiotic polyps, we conducted a thermal performance experiment.

Photosynthesis and respiration of *G. fascicularis* polyps were measured across a 12 ˚C temperature

range over 10 days. For this, the polyps were acclimated to a different temperature each day, in

the following order: 20, 22, 23, 24, 25, 26, 27, 28, 30 to 32 °C. The temperature of the water bath

was changed overnight, and incubations took place the subsequent day at the respective

temperature.

Photosynthesis and respiration of symbiotic and bleached polyps

We measured oxygen evolution of individual *G. fascicularis* polyps in light and dark incubations

to calculate net photosynthesis (PN) and dark respiration (R) respectively. For these measurements

four bleached and four symbiotic polyps from three colonies were used (n = 12). Every day, each

polyp was placed in individual 160 mL glass incubation jars (Weck, Germany), equipped with a

magnetic stirrer (~5 cm s^{-1}). Three additional jars containing only seawater and stirring bars were

used to control background biological activity. Light intensity of 230 - 280 µmol photons m^{-2} s^{-1}

during light incubations (ATI 80W Aquablue Special) and temperatures were maintained through

thermostat-controlled water baths placed on a custom-made multiplate stirring system (Rades et

al. 2022).

Dissolved oxygen (DO) was measured at the start and at the end of each incubation using a

handheld multiparameter probe (WTW Multi 3620 IDS set). Light incubations lasted 2 h around

mid-day followed by 1.5 h of dark incubation. Net photosynthesis and respiration were calculated

using the following formula (Schneider and Erez 2006):

215
$$PN \text{ or } R[mgO_2 \times cm^{-2} \times h^{-1}] \ = \ \frac{\Delta O_2[mg \times L^{-1}] \ \times \ V_{incubation}[L]}{T[h] \times SA[cm^2]}$$

216   Where ΔO_2 is the difference in dissolved oxygen between the end and the start of the incubation

217   (DO_{end} - DO_{start}) corrected for the controls' ΔO_2. This is multiplied by the volume of the incubation

218   chambers in L ($V_{incubation}$), and divided by incubation time (T) in hours and polyp surface area (SA)

219   in cm^2.

Surface area measurements

221   Polyp surface area was measured through photogrammetry. Between 40 to 50 pictures of each

222   polyp were taken with a phone camera two days before the beginning of the incubations. These

223   were used to create 3D models with 3DF Zephyr Free (v.4.523), cleaned on Artec3D (Studio 11

224   Professional v.11.2.2.16), and loaded on MeshLab (v.2016.12, Cignoni et al. 2008) for size scaling

225   and calculation of live coral surface area.

Thermal performance data analysis

All analyses were conducted in the R statistical environment (v.4.1.0, R Core Team 2021) and

plotted with ggplot2 (v.3.3.5, Wickham 2016). The effects of symbiotic state ('symbiotic' vs.

'bleached') or colony identity on net photosynthesis (PN) and respiration (R) were analyzed using

linear mixed-effect models considering light and dark incubations separately. For this, 'symbiotic

state', 'colony identity', and 'temperature' were set as fixed factors, and 'polyp identity' as random

factor. The package 'lmerTest' (v.3.1-3, Kuznetsova et al. 2017) was used to construct the models

and calculate p-values. The package 'performance' was used to check the residuals and to compare

alternative models (v.0.7.3, Lüdecke et al. 2021). Differences between colonies were tested as

pairwise comparisons of estimated marginal means (Searle et al. 1980) with Bonferroni correction

with the 'emmeans' package (v.1.6.2-1).

Symbiosis reestablishment after menthol bleaching

Native Symbiodiniaceae characterization

G. fascicularis colonies were collected in March 2019 from Crescent Bay in Hong Kong. DNA

was successfully extracted from two colonies using the Qiagen DNeasy 96 Blood & Tissue kit.

The ITS2 marker region was amplified with the primers SYM_VAR_5.8S2 and SYM_VAR

following the PCR protocol by Hume et al. (2018), followed by paired-end sequencing on the

Illumina MiSeq platform (2 × 300 bp). Raw sequencing data was analyzed using the SymPortal

workflow remote instance (Hume et al. 2019), with results reported as post-MED ITS2 sequences

in relative abundances.

Menthol bleaching and inoculation of cultured Symbiodiniaceae

We explored the possibility of returning adult corals to the symbiotic state after menthol bleaching.

For this, we inoculated menthol-bleached coral fragments from Hong Kong with cultured

Symbiodiniaceae. Bleaching followed the same protocol described for the Red Sea colonies. Two

species of symbionts, *Cladocopium goreaui* and *Durusdinium trenchii* were chosen for targeted

exposure of the bleached fragments, as species from both of these genera are known to form stable

symbiosis with *G. fascicularis* across the South China Sea (Tong et al. 2017). Batch cultures of *C.*

goreaui (AIMS-SCF-055, ITS2 type-sequence C1, isolated from *Acropora tenuis*) and *D. trenchii*

(AIMS-SCF-088, ITS2 type-sequence D1-4, isolated from *A. muricata*) were obtained from the

Symbiont Culture Facility at the Australian Institute of Marine Science and maintained in f/2

media (Guillard 1975) under the same conditions as the corals. A total of 16 fragments were

inoculated with a concentration of 200 symbiont cells ml^{-1} in each 6-L aquarium. The

concentration of symbionts was divided in a 90:10 ratio (180 cells ml^{-1} *D. trenchii* : 20 cells ml^{-1}

C. goreaui). This ratio was chosen because members of *Cladocopium* are the most prevalent

symbionts found in *G. fascicularis* in Hong Kong (Huang et al. 2011; Tong et al. 2017) and across

its wide geographic range from the Red Sea to the South China Sea, where *Durusdinium* is only

occasionally found and only in comparably warm habitats (Dong et al. 2009; Ziegler et al. 2017;

Zhou et al. 2017). The symbiont aliquots were centrifuged to remove the culture media and washed

once with FASW before final resuspension in 10 mL FASW. The bleached fragments were first

fed before being exposed to the chosen symbionts. For this, a suspension of ReefRoids in ASW

was pipetted directly into the oral opening of each polyp. The same process was then immediately

repeated with the suspended symbionts. Aquarium filters were turned off for 1 h to allow the corals

to feed and to prevent removal of symbiont cells from the water column. Fragments were

inoculated twice a week for 6 weeks, until symbiosis reestablishment was visually confirmed.

Symbiodiniaceae populations *in hospite* were then characterized with fluorescent in-situ

hybridization (FISH) and flow cytometry (McIlroy et al. 2020).

Fluorescent In-Situ Hybridisation (FISH)

After 6 weeks of inoculation, a single polyp was removed from each fragment with a chisel. Tissue

was removed from the skeleton with an airbrush containing deionized water. The resulting slurry

was homogenized using a 20-µl pipette tip. 1.5 mL of homogenate was transferred to a 2-mL

Eppendorf tube and centrifuged at 500 rpm for seven minutes, after which the supernatant was

discarded. The remaining pellet was washed once before being preserved in 5X SET buffer and

stored at -80 °C for further analysis.

Samples were prepared following an adapted protocol from McIlroy et al. (2020). Briefly, after

washing with 5X SET in IGEPAL at 0.4 and 0.1 % vol/vol samples were split equally between

three 1.5-mL black Eppendorf tubes. Each sample had one tube treated with a SymC probe, one

tube with a SymD probe and one without any probe as a control. Both SymC (5'-

CTACCCAAGAACTTGCAGG-3') and SymD (5'-CCACCCAAGAACTCGCGTG-3') probes

were designed to have genus-level specificity for *Cladocopium* and *Durusdinium*, respectively,

and were labelled at 5' end with Alexa Fluor ® 532. After overnight hybridization at 45 °C (5X

SET, 0.1 % vol/vol IGE- PAL, 10 % vol/vol formamide and 100 pmol of the relevant probe),

samples were washed once in warm 1X SET, vortexed and resuspended in 500 µl of 1X SET for

same-day cytometric analysis on a BD FACSAria Fusion flow cytometer (BD Biosciences, CA).

A gating hierarchy was used to isolate cells of interest. Cells were screened for size and presence

of doublets (multiple cells stuck together) based on light-scatter parameters (forward-scatter (FSC)

and side-scatter (SSC) (Fig. S6a-c). Probe-positive cells were then distinguished based on their

position within the 582m vs SSC-A scatterplot (Fig. S6d,e). FlowJo™ Software (v.10.8.1, BD Life

Sciences) was used to determine the percentage of probe-positive cells (either *Cladocopium* or

Durusdinium), to identify the presence and the relative abundance of each algal genus within the

sample.

Completing gametogenic cycles and *ex-situ* spawning

Galaxea fascicularis in the GBR spawns between October and December, 0.05-200 minutes after

sunset (MAS) and 1 – 8 nights after full moon (NAFM) (Babcock et al. 1986; Baird et al. 2021).

To determine if the *ex situ* reproductive pattern remained in synchrony with the wild we followed

our colonies over three consecutive reproductive seasons (2019 – 2021).

To ascertain gamete development in each colony, two months prior to predicted spawning, and

two to four days prior to the full moon, whole polyps were removed from each colony and

longitudinal sections imaged (Canon 5d MKIII, Fig. 5a,d). Based on the stage of gamete

development in these sections *ex situ* spawning activity was predicted.

To record spawning observations each year, colonies were observed with red head torches during

the predicted spawning months. Observations commenced two NAFM and 30 mins prior to the

predicted spawning time, during which time the broodstock aquariums were isolated from the

filtration system. In addition, all internal water pumps were also turned off leaving the aquariums'

water static. Observations continued over multiple consecutive nights until all colonies that had

developed gametes, during that reproductive season, spawned. Spawning date and time of each

colony was recorded, with onset of spawning being denoted as the time of first gamete release.

Following release, gametes were pooled to allow fertilization to occur, and subsequent embryos

were reared and larvae settled following the method described by Craggs et al. (2019).

Results

Production and long-term rearing of bleached polyps

We produced single polyps (clonal replicates) of *G. fascicularis* colonies from the Red Sea and

bleached them with menthol. All polyps (n = 24, including additional polyps not used in this

experiment) survived the menthol treatment and appeared visually completely white. Similarly, all

menthol-bleached coral fragments from Hong Kong (n = 16) survived the treatment. Visual

inspection under a fluorescent stereomicroscope revealed no detectable algal cells in the bleached

polyps (Red Sea colonies, Fig. 2a), but inspection at higher magnification (compound fluorescent

microscope, ×400) revealed the presence of few scattered algal cells, limited to the base of the tentacles (Fig. 2b).

All polyps that were employed in the thermal performance experiment (n = 12) remained visually completely bleached and viable throughout the 10 days of temperature treatment, with the exception of two polyps that appeared dead on the 7th and 8th day of incubation, respectively. These polyps showed atrophic and unresponsive tentacles, and overall thinning of the soft tissue until it was not recognizable. Overall, we successfully maintained the bleached polyps for three weeks from the termination of the menthol treatment.

Fig. 2 Microscopic comparison between symbiotic and menthol-bleached polyps. Representative stereo and compound micrographs in bright field and UV light, with filters for chlorophyll (red) and coral tissue (green) autofluorescence. **a** Symbiodiniaceae cells are abundant in symbiotic polyps while not detectable in bleached polyps. **b** At higher magnifications, few algal cells are still detectable in bleached polyps only at the base of the tentacles (arrows).

Amenability to experimental manipulation and broad thermal tolerance

Symbiotic (untreated) polyps had positive net photosynthesis across all temperatures from 21 to 32 °C (Fig. 3). Net photosynthesis peaked at 23 °C (mean 13.05 µg O_2 cm^{-2} h^{-1}) and reached the minimum at 32 °C (mean 3.26 µg O_2 cm^{-2} h^{-1}). In contrast, respiration overall increased with temperature with minimum values at 21 °C (mean 5.77 µg O_2 cm^{-2} h^{-1}) and maximum at 32 °C (mean 25.70 µg O_2 cm^{-2} h^{-1}; Fig. 3). Bleached polyps, as expected by their lack of photosymbionts, exhibited a net oxygen depletion. Similar to symbiotic polyps, respiration in bleached polyps increased with temperature, both in light and dark incubations (Fig. 3). Overall, respiration in symbiotic polyps was higher than in bleached polyps ($P_{lmer} < 0.0001$).

While net photosynthesis was similar between colonies ($P_{lmer} > 0.05$), respiration was significantly higher in one colony ($P_{lmer} < 0.001$) compared to the other two (RS2-RS1: $P_{Bonf} < 0.01$; RS2-RS3: $P_{Bonf} < 0.01$; RS1-RS3: $P_{Bonf} > 0.05$; Fig. S4).

Fig. 3 Net photosynthesis and respiration rate of symbiotic states in *Galaxea fascicularis.*
Symbiotic (brown symbols) and bleached (white symbols) polyps in light and dark incubations.
Noise added (jittering) for ease of visualization, means per group depicted as diamonds. n = 12

Symbiosis re-establishment after bleaching

Inoculation with cultured symbionts resulted in successful symbiosis re-establishment for 100 %
of the samples (n = 16), which were macroscopically visibly symbiotic. Of these, nine samples
were successfully FISH hybridized, with the genus-specific probes identifying the presence of both
target genera in each sample. Individual variation in the relative ratio of each genus was evident
among the fragments, with *Durusdinium* contributing to 4 – 63 % of cells and *Cladocopium* to 37
– 82 % (Fig. 4, Tab. S2). A small, but significant proportion of symbionts remained unlabeled in
some samples. These may represent additional symbiont genera or be the result of variable probe
efficiency. *G. fascicularis* colonies from the same location were found to harbor *Cladocopium*
symbionts only (dominant ITS2 sequence types: C1 and C1c, Tab. S3).

Fig. 4 Symbiosis reestablishment in menthol-bleached adult corals inoculated with cultured

Symbiodiniaceae of the genera *Cladocopium* and *Durusdinium*. The presence and relative

abundance (%) of both genera was confirmed and quantified through fluorescent in-situ

hybridization (FISH) and flow cytometry.

Ex-situ spawning

Spawning was monitored each year over the three-year observation period for all 35 colonies.

Annual fecundity, as a percentage of colonies completing a full gametogenic cycle, was on average

56.2 % during these three spawning cycles (2019: 48.6 %, 2020: 45.7 %, and 2021: 74.3 %).

Spawning occurred in synchrony with wild predictions, with gametes being released between

October and January each year, between 5 - 259 MAS and 4 - 13 NAFM (Fig. 5b-c & e-f, Fig. 6).

Following embryo rearing, Symbiodiniaceae were observed within primary polyps 14 days post

settlement (Fig. 5g). These originated from the adult colonies housed in the same system and taken

up via the water column. Polyps were fully developed 40 days post settlement (Fig. 5h) and grew

into small colonies within the first six months (Fig. 5i).

Fig. 5 Representative images of female and male colonies of *Galaxea fascicularis* spawning

***ex situ* and gamete development**. **a** Longitudinal section of female polyp with pinkish-red

pigmented oocytes. **b** Female colony releasing oocyte bundle *ex situ*. **c** Close-up of female bundle

post release. **d** Longitudinal section of male polyp with white oocytes. **e** Male colony releasing

oocytes/sperm bundle *ex situ*. **f** Close-up of male gamete bundles post release undergoing

disassociation. **g** Newly settled *G. fascicularis* primary polyps showing initial Symbiodiniaceae

(arrow) uptake 14 days post settlement. **h** Primary polyp 40 days post settlement. **i** Multiple

colonies six months post settlement. Scale bar: a-h = 1 mm, i = 1 cm.

Fig. 6 *Galaxea fascicularis ex situ* **spawning periodicity over three spawning cycles (2019 –**
2021). **a** Spawning time, recorded as the first gamete release in Minutes After Sunset (MAS) by
date. **b** Night After Full Moon (NAFM) of gamete release by date. **c** Comparison of MAS and
NAFM values over three *ex situ* spawning cycles. 2019 n = 28, 2020 n = 19, 2021 n = 69.

Discussion

Here, we explored the potential of *G. fascicularis* as a novel coral model for symbiosis research
by assessing its amenability to aquarium rearing, experimental handling, and symbiosis
manipulation.

Successful bleaching with menthol

We showed that *G. fascicularis* can be readily and reliably rendered aposymbiotic through menthol
bleaching, similar to other cnidarian model systems (e.g., Matthews et al. 2015; Röthig et al. 2021).
We lowered the menthol concentration in the protocol developed by Wang et al. (2012) to further
limit stress on the host, and indeed all coral polyps survived this bleaching procedure on two
independently replicated occasions. Overall, menthol treatment was very effective, with only a
few algal cells remaining at the base of the tentacles. From a photophysiological perspective, these

scant remnants can be considered negligible, as illustrated by the light and dark oxygen production

data. Furthermore, bleached polyps remained aposymbiotic for weeks and we therefore considered

this bleaching protocol suitable for experimental investigation of aposymbiotic *G. fascicularis* in

physiological studies and symbiosis manipulation.

Notably, besides Symbiodiniaceae, reef-building corals also associate with other microorganisms

(i.e., bacteria, viruses, and other unicellular algae and protists) which constitute the so-called coral

holobiont (Rosenberg et al. 2007). Although this has only been appreciated more recently,

holobiont composition and dynamics are now understood to underlie coral health and adaptability

(Pogoreutz et al. 2020). As menthol has photo-inhibitory (Wang et al. 2012; Clowez et al. 2021)

and antimicrobial properties (İşcan et al. 2002), further studies should investigate the effects of

menthol bleaching on the remaining members of the microbiome.

Tractability of symbiotic and aposymbiotic *G. fascicularis*

Tractability, as amenability to experimental work and laboratory conditions, is one of the most

important requirements for model organisms. The *G. fascicularis* polyps and fragments could be

maintained for three and ten weeks respectively in simple systems consisting of small independent

tanks (5-6 L) with regular feeding and basic water quality care. These time frames match the

requirements for most experimental designs, and we believe could be further extended. Indeed,

symbiotic individuals looked healthy and responsive throughout the whole period, testifying to a

comparatively hardy coral with only low demands on rearing conditions. Additionally, the positive

net photosynthetic rate across a large temperature range confirmed the broad thermal tolerance of

this species (Al-Horani 2005; McIlroy et al. 2019), which was retained in the simplified single

polyp application.

Moreover, bleached polyps could be maintained in the simple experimental system for three weeks

even with the additional stress of the thermal experiment. However, they eventually showed signs

of deteriorating condition indicating that not all nutritional needs were met. Thus one priority is to

develop bespoke feed for long-term rearing of bleached corals (Wang et al. 2012).

Thermal performance of symbiotic and aposymbiotic *G. fascicularis*

Studies that compare symbiotic and aposymbiotic coral holobionts are particularly informative for

understanding coral response to climate change, and therefore we investigated the comparability

between symbiotic and aposymbiotic *G. fascicularis* polyps. As expected, these differed in their

photosynthetic rates, where the latter were net oxygen consumers even under illumination. Overall,

symbiotic polyps had higher respiration rates than aposymbiotic polyps, which could be ascribed

to the additional metabolic burden of the associated Symbiodiniaceae, as well as a greater

metabolic capacity derived from the availability of photosynthates (Muscatine et al. 1981; Al-

Horani et al. 2003). For all polyps, respiration rates overall increased with temperature in line with

optimum thermal performance of corals and macroalgae from the central Red Sea (27-33 ˚C)

(Anton et al. 2020).

Interestingly, we observed intraspecific variability in physiological performance. One colony

(RS2) had significantly higher respiration rates than the others (Fig. S4). This manifested

exclusively under dark conditions and was therefore more likely linked to host rather than

symbiont identity. This variability warrants further investigations to better characterize host

genotypes, as in the long term it will be necessary to establish representative host clonal lines.

Symbiosis reestablishment in adult corals

We showed that adult bleached *G. fascicularis* can successfully reestablished symbiosis with

cultured Symbiodiniaceae. The experimental repopulation of adult corals with cultured non-native

symbionts has only recently been demonstrated (Scharfenstein et al. 2022), and it is considered

particularly relevant in the context of symbiosis manipulations to enhance heat tolerance in reef-

building corals (van Oppen et al. 2015). Notably, here this was further explored with two non-

native algal cultures during simultaneous exposure. The first, *D. trenchii* (ITS2 type D1-4), has

never been found in *G. fascicularis* from Hong Kong (Huang et al. 2011; Tong et al. 2017) nor in

our characterization of Symbiodiniaceae from that site specifically (two other colonies from the

same location, Tab. S3) and can therefore confidently be considered to be non-native. The second,

C. goreaui (ITS2 type C1), was isolated from the coral species *A. tenuis* from Australia. Even

though the dominant C1 ITS2-sequence is also present in *G. fascicularis* from Hong Kong, the

high coral host-Symbiodiniaceae specificity, large radiation of C1-like Symbiodiniaceae

(LaJeunesse 2005; Thornhill et al. 2014), and the consistent occurrence of the minor sequence C1c

in addition to C1 in our samples, implicate the *C. goreaui* culture as non-native in *G. fascicularis*.

However, the higher similarity with the native symbionts likely explains the more efficient

establishment of the *C. goreaui* culture than the *D. trenchii* culture despite a reversed (10:90) ratio

in supply.

Galaxea fascicularis has generally been reported to host taxa from the genera *Cladocopium* and

Durusdinium, both exclusively and simultaneously, and with an equatorial-ward shift from

Cladocopium dominance to *Durusdinium* dominance (Huang et al. 2011, and references therein).

At Hong Kong's latitude and at our rearing temperature, *G. fascicularis* is expected to be

exclusively associated with *Cladocopium* (Huang et al. 2011; Tong et al. 2017; Zhou et al. 2017),

yet the targeted inoculation produced mixed-genus associations. This demonstrated how targeted

inoculation can successfully shift Symbiodiniaceae community composition towards a higher

proportion of heat tolerant strains without thermal treatment (Silverstein et al. 2015). Although the

longevity of these partnerships is still unclear, these are promising results for future development

of *in hospite* studies on the role of temperature on inter-partner symbiosis dynamics and symbiont

recognition mechanisms (Bove et al. 2022). The success and potential insights from such

experimental approaches will rely on future efforts to establish a variety of coral host-specific

Symbiodiniaceae cultures (including the establishment of specific cultures from *G. fascicularis*)

for further study.

Of note, repopulation is possible even with lower concentrations of Symbiodiniaceae, as we

observed from earlier qualitative trials (details in Supplementary Text). There, we exposed the

menthol bleached polyps after they were used in the thermal performance experiment to

Symbiodiniaceae through the water medium (symbionts shed by nearby symbiotic colonies) or

additionally in combination with a one-time inoculum of freshly isolated symbionts from

symbiotic conspecifics. The success rates of these qualitative trials of ~30 % and 43 %,

respectively, were lower than in the Hong Kong experiment (100 %) yet remarkable considering

the poorer health conditions of the polyps at the time of re-exposure to symbionts (i.e., after

repeated incubations through the 21-32 °C temperature excursion), and the expectedly smaller

number of symbionts available for uptake. In these trials, bleached hosts retained the ability to

feed, which is known to facilitate symbiont uptake (Fitt and Trench 1983) and likely contributed

to symbiosis re-establishment. In the long-term, the polyps that reestablished symbiosis grew into

small colonies and became visually indistinguishable from the untreated controls (Fig. S5). Taken

together, these observations attest to the suitability of *G. fascicularis* to experimental symbiosis

manipulation.

Ex situ **spawning**

Galaxea fascicularis is a good model species candidate for sexual reproductive work due its high

survivorship in aquarium conditions. As demonstrated herein, a combination of seasonal

programming and husbandry enabled the species to complete a full gametogenic cycle *ex situ* over

multiple years, which closely mimics that observed in the wild (Babcock et al. 1986; Baird et al.

2021). Furthermore, we showed *in vitro* fertilization, embryological development, and *ex situ*

larval settlement resulting in spat of known ages. As the age at which the species reaches sexual

maturity is currently unknown, future studies should focus on this aspect, which is also important

for producing an F2 generation and for closing the species life cycle *ex situ* (Craggs et al. 2020).

Access to a F2 generation of known parental lineage will provide a platform to conduct

experiments in areas such as mapping quantitative trait loci (Zhang 2012) or phenotypic traits such

as growth, disease resistance, or thermal tolerance (van Oppen et al. 2015). As *Galaxea* must

acquire their symbionts from the seawater (horizontal transmission, with symbionts visible after

the development of tentacles in settled polyps, Lin et al. (2022)), *ex situ* spawning also opens

possibilities for studies on the dynamics of initial symbiosis establishment and how this is affected

by symbiont identity, heritability in symbiont selection, and for rearing of axenic or gnotobiotic

coral lineages, paralleling procedures established for *Hydra* (Fraune et al. 2015).

Conclusions

The study of coral-algal symbiosis has been constrained by the difficulties of maintaining corals

ex situ and of manipulating the coral-algal association to unravel symbiosis functioning. Here, we

showed that adopting *G. fascicularis* as a model system can alleviate these limitations. Its

demonstrated compatibility to rearing in simplified systems and experimental applications,

together with the repeated spawning in aquaria, confirm the high tractability of this coral species.

Further, the possibility to readily obtain aposymbiotic individuals suitable for detailed study and

symbiosis re-establishment set the stage for exciting future developments in coral symbiosis

research which is necessary to understand corals' potential to cope with changing oceans.

Besides these strengths, our study also highlights areas worth future efforts on the path to

developing the *G. fascicularis* model system further. These include: i) better characterization of

the effect of menthol on the rest of the coral microbiome (e.g., bacteria); ii) development of custom

feed for long-term maintenance of bleached individuals; iii) assessment of long-term persistence

of non-native Symbiodiniaceae, and testing of a variety of symbiont strains to understand the

breadth of symbiont diversity that can be accommodated by the *G. fascicularis* host; iv) further

characterization of host colonies and genotypes with the aim of establishing clonal lines, and from

these, isolate and establish Symbiodiniaceae cultures, mirroring for example the development of

the Aiptasia model; and v) development of collaborative platforms and open-access community

resources around this emerging model system to accelerate research and discovery.

Acknowledgements

We thank Dr. Jafargholi Imani and Dr. Kathrin Ehlers (both JLU) for access and guidance in the

fluorescence microscope facilities, André Dietzmann and Catarina P. P. Martins (both JLU) for

assistance during experiments, and the Li Ka Shing Faculty of Medicine Core Facility (HKU) for

the flow cytometry equipment.

This study is part of the 'Ocean2100' global change simulation project of the Colombian-German

Center of Excellence in Marine Sciences (CEMarin) funded by the German Academic Exchange

Service. We acknowledge financial support by the German Research Foundation (DFG, Project:

469364832) to MZ and Hong Kong Research Grants Council #17117221 to SEM and #17108620

to DMB.

Data and code availability

Data and R scripts used for this study are available at:

https://github.com/sPuntinG/Galaxea_Coral_Model/

Author contributions

GP and MZ conceived and designed the study. GP, JC, RH, KEE, SM produced and analyzed data.

MS, DMB, MZ contributed reagents/materials/analysis tools. GP and MZ wrote the first draft with

contributions from all authors.

Conflict of Interest

The authors declare no conflict of interest for this submission.

References

Al-Horani FA (2005) Effects of changing seawater temperature on photosynthesis and
    calcification in the scleractinian coral *Galaxea fascicularis*, measured with O_2, Ca^{2+} and pH
    microsensors. Sci Mar 69:347–354
    http://scientiamarina.revistas.csic.es/index.php/scientiamarina/article/view/263/261

Al-Horani FA, Al-Moghrabi SM, de Beer D (2003) The mechanism of calcification and its
    relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*.
    Mar Biol 142:419–426 http://link.springer.com/10.1007/s00227-002-0981-8

Anton A, Randle JL, Garcia FC, Rossbach S, Ellis JI, Weinzierl M, Duarte CM (2020)
    Differential thermal tolerance between algae and corals may trigger the proliferation of
    algae in coral reefs. Glob Chang Biol 26:4316–4327
    https://onlinelibrary.wiley.com/doi/10.1111/gcb.15141

Babcock RC, Bull GD, Harrison PL, Heyward AJ, Oliver JK, Wallace CC, Willis BL (1986)
    Synchronous spawnings of 105 scleractinian coral species on the Great Barrier Reef. Mar
    Biol 90:379–394 http://link.springer.com/10.1007/BF00428562

Baird AH, Guest JR, Edwards AJ, Bauman AG, Bouwmeester J, Mera H, Abrego D, Alvarez-
    Noriega M, Babcock RC, Barbosa MB, Bonito V, Burt J, Cabaitan PC, Chang C-F,
    Chavanich S, Chen CA, Chen C-J, Chen W-J, Chung F-C, Connolly SR, Cumbo VR,
    Dornelas M, Doropoulos C, Eyal G, Eyal-Sham L, Fadli N, Figueiredo J, Flot J-F, Gan S-
    H, Gomez E, Graham EM, Grinblat M, Gutiérrez-Isaza N, Harii S, Harrison PL, Hatta M,
    Ho NAJ, Hoarau G, Hoogenboom M, Howells EJ, Iguchi A, Isomura N, Jamodiong EA,
    Jandang S, Keyse J, Kitanobo S, Kongjandtre N, Kuo C-Y, Ligson C, Lin C-H, Low J, Loya
    Y, Maboloc EA, Madin JS, Mezaki T, Min C, Morita M, Moya A, Neo S-H, Nitschke MR,
    Nojima S, Nozawa Y, Piromvaragorn S, Plathong S, Puill-Stephan E, Quigley K, Ramirez-
    Portilla C, Ricardo G, Sakai K, Sampayo E, Shlesinger T, Sikim L, Simpson C, Sims CA,
    Sinniger F, Spiji DA, Tabalanza T, Tan C-H, Terraneo TI, Torda G, True J, Tun K,
    Vicentuan K, Viyakarn V, Waheed Z, Ward S, Willis B, Woods RM, Woolsey ES,
    Yamamoto HH, Yusuf S (2021) An Indo-Pacific coral spawning database. Sci Data 8:35
    http://www.nature.com/articles/s41597-020-00793-8

Bove CB, Ingersoll MV, Davies SW (2022) Help Me, Symbionts, You're My Only Hope:
Approaches to Accelerate our Understanding of Coral Holobiont Interactions. Integr Comp
Biol https://academic.oup.com/icb/advance-article/doi/10.1093/icb/icac141/6696962

Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G (2008) MeshLab: an
Open-Source Mesh Processing Tool.

Clowez S, Renicke C, Pringle JR, Grossman AR (2021) Impact of menthol on growth and
photosynthetic function of *Breviolum minutum* (Dinoflagellata, Dinophyceae,
Symbiodiniaceae) and interactions with its *Aiptasia* host. J Phycol 57:245–257
https://onlinelibrary.wiley.com/doi/10.1111/jpy.13081

Craggs J, Guest J, Brett A, Davis M, Sweet M (2018) Maintaining natural spawning timing in
Acropora corals following long distance inter-continental transportation. J Zoo Aquarium
Res 6:30–36

Craggs J, Guest JR, Bulling M, Sweet M (2019) Ex situ co culturing of the sea urchin, *Mespilia*
globulus and the coral *Acropora millepora* enhances early post-settlement survivorship. Sci
Rep 9:12984 http://www.nature.com/articles/s41598-019-49447-9

Craggs J, Guest JR, Davis M, Simmons J, Dashti E, Sweet M (2017) Inducing broadcast coral
spawning ex situ: closed system mesocosm design and husbandry protocol. Ecol Evol
7:11066 https://onlinelibrary.wiley.com/doi/10.1002/ece3.3538

Craggs J, Guest JR, Davis M, Sweet M (2020) Completing the life cycle of a broadcast spawning
coral in a closed mesocosm. Invertebr Reprod Dev 64:244–247
https://www.tandfonline.com/doi/full/10.1080/07924259.2020.1759704

Dimond J, Carrington E (2007) Temporal variation in the symbiosis and growth of the temperate
scleractinian coral *Astrangia poculata*. Mar Ecol Prog Ser 348:161–172 http://www.int-
res.com/abstracts/meps/v348/p161-172/

Dong Z-J, Huang H, Huang L-M, Li Y-C (2009) Diversity of symbiotic algae of the genus
Symbiodinium in scleractinian corals of the Xisha Islands in the South China Sea. J Syst
Evol 47:321–326 https://onlinelibrary.wiley.com/doi/10.1111/j.1759-6831.2009.00034.x

Falkowski PG, Dubinsky Z, Muscatine L, Porter JW (1984) Light and the bioenergetics of a

symbiotic coral. Bioscience 34:705–709 https://academic.oup.com/bioscience/article-

lookup/doi/10.2307/1309663

Fitt WK, Trench RK (1983) Endocytosis of the symbiotic dinoflagellate *Symbiodinium*

microadriaticum Freudenthal by endodermal cells of the scyphistomae of *Cassiopeia*

xamachana and resistance of the algae to host digestion. J Cell Sci 64:195–212

Fraune S, Anton-Erxleben F, Augustin R, Franzenburg S, Knop M, Schröder K, Willoweit-Ohl

D, Bosch TCG (2015) Bacteria–bacteria interactions within the microbiota of the ancestral

metazoan *Hydra* contribute to fungal resistance. ISME J 9:1543–1556

http://www.nature.com/articles/ismej2014239

Galliot B (2012) *Hydra*, a fruitful model system for 270 years. Int J Dev Biol 56:411–423

http://www.intjdevbiol.com/paper.php?doi=120086bg

Gates RD, Baghdasarian G, Muscatine L (1992) Temperature stress causes host cell detachment

in symbiotic cnidarians: implications for coral bleaching. Biol Bull 182:324–332

https://www.journals.uchicago.edu/doi/10.2307/1542252

Gattuso J-P, Allemand D, Frankignoulle M (1999) Photosynthesis and Calcification at Cellular,

Organismal and Community Levels in Coral Reefs: A Review on Interactions and Control

by Carbonate Chemistry. Am Zool 39:160–183 https://academic.oup.com/icb/article-

lookup/doi/10.1093/icb/39.1.160

Glynn PW, D'Croz L (1990) Experimental evidence for high temperature stress as the cause of

El Niño-coincident coral mortality. Coral Reefs 8:181–191

Guillard RRL (1975) Culture of Phytoplankton for Feeding Marine Invertebrates. In: Smith,

W.L., Chanley M.H. (eds) Culture of Marine Invertebrate Animals. Springer US, Boston,

MA, pp 29–60 http://link.springer.com/10.1007/978-1-4615-8714-9_3

Harrison PL (1988) Pseudo-gynodioecy: an unusual breeding system in the scleractinian coral

Galaxea fascicularis. 699–

705https://researchportal.scu.edu.au/esploro/outputs/conferencePaper/Pseudo-gynodioecy-

an-unusual-breeding-system-in-the-scleractinian-coral-Galaxea-

fascicularia/991012821519702368

Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral

reefs. Mar Freshw Res 50:839–866 http://www.publish.csiro.au/?paper=MF99078

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadia G, Raymundo L, Williams G,

Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of

the Paris Agreement. Sci Rep 6:39666 http://www.nature.com/articles/srep39666

Huang H, Dong Z, Huang L, Yang J, Di B, Li Y, Zhou G, Zhang C (2011) Latitudinal variation

in algal symbionts within the scleractinian coral Galaxea fascicularis in the South China

Sea. Mar Biol Res 7:208–211

http://www.tandfonline.com/doi/abs/10.1080/17451000.2010.489616

Hume BCC, Smith EG, Ziegler M, Warrington HJM, Burt JA, LaJeunesse TC, Wiedenmann J,

Voolstra CR (2019) SymPortal: A novel analytical framework and platform for coral algal

symbiont next-generation sequencing ITS2 profiling. Mol Ecol Resour 19:1063–1080

Hume BCC, Ziegler M, Poulain J, Pochon X, Romac S, Boissin E, de Vargas C, Planes S,

Wincker P, Voolstra CR (2018) An improved primer set and amplification protocol with

increased specificity and sensitivity targeting the *Symbiodinium* ITS2 region. PeerJ 6:e4816

https://peerj.com/articles/4816

İşcan G, Kirimer N, Kürkcüoğlu M, Hüsnü Can Başer, Demirci F (2002) Antimicrobial

screening of *Mentha piperita* essential oils. J Agric Food Chem 50:3943–3946

https://doi.org/10.1021/jf011476k

Keshavmurthy S, Hsu CM, Kuo CY, Denis V, Leung JKL, Fontana S, Hsieh HJ, Tsai W Sen, Su

WC, Chen CA (2012) Larval development of fertilized "pseudo-gynodioecious" eggs

suggests a sexual pattern of gynodioecy in *Galaxea fascicularis* (Scleractinia: Euphyllidae).

Zool Stud 51:143–149

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. J Stat Softw 82:1–26

https://www.jstatsoft.org/index.php/jss/article/view/v082i13

LaJeunesse TC (2005) "Species" Radiations of Symbiotic Dinoflagellates in the Atlantic and

Indo-Pacific Since the Miocene-Pliocene Transition. Mol Biol Evol 22:570–581

https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msi042

Liew YJ, Aranda M, Voolstra CR (2016) Reefgenomics.Org - a repository for marine genomics
data. Database 2016:baw152 https://academic.oup.com/database/article-
lookup/doi/10.1093/database/baw152

Lin C, Kang C-M, Huang C-Y, Li H-H, Tsai S (2022) Study on the Development and Growth of
Coral Larvae. Appl Sci 12:5255 https://www.mdpi.com/2076-3417/12/10/5255

Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D (2021) performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. J Open Source
Softw 6:3139 https://joss.theoj.org/papers/10.21105/joss.03139

Marshall AT, Clode P (2004) Calcification rate and the effect of temperature in a zooxanthellate
and an azooxanthellate scleractinian reef coral. Coral Reefs 23:218–224
http://link.springer.com/10.1007/s00338-004-0369-y

Marshall PA, Baird AH (2000) Bleaching of corals on the Great Barrier Reef: differential
susceptibilities among taxa. Coral Reefs 19:155–163
http://link.springer.com/10.1007/s003380000086

Matthews JL, Sproles AE, Oakley CA, Grossman AR, Weis VM, Davy SK (2015) Menthol-
induced bleaching rapidly and effectively provides experimental aposymbiotic sea
anemones (*Aiptasia* sp.) for symbiosis investigations. J Exp Biol 219:306–310
http://jeb.biologists.org/cgi/doi/10.1242/jeb.128934

McIlroy SE, Thompson PD, Yuan FL, Bonebrake TC, Baker DM (2019) Subtropical thermal
variation supports persistence of corals but limits productivity of coral reefs. Proc R Soc B
Biol Sci 286:20190882 https://royalsocietypublishing.org/doi/10.1098/rspb.2019.0882

McIlroy SE, Wong JCY, Baker DM (2020) Competitive traits of coral symbionts may alter the
structure and function of the microbiome. ISME J 14:2424–2432
https://www.nature.com/articles/s41396-020-0697-0

McLachlan RH, Price JT, Solomon SL, Grottoli AG (2020) Thirty years of coral heat-stress
experiments: a review of methods. Coral Reefs 39:885–902
https://link.springer.com/10.1007/s00338-020-01931-9

Muscatine L, R. McCloskey L, E. Marian R (1981) Estimating the daily contribution of carbon

from zooxanthellae to coral animal respiration. Limnol Oceanogr 26:601–611

http://doi.wiley.com/10.4319/lo.1981.26.4.0601

Niu W, Huang H, Lin R, Chen C-H, Shen K-N, Hsiao C-D (2016) The complete mitogenome of

the Galaxy Coral, *Galaxea fascicularis* (Cnidaria: Oculinidae). Mitochondrial DNA Part B

1:10–11 https://www.tandfonline.com/doi/full/10.1080/23802359.2015.1137796

van Oppen MJH, Oliver JK, Putnam HM, Gates RD (2015) Building coral reef resilience

through assisted evolution. Proc Natl Acad Sci 112:2307–2313

http://www.pnas.org/lookup/doi/10.1073/pnas.1422301112

Pavia RTB, Estacion JS (2019) Survival and growth of isolated polyps of *Galaxea fascicularis*

(Linnaeus 1767) on six kinds of culture substrates: implications for mariculture, aquarium

culture, and conservation. J World Aquac Soc 50:219–230

https://onlinelibrary.wiley.com/doi/10.1111/jwas.12538

Pogoreutz C, Voolstra CR, Rädecker N, Weis V (2020) The coral holobiont highlights the

dependence of cnidarian animal hosts on their associated microbes. Cellular dialogues in the

holobiont. pp 91–118

Puntin G, Sweet M, Fraune S, Medina M, Sharp K, Weis VM, Ziegler M (2022) Harnessing the

Power of Model Organisms To Unravel Microbial Functions in the Coral Holobiont.

Microbiol Mol Biol Rev https://journals.asm.org/doi/10.1128/mmbr.00053-22

R Core Team (2021) R: A Language and Environment for Statistical Computing. https://www.r-

project.org/ https://www.r-project.org/

Rades M, Schubert P, Ziegler M, Kröckel M, Reichert J (2022) Building plan for a temperature-

controlled multi-point stirring incubator. https://www.protocols.io/view/building-plan-for-a-

temperature-controlled-multi-p-dm6gpb34dlzp/v1 https://www.protocols.io/view/building-

plan-for-a-temperature-controlled-multi-p-dm6gpb34dlzp/v1

Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I (2007) The role of

microorganisms in coral health, disease and evolution. Nat Rev Microbiol 5:355–362

Röthig T, Puntin G, Wong JCY, Burian A, McLeod W, Baker DM (2021) Holobiont nitrogen

control and its potential for eutrophication resistance in an obligate photosymbiotic

jellyfish. Microbiome 9:127

https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-021-01075-0

Scharfenstein HJ, Chan WY, Buerger P, Humphrey C, van Oppen MJH (2022) Evidence for de

novo acquisition of microalgal symbionts by bleached adult corals. ISME J 16:1676–1679

https://www.nature.com/articles/s41396-022-01203-0

Schneider K, Erez J (2006) The effect of carbonate chemistry on calcification and photosynthesis

in the hermatypic coral *Acropora eurystoma*. Limnol Oceanogr 51:1284–1293

https://doi.org/10.4319/lo.2006.51.3.1284

Schubert P, Wilke T (2018) Coral Microcosms: Challenges and Opportunities for Global Change

Biology. Corals in a Changing World. InTech, pp 143–175

http://www.intechopen.com/books/corals-in-a-changing-world/coral-microcosms-

challenges-and-opportunities-for-global-change-biology

Searle SR, Speed FM, Milliken GA (1980) Population Marginal Means in the Linear Model: An

Alternative to Least Squares Means. Am Stat 34:216–221

http://www.tandfonline.com/doi/abs/10.1080/00031305.1980.10483031

Silverstein RN, Cunning R, Baker AC (2015) Change in algal symbiont communities after

bleaching, not prior heat exposure, increases heat tolerance of reef corals. Glob Chang Biol

21:236–249 https://onlinelibrary.wiley.com/doi/10.1111/gcb.12706

Thornhill DJ, Lewis AM, Wham DC, LaJeunesse TC (2014) HOST-SPECIALIST LINEAGES

DOMINATE THE ADAPTIVE RADIATION OF REEF CORAL ENDOSYMBIONTS.

Evolution (N Y) 68:352–367 https://onlinelibrary.wiley.com/doi/10.1111/evo.12270

Tong H, Cai L, Zhou G, Yuan T, Zhang W, Tian R, Huang H, Qian P-Y (2017) Temperature

shapes coral-algal symbiosis in the South China Sea. Sci Rep 7:1–12

http://dx.doi.org/10.1038/srep40118

Veron JEN, Stafford-Smith MG, Turak E, DeVantier LM (2016) Corals of the World.

http://www.coralsoftheworld.org/species_factsheets/species_factsheet_distribution/galaxea-

fascicularis/?version=0.01

http://www.coralsoftheworld.org/species_factsheets/species_factsheet_distribution/galaxea-
fascicularis/?version=0.01

Voolstra CR (2013) A journey into the wild of the cnidarian model system *Aiptasia* and its
symbionts. Mol Ecol 22:4366–4368

Wang J-T, Chen Y-Y, Tew KS, Meng P-J, Chen CA (2012) Physiological and biochemical
performances of menthol-induced aposymbiotic corals. PLoS One 7:e46406
https://dx.plos.org/10.1371/journal.pone.0046406

Weis VM, Davy SK, Hoegh-Guldberg O, Rodriguez-Lanetty M, Pringle JR (2008) Cell biology
in model systems as the key to understanding corals. Trends Ecol Evol 23:369–376
https://linkinghub.elsevier.com/retrieve/pii/S0169534708001614

Wepfer P, Nakajima Y, Hui F, Mitarai S, Economo E (2020a) Metacommunity ecology of
Symbiodiniaceae hosted by the coral *Galaxea fascicularis*. Mar Ecol Prog Ser 633:71–87
https://www.int-res.com/abstracts/meps/v633/p71-87/

Wepfer PH, Nakajima Y, Sutthacheep M, Radice VZ, Richards Z, Ang P, Terraneo T, Sudek M,
Fujimura A, Toonen RJ, Mikheyev AS, Economo EP, Mitarai S (2020b) Evolutionary
biogeography of the reef-building coral genus *Galaxea* across the Indo-Pacific ocean. Mol
Phylogenet Evol 151:106905 https://doi.org/10.1016/j.ympev.2020.106905

Wickham H (2016) Ggplot2: Elegant graphics for data analysis. Springer International
Publishing, Cham, Switzerland https://books.google.de/books?id=RTMFswEACAAJ

Zhang Y (2012) Quantitative Trait Loci (QTL). Humana Press, Totowa, NJ
http://link.springer.com/10.1007/978-1-61779-785-9

Zhou G, Cai L, Li Y, Tong H, Jiang L, Zhang Y, Lei X, Guo M, Liu S, Qian P-Y, Huang H
(2017) Temperature-driven local acclimatization of *Symbiodnium* hosted by the coral
Galaxea fascicularis at Hainan Island, China. Front Microbiol 8:1–9
http://journal.frontiersin.org/article/10.3389/fmicb.2017.02487/full

Ziegler M, Arif C, Burt JA, Dobretsov S, Roder C, LaJeunesse TC, Voolstra CR (2017)
Biogeography and molecular diversity of coral symbionts in the genus Symbiodinium
around the Arabian Peninsula. J Biogeogr 44:674–686

https://onlinelibrary.wiley.com/doi/10.1111/jbi.12913

Ziegler M, Roder C, Büchel C, Voolstra C (2015) Niche acclimatization in Red Sea corals is

dependent on flexibility of host-symbiont association. Mar Ecol Prog Ser 533:149–161

http://www.int-res.com/abstracts/meps/v533/p149-161/

775