1
2
Temperature sensitivity of the interspecific interaction strength of coastal marine fish communities

Authors:

Masayuki Ushio[1,2,3*], Tetsuya Sado[4], Takehiko Fukuchi[4], Sachia Sasano[5,6], Reiji Masuda[5],

Yutaka Osada[7], Masaki Miya[4,*]

Email: MU, ong8181@gmail.com; TS, zacco_evolans@yahoo.co.jp; TF,

fukuchi_takehiko0728@yahoo.co.jp; SS, sasano.sachia.75e@gmail.com; RM,

masuda.reiji.3w@kyoto-u.ac.jp; YO, ytosada@gmail.com; MM, masaki_miya@me.com

Author Affiliations:

[1]Hakubi Center, Kyoto University, Kyoto 606-8501, Japan,

[2]Center for Ecological Research, Kyoto University, Otsu 520-2113, Japan,

[3]Department of Ocean Science, The Hong Kong University of Science and Technology, Clear

Water Bay, Kowloon, Hong Kong SAR, China

[4]Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan,

[5]Maizuru Fisheries Research Station, Kyoto University, Maizuru, Kyoto 625-0086, Japan,

[6]Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Ishigaki,

Okinawa 907-0451, Japan,

[7]Graduate School of Life Sciences, Tohoku University, Sendai 980-8578, Japan

Keywords: Community dynamics; Information flow; Environmental DNA; Fish; Interaction

network; Quantitative sequencing

Data availability statement:

Source scripts for the analyses and figure generations are currently publicly available in

Github (https://github.com/ong8181/eDNA-BosoFish-network). DNA data is deposited

DDBJ Sequence Read Archive (DRA submission ID = DRA014111) and will make public

upon acceptance of the manuscript.

Abstract:

The effects of temperature on interaction strengths are important for understanding and

forecasting how global climate change impacts marine ecosystems; however, tracking and

quantifying interactions of marine fish species is practically difficult especially under field

conditions, and thus, how temperature influences their interaction strengths under field

conditions remains poorly understood. We herein performed quantitative fish environmental

DNA (eDNA) metabarcoding on 550 seawater samples that were collected twice a month

from 11 coastal sites for two years in the Boso Peninsula, Japan, and analyzed eDNA

monitoring data using nonlinear time-series analytical tools. We detected fish-fish

interactions as information flow between eDNA time series, reconstructed interaction

networks for the top 50 frequently detected species, and quantified pairwise, fluctuating

interaction strengths using nonlinear time series analysis methods. Although there was a large

variation, water temperature generally influenced fish-fish interaction strengths. The impact

of water temperature on interspecific interaction strengths varied among fish species,

suggesting that fish species identity influences the temperature effects on interactions. For

example, interaction strengths that *Halichoeres tenuispinis* and *Microcanthus stringatus*

received strongly increased with water temperature, while those of *Engraulis japonicus* and

Girella punctata decreased with water temperature. An increase in water temperature induced

by global climate change may change fish interactions in a complex way, which consequently

influences marine community dynamics and stability. Our research demonstrates a practical

research framework to study the effects of environmental variables on interaction strengths of

marine communities under field conditions, and thus, would contribute to further

understanding and predicting natural marine ecosystem dynamics.

Introduction

Interspecific interactions are key to understanding and predicting the dynamics of ecological communities (May 1972; Wootton & Emmerson 2005; Tang *et al.* 2014; Wootton & Stouffer 2015). Theoretical and empirical studies have shown that various properties of interspecific interactions, namely, the number of interactions, sign (positive or negative), strength, and pair-wise correlations of interactions (e.g., prey-predator interactions), influence the dynamics, stability, and diversity of ecological communities in terrestrial and aquatic ecosystems (Mougi & Kondoh 2012; Tang *et al.* 2014; Allesina *et al.* 2015; Ushio *et al.* 2018a; Ratzke *et al.* 2020; Ushio 2022a). Interaction strength is a fundamental property, and previous studies examined the relationship between interaction strengths and ecological community properties (Wootton & Emmerson 2005; Wootton & Stouffer 2015). The dominance of weak interactions stabilizes the dynamics of a natural fish community (Ushio *et al.* 2018a). Interaction strengths have been reported to decrease with increases in the diversity of experimental microbial communities (Ratzke *et al.* 2020), and this holds true even for more diverse ecological communities under field conditions (Ushio 2022a).

Environmental variables exert significant effects on interspecific interaction strengths. The effects of temperature on interaction strengths are important for understanding and forecasting the impact of the ongoing global climate change on ecosystems. The relationship between temperature and interaction strengths has been investigated in terrestrial and aquatic ecosystems for decades (Coley & Aide 1991; Kishi *et al.* 2005; Adams & Zhang 2009; Rall *et al.* 2010; Kordas *et al.* 2011; Kratina *et al.* 2012; Hein *et al.* 2014; Allan *et al.* 2015; Thakur *et al.* 2017). In terrestrial ecosystems, Coley & Aide (1991) proposed that interactions between plants and insect herbivores were generally stronger in warmer regions, which may result in stronger negative density dependence for tree species and contribute to higher plant diversity in a tropical region (Forrister *et al.* 2019). The influence of temperature on interaction strengths has also been investigated in other systems: terrestrial arthropods interactions (Rall *et al.* 2010), fish-fish interactions (Hein *et al.* 2014; Allan *et al.* 2015), and fish-prey interactions (Kishi *et al.* 2005). In addition, Wieczynski *et al.* (2021) showed that species-level functional traits (e.g., body size and shape) may underlie the relationships between temperature and interaction strengths. Therefore, interplays among temperature, interaction strengths, species identity, and community-level function (e.g., primary production) is prevalent, and a more detailed understanding of the complex interactions among them is critical for predicting ecosystem responses to climate change.

In marine ecosystems, interspecific interactions in ecological communities play a
fundamental role in system dynamics, such as population dynamics, primary production, and
nutrient cycling (Hannisdal *et al.* 2017; Ushio *et al.* 2018a; Penn *et al.* 2019), as well as
ecosystem services, including food supply (Smith *et al.* 2019). Although many studies have
evaluated the effects of temperature on interaction strengths of marine organisms (Kordas *et*
al. 2011), most of these studies have been performed targeting relatively immobile or small
organisms (Bertness & Ewanchuk 2002; Chen *et al.* 2012) or under laboratory conditions
(e.g., mesocosm experiments; Allan *et al.* 2015). While the previous studies have provided
invaluable information, it currently remains unclear how temperature influences the strengths
of interspecific interactions of larger, more mobile marine organisms, such as fish, which may
exert strong top-down regulations on ecological communities, especially under field
condition. This may be due to the difficulties associated with detecting and measuring
interaction strengths among multiple, relatively large, mobile species under field conditions;
the identification of quickly-moving fish species and the quantification of their abundance
under field conditions are challenging, and the quantification of their interactions is even
more difficult. Overcoming these difficulties and understanding how temperature influences
the strength of interspecific interactions of marine fish communities will provide insights into
how marine fish communities are assembled, how they may respond to the ongoing and
future global climate change, and how the changes in fish communities may transmit to other
trophic levels.

Environmental DNA (eDNA), defined here as extra-organismal DNA left behind by
macro-organisms (Bohmann *et al.* 2014), has been attracting increasing attention as an
indirect genetic marker for inferring the presence of species for biodiversity monitoring
(Deiner *et al.* 2017; Cristescu & Hebert 2018). A simple protocol for collecting eDNA
samples from aquatic environments facilitates continuous biodiversity monitoring at multiple
sites (Deiner *et al.* 2017), and eDNA metabarcoding (the simultaneous detection of multiple
species using universal primers and a high-throughput sequencer) provides useful
information on the dynamics of ecological communities (Bálint *et al.* 2018; Miya *et al.* 2020;
Miya 2022; Ushio 2022a). Recent studies demonstrated that eDNA metabarcoding combined
with frequent water sampling enabled the efficient monitoring of high-diversity ecological
communities (Bista *et al.* 2017; Djurhuus *et al.* 2020; Ushio 2022a). Furthermore, if eDNA
metabarcoding is performed quantitatively (e.g., by including spike-in DNAs; Ushio *et al.*
2018b), these data may contain information on species abundance.

More importantly, recent studies have shown that information on interspecific

interactions may be embedded in multispecies eDNA time series, particularly when the data

is "quantitative" (Ushio 2022a). Advances in nonlinear time series analyses have enabled the

quantification of interspecific interactions only from time series data. For example, transfer

entropy (TE) is a method based on the information theory that quantifies information flow

between two variables (Schreiber 2000; Runge *et al.* 2012). Information flow may be an

index of interspecific interactions when applied to ecological data. Convergent cross mapping

(CCM) is a causality detection tool in empirical dynamic modeling (Sugihara *et al.* 2012),

which is based on the dynamical theory. TE and CCM have more recently been understood

under the information theory framework, and unified information-theoretic causality (UIC)

may also quantify information flow between variables (rUIC package; Osada & Ushio 2021).

In addition, a recently developed, improved version of a sequentially-weighted global linear

map (S-map), called the MDR S-map, enables accurate quantifications of interaction

strengths of a large interaction network even when the number of network nodes exceeds the

time series length (Chang *et al.* 2021). These advanced statistical tools facilitate the detection

and quantification of interspecific interactions from quantitative, multispecies eDNA

time-series data.

In the present study, we collected eDNA samples twice a month from 11 coastal sites at

the southern tip of the Boso Peninsula, central Japan (Fig. 1a) for two years (a total of 550

samples). This region is located on the Pacific side of Honshu Island around 35°N and is

markedly affected by the warm Kuroshio Current, cold Oyashio Current, and inner-bay water

from Tokyo Bay. These geographic and oceanographic characteristics form latitudinal

temperature gradients, allowing us to investigate temperature effects on interspecific

interactions. We obtained quantitative eDNA metabarcoding data by combining MiFish

eDNA metabarcoding data (relative abundance data) and the eDNA concentration data

obtained through quantitative PCR (qPCR) for one of the most common fish species,

Acanthopagrus schlegelii (e.g., absolute abundance data of *A. schlegelii* eDNA was utilized

as an internal spike-in DNA; see Methods). Based on quantitative, multispecies eDNA

time-series data, pairwise species interactions were detected as information flow between

species using a nonlinear time series analysis (Osada & Ushio 2021). In addition, interaction

strengths at each time point were quantified for the detected fish-fish interactions using the

MDR S-map method (Chang *et al.* 2021). As our eDNA time series was taken twice a month,

the interactions detected should also occur at the same time scale (i.e., about two-week),

which means that we focus on behavior-level interactions rather than birth-death process in

the present study. In addition, the interactions we detected by the time series analysis could

include various types of interactions such as competition and mutualism that could be

involved in the population dynamics. We hypothesized that (1) a positive relationship exists

between temperature and interaction strengths at the community level, as found in other

ecosystems and many types of interactions, and (2) the relationship between the patterns of

temperature and interaction strengths varies among fish species because of species-specific

ecologies and the behavior of component species.

Materials and Methods

Study sites and environmental observations

We selected 11 sites (Stations [Sts.] 1–11) for seawater sampling along the shoreline of

the southern tip of the Boso Peninsula (Fig. 1a). The 11 stations are located within or

surrounded by rocky shores with intricate coastlines. They are arranged such that the five

stations on the Pacific and Tokyo Bay sides are approximately at the same latitudinal

intervals, with St. 6 in between. The northernmost stations on both sides (Sts. 1 and 11) are

markedly affected by the cold Oyashio Current (Yang *et al.* 1993) and inner-bay water from

Tokyo Bay (Fujiwara & Yamada 2002), respectively, while the remaining stations (Sts. 2–10)

are markedly affected by the warm Kuroshio Current flowing northward along the Pacific

coast as well as its branches (Soh 2003). Before seawater sampling at each station, we

measured seawater temperature (˚C) and salinity (‰) using a portable water-quality meter

(WQC-30, Toa DKK, Tokyo, Japan). In addition, we recorded the sampling start time,

latitude/longitude, weather and sea conditions, and turbidity.

Collection of eDNA samples

Detailed information on seawater sampling and on-site filtration is provided in

Supporting Information. We collected seawater samples twice a month (i.e., around the first

and second quarter moons) for two years between August 2017 and August 2019 by casting a

bucket fastened to a rope at the 11 sites. We filtered the collected seawater using a

combination of a filter cartridge and disposable syringe until the final filtration volume

reached 1,000 mL for each replicate and obtained duplicate samples at each site. We added

RNAlater (Thermo Fisher Scientific, DE, USA) into the cartridge to prevent eDNA

degradation. We made a filtration blank (FB) by filtering 500 mL of purified water in the

same manner at the end of each day of water sampling.

Laboratory protocol

Detailed information on the laboratory protocol is provided in Supporting Information.

We thoroughly sterilized the workspace and equipment in the DNA extraction, pre-PCR, and

post-PCR rooms before all experiments. We used filtered pipette tips and conducted all

eDNA extraction, pre-PCR, and post-PCR manipulations in three different dedicated rooms

that were separate from each other to safeguard against cross-contamination.

We extracted eDNA from filter cartridges using the DNeasy Blood & Tissue kit

(Qiagen, Hilden, Germany) following the methods developed by Miya *et al.* (2016) with

slight modifications (Miya & Sado 2019a; Minamoto *et al.* 2021). After removing RNAlater

from filter cartridges, they were subjected to lysis using proteinase K. We purified the

collected DNA extracts using the above kit following the manufacturer's protocol and the

final elution volume was set to 200 µL. An extraction blank (EB) was also created during this

process.

We employed two-step PCR for paired-end library preparation using the MiSeq

platform (Illumina, CA, USA). We generally followed the methods developed by Miya et al.

(2015) and subsequently modified by Miya & Sado (2019b). Detailed information on library

preparation is provided in Supporting Information.

We performed the first-round PCR (1st PCR) using a mixture of the following six

primers: MiFish-U-forward, MiFish-U-reverse, MiFish-E-forward-v2, MiFish-E-reverse-v2,

MiFish-U2-forward, and MiFish-U2-reverse (Table 1). The 1st PCR amplified a

hypervariable region of the mitochondrial 12S rRNA gene (*ca*. 172 bp; hereafter called the

"MiFish sequence"). We also prepared a 1st PCR blank (1B) during this process, in addition

to FB and EB. After the 1st PCR, PCR products were purified, quantified, diluted to 0.1

ng/µL, and used as templates for the second-round PCR (2nd PCR).

We performed the 2nd PCR to append dual-indexed sequences and flow cell-binding

sites for the MiSeq platform. We prepared a 2nd PCR blank (2B) during this process in

addition to FB, EB, and 1B. In total, we made 195 blanks (FB = 60, EB = 50, 1B = 50, 2B =

35) to monitor contamination during the on-site filtration, subsequent DNA extraction, and

1st and 2nd PCR of 550 samples.

We pooled each of the individual paired-end libraries in an equal volume, performed

electrophoresis on an agarose gel (Invitrogen, CA, USA), and excised the target amplicons

(*ca*. 370 bp). The concentrations of the size-selected libraries were measured, diluted to

10–12.0 pM, and sequenced on the MiSeq platform using the MiSeq v2 Reagent Kit for $2 \times$

150 bp PE (Illumina, CA, USA) following the manufacturer's protocol.

Sequence analysis

We performed data preprocessing and analyses of raw MiSeq reads from the MiSeq run

using PMiFish ver. 2.4 (https://github.com/rogotoh/PMiFish; Miya et al. 2020). Detailed

information on sequence data processing is provided in Supporting Information. Forward

(R1) and reverse (R2) reads were merged while discarding short reads (<100 bp) after tail

trimming and paired reads with too many differences (>5 positions) in the aligned region (*ca*.

65 bp). Primer sequences were removed from merged reads and reads without the primer

sequences underwent quality filtering to remove low quality reads. Preprocessed reads were

dereplicated, and all singletons, doubletons, and tripletons were removed from subsequent

analyses to avoid false positives (Edgar 2010). Dereplicated reads were denoised to generate

amplicon sequence variants (ASVs) that removed all putatively chimeric and erroneous

sequences (Edgar 2016; Callahan *et al.* 2017).

ASVs were subjected to taxon assignments to species names with a sequence identity

of >98.5% with the reference sequences (two nucleotide differences allowed) and a query

coverage of ≥90%. ASVs with identical fish species names were clustered as molecular

operational taxonomic units (MOTUs) following the procedure described in Supporting

Information. MiFish DB ver. 43 was used for taxon assignment, comprising 7,973 species

distributed across 464 families and 2,675 genera. The ASV table was rarefied to the

approximate minimum read number (*ca*. 20,000 reads). To refine the above taxon

assignments, family-level phylogenies were reproduced from MiFish sequences from

MOTUs and reference sequences (contained in MiFish DB ver. 43) belonging to these

families. A total of 103 family-level trees were visually inspected and taxon assignments

were revised. The final list of detected species is provided in Table S1.

Estimation of DNA copy numbers

The nonlinear time series analytical tools used in the present study require quantitative

time-series data. To estimate fish eDNA concentrations, we initially measured the eDNA

concentrations of the most common fish species in the region, Japanese Black Seabream (*A*.

schlegelii), using qPCR, which was detected in 504 out of 550 seawater samples (91.6%). By

dividing *A. schlegelii* sequence reads by the *A. schlegelii* eDNA quantity in each sample, we

estimated the number of sequence reads generated per *A. schlegelii* eDNA copy for each

sample, and we applied the same conversion factor (i.e., sequence reads /*A. schlegelii* eDNA

copy) to other fish species. Note that sequence reads generated per eDNA copy may vary

depending on factors such as the level of PCR inhibition and PCR amplification efficiency;

however, in general, this "internal spike-in DNA" method has been shown to reasonably

estimate the quantity of eDNA concentrations (Ushio *et al*. 2018b; Ushio 2022a). More

importantly, species-specific biases that may be introduced in the internal spike-in DNA

method do not cause serious biases in the outcomes of our nonlinear time-series analysis

because it standardizes time series to have a zero mean and a unit variance before analyses

and also only utilizes the fluctuation patterns of time series. When we did not detect any *A.*

schlegelii eDNA by qPCR, we replaced the "zero" value with the minimum eDNA copy

numbers of *A. schlegelii* (0.346 copies/µl DNA). Similarly, when we did not detect any *A.*

schlegelii eDNA sequence reads by metabarcoding, we replaced the "zero" value with the

minimum eDNA sequence reads of *A. schlegelii* (12 reads/sample). These corrections

estimated how many sequence reads were generated per eDNA copy for all samples. Details

are provided in Supporting Information.

Nonlinear time series analysis to detect fish-fish interactions

We detected fish-fish interactions using a nonlinear time series analysis based on

quantitative eDNA time-series data from multiple species and sites. Since the reliable

detection of interspecific interactions requires sufficient information in the target time series,

we arbitrarily selected the top 50 most frequently detected species and excluded other rarer

fish species from the analysis. We quantified information flow between two fish species'

eDNA time series by the "unified information-theoretic causality (UIC)" method

implemented in the "rUIC" v0.1.5 package (Osada & Ushio 2021) of R (R Core Team 2022).

UIC tests the statistical clarity of information flow between variables in terms of TE

(Schreiber 2000) computed by nearest neighbor regression based on time-delay embedding of

explanatory variables (i.e., cross mapping; Sugihara *et al*. 2012). In contrast to the standard

method used to measure TE, UIC quantifies information flow as follows:

$$TE = \frac{1}{T} \sum_{t=1}^{T} \log \left(\frac{p\left(y_{t+tp} \mid x_t, x_{t-\tau}, \cdots, x_{t-(E-1)\tau}, z_t\right)}{p\left(y_{t+tp} \mid x_{t-\tau}, x_{t-2\tau}, \cdots, x_{t-(E-1)\tau}, z_t\right)} \right), \# \dots [1]$$

where x, y, and z represent a potential causal variable, effect variable, conditional variable (if

available), respectively. $p(A|B)$ represents conditional probability: the probability of A

conditioned on B. t, tp, τ, and E represent the time index, time step, a unit of time lag, and the

optimal embedding dimension, respectively. T is the total number of points in the

reconstructed state space (this is equivalent to the total number of time points minus (E–1)).

For example, if TE between y_{t+1} and $\{x_t, x_{t-\tau}, \ldots, x_{t-(E-1)\tau}\}$ is measured (tp is set to 1), it tests

the causal relationship between y_{t+1} and x_t. Optimal E was selected by measuring TE as

follows:

$$TE = \frac{1}{T}\sum_{t=1}^{T} \log\left(\frac{p\left(x_{t+tp} \mid y_t, x_t, x_{t-\tau}, \cdots, x_{t-(E-1)\tau}\right)}{p\left(x_{t+tp} \mid y_t, x_t, x_{t-\tau}, \cdots, x_{t-(E_R-1)\tau}\right)}\right), \# \ldots [2]$$

where E_R ($< E$) is the optimal embedding dimension of lower dimensional models.

Significance tests were conducted by bootstrapping data after embedding (the significance

levels were set to 0.05). Eqn. [2] is a TE version of simplex projection (Sugihara & May

1990). TE measured according to Eqn. [1] gains the advantage of previous causality tests, *i.e.*,

standard TE methods (Schreiber 2000; Runge *et al.* 2012) and convergent cross mapping

(CCM) (Sugihara *et al.* 2012), the algorithm of which is explained and implemented in the

rUIC website (https://github.com/yutakaos/rUIC).

By using UIC, we quantified TE between fish eDNA time series. As environmental

variables, water temperature (℃), salinity (‰), wave height (m), and tide level (cm) were

considered as conditional variables. This means that the effects of the environmental

variables on the fish eDNA abundance were removed in the analysis when quantifying

interspecific interaction strengths. Importantly, in most cases, water temperature had

statistically clear influence on fish eDNA dynamics and included as a conditional variable in

the embedding. Because water temperature showed clear seasonality in the region (Fig. S1),

the effect of the seasonality in detecting causation was taken into account in this analysis. We

merged all eDNA time series across the 11 study sites and standardized the eDNA time series

to have zero means and a unit of variance before the analysis. If TE between two fish eDNA

time series was statistically clearly higher than zero, we interpreted it as a sign of an

interspecific interaction. After the detections of interspecific interactions among the top 50

fish species, we visualized the interaction network (i.e., the reconstruction of fish-fish

interaction network). Importantly, UIC quantifies the average information flow between two

time series (see Eqn. [1]), and thus there is only one TE value for each pair of fish species,

which is critically different from interaction strengths quantified by the MDR S-map (see the

following section).

Nonlinear time series analysis to quantify interaction strengths

We quantified fish-fish interaction strengths for the interspecific interactions detected

by UIC. For this purpose, we used an improved version of the sequential locally-weighted

global linear map (S-map) (Sugihara 1994), called the multiview-distance, regularized S-map

(MDR S-map) (Chang *et al*. 2021). Consider a system that has E different interacting

variables, and assume that the state space at time t is given by $x(t) = \{x_1(t), x_2(t), \ldots, x_E(t)\}$.

For each target time point t^*, the S-map method produces a local linear model that predicts

the future value $x_1(t^*+p)$ from the multivariate reconstructed state space vector $x(t^*)$. That is,

$$\hat{x}_1(t^* + p) = IS_0 + \sum_{j=1}^{E} \boldsymbol{IS}_j \, x_j(t^*) \# \ldots [3]$$

328
where $\hat{x}_1(t^* + p)$ is a predicted value of x_1 at time t^*+p, IS_j is a regression coefficient and

interpreted as interaction strength (or called S-map coefficient), and IS_0 is an intercept of the

linear model. The linear model is fit to the other vectors in the state space. However, points

that are close to the target point, $x(t^*)$, are given greater weighting (i.e., locally weighted

linear regression). In the standard S-map, the distances between the target point, $x(t^*)$, and

other points are measured by Euclidean distance. However, Euclidean distance cannot be a

good measure if the dimension of the state space is high, and in this case, it is impossible to

identify nearest neighbors correctly if we rely on Euclidean distance. In the MDR S-map, the

distance between the target point and other points are measured by the multiview distance

(Chang *et al*. 2021), which can be calculated by ensembling distances measured in various

low-dimensional state spaces (multiview embedding; see Ye & Sugihara 2016). In addition,

to reduce the possibility of overestimation and to improve forecasting skill, regularization

(i.e., ridge regression) is also applied (Cenci *et al*. 2019). Chang et al. (2021) showed that the

MDR S-map outperformed other S-map methods and it enables improved estimations of

interaction strengths. Importantly, the MDR S-map quantifies interaction strength at each

time point (i.e., the maximum possible number of interaction strength values is the number of

fish-fish interactions $\times$ the number of time points – the optimal embedding dimension + 1).

In the present study, we implemented the MDR S-map in our custom R package, "macam"

v0.0.10 (Ushio 2022b) and used it for computation.

Quantification of the temperature sensitivity of fish species interactions

After reconstructing fish interaction networks and quantifying interaction strengths, we

analyzed the relationships among fish species interaction strengths and biotic and abiotic

variables. We performed all analyses using R v4.2.1 (R Core Team 2022). Generalized

additive mixed models (GAMMs) were performed using the "mgcv" package of R (Wood

2004). We visualized results with "ggplot2" (Wickham 2009) and "cowplot" (Wilke 2017). In

the present study, we used the term "statistical clarity" instead of "statistical significance" to

avoid misinterpretations, according to the recommendations by Dushoff et al. (2019).

In the first analysis, the relationships between fish-fish interaction strengths and the

characteristics of the study sites were analyzed by GAMM. In the analysis, in-strength

(interaction strengths that a species receives) and out-strength (interaction strengths that a

species gives) were separately calculated. In GAMM, in-strength or out-strength were an

explained variable, and an environmental or ecological variable was an explaining variable.

Explaining variables included water temperature, species richness, total fish eDNA

concentrations, salinity, tide level, and wave height. A gamma distribution was assumed as

the error distribution, the "log" function was used as a link function, and study site and fish

species were used as a random effect (i.e., in R, `gamm(abs(IS) ~ s(explaining_variable)`,

`family = Gamma(link="log"), random = list(site = ~1, fish_species = ~1))`, where

`s()` indicates a smoothing term). The biological assumption behind this modeling is that

explaining variables, such as water temperature, may linearly or nonlinearly influence fish

species interactions. Moreover, the effects of explaining variables may randomly vary

depending on the study site and fish species. The effect was considered to be statistically

clear at $P < 0.05$.

In the second analysis, we focused on the effects of water temperature on the

interaction strengths of each fish. In the analysis, GAMM was used again. A gamma

distribution was assumed as the error distribution and the "log" function was used as a link

function, and the study site was used as a random effect (i.e., in R, `gamm(abs(IS) ~`

`s(explaining_variable), family = Gamma(link="log"), random = list(site = ~1))`,

where `s()` indicates a smoothing term). GAMM was separately performed for each fish

species. We also analyzed the effects of other properties on the interaction strengths of each

fish species, which are provided in the Supporting Information. Again, the effect was

considered to be statistically clear at $P < 0.05$.

Code and data availability

384   Sequence data were deposited in DDBJ Sequence Read Archives (DRA) (DRA accession

385   number = DRA014111). Analysis scripts are available at Github

386   (https://github.com/ong8181/eDNA-BosoFish-network).

Results and Discussion

Taxonomic diversity and dynamics of fish eDNA

Our multiple MiSeq runs generated 71,892,685 reads in total and detected 1,130 molecular operational taxonomic units (MOTUs). We inspected their family-level phylogenies to increase the accuracy of taxonomic assignments, recognizing 856 MOTUs across 33 orders, 167 families, and 466 genera (Table S1). This taxonomic diversity was similar to that of the local fauna (948 species across 33 orders, 158 families, and 493 genera) compiled from a literature survey and museum collections (Table S2).

Sequence reads in the ASV table were converted to estimated eDNA concentrations by using the eDNA concentrations of a common fish species detected across most samples (Japanese Black Seabream, *A. schlegelii*) (i.e., an analog of the internal spike-in DNA method; see Methods and Fig. S1). The converted ASV table included 23,863 detections and we selected the top 50 most frequently detected species for our time-series analysis. The detection of these 50 species ranged between 148 and 532 with a mean of 296, and their total detection reached 14,793 (62.0%).

Fish eDNA concentrations and fish species richness showed a clear intra-annual pattern (i.e., seasonality; Fig. 1b, c), which were higher in warmer months (e.g., between July and October) than in colder months (e.g., between January and March) at all sites. Total eDNA concentrations and species richness positively correlated with water temperature (Fig. S2). Seasonal changes in eDNA concentrations were consistent with the patterns of seasonal occurrences in tropical and subtropical fish species in the Boso Peninsula, to which they are transported by the warm Kuroshio Current, settling on the coastal waters during the warmer months and subsequently disappearing during the colder months (Senou *et al.* 2006; Saito 2019).

Reconstruction of the fish species interaction network

We reconstructed fish species interaction networks based on the quantitative, multispecies fish eDNA time series. Regarding the 50 frequently detected fish species, we quantified pairwise information flow between fish species. Figure 2 shows reconstructed fish interaction networks for the 50 frequently detected fish species in the Boso Peninsula. At the regional scale, the linear correlations among water temperature, total eDNA concentrations, interaction strengths, the number of interactions, and species richness was all statistically clear ($P < 0.05$; Fig. S3) except for the linear correlation between water temperature and

mean interaction strengths ($P > 0.05$). The interaction strengths became weak as species

richness increased, which is consistent with a previous study (Ratzke *et al.* 2020; Ushio

2022a), suggesting that weak interactions are key to the maintenance of species-rich

communities.

Most of the statistically clear information flow may be interpreted from the viewpoint

of fish-fish behavior-level interspecific interactions (Table S3 and Supporting Discussion),

which is convincing considering the time resolution of our eDNA time series (but see the

section "*Potential limitations of the present study*" below). The largest information flow was

detected from *Pseudoblennius marmoratus* to *Pseudolabrus eoethinus* (Table S3). These fish

species have overlapping habitats, and thus, they may interact. Bidirectional information flow

was detected between *P. eoethinus* and *Gymnothorax kidako*. They are both carnivorous fish

species and may conduct joint hunting. *Chaenogobius annularis* and *Takifugu niphobles* are

often found together in the sand between reefs. *T. niphobles* frequently dive in the sand, and

benthos and mysids that are dug up during the dive may be prey for *C. annularis*. Further

discussions about the detected information flow are described in Supporting Information

(Table S3 and Supplementary Discussion).

Interaction strengths and environmental variables

We investigated how interaction strengths (i.e., regression coefficients estimated by the

MDR S-map) changed with environmental variables (e.g., water temperature) and ecological

properties (e.g., species richness and total eDNA concentrations) (Figs. 3 and S4). The

in-strengths and out-strengths of fish species interactions were statistically clearly associated

with water temperatures, species richness, and total eDNA concentrations (GAMM, $P < 0.05$;

Fig. 3 and Table S4) except for the effects on species richness on the in-strengths (Fig. 3b).

In-strengths of the fish-fish interactions increased with water temperature (Fig. 3a),

supporting our first hypothesis while out-strengths of the interactions showed an opposite

pattern (Fig. 3d), which might suggest there is a difference in the temperature dependence

between the in-strengths and out-strengths of the interactions. Indeed, water temperature may

influence fish physiological activity (Kishi *et al.* 2005; Claireaux *et al.* 2006; Oyugi *et al.*

2012) often in a complex way, and thus, the community-level influence of water temperature

may also be complex as they should arise from the individual-level influence of water

temperatures on fish. Interaction strengths were also statistically clearly influenced by species

richness and total eDNA concentrations (except for Fig. 3b); the interaction strengths

decreased with increasing water temperature. The effects of salinity, tide and wave were less

clear, although the effects were statistically clear except for the effects of tide level on the

out-strength (Figs. S4 and S6). Overall, environmental and ecological variables influenced

the interaction strengths statistically clearly, but large variations remained unexplained (Figs.

S5–S6), suggesting that other factors (e.g., fish ecology, nutrient, and physiological status)

may influence the interaction strengths.

We also investigated how temperature influenced the interaction strength of individual

fish species. Figure 4 shows how interaction strengths among fish species changed with water

temperature. For visualization purpose, we show fish species of which interaction strengths

changed with water temperature highly statistically clearly ($P < 0.0001$). The in-strengths of

several fish species clearly increased at higher water temperatures: for example, *Halichoeres*

tenuispinis, *Macrocanthus strigatus*, *Pempheris schwenkii*, *Stethojulis interrupta terina*, and

Thalassoma cupido (Fig. 4a). On the other hand, in-strengths of some fish species and

out-strengths decreased at higher water temperatures: for example, in-strengths of *Ditrema*

temminckii temminckii, *Engraulis japonicus*, and *Girella punctata*, and out-strengths of five

fish species (Fig. 4a,b). These results support our second hypothesis that the relationship

between the patterns of temperature and interaction strengths varies among fish species.

These results suggest that, although water temperature may have strong influences on

fish-fish interaction strengths in general, the sign of temperature effect (i.e., positive or

negative) can vary depending on fish species and environmental conditions. Previous studies

showed that fish physiological activities, such as feeding rates, growth rates, and swimming

speed, were influenced by water temperature (Kishi *et al*. 2005; Claireaux *et al*. 2006; Oyugi

et al. 2012). In addition, the direction (i.e., positive or negative) of the temperature effects on

fish metabolic activities may be species-specific (Oyugi *et al*. 2012), and this species

specificity may underlie the species-specific effects of temperature on the interaction strength.

Also, importantly, other environmental conditions may influence fish-fish interactions (Figs.

S7–S8), suggesting that the interaction strengths may fluctuate through time and space as

shown in previous studies (Ushio *et al*. 2018a; Ushio 2022a). Conversely, interactions

detected and quantified under a controlled environment might not necessarily be observed

under field conditions, and thus, our research framework that enables the detection and

quantification of interactions under field conditions would play a critical role in

understanding the effects of temperature on fish-fish interactions under field conditions.

Potential limitations of the present study

The present results showing that the strengths of fish-fish interactions depend on water

temperature rely on several assumptions that were not fully investigated, and thus, careful

interpretations and discussions are required. First, the extent to which fish eDNA

concentrations accurately represent fish abundance is an open question. Previous studies

demonstrated that the quantity of eDNA may be a proxy of fish abundance and/or biomass

(i.e., a positive correlation between the quantity of eDNA and fish abundance/biomass)

(Takahara *et al.* 2012; Thomsen *et al.* 2012; Yamamoto *et al.* 2016). Our nonlinear time series

analysis used the information embedded in fluctuations in time series, and time series are

always standardized before the analysis. Therefore, species-specific differences in the copy

numbers of the genetic marker and eDNA release rates, which may bias the estimation of

absolute abundance, do not cause serious biases in the reconstruction of the networks and

estimations of interaction strengths. However, accurate estimations of the absolute abundance

of fish will improve the accuracy of analyses, and the integration of eDNA concentration data,

knowledge on eDNA dynamics (i.e., release and degradation rates), and the hydrodynamic

modeling of ocean water flow will be a promising approach (e.g., as in Fukaya *et al.* 2021).

Another limitation is that the reconstructed interaction network was not fully validated

in the present study. Although previous studies demonstrated that network reconstructions

based on a time series analysis work reasonably well and show meaningful patterns (Sugihara

et al. 2012; Ushio *et al.* 2018a; Runge *et al.* 2019; Ushio 2022a),

experiment/observation-based validations of the reconstructed network have not yet been

performed. Potential causal interactions between fish species may be reasonably interpreted

(Supplementary Discussion); however, developing a framework to efficiently validate

"previously unknown" interactions detected by eDNA data and a nonlinear time series

analysis is an important future direction.

Furthermore, although we showed that temperature effects on interaction strength are

relatively clear and common in the Boso peninsula coastal region, the range of water

temperatures in the present analysis was still narrow and there is currently no evidence that

we can generalize these results to other regions. Nonetheless, the advantages of the eDNA

method are the low cost, time, and labor needed for field sampling as well as the potential

scalability of the library preparation process (Buchner *et al.* 2021; Ushio *et al.* 2022).

Therefore, frequent (Ushio 2022a) and large spatial-scale ecological monitoring (e.g.,

ANEMONE DB; https://db.anemone.bio/) is possible, which will provide a more detailed

understanding of the temperature-interaction strength relationships of fish on a larger spatial

scale.

Conclusions

The present study demonstrated that the strengths of fish species interactions changed

with water temperature under field conditions. For several fish species, species interactions

were intensified in warmer water and for some other fish species, species interactions were

weakened in warmer water. This may change the correlation and distribution of interaction

strengths in a community, which may consequently influence community dynamics and

stability (Tang *et al*. 2014; Allesina *et al*. 2015; Ushio *et al*. 2018a). Therefore, a more

detailed understanding of the effects of environmental conditions on species interactions

under field conditions will be required to improve our capability to understand, forecast, and

even manage natural ecological communities and dynamics, which is particularly important

under ongoing global climate change. Developments and improvements in eDNA techniques

and statistical analyses are still required; however, because eDNA analysis is potentially

applicable to any type of organisms even if they are difficult to be detected using traditional

methods, our framework integrating an eDNA analysis and advanced statistical analyses pave

a way to understand and forecast dynamics of various ecological communities under field

conditions.

Author contributions

All authors agreed to the submission. MU and MM conceived the idea and designed the

research; MM, TS, and TF collected the samples, performed eDNA metabarcoding and

analyzed sequence data; SS and RM performed quantitative PCR analysis; MU compiled

data; MU and YO analyzed data; MU and MM wrote the first draft and completed the final

manuscript with contributions from all coauthors.

Acknowledgments

We thank T. Komai, R.O. Gotoh, T. Sunobe, and K. Takiguchi for assisting with the

bimonthly collection of eDNA samples. This research was supported by JSPS KAKENHI (B)

Grant Numbers JP19H03291, JP22H02691, and MEXT OGAP Project Grant Number

JPMXD0618068274 to MM, JSPS KAKENHI (B) Grant Number 20H03323, and the Hakubi

Project in Kyoto University to MU.

References:

Adams, J.M. & Zhang, Y. (2009). Is there more insect folivory in warmer temperate climates? A latitudinal comparison of insect folivory in eastern North America. *J. Ecol.*, 97, 933–940.

Allan, B.J.M., Domenici, P., Munday, P.L. & McCormick, M.I. (2015). Feeling the heat: the effect of acute temperature changes on predator–prey interactions in coral reef fish. *Conserv. Physiol.*, 3, cov011.

Allesina, S., Grilli, J., Barabás, G., Tang, S., Aljadeff, J. & Maritan, A. (2015). Predicting the stability of large structured food webs. *Nat. Commun.*, 6, 7842.

Bálint, M., Pfenninger, M., Grossart, H.-P., Taberlet, P., Vellend, M., Leibold, M.A., *et al.* (2018). Environmental DNA Time Series in Ecology. *Trends Ecol. Evol.*, 33, 945–957.

Bertness, M.D. & Ewanchuk, P.J. (2002). Latitudinal and climate-driven variation in the strength and nature of biological interactions in New England salt marshes. *Oecologia*, 132, 392–401.

Bista, I., Carvalho, G.R., Walsh, K., Seymour, M., Hajibabaei, M., Lallias, D., *et al.* (2017). Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics of lake ecosystem biodiversity. *Nat. Commun.*, 8, 14087.

Bohmann, K., Evans, A., Gilbert, M.T.P., Carvalho, G.R., Creer, S., Knapp, M., *et al.* (2014). Environmental DNA for wildlife biology and biodiversity monitoring. *Trends Ecol. Evol.*, 29, 358–367.

Buchner, D., Macher, T.-H., Beermann, A.J., Werner, M.-T. & Leese, F. (2021). Standardized high-throughput biomonitoring using DNA metabarcoding: Strategies for the adoption of automated liquid handlers. *Environ. Sci. Ecotechnology*, 8, 100122.

Callahan, B.J., McMurdie, P.J. & Holmes, S.P. (2017). Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *Isme J.*

Cenci, S., Sugihara, G. & Saavedra, S. (2019). Regularized S-map for inference and forecasting with noisy ecological time series. *Methods Ecol. Evol.*, 10, 650–660.

Chang, C.-W., Miki, T., Ushio, M., Lu, H.-P., Shiah, F.-K. & Hsieh, C. (2021). Reconstructing large networks with time-varying interactions. *ArXiv210204069 Q-Bio*.

Chen, B., Landry, M.R., Huang, B. & Liu, H. (2012). Does warming enhance the effect of microzooplankton grazing on marine phytoplankton in the ocean? *Limnol. Oceanogr.*, 57, 519–526.

Claireaux, G., Couturier, C. & Groison, A.-L. (2006). Effect of temperature on maximum swimming speed and cost of transport in juvenile European sea bass (Dicentrarchus labrax). *J. Exp. Biol.*, 209, 3420–3428.

Coley, P.D. & Aide, T.M. (1991). *A comparison of herbivory and plant defenses in temperate and tropical broad-leaved forests*. Wiley & Sons, New York.

Cristescu, M.E. & Hebert, P.D.N. (2018). Uses and Misuses of Environmental DNA in Biodiversity Science and Conservation. *Annu. Rev. Ecol. Evol. Syst.*, 49, 209–230.

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., *et al.*

597   (2017). Environmental DNA metabarcoding: Transforming how we survey animal
598     and plant communities. *Mol. Ecol.*, 26, 5872–5895.

599 Djurhuus, A., Closek, C.J., Kelly, R.P., Pitz, K.J., Michisaki, R.P., Starks, H.A., *et al.* (2020).
600     Environmental DNA reveals seasonal shifts and potential interactions in a marine
601     community. *Nat. Commun.*, 11, 254.

602 Dushoff, J., Kain, M.P. & Bolker, B.M. (2019). I can see clearly now: Reinterpreting
603     statistical significance. *Methods Ecol. Evol.*, 10, 756–759.

604 Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST.
605     *Bioinformatics*, 26, 2460–2461.

606 Edgar, R.C. (2016). UNOISE2: improved error-correction for Illumina 16S and ITS amplicon
607     sequencing. *bioRxiv*.

608 Forrister, D.L., Endara, M.-J., Younkin, G.C., Coley, P.D. & Kursar, T.A. (2019). Herbivores
609     as drivers of negative density dependence in tropical forest saplings. *Science*, 363,
610     1213 LP – 1216.

611 Fujiwara, T. & Yamada, Y. (2002). Inflow of oceanic water into Tokyo Bay and generation of
612     a subsurface hypoxic water mass. *J. Geophys. Res. Oceans*, 107, 13-1-13–9.

613 Fukaya, K., Murakami, H., Yoon, S., Minami, K., Osada, Y., Yamamoto, S., *et al.* (2021).
614     Estimating fish population abundance by integrating quantitative data on
615     environmental DNA and hydrodynamic modelling. *Mol. Ecol.*, 30, 3057–3067.

616 Hannisdal, B., Haaga, K.A., Reitan, T., Diego, D. & Liow, L.H. (2017). Common species link
617     global ecosystems to climate change: dynamical evidence in the planktonic fossil
618     record. *Proc. R. Soc. B Biol. Sci.*, 284.

619 Hein, C.L., Öhlund, G. & Englund, G. (2014). Fish introductions reveal the temperature
620     dependence of species interactions. *Proc. R. Soc. B Biol. Sci.*, 281, 20132641.

621 Kishi, D., Murakami, M., Nakano, S. & Maekawa, K. (2005). Water temperature determines
622     strength of top-down control in a stream food web. *Freshw. Biol.*, 50, 1315–1322.

623 Kordas, R.L., Harley, C.D.G. & O'Connor, M.I. (2011). Community ecology in a warming
624     world: The influence of temperature on interspecific interactions in marine systems. *J.
625     Exp. Mar. Biol. Ecol.*, Global change in marine ecosystems, 400, 218–226.

626 Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A. & Shurin, J.B. (2012).
627     Warming modifies trophic cascades and eutrophication in experimental freshwater
628     communities. *Ecology*, 93, 1421–1430.

629 May, R.M. (1972). Will a Large Complex System be Stable? *Nature*, 238, 413–414.

630 Minamoto, T., Miya, M., Sado, T., Seino, S., Doi, H., Kondoh, M., *et al.* (2021). An
631     illustrated manual for environmental DNA research: Water sampling guidelines and
632     experimental protocols. *Environ. DNA*, 3, 8–13.

633 Miya, M. (2022). Environmental DNA Metabarcoding: A Novel Method for Biodiversity
634     Monitoring of Marine Fish Communities. *Annu. Rev. Mar. Sci.*, 14, 161–185.

635 Miya, M., Gotoh, R.O. & Sado, T. (2020). MiFish metabarcoding: a high-throughput
636     approach for simultaneous detection of multiple fish species from environmental
637     DNA and other samples. *Fish. Sci.*, 86, 939–970.

638 Miya, M., Minamoto, T., Yamanaka, H., Oka, S., Sato, K., Yamamoto, S., *et al.* (2016). Use

of a filter cartridge for filtration of water samples and extraction of environmental
DNA. *J. Vis. Exp.*, e54741–e54741.

Miya, M. & Sado, T. (2019a). DNA extraction from a filter cartridge. In: *Environmental DNA*
sampling and experimental manual version 2.1. eDNA Methods Standardization
Committee, The eDNA Society, Otsu, Japan, pp. 31–42.

Miya, M. & Sado, T. (2019b). Multiple species detection using MiFish primers. In:
Environmental DNA sampling and experimental manual version 2.1. eDNA Methods
Standardization Committee, The eDNA Society, Otsu, Japan, pp. 55–92.

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y., Sato, K., *et al.* (2015). MiFish, a set
of universal PCR primers for metabarcoding environmental DNA from fishes:
detection of more than 230 subtropical marine species. *R. Soc. Open Sci.*, 2, 150088.

Mougi, A. & Kondoh, M. (2012). Diversity of interaction types and ecological community
stability. *Science*, 337, 349–351.

Osada, Y. & Ushio, M. (2021). R package for Unified Information-theoretic Causality.

Oyugi, D.O., Cucherousset, J. & Robert Britton, J. (2012). Temperature-dependent feeding
interactions between two invasive fishes competing through interference and
exploitation. *Rev. Fish Biol. Fish.*, 22, 499–508.

Penn, J.L., Weber, T., Chang, B.X. & Deutsch, C. (2019). Microbial ecosystem dynamics
drive fluctuating nitrogen loss in marine anoxic zones. *Proc. Natl. Acad. Sci.*, 116,
7220 LP – 7225.

R Core Team. (2022). R: A language and environment for statistical computing.

Rall, B.C., Vucic-Pestic, O., Ehnes, R.B., Emmerson, M. & Brose, U. (2010). Temperature,
predator–prey interaction strength and population stability. *Glob. Change Biol.*, 16,
2145–2157.

Ratzke, C., Barrere, J. & Gore, J. (2020). Strength of species interactions determines
biodiversity and stability in microbial communities. *Nat. Ecol. Evol.*

Runge, J., Bathiany, S., Bollt, E., Camps-Valls, G., Coumou, D., Deyle, E., *et al.* (2019).
Inferring causation from time series in Earth system sciences. *Nat. Commun.*, 10,
1–13.

Runge, J., Heitzig, J., Petoukhov, V. & Kurths, J. (2012). Escaping the Curse of
Dimensionality in Estimating Multivariate Transfer Entropy. *Phys. Rev. Lett.*, 108,
258701.

Saito, H. (2019). The Kuroshio: its recognition, scientific activities and emerging issues.
Kuroshio current: Physical, biogeochemical, and ecosystem dynamics. In: *Kuroshio*
Current: Physical, Biogeochemical, and Ecosystem Dynamics, Geophysical
monograph series (eds. Nagai, T., Saito, H., Suzuki, K. & Takahashi, M.). John Wiley
& Sons, Inc., Hoboken, NJ.

Schreiber, T. (2000). Measuring Information Transfer. *Phys. Rev. Lett.*, 85, 461–464.

Senou, H., Matsuura, K. & Shinohara, G. (2006). Checklist of Fishes in the Sagami Sea with
Zoogeographical Comments on Shallow Water Fishes Occurring along the Coastlines
under the Influence of the Kuroshio Current. 国立科学博物館専報, 389–542.

Smith, P., Adams, J., Beerling, D.J., Beringer, T., Calvin, K.V., Fuss, S., *et al.* (2019).

681   Land-management options for greenhouse gas removal and their impacts on
682   ecosystem services and the sustainable development goals. *Annu. Rev. Environ.*
683   *Resour.*, 44, 255–286.

684   Soh, W. (2003). Transport processes deduced from geochemistry and the void ratio of surface
685   core samples, deep sea Sagami Bay, central Japan. *Prog. Oceanogr.*, Long-Term
686   Monitoring of Sedimentary Processes in Central Sagami Bay Japan, 57, 109–124.

687   Sugihara, G. (1994). Nonlinear Forecasting for the Classification of Natural Time Series.
688   *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 348, 477–495.

689   Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M., *et al.* (2012). Detecting
690   causality in complex ecosystems. *Science*, 338, 496–500.

691   Sugihara, G. & May, R.M. (1990). Nonlinear forecasting as a way of distinguishing chaos
692   from measurement error in time series. *Nature*, 344, 734–41.

693   Takahara, T., Minamoto, T., Yamanaka, H., Doi, H. & Kawabata, Z. (2012). Estimation of
694   fish biomass using environmental DNA. *PLOS ONE*, 7, e35868.

695   Tang, S., Pawar, S. & Allesina, S. (2014). Correlation between interaction strengths drives
696   stability in large ecological networks. *Ecol. Lett.*, 17, 1094–100.

697   Thakur, M.P., Künne, T., Griffin, J.N. & Eisenhauer, N. (2017). Warming magnifies predation
698   and reduces prey coexistence in a model litter arthropod system. *Proc. R. Soc. B Biol.*
699   *Sci.*, 284, 20162570.

700   Thomsen, P.F., Kielgast, J.O.S., Iversen, L.L., Wiuf, C., Rasmussen, M., Gilbert, M.T.P., *et al.*
701   (2012). Monitoring endangered freshwater biodiversity using environmental DNA.
702   *Mol. Ecol.*, 21, 2565–2573.

703   Ushio, M. (2022a). Interaction capacity as a potential driver of community diversity. *Proc. R.*
704   *Soc. B Biol. Sci.*, 289, 20212690.

705   Ushio, M. (2022b). macam: A collection of miscellaneous functions for ecology in R.

706   Ushio, M., Furukawa, S., Murakami, H., Masuda, R. & Nagano, A.J. (2022). An efficient
707   early-pooling protocol for environmental DNA metabarcoding. *bioRxiv*.

708   Ushio, M., Hsieh, C., Masuda, R., Deyle, E.R., Ye, H., Chang, C.-W., *et al.* (2018a).
709   Fluctuating interaction network and time-varying stability of a natural fish community.
710   *Nature*, 554, 360–363.

711   Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., *et al.* (2018b).
712   Quantitative monitoring of multispecies fish environmental DNA using
713   high-throughput sequencing. *Metabarcoding Metagenomics*, 2, e23297.

714   Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
715   York.

716   Wieczynski, D.J., Singla, P., Doan, A., Singleton, A., Han, Z.-Y., Votzke, S., *et al.* (2021).
717   Linking species traits and demography to explain complex temperature responses
718   across levels of organization. *Proc. Natl. Acad. Sci.*, 118.

719   Wilke, C.O. (2017). *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."*

720   Wood, S.N. (2004). Stable and efficient multiple smoothing parameter estimation for
721   generalized additive models. *J. Am. Stat. Assoc.*, 99, 673–686.

Wootton, J.T. & Emmerson, M. (2005). Measurement of interaction strength in nature. *Annu.*

Rev. Ecol. Evol. Syst., 36, 419–444.

Wootton, K.L. & Stouffer, D.B. (2015). Many weak interactions and few strong; food-web

feasibility depends on the combination of the strength of species' interactions and

their correct arrangement. *Theor. Ecol.*

Yamamoto, S., Minami, K., Fukaya, K., Takahashi, K., Sawada, H., Murakami, H., *et al.*

(2016). Environmental DNA as a 'snapshot' of fish distribution: A case study of

Japanese jack mackerel in Maizuru bay, Sea of Japan. *PLOS ONE*, 11, e0149786.

Yang, S.-K., Nagata, Y., Taira, K. & Kawabe, M. (1993). Southward intrusion of the

intermediate Oyashio water along the east coast of the Boso Peninsula, Japan II.

Intrusion events into Sagami Bay. *J. Oceanogr.*, 49, 173–191.

Ye, H. & Sugihara, G. (2016). Information leverage in interconnected ecosystems:

Overcoming the curse of dimensionality. *Science*, 353, 922 LP – 925.

Tables

Table 1. Primer sequences used in the present study

Primer information	Primer sequence (5′– 3′)[1,2,3,4,5]	Length
1st PCR primers		
MiFish-U-forward	*ACACTCTTTCCCTACACGACGCTCTTCCGATCT* NNNNNN GTCGGTAAAACTCGTGCCAGC	60
MiFish-U-reverse	*GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT* NNNNNN CATAGTGGGGTATCTAATCCCAGTTTG	67
MiFish-E-forward-v2	*ACACTCTTTCCCTACACGACGCTCTTCCGATCT* NNNNNN RGTTGGTAAATCTCGTGCCAGC	61
MiFish-E-reverse-v2	*GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT* NNNNNN GCATAGTGGGGTATCTAATCCTAGTTTG	63
MiFish-U2-forward	*ACACTCTTTCCCTACACGACGCTCTTCCGATCT* NNNNNN GCCGGTAAAACTCGTGCC	57
MiFish-U2-reverse	*GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT* NNNNNN CATAGGAGGGTGTCTAATCCCCGTTTG	67
2nd PCR primers		
2nd-PCR-forward	<u>AATGATACGGCGACCACCGAGATCTACAC</u> **XXXXXXXX** *ACACTCTTTCCCTACACGACGCTCTTCCGATCT*	70
2nd-PCR-reverse	<u>CAAGCAGAAGACGGCATACGAGAT</u> **XXXXXXXX** *GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT*	66

[1]Normal characters indicate target-specific universal primers (i.e., MiFish primers).

[2]The six random bases (Ns) in the middle of the 1st PCR primers were appended to enhance cluster separation on flow cells.

[3]Italic characters indicate Illumina sequencing primers.

[4]**X** indicates index sequences to identify each sample.

[5]Underlined characters indicate P5/P7 adapter sequences for Illumina sequencing.

Figure legends:

Figure 1. Study sites and overall dynamics of environmental DNA (eDNA)

concentrations and the number of fish species detected. (**a**) Study sites in the Boso

Peninsula. The study sites are influenced by the Kuroshio current (red arrow; left panel) and

distributed along the coastal line in the Boso Peninsula (right panel). (**b**) Total eDNA copy

numbers estimated by quantitative eDNA metabarcoding (see Methods for detail). (**c**) Fish

species richness detected by eDNA metabarcoding. Points and lines indicate raw values and

LOESS smoothing lines, respectively. The line color indicates the sampling site. Warmer

colors generally correspond to study sites with a higher mean water temperature.

Figure 2. Interaction networks of the fish community in the Boso Peninsula coastal

region. The "average" interaction network reconstructed by quantifying information transfer

between eDNA time series. Transfer entropy (TE) was quantified by leveraging all eDNA

time series from multiple study sites to draw this network. Only information flow larger than

80% quantiles (i.e., strong interaction) was shown as interspecific interactions for

visualization. The edge color indicates scaled transfer entropy values, and fish illustration

colors represent their ecology (e.g., habitat and feeding behavior). Node colors and node sizes

indicate the fish family and fish abundance (total eDNA copy numbers of the fish species),

respectively.

Figure 3. Dependence of interaction strengths on biotic and abiotic variables (50

dominant fish species and 11 study sites were leveraged). The panels show the overall

effects of biotic and abiotic variables on interaction strengths of the 50 dominant fish species:

Effects of (**a**, **d**) water temperature, (**b**, **e**) species richness, and (**c**, **f**) total eDNA copy

numbers. The *y*-axis indicates the effects of the variables on fish-fish interaction strengths

quantified by the MDR S-map method. **a**–**c** show the effects on the species interactions that a

focal species receives (i.e., In-strength), and **d**–**f** show the effects on the species interactions

that a focal species gives (i.e., Out-strength). The line indicates the average effects estimated

by the general additive model (GAM), and the grey region indicates 95% confidential

intervals. LME and GAM indicate the statistical clarity of the linear mixed model portion and

GAM portion, respectively. Detailed statistical results and raw data are shown in Table S4

and Fig. S5, respectively.

Figure 4. Temperature dependence of fish species interactions at the species level. a and

b show temperature effects on fish species interactions quantified by the MDR S-map method.

Note that the MDR S-map enables quantifications of interaction strengths at each time point,

and thus the number of data points is large. (**a**) Points indicate the species interactions that a

focal species (indicated by the strip label and fish image) receives (i.e., In-strength). (**b**)

Points indicate the species interactions that a focal species (indicated by the strip label and

fish image) gives (i.e., Out-strength). For **a** and **b**, only fish species of which interactions are

statistically clearly affected by water temperature are shown (to exclude fish species with

relatively weak temperature effects, $P < 0.0001$ was used as a criterion here). Point color

indicates the study site. Gray line is drawn by GAM (the study sites were averaged for

visualization purpose).

Figure 1 | Study sites and overall dynamics of environmental DNA (eDNA) concentrations and the number of fish species detected. (**a**) Study sites in the Boso Peninsula. The study sites are influenced by the Kuroshio current (red arrow; left panel) and distributed along the coastal line in the Boso Peninsula (right panel). (**b**) Total eDNA copy numbers estimated by quantitative eDNA metabarcoding (see Methods for detail). (**c**) Fish species richness detected by eDNA metabarcoding. Points and lines indicate raw values and LOESS smoothing lines, respectively. The line color indicates the sampling site. Warmer colors generally correspond to study sites with a higher mean water temperature. For the high resolution figure, see https://bit.ly/3uUZKHE.

Figure 2 | Interaction networks of the fish community in the Boso Peninsula coastal region. The "average" interaction network reconstructed by quantifying information transfer between eDNA time series. Transfer entropy (TE) was quantified by leveraging all eDNA time series from multiple study sites to draw this network. Only information flow larger than 80% quantiles (i.e., strong interaction) was shown as interspecific interactions for visualization. The edge color indicates scaled transfer entropy values, and fish illustration colors represent their ecology (e.g., habitat and feeding behavior). Node colors and node sizes indicate the fish family and fish abundance (total eDNA copy numbers of the fish species), respectively. For the high resolution figure, see https://bit.ly/3uUZKHE.

Figure 3 | Dependence of interaction strengths on biotic and abiotic variables (50 dominant fish species and 11 study sites were leveraged). The panels show the overall effects of biotic and abiotic variables on interaction strengths of the 50 dominant fish species: Effects of (**a**, **d**) water temperature, (**b**, **e**) species richness, and (**c**, **f**) total eDNA copy numbers. The y-axis indicates the effects of the variables on fish-fish interaction strengths quantified by the MDR S-map method. **a-c** show the effects on the species interactions that a focal species receives (i.e., In-strength), and **d-f** show the effects on the species interactions that a focal species gives (i.e., Out-strength). The line indicates the average effects estimated by the general additive model (GAM), and the grey region indicates 95% confidential intervals. LME and GAM indicate the statistical clarity of the linear mixed model portion and GAM portion, respectively. Detailed statistical results and raw data are shown in Table S4 and Fig. S5, respectively.

Figure 4 | Temperature dependence of fish species interactions at the species level. **a** and **b** show temperature effects on fish species interactions quantified by the MDR S-map method. Note that the MDR S-map enables quantifications of interaction strengths at each time point, and thus the number of data points is large. (**a**) Points indicate the species interactions that a focal species (indicated by the strip label and fish image) receives (i.e., In-strength). (**b**) Points indicate the species interactions that a focal species (indicated by the strip label and fish image) gives (i.e., Out-strength). For **a** and **b**, only fish species of which interactions are statistically clearly affected by water temperature are shown (to exclude fish species with relatively weak temperature effects, $P < 0.0001$ was used as a criterion here). Point color indicates the study site. Gray line is drawn by GAM (the study sites were averaged for visualization purpose).