MCRS1 modulates the heterogeneity of microtubule minus-end morphologies in mitotic spindles

Alejandra Laguillo-Diego[1,§*], Robert Kiewisz[2,#*], Carlos Martí-Gómez[3], Daniel Baum[4], Thomas Müller-Reichert[2,+] and Isabelle Vernos[1,5,6,+]

[1] Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, 08003, Spain

[2] Experimental Center, Faculty of Medicine Carl Gustav Carus, Technische Universität Dresden, 01307 Dresden, Germany

[3] Simons Center for Quantitative Biology, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 11724, USA

[4] Department of Visual and Data-Centric Computing, Zuse Institute Berlin, 14195 Berlin, Germany

[5] Universitat Pompeu Fabra (PF), Barcelona, 08003, Spain

[6] ICREA, Pg. Lluís Companys 23, Barcelona 08010, Spain

* These authors contributed equally.

+ Shared last authors

[§] Current address: Sanford I. Weill Department of Medicine, Sandra and Edward Meyer Cancer Center, Weill Cornell Medicine, New York, NY 10021, USA

[#] Current addresses:

Simons Machine Learning Center, New York Structural Biology Center, New York, NY 10027, USA

Biocomputing Unit, Centro Nacional de Biotecnologia (CNB-CSIC), Darwin, 3, Campus Universidad Autonoma, 28049 Cantoblanco, Madrid, Spain

27 Abstract

Faithful chromosome segregation requires the assembly of a bipolar spindle, consisting of two antiparallel
microtubule (MT) arrays having most of their minus ends focused at the spindle poles and their plus ends
overlapping in the spindle midzone. In mammalian cells, bipolar spindles have bundles of kinetochore
microtubules (KMTs), called kinetochore (k)-fibers that attach the chromosomes to both spindle poles.
Spindle assembly, chromosome alignment and segregation require highly dynamic MTs. The plus end
dynamics of MTs has been extensively investigated; instead, their minus end dynamics remains poorly
characterized. Here, we used large-scale electron tomography of HeLa cells to study the morphology of
the MT minus ends in 3D-reconstructed metaphase spindles. We found that the MT minus ends are
heterogeneous showing either open or closed morphologies. Silencing the minus-end specific stabilizer,
MCRS1 increases the proportion of open MT minus ends, suggesting that this morphology corresponds
to dynamic MT ends. Altogether, our data indicate that the dynamics of the spindle MT minus ends is
more complex than anticipated with unsynchronized dynamic states within the k-fibers.

Introduction

The mitotic spindle is a highly complex molecular machinery that assembles during cell division to segregate chromosomes to the daughter cells. It is structurally and functionally defined by its main component, the microtubules (MTs). During mitosis, MTs are nucleated through different pathways, involving the centrosomes, the chromosomes and pre-existing MTs (Lüders, 2016; Meunier and Vernos, 2012; Teixido-Travesa et al., 2012). Altogether, these MTs organize the bipolar spindle, which consists of two anti-parallel arrays having MT plus ends at the spindle equator and the minus ends mainly associated with the two spindle poles (i.e., the centrosomes). Within the spindle of mammalian cells, kinetochore MTs (KMTs) organize into bundles, so-called kinetochore-fibers (k-fibers) that connect chromosomes to the two spindle poles. In addition, astral (AMTs) and interpolar MTs (IMTs; both termed non-KMTs here) constitute the main body of the spindle, providing additional forces for spindle positioning and chromosome movements by interaction with molecular motors (Meunier and Vernos, 2012; Meunier and Vernos, 2016).

The regulation of MT dynamics is fundamental for the overall organization of the spindle and the segregation of the chromosomes. Dynamics can be observed at both MT ends that can stochastically switch between phases of growth and shrinkage (Kirschner and Mitchison, 1986; Mitchison and Kirschner, 1984). MT plus-end dynamics promotes the attachment of MTs to the kinetochores, thus driving the assembly of the k-fibers (Mimori-Kiyosue and Tsukita, 2003). The dynamic nature of these attachments is key for error correction, the activity of the spindle assembly checkpoint (SAC) and chromosome segregation (Foley and Kapoor, 2013; Lampson and Grishchuk, 2017; Maiato et al., 2004; Monda and Cheeseman, 2018). Although MT minus-end dynamics has been described as essential for spindle assembly and for chromosome movements, little information is currently available on the structure and dynamics of the MT minus ends in the mammalian spindle, as most of them are focused at the spindle poles, an extremely crowded region difficult to observe and manipulate (Akhmanova and Steinmetz, 2019).

In mammalian cells, most MTs are nucleated by the γ-TuRC, a dedicated multiprotein complex, Although MTs in mitosis are nucleated through various pathways, spindle pole-located γ-TuRC plays a crucial role in MT nucleation. In general, MTs nucleated from γ-TuRC are "capped" and their minus ends and they are therefore stabilized (Consolati et al., 2020; Wieczorek et al., 2020; Zimmermann et al., 2020). However, various studies indicate that MT minus ends depolymerize at the spindle poles (Jiang et al., 2017). Indeed, a movement of tubulin subunits from the spindle equator to the spindle poles was observed by live-cell imaging upon photoactivation of a tubulin stripe in the close proximity to the aligned chromosomes in the metaphase spindle (Mitchison, 1989). This phenomenon, named spindle flux, was proposed to play a role in the control of spindle length and in chromosome segregation (Ganem and Compton, 2006; Ganem et al., 2005; Steblyanko et al., 2020). The current view is that the spindle flux is

the result of a combination of MT transport towards the spindle poles and a net incorporation of tubulin at
the MT plus ends. This incorporation is compensated by the removal of tubulin dimers at MT minus ends
at the spindle poles (Kirschner and Mitchison, 1986; Mitchison, 1989; Waters et al., 1996) caused by
kinesin-13 depolymerases (Ems-McClung and Walczak, 2010). Consistently, silencing the spindle pole-
localized kinesin-13 depolymerase, kif2a leads to the formation of monopolar spindles, indicating that the
regulation of MT minus end dynamics is essential for bipolar spindle assembly (Ganem and Compton,
2004). Additional support for this idea was also provided by the identification of Microspherule Protein 1
(MCRS1), a RanGTP-regulated protein that localizes to the KMT minus ends and regulates their
depolymerization rate at the spindle poles. Spindles assembled in MCRS1-silenced cells have a faster
poleward flux, show hyper-stretched kinetochores and form unstable spindles (Meunier and Vernos,
2011).

Several studies indicate that the dynamic state of MTs is associated with specific end morphologies.
Cryo-electron microscopy studies of *in vitro*-assembled MTs showed that fast growing MTs have flared
ends with curved sheet-like protofilaments at their tips, while slowly growing MTs have blunt ends. In
contrast, depolymerizing MT ends display outward-curled (also called ramshorn-like) protofilaments
(Chretien et al., 1995; Mandelkow et al., 1991; Muller-Reichert et al., 1998; Rice, 2018; Simon and
Salmon, 1990) . Importantly, closed MT ends have also been observed. MTs nucleated by the yTuRC
complex *in vitro* show an electron-dense material 'closing' their minus-ends (Zheng et al., 1995).

Although MT dynamics in cells is more complex than *in vitro* due to the presence of a large variety of MT-
associated proteins (MAPs), including some that specifically bind to the MT ends, the morphologies of
both growing and depolymerizing ends were found to be very similar to those observed *in vitro*
(VandenBeldt et al., 2006). Moreover, recent electron microscopy studies of plastic-embedded spindles
also revealed flared and curled MT end morphologies (O'Toole et al., 2003)., suggesting that the spindle
MTs may be in different phases of polymerization and depolymerization, respectively. MTs having a blunt
end may be pausing, polymerizing or depolymerizing, and therefore it is difficult to categorize them in any
specific dynamic state Interestingly, studies performed in early *C. elegans* embryos showed that the MT
minus ends associated with the mitotic centrosomes had heterogeneous morphologies described as
either closed or open (O'Toole et al., 2003). The presence of a "cap-like" structure at the MT minus ends
associated with either spindle pole bodies in S. pombe or mitotic centrosomes in C. elegans embryos was
interpreted as the yTuRC (Hoog et al., 2011; O'Toole et al., 2003; Teixido-Travesa et al., 2012). However,
there is currently little information concerning the MT minus ends in the mitotic spindle poles of
mammalian cells.

Here, we set out to gain novel insights into the MT minus-end morphologies in the metaphase spindle of
human cells using electron tomography. We found that MT minus ends can have either closed or open
morphologies in a proportion that is modified upon silencing the MT minus end regulator MCRS1. These

data confirm the role of MCRS1 at the MT minus ends during mitosis. The heterogeneity of KMT minus-
end morphologies suggests a complex mechanism of regulation of their dynamics.

Results

Different MT minus-end morphologies co-exist at the spindle poles

We used 3D reconstructions of whole mitotic spindles assembled in HeLa cells (data from Kiewisz et al. 2021) to analyze the morphology of the MT ends (**Figure 1A**; **Table 1**). In these control cells, we defined two main classes of ends: closed ends that have an electron-dense "cap" (**Figure 1B**, left panels; **Figure 1-figure supplement 1A**) and open ends that typically have a flared morphology (**Figure 1B**, right panels; **Figure 1-figure supplement 1B**). Other types of MT ends were classified as 'undefined'.

We first focused on the MTs forming the k-fibers (i.e., the KMTs), strictly defined as those directly associated through their plus ends to the kinetochores at the chromosome surface (Kiewisz et al., 2021). In agreement with the dynamic nature described for KMT plus ends (Cheeseman and Desai, 2008), the close examination of the KMT end morphology at the kinetochore showed that they were open (green circles) and flared (**Figure 1C**; **Figure 1-movie supplement 1-2; Table 2**). These results indicate that there is a correlation between the morphology and dynamics of MT ends, as previously described, (Hoog et al., 2011). We then focused on the opposite end of the KMTs away from the kinetochore defining this one as the minus end. KMT minus ends had either closed (**Figure 1C**, circles in purple) or open morphologies (**Figure 1C**, circles in green). The heterogeneous morphologies of the KMT minus ends suggest that they may co-exist in different dynamic states.

To determine whether there was any correlation between the KMT minus-end morphologies and their position along the half-spindle axis, we then looked at their distribution along this axis. About half of the KMT minus ends were located at the spindle poles, while the other half was distributed along the half-spindle (Kiewisz et al., 2021). We went on to determine the relative position of each KMT minus end on the half-spindle axis (**Figure 1D**). The half-spindle length was defined as the distance between each of the spindle poles (position = 0) and the individual kinetochores (position = 1). We then estimated the proportion of open minus ends at each relative distance along the half-spindle axis (**Figure 1E**). The inferred proportion of the open minus ends in the proximity of the centriole pair (defining the spindle pole) was 52.97% (95% credible interval (CI) = [21.48, 77.43]%) (**Figure 1E,** gray area D< 0.2, region where the majority of KMT minus ends are located). Therefore, half of the KMT minus ends at the spindle poles is open, while the other half is closed (i.e., capped), suggesting that they are in different dynamic states.

We then asked whether KMTs belonging to the same k-fiber might have similar minus end morphologies indicative of a synchronized dynamic state. To address this question, we performed a model comparison using a likelihood ratio test between models that included or ignored the k-fiber as a random effect. We found no evidence of a clustering of open minus ends at specific k-fibers (p-value = 1). This suggested that the minus-end morphology of any given KMT within a k-fiber is independent from the minus end morphologies of the other KMTs in the same k-fiber.

To determine whether the mixture of minus end morphologies suggestive of mixed dynamic states is
specific for the KMTs or a more general feature of the spindle MTs, we extended our studies to the non-
KMTs. For these MTs, we defined the average position of all kinetochores (**Figure 1D,** aK) as position 1
on the half-spindle axis. In addition, the polarity of the non-KMTs was determined by the positioning of the
ends along the half-spindle axis. The end of each non-KMT closer to the spindle pole was defined as the
minus end, with the other end assigned as the plus end. The end morphologies of non-KMTs were also
classified as either closed or open. We then estimated the proportion of open minus ends at each relative
distance. Interestingly, we found that 64.04% (CI = [33.00, 86.22]%) of the non-KMT minus ends were
open in the region near the centriole pair (**Figure 1F**, gray area). Again, these data showed that MT
minus ends have heterogeneous morphologies, overall suggesting the coexistence of different dynamic
states. This further suggests previously unrecognized mechanisms that regulate the MT minus end
dynamics.

MCRS1 silencing induces changes in k-fiber ultrastructure and spindle shape

MCRS1 was shown to associate with the k-fiber minus ends in metaphase and regulate their dynamics
(Meunier and Vernos, 2011). Therefore, we decided to examine the putative changes in the MT end
morphologies in spindles assembled in MCRS1-silenced cells by electron tomography. Since siRNA-
based gene silencing may not be homogeneous in a cell population, we first carefully looked for
morphological spindle features that could be altered in MCRS1-silenced cells and used as a signature for
selecting the spindles to be processed for electron tomography. Western blot analysis of MCRS1-silenced
cells showed a reduction in the target protein level of close to 90% (**Figure 2A-B**). High-contrast light
microscopy images revealed specific changes in the half-spindle shape and the outer spindle MT angle
(**Figure 2C**). siScramble spindles showed an average half-spindle angle of 80.81º ±10.60 (**Figure 2E**, n =
105,). In contrast, MCRS1-silenced cells had an average half-spindle angle of 71.5º ±8.05 (n= 110; p-
value = 4.729 x 10^{-8}), significantly narrower than in siScramble cells. In addition, siScramble cells showed
an average angle for the outer MTs of 152.34º ± 9.62 (**Figure 2G**, n = 201,), whereas siMCRS1-depleted
cells displayed an average angle for the outer MTs of 167.51º ±10.72 (n = 161, p-value = 7.486 x 10^{-26}).
Thus, the outer MTs in MCRS1-depleted cells were significantly straighter than in control cells. Similar
differences were also clearly observed in 3D models after processing both siScramble and MCRS1-
silenced cells for electron tomography. In stacked serial plastic sections (**Figure 2D**; **Figure 2-movie
supplement 1-2**), siScramble cells had a half-spindle angle of 90.62º ±6.41 (**Figure 2F**, n = 4), whereas
MCRS1-silenced spindles had a significantly lower half-spindle angle of 80.13º ±5.77 (n = 8, p-value=
0.017). Moreover, siScramble spindles showed an average outer MT angle of 145.77º ±5.55 (**Figure 2H,**
n = 8,), whereas siMCRS1 spindles displayed a significantly higher average outer MT angle of 158.00º
±8.74 (n = 16, p-value = 0.002). From these data, we concluded that depletion of MCRS1 caused a
reduction in the half-spindle angle and a simultaneous increase in the MT angle.

These parameters were then used for the selection of two spindles from siScramble cells and five
spindles from MCRS1-silenced cells for 3D reconstruction of quarters of spindle volumes that we
validated as representative of full spindles (**Figure 3-figure supplement 1**). To determine whether all the
selected spindles were at a similar stage in mitosis, we compared the average outer-kinetochore distance
for all our data sets. The average inter-kinetochore distance in the selected spindles was 1.06± 0.21 µm
(n = 28) in siScramble cells and 1.06± 0.21 µm (n = 55) in MCRS1-silenced cells. Control spindles
showed an average value of 1.07± 0.21 µm (n = 146; siScramble *versus* Control, p-value = 0.108;
siScramble *versus* siMCRS1, p-value = 0.240). Additionally, the values measured for Control and
siScramble cells were in agreement with previous studies (Kiewisz et al., 2021). The high similarity in the
inter-kinetochore distance in all these conditions indicated that the selected spindles were captured at a
similar mitotic stage.

We then analyzed the 3D reconstructions of the selected spindles. Spindles assembled in MCRS1-
silenced cells showed distinct ultrastructural features. The k-fibers were indeed straighter in siMCRS1
spindles than in the siScramble ones (**Figure 3A; Figure-movie supplement 1-2**). Interestingly, the
KMTs did not reach the centriole pair area at the spindle poles in siMCRS1 spindles, whereas the
centriole pair was embedded within the mass of KMT minus ends in the siScramble spindles.
Consistently, the quantification of MT minus end distribution revealed that the majority reached the
centrioles in control (**Figure 3B**, mean peak at position = 0.03) and siScramble spindles (**Figure 3B**,
mean peak at position = 0.04, p-value= 0.201). In contrast, the MT minus ends in siMCRS1 spindles
peaked at a mean position of 0.07, further away from the centrioles (**Figure 3B**, p-value = 0.018;
siScramble vs siMCRS1 at D < 0.2). This increased distance between the centrioles and the KMT minus
ends in MCRS1-silenced cells is fully consistent with the proposed role of MCRS1 in controlling the rate of
MT minus end depolymerization. We then determined the number of KMTs per k-fiber in the three
different conditions (**Figure 3C**) and evaluated the putative differences using a Generalized Linear Model
(GLM) with Poisson likelihood and log link function, using siMCRS1 and siScramble data as covariates.
We found that the number of KMTs per k-fiber was reduced in MCRS1-silenced cells with an average of
7.27 KMT (n = 106, p-value = 0.0002) instead of 9.19 KMTs per k-fiber (n = 44) in siScramble spindles.
No significant differences were found between this value in siScramble and control cells (8.93 KMTs per
k-fiber in control cells, n= 226, p-value = 0.599). Altogether, our analysis revealed a series of
morphological alterations in the spindles assembled in MCRS1-silenced cells consistent with changes in
the dynamics of the MT minus ends in these cells as previously proposed (Meunier and Vernos, 2011).

Silencing of MCRS1 increases the proportion of open minus ends of all spindle MTs

To explore more specifically the impact of MCRS1 silencing on the spindle MTs, we then quantified the
morphology and relative distribution of their minus ends within the spindle. We selected the KMTs in the

reconstructions of the siMCRS1 spindles as described above and manually classified their minus end
morphologies into the three categories: open, closed and undefined (**Figure 4E**; **Figure 1B**). Next, we
determined the relative distance of each KMT minus end along the half-spindle axis. The percentage of
open KMT minus ends in the siMCRS1 spindles was 75.28% (CI= [44.83, 92.32]%, n = 386) in the region
surrounding the centriole pair (d < 0.2; **Figure 4A; Figure 4-movie supplement 1**, green line), whereas it
was 56.17% in control cells (CI= [25.05, 79.23]%, n = 2071). To compare both percentages, as
percentages are bound between 0 and 100, we used the log2 transformation of the ratio open/closed. In
this new scale, 0 means that there are the same number of open and closed minus ends;
log2(open/closed) = 1 indicates that there are 2 times more open than closed minus ends, and -1 that
there are 2 times more closed than open minus ends. This quantity is now unbound and symmetric
regardless of the group order, so it is more appropriate to represent differences between 2 groups in
percentages. The log2(open/closed) was 1.38 higher in siMCRS1 cells compared to control cells around
the centrioles (CI = [0.31, 2.39]; **Figure 4B**), indicating that MCSR1 modulates KMTs minus ends
capping.

We then examined the minus end morphologies of a randomly selected group of non-KMTs at the spindle
poles following the procedure described above. We found that the estimated percentage of open non-
KMT minus ends in the region near the centriole pair (d < 0.2) was 80.76% (CI = [53.16, 93.31]%) in
MCRS1-silenced cells and 64.04% (CI = [33.00, 86.21]%) in control cells (**Figure 4C**). As before, we used
the log2 transformation of the ratio open/closed to compare both percentages. The log2(open/closed) was
1.35 higher in siMCRS1 cells compared to control cells for non-KMTs (CI = [0.56, 2.14], **Figure 4D**) for
the non-KMT minus ends. This analysis revealed that the non-KMT minus end morphologies are also
altered close to the spindle pole in the absence of MCRS1.

Previous experiments suggested that MCRS1 function was specific for the regulation of the KMT minus
end dynamics (Meunier and Vernos, 2011). To directly address this hypothesis, we analyzed our data to
determine whether the change in the ratios of close and open minus ends for KMTs and non-KMTs in
control *versus* siMCRS1 cells was similar or not. We eliminated the distance-dependence of our analysis
and restricted our quantification to the spindle-pole region (d < 0.2) for both MT populations (**Figure 4F**).
Considering both categories together, the change in log(open/closed) between siMCRS1 and control cells
for non-KMTs was not significantly different from that in KMTs (Δlog(open/closed) = -0.07, p-value =
0.79). Altogether, these results suggest that MCRS1 silencing has a similar impact on the morphology of
both KMT and non-KMT minus ends at the spindle poles with an increase in the percentage of open ends
in the absence of MCRS1.

248 Discussion

The depolymerization of MT minus ends has been postulated to play an essential role in the control of
spindle size, k-fiber dynamics and chromosome movements (Barisic et al., 2021; Waters et al., 1996).
However, there is currently little information on the precise mechanism that establishes and controls MT
minus-end dynamics at the spindle poles. A direct visualization of the MT minus ends at any given point
may thus provide insightful clues about their actual dynamics. Here, we applied large-scale electron
tomography (Redemann et al. 2017; Kiewisz et al. 2021) to directly visualize MT minus end morphologies
at mitotic spindle poles in human cells in metaphase. We found that MT minus ends do not have
homogeneous morphologies. Rather, KMTs and non-KMTs display a mixture of open and closed minus
ends, in similar proportions (**Figure 5**). Previous studies in cells established a correlation between the
ultrastructure of MT ends and their dynamic state (Hoog et al., 2011) (Höög et al. 2011). Here, we show
that the dynamic KMT plus ends associated with the kinetochores are mainly open and flared. This is in
agreement with previous studies suggesting that dynamic MTs have these types of end morphologies
(McIntosh et al., 2008; VandenBeldt et al., 2006). The closed conformation that we observed for a large
proportion of MT minus ends is reminiscent of previous reports on non-dynamic MT minus ends capped
by the y-TuRC (Zheng et al., 1995). This suggests that the closed end morphology may correspond to
stable and/or anchored MTs. The mixture of morphologies that we observed for the MT minus ends in the
spindle also indicate that their depolymerization at the spindle pole is not synchronous, not even for KMTs
within the same fiber.

We observed a significant increase in the proportion of open minus ends at the spindle poles upon
silencing of the proposed regulator of MT minus-end dynamics, MCRS1 (Meunier and Vernos, 2011). In
addition, we observed an increase in the relative distance of the KMT minus ends to the centriole pair in
these cells (**Figure 4F**), also consistent with an increase in MT minus-end depolymerization (Meunier and
Vernos, 2011). Moreover, the loss of curvature of the KMTs (**Figure 3A**) suggests an increase in the
tension that could also fit with previous reports about the forces exerted by MT minus end
depolymerization on the attached chromosomes (DeLuca et al., 2006; Waters et al., 1996).

MCRS1 was identified as a novel RanGTP-regulated factor that associates specifically with the k-fiber
minus ends in cold-treated cells (Meunier and Vernos, 2011). We found here that both KMT and non-KMT
minus-end morphologies change in MCRS1-silenced cells, suggesting that MCRS1 may associate with
both MT populations in the spindle. However, since it was recently reported that KMTs directly interact
with many non-KMTs near the spindle pole it is possible that some MTs we classified as non-KMT are in
fact considered as part of the k-fibers (Kiewisz et al., 2021). The impact of MCRS1 silencing on the non-
KMT minus end morphology could therefore indicate that MCRS1 could act on the non-KMTs that are
included in the k-fiber. Alternatively, MCRS1 may have a more general function in regulating MT minus
end dynamics within the spindle than previously proposed.

It is interesting to speculate about the nature of the complex that associates with the MT minus ends to
generate a cap structure. The γ-TuRC-ring complex with an approximate molecular weight of 2353.97
kDa appears as a cone-like structure as observed *in vitro* (Consolati et al., 2020; Drutovic et al., 2020;
Wieczorek et al., 2020; Zimmermann et al., 2020). Some MT minus ends may be 'capped' by this
complex but since they are thought to depolymerize when reaching the spindle pole, we can speculate
that the electron-dense material could correspond to other MT minus-end binding complexes such as
ASPM-katanin (Jiang et al. 2017), the MCRS1-KANSL-complex (Meunier et al., 2015) and/or
NuMA/dynein (Elting et al., 2014) (**Figure 4F**).

Recent data suggested that MT minus end depolymerization and the spindle flux can be uncoupled
(Steblyanko et al., 2020). Instead, our data suggest that there is a direct correlation between the
proportion of open MT minus ends and the speed of the spindle flux that was reported to increase in
absence of MCRS1 (Meunier and Vernos, 2011). The question remains, however, how can cells undergo
constant spindle flux, when MT minus ends are not synchronously depolymerizing? Our data can provide
only a snapshot of the MT minus end morphologies in spindles. Most likely, the binding/unbinding of
components at the MT minus ends is stochastic. In such a model, MT minus ends would overall
depolymerize, although individual MTs may pause, which would be visible as a heterogeneity in the
morphology of individual ends (**Figure 4F**). A similar situation but with a net polymerization instead of a
depolymerization has been described for MT plus ends (VandenBeldt et al., 2006). The putative
mechanism underlying MT minus end dynamics at the spindle poles still needs to be addressed in the
future (Chen et al., 2019; Dudka et al., 2019).

Experimental procedures

Standard techniques

Cell culture

HeLa (Kyoto) cells were grown at 37ºC in a 5% CO_2 atmosphere. The cells were cultured in DMEM medium (Thermo Fisher Scientific, Waltham, MA) containing 4.5 g/L glucose and supplemented with L-glutamine with sodium pyruvate (Lonza, Basel, Switzerland), 10% fetal bovine serum (Thermo Fisher Scientific, Waltham, MA) and an antibiotic cocktail containing 100 µg/ml streptomycin and 100 units/ml penicillin (Sigma-Aldrich, St Louis, MO). For regular maintenance, HeLa cells were split with 0.25% Trypsin (Sigma-Aldrich, St Louis, MO) when reaching around 80-90% confluency.

RNA interference

Cells were seeded at 150.000 cells/ml in 75 cm^2 flasks (Sigma-Aldrich, St Louis, MO) the day before transfection. Per flask, 500 pmol of siRNA and 25 µl Lipofectamine RNAiMax transfection reagent (Life Technologies, Carslbad, CA) were used and scaled accordingly for smaller volumes. Transfection was performed following the manufacturer's protocol. After 48 hours post-transfection, cells were split and seeded. At 60 hours, cells were transfected as described above using siRNA purchased from Dharmacon (Lafayette, CO).

Gel electrophoresis and Western Blots

Protein lysates from HeLa cells were prepared by resuspending cell pellets in RIPA buffer containing 50mM Tris-HCl pH 7.4, 150 mM NaCl, 2 mM EGTA, 1% Triton X-100 (Sigma-Aldrich, St Louis, MO), 0.5% DOC (Sigma-Aldrich, St Louis, MO) and 0.1% SDS supplemented with protease inhibitors (Sigma-Aldrich, St Louis, MO). After incubation on ice for 15 minutes, the lysates were centrifuged at 4ºC at maximum speed for 15 minutes. The protein content of the lysates was determined by using the Pierce BCA kit (Thermo Fisher Scientific) and following the manufacturer's instructions. For each run, 50 µg of protein lysate was diluted in loading buffer 5x, boiled for 10 minutes and loaded in a 10% SDS-PAGE gel. Gels were run at 120 V for 90 minutes.

For Western Blots, a semi-dry transfer was done to blot the proteins in a 0.45 µm nitrocellulose membrane (GE Healthcare, Chicago, IL). Proteins were transferred for 90 minutes at 65 mA. The membrane was blocked using 5% milk (Sigma-Aldrich, St Louis, MO) in TBS at 4ºC overnight. Primary antibodies were incubated in 0.1% Tween20 (Sigma-Aldrich, St Louis, MO) in TBS for 1 hour at room temperature. Secondary antibodies conjugated with AlexaFluors (AlexaFluors800 – MCRS1;

333 AlexaFluors680 - α-Tubulin) were incubated in the same buffer at room temperature for 45 minutes. Blots
334 were developed using an Odyssey Infrarred Imaging System (Li-cor, Lincoln, NE).

Light microscopy

For immunofluorescence, HeLa cells were grown on 18-mm round coverslips (Marienfeld, Germany) in
six-well plates. Cells were fixed by immersion in cold methanol (-20ºC) for 10 minutes. Cells were then
blocked and permeabilized in 0.5% BSA (Panreac, Barcelona, Spain) and 0.1% Triton X-100 (Sigma-
Aldrich, St Louis, MO) in 1x PBS for 30 minutes at room temperature. Primary antibodies were incubated
in the same blocking buffer for 1 hour at room temperature. Secondary antibodies conjugated with
AlexaFluors 488 and Hoechst to stain DNA (Life Technologies, Carlsbad, CA) were diluted in blocking
buffer 1:1000 and incubated at room temperature for 45 minutes. Coverslips were mounted in 9.6% (w/v)
homemade Mowiol, containing 24% (w/v) glycerol (Sigma-Aldrich, St Louis, MO) and 0.1 M Tris-Cl).

After immunostaining, samples were visualized by light microscopy to select MCRS1-depleted cells. The
samples were visualized with a LEICA SP5 microscope (Leica Microsystems, Austria) equipped with a
water 63x objective lens (Leica HCX PL APO 63x/1.2 W) and a PMT Leica hybrid detector camera (HyD)
(Dresden). Images were acquired using a 488 nm argon laser with a HyD sensor set to collect light from
500nm to 550 nm and a 594 nm diode laser with a sensor set from 620 nm to 730 nm. Chromosome
alignment at the metaphase plate was considered as a readout for MCRS1 depletion. Selected mitotic
cells in the metaphase stage were further characterized by measuring both the half-spindle and the MT
angle using Fiji (Schindelin et al. 2012) (**Figure 2B**). For this, the maximum intensity projection tool was
used for the z-stacks corresponding to the α-tubulin staining in the selected spindles. A threshold was set
for the projected images to create a binary mask. The angles were measured manually using the angle
tool. The half-spindle angle was measured to analyze the shape of the spindle poles, and it was
measured the outline of the half-spindle by calculating an angle between the metaphase plate and spindle
pole (**Figure 2B**). The MT angle was measured to analyze the degree of bending of the outer MTs. We
measured the outline of the quarter-spindle by calculating the angle between the metaphase plate and
the spindle pole (**Figure 2B**). Differences in the measured angles in both siScramble and siMCRS1
spindles were assessed by applying a linear regression model.

Electron microscopy

Sample preparation

In preparation for electron microscopy, 3-mm and 6-mm sapphire discs (M. Wohlwend GmbH,
Switzerland) were prepared for the attachment of cells. The sapphire discs were carefully cleaned by
immersion in a Piranha solution (1:1 H2SO4 and H202, v/v), coated with poly-L-lysine 0.1% (w/v) and
dried at 60ºC for 2 hours. Finally, the discs were incubated with a 1:10 solution of fibronectin (Sigma-

Aldrich, St Louis, MO) in PBS at 37ºC for 2 hours. The mitotic fraction of the cell culture was then
collected by applying the "shake-off" technique (Kiewisz, Müller-Reichert, and Fabig 2020). Briefly, flasks
with HeLa cells were subjected to two "shake-off" rounds. Flasks were hit against the surface of the
bench to detach mitotic cells. The collected cells were then centrifuged at 1200 rpm at 37ºC for 4 minutes
and the cell pellets were resuspended in 1 ml of DMEM medium (Thermo Fisher Scientific, Waltham, MA)
supplemented with 10% bovine serum albumin (BSA; Thermo Fisher Scientific, Waltham, MA) and 10%
of fetal bovine serum (Thermo Fisher Scientific, Waltham, MA). Next, the cells were allowed to attach to
the sapphire discs for 10 minutes at 37ºC.

High-pressure freezing and freeze substitution

Cells attached to 3-mm sapphire discs were cryo-immobilized using an EM ICE (Leica Microsystems,
Austria) and 6-mm sapphire discs were frozen by using a Compact 03 high-pressure freezer (M.
Wohlwend GmbH, Switzerland). For each freezing round with the EM ICE, a type-A aluminum carrier
(Leica Microsystems, Austria) with the 100 µm-indentation facing-up was placed in the specimen loading
device of the freezer. The cavity of the carrier was then filled with 5 µl of DMEM containing 10% BSA and
the sapphire disc with the attached HeLa cells facing down was placed onto the carrier. Next, a spacer
ring was mounted on top of the sample and the freezing started immediately. For using the Compact 03
high-pressure freezer, the 6-mm sapphire discs with the mitotic cells facing down were placed on a type-A
aluminum planchette with the 40µm-deep cavity pre-filled with warm DMEM medium supplemented with
10% BSA. The closed carrier was then placed in the specimen holder, clamped with the holder arm and
immediately cryo-immobilized. Using both freezers, samples were frozen at ~2,000 bar with a cooling rate
of ~20,000ºC/s (Reipert, Fischer, and Wiche 2004). After freezing, samples were stored in liquid nitrogen
until further use. Freeze substitution was carried out as previously described (Muller-Reichert et al.,
2003). Briefly, samples were transferred to cryo-vials filled with anhydrous acetone containing 1%
osmium tetroxide (EMS, USA) and 0.1% uranyl acetate (Polysciences, USA). Freeze-substitution was
done using an automatic freeze substitution machine (EM AFS, Leica Microsystems, Germany). Samples
were kept at -90ºC for 1 hour, then warmed up to -30ºC in steps of 5ºC per hour and maintained at -30ºC
for 5 hours. Next, the temperature was increased to 0ºC in steps of 5ºC per hour. Finally, after reaching
0ºC samples were washed three times with pure anhydrous acetone at room temperature.

Sample embedding, pre-selection of mitotic cells and ultramicrotomy

For resin embedding, the samples were placed in flow-through chambers (Leica, Germany) and infiltrated
with Epon/Araldite (EMS, Hatfield, PA) in three steps of 1 hour each with increasing concentration of
resin: 1:3, 1:1 and 3:1 (resin:acetone, w/v), followed by a single step overnight in pure resin at room
temperature (Müller-Reichert et al. 2003). Then, samples were polymerized at 60ºC for 48 hours. After
polymerization the plastic samples were removed from the flow-through chambers. Using a razor blade,

the sapphire discs were then removed from the resin blocks to expose the embedded cells for further
processing as described previously (Kiewisz et al., 2021).

In order to screen for cells in metaphase, the resin blocks were observed from the top using an upright
brightfield microscope (Zeiss, Germany). The two main features used to select cells in metaphase were a
rounded shape and a distinguishable metaphase plate (Kiewisz et al., 2021). Serial semi-thick (300 nm)
sections were cut using an EM UC6 ultramicrotome (Leica Microsystems, Austria) and collected on
Formvar-coated slot grids. Samples were post-stained with 2% uranyl acetate for 10 minutes followed by
a 0.4% of Reynold's lead citrate solution (Science Services, Germany). Colloidal 15nm-gold particles
(British Biocell International, UK) were attached to both sides of the sections mounted on the grids to
serve as fiducial markers for tomographic reconstruction.

Final staging of pre-selected spindles

The serial sections were imaged using a TECNAI T12 Biotwin transmission electron microscope
(ThermoFisher Scientific, Waltham, MA) operated at 120kV and equipped with an F214 CCD camera
(TVIPS GmbH, Germany). Images of whole cells in metaphase were acquired at 1200x magnification
using the EMMenu Software (TVIPS GmbH, Germany). To choose a cell for electron tomography, the
metaphase plate had to be correctly formed when looking at the chromosome area in 3D, and for each
chosen cell the half spindle and the MT angle were measured in the 3D volumes as described (see the
section on immunofluorescence). For this, the EM stacks were projected in 3D. The chromosome and
microtubule area were estimated by manually labeling the chromosome and MT area. Both the half-
spindle angle and MT angle were finally calculated using the ZIB Amira software (Zuse Institute Berlin,
Germany).

Three-dimensional reconstruction by electron tomography

Data acquisition and calculation of tomograms

Electron tomography was performed on the selected metaphase cells as previously described (Kiewisz et
al. 2021). Briefly, a series of tilted views were recorded using a TECNAI F30 transmission electron
microscope (ThermoFisher Scientific, Waltham, MA) operated at 300 kV and equipped with a Gatan
US1000 2K x 2K CCD camera. The SerialEM software package was used for the acquisition and
montaging of data sets (Mastronarde, 2003; Mastronarde, 2005). For dual-axis electron tomography,
images were captured every 1.0° over a ±60° range at a pixel size of 2.32 nm. For a recording of the
second axis, grids were rotated by 90º and another series of tilted views was acquired (Mastronarde,
1997). Tomograms were calculated by using the IMOD software package (Mastronarde and Held, 2017).
To increase the sample number, we acquired volumes of quarters of the cells (see **Table 1**).

 Segmentation of MTs and stitching of serial tomograms

MTs were semi-automatically segmented using the ZIB Amira (Zuse Institute Berlin, Germany) software
package (Lindow et al., 2021) as previously described (Lindow et al., 2021; Redemann et al., 2014;
Weber et al., 2012). After manual correction of MT segmentation, the serial tomograms of each recorded
cell were stitched using the segmented MTs as alignment markers (Lindow et al., 2021; Weber et al.,
2014).

Z-correction of stacked tomograms

Each stack of serial tomograms was expanded in *Z* to correct for a sample collapse during the data
acquisition (McEwen and Marko, 1999). We corrected this shrinkage by applying a Z-factor to the stacked
tomograms (Kiewisz et al., 2021; O'Toole et al., 2020). Taking the microtome setting of around 300 nm,
we multiplied this value by the number of serial sections. For each spindle, we also determined the
thickness of each serial tomogram and then calculated the total thickness of the reconstruction. The *Z*-
factor was then determined by dividing the actual thickness of each stack of tomograms by the total
thickness as determined by the microtome setting. Such calculated Z-factors were then applied to our full
spindle reconstructions.

Quantitative analysis of tomographic data

For quantitative analysis of the tomographic data, the automatic spatial graph analysis (ASGA) software
tool (Kiewisz and Muller-Reichert, 2022) was used to measure the outer-kinetochore distance, the
number of MTs per k-fiber, and the distribution and morphology of the MT minus ends. Quantitative
analyses were carried out essentially as described (Kiewisz et al., 2021).

Outer-kinetochore distance

The inter-kinetochore distance was used as a readout of the mitotic stage of the selected cells (Maresca
and Salmon, 2009). To measure the inter-kinetochore distance, the neighboring sister kinetochores were
identified in the 3D models. The center of each kinetochore was defined as the median position of all
KMT plus ends associated with each selected outer kinetochore, and the inter-kinetochore distance was
then calculated as the 3DEuclidean distance between the defined median centers of each kinetochore
pair. The inter-kinetochore distances were determined for control, siScramble and siMCRS1 cells, and
values were compared using a linear regression model.

Number of KMTs per k-fiber

461   MTs associated with the kinetochores were defined as KMTs. Accordingly, the average number of KMTs
462   was determined for each experimental condition. Analysis of the number of KMTs per k-fiber was done
463   using a Generalized Linear Model (GLM) in R, applying a Poisson likelihood and a log link function. The
464   Poisson distribution is generally used to model discrete positive outcomes such as the number of
465   microtubules, when the numbers are small and cannot be approximated by a normal distribution ($n < 20$).

Distribution of MT minus ends

The polarity of KMTs was assigned as follows. The end of each KMT associated with a kinetochore was
assumed to be the plus end and the other end the minus end. To analyze the position of the KMT minus
ends in the metaphase spindles, the 3D Euclidean distance of each KMT end to both spindle poles (i.e.,
to the center of the mother centriole of the respective pole) was determined. Then, the KMT minus-end
was determined as the end closest to one of the spindle poles. In addition, the relative position of KMT
minus ends along the pole-to-pole axis was calculated. The relative position of each minus end is given
as the normalized position between the kinetochore (position = 1) and the mother centriole (position = 0)
along the pole-to-pole axis. The distribution of relative positions of KMT minus ends (mean, ± STD) is as
an average density distribution for each condition. For non-KMTs, the end closer to the nearest centriole
pair was defined as the minus end. The relative distance for the non-KMTs was calculated between the
average kinetochore (position = 1) and the mother centriole (position = 0).

Morphology of MT ends

MT ends were annotated by manual segmentation. MT ends were manually classified as open, closed or
undefined by two different observers using the end classifier in the ZIB Amira software package (Detlev et
al., 2005). End annotation was blind as the MT end morphology was only correlated to MT identity (either
KMT or non-KMT). The polarity of the MT ends (either plus or minus) and location (relative distance)
within the spindle was determined after ending the classification.

Analysis of the MT end morphology was modeled as a binary outcome (open *vs.* closed), such that the
number of open ends was naturally drawn from a binomial distribution depending on the true unobserved
proportion of open ends. To model the dependency of the proportion of open ends on the distance to the
spindle pole, we discretized the data into 16 different intervals and counted the number of open ends
called by each observer. We used a third order b-splines on the underlying logit transformation of the
proportion of open ends to jointly infer the proportion of open ends at each possible distance from the
pole. We then added the observer as a random effect to consider the differences in the classification
made by the two independent observers.

Error analysis of tomographic data

Errors in automatic MT segmentation and in the stitching of serial tomograms have been discussed previously (Kiewisz et al., 2021; Lindow et al., 2021; Redemann et al., 2014). As for the segmentation of MTs, the error associated for our approach is in the range of 5-10% (Weber et al. 2012). Each individual MT in our reconstruction has been checked manually for the correct tracing of both ends.

In previous publications (Lindow et al., 2021; Redemann et al., 2017; Weber et al., 2014), we estimated the overall quality of the MT stitching by analyzing the distribution of MT endpoints in the Z-direction (i.e., normal to the plane of the slice). We expect to find approximately the same density of MT endpoints along the Z-direction of each serial-section tomogram. Therefore, if the density of endpoints after matching is approximately the same along the Z-direction of the serial-section tomograms, we can assume that the number of artificial points that have been introduced at the interfaces of the serial sections are negligible (Kiewisz et al., 2021).

504 Data availability

Tomographic data has been uploaded to the TU Dresden Open Access Repository and Archive system
(OpARA) and is available as open access:

http://doi.org/10.25532/OPARA-XX

The code used to perform quantitative analysis of MT organization in spindles has been uploaded to the
GitHub   repository   and   is   available   as   open   access   under   the   GPL   v3.0   license:
https://github.com/RRobert92/ASGA; https://github.com/RRobert92/ASGA_3DViewer

Data and code for statistical analysis of MT ends can be accessed at

https://bitbucket.org/cmartiga/kfibers/src/master/

Acknowledgments

The authors would like to thank support form the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 675737 (DiviDE ITN network) to AL, RK, IV and and TMR. Research in the Müller-Reichert laboratory is supported by funds from the Deutsche Forschungsgemeinschaft (MU 1423/8-2). Work in the Vernos lab was supported by the Spanish Ministry of Economy (MINECO) I+D grant BFU2012-37163 and BFU2015-68726-P. A.L.D also received an EMBO short-term fellowship to visit the Müller-Reichert lab, grant agreement No. 8704. We thank Dr. Tobias Fürstenhaupt (Electron Microscopy Facility at the MPI-CBG, Dresden, Germany) for technical support. We thank the Vernos and Müller-Reichert groups and the members of the DiviDE ITN for discussions. We acknowledge the Spanish Ministry of Economy, Industry and Competitiveness (MEIC) to the EMBL partnership and support of the Spanish Ministry of Economy and Competitiveness, 'Centro de Excelencia Severo Ochoa' as well as support of the CERCA Programme / Generalitat de Catalunya.

Declaration of interests

The authors declare no competing financial interests.

References

Akhmanova, A., and M.O. Steinmetz. 2019. Microtubule minus-end regulation at a glance. *J Cell Sci*. 132.

Barisic, M., G. Rajendraprasad, and Y. Steblyanko. 2021. The metaphase spindle at steady state - Mechanism and functions of microtubule poleward flux. *Semin Cell Dev Biol*. 117:99-117.

Cheeseman, I.M., and A. Desai. 2008. Molecular architecture of the kinetochore-microtubule interface. *Nat Rev Mol Cell Biol*. 9:33-46.

Chen, X., L.A. Widmer, M.M. Stangier, M.O. Steinmetz, J. Stelling, and Y. Barral. 2019. Remote control of microtubule plus-end dynamics and function from the minus-end. *Elife*. 8.

Chretien, D., S.D. Fuller, and E. Karsenti. 1995. Structure of growing microtubule ends: two-dimensional sheets close into tubes at variable rates. *J Cell Biol*. 129:1311-1328.

Consolati, T., J. Locke, J. Roostalu, Z.A. Chen, J. Gannon, J. Asthana, W.M. Lim, F. Martino, M.A. Cvetkovic, J. Rappsilber, A. Costa, and T. Surrey. 2020. Microtubule Nucleation Properties of Single Human gammaTuRCs Explained by Their Cryo-EM Structure. *Dev Cell*. 53:603-617 e608.

DeLuca, J.G., W.E. Gall, C. Ciferri, D. Cimini, A. Musacchio, and E.D. Salmon. 2006. Kinetochore microtubule dynamics and attachment stability are regulated by Hec1. *Cell*. 127:969-982.

Detlev, S., M. Westerhoff, and H.-C. Hege. 2005. Amira: A Highly Interactive System for Visual Data Analysis *"The visualization handbook."*.

Drutovic, D., X. Duan, R. Li, P. Kalab, and P. Solc. 2020. RanGTP and importin beta regulate meiosis I spindle assembly and function in mouse oocytes. *EMBO J*. 39:e101689.

Dudka, D., C. Castrogiovanni, N. Liaudet, H. Vassal, and P. Meraldi. 2019. Spindle-Length-Dependent HURP Localization Allows Centrosomes to Control Kinetochore-Fiber Plus-End Dynamics. *Curr Biol*. 29:3563-3578 e3566.

Elting, M.W., C.L. Hueschen, D.B. Udy, and S. Dumont. 2014. Force on spindle microtubule minus ends moves chromosomes. *J Cell Biol*. 206:245-256.

Ems-McClung, S.C., and C.E. Walczak. 2010. Kinesin-13s in mitosis: Key players in the spatial and temporal organization of spindle microtubules. *Semin Cell Dev Biol*. 21:276-282.

Foley, E.A., and T.M. Kapoor. 2013. Microtubule attachment and spindle assembly checkpoint signalling at the kinetochore. *Nat Rev Mol Cell Biol*. 14:25-37.

Ganem, N.J., and D.A. Compton. 2004. The KinI kinesin Kif2a is required for bipolar spindle assembly through a functional relationship with MCAK. *J Cell Biol*. 166:473-478.

Ganem, N.J., and D.A. Compton. 2006. Functional roles of poleward microtubule flux during mitosis. *Cell Cycle*. 5:481-485.

Ganem, N.J., K. Upton, and D.A. Compton. 2005. Efficient mitosis in human cells lacking poleward microtubule flux. *Curr Biol*. 15:1827-1832.

Hoog, J.L., S.M. Huisman, Z. Sebo-Lemke, L. Sandblad, J.R. McIntosh, C. Antony, and D. Brunner. 2011. Electron tomography reveals a flared morphology on growing microtubule ends. *J Cell Sci*. 124:693-698.

Jiang, K., L. Rezabkova, S. Hua, Q. Liu, G. Capitani, A.F.M. Altelaar, A.J.R. Heck, R.A. Kammerer, M.O.
Steinmetz, and A. Akhmanova. 2017. Microtubule minus-end regulation at spindle poles by an
ASPM-katanin complex. *Nat Cell Biol*. 19:480-492.

Kiewisz, R., T. Muller-Reichert, and G. Fabig. 2021. High-throughput screening of mitotic mammalian
cells for electron microscopy using classic histological dyes. *Methods Cell Biol*. 162:151-170.

Kiewisz   R,   Müller-Reichert   T.   2022.   Automatic   Spatial-Graph   Analysis   (ASGA).
doi:https://doi.org/10.5281/zenodo.3732108

Kirschner, M.W., and T. Mitchison. 1986. Microtubule dynamics. *Nature*. 324:621.

Lampson, M.A., and E.L. Grishchuk. 2017. Mechanisms to Avoid and Correct Erroneous Kinetochore-
Microtubule Attachments. *Biology (Basel)*. 6.

Lindow, N., F.N. Brunig, V.J. Dercksen, G. Fabig, R. Kiewisz, S. Redemann, T. Muller-Reichert, S.
Prohaska, and D. Baum. 2021. Semi-automatic stitching of filamentous structures in image stacks
from serial-section electron tomography. *J Microsc*. 284:25-44.

Lüders, J. 2016. The Microtubule Cytoskeleton: Organisation, Function and Role in Disease. The
Microtubule Cytoskeleton: Organisation, Function and Role in Disease. Springer.

Maiato, H., J. DeLuca, E.D. Salmon, and W.C. Earnshaw. 2004. The dynamic kinetochore-microtubule
interface. *J Cell Sci*. 117:5461-5477.

Mandelkow, E.M., E. Mandelkow, and R.A. Milligan. 1991. Microtubule dynamics and microtubule caps: a
time-resolved cryo-electron microscopy study. *J Cell Biol*. 114:977-991.

Maresca, T.J., and E.D. Salmon. 2009. Intrakinetochore stretch is associated with changes in kinetochore
phosphorylation and spindle assembly checkpoint activity. *J Cell Biol*. 184:373-381.

Mastronarde, D.N. 1997. Dual-axis tomography: an approach with alignment methods that preserve
resolution. *J Struct Biol*. 120:343-352.

Mastronarde, D.N. 2003. SerialEM: A Program for Automated Tilt Series Acquisition on Tecnai
Microscopes Using

Prediction of Specimen Position. *Micros Microanal*. 9:1182-| 1183.

Mastronarde, D.N. 2005. Automated electron microscope tomography using robust prediction of
specimen movements. *J Struct Biol*. 152:36-51.

Mastronarde, D.N., and S.R. Held. 2017. Automated tilt series alignment and tomographic reconstruction
in IMOD. *J Struct Biol*. 197:102-113.

McEwen, B.F., and M. Marko. 1999. Three-dimensional transmission electron microscopy and its
application to mitosis research. *Methods Cell Biol*. 61:81-111.

McIntosh, J.R., E.L. Grishchuk, M.K. Morphew, A.K. Efremov, K. Zhudenkov, V.A. Volkov, I.M.
Cheeseman, A. Desai, D.N. Mastronarde, and F.I. Ataullakhanov. 2008. Fibrils connect
microtubule tips with kinetochores: a mechanism to couple tubulin dynamics to chromosome
motion. *Cell*. 135:322-333.

Meunier, S., M. Shvedunova, N. Van Nguyen, L. Avila, I. Vernos, and A. Akhtar. 2015. An epigenetic
regulator emerges as microtubule minus-end binding and stabilizing factor in mitosis. *Nat*
Commun. 6:7889.

Meunier, S., and I. Vernos. 2011. K-fibre minus ends are stabilized by a RanGTP-dependent mechanism
essential for functional spindle assembly. *Nat Cell Biol.* 13:1406-1414.

Meunier, S., and I. Vernos. 2012. Microtubule assembly during mitosis - from distinct origins to distinct
functions? *J Cell Sci.* 125:2805-2814.

Meunier, S., and I. Vernos. 2016. Acentrosomal Microtubule Assembly in Mitosis: The Where, When, and
How. *Trends Cell Biol.* 26:80-87.

Mimori-Kiyosue, Y., and S. Tsukita. 2003. "Search-and-capture" of microtubules through plus-end-binding
proteins (+TIPs). *J Biochem.* 134:321-326.

Mitchison, T., and M. Kirschner. 1984. Dynamic instability of microtubule growth. *Nature.* 312:237-242.

Mitchison, T.J. 1989. Polewards microtubule flux in the mitotic spindle: evidence from photoactivation of
fluorescence. *J Cell Biol.* 109:637-652.

Monda, J.K., and I.M. Cheeseman. 2018. The kinetochore-microtubule interface at a glance. *J Cell Sci.*
131.

Muller-Reichert, T., D. Chretien, F. Severin, and A.A. Hyman. 1998. Structural changes at microtubule
ends accompanying GTP hydrolysis: information from a slowly hydrolyzable analogue of GTP,
guanylyl (alpha,beta)methylenediphosphonate. *Proc Natl Acad Sci U S A.* 95:3661-3666.

Muller-Reichert, T., H. Hohenberg, E.T. O'Toole, and K. McDonald. 2003. Cryoimmobilization and three-
dimensional visualization of C. elegans ultrastructure. *J Microsc.* 212:71-80.

O'Toole, E., M. Morphew, and J.R. McIntosh. 2020. Electron tomography reveals aspects of spindle
structure important for mechanical stability at metaphase. *Mol Biol Cell.* 31:184-195.

O'Toole, E.T., K.L. McDonald, J. Mantler, J.R. McIntosh, A.A. Hyman, and T. Muller-Reichert. 2003.
Morphologically distinct microtubule ends in the mitotic centrosome of Caenorhabditis elegans. *J*
Cell Biol. 163:451-456.

Redemann, S., J. Baumgart, N. Lindow, M. Shelley, E. Nazockdast, A. Kratz, S. Prohaska, J. Brugues, S.
Furthauer, and T. Muller-Reichert. 2017. C. elegans chromosomes connect to centrosomes by
anchoring into the spindle network. *Nat Commun.* 8:15288.

Redemann, S., B. Weber, M. Moller, J.M. Verbavatz, A.A. Hyman, D. Baum, S. Prohaska, and T. Muller-
Reichert. 2014. The segmentation of microtubules in electron tomograms using Amira. *Methods*
Mol Biol. 1136:261-278.

Rice, L.M. 2018. A new look for the growing microtubule end? *J Cell Biol.* 217:2609-2611.

Simon, J.R., and E.D. Salmon. 1990. The structure of microtubule ends during the elongation and
shortening phases of dynamic instability examined by negative-stain electron microscopy. *J Cell*
Sci. 96 (Pt 4):571-582.

Steblyanko, Y., G. Rajendraprasad, M. Osswald, S. Eibes, A. Jacome, S. Geley, A.J. Pereira, H. Maiato,
and M. Barisic. 2020. Microtubule poleward flux in human cells is driven by the coordinated action
of four kinesins. *EMBO J.* 39:e105432.

Teixido-Travesa, N., J. Roig, and J. Luders. 2012. The where, when and how of microtubule nucleation -
one ring to rule them all. *J Cell Sci*. 125:4445-4456.

VandenBeldt, K.J., R.M. Barnard, P.J. Hergert, X. Meng, H. Maiato, and B.F. McEwen. 2006.
Kinetochores use a novel mechanism for coordinating the dynamics of individual microtubules.
Curr Biol. 16:1217-1223.

Waters, J.C., T.J. Mitchison, C.L. Rieder, and E.D. Salmon. 1996. The kinetochore microtubule minus-
end disassembly associated with poleward flux produces a force that can do work. *Mol Biol Cell*.
7:1547-1558.

Weber, B., G. Greenan, S. Prohaska, D. Baum, H.C. Hege, T. Muller-Reichert, A.A. Hyman, and J.M.
Verbavatz. 2012. Automated tracing of microtubules in electron tomograms of plastic embedded
samples of Caenorhabditis elegans embryos. *J Struct Biol*. 178:129-138.

Weber, B., E.M. Tranfield, J.L. Hoog, D. Baum, C. Antony, T. Hyman, J.M. Verbavatz, and S. Prohaska.
2014. Automated stitching of microtubule centerlines across serial electron tomograms. *PLoS
One*. 9:e113222.

Wieczorek, M., L. Urnavicius, S.C. Ti, K.R. Molloy, B.T. Chait, and T.M. Kapoor. 2020. Asymmetric
Molecular Architecture of the Human gamma-Tubulin Ring Complex. *Cell*. 180:165-175 e116.

Zheng, Y., M.L. Wong, B. Alberts, and T. Mitchison. 1995. Nucleation of microtubule assembly by a
gamma-tubulin-containing ring complex. *Nature*. 378:578-583.

Zimmermann, F., M. Serna, A. Ezquerra, R. Fernandez-Leiro, O. Llorca, and J. Luders. 2020. Assembly
of the asymmetric human gamma-tubulin ring complex by RUVBL1-RUVBL2 AAA ATPase. *Sci
Adv*. 6.

Figures

Figure 1. Quantitative analysis of MT minus-end morphologies

(**A**) Illustration of a MT (arrow) associated with a chromosome as visualized in a HeLa cell at metaphase.
In this study, such an MT is defined as a KMT (top view, tomographic slice; bottom view, tomographic
slice with overlay). Scale bar, 100 nm. (**B**) Visualization of a closed and an open MT end morphology
(top view, tomographic slice; bottom view, tomographic slice with overlay for better illustration). MT ends
showing an electron-dense cap are classified as closed (left panels). Curved or sheet-like ends are
classified as open (right panels). Scale bar, 25 nm. (**C**) Three-dimensional model (top view) showing
KMTs in a quarter volume reconstruction of a control spindle (Ctrl #3). KMTs are depicted as black lines
and their end morphologies are indicated as open (green circles), closed (purple circles) or undefined
(open circles). The position of the spindle pole is indicated by an arrow. Scale bar, 500 nm. (**D**) Schematic
drawing illustrating the positional analysis of both KMT and non-KMT minus ends. The position of each
MT minus end along the spindle pole axis is given as a relative distance. The relative distance of each
KMT minus end along the half-spindle axis is defined by the position between the pole [P1, 0] and the
kinetochore [K, 1]. The relative position of non-KMT minus ends is defined by the position of the pole and
the average position of all kinetochores in the spindle [aK, 1]. The absolute distance is determined by
measuring the 3D distance of each MT minus end to the nearest mother centriole (m). A KMT is indicated
by a solid purple line, a non-KMT by a dashed purple line. (**E**) Histogram showing the absolute number of
KMT minus ends (right axis) at different relative distances. The shadow around the line (purple)
represents the credible interval (CI) with a 95% probability to observe the true value within this interval.
The graph also shows the estimated percentage of open minus ends at each given relative position (left
axis). The shaded area (grey) represents the MT-spindle pole association area. This area was defined as
the region of the spindle pole with the highest concentration of KMT minus ends as observed in the
histogram. The total number of classified KMT minus ends for control spindles was n = 2071. (**F**)
Histogram showing the absolute number of non-KMT minus the MT spindle pole association area. The
total number of classified non-KMT minus-ends is n = 1516.

Figure 1-figure supplement 1. Gallery of MT end morphologies as observed by electron tomography

(**A**) Representative collection of images showing closed MT ends (top row, tomographic slices; lower row, tomographic slices with overlays). The polarity of the MTs is not assigned. Scale bar, 25 μm. (**B**) Collection of images showing open MT ends. Scale bar, 25 μm.

Figure 2. Characterization of metaphase spindles in MCRS1-silenced cells

(**A**) Western Blot analysis showing the levels of MCRS1 in control and MCRS1-silenced cells. Tubulin
levels were used as loading control. (**B**) Immunofluorescence staining of siScramble (top panel) and
MCRS1-silenced cells (lower panel). MCRS1 is shown in green, DNA in blue and tubulin in magenta.
MCRS1 staining can be observed as tiny spots at spindle poles in the control cell. Upon silencing,
MCRS1 cannot be detected anymore. Scale bars, 5 µm. (**C**) Black/white images of a control (top, left) and
a silenced spindle (bottom, left). The magenta lines show the angles that were measured for the half
spindle (top, right) and for the MTs in the outward position of the spindle (bottom, right). Scale bars, 5 µm.
(**D**) Models of spindle shape in siScramble and siMCRS1 cells after screening for electron microscopy.
MT density is shown in gray, chromosomes in blue. Scale bar, 1 µm. (**E**) Boxplot showing the
quantification of the half-spindle angle obtained from immunofluorescence images. The boxes show the
upper and lower quartiles, the whiskers show the minimum and maximal values excluding outliers; the
line inside the box indicates the median. Asterisks show significant difference according to a linear
regression model considering two different experiments with a p-value = 4.729×10^{-8} (siScramble: n =
105; siMCRS1: n = 110). (**F**) Scatterplot showing the quantification of the MT angle from electron
microscopic images. The line shows the mean, and the error bars represent ± SD. Asterisks show
significant difference according to a two-tailed student t-test with a p-value = 0.017. (siScramble: n = 4;
siMCRS1: n = 8). (**G**) Boxplot showing the quantification of the half-spindle angle obtained from
immunofluorescence images. Asterisks show significant difference according to a linear regression model
considering two different experiments with a p-value = 7.486×10^{-26} (siScramble: n = 201; siMCRS1: n =
161). (**H**) Scatterplot showing the quantification of the MT angle obtained from electron microscopic
images. to a two-tailed student t-test with a p-value = 0.002 (siScramble: n = 8; siMCRS1: n = 16).

Figure 3. Shape of k-fibers in siScramble and MCRS1-silenced spindles

(**A**) Un-tilted top views of quarter spindles in a siScramble (left, Scramble spindle #2) and a siMCRS1 cell (right, MCRS1 spindle #5) showing k-fibers as black lines. The spindle poles are indicated as green circles (the position of the spindle poles is given by the position of the centrioles, a green circle at the bottom). Scale bar, 1µm. (**B**) Line-plot representing the Gaussian kernel density distribution of k-fiber minus ends at different relative distances from the mother centriole (position = 0) to the kinetochores

(position = 1). The shaded area indicates the MT-centrosome association area. The shaded area
represents the MT-spindle pole association area with the MT spindle pole association area. (**C**) Boxplot
representing the number of KMTs per k-fiber in the three conditions. The boxes show the upper and lower
quartiles, the whiskers show the minimum and maximal values excluding outliers; the line inside the box
indicates the median. Asterisks show significant differences according to a generalized linear model with
a p-value = 2×10^{-4}. N.s stands for non-significant with a p-value = 0.599 (Control: n = 226, siScramble: n
= 44, siMCRS1: n = 106).

Figure 3-figure supplement 1. Comparison of the characteristics of a full spindle *versus* its quarters

(**A**) Perspective view of a full 3D model showing the KMTs in a control spindle (Ctrl #1). KMTs are shown as black lines, chromosomes in gray. (**B**) Perspective view of a 3D model showing the KMTs in a quarter spindle (Ctrl #1). (**C**) Scatterplots showing the number of KMTs per k-fiber for a full spindle (Ctrl #1) and the four quarters of the same spindle. The line shows the mean, and the error bars represent ± SD. The mean numbers of KMTs for full spindle 8.04 ± 1.86 (mean ± SD, n = 93); quarter #1, 7.84 ± 2.15, n = 25; #2, 8.03 ± 2.20, n = 29; #3, 7.24 ± 2.21, n = 29; and #4, 8.19 ± 2.15, n = 32). There are no significant differences among the mean values for the full spindle and the quarter spindles after ANOVA analysis (p-value = 0.3791). (**D**) Scatterplot of the outer-kinetochore distance in the full spindle *versus* the spindle quarters. The line shows the mean, and the error bars represent ± SD. The mean outer-kinetochore distances for full spindle, 1.07 µm ± 0.20, n = 43; quarter #1, 1.06 µm ± 0.20, n = 21; #2, 1.05 µm ± 0.23, n = 24; #3, 1.06 µm ± 0.24, n = 26; and #4, 1.03 µm ± 0.18, n = 28. There are no significant differences among the mean values for the full spindle and the quarter spindles after ANOVA analysis (p-value = 0.9192).

Figure 3-figure supplement 2. Comparative analysis of outer-kinetochore distances in 3D models of control, siScramble and siMCRS1-depleted cells.

Scatter plots of the inter-kinetochore distances as measured for each of the 3D-reconstructed spindles. The mean inter-kinetochore distance in control cells is 1.14 µm (n = 114), in siScramble cells is 1.05 µm (n = 44) and in siMCRS1 0.99 µm (n = 98). A linear regression model did not reveal significant differences between siScramble versus Control p-value = 0.108 and siScramble versus siMCRS1 = 0.240. The numbers for each cell are matching the data sets in **Table 1**. Cells #1-2 correspond to full spindle reconstructions and cells #3-10 to quarter spindles.

Figure 4. MT minus-end morphology in MCRS1-silenced cells

(**A**) Estimated percentage of open KMT minus ends plotted against the relative distance on the half-
spindle length (position 0 = mother centriole, position 1 = kinetochores) in control (purple line; n = 1707)
and siMCRS1 cells (green line; n = 541). The shadow around the lines (purple and green) represents the
credible interval (CI) with a 95% probability to observe the true value within this interval. The shaded area
(grey) represents the MT-spindle pole association area. (**B**) Posterior distribution of the comparison of the
log2(open/closed) of siMCRS1 *versus* control (as shown in (**A**)) in KMTs (green line). The shadow around
the line (green) represents the 95% credible interval (CI). (**C**) Estimated percentage of open non-KMT
minus ends plotted against the relative distance on the half-spindle length in control (purple line, n= 1084)
and siMCRS1 cells (green line, n = 3115). The shadow around the line (green) represents the 95%
credible interval (CI). (**D**) Posterior distribution of the comparison of the log2(open/closed) of siMCRS1
versus control (as shown in **C**) in non-KMTs (green line). The shadow around the line (green) represents
the 95% credible interval (CI). (**E**) Three-dimensional model (top view) showing KMTs in a quarter
reconstruction of a siMCRS1 cell (siMCRS1 #5). KMTs are depicted as black lines and their end
morphologies are indicated as open (green circles), closed (purple circles) or undefined (open circles).
The position of the spindle pole is indicated by an arrow. Scale bar, 500 nm. (**F**) Bar representation of the
estimated percentage of open KMTs (Control: 60.21%, = 2064; siMCRS1: 72.66%, n = 848) and non-
KMT minus ends (Control: 64.30%, siMCRS1: 74.51%, n = 893; siMCRS1 = 4119) in control (purple) and
siMCRS1 cells (green) in the spindle pole region (d< 0.2).

Figure 5. Proposed model of KMT minus-end dynamics regulation in mammalian metaphase.
(**A**) Left panel illustrates the regulation of KMT minus-end dynamics in metaphase of control cells. (**B**)
Right panel illustrates minus-end dynamics in response to MCRS1 silencing.

Movie supplement

Figure 1. Movie supplement 1. Generation of a 3D model from joined serial electron tomograms displaying spindle control #3

Series of stitched tomograms and corresponding 3D model of spindle CTRL #3. The stacking of serial tomograms to increase the tomographic volume is illustrated. KMTs are shown as red lines.

Figure 1. Movie supplement 2. 3D model of KMTs end morphology from siMCRS1 #5

3D model of KMTs with end morphology indicated. KMTs ends were labeled as green (open end), purple (close end) and white (undefined end) circles. The KMTs are indicated as black lines.

Figure 2. Movie supplement 1. 3D model of a low-resolution spindle used for angle estimation from siScramble #2

Series of stitched screening EM micrographs of spindle siScramble #2. The stacking of serial tomograms to increase the tomographic volume is illustrated. Chromosomes are shown as blue, and the spindle is indicated with transparent gray area.

Figure 2. Movie supplement 2. 3D model of a low-resolution spindle used for angle estimation from siMCRS1 #5

Series of stitched screening EM micrographs of spindle siMCRS1#5. The stacking of serial tomograms to increase the tomographic volume is illustrated. Chromosomes are shown as blue, and the spindle is indicated with transparent gray area.

Figure 4. Movie supplement 1. 3D model of KMTs end morphology from siScramble #2

3D model of KMTs with end morphology indicated. KMTs ends were labeled as green (open end), purple (close end) and white (undefined end) spheres. The KMTs were indicated as black lines.

Figure 4. Movie supplement 2. 3D model of KMTs end morphology from siMCRS1 #5

3D model of KMTs with end morphology indicated. KMTs ends were labeled as green (open end), purple (close end) and white (undefined end) spheres. The KMTs were indicated as black lines.

 Tables

 Table 1. Tomographic data sets as used throughout this study

Metadata/ data set	Original data set	Spindle reconstruction	No. of serial sections[1]/ Z-factor	Estimated tomographic volume (μm^3)	Data size (Gb)
Ctrl #1 (Kiewisz et. al., 2021)	T_0475	Full (divided into quarters)	22 / 1.30	144	44.3
Ctrl #2 (Kiewisz et. al., 2021)	T_0479	Full (divided into quarters)	29 / 1.40	263	74.2
Ctrl #3	T_0550_wt_1	Quarter	17 / 1.30	114	30
siScr #1	T_0506	Quarter	12 / 1.25	58	22.4
siScr #2	T_0550_ctrl_3	Quarter	14 / 1.30	78	28.3
siMCRS1 #1	T_0506_mcrs1_2	Quarter	18 / 1.32	94	26.8
siMCRS1 #2	T_0506_mcrs1_4	Quarter	18 / 1.35	87	24.9
siMCRS1 #3	T_0506_mcrs1_5	Quarter	17 / 1.30	79	22.8
siMCRS1 #4	T_0550_mcrs1_2	Quarter	16 / 1.35	112	32.1
siMCRS1 #5	T_0550_mcrs1_3	Quarter	21 / 1.36	110	31.6

815 [1] Average section thickness is 300 nm.
816

817 **Table 2. Quantitative analysis of the tomographic reconstructions as used**

818 **throughout this study**

Data set	No. of MTs	No. of KMTs	No. of KMTs in the volume	No. of k-fibers	Pole-to-pole distance (µm)	Outer-kinetochore distance (µm)
Ctrl #1	4884	797	748	92	7.16	1.08 ± 0.20 (n = 43)
Ctrl #2	8047	1102	1072	110	10.39	1.24 ± 0.21 (n = 50)
Ctrl #3	-	407	205	22	-	1.15 ± 0.22 (n = 22)
siScr #1	2913	397	203	20	-	1.02 ± 0.20 (n = 17)
siScr #2	6221	344	205	24	-	1.06 ± 0.21 (n = 28)
siMCRS1 #1	2884	224	50	8	-	1.10 ± 0.24 (n = 20)
siMCRS1 #2	2451	169	57	10	-	1.12 ± 0.23 (n = 10)
siMCRS1 #3	3635	301	159	18	-	1.06 ± 0.11 (n = 6)
siMCRS1 #4	5624	348	180	29	-	1.02 ± 0.18 (n = 28)
siMCRS1 #5	3811	484	273	39	-	1.13 ± 0.20 (n = 35)

820 **Table 3. Quantitative analysis of MT minus-end morphologies**

End morphology	KMT minus ends				KMT plus ends			
	Total no.	Matching open ends	Matching closed ends	Undefined ends	Total no.	Matching open ends	Matching closed ends	Undefined ends
Control (Ctrl #1, #2, #3)	2239	682	385	532	1841	1509	6	218
siScramble (siScr #1, #2)	407	109	66	137	432	350	2	47
siMCRS1 (siMCRS1 #1, #2, #3, #4, #5)	726	297	90	185	847	692	2	108
End morphology	**non-KMT minus ends**							
	Total no. of ends	Matching open ends	Matching closed ends	Undefined ends				
Control (Ctrl #1, #2, #3)	1511	521	209	427				
siScramble (siScr #1, #2)	1155	278	59	591				
siMCRS1 (siMCRS1 #1, #2, #3, #4, #5)	6654	1876	509	3539				