**Title**

Composition of a low erucic acid, low fiber field pennycress (*Thlaspi arvense* L) grain referred

to as CoverCress[TM] developed through breeding and gene editing

**Author List**

Gary Hartnell[1*,] Shawna Lemke[2], and Chris Aulbach[3]

*Corresponding author (email): gary.hartnell@gmail.com

**Author Affiliation**

[1]President/Owner Hartnell International Consulting, LLC (St. Peters, MO)

[2]Principal/Owner SLL Consulting & Services, LLC (St. Louis, MO)

[3]Vice President Agronomy, CoverCress Inc. (St. Louis, MO)

**Abstract**

Field pennycress (*Thlaspi arvense* L.) can be domesticated and cultivated as an annual in a corn /

pennycress / soybean rotation where pennycress is sown and harvested as a cash cover crop. To

improve the nutritional profile, pennycress was modified in two ways to achieve the same

alterations in characteristics: 1) through selection of mutants and 2) through gene editing. These

alterations resulted in a low erucic acid, lower fiber phenotype and the resulting products from

these combinations are referred to as CoverCress CCWG and CoverCress B3WG, respectively.

CCWG and B3WG were planted as cover crops in five U.S. locations in the fall and the grain

was harvested in the subsequent June. The grain was treated with 83 mM copper sulfate solution

as a potential palatability agent for the naturally high glucosinolate levels, and was subsequently

analyzed for nutrient (proximates, minerals, fatty acids, amino acids, and vitamins) and anti-

nutrient (sinapine, glucosinolates, mold) content. The low erucic acid, lower fiber phenotype

was consistently achieved across five lots. Generally, the nutrient content for both CCWG and

B3WG were similar to canola grain. Canola grain contains the anti-nutrient sinapine but is

significantly reduced to below the level of detection in CoverCress grain.  As expected,

CoverCress grain contains about 10 times more glucosinolates than canola.  Based on the

composition of CoverCress grain, it may provide a source of energy and amino acids to animals

with restricted inclusion based on the glucosinolate content.

**Keywords**

Pennycress, CoverCress, low-erucic acid, cover crop, gene editing

**Introduction**

Field pennycress (*Thlaspi arvense L.*) is an annual weed that grows over the winter in

Eurasia and North America. It is a member of the Brassicacae family, commonly known as the

mustard family. Field pennycress contains high levels of oil (~25-30%) that makes it a desirable

ultra-low carbon fuel feedstock. The potential commercial value of field pennycress production

has been researched extensively in a variety of leading laboratories including the National Center

for Agricultural Innovation, USDA ARS, Peoria, IL, Western Illinois University, Illinois State

University and the University of Minnesota. The cumulative literature and experience establish

that field pennycress can be domesticated and cultivated as a winter annual in a corn / pennycress

/ soybean rotation where pennycress is sown and harvested as a cash winter cover crop as

(Phipphen and Phippen, 2012; Sedbrook *et al.*, 2014).

In addition to this primary value for fuel, the seed could provide an energy source for

animal feeds such as chicken feed, however, the oil typically contains >35% erucic acid. Erucic

acid is a 22-carbon monounsaturated acid that is absorbed, distributed and metabolized like other

fatty acids involving primarily metabolism via mitochondrial beta-oxidation and, to a lesser

extent, peroxisomal beta-oxidation. Like other longer-chain fatty acids, the rate of mitochondrial

beta oxidation is comparatively lower for erucic acid; however, elevated erucic acid levels

induce liver peroxisomal oxidation pathways as a mechanism of compensation.  Interest in the

safety of erucic acid occurred when results of studies in rats associated the dietary intake of high

doses of erucic acid with myocardial lipidosis and heart lesions (Bremer and Norum, 1982).

Oilseed rape conventionally contains similarly high levels of erucic acid. Low erucic acid

varieties were identified and marketed as canola, which have been shown to be safe to broiler

chickens when incorporated in diets up to 25% inclusion. Reduction in erucic acid is achieved

through disruption of Fatty Acid Elongation 1 (FAE1), resulting in higher levels of oleic (18:1).

It is through this same mechanism that erucic acid levels has been lowered in pennycress

(McGinn et al., 2019).

Field pennycress is also high in fiber, which can impact digestibility as a feed ingredient.

The production of seed coat fiber was first characterized in the model plant *Arabidopsis*.

*Arabidopsis* seed coats derive their brown color from the accumulation of proanthocyanidins

(PAs), a class of flavonoid chemicals (polymerized flavan-3-ols, or condensed tannins) that

protect against a variety of biotic and abiotic stresses and help maintain seed dormancy and

viability (Debeaujon *et al*., 2003). PAs start out as colorless epicatechin compounds until they

are transported to the vacuole where they are polymerized and oxidized as the seed desiccates. In

*Arabidopsis,* PAs are only produced in a narrowly defined cell layer in the endothelium of the

seed, and the genes TTG1, TT8/bHLH042, and TT2/MYB123 and have been demonstrated as

being the three main regulators of PA biosynthesis in seed coat (Baudry *et al*., 2004; Lepiniec *et*

*al*., 2006).  Gonzalez *et al*. (2009) described how the TTG1 works in a complex with a particular

combination of MYB class and bHLH class transcription factors to regulate epidermal

development of the seed coat.  Loss-of-function mutants in these genes exhibit the transparent

"testa" phenotype as a result of low levels of oxidized PAs in the seed coat.  Similarly, the

transparent testa phenotype was observed with loss-of-function mutations in orthologs of these

genes in pennycress (Chopra *et al*, 2018), resulting in reduced fiber content.  This phenomenon

has been observed in other brassicas such as canola and is characterized by yellow seeds that

have more oil because of the resulting thinner seed coat and larger embryo (Abraham & Bhatia,

1986). Meal from these brassicas have also been shown to be useful in animal feed because of

the relatively lower fiber and higher metabolizable energy (Slominski *et al*., 1994, Slominski,

2015;).

Reduction of erucic acid and fiber levels in pennycress has been achieved through

conventional breeding and gene editing for loss of function in the associated gene pathways. A

low-erucic acid, lower-fiber pennycress is being developed under the trade name of

CoverCress[TM] .

CoverCress[1]  represents  a clear opportunity for sustainable optimization of certain

agricultural systems.  It serves as an important winter cover, working within the no/low-till

cropping systems to prevent soil erosion from fallow fields and improve soil nitrogen

management. As part of the safety assessment of this new grain for animal feed, a

comprehensive compositional study was conducted on both the mutant breeding line and gene

edited lines.

---

[1] For simplicity, CoverCress is not marked with ™ for the remainder of the document.

**Materials and Methods**

*CoverCress Line Development*

　　　　Two studies were conducted to assess the composition of CoverCress grain. The first study utilized grain from a low-erucic acid, lower-fiber variety of CoverCress developed through breeding by identification of mutants. One parent line, referred to as MN106-V300, was developed via ethyl methanesulfonate (EMS) mutation breeding. This line contains a mutation to the FAE1 gene (specifically, a cytosine to thymine change at base position 1018bp), resulting in a premature stop codon and complete loss of function of the gene, causing a reduction in erucic acid level (Chopra *et al*., 2020). The second parent line, referred to as Y1126, was isolated from a cultivated field in Grantfork IL and was identified as a lower-fiber phenotype with a yellow seed coat. This line contains a naturally-occurring deletion of 21bp in the TRANSPARENT TESTA GLABRA 1 (TTG1) gene. This mutation results in a deletion of 7 amino acids in the conserved area of TTG1 protein, leading to a complete loss of function.  The MN106-V300 and Y1126 lines were crossed in fall 2017, and F2 plants carrying homozygous mutant alleles for these genes were further propagated in bulk, first in local greenhouses in the spring of 2018 and then in the field in an 0.3-acre increase planted at Sigel, IL in the fall of 2018. Seeds from this increase were harvested in bulk in the spring of 2019, and this seed formed the foundation source. It should be noted that this source is segregating for all background genetics that differ between Y1126 and MN106-V300 and fixed only for FAE1 and TTG1. This line is referred to as CoverCress Whole Grain (CCWG).

　　　　The second study used a line that was generated through application of gene editing techniques to obtain lines of low-erucic acid, lower-fiber pennycress and is referred to as B3WG. To generate this line, mutations were introduced into pennycress cultivar B3 using a CRISPR/SpCas9 DNA construct designed to target genomic edits to the FAE1 and TT8 genes. This transgene construct was delivered to B3 using a disarmed Agrobacterium tumefaciens strain (GV3101) and a standard floral dip transformation method. Presence of the edits in T1 plants was confirmed through multiple methods including confirmatory PCR screening of a fragment of the T-DNA.  Seed from the progeny T2 generation was screened for segregants that did not have the transgene.

Resulting progeny in the T3 generation were screened again for negative presence of

DsRED and Cas as well as homozygous edits to TT8 and FAE1.  Three types of plants were

identified. B3 [WG].1 harbors a single G insertion in the TT8 gene and a single A insertion in the

FAE1 gene. B3 [WG].2 harbors a single base deletion in the TT8 gene and a single A insertion in

the FAE1 gene. In both iterations, B3WG plants produced yellow seed (a marker for low fiber)

and low accumulation of erucic acid in seeds. The B3WG.1 and B3WG.2 plants were taken

forward for subsequent characterization and seed bulk up.

*Grain production*

Two studies were conducted to measure the nutrient and antinutrient content of

CoverCress grain.  In Study A, CoverCress seed developed through breeding (CCWG) and

grown in five locations (Venedy, IL; Sigel, IL; Havana, IL; and two locations in Mt. Pulaski, IL)

from September 2019 and harvested in June of 2020.  In Study B, CoverCress seed was

developed through gene editing (B3WG) and grown in five locations (Havana, IL; Arenzville,

IL; Mt. Pulaski, IL: and two locations in Martinsville, IL) from September 2020 and harvested in

June of 2021.  Seeds were transported to CoverCress Inc. (St. Louis) and cleaned.

*Treatment of grain with copper sulfate*

Native pennycress and CoverCress grain contain high levels of glucosinolates in the form

of aliphatic sinigrin (approximately 35-110 µ moles/g meal). Glucosinolates are considered an

antinutrient and can prevent the absorption of iodine and may affect thyroid function in target

species (Schöne et al., 1993). Glucosinolates can also impart a bitter flavor which may affect

feed consumption (Bischoff, 2021). Copper sulfate treatment has been suggested to overcome the

deleterious effects of glucosinolates for other high glucosinolate grains (Schöne *et al*., 1993;

Singhal & Sinha 2000; Payvastagan, 2013). It was theorized that copper may reduce or mask the

bitter taste of glucosinolates, improving palatability. Therefore, both the CCWG and B3WG

grains were pre-treated with copper sulfate prior to characterization of the grain composition and

use in subsequent broiler feeding studies.

Three hundred grams of CCWG grain from each of the five locations was blended with

45 ml of 83.3 mM copper sulfate pentahydrate solution.  The samples were labelled CCWG-15-

32-A, CCWG-15-32-B, CCWG-15-32-C, CCWG-15-32-D, and CCWG-15-32-E. Four hundred

and sixty grams of CCWG and B3WG grain from each of the five locations were blended with

60 ml of 83.3 mM copper sulfate pentahydrate solution. The samples were labelled, B3WG-15-

32-A, B3WG-15-32-B, B3WG-15-32-C, B3WG-15-32-D, and B3WG-15-32-E.  Samples were

stored at 2 to 8ºC.

***Methods of Analysis***

169   The five lots of CCWWG-15-32 and B3WG-15-32 were analyzed duplicate by EPL Bio

Analytical Services (Niantic, IL) following Good Laboratory Practices (GLP) and by Dairyland

Laboratories (Arcadia, WI) a commercial laboratory that uses methods on the latest research

findings and technical methodologies available to the feed industry. Dairyland Laboratories

participates (DLL) in the National Forage Testing Association, North American Proficiency

Testing Program (NAPT), and American Association of Feed Control Officials (AAFCO)

Proficiency Testing Program.  These programs help laboratories generate accurate and precise

analyses.

177   The list of analytes and the respective Standard Operating Procedure (SOP) used by each

178 lab are shown in Table 1.  The details of the specific analytical methodology used can be

179 obtained from the respective laboratories.

180 Table 1. List of analytes measured by EPL Bio Analytical Services (EPL) and Dairyland
181 Laboratories and Standard Operating Procedures (SOP).

| | EPL | Dairyland Laboratories |
|---|---|---|
| Analyte | SOP | SOP |
| Moisture/Dry Matter | GSOP-NC-4 | DLARC.SOP.FEED.WCFEED.Lab Dry Matter 105 3hr.001 |
| Crude Protein | GSOP-NC-20 | DLARC.SOP.FEED.WCFEED.Leco528.008 |
| Ash | GSOP-NC-2 | DLARC.SOP.FEED.WCFEED.Ash.003 |
| Crude Fiber | GSOP-NC-5 | DLARC.SOP.FEED.WCFEED.Crude Fiber.006 |
| Acid Detergent Fiber (ADF) | GSOP-NC-3 | DLARC.SOP.FEED.WCFEED.Sequential NDF/ADF/Lignin Analysis.000 |
| Neutral Detergent Fiber (NDF) | GSOP-NC-9 | DLARC.SOP.FEED.WCFEED.Sequential NDF/ADF/Lignin Analysis.000 |
| Total Dietary Fiber (TDF) | GSOP-NC-359 | Not measured |
| Crude Fat (Acid hydrolysis) | GSOP-NC-141 | DLARC.SOP.FEED.WCFEED.AH Fat.002 |
| Minerals (Ca, P, Mg, K, Na, S, Cu, Mn, Fe, Zn) | GSOP-NC-60 | DLARC.SOP.FEED.WCFEED.ICP Analysis.006 |
| Chloride | GSOP-NC-328 | DLARC.SOP.FEED.WCFEED.Salt.006 |
| Carbohydrates | GSOP-NC-494 | Not Measured |

| | | |
|---|---|---|
| Cysteine and Methionine | GSOP-NC-279 | DLARC.SOP.FEED.WCFEED.Cysteine and Methionine.000 |
| Tryptophan | GSOP-NC-22 | DLARC.SOP.FEED.WCFEED.Tryptophan Analysis.001 |
| 15 Amino Acids | GSOP-NC-58 | DLARC.SOP.FEED.WCFEED. Amino Acid Analysis.000 |
| Fatty Acids | GSOP-NC-319 | DLARC.SOP.FEED.WCFEED.Fatty Acid Analysis.004 |
| Vitamin B1 (Thiamine) | GSOP-NC-262 | Not Measured |
| Vitamin B2 (Riboflavin) | GSOP-NC-46 | Not Measured |
| Vitamin B6 (Pyridoxine) | GSOP-NC-220 | Not Measured |
| Folic Acid | GSOP-NC-26 | Not Measured |
| Niacin | GSOP-NC-28 | Not Measured |
| Tocopherols (Alpha, beta, gamma, delta) | GSOP-NC-341 | Not Measured |
| Fumonisins (B1, B2, B3) | GSOP-NC-336 | Not Measured |
| Aflatoxin (B1, B2, G1, G2), T-2 Toxin, Ochratoxin-A | GSOP-NC-337 | Not Measured |
| Deoxynivalenol, 3/15-acetyl-deoxynivalenol, Zearalenone | GSOP-NC-338 | Not Measured |
| Ergosine, Ergotamine, Ergoocornine, Ergocryptine, Ergocristine | GSOP-NC-456 | Not Measured |
| | | Not Measured |
| Sinapine | GSOP-NC-208 | Not Measured |
| Sinigrin | GSOP-NC-650 | Not Measured |

### *Data Handling*

The data for each study consisted of five samples each from one of five locations grown
in Illinois with duplicate values for each sample.  The duplicates were averaged to give one value
for each of the five samples.

For the data generated by EPL, Statistical Consultants Plus, LLC (Fenton, MO)
calculated measures of data dispersion (variability/standard deviation and data ranges) and
measures of central tendency (mean and median) for each analyte using the analytical data from
the five test substances each representing five different field locations using SAS, version 9.4.
For the data generated by Dairyland Laboratories, Hartnell International Consulting LLC (St.
Peters, MO) calculated measures of data dispersion (variability/standard deviation and data
ranges) and measures of central tendency (mean and median) for each analyte. Measures were
calculated using Microsoft Excel.

### Results and Discussion

### Study A – CCWG-15-32

Proximates

Table 2 contains the results of proximate analyses. In general, data were in
agreement between laboratories except for acid detergent fiber (ADF), neutral detergent
fiber (NDF) and crude fat.  The two labs used different fat extraction methods. Methodology
was also different for the analysis of ADF and NDF, however, in both cases the ADF and
NDF analyses were conducted on the residue after fat extraction. It is postulated that
Dairyland Laboratories's method extracted more fat from the product than EPL's method
and that EPL's assessment of ADF and NDF may be higher because of residual fat, which
inflated the ADF and NDF values. In addition, there was greater variation noted with EPL's
fat values than with Dairyland Labs.  The Fiber Best Practices Working Group under
AAFCO's Laboratory and Services Committee provides a detailed discussion on the critical
factors in determining fiber in feed and forages (AAFCO, 2017). As expected, mean
moisture levels were 15.6 -17.1% because of the addition of the copper sulfate solution to
the CoverCress grain.  Mean and median values are similar for all parameters, indicating
that the values from the five lots are evenly distributed, i.e. the data are symmetrical in
distribution.

216
217 Table 2. CCWG-15-32 Proximates (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Moisture | % | 17.1 | 0.23 | 17.0 | 16.9 | 17.4 |
| DLL | Moisture | % | 15.6 | 0.34 | 15.5 | 15.1 | 16.1 |
| EPL | Dry Matter | % | 82.9 | 0.23 | 83.1 | 82.7 | 83.1 |
| DLL | Dry Matter | % | 84.4 | 0.34 | 84.5 | 83.9 | 84.9 |
| EPL | Crude Fiber | % | 23.1 | 1.28 | 22.8 | 21.5 | 25.0 |
| DLL | Crude Fiber | % | 23.0 | 1.72 | 22.7 | 20.9 | 24.9 |
| EPL | Acid Detergent Fiber | % | 30.1 | 1.54 | 29.9 | 28.0 | 32.1 |
| DLL | Acid Detergent Fiber | % | 13.9 | 1.58 | 13.5 | 12.1 | 15.8 |
| EPL | Neutral Detergent Fiber | % | 29.5 | 1.55 | 29.2 | 27.7 | 32.0 |
| DLL | Neutral Detergent Fiber | % | 20.3 | 2.76 | 19.3 | 17.3 | 24.5 |
| EPL | Total Dietary Fiber | % | 30.0 | 0.97 | 29.8 | 28.9 | 31.5 |

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Crude Protein | % | 25.1 | 1.53 | 25.5 | 22.6 | 26.7 |
| DLL | Crude Protein | % | 25.5 | 1.63 | 25.6 | 22.7 | 26.7 |
| EPL | Crude Fat | % | 32.7 | 1.26 | 33.1 | 30.6 | 33.7 |
| DLL | Crude Fat | % | 36.3 | 0.52 | 36.5 | 35.5 | 36.7 |
| EPL | Carbohydrates | % | 36.9 | 1.61 | 36.9 | 34.6 | 38.5 |
| EPL | Ash | % | 5.3 | 0.39 | 5.5 | 4.9 | 5.7 |
| DLL | Ash | % | 5.8 | 0.40 | 5.9 | 5.3 | 6.2 |

[1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories

<u>Amino Acid analysis</u>

Table 3 contains the amino acid profile of CCWG-15-32 when expressed on a 100%

dry matter basis and Table 4 contains the amino acid profile of CCWG-15-32 when

expressed on a % of total protein basis. The amino acid values from the five lots showed a

symmetrical distribution with the means and median values being similar. EPL consistently

had lower variability (smaller standard deviations) than Dairyland Laboratories. Dairyland

reported numerically higher values for most amino acids whether expressed as a percent of

dry matter or as a percent of protein. Dairyland had a higher percentage of total amino acids

when expressed as total amino acids as a percent of crude protein (97% versus 89.4%).

Table 3. CCWG-15-32 Amino Acids (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Alanine | % | 1.18 | 0.074 | 1.18 | 1.08 | 1.28 |
| DLL | Alanine | % | 1.20 | 0.107 | 1.20 | 1.09 | 1.38 |
| EPL | Arginine | % | 1.53 | 0.071 | 1.55 | 1.42 | 1.60 |
| DLL | Arginine | % | 1.82 | 0.159 | 1.86 | 1.58 | 1.98 |
| EPL | Aspartic Acid | % | 2.03 | 0.245 | 1.99 | 1.74 | 2.42 |
| DLL | Aspartic Acid | % | 2.34 | 0.335 | 2.30 | 2.01 | 2.89 |
| EPL | Cystine | % | 0.41 | 0.053 | 0.44 | 0.32 | 0.45 |
| DLL | Cysteine | % | 0.41 | 0.058 | 0.42 | 0.31 | 0.47 |
| EPL | Glutamic Acid | % | 3.79 | 0.289 | 3.82 | 3.34 | 4.13 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| DLL | Glutamic Acid | % | 4.16 | 0.407 | 4.19 | 3.65 | 4.77 |
| EPL | Glycine | % | 1.56 | 0.087 | 1.57 | 1.42 | 1.66 |
| DLL | Glycine | % | 1.69 | 0.159 | 1.68 | 1.52 | 1.95 |
| EPL | Histidine | % | 0.69 | 0.026 | 0.70 | 0.64 | 0.71 |
| DLL | Histidine | % | 0.59 | 0.058 | 0.59 | 0.51 | 0.65 |
| EPL | Isoleucine | % | 1.01 | 0.050 | 1.03 | 0.94 | 1.07 |
| DLL | Isoleucine | % | 0.93 | 0.111 | 0.92 | 0.77 | 1.08 |
| EPL | Leucine | % | 1.72 | 0.125 | 1.71 | 1.54 | 1.89 |
| DLL | Leucine | % | 1.91 | 0.220 | 1.90 | 1.65 | 2.26 |
| EPL | Lysine | % | 1.28 | 0.050 | 1.29 | 1.20 | 1.33 |
| DLL | Lysine | % | 1.40 | 0.077 | 1.43 | 1.29 | 1.48 |
| EPL | Methionine | % | 0.41 | 0.026 | 0.42 | 0.36 | 0.43 |
| DLL | Methionine | % | 0.48 | 0.047 | 0.47 | 0.42 | 0.53 |
| EPL | Phenylalanine | % | 1.13 | 0.060 | 1.15 | 1.03 | 1.18 |
| DLL | Phenylalanine | % | 1.20 | 0.135 | 1.20 | 0.99 | 1.36 |
| EPL | Proline | % | 1.25 | 0.060 | 1.27 | 1.17 | 1.32 |
| DLL | Proline | % | 1.38 | 0.079 | 1.20 | 1.27 | 1.48 |
| EPL | Serine | % | 1.01 | 0.072 | 1.02 | 0.92 | 1.12 |
| DLL | Serine | % | 1.13 | 0.116 | 1.12 | 1.03 | 1.32 |
| EPL | Threonine | % | 1.14 | 0.053 | 1.15 | 1.06 | 1.21 |
| DLL | Threonine | % | 1.25 | 0.106 | 1.25 | 1.14 | 1.42 |
| EPL | Tryptophan | % | 0.35 | 0.011 | 0.35 | 0.34 | 0.37 |
| DLL | Tryptophan | % | 0.52 | 0.035 | 0.52 | 0.48 | 0.57 |
| EPL | Tyrosine | % | 0.64 | 0.026 | 0.63 | 0.61 | 0.67 |
| DLL | Tyrosine | % | 0.88 | 0.102 | 0.86 | 0.74 | 1.02 |
| EPL | Valine | % | 1.30 | 0.063 | 1.32 | 1.21 | 1.37 |
| DLL | Valine | % | 1.45 | 0.148 | 1.47 | 1.24 | 1.65 |

230 [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
231

232
233

234    Table 4. CCWG-15-32 Amino Acids (% of protein basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Alanine | % | 4.70 | 0.082 | 4.75 | 4.59 | 4.78 |
| DLL | Alanine | % | 4.73 | 0.251 | 4.68 | 4.46 | 5.11 |
| EPL | Arginine | % | 6.11 | 0.232 | 6.21 | 5.71 | 6.26 |
| DLL | Arginine | % | 7.12 | 0.241 | 7.22 | 6.79 | 7.36 |
| EPL | Aspartic Acid | % | 8.09 | 0.561 | 7.82 | 7.68 | 9.06 |
| DLL | Aspartic Acid | % | 9.15 | 0.914 | 8.89 | 8.38 | 10.73 |
| EPL | Cystine | % | 1.65 | 0.245 | 1.75 | 1.21 | 1.78 |
| DLL | Cysteine | % | 1.6 | 0.268 | 1.65 | 1.16 | 1.83 |
| EPL | Glutamic Acid | % | 15.10 | 0.292 | 15.00 | 14.75 | 15.45 |
| DLL | Glutamic Acid | % | 16.32 | 0.822 | 16.11 | 15.50 | 17.69 |
| EPL | Glycine | % | 6.20 | 0.062 | 6.21 | 6.13 | 6.29 |
| DLL | Glycine | % | 6.63 | 0.366 | 6.56 | 6.27 | 7.22 |
| EPL | Histidine | % | 2.75 | 0.113 | 2.78 | 2.56 | 2.86 |
| DLL | Histidine | % | 2.32 | 0.122 | 2.30 | 2.16 | 2.48 |
| EPL | Isoleucine | % | 4.03 | 0.077 | 4.02 | 3.93 | 4.14 |
| DLL | Isoleucine | % | 3.62 | 0.233 | 3.57 | 3.40 | 4.02 |
| EPL | Leucine | % | 6.84 | 0.149 | 6.82 | 6.72 | 7.08 |
| DLL | Leucine | % | 7.47 | 0.512 | 7.29 | 7.08 | 8.37 |
| EPL | Lysine | % | 5.10 | 0.171 | 5.07 | 4.85 | 5.28 |
| DLL | Lysine | % | 5.51 | 0.139 | 5.50 | 5.30 | 5.68 |
| EPL | Methionine | % | 1.62 | 0.014 | 1.62 | 1.61 | 1.64 |
| DLL | Methionine | % | 1.88 | 0.121 | 1.86 | 1.74 | 2.03 |
| EPL | Phenylalanine | % | 4.50 | 0.051 | 4.52 | 4.42 | 4.54 |
| DLL | Phenylalanine | % | 4.69 | 0.249 | 4.64 | 4.37 | 5.06 |
| EPL | Proline | % | 4.98 | 0.254 | 5.06 | 4.55 | 5.17 |
| DLL | Proline | % | 5.42 | 0.399 | 5.34 | 4.93 | 5.90 |
| EPL | Serine | % | 4.04 | 0.103 | 4.03 | 3.91 | 4.19 |

| | | | | Standard | | Range | |
|---|---|---|---|---|---|---|---|
| DLL | Serine | % | 4.42 | 0.313 | 4.35 | 4.05 | 4.89 |
| EPL | Threonine | % | 4.55 | 0.076 | 4.54 | 4.48 | 4.68 |
| DLL | Threonine | % | 4.91 | 0.243 | 4.85 | 4.62 | 5.26 |
| EPL | Tryptophan | % | 1.41 | 0.052 | 1.39 | 1.37 | 1.50 |
| DLL | Tryptophan | % | 2.05 | 0.066 | 2.06 | 1.96 | 2.12 |
| EPL | Tyrosine | % | 2.55 | 0.139 | 2.49 | 2.41 | 2.75 |
| DLL | Tyrosine | % | 3.43 | 0.215 | 3.34 | 3.27 | 3.80 |
| EPL | Valine | % | 5.20 | 0.102 | 5.20 | 5.08 | 5.34 |
| DLL | Valine | % | 5.69 | 0.267 | 5.61 | 5.48 | 6.14 |

235 [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
236

Fatty Acid Characterization

Tables 5 and 6 contain the fatty acid profile of CCWG-15-32 when expressed on a

100% dry matter basis and percent of total fatty acids, respectively.  In general, there was

good agreement between laboratories. The fatty acid values from the five lots showed a

symmetrical distribution with the means and median values being similar.  The major fatty

acids are oleic (40% of the fatty acids) followed by linoleic (34.5% of the fatty acids) and

linolenic (~18% of fatty acids).  These three fatty acids comprise over 90% of the fatty acids

in CCWG-15-32.  No long chain polyunsaturated long chain fatty acids such as EPA and

DHA were detected.  Results confirm that CCWG-15-32 contains negligible levels of erucic

acid. Dairyland Laboratories did not detect it and EPL reported erucic acid to be less than

0.1% of the total fatty acids.

Table 5.  CCWG-15-32 Fatty Acids (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Myristic (C14:0) | % | 0.02 | 0.001 | 0.02 | 0.02 | 0.02 |
| DLL | Myristic (C14:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Palmitic (C16:0) | % | 0.79 | 0.085 | 0.82 | 0.67 | 0.88 |
| DLL | Palmitic (C16:0) | % | 0.75 | 0.063 | 0.74 | 0.67 | 0.82 |
| EPL | Palmitoleic (C16:1) | % | 0.03 | 0.005 | 0.03 | 0.02 | 0.03 |

| Lab | Fatty Acid | Unit | | | | | |
|---|---|---|---|---|---|---|---|
| DLL | Palmitoleic (C16:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecanoic (C17:0) | % | 0.01 | 0.001 | 0.01 | 0.01 | 0.01 |
| DLL | Heptadecanoic (C17:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecenoic (C17:1) | % | 0.01 | 0.001 | 0.01 | 0.01 | 0.02 |
| DLL | Heptadecenoic (C17:1) | % | NA[2] | NA | NA | NA | NA |
| EPL | Stearic (C18:0) | % | 0.20 | 0.024 | 0.19 | 0.17 | 0.23 |
| DLL | Stearic (C18:0) | % | 0.18 | 0.020 | 0.18 | 0.16 | 0.21 |
| EPL | Oleic (C18:1) | % | 8.04 | 0.962 | 8.04 | 6.73 | 9.19 |
| DLL | Oleic (C18:1) | % | 7.45 | 0.815 | 7.35 | 6.36 | 8.28 |
| EPL | Linoleic (C18:2) | % | 6.87 | 0.605 | 7.19 | 6.08 | 7.46 |
| DLL | Linoleic (C18:2) | % | 6.40 | 0.570 | 6.35 | 5.55 | 6.93 |
| EPL | Linolenic (C18:3) | % | 3.62 | 0.241 | 3.68 | 3.34 | 3.87 |
| DLL | Linolenic (C18:3) | % | 3.07 | 0.196 | 2.98 | 2.82 | 3.27 |
| EPL | Nonadecenoic (C19:1) | % | 0.01 | 0.001 | 0.01 | 0.01 | 0.01 |
| DLL | Nonadecenoic (C19:1) | % | NA | NA | NA | NA | NA |
| EPL | Arachidic (C20:0) | % | 0.03 | 0.004 | 0.03 | 0.03 | 0.04 |
| DLL | Arachidic (C20:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Eicosenoic (C20:1) | % | 0.16 | 0.011 | 0.16 | 0.15 | 0.17 |
| DLL | Eicosenoic (C20:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Eicosadienoic (C20:2) | % | 0.03 | 0.002 | 0.03 | 0.03 | 0.03 |
| DLL | Eicosadienoic (C20:2), Eicosatrienoic(C20:3) | % | 0.45 | 0.179 | 0.51 | 0.17 | 0.64 |
| EPL | Behenic (C22:0) | % | 0.02 | 0.002 | 0.02 | 0.02 | 0.02 |
| DLL | Behenic (C22:0) | % | NA | NA | NA | NA | NA |
| EPL | Erucic (C22:1) | % | 0.06 | 0.013 | 0.07 | 0.04 | 0.08 |
| DLL | Erucic (C22:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Lignoceric (C24:0) | % | 0.02 | 0.002 | 0.02 | 0.01 | 0.02 |
| DLL | Lignoceric (C24:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Nervonic (C24:1) | % | 0.08 | 0.007 | 0.07 | 0.07 | 0.08 |
| DLL | Nervonic (C24:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

250  [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories

  2 NA: Not Analyzed
252 The following fatty acids were below the level of detection for Lab EPL: Arachidonic (C20:4), Capric (C10:0),
253 Caproic (C6:0), Caprylic (C8:0), Docosadienoic (C22:2), Eicosatrienoic (C20:3), Gamma Linolenic (C18:3),
254 Heneicosanoic (C21:0), Lauric (C12:0), Myristoleic (C14:1), Nonadecanoic (C19:0), Pentadecanoic (C15:0),
255 Pentadecenoic (C15:1), Tricosanoic (C23:0); for Lab DLL: Arachidonic (C20:4), Docosahexanoic acid (C22:6);
256 Eicosapentaenoic acid (C20:5).
257

258 Table 6. CCWG-15-32 Fatty Acids as a Percent of Total Fatty Acids

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Myristic (C14:0) | % | 0.09 | 0.006 | 0.09 | 0.09 | 0.10 |
| DLL | Myristic (C14:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Palmitic (C16:0) | % | 3.98 | 0.066 | 4.00 | 3.86 | 4.02 |
| DLL | Palmitic (C16:0) | % | 4.06 | 0.076 | 4.04 | 4.00 | 4.20 |
| EPL | Palmitoleic (C16:1) | % | 0.15 | 0.012 | 0.16 | 0.13 | 0.16 |
| DLL | Palmitoleic (C16:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecanoic (C17:0) | % | 0.03 | 0.018 | 0.03 | 0.00 | 0.05 |
| DLL | Heptadecanoic (C17:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecenoic (C17:1) | % | 0.07 | 0.011 | 0.08 | 0.06 | 0.09 |
| DLL | Heptadecenoic (C17:1) | % | NA[2] | NA | NA | NA | NA |
| EPL | Stearic (C18:0) | % | 1.00 | 0.044 | 1.02 | 0.94 | 1.04 |
| DLL | Stearic (C18:0) | % | 0.99 | 0.050 | 1.01 | 0.91 | 1.03 |
| EPL | Oleic (C18:1) | % | 40.14 | 1.223 | 40.55 | 38.50 | 41.55 |
| DLL | Oleic (C18:1) | % | 40.07 | 0.943 | 40.19 | 38.61 | 41.06 |
| EPL | Linoleic (C18:2) | % | 34.41 | 0.611 | 34.20 | 33.80 | 35.30 |
| DLL | Linoleic (C18:2) | % | 34.48 | 0.337 | 34.39 | 34.17 | 34.94 |
| EPL | Linolenic (C18:3) | % | 18.15 | 0.722 | 17.90 | 17.55 | 19.40 |
| DLL | Linolenic (C18:3) | % | 16.55 | 0.632 | 16.32 | 16.18 | 17.67 |
| EPL | Nonadecenoic (C19:1) | % | 0.03 | 0.018 | 0.03 | 0.00 | 0.05 |
| DLL | Nonadecenoic (C19:1) | % | NA | NA | NA | NA | NA |
| EPL | Arachidic (C20:0) | % | 0.03 | 0.004 | 0.03 | 0.03 | 0.04 |
| DLL | Arachidic (C20:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

| | | | | | | Range | |
|---|---|---|---|---|---|---|---|
| EPL | Eicosenoic (C20:1) | % | 0.16 | 0.011 | 0.16 | 0.15 | 0.17 |
| DLL | Eicosenoic (C20:1) | % | 1.47 | 0.088 | 1.44 | 1.38 | 1.60 |
| EPL | Eicosadienoic (C20:2) | % | 0.03 | 0.002 | 0.03 | 0.03 | 0.03 |
| DLL | Eicosadienoic (C20:2), Eicosatrienoic(C20:3) | % | 2.38 | 0.939 | 2.54 | 1.09 | 3.54 |
| EPL | Behenic (C22:0) | % | 0.02 | 0.002 | 0.02 | 0.02 | 0.02 |
| DLL | Behenic (C22:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Erucic (C22:1) | % | 0.06 | 0.013 | 0.07 | 0.04 | 0.08 |
| DLL | Erucic (C22:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Lignoceric (C24:0) | % | 0.07 | 0.021 | 0.08 | 0.04 | 0.09 |
| DLL | Lignoceric (C24:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Nervonic (C24:1) | % | 0.37 | 0.024 | 0.37 | 0.34 | 0.40 |
| DLL | Nervonic (C24:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

[1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
[2] NA: Not Analyzed
The following fatty acids were below the level of detection for Lab A: Arachidonic (C20:4), Capric (C10:0),
Caproic (C6:0), Caprylic (C8:0), Docosadienoic (C22:2), Eicosatrienoic (C20:3), Gamma Linolenic (C18:3),
Heneicosanoic (C21:0), Lauric (C12:0), Myristoleic (C14:1), Nonadecanoic (C19:0), Pentadecanoic (C15:0),
Pentadecenoic (C15:1), Tricosanoic (C23:0); for Lab B: Arachidonic (C20:4), Docosahexanoic acid (C22:6);
Eicosapentaenoic acid (C20:5).
266

267 <u>Glucosinolate Characterization and Quantification</u>
268
269     The sole glucosinolate, sinigrin, averaged 85.7 and 86.9 µmoles/g of CCWG-15-32 for

270 the two detection methods (Table 7).  The median value was higher than the mean due to one lot

271 containing almost half the amount of sinigrin as the other four lots.  This is reflected by the low

272 minimum value as compared to the maximum value.  The much lower sinigrin level in the one

273 lot may have resulted from the soil conditions at that site.

274
275 Table 7.  CoverCress CS (CCWG-15-32) glucosinolates (Sinigrin, 100% dry matter basis) – EPL
276 Bio Analytical Services

| Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|
| Sinigrin – MS | µmoles/g | 85.7 | 16.83 | 90.7 | 56.2 | 98.6 |
| Sinigrin – UV | µmoles/g | 86.9 | 23.97 | 95.7 | 45 | 102 |

<u>Minerals and Vitamins</u>

Table 8 contains the results of the mineral analyses.  EPL reported consistently

higher mineral levels than Dairyland Laboratories, with the exception of zinc.  The

discrepancy between labs is unexplained.  Generally, the values for the five lots had

symmetric distribution.  One location had a sulfur level that was 50% of the others.  This

lower sulfur value corresponds to the lower glucosinolate level of the sample from the same

location.  Glucosinolates are secondary sulfur compounds commonly found in brassica

species (Aghajanzada *et al*., 2014), which suggests there is a plausible relationship between

both values being lower at this site.  The reason for the lower level at one location as

compared to the other four locations is not clear but may have been a result of the soil

fertilization.   As expected, mean copper values were approximately 800 ppm because of the

addition of copper sulfate.  The analytical results confirm that the target level of 800 ppm

copper was achieved.

Table 8. CCWG-15-32 minerals (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Calcium | % | 0.849 | 0.079 | 0.826 | 0.791 | 0.982 |
| DLL | Calcium | % | 0.72 | 0.108 | 0.69 | 0.60 | 0.88 |
| EPL | Phosphorus | % | 0.991 | 0.138 | 1.060 | 0.840 | 1.135 |
| DLL | Phosphorus | % | 0.70 | 0.123 | 0.78 | 0.56 | 0.81 |
| EPL | Magnesium | % | 0.378 | 0.039 | 0.388 | 0.335 | 0.428 |
| DLL | Magnesium | % | 0.29 | 0.010 | 0.30 | 0.28 | 0.30 |
| EPL | Potassium | % | 0.951 | 0.073 | 0.923 | 0.909 | 1.080 |
| DLL | Potassium | % | 0.74 | 0.114 | 0.71 | 0.63 | 0.93 |
| EPL | Sulfur | % | 1.008 | 0.230 | 1.120 | 0.600 | 1.140 |
| DLL | Sulfur | % | 0.87 | 0.230 | 0.98 | 0.47 | 1.05 |
| EPL | Sodium | % | 0.005 | 0.003 | 0.004 | 0.003 | 0.009 |
| DLL | Sodium | % | 0.00 | 0.003 | 0.00 | 0.00 | 0.01 |
| EPL | Chloride | ppm | 185.10 | 35.379 | 186.50 | 143.00 | 238.50 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| DLL | Chloride | ppm | 400 | 90 | 500 | 300 | 500 |
| EPL | Iron | ppm | 102.18 | 12.350 | 108.50 | 83.850 | 112.00 |
| DLL | Iron | ppm | 40 | 55.1 | 29 | 16 | 98.5 |
| EPL | Copper | ppm | 820.70 | 17.101 | 814.00 | 806.00 | 845.50 |
| DLL | Copper | ppm | 808 | 68.7 | 811 | 704 | 890 |
| EPL | Manganese | ppm | 31.150 | 4.019 | 29.650 | 27.000 | 37.650 |
| DLL | Manganese | ppm | 27 | 2.6 | 26 | 23.5 | 30 |
| EPL | Zinc | ppm | 37.710 | 2.086 | 39.200 | 35.400 | 39.250 |
| DLL | Zinc | ppm | 55 | 11.4 | 55 | 43 | 72 |

293
294

298  [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
299

Table 9 contains the vitamins as analyzed by EPL for the five lots. Generally, the

values for the five lots had symmetric distribution.  Nutritionists and feed formulators, as

common practice, do not use the vitamin content of feedstuffs in formulating diets due to the

variability, stability, low levels in the feed ingredient and the low cost of supplementing

vitamins. The Canola Council of Canada in their Canola Feed Guide (2019) state, "As is

recommended with most natural sources of vitamins in animal feeds, users should not place

too much reliance on these values and use supplemental vitamin premixes instead."

Table 9. CCWG-15-32 vitamins (100% dry matter basis) – EPL Bio Analytical Services

| Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|
| Folic Acid | mg/kg | 3.3 | 1.35 | 3.2 | 1.8 | 5.3 |
| Niacin | mg/kg | 51.2 | 10.52 | 48.6 | 42.1 | 69.3 |
| Vitamin B1 (Thiamine) | mg/kg | 5.4 | 0.25 | 5.3 | 5.2 | 5.8 |
| Vitamin B2 (Riboflavin) | mg/kg | 2.7 | 0.54 | 2.6 | 2.3 | 3.6 |
| Vitamin B6 (Pyridoxine) | mg/kg | 10.6 | 1.69 | 10.3 | 8.8 | 13.4 |
| Alpha-Tocopherol | mg/kg | 118.0 | 10.86 | 121.5 | 99.0 | 125.5 |
| Beta-Tocopherol | mg/kg | 1.8 | 0.07 | 1.8 | 1.7 | 1.9 |
| Delta-Tocopherol | mg/kg | 2.3 | 0.16 | 2.3 | 2.1 | 2.5 |
| Gamma-Tocopherol | mg/kg | 53.4 | 14.65 | 48.6 | 40.3 | 78.5 |

<u>Sinapine and Mycotoxins</u>

Sinapine, like glucosinolates, is an anti-nutrient found in modern oilseed rape and

canola meals. Sinapine is metabolized in animals to trimethylamine, which is further

metabolized by trimethyl oxidase. Certain brown egg laying strains of hens have been

reported to have lower hepatic trimethyl oxidase activity apparently leading to accumulation

of trimethylamine in some brown eggs and an associated "fishy taint". As summarized by

Rymer & Short (2003), sinapines are present in modern rapeseed at levels of 12-23 g/kg

seed (Huisman & Tolman, 2001). They are converted in the large intestine to

trimethylamine, which apparently produces an undesirable fishy odor in the eggs of certain

brown egg chickens. Some of these types of birds lack the ability to produce trimethylamine

oxidase (Jeroch, 2008), and as a result the trimethylamine accumulates in the eggs of these

birds. This odor in eggs occurs at levels of 0.8 mg/kg diet (Fenwick, 1982). The

concentrations of sinapine in CCWG-15-32 were  <0.05% DM basis  as reported by EPL in

all five lots.  This is much lower in comparison to canola meal where the Canola Council of

Canada (2015) reports sinapine levels of 1.0% as is basis (with an average 12% moisture

content) or 1.14% on a 100% DM basis.

The following mycotoxins were below the limits of detection in all samples from all of

the five lots:  Aflatoxin B1, B2, G1, G2; T-2 Toxin; Ochratoxin A; Deoxynivalenol (DON), 3-

Acetyl-DON; 15-Acetyl-DON; Zearalenone; Fumonisin B1, B2, B3; Ergosine; Ergotamine;

Ergocornine; and Ergocryptine.   For Ergocristine, which is a natural ergot alkaloid, four lots

were below the levels of detection. For one lot, one of the two replicates was below the level of

detection and the second had 15.9 ppb on a dry matter basis with level of detection at 1.5 ppb.

**Study B – B3WG-15-32**

<u>Proximate Analyses</u>

Table 10 contains the results of proximate analyses.  Just as with the CCWG samples, it

was postulated that Dairyland Laboratories's method extracted more fat from the product than

EPL's method. In addition, there was greater variation noted with EPL's fat values than with

Dairyland Labs.  Fat in the sample may affect the fiber analysis with higher fat resulting in

inflated crude fiber, ADF and NDF values.  Moisture levels are high due to the addition of the

copper sulfate solution to the CoverCress grain.  Since the mean and median values are similar

for all proximates, the values from the five lots are evenly distributed or symmetrical in

distribution.

Table 10.  CoverCress CS (B3WG-15-32) Proximates (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Moisture | % | 19.6 | 0.24 | 19.6 | 19.3 | 19.9 |
| DLL | Moisture | % | 16.6 | 0.38 | 16.7 | 16.1 | 17.1 |
| EPL | Dry Matter | % | 80.4 | 0.24 | 80.5 | 80.1 | 80.7 |
| DLL | Dry Matter | % | 83.4 | 0.38 | 83.3 | 82.9 | 83.9 |
| EPL | Crude Fiber | % | 20.0 | 2.82 | 21.0 | 15.6 | 22.6 |
| DLL | Crude Fiber | % | 12.1 | 0.71 | 12.5 | 10.9 | 12.5 |
| EPL | Acid Detergent Fiber | % | 25.8 | 2.73 | 25.3 | 22.1 | 28.7 |
| DLL | Acid Detergent Fiber | % | 17.8 | 2.04 | 16.8 | 15.9 | 20.2 |
| EPL | Neutral Detergent Fiber | % | 26.9 | 3.32 | 27.7 | 22.0 | 30.2 |
| DLL | Neutral Detergent Fiber | % | 22.6 | 1.48 | 21.9 | 21.3 | 24.4 |
| EPL | Total Dietary Fiber | % | 28.8 | 1.04 | 28.7 | 27.7 | 30.3 |
| EPL | Crude Protein | % | 25.4 | 1.27 | 25.0 | 24.1 | 27.4 |
| DLL | Crude Protein | % | 23.9 | 0.97 | 24.1 | 22.5 | 25.1 |
| EPL | Crude Fat | % | 32.7 | 1.62 | 33.0 | 30.4 | 34.8 |
| DLL | Crude Fat | % | 37.0 | 1.07 | 37.0 | 35.7 | 38.5 |
| EPL | Carbohydrates | % | 36.3 | 2.12 | 36.8 | 32.7 | 38.1 |
| EPL | Ash | % | 5.6 | 0.45 | 5.4 | 5.1 | 6.2 |
| DLL | Ash | % | 6.0 | 0.86 | 6.1 | 4.6 | 6.7 |

346 [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
347

<u>Amino Acid Analyses</u>

Table 11 contains the amino acid profile of B3WG-15-32 when expressed on a 100% dry

matter basis and Table 12 contains the amino acid profile of B3WG-15-32 when expressed on a

protein basis.   The amino acid values from the five lots showed a symmetrical distribution with

the means and median values being similar.  Dairyland had a higher percentage of total amino

acids when total amino acids were expressed as a percent of crude protein (88.1% versus 83.1%).

Table 11.  CoverCress CS (B3WG-15-32) Amino Acids (100% dry matter basis)

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Alanine | % | 1.13 | 0.023 | 1.14 | 1.11 | 1.15 |
| DLL | Alanine | % | 1.00 | 0.034 | 0.99 | 0.97 | 1.06 |
| EPL | Arginine | % | 1.39 | 0.067 | 1.39 | 1.30 | 1.46 |
| DLL | Arginine | % | 1.57 | 0.052 | 1.55 | 1.51 | 1.63 |
| EPL | Aspartic Acid | % | 1.93 | 0.054 | 1.91 | 1.91 | 2.03 |
| DLL | Aspartic Acid | % | 1.92 | 0.031 | 1.93 | 1.88 | 1.96 |
| EPL | Cystine | % | 0.38 | 0.022 | 0.38 | 0.36 | 0.41 |
| DLL | Cysteine | % | 0.38 | 0.017 | 0.39 | 0.36 | 0.41 |
| EPL | Glutamic Acid | % | 3.83 | 0.110 | 3.81 | 3.68 | 3.95 |
| DLL | Glutamic Acid | % | 3.59 | 0.133 | 3.59 | 3.40 | 3.75 |
| EPL | Glycine | % | 1.38 | 0.064 | 1.38 | 1.30 | 1.46 |
| DLL | Glycine | % | 1.45 | 0.041 | 1.45 | 1.39 | 1.51 |
| EPL | Histidine | % | 0.54 | 0.022 | 0.54 | 0.52 | 0.57 |
| DLL | Histidine | % | 0.47 | 0.015 | 0.47 | 0.45 | 0.49 |
| EPL | Isoleucine | % | 0.93 | 0.019 | 0.93 | 0.90 | 0.94 |
| DLL | Isoleucine | % | 0.73 | 0.047 | 0.74 | 0.68 | 0.80 |
| EPL | Leucine | % | 1.63 | 0.046 | 1.63 | 1.56 | 1.68 |
| DLL | Leucine | % | 1.59 | 0.059 | 1.58 | 1.53 | 1.68 |
| EPL | Lysine | % | 1.28 | 0.035 | 1.28 | 1.24 | 1.34 |
| DLL | Lysine | % | 1.12 | 0.032 | 1.13 | 1.09 | 1.17 |
| EPL | Methionine | % | 0.37 | 0.029 | 0.38 | 0.33 | 0.41 |
| DLL | Methionine | % | 0.33 | 0.019 | 0.33 | 0.31 | 0.36 |
| EPL | Phenylalanine | % | 0.99 | 0.060 | 0.98 | 0.92 | 1.08 |
| DLL | Phenylalanine | % | 1.04 | 0.033 | 1.04 | 0.99 | 1.08 |
| EPL | Proline | % | 1.22 | 0.043 | 1.23 | 1.17 | 1.27 |
| DLL | Proline | % | 1.36 | 0.020 | 1.37 | 1.33 | 1.38 |
| EPL | Serine | % | 0.93 | 0.035 | 0.93 | 0.89 | 0.99 |
| DLL | Serine | % | 0.99 | 0.019 | 0.99 | 0.97 | 1.02 |
| EPL | Threonine | % | 1.03 | 0.030 | 1.04 | 0.99 | 1.07 |
| DLL | Threonine | % | 1.06 | 0.025 | 1.05 | 1.03 | 1.10 |
| EPL | Tryptophan | % | 0.36 | 0.016 | 0.37 | 0.33 | 0.37 |
| DLL | Tryptophan | % | 0.47 | 0.059 | 0.47 | 0.42 | 0.56 |
| EPL | Tyrosine | % | 0.49 | 0.032 | 0.49 | 0.45 | 0.54 |
| DLL | Tyrosine | % | 0.74 | 0.018 | 0.74 | 0.72 | 0.75 |
| EPL | Valine | % | 1.23 | 0.019 | 1.23 | 1.20 | 1.26 |
| DLL | Valine | % | 1.17 | 0.063 | 1.17 | 1.11 | 1.27 |

355  [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories

357    Table 12. CoverCress CS (B3WG-15-32) Amino Acids as Percent of Protein

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Alanine | % | 4.46 | 0.164 | 4.48 | 4.20 | 4.60 |
| DLL | Alanine | % | 4.20 | 0.118 | 4.19 | 4.05 | 4.38 |
| EPL | Arginine | % | 5.49 | 0.160 | 5.44 | 5.31 | 5.70 |
| DLL | Arginine | % | 6.59 | 0.138 | 6.59 | 6.43 | 6.74 |
| EPL | Aspartic Acid | % | 7.64 | 0.450 | 7.73 | 6.97 | 8.12 |
| DLL | Aspartic Acid | % | 8.05 | 0.220 | 7.99 | 7.81 | 8.33 |
| EPL | Cystine | % | 1.50 | 0.104 | 1.50 | 1.41 | 1.67 |
| DLL | Cysteine | % | 1.62 | 0.076 | 1.62 | 1.49 | 1.69 |
| EPL | Glutamic Acid | % | 15.13 | 0.711 | 15.35 | 13.90 | 15.75 |
| DLL | Glutamic Acid | % | 15.07 | 0.172 | 15.09 | 14.88 | 15.27 |
| EPL | Glycine | % | 5.45 | 0.094 | 5.41 | 5.36 | 5.56 |
| DLL | Glycine | % | 6.09 | 0.087 | 6.06 | 6.01 | 6.20 |
| EPL | Histidine | % | 2.14 | 0.043 | 2.14 | 2.08 | 2.19 |
| DLL | Histidine | % | 1.99 | 0.027 | 1.99 | 1.96 | 2.02 |
| EPL | Isoleucine | % | 3.66 | 0.124 | 3.72 | 3.45 | 3.74 |
| DLL | Isoleucine | % | 3.07 | 0.125 | 3.09 | 2.87 | 3.19 |
| EPL | Leucine | % | 6.42 | 0.162 | 6.50 | 6.13 | 6.51 |
| DLL | Leucine | % | 6.69 | 0.081 | 6.71 | 6.56 | 6.78 |
| EPL | Lysine | % | 5.05 | 0.330 | 5.08 | 4.53 | 5.35 |
| DLL | Lysine | % | 4.71 | 0.121 | 4.69 | 4.55 | 4.85 |
| EPL | Methionine | % | 1.47 | 0.083 | 1.49 | 1.38 | 1.57 |
| DLL | Methionine | % | 1.39 | 0.050 | 1.38 | 1.33 | 1.45 |
| EPL | Phenylalanine | % | 3.89 | 0.072 | 3.91 | 3.80 | 3.96 |
| DLL | Phenylalanine | % | 4.35 | 0.044 | 4.38 | 4.30 | 4.40 |
| EPL | Proline | % | 4.83 | 0.136 | 4.87 | 4.60 | 4.95 |
| DLL | Proline | % | 5.73 | 0.164 | 5.71 | 5.48 | 5.91 |
| EPL | Serine | % | 3.68 | 0.062 | 3.69 | 3.60 | 3.76 |
| DLL | Serine | % | 4.16 | 0.116 | 4.13 | 4.06 | 4.36 |
| EPL | Threonine | % | 4.08 | 0.103 | 4.11 | 3.91 | 4.18 |
| DLL | Threonine | % | 4.44 | 0.098 | 4.38 | 4.36 | 4.58 |
| EPL | Tryptophan | % | 1.41 | 0.058 | 1.44 | 1.34 | 1.48 |
| DLL | Tryptophan | % | 1.98 | 0.174 | 1.93 | 1.80 | 2.21 |
| EPL | Tyrosine | % | 1.93 | 0.051 | 1.97 | 1.87 | 1.97 |
| DLL | Tyrosine | % | 3.09 | 0.073 | 3.08 | 2.99 | 3.18 |
| EPL | Valine | % | 4.85 | 0.170 | 4.92 | 4.56 | 4.98 |
| DLL | Valine | % | 4.89 | 0.180 | 4.98 | 4.59 | 5.04 |

358    [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
359

<u>Fatty Acid Characterization</u>

Tables 13 and 14 contains the fatty acid profile of B3WG-15-32 when expressed on

a 100% dry matter basis and percent of total fatty acids, respectively.  In general, there was

good agreement between laboratories. The fatty acid values from the five lots showed a

symmetrical distribution with the means and median values being similar.  The major fatty

acids are oleic (~43% of the total fatty acids) followed by linoleic (~32% of the total fatty

acids) and linolenic (~18% of the total fatty acids).  These three fatty acids comprise over

90% of the fatty acids in B3WG-15-32.  No high polyunsaturated long chain fatty acids

such as EPA and DHA were detected. Results confirm that B3WG-15-32 contains

negligible levels of erucic acid. Dairyland Laboratories did not detect it and EPL reported

erucic acid to be less than 0.5% of the total fatty acids.

Table 13.  CoverCress CS (B3WG-15-32) Fatty Acids (100% dry matter basis)

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Myristic (C1 4:0) | % | 0.02 | 0.002 | 0.02 | 0.01 | 0.02 |
| DLL | Myristic (C14:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Palmitic (C16:0) | % | 0.62 | 0.039 | 0.62 | 0.56 | 0.67 |
| DLL | Palmitic (C16:0) | % | 1.17 | 0.051 | 1.16 | 1.11 | 1.24 |
| EPL | Palmitoleic (C16:1) | % | 0.02 | 0.002 | 0.02 | 0.02 | 0.03 |
| DLL | Palmitoleic (C16:1) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecanoic (C17:0) | % | 0.01 | 0.001 | 0.01 | 0.01 | 0.01 |
| DLL | Heptadecanoic (C17:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecenoic (C17:1) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Heptadecenoic (C17:1) | % | NA[2] | NA | NA | NA | NA |
| EPL | Stearic (C18:0) | % | 0.13 | 0.010 | 0.13 | 0.12 | 0.14 |
| DLL | Stearic (C18:0) | % | 0.26 | 0.017 | 0.27 | 0.24 | 0.28 |
| EPL | Oleic (C18:1) | % | 6.75 | 0.430 | 6.74 | 6.17 | 7.38 |
| DLL | Oleic (C18:1) | % | 13.61 | 0.930 | 13.84 | 12.35 | 14.86 |
| EPL | Linoleic (C18:2) | % | 5.16 | 0.411 | 5.14 | 4.77 | 5.84 |
| DLL | Linoleic (C18:2) | % | 9.57 | 0.733 | 9.83 | 8.57 | 10.44 |
| EPL | Linolenic (C18:3) | % | 2.91 | 0.250 | 2.83 | 2.68 | 3.34 |
| DLL | Linolenic (C18:3) | % | 5.16 | 0.423 | 5.28 | 4.72 | 5.70 |
| EPL | Nonadecenoic (C19:1) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Nonadecenoic (C19:1) | % | NA | NA | NA | NA | NA |
| EPL | Arachidic (C20:0) | % | 0.02 | 0.001 | 0.02 | 0.02 | 0.02 |
| DLL | Arachidic (C20:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Eicosenoic (C20:1) | % | 0.14 | 0.009 | 0.14 | 0.12 | 0.14 |

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| DLL | Eicosenoic (C20:1) | % | 0.23 | 0.014 | 0.23 | 0.21 | 0.25 |
| EPL | Eicosadienoic (C20:2) | % | 0.02 | 0.003 | 0.02 | 0.02 | 0.03 |
| DLL | Eicosadienoic (C20:2), Eicosatrienoic(C20:3) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Behenic (C22:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Behenic (C22:0) | % | NA | NA | NA | NA | NA |
| EPL | Erucic (C22:1) | % | 0.07 | 0.022 | 0.06 | 0.05 | 0.11 |
| DLL | Erucic (C22:1) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Lignoceric (C24:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Lignoceric (C24:0) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| EPL | Nervonic (C24:1) | % | 0.05 | 0.005 | 0.05 | 0.04 | 0.06 |
| DLL | Nervonic (C24:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

[1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
[2] NA: Not Analyzed
The following fatty acids were below the level of detection for Lab EPL: Capric (C10:0), Caproic (C6:0), Caprylic
(C8:0), Docosadienoic (C22:2), Docosahexaenoic (C22:6), Eicosatrienoic (C20:3), Eicosapentaenoic (C20:5),
Gamma Linolenic (C18:3), Heneicosanoic (C21:0), Heptadecenoic (C17:1), Lauric (C12:0), Myristoleic (C14:1),
Nonadecanoic (C19:0), Nonadecenoic (C19:1), Pentadecanoic (C15:0), Pentadecenoic (C15:1), Tricosanoic (C23:0);
for Lab DLL: Arachidonic (C20:4), Docosahexanoic acid (C22:6); Eicosapentaenoic acid (C20:5).

381
382 Table 14.  CoverCress CS (B3WG-15-32) Fatty Acids as a Percent of Total Fatty Acids

| Analytical Lab | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Range Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Myristic (C14:0) | % | 0.10 | 0.011 | 0.10 | 0.09 | 0.11 |
| DLL | Myristic (C14:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Palmitic (C16:0) | % | 3.88 | 0.042 | 3.89 | 3.84 | 3.95 |
| DLL | Palmitic (C16:0) | % | 3.89 | 0.079 | 3.91 | 3.77 | 3.96 |
| EPL | Palmitoleic (C16:1) | % | 0.15 | 0.018 | 0.16 | 0.12 | 0.17 |
| DLL | Palmitoleic (C16:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecanoic (C17:0) | % | 0.03 | 0.018 | 0.03 | 0.00 | 0.05 |
| DLL | Heptadecanoic (C17:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Heptadecenoic (C17:1) | % | 0.07 | 0.011 | 0.08 | 0.06 | 0.09 |
| DLL | Heptadecenoic (C17:1) | % | NA | NA | NA | NA | NA |
| EPL | Stearic (C18:0) | % | 0.84 | 0.039 | 0.84 | 0.77 | 0.88 |
| DLL | Stearic (C18:0) | % | 0.88 | 0.040 | 0.87 | 0.83 | 0.93 |
| EPL | Oleic (C18:1) | % | 42.32 | 2.261 | 42.40 | 39.00 | 45.35 |
| DLL | Oleic (C18:1) | % | 45.40 | 2.134 | 45.34 | 42.22 | 48.16 |
| EPL | Linoleic (C18:2) | % | 32.42 | 1.425 | 32.50 | 30.25 | 34.25 |
| DLL | Linoleic (C18:2) | % | 31.89 | 1.431 | 31.86 | 29.84 | 33.87 |
| EPL | Linolenic (C18:3) | % | 18.25 | 0.793 | 18.15 | 17.35 | 19.50 |
| DLL | Linolenic (C18:3) | % | 17.18 | 0.737 | 17.24 | 16.44 | 18.26 |
| EPL | Nonadecenoic (C19:1) | % | 0.03 | 0.018 | 0.03 | 0.00 | 0.05 |
| DLL | Nonadecenoic (C19:1) | % | NA | NA | NA | NA | NA |
| EPL | Arachidic (C20:0) | % | 0.13 | 0.013 | 0.13 | 0.13 | 0.016 |
| DLL | Arachidic (C20:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Eicosenoic (C20:1) | % | 0.85 | 0.024 | 0.84 | 0.82 | 0.87 |

| | | | Mean | Standard Deviation | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| DLL | Eicosenoic (C20:1) | % | 0.77 | 0.042 | 0.78 | 0.70 | 0.82 |
| EPL | Eicosadienoic (C20:2) | % | 0.03 | 0.002 | 0.03 | 0.03 | 0.03 |
| DLL | Eicosadienoic (C20:2), Eicosatrienoic(C20:3) | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Arachidonic (20:4) | % | 0.02 | 0.018 | 0.02 | 0.00 | 0.05 |
| EPL | Behenic (C22:0) | % | 0.06 | 0.003 | 0.06 | 0.06 | 0.07 |
| DLL | Behenic (C22:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Erucic (C22:1) | % | 0.43 | 0.109 | 0.38 | 0.36 | 0.62 |
| DLL | Erucic (C22:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Lignoceric (C24:0) | % | 0.07 | 0.005 | 0.07 | 0.06 | 0.07 |
| DLL | Lignoceric (C24:0) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| EPL | Nervonic (C24:1) | % | 0.31 | 0.025 | 0.30 | 0.30 | 0.36 |
| DLL | Nervonic (C24:1) | % | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

[1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories

[2] NA: Not Analyzed

The following fatty acids were below the level of detection for Lab A: Arachidonic (C20:4), Capric (C10:0), Caproic (C6:0), Caprylic (C8:0), Docosadienoic (C22:2), Eicosatrienoic (C20:3), Gamma Linolenic (C18:3), Heneicosanoic (C21:0), Lauric (C12:0), Myristoleic (C14:1), Nonadecanoic (C19:0), Pentadecanoic (C15:0), Pentadecenoic (C15:1), Tricosanoic (C23:0); for Lab B: Arachidonic (C20:4), Docosahexanoic acid (C22:6); Eicosapentaenoic acid (C20:5).

## Glucosinolate Characterization and Quantification

Sinigrin averaged 105.4 µmoles/g of B3WG-15-32 on a 100% dry matter basis and was similar among lots (Table 15). Sinigrin was measured using the Ultraviolet method (UV).

Table 15.  CoverCress CS (B3WG-15-32) Glucosinolates (Sinigrin, 100% dry matter basis) -EPL Bio Analytical Services

| | | | Standard | | Range | |
|---|---|---|---|---|---|---|
| Variable | Units | Mean | Deviation | Median | Minimum | Maximum |
| Sinigrin - UV | µmoles/g | 105.4 | 4.21 | 104.3 | 101.4 | 110.7 |

## Minerals, Sinapine, and Mycotoxins

Table 16 contains the results to the mineral analyses.  EPL and Dairyland Laboratories results varied.  The discrepancy between labs is unexplained.  Generally, the values for the five lots had symmetric distribution.  The high level of copper is due to the addition of copper sulfate.  Target addition of copper was 800 ppm.  The analytical results confirm that the target levels were achieved.

407
408 Table 16. CoverCress CS (B3WG-15-32) Minerals (100% dry matter basis)

| Analytical Lab[1] | Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| EPL | Calcium | % | 0.93 | 0.074 | 0.95 | 0.84 | 1.01 |
| DLL | Calcium | % | 1.25 | 0.154 | 1.31 | 0.99 | 1.39 |
| EPL | Phosphorus | % | 0.99 | 0.135 | 0.96 | 0.80 | 1.14 |
| DLL | Phosphorus | % | 0.68 | 0.134 | 0.71 | 0.48 | 0.81 |
| EPL | Magnesium | % | 0.38 | 0.091 | 0.37 | 0.30 | 0.48 |
| DLL | Magnesium | % | 0.35 | 0.087 | 0.30 | 0.27 | 0.44 |
| EPL | Potassium | % | 1.11 | 0.083 | 1.05 | 1.05 | 1.23 |
| DLL | Potassium | % | 1.04 | 0.117 | 1.05 | 0.89 | 1.17 |
| EPL | Sulfur | % | 1.17 | 0.061 | 1.15 | 1.12 | 1.26 |
| DLL | Sulfur | % | 0.94 | 0.064 | 0.96 | 0.86 | 1.01 |
| EPL | Sodium | % | 0.00 | 0.000 | 0.00 | 0.00 | 0.00 |
| DLL | Sodium | % | 0.01 | 0.004 | 0.01 | 0.01 | 0.02 |
| EPL | Chloride | ppm | 399.4 | 85.334 | 432.00 | 275.50 | 473.00 |
| DLL | Chloride | ppm | 800 | 25 | 700 | 600 | 1200 |
| EPL | Iron | ppm | 107.85 | 48.471 | 84.95 | 63.90 | 162.50 |
| DLL | Iron | ppm | 72 | 30.7 | 57 | 40.5 | 113 |
| EPL | Copper | ppm | 959.70 | 29.641 | 955.00 | 930.50 | 1001.5 |
| DLL | Copper | ppm | 1052 | 257.8 | 1087.5 | 633 | 1315 |
| EPL | Manganese | ppm | 36.96 | 3.254 | 38.30 | 31.80 | 39.75 |
| DLL | Manganese | ppm | 46 | 6.5 | 47.5 | 39 | 55 |
| EPL | Zinc | ppm | 38.88 | 4.672 | 40.20 | 33.65 | 45.00 |
| DLL | Zinc | ppm | 46 | 15.6 | 43 | 34 | 73 |

409 [1] Analytical Lab: EPL – EPL Bio Analytical Services; DLL – Dairyland Laboratories
410

The concentrations of sinapine in B3WG-15-32 were <0.05% as reported by EPL in all

five lots.  This is in comparison to canola meal where the Canola Council of Canada (2015)

reports sinapine levels of 1.0% as is basis (with an average 12% moisture content) or 1.14% on a

100% DM basis.

No mycotoxins were detected in the five lots.  All mycotoxins were below the limits of

detection in all samples from all of the five lots: Aflatoxin B1, B2, G1, G2; T-2 Toxin;

Ochratoxin A; Deoxynivalenol (DON), 3-Acetyl-DON; 15-Acetyl-DON; Zearalenone;

Fumonisin B1, B2, B3; Ergosine; Ergotamine; Ergocristine; and Ergocornine; and

Ergocryptine.

<u>Vitamins</u>

424      Table 17 contains the vitamins as analyzed by EPL for the five lots. Generally, the

425  values for the five lots had symmetric distribution.

427  Table 17. CoverCress CS (B3WG-15-32) Vitamins (100% dry matter basis) – EPL Bio Analytical
428  Services

| Variable | Units | Mean | Standard Deviation | Median | Range Minimum | Maximum |
|---|---|---|---|---|---|---|
| Folic Acid | mg/kg | 5.6 | 0.67 | 5.2 | 5.0 | 6.6 |
| Niacin | mg/kg | 40.2 | 2.06 | 40.6 | 37.1 | 42.5 |
| Vitamin B1 (Thiamine) | mg/kg | 5.2 | 0.06 | 5.2 | 5.1 | 5.3 |
| Vitamin B2 (Riboflavin) | mg/kg | 4.7 | 0.29 | 4.6 | 4.3 | 5.0 |
| Vitamin B6 (Pyridoxine) | mg/kg | 6.1 | 0.17 | 6.1 | 5.9 | 6.3 |
| Alpha-Tocopherol | mg/kg | 181.8 | 24.28 | 174.5 | 160.5 | 218.0 |
| Beta-Tocopherol | mg/kg | 3.1 | 0.13 | 3.1 | 2.9 | 3.4 |
| Delta-Tocopherol | mg/kg | <1.25 | | <1.25 | <1.25 | <1.25 |
| Gamma-Tocopherol | mg/kg | 13.4 | 1.75 | 12.5 | 11.9 | 15.8 |

**Conclusions**

The results of these extensive compositional and nutritional analyses show:

●    The composition and nutritional components of CCWG-15-32 and B3WG-15-32

lots were generally consistent with very good process control and low inter-lot

variability

●    As expected by loss of function of the FAE1 gene, the fatty acid profile of the

CCWG-15-32 and B3WG-15-32 lots showed negligible erucic acid levels

●    As expected by loss of function of the TT8 gene, mean ADF and crude fibers

levels were in acceptable ranges for broiler diets

●    Sinigrin (2-propenyl glucosinolate) was the only detectable glucosinolate

consistent with previous published results in field pennycress seed or meals and

total glucosinolate concentration in CCWG-15-32 ranged from 45 to 102 µmoles/g

on 100% DM basis. Total glucosinolate concentration in B3WG-15-32 ranged

from 101.4 to 110.7 µmoles/g on 100% DM basis

●    Total copper levels achieved target levels of 800 ppm

●      Other anti-nutrients (sinapine) and mycotoxins were below the imit of detection or

447           quantification

The main purpose of CoverCress whole grain as a feed ingredient in animal diets (e.g.

broilers) is as an energy source. Determination of energy value is complex, but is related to the

fat, protein, and fiber content. Further, the degree of unsaturation of the fatty acids is a factor in

energy. While there are some slight differences in total fat and the fiber profile, the overall

nutritional value of CCWG for energy is expected to be similar to that of canola seed (see Table

18 below). Levels of the four key limiting amino acids in broiler diets, methionine, lysine,

threonine and valine are similar between CCWG and canola. The low-erucic acid phenotype

expected with loss of function of FAE1 was achieved as shown by the low to non-detectable

levels in CCWG. The nutritional value in terms of energy (total fat, profile of fatty acids, crude

fiber, ADF, NDF) of B3WG is comparable to that of CCWG.  Crude fiber levels were variable

across lots and between CCWG and B3WG. This may be due to assay or natural variability

within the plants. Levels of the four key limiting amino acids in broiler diets, methionine, lysine,

threonine and valine were lower for the B3WG as compared to the CCWG.  The difference is

mostly like due to the lower amino acids as a percent of protein in the B3WG as compared to the

CCWG (88.1% versus 97% of the protein for B3WG and CCWG, respectively.

Table 18**.** Key Compositional Parameters for Canola Whole Grain, CCWG and B3WG

| Component | Canola[1] | CCWG-15-32 Mean[2] [range] | B3WG-15-32 Mean[2] [range] |
|---|---|---|---|
| Fat % DM | 39.1 | 36.3 [35.5-36.7] | 37.0 [35.7-38.5] |
| Erucic acid % TFA | <0.1 | <0.1 | <0.1 |
| Oleic % of TFA | 61.4 | 40.07 [38.61-41.06] | 45.4 [42.22-48.16] |
| Linoleic % TFA | 20.1 | 34.48 [34.17-34.94] | 31.89 [29.84-33.87] |
| Linolenic % TFA | 9.3 | 16.55 | 17.18 |

| | | | |
|---|---|---|---|
| | | [16.18-17.67] | [16.44-18.26] |
| Total saturated %TFA | 7.0 | 5.0 | |
| Total mono-unsaturated %TFA | 64.4 | 41.6 | |
| Total poly-unsaturated % TFA | 28.6 | 53.4 | |
| | | | |
| Protein %DM | 24.5 | 25.5 [22.7-26.7] | 23.9 [22.5-25.1] |
| Methionine % protein | 1.93 | 1.88 [1.74-2.03] | 1.39 [1.33-1.45] |
| Lysine % protein | 5.66 | 5.51 [5.30-5.68] | 4.71 [4.55-4.85] |
| Threonine % protein | 3.97 | 4.91 [4.62-5.26] | 4.44 [4.36-4.58] |
| | | | |
| Valine %protein | 4.40 | 5.69 [5.48-6.14] | 4.89 [4.59-5.04] |
| Crude Fiber %DM | 13.3 | 23.0 [20.9-24.9] | 12.1 [10.9-12.5] |
| ADF | 18.1 | 13.9 [12.1-15.8] | 17.8 [15.9-20.2] |
| NDF | 25.7 | 20.3 [17.3-24.5] | 22.6 [21.3-24.4] |
| | | | |
| Sulfur % DM | 0.45[3] [0.36-0.55] | 0.87 [0.47-1.05] | 0.94 [0.86-1.01] |
| Total glucosinolates μmoles/g on DM basis UV method | 9.8 | 86.9 [45-102] | 105.4 [101.4-110.7] |

[1]Values from Dairy One feed Composition Laboratory; means of whole canola seed samples from crop years 2004 -

2020. https://www.dairyoneservices.com/feedcomposition/

[2]From the respective 5-lot study, DLL values

[3]From https://www.grainscanada.gc.ca/en/grain-research/export-quality/oilseeds/canola/2020/index.htm

The composition and nutritional components of C3WG-15-32 lots were generally

consistent with very good process control and inter-lot variability. These results will enable

eventual development of permissible ranges for key nutritional analytes and guaranteed levels for

specific components (i.e. commercial specifications) like total crude protein, crude fiber, crude

fat, copper and sulfur.  A glucosinolate specification will be needed to ensure safe consumption

depending on the level of inclusion in the diet and sensitivity of the animal species. In these lots,

the mean level of glucosinolates was slightly higher in B3WG than CCWG, but ranges for the

two overlapped, indicating the difference is likely due to natural variability.

The current study shows that low-erucic acid, lower-fiber pennycress (CoverCress)

produces a consistent compositional phenotype that may enable the seed to be consumed as an

energy source for various animal species.

**Acknowledgements**

The authors would like to acknowledge Shengjun Liu (CoverCress Inc; St. Louis, MO) for

treatment of the CoverCress grain with copper sulfate solution, Margaret Nemeth (Statistical

Consultants Plus; Fenton, MO) for statistical compilation and CoverCress Inc. (St. Louis MO)

for financial support.

**Conflict of Interest**

Shawna Lemke and Gary Hartnell were paid by CoverCress Inc for their consulting services.

Chris Aubach is an employee of CoverCress Inc.

**References**

AAFCO. (2017). Critical factors in determining fiber in feeds and forages.
https://www.aafco.org/Portals/0/SiteContent/Laboratory/Fiber_Best_Practices_Working_Group/
Fiber-Critical-Conditions-Final.pdf

Abraham, V., Bhatia, C.R. (1986). Development of strains with yellow-seed coat in Indian
mustard (Brassica juncea Czern & Coss). Plant Breed. 97, 86-88.

Aghajanzadeh, T., Hawkesford, M. J., & De Kok, L. J. (2014). The significance of
glucosinolates for sulfur storage in Brassicaceae seedlings. Frontiers in Plant Science, 5, 704.
https://doi.org/10.3389/fpls.2014.00704

Baudry, A., Heim, M. A., Dubreucq, B., Caboche, M., Weisshaar, B., Lepiniec, L. (2004). TT2,
TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin
biosynthesis in Arabidopsis thaliana. Plant J., 39(3), 366-80. doi: 10.1111/j.1365-
313X.2004.02138.x. PMID: 15255866.

Bischoff, K. L. (2021). Glucosinolates. Nutraceuticals, 903–909. https://doi.org/10.1016/b978-0-
12-821038-3.00053-7

Bremer. J. and K. R. Norum. (1982). Metabolism of very long-chain monounsaturated fatty acids
(22: 1) and the adaptation to their presence in the diet J. Journal of Lipid Research Volume 23,
243-256.

Canola Council of Canada. (2015). Canola meal feeding guide. Feed Industry Guide (5th
edititon). https://www.canolacouncil.org/canolamazing/wordpress/wp-
content/uploads/2016/08/2015-Feed-Guide.pdf

Canola Council of Canada. (2019). Canola meal feeding guide. Feed Industry Guide (6th
edition). https://www.canolacouncil.org/canolamazing/wordpress/wp-
content/uploads/2019/10/2019-Canola-Meal-Feed-Guide-Web.pdf

Chopra, R., Johnson, E. B., Daniels, E., McGinn, M., Dorn, K. M.,Esfahanian, M., Folstad, N.,
Amundson, K.,  Altendorf, K., Betts, K., Freis, K., Anderson, J. A., Wyse, D.L., Sebdrook, J.C.
and David Marks, M. (2018). Translational genomics using Arabidopsis as a model enables the
characterization of pennycress genes through forward and reverse genetics. Plant J. 96(6),1093-
1105. Doi: 10.1111/tpj,14147.PMID: 30394623.

Chopra, R., Johnson, E. B., Emenecker, R., Cahoon, E. B., Lyons, J., Kliebenstein, D. J.,  …
David Marks, M. (2020). Identification and stacking of crucial traits required for the
domestication of pennycress. Nature Food, 1(1), 84–91. https://doi.org/10.1038/s43016-019-
0007-z

Debeaujon, I., Nesi, N., Perez, P. (2003). Proanthocyanidin-accumulating cells in Arabidopsis
testa: regulation of differentiation and role in seed development. *Plant Cell*, 15(11), 2514-2531.
doi:10.1105/tpc.014043

Fenwick, G. R. (1982). The assessment of a new protein source--rapeseed. The Proceedings of
the Nutrition Society, 41(3), 277–288.

Gonzalez, A., Mendenhall, J., Huo, Y., and Lloyd, A. (2009). TTG1 complex MYBs, MYB5 and
TT2, control outer seed coat differentiation. *Dev. Biol.* 325, 412–421. doi:
10.1016/j.ydbio.2008.10.005

Huisman, J., & Tolman, G. H. (2001). Antinutritional factors in the plant proteins of diets for
non-ruminants. Recent developments in Pig Nutrition 3, 261–291.

Jerroch, J., Jankowski, J., Schone, F. (2008). Rapeseed products in the feeding of broiler and
laying hens. © Verlag Eugen Ulmer, Stuttgart Arch.Geflügelk. 72 (2). S. 49–55, 2008, ISSN
0003-9098.

Lepiniec, L., Debeaujon, I., Routaboul, J. M., Baudry, A., Pourcel, L., Nesi, N., Caboche, M.
(2006). Genetics and biochemistry of seed flavonoids. Annu Rev Plant Biol., 57:405-30. doi:
10.1146/annurev.arplant.57.032905.105252. PMID: 16669768.

McGinn M, Phippen WB, Chopra R, Bansal S, Jarvis BA, Phippen ME, Dorn KM, Esfahanian
M, Nazarenus TJ, Cahoon EB, Durrett TP, Marks MD, Sedbrook JC. Molecular tools enabling
pennycress (Thlaspi arvense) as a model plant and oilseed cash cover crop. Plant Biotechnol J.
2019 Apr;17(4):776-788. doi: 10.1111/pbi.13014. Epub 2018 Oct 25. PMID: 30230695;
PMCID: PMC6419581.

Payvastagan, S., Farhoomand, P., Shahrooze, R., Delfani, N., & Talatapeh, A. (2012). The
effects of different levels of canola meal and copper on performance, susceptibility to ascites and
plasma enzyme activities in broiler chickens. Ann. Biol. Res, 3(11), 5252-5258.

Phippen, W.B., & Phippen, M.E.. 2012. Soybean seed yield and quality as a response to field
pennycress residue. Crop Science 52, 2767-2773. https://doi.org/10.2135/cropsci2012.03.0192

Rymer, C., & Short, F. (2003). The nutritive value for livestock of UK oilseed rape and rapeseed
meal. Home-Grown Cereals Authority. Retrieved from
http://cereals.ahdb.org.uk/media/376883/rr_os14_-_complete_final_report.pdf

SAS Software Release 9.4 (TS1M7). Copyright© 2016 by SAS Institute Inc., Cary NC.

Schöne, F., Jahreis, G., Richter, G., & Lange, R. (1993). Evaluation of rapeseed meals in broiler
chicks: effect of iodine supply and glucosinolate degradation by myrosinase or copper. Journal of
the Science of Food and Agriculture, 61(2), 245-252.

Sedbrook, J. C., Phippen, W. B., & Marks, M. D. (2014). New approaches to facilitate rapid
domestication of a wild plant to an oilseed crop: example pennycress (Thlaspi arvense L.). Plant
Science, 227, 122-132.

Singhal, K. K., & Sinha, A. K. (2000). Effect of copper sulphate treatment of mustard cake on its
glucosinolate content, palatability and nutrient utilization in kids. Indian Journal of Dairy
Science, 53(3), 210-215.

Slominski, A. B., Campbell, L. D., Guenter, W. (1994). *Journal of Agricultural and Food*
*Chemistry*, *42*(3), 704-707. DOI: 10.1021/jf00039a020

Slominski, A. B. (2015). Nutritive Value of Canola Meal: The Dietary Fibre Story. University of
Manitoba. https://www.agwest.sk.ca/IRC2015/BSlominskiCanolameal.pdf
592
593