Cortical-brainstem interplay during speech perception in older adults with and without hearing loss

Jesyin Lai[1,2,3], Claude Alain[4], Gavin M. Bidelman[1,2,5,6,*]

[1] Institute for Intelligent Systems, University of Memphis, Memphis, TN, USA

[2] School of Communication Sciences and Disorders, University of Memphis, Memphis, TN, USA

[3] Diagnostic Imaging Department, St. Jude Children's Research Hospital, Memphis, TN, USA

[4] Rotman Research Institute, Baycrest Centre for Geriatric Care, Toronto, ON, Canada

[5] Department of Speech, Language and Hearing Sciences, Indiana University, Bloomington, IN, USA

[6] Program in Neuroscience, Indiana University, Bloomington, IN, USA

* Correspondence:
Corresponding Author
gbidel@indiana.edu (G.M.B.)

Keywords: age-related hearing loss, brainstem, frequency-following responses, arousal, cortical
α, EEG

Abstract

Real-time modulation of brainstem frequency-following responses (FFRs) by online changes in

cortical arousal state via the corticofugal (top-down) pathway has been demonstrated previously in

young adults and is more prominent in the presence of background noise. FFRs during high cortical

arousal states also have a stronger relationship with speech perception. Aging is associated with

increased auditory brain responses, which might reflect degraded inhibitory processing within the

peripheral and ascending pathways, or changes in attentional control regulation via descending

auditory pathways. Here, we tested the hypothesis that online corticofugal interplay is impacted by

age-related hearing loss. We measured EEG in older adults with normal-hearing (NH) and mild to

moderate hearing-loss (HL) while they performed speech identification tasks in different noise

backgrounds. We measured α power to index online cortical arousal states during task engagement.

Subsequently, we split brainstem speech-FFRs, on a trial-by-trial basis, according to fluctuations in

concomitant cortical α power into low or high α FFRs to index cortical-brainstem modulation. We

found cortical α power was smaller in the HL than NH group. In NH listeners, α-FFRs modulation for

clear speech (i.e., without noise) also resembled that previously observed in younger adults for speech

in noise. Cortical-brainstem modulation was further diminished in HL older adults in the clear

condition and by noise in NH older adults. Machine learning classification showed low α FFR

frequency spectra yielded higher accuracy for classifying listeners' perceptual performance in both NH

and HL participants. Moreover, low α FFRs decreased with increased hearing thresholds at 0.5-2 kHz

for clear speech but noise generally reduced low α FFRs in the HL group. Collectively, our study

reveals cortical arousal state actively shapes brainstem speech representations and provides a potential

new mechanism for older listeners' difficulties perceiving speech in cocktail party-like listening

situations in the form of a miss-coordination between cortical and subcortical levels of auditory

processing.

1   Introduction

Declines in auditory processing (Poth et al., 2001; Parthasarathy et al., 2010; Kortlang et al.,

2016; Lai and Bartlett, 2018) and speech comprehension (Anderson et al., 2012) (Schneider et al.,

2005; Peelle et al., 2010)—especially in the presence of background noise (Dubno, 1984; Takahashi

and Bacon, 1992; Souza et al., 2007; Anderson et al., 2011; Song et al., 2011; Jin et al., 2014; Presacco

et al., 2016; Vermeire et al., 2016)—are ubiquitous during aging and age-related hearing loss. Age-

related declines in the sensory (auditory) system (Parthasarathy and Bartlett, 2011, 2012; Fostick et al.,

2013; Parthasarathy et al., 2014, 2016; Lai and Bartlett, 2015, 2018; Lai et al., 2017)[e.g., age-related

impairments in sound source segregation (Alain et al., 1996; Alain and McDonald, 2007; Gallun and

Best, 2020)], changes in cognitive function (Park et al., 2003), or a combination of both (Pichora-Fuller

and Singh, 2006; Wayne and Johnsrude, 2015; Wingfield et al., 2016) could lead to listening and

comprehension difficulties in elderly listeners.

Evidence suggests that speech-in-noise (SiN) problems could be related to dysfunctional

connections and changes in speech processing between cortical and subcortical levels of the auditory

system that emerge with age and age-related hearing impairment. It is well established that SiN

processing can be affected by many factors, such as attention (Saiz-Alía et al., 2019; Price and

Bidelman, 2021) and arousal state (Mai et al., 2019; Saderi et al., 2021). Several studies have shown

neural correlates of these phenomena. For example, findings from EEG studies on emotion suggest

that power in the cortical α band (8-12 Hz) is a useful indicator of arousal state (Aftanas et al., 2002;

Uusberg et al., 2013). Moreover, parieto-occipital α power was shown to index cognitive processing,

effortful listening (Wöstmann et al., 2015; McMahon et al., 2016; Dimitrijevic et al., 2017), the state

of wakefulness (Pfurtscheller et al., 1996) and top-down processing (Henry et al., 2017). Alpha

oscillatory activity has also been associated with adaptive, intentional, and top-down suppression of

task-irrelevant information (Rihs et al., 2007; Jensen and Mazaheri, 2010; Händel et al., 2011; Klatt et

al., 2020). Increased α power has been proposed to index inhibitory processing across sensory

modalities (Klimesch et al., 2007; Weisz et al., 2007, 2011; Strauß et al., 2014), while decreased α

oscillations is thought to facilitate sensory processing or neural firing (Haegens et al., 2011; Klatt et

al., 2020). There is, however, no consensus regarding the mechanisms underlying α oscillations

reported in these studies. This could be due to the fact that in most studies α is treated as a unitary

measure rather than reflecting different underlying processes. Meanwhile, evidence suggests that

cortical α oscillations changes with aging (Yordanova et al., 1998; Böttger et al., 2002), such as a

decrease in α frequency (Chiang et al., 2011) and reduced spontaneous entrainment of resting-state α

oscillations (Gaál et al., 2010). Studying α power during SiN perception in older adults may reveal the

impacts of aging in top-down attentional control that help facilitate the processing of target vs. non-

target sounds, thus providing insight concerning why cocktail party-like situations are more difficult

in older listeners (Pichora-Fuller et al., 2017).

In addition to cortical changes, age-related declines in speech coding have been widely

observed at subcortical levels of the auditory system, both in terms of local processing within the

brainstem but also its functional signaling to and from the cortex (Bidelman et al., 2019). In young

adults, we recently observed that speech-evoked brainstem frequency-following response (FFR)

amplitude varied as a function of α power (Lai et al, 2022). That is, low FFR amplitude coincided with

low α power whereas high FFR amplitude was associated with high α states. Notably, low α FFRs

correlated positively with behavioral response times (RTs) for speech discrimination and more

accurately decoded the input speech stimuli as revealed by neural classifiers. Extending this approach

here to address questions of auditory aging, we analyzed neuroelectric FFRs recorded during active

speech perception in age-matched older adults with normal (NH) or mild hearing loss (HL). This

allowed us to investigate the effects of age-related hearing loss on cortical α state and its modulation

of brainstem speech processing in real-time. We aimed to determine the nature of auditory cortical-

brainstem interplay in older adults, and more critically, whether such online corticofugal engagement

during SiN listening is altered due to hearing loss, as suggested in prior work (Bidelman et al., 2019).

Our collective results reveal that brainstem speech-FFRs are dynamically modulated by fluctuations in

cortical α state in normal-hearing listeners but this cortical-subcortical interplay declines in age-related

hearing loss.

2   Methods and Materials

2.1   Participants

Detailed information on participants, informed consent, and demographics are reported in our

original report detailing age-related changes in the brainstem and cortical evoked potentials (Bidelman

et al., 2019). New analyses reported herein examine online changes in FFRs as a function of the

simultaneous cortical state. All participants had no reported history of neurological or psychiatric

illness. Participants were aged between 52 to 75 (69 ± 5.8 years; 16/16 M/F). There were divided into

normal (NH) and hearing-impaired (HL) groups based on their pure-tone audiometry hearing

thresholds. We used 25 dB HL as the cutoff to define normal hearing, which has been used in the clinic

since 1970s (Gatlin and Dhar, 2021). NH listeners (n = 13) listeners had average thresholds (250-8000

Hz) better than 25 dB HL across both ears whereas HL listeners (n = 19) had average thresholds poorer

than 25 dB HL. The pure-tone averages (PTAs) (i.e., mean of 500, 1000, 2000 Hz) of NH listeners

were ~10 dB better than in HL listeners (mean ± SD; NH: 15.3 ± 3.27 dB HL, HL: 26.4 ± 7.1 dB HL;

$t_{2.71}$ = -5.95, p < 0.0001; NH range = 8.3-20.83 dB HL, HL range = 15.8-45 dB HL). Both NH (t_{12} =

0.15, p = 0.89) and HL (t_{18} = -2.02, p = 0.06) groups otherwise had symmetric PTA between ears. Both

NH and HL groups had elevated hearing thresholds at very high frequencies (≥ 8000 Hz), which is

typical of age-related presbycusis in older adults. Besides hearing, the two groups were matched in age

(NH: 66.2 ± 6.1 years, HL: 70.4 ± 4.9 years; $t_{2.22}$ =-2.05, p = 0.052) and sex balance (NH: 5/8 M/F;

HL: 11/8; Fisher's exact test, p = 0.47). Age and hearing loss were not correlated (Pearson's r = 0.29, p

= 0.10), suggesting these factors of aging were largely independent in our sample.

2.2   QuickSiN Test

The Quick Speech-in-Noise (QuickSiN) test was used to measure listeners' speech reception

thresholds in noise (Killion et al., 2004). A list of six sentences with five keywords per sentence spoken

by a female talker in a background of four-talker babble noise was heard by listeners during the test.

Target sentences were presented at 70 dB sound pressure level (SPL) (binaurally) at signal-to-noise

ratios (SNRs) decreasing from 25 dB (relatively easy) to 0 dB (relatively difficult) in 5 dB steps. The

number of keywords correctly recalled was logged, and a score was computed for each listener. The

SNR-loss score indexes the difference between a listener's SNR-50 (i.e., the SNR required to identify

50 % of the keywords correctly) and the average SNR threshold for normal-hearing adults (i.e., 2 dB)

(Killion et al., 2004). A higher score reflects poorer SiN perception. Each listener's SNR-loss score

was averaged from four lists of sentence presentations. In this study, NH listeners' scores ranged from

-0.25 to 2.5 dB of SNR-loss (M = 1.1, SD = 0.8) while HL listeners' scores ranged from -2.5 to 8.5 of

SNR-loss (M = 2, SD= 2.5) (see Fig. 1D in Bidelman et al., 2019).

2.3   EEG Stimuli and Task

The stimuli and task are described fully in Bidelman et al. (2019) and illustrated here in Fig. 1A. Three

naturally produced English consonant-vowel phonemes (/ba/, /pa/ and /ta/), from the standardized

UCLA version of the Nonsense Syllable Test (Dubno and Schaefer, 1992), were generated by a female

talker. The duration of each phoneme was 100-ms and the average root mean square SPL of each

phoneme matched. All three tokens had a common voice fundamental frequency (mean F0 = 150 Hz),

first and second formants (F1 = 885, F2 = 1389 Hz). The resulting stimulus-evoked response (i.e., FFR)

should predominantly originate from the subcortex (Brugge et al., 2009; Bidelman, 2018) since the

stimulus F0 is above the phase-locking limit of the cortical neurons and "cortical FFRs" (Coffey et al.,

2016; Bidelman, 2018; Bidelman and Momtaz, 2021). Speech tokens were delivered binaurally to

listeners in either clear (i.e., no noise) or noise-degraded conditions. A complete set of stimulus

presentation in each condition contained a total of 3000 /ba/, 3000 /pa/, and 210 /ta/tokens (spread

evenly over 3 blocks to allow for breaks). The interstimulus interval between tokens was randomly

jittered within the range of 95-155 ms (5ms steps, uniform distribution). The /ba/ and /pa/ tokens were

presented more frequently than the /ta/ token in a pseudo-random manner such that at least two frequent

tokens intervened between infrequent tokens. The infrequent /ta/ token was denoted as the target in

which listeners were required to provide a response by pressing a button on the computer whenever

they detected it. Both reaction time (RT) and detection accuracy (%) were recorded. For the noise-

degraded condition, the same procedures as the clear condition were repeated, but the tokens were

presented in an identical speech triplet mixed with eight talker noise babble (Killion et al., 2004) at a

signal-to-noise ratio (SNR) of 10 dB. There were 6 blocks (3 clear & 3 noise) collected from each

participant. Having the clear and noise conditions allowed us to compare behavioral performance in

different backgrounds and evaluate the impact of noise on speech perception in NH vs. HL listeners,

respectively. The task ensured that listeners were actively engaged during speech perception and online

EEG recording. Stimuli were controlled by a MATLAB program (The Mathworks, Inc.; Natick, MA)

routed to a TDT RP2 interface (Tucker-Davis Technologies; Alachua, FL) and delivered binaurally

through insert earphones (ER-3; Etymotic Research; Elk Grove Village, IL). The speech stimuli were

presented at 75 dB SPL (noise at 65 dB SPL) with alternating polarity.

2.4   EEG recording and preprocessing

The neuroelectric activity was recorded from 32 channels at standard 10-20 electrode locations

on the scalp (Oostenveld and Praamstra, 2001) during the target speech detection task. Electrode

impedances were $\leq$ 5 kΩ. EEGs were digitized at 20 kHz using SynAmps RT amplifiers

(Compumedics Neuroscan; Charlotte, NC). After EEG acquisition, the data were processed using the

'mne' package in Python 3.9.7. EEG data were re-referenced offline to the mastoids (TP9/10) for

sensor (channel-level) analyses. For source analysis of brainstem FFRs, we used a common average

reference prior to source transformation (detailed below). Responses were then filtered 100-1000 Hz

[finite impulse response (FIR) filters; hamming window with 0.02 dB passband ripple, 53 dB stopband

attenuation, -6 dB cutoff frequency] to further isolate brainstem activity (Musacchia et al., 2008;

Bidelman et al., 2013).

2.5   Derivation of Source FFRs and Cortical Activities

The derivation of source FFR waveforms and isolation of cortical activities are similar to the

methods described in Lai et al. (2022) for young adults. The 32-ch sensor data were transformed into

source space using a virtual source montage. The source montage comprised of a single regional source

(i.e., current flow in x, y, z planes) positioned in the brainstem and midbrain (i.e., inferior colliculus)

[details refer to (Bidelman, 2018b; Bidelman and Momtaz, 2021; Price and Bidelman, 2021)]. Source

current waveforms (SWF) from the brainstem source were obtained using the formula: $SWF = L^{-1}$ x

FFR, where L is the brainstem source leadfield matrix (size 3x64) and FFR is the 32-ch sensor data

(64 x NSamples). This essentially applied a spatial filter to all electrodes that calculated their weighted

contribution to the scalp-recorded FFRs in order to estimate source activity within the midbrain in the

x, y, and z directions (Scherg and Ebersole, 1994; Scherg et al., 2002). This model explains ~90 % of

the scalp-recorded FFR (Bidelman et al., 2019; Price and Bidelman, 2021). Only the z-oriented SWF

was used for further analysis (x and y SWFs were not analyzed) given the predominantly vertical

orientation of current flow in the auditory midbrain pathways relative to the scalp [x- and y-orientations

contribute little to the FFR (Bidelman, 2018b)].

We isolated cortical α band activity from the EEG and used it as a running index of arousal

state (high or low) during the target speech detection task. EEG at the Pz and Oz channels were filtered

at 8-12 Hz (FIR filters, -6 dB cutoff frequency at 7 Hz and 13.5 Hz) and averaged (i.e., equivalent to

POz) to obtain cortical α-band activity at a posterior scalp region. Filtered α activities were epoched

with a time window of 195 ms (-50 to 145 ms in which 0 ms corresponded to the onset of a /ba/ or /pa/

token) to capture approximately 1-2 cycles of α band. This epoch window encapsulated the entirety of

the evoked FFR within the immediate trial with no spillover from the preceding or subsequent trial(s).

Infrequent /ta/ tokens were excluded from analysis due to their limited trials. The root mean square

(RMS) amplitude of single trial α activity was computed to quantify cortical arousal level over the

duration of the target speech detection task. We then normalized RMS values to the median of RMS

values of each run, respectively. Next, the distribution of trial-wise normalized α RMS was visualized

using a histogram. We categorized trials of each participant per condition falling within the 0-35th

percentile as "low α" power and those falling within the 65-100th percentile as "high α" power. This

categorization was used because it provided ~2100 trials for each low or high α power in each

participant per condition, which is reasonable to obtain an average FFR with robust signal-to -noise

ratio (i.e., ≥3 dB SNR) (Bidelman, 2018a). More detailed information of this methodology can be

found in Lai et al. (2022) (see their Fig. 2). We similarly measured cortical activity in another frequency

band (e.g., β band; 18-22 Hz) from the same location (i.e., POz β) and α band from a different electrode

site (i.e., Fz α). These control analyses allowed us to ensure that the observed changes in speech-evoked

FFRs were specifically associated with cortical arousal level (indexed by α power) rather than general

fluctuations in the EEG, per se.

2.6   Analysis of Brainstem FFRs

We categorized source FFRs based on whether α amplitude in the same epoch was either high

or low power, thus deriving FFRs according to the trial-by-trial cortical state. Source FFR waveforms

(from the z-orientated dipole) were averaged for each α category and noise condition per participant.

Subsequently, we analyzed the steady-state portion (10–100 ms) of FFR waveforms using the FFT

(Blackman window; 11.1 Hz frequency resolution) to capture the spectral composition of the response.

F0 amplitude was measured as the peak spectral maximum within an 11 Hz bin centered around 150

Hz (i.e., F0 of the stimuli). To compare FFR F0 amplitudes during low vs. high α power, a normalized

(within-subject) F0 ratio was calculated as follows:

208
$$F0\ ratio = \frac{F0amp_{high\alpha}}{F0amp_{low\alpha}} \tag{1}$$

209   Where F0 ratios > 1 indicate stronger brainstem FFRs during states of high cortical α power

210   and F0 ratios < 1 indicate stronger FFRs during states of low cortical α power (Lai et al., 2022).

2.7   Statistical Analysis

We used rmANOVAs to compare brainstem F0 ratios among the clear vs. noise condition, and

NH vs. HL group. Multiple pairwise comparisons (Mann-Whitney U test with Bonferonni corrections)

between the NH and HL groups were performed using the 'pingouin' package in Python. One sample

t-tests ('scipy' package in Python) were also used to evaluate whether FFR F0 ratios were significantly

different from 1 (and thereby indicated the significance of α modulation). Wilcoxon signed-rank test

was used when comparing raw F0 amplitudes at low vs. high α power within participants for each SNR

condition and hearing group. To compare differences in α RMS values of all participants across (clear

vs. noise) conditions, we performed post-hoc Conover's test ("scikit_posthocs" package in Python),

which is a non-parametric pairwise test, with Bonferroni adjustment. To assess differences in raw F0

amplitudes (log-transformed) across factors for NH and HL groups, we first performed a 2 x 2 x 2

(SNR x α power x hearing group) mixed model (participants = random factor) ANOVA ('lme4'

package in Rstudio). Following a significant interaction, we then ran separate 2 x 2 (α power x SNR)

mixed-model ANOVAs for the NH and HL groups, respectively. Initial diagnostics were performed

using residual and Q-Q plots to assess the heteroscedasticity and normality of data. F0 amplitudes were

log-transformed to improve normality and homogeneity of variance assumptions. Effect sizes are

reported as η_p^2. To check if low or high α FFR F0 amplitudes were associated with behavioral

performance (i.e., QuickSiN, PTA, %- correct, and RTs) by pooling all NH and HL participants, we

performed Spearman's correlations ('scipy' package in Python) to test their pairwise correlations of

brain and behavior measures. Spearman's correlation was used because these measures were found to

be not normally distributed ($p < 0.05$) from the test of normality using Shapiro-Wilk test.

2.8   Classification of Performance Level from FFR Frequency Spectra via Machine Learning

All participants' performance in the clear condition was categorized into three levels (poor, average,

and good) based on their percent correct of /ta/ detections. Participants with poor performance had %-

correct ≤ 30th-percentile of overall %-correct while good-performance participants had %-correct ≥

70th-percentile of overall %-correct. Subsequently, we classified poor- and good-performing

participants using frequency spectra of their low or high α FFRs and a support vector machine (SVM)

classifier (kernel = radius basis function, C=1000, gamma = scale) in the 'scikit-learn' package in

python. Due to the limitation in sample size (32 total observations), we were only able to perform a

two-group rather than three-group machine learning (ML) classification. There were ten participants

falling into either poor- or good-performance category, which provided twenty participants' frequency

spectra to be used as data input for SVM. Frequency spectra were obtained from the FFT of average

FFR waveforms (10 to 100 ms steady state portion) across ~2100 trials of low or high α FFRs per

participant. The absolute amplitudes of frequency spectra, which consisted of 226 amplitude-by-

frequency points, were the input features; performance level (i.e., poor vs. good) served as the ground

truth class labels. The ML classification procedures on FFRs were similar to those described by Xie et

al. (2009) and Lai et al. (2022). During one iteration of training and testing, a four-fold cross-validation

approach was used to train and evaluate the performance of the SVM classifier to obtain a mean

classification accuracy [Fig. 1 of Xie et al. (2009)]. In this process, a 4-fold stratification was performed

to randomly and equally divide participants into 4 subgroups with 5 unique participants (almost similar

number of poor- and good-performing participants) in each subgroup. Three of the 4 subgroups were

selected as the training data while the remaining subgroup was used as the hold-out testing data. To

mitigate the problem of imbalanced participant numbers in the two classes of training set, we randomly

over-sampled the minority class using the 'imblearn' package in python. These steps were repeated

within each iteration so that each subgroup was held-out as the test data whereas the other 3 subgroups

were used to train the SVM classifier. Mean classification accuracy of poor- vs. good-performance was

calculated across cross-validated iterations. We performed a total of N=5000 iterations to examine

group classification for low vs. high α power FFRs.

To evaluate if the classifier accuracy (mean of N=5000 iterations) was statistically significant,

we randomly shuffled the 226 data points of frequency spectra in each participant, and the same training

and testing procedures described above were repeated to derive a null distribution of classification

accuracies. We then calculated the p-value to determine the statistical significance of "true" classifier

performance using the formula described in Phipson and Smyth (2010): $p = (a + 1)/(n + 1)$, where a

is the number of classification accuracies from the null distribution that exceeds the median of the

actual distribution of classification accuracies and n is the total number of classification accuracies

from the null distribution.

2.9   Fitting Linear Regression Models with Brain and Behavior Measures

To compare the changes in low α FFR F0 amplitudes as behavior performance changed in the

NH and HL groups, we fitted linear regression models for pooled SNR, clear and noise conditions,

respectively, by using behavior performance and group as main factors, and their interaction factor.

$$Y = \beta_0 + \beta_1 \cdot X + \beta_2 \cdot group + \beta_3 \cdot X \cdot group \qquad (2)$$

Y represents FFR F0 amplitudes during low α power, X is one of the behavior measures

(QuickSiN, PTA, %- correct, or RTs), and *group* is the dummy variable for NH (*group=0*) and HL

(*group=1*). Meanwhile, β_0 is a constant and β_1, β_2, or β_3 is the coefficient or slope for the respective

variable. After fitting regression models, we checked assumptions for the normality of residuals,

homoscedasticity of residuals and linearity of the models. For residual normality, we tested if model

residuals were normally distributed using Shapiro-Wilk test and visualized the Q-Q plots of the

residuals. For residual homoscedasticity, we used Breusch-Pagan and Goldfeld-Quandt tests. For

model linearity, we visualized residual distributions by plotting residuals vs. predicted values. Result

interpretation was then carried out after confirming the above three assumptions.

The fitted linear regression models allowed us to study if the slopes (i.e., the change of FFR

amplitudes) of the NH group was significant as well as if the slope of the HL group was significantly

different from the NH group. In the NH group, i.e., *group=0*, Eq. (2) can be written as $Y = \beta_0 + \beta_1 \cdot X$

and β_1 is the slope for the NH group. In contrast, in the HL group, i.e., *group=1*, Eq. (2) can be written

as $Y = \beta_0 + \beta_1 \cdot X + \beta_2 + \beta_3 \cdot X = (\beta_0 + \beta_2) + (\beta_1 + \beta_3)X$, where $(\beta_1 + \beta_3)$ is the slope for the HL group and

β_3 represents the difference in slope between the HL and NH group.

3   Results

3.1   Behavior performance of target speech detection

Behavioral responses during the EEG task (%-correct /ta/ detections vs. RTs) showed a negative

correlation for the clear but not noise condition (Spearman's r = -0.45, p = 0.01, Fig. 1B); participants

with slower response speeds showed poorer speech detection accuracies. This is consistent with

previous findings showing negative associations between hit responses and RTs in younger listeners

(e.g., Lai et al. 2022). When separated into the NH and HL groups, we found a negative relationship

between behavioral hit responses and decision speeds but only in the HL group (Fig. 1C).

3.2   Cortical α band and brainstem speech-FFRs

Differences in cortical α-band amplitudes during low vs. high α states were prominent at the

single participant (Fig. 2A) as well as group level (Fig. 2C). Spectral differences in the corresponding

brainstem FFR for these same low vs. high cortical trials were also notable (Fig. 2B). Cortical α (both

low and high levels) was overall higher in the NH listeners (p<0.01, non-parametric post-hoc

Conover's test with Bonferroni adjustment) but both groups showed clear separability of "low" vs.

"high" α states during the speech detection task.

In response to clear speech (Fig. 2D), brainstem F0 ratios (indexing cortical α-related FFR

enhancement) in the NH group were significantly higher than 1 (t_{12} = 2.64, p = 0.02, 1-sample t-test)

and higher than the HL group overall (U = 192, p = 0.01, Mann-Whitney U test). In response to the

noise-degraded speech, this cortical-FFR enhancement was observed in the HL group (t_{18} = 2.67, p =

0.02, 1-sample t-test), but was not significantly different than the FFR enhancement observed in the

NH group. Repeating the same analysis for both controls (POz β and Fz α) revealed no difference in

308 F0 ratios for the NH vs. HL group (Fig. 1 in Supplementary), indicating cortical modulation of the FFR

309 was restricted to POz α.

310  A 3-way mixed-model ANOVA performed on log F0 amplitudes revealed a significant SNR x

311 α power x group interaction ($F_{1,96} = 9.5$, $p = 0.003$, $\eta_p^2 = 0.09$) (Fig. 2E). To make sense of this complex

312 interaction, we performed separate 2-way (SNR x α power) mixed-model ANOVAs by hearing group.

313 The SNR x α power interaction was significant in both the NH ($F_{1,39} = 5.17$, $p = 0.03$, $\eta_p^2 = 0.12$) as

314 well as the HL group ($F_{1,57} = 4.11$, $p = 0.05$, $\eta_p^2 = 0.07$). Though the comparison of effect sizes suggests

315 this interaction was stronger in NH listeners, the interaction was distinct in direction compared to the

316 HL group. In the NH group, FFR F0 amplitudes were significantly higher during high α power for clear

317 speech. This pattern was dampened and reversed in the HL group.

318 **3.3 Brain-behavior relations in both the NH and HL groups**

319  We next assessed associations between low or high α FFR F0 amplitudes) and behavioral

320 performance (QuickSiN, PTA, %-correct, and RTs) by performing Spearman's correlation analyses.

321 The positive association between F0 amplitudes and percent correct was significant at low but not high

322 α power (Spearman's $r = 0.3$, $p = 0.02$, Fig. 3A) while the associations between F0 amplitudes (either

323 during low or high α power) and other behavior measures were not significant. To further assess if

324 FFRs during low or high α power in either clear or noise condition were more predictive of behavioral

325 responses, we performed ML classification of percent correct (or perceptual performance level) into

326 poor (≤ 30th of overall percent correct) or good (≥ 70th of overall percent correct) using participants'

327 FFR frequency spectra as input for the SVM classifiers. In the clear condition (Fig. 3B), the classifier

328 was significantly better in decoding participants' perceptual performance level using low α FFRs

329 compared to high α FFRs. In contrast, in the noise condition (Fig. 3C), classification accuracies did not

330 differ between low or high α FFRs nor where they above the null distribution. These results provide

331 evidence that adding background noise disrupted the relationship of low α FFRs with behavioral

332 measures potentially as a result of compromising the SNR of the neural responses.

3.4 Comparison of brain-behavior relations in the NH and HL groups

The aforementioned analyses showed that during high arousal states (i.e., with low α power),

FFRs have a stronger relation with behavior compared to low arousal states, especially in the clear

condition, when pooling the NH and HL groups. Hence, we studied the changes in low α FFR

amplitudes with behavioral performance (QuickSiN, PTA, %-correct, and RTs) systematically by

fitting linear regression models [Eq. (2)]. We observed significant coefficients or slopes (β) when PTA

was used as X in Eq (2). For both the clear and noise-degraded speech, we found a significant negative

slope between low α F0 amplitudes and PTAs in the NH group ($\beta1 = -4.82 \times 10^{-9}$, $p = 0.01$, Fig. 4A) but

the slope of the HL group was not significantly different from the NH group. This indicates that even

with clinically "normal" hearing, participants with slightly poorer thresholds have smaller FFRs during

low α states. When separating the data by SNR, we observed similar trends of low α FFR amplitudes

decreased with increased PTA in the NH and HL groups for clear speech (Fig. 4B) though the slopes

were not significant. For noisy speech (Fig. 4C), we found that low α FFR amplitudes decreased

significantly with increased PTA in the NH group ($\beta1 = -6.53 \times 10^{-9}$, $p = 0.01$) and the slope of the HL

group was also significantly different ($\beta3 = 6.77 \times 10^{-9}$, $p = 0.02$) from the NH group. The fitted

regression lines in Fig. 4C showed that low α FFR amplitudes were generally diminished by noise in

the HL group.

4 Discussion

Previous neuroimaging work reveals weaker functional connectivity between brainstem and

cortex in older listeners with mild hearing loss than older adults with normal hearing for their age, and

this interplay robustly predicts their SiN perceptual performance (Bidelman et al., 2019). Adding to

these findings, we show the existence of active and dynamic modulation of brainstem speech

processing in NH older listeners, which was dependent on online changes in listeners' cortical state.

This active and dynamic cortical-brainstem modulation, however, is diminished when processing

speech in noise and in older adults with HL. Compared to NH listeners, HL listeners showed weaker

parieto-occipital α power but those with minimal hearing loss (i.e., smaller PTA) had unusually large

FFRs during low α states (gray dashed box in Fig. 4B). Although FFRs were smaller during low α

power, they were predictive of perceptual speech measures (Fig. 3A) when pooling NH and HL

participants and especially for clear speech (Fig. 3B). Collectively, our findings suggest that (i) FFRs

during low α power (i.e., high cortical arousal states) are more predictive of behavioral performance,

(ii) decreased low α FFRs with increased PTA in both NH and HL participants, and (iii) increased low

α FFRs in older adults with mild hearing loss for clear speech suggesting an increase in central gain.

4.1   Effects of age on cortical α power and cortical modulation of brainstem speech processing

Cortical α indexes states of wakefulness and arousal (Pfurtscheller et al., 1996; Aftanas et al., 2002;

Uusberg et al., 2013). Still, there is also evidence showing that α power may vary or index mind

wandering during cognitive tasks (Compton et al., 2019; Maillet et al., 2020). Calculating the span

length of low or high α trials for each listener showed averages of ~ 1.7 trials in both hearing groups,

equating to several hundred milliseconds during our task. The relative speed of these fluctuations

suggests that the α-modulations observed here are unlikely related to mind wandering *per se*, which

presumably develops over longer time courses [tens of seconds (Pelagatti et al., 2020)]. Instead, we

infer low α power tracks high arousal state while high α power reflects task focus but in a state of

wakeful relaxation. Induced α activity is crucial for SiN perception as it might suppresses irrelevant

information like noise to aid target speech processing (Strauß et al., 2014). In our previous study

conducted in younger listeners (18-35 years) using similar EEG tasks, we observed larger α power to

noise-degraded compared to clear speech during active engagement (see Fig. 2F in Lai et al., 2022).

However, here in both NH and HL older adults, we do not find this same noise-related α effect. In

general, α power was larger in NH than HL older listeners (Fig. 2C). We also observed that high α

RMS values in young NH listeners (Fig. 2F in Lai et al., 2022) had a larger range than NH older adults.

This observation advocates a reduction of α power with aging which has also been shown in other

studies (Babiloni et al., 2006; Purdon et al., 2015). Decreased α activity is also related to declines in

cognitive functions with increasing age (Klimesch, 1997, 1999).

384   More critically, we demonstrate the presence of dynamic and online modulation of brainstem

speech encoding by fluctuations in cortical α state in older NH adults that are fundamentally different

from those observed in younger, normal-hearing listeners (cf. Lai et al., 2022). In younger listeners,

lower cortical α states positively correlate with smaller FFRs during SiN perception (Lai et al., 2022).

Furthermore, low-α-indexed FFRs recorded in noisy backgrounds are predictive of behavioral RTs for

rapid speech detection and have higher accuracies in token classification (Lai et al., 2022). Here, unlike

younger listeners which require more difficult perceptual tasks (i.e., SiN perception) to tax the system

and reveal effects of cortical arousal state on brainstem FFRs, we observed cortical modulation of FFRs

in NH older adults during the perception of *clear* speech (Fig. 2D & E). Moreover, low-α-indexed

FFRs associated more with behavior [speech detection (Fig. 3) and PTAs (Figs 4)] than high-α-indexed

FFRs. Contrastively, in noise, low α and high α FFR amplitudes were not classifiable in terms of

perceptual performance level. Taken together, the pattern of cortical-brainstem interactions in speech

processing we found here in older NH listeners appears similar to what is found in younger listeners

under challenging listening environments (cf. Lai et al, 2022). This indicates that aging might alter

cross-talk between functional levels of the auditory system under challenging listening conditions as a

means of compensatory processing. Similar maladaptive plasticity has been previously observed at

higher cortical levels where frontal brain regions are more strongly engaged to aid auditory-sensory

coding in superior temporal gyrus (Price et al., 2019). This further suggests the presence of age-related

deficits in top-down modulation of brainstem speech processing by cortex and provides an explanation

to why older listeners find it more exhausting to participate in cocktail party-like listening situations

compared to younger listeners (Pichora-Fuller et al., 2017).

4.2   Effects of hearing loss on cortical α power and cortical modulation of brainstem speech

processing

Compared to the NH group, we observed decreases in parieto-occipital α power in the HL group

in both SNR conditions (Fig. 2C). Lower α power is reported in listeners with moderate hearing-loss

across the age spectrum (Petersen et al., 2015). Moreover, PTA correlates negatively with pre-stimulus

α power in older listeners (Alhanbali et al., 2021). These findings are partly concordant with our data

since we found lower α power (during stimuli) in older listeners with mild to moderate hearing loss.

Furthermore, in the HL group, we found no cortical-related enhancements of FFRs (i.e., F0 ratio ≈1)

for clear speech, and responses were not different from the NH group in the noise condition though F0

ratio was > 1 (Fig. 2D). The interaction effect of SNR x α power was also distinct in direction between

hearing groups. This finding implies that modulation of brainstem speech processing by cortical α state

is altered in older listeners with mild hearing loss for both clear and noise-degraded speech processing.

In addition to parieto-occipital α power, trends of low-α-indexed FFRs to clear speech

decreased with increased PTAs were observed in NH and HL listeners (Fig. 4B) and the slope of

decrease was significant for noise-degraded speech in NH listeners (Fig 4C). The reduction in low α

FFRs with poorer PTAs is probably related to the decrease in peripheral hearing ability. However,

when comparing across groups at comparable hearing loss (PTA = 15-22 dB HL), we found enhanced

FFR amplitudes in HL listeners (gray dashed box in Fig. 4B). Speculatively, this could indicate an

increase in central gain, probably related to high-frequency (> 4 kHz) hearing loss in the HL group

(our NH listeners had normal audiometric thresholds up to 4 kHz). Similar central gain compensation

secondary to peripheral hearing loss has been observed previously in both animal and human

neuroimaging studies (Bidelman et al., 2014; Chambers et al., 2016). These phenomena were

completely collapsed by noise in the HL group where low α FFRs were relatively smaller in most HL

listeners (Fig. 4C).

4.3   Association of brainstem speech processing during high arousal states with behaviors

In younger listeners and under noisy backgrounds, we previously showed that neural decoding

applied to low α FFRs offered higher accuracies in token classification as compared to high α FFRs

(see Fig. 5 in Lai et al., 2022). In this study, under no background noise, we observed frequency spectra

of low α FFR had better classification accuracies for perceptual performance level than high α FFR.

Thus, better speech token discrimination is consistently observed in FFRs during high arousal states in

both younger and older adults. Furthermore, low α FFRs were also observed to be decreased as PTA

increased, especially in the NH group for noise-degraded speech. These observations suggested that

brainstem FFRs during high arousal states have a strong association with behavior perception.

5   Conclusion

Collectively, our study reveals age-related hearing loss not only reduces cortical α power but

differentially alters its dynamic relationship with the subcortical speech processing especially in the

presence of noise. While brainstem speech processing is actively modulated by cortical arousal state

in normal-hearing older adults, this modulation is disrupted by signal degradations (i.e., noise) and

hearing loss. Speech-FFRs during low α states also offer a higher fidelity representation of the acoustic

speech signature and are more predictive of perceptual performance than FFRs yoked to states of high

cortical α. Enhanced FFRs in older adults with near-normal hearing (i.e., very mild hearing loss)

suggest the presence of increased central gain compensation for reduced auditory input (Bidelman et

al., 2014; Chambers et al., 2016).

6   Funding

This work was supported by the National Institutes of Health (NIH/NIDCD R01DC016267)

(G.M.B.), the Canadian Institutes of Health Research (MOP 106619) (C.A.), and the Natural

Sciences and Engineering Research Council of Canada (NSERC, 194536) (C.A).

7   Acknowledgments

We would like to thank Dawei Shen and Stephen R. Arnott for their assistance with data collection as
well as Yimei Li and Tushar Patni for providing consultation in statistical linear regression analysis.
455

8   Conflict of Interest

The authors declare no competing financial interests.

9   References

Aftanas LI, Varlamov AA, Pavlov S v., Makhnev VP, Reva N v. (2002) Time-dependent cortical
asymmetries induced by emotional arousal: EEG analysis of event-related synchronization and
desynchronization in individually defined frequency bands. Int J Psychophysiol 44:67–82.

Alain C, McDonald KL (2007) Age-Related Differences in Neuromagnetic Brain Activity
Underlying Concurrent Sound Perception. The Journal of Neuroscience 27:1308.

Alain C, Ogawa KH, Woods DL (1996) Aging and the Segregation of Auditory Stimulus Sequences.
Journal of Gerontology: PSYCHOLOGICAL SCIENCES 51:91–9.

Alhanbali S, Munro KJ, Dawes P, Perugia E, Millman RE (2021) Associations between pre-stimulus
alpha power, hearing level and performance in a digits-in-noise task. International journal of
audiology 61(3): 197–204.

Anderson S, Parbery-Clark A, Yi HG, Kraus N (2011) A neural basis of speech-in-noise perception
in older adults. Ear and Hearing 32:750–757.

Babiloni C, Binetti G, Cassarino A, Dal Forno G, del Percio C, Ferreri F, Ferri R, Frisoni G,
Galderisi S, Hirata K, Lanuzza B, Miniussi C, Mucci A, Nobili F, Rodriguez G, Romani GL, Rossini
PM (2006) Sources of cortical rhythms in adults during physiological aging: a multicentric EEG
study. Hum Brain Mapp 27:162–172.

Bidelman GM (2018a) Sonification of scalp-recorded frequency-following responses (FFRs) offers
improved response detection over conventional statistical metrics. Journal of neuroscience
methods 293: 59–66.

Bidelman GM (2018b) Subcortical sources dominate the neuroelectric auditory frequency-following
response to speech. Neuroimage 175:56–69.

Bidelman GM, Momtaz S (2021) Subcortical rather than cortical sources of the frequency-following
response (FFR) relate to speech-in-noise perception in normal-hearing listeners. Neuroscience
letters, 746, 135664.

Bidelman GM, Moreno S, Alain C (2013) Tracing the emergence of categorical speech perception in
the human auditory system. Neuroimage 79:201–212.

Bidelman GM, Price CN, Shen D, Arnott SR, Alain C (2019) Afferent-efferent connectivity between
auditory brainstem and cortex accounts for poorer speech-in-noise comprehension in older adults.
Hearing Research 382:107795.

Bidelman GM, Villafuerte JW, Moreno S, Alain C (2014) Age-related changes in the subcortical-
cortical encoding and categorical perception of speech. Neurobiol Aging 35:2526–2540.

Böttger D, Herrmann CS, von Cramon DY (2002) Amplitude differences of evoked alpha and
gamma oscillations in two different age groups. Int J Psychophysiol 45:245–251.

Brugge JF, Nourski K v., Oya H, Reale RA, Kawasaki H, Steinschneider M, Howard MA (2009)
Coding of Repetitive Transients by Auditory Cortex on Heschl's Gyrus. Journal of Neurophysiology
102:2358.

Chambers AR, Resnik J, Yuan Y, Whitton JP, Edge AS, Liberman MC, Polley DB (2016) Central
Gain Restores Auditory Processing following Near-Complete Cochlear Denervation. Neuron 89:867–
879.

Chiang AKI, Rennie CJ, Robinson PA, van Albada SJ, Kerr CC (2011) Age trends and sex
differences of alpha rhythms including split alpha peaks. Clin Neurophysiol 122:1505–1517.

Coffey EBJ, Herholz SC, Chepesiuk AMP, Baillet S, Zatorre RJ (2016) Cortical contributions to the
auditory frequency-following response revealed by MEG. Nature Communications 7:11070.

Compton RJ, Gearinger D, Wild H (2019) The wandering mind oscillates: EEG alpha power is
enhanced during moments of mind-wandering. Cognitive, Affective and Behavioral Neuroscience
19:1184–1191.

Dimitrijevic A, Smith ML, Kadis DS, Moore DR (2017) Cortical alpha oscillations predict speech
intelligibility. Frontiers in Human Neuroscience 11:88.

Dubno JR (1984) Effects of age and mild hearing loss on speech recognition in noise. J Acoust Soc
Am 76:87–96.

Dubno JR, Schaefer AB (1992) Comparison of frequency selectivity and consonant recognition
among hearing-impaired and masked normal-hearing listeners. J Acoust Soc Am 91:2110–2121.

Fostick L, Ben-Artzi E, Babkoff H (2013) Aging and speech perception: beyond hearing threshold
and cognitive ability. Journal of Basic and Clinical Physiology and Pharmacology 24:175–183.

Gaál ZA, Boha R, Stam CJ, Molnár M (2010) Age-dependent features of EEG-reactivity--spectral,
complexity, and network characteristics. Neurosci Lett 479:79–84.

Galbraith GC, Brown WS (1990) Cross-correlation and latency compensation analysis of click-
evoked and frequency-following brain-stem responses in man. Electroencephalogr Clin Neurophysiol
77:295–308.

Gallun FJ, Best V (2020) Age-Related Changes in Segregation of Sound Sources. Auditory Research
72:143–171.

Haegens S, Nácher V, Luna R, Romo R, Jensen O (2011) α-Oscillations in the monkey sensorimotor
network influence discrimination performance by rhythmical inhibition of neuronal spiking. Proc
Natl Acad Sci U S A 108:19377–19382.

Händel BF, Haarmeier T, Jensen O (2011) Alpha oscillations correlate with the successful inhibition
of unattended stimuli. J Cogn Neurosci 23:2494–2502.

Henry MJ, Herrmann B, Kunke D, Obleser J (2017) Aging affects the balance of neural entrainment
and top-down neural modulation in the listening brain. Nature Communications 2017 8:1 8:1–11.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: Gating by
inhibition. Frontiers in Human Neuroscience 4:186.

Jin S-HH, Liu C, Sladen DP (2014) The effects of aging on speech perception in noise: comparison
between normal-hearing and cochlear-implant listeners. J Am Acad Audiol 25:656–665.

Killion MC, Niquette PA, Gudmundsen GI, Revit LJ, Banerjee S (2004) Development of a quick
speech-in-noise test for measuring signal-to-noise ratio loss in normal-hearing and hearing-impaired
listeners. J Acoust Soc Am 116:2395.

Klatt LI, Getzmann S, Begau A, Schneider D (2020) A dual mechanism underlying retroactive shifts
of auditory spatial attention: dissociating target- and distractor-related modulations of alpha
lateralization. Scientific Reports 2020 10:1 10:1–13.

Klimesch W (1997) EEG-alpha rhythms and memory processes. Int J Psychophysiol 26:319–340.

Klimesch W (1999) EEG alpha and theta oscillations reflect cognitive and memory performance: a
review and analysis. Brain Res Brain Res Rev 29:169–195.

Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: the inhibition-timing
hypothesis. Brain Res Rev 53:63–88.

Kortlang S, Mauermann M, Ewert SD (2016) Suprathreshold auditory processing deficits in noise:
Effects of hearing loss and age. Hearing Research 331:27–40.

Krishnan A, Xu Y, Gandour J, Cariani P (2005) Encoding of pitch in the human brainstem is
sensitive to language experience. Brain Res Cogn Brain Res 25:161–168.

Lai J, Bartlett EL (2015) Age-related shifts in distortion product otoacoustic emissions peak-ratios
and amplitude modulation spectra. Hearing Research 327:186–198.

Lai J, Bartlett EL (2018) Masking Differentially Affects Envelope-following Responses in Young
and Aged Animals. Neuroscience 386:150–165.

Lai, J., Price, C. N., & Bidelman, G. M. (2022). Brainstem speech encoding is dynamically shaped
online by fluctuations in cortical α state. Neuroimage 263: 119627.

Lai J, Sommer AL, Bartlett EL (2017) Age-related changes in envelope-following responses at
equalized peripheral or central activation. Neurobiol Aging 58:191.

Mai G, Schoof T, Howell P (2019) Modulation of phase-locked neural responses to speech during
different arousal states is age-dependent. Neuroimage 189:734–744.

Maillet D, Yu L, Lau B, Chow R, Alain C, Grady CL (2020) Differential effects of mind-wandering
and visual distraction on age-related changes in neuro-electric brain activity and variability.
Neuropsychologia 146:107565.

McMahon CM, Boisvert I, de Lissa P, Granger L, Ibrahim R, Lo CY, Miles K, Graham PL (2016)
Monitoring Alpha Oscillations and Pupil Dilation across a Performance-Intensity Function. Frontiers
in Psychology 7: 745.

Musacchia G, Strait D, Kraus N (2008) Relationships between behavior, brainstem and cortical
encoding of seen and heard speech in musicians and non-musicians. Hear Res 241:34–42.

Oostenveld R, Praamstra P (2001) The five percent electrode system for high-resolution EEG and
ERP measurements. Clinical Neurophysiology 112:713–719.

Park HL, O'Connell JE, Thomson RG (2003) A systematic review of cognitive decline in the general
elderly population. Int J Geriatr Psychiatry 18:1121–1134.

Parthasarathy A, Bartlett E (2012) Two-channel recording of auditory-evoked potentials to detect
age-related deficits in temporal processing. Hearing Research 289:52–62.

Parthasarathy A, Bartlett EL (2011) Age-related auditory deficits in temporal processing in F-344
rats. Neuroscience 192:619–630.

Parthasarathy A, Cunningham PA, Bartlett EL (2010) Age-related differences in auditory processing
as assessed by amplitude-modulation following responses in quiet and in noise. Frontiers in Aging
Neuroscience 2:1–10.

Parthasarathy A, Datta J, Torres JA, Hopkins C, Bartlett EL (2014) Age-related changes in the
relationship between auditory brainstem responses and envelope-following responses. Journal of the
Association for Research in Otolaryngology 15:649–66.

Parthasarathy A, Lai J, Bartlett EL (2016) Age-Related Changes in Processing Simultaneous
Amplitude Modulated Sounds Assessed Using Envelope Following Responses. Journal of the
Association for Research in Otolaryngology 17:119–132.

Peelle JE, Troiani V, Wingfield A, Grossman M (2010) Neural Processing during Older Adults'
Comprehension of Spoken Sentences: Age Differences in Resource Allocation and Connectivity.
Cerebral Cortex (New York, NY) 20:773.

Pelagatti C, Binda P, Vannucci M (2020) A closer look at the timecourse of mind wandering:
Pupillary responses and behaviour. PLOS ONE 15:e0226792.

Petersen EB, Wöstmann M, Obleser J, Stenfelt S, Lunner T (2015) Hearing loss impacts neural alpha
oscillations under adverse listening conditions. Frontiers in psychology, 6, 177.

Pfurtscheller G, Stancák A, Neuper C (1996) Event-related synchronization (ERS) in the alpha band
— an electrophysiological correlate of cortical idling: A review. International Journal of
Psychophysiology 24:39–46.

Pichora-Fuller MK, Alain C, Schneider BA (2017) Older Adults at the Cocktail Party. :227–259.

Poth EA, Boettcher FA, Mills JH, Dubno JR (2001) Auditory brainstem responses in younger and
older adults for broadband noises separated by a silent gap. Hearing Research 161:81–86.

Presacco A, Simon JZ, Anderson S (2016) Evidence of degraded representation of speech in noise, in
the aging midbrain and cortex. Journal of Neurophysiology 116.

Price CN, Alain C, Bidelman GM (2019) Auditory-frontal Channeling in α and β Bands is Altered by
Age-related Hearing Loss and Relates to Speech Perception in Noise. Neuroscience 423:18–28.

Price CN, Bidelman GM (2021) Attention reinforces human corticofugal system to aid speech
perception in noise. Neuroimage 235:118014.

Purdon PL, Pavone KJ, Akeju O, Smith AC, Sampson AL, Lee J, Zhou DW, Solt K, Brown EN
(2015) The Ageing Brain: Age-dependent changes in the electroencephalogram during propofol and
sevoflurane general anaesthesia. British Journal of Anaesthesia 115:i46–i57.

Rihs TA, Michel CM, Thut G (2007) Mechanisms of selective inhibition in visual spatial attention
are indexed by alpha-band EEG synchronization. Eur J Neurosci 25:603–610.

Saderi D, Schwartz ZP, Heller CR, Pennington JR, David S v. (2021) Dissociation of task
engagement and arousal effects in auditory cortex and midbrain. Elife 10:1–25.

Saiz-Alía M, Forte AE, Reichenbach T (2019) Individual differences in the attentional modulation of
the human auditory brainstem response to speech inform on speech-in-noise deficits. Scientific
Reports 2019 9:1 9:1–10.

Scherg M, Ebersole JS (1994) Brain source imaging of focal and multifocal epileptiform EEG
activity. Neurophysiol Clin 24:51–60.

Scherg M, Ille N, Bornfleth H, Berg P (2002) Advanced tools for digital EEG review: virtual source
montages, whole-head mapping, correlation, and phase analysis. J Clin Neurophysiol 19:91–112.

Schneider BA, Daneman M, Murphy DR (2005) Speech comprehension difficulties in older adults:
cognitive slowing or age-related changes in hearing? Psychology and Aging 20:261–271.

Skoe E, Kraus N (2010) Hearing it again and again: on-line subcortical plasticity in humans. PloS
one, 5(10), e13645.

Song JH, Skoe E, Banai K, Kraus N (2011) Perception of Speech in Noise: Neural Correlates.
Journal of Cognitive Neuroscience 23:2268.

Souza PE, Boike KT, Witherell K, Tremblay K (2007) Prediction of speech recognition from
audibility in older listeners with hearing loss: effects of age, amplification, and background noise. J
Am Acad Audiol 18:54–65.

Strauß A, Wöstmann M, Obleser J (2014) Cortical alpha oscillations as a tool for auditory selective
inhibition. Front Hum Neurosci 8: 350.

Takahashi GA, Bacon SP (1992) Modulation detection, modulation masking, and speech
understanding in noise in the elderly. Journal of Speech and Hearing Research 35:1410–1421.

Tichko P, Skoe E (2017) Frequency-dependent fine structure in the frequency-following response:
The byproduct of multiple generators. Hear Res 348:1–15.

Uusberg A, Uibo H, Kreegipuu K, Allik J (2013) EEG alpha and cortical inhibition in affective
attention. International Journal of Psychophysiology 89:26–36.

Vermeire K, Knoop A, Boel C, Auwers S, Schenus L, Talaveron-Rodriguez M, de Boom C, de
Sloovere M (2016) Speech Recognition in Noise by Younger and Older Adults: Effects of Age,
Hearing Loss, and Temporal Resolution. Annals of Otology, Rhinology & Laryngology 125:297–
302.

Weisz N, Dohrmann K, Elbert T (2007) The relevance of spontaneous activity for the coding of the
tinnitus sensation. Prog Brain Res 166:61–70.

Weisz N, Hartmann T, Müller N, Lorenz I, Obleser J (2011) Alpha Rhythms in Audition: Cognitive
and Clinical Perspectives. Frontiers in Psychology 2: 73.

Wöstmann M, Herrmann B, Wilsch A, Obleser J (2015) Neural Alpha Dynamics in Younger and
Older Listeners Reflect Acoustic Challenges and Predictive Benefits. Journal of Neuroscience
35:1458–1467.

Yordanova JY, Kolev VN, Başar E (1998) EEG theta and frontal alpha oscillations during auditory
processing change with aging. Electroencephalogr Clin Neurophysiol 108:497–505.

10   Figures

Figure 1. Target speech detection performance correlates with average response time collected
during an active EEG task. (A) Prior to EEG recordings, all participants' pure-tone audiometry
tested at 250-8000 Hz were obtained and speech-in-noise perception was assessed with QuickSiN.
Subsequently, speech-EEGs to consonant-vowel phonemes (/ba/, /pa/, and /ta/) were recorded under
clear or noisy (+10 dB SNR) backgrounds while participants actively engaged in the target speech
detection task. (B) Correct responses to target speech (i.e., /ta/) are predictive of reaction times in
quite (clear) but not noisy backgrounds. (C) When participants were divided into normal (NH) and
hearing loss (HL) groups, the correlation of speech detection accuracy and reaction time obtained in
quiet backgrounds is significant only in the HL group. r = Spearman's correlation; shaded area = 95
% CI of the regression line.

Figure 2. Normal hearing (NH) participants have a higher parieto-occipital α power and their
brainstem speech processing is more strongly modulated by α band in clear backgrounds. (A)
Average α waveform of low- and high-power trials plotted from a representative NH participant. (B)
Frequency spectra of the steady state (10-100 ms) portion of low and high α brainstem FFRs from the
same participant in (A). Note the distinct response at ~150 Hz, corresponding to the voice pitch. (C)
Root-mean-square (RMS) values of both low and high α of the NH group were significantly higher
than the HL group. # p< 0.01 (Conover's test, non-parametric pairwise test, with Bonferroni
adjustment) (D) FFR F0 ratios during low and high α trials. FFR F0 ratios were higher in the NH vs.
HL group (Mann-Whitney U test) in the clear condition. Bars marked (<0.05) are significantly larger
than 1 (1-sample t-test) indicating enhancement of the FFR with changes in cortical α. (E) Grand
average FFR F0 amplitudes as a function of SNR (clear vs. noise), α power (low vs. high), and group
(NH vs. HL). Error bars= ± s.e.m., *p<0.05 (Wilcoxon signed-rank test).

Figure 3. Low α FFRs better predict behavioral performance than high α FFRs during clear speech perception when pooling across background conditions. (A) Low α FFRs correlated positively with percent correct of speech target detection. r = Spearman's correlation. Shaded area=95 % CI of the regression line. (B) For clear speech, SVM classifier accuracy was significantly better using low α (but not high α) FFR frequency spectra to classify subjects' perceptual performance level (i.e., poor vs. good) compared to the null classification accuracies. (C) For noise-degraded speech, classification accuracies were similar when using low or high α FFR spectra and did not differ from the null classification accuracies. Upper/lower ticks=max/min; center tick=medians. * p< 0.01

Figure 4. Hearing loss is associated with smaller speech-FFR amplitude under low cortical α (i.e.,
high arousal) states. (A) When pooling SNRs, the decrease (slope) of low α FFR F0 amplitudes as a
function of PTA in the NH group was significant ($\beta1 = -4.82 \times 10^{-9}$, p = 0.01) and the slope of the HL
group was not significantly different from the NH group. (B) For clear speech, similar trends of low α
FFR amplitudes decreased with increased PTA were observed in the NH and HL groups though the
slopes were not significant. The gray dashed box marks the overlapping PTA region for both groups.
(C) For noise-degraded speech, the slope of low α FFR amplitudes decreased with increased PTA was
significant ($\beta1 = -6.53 \times 10^{-9}$, p = 0.01) in the NH group and the slope of the HL group was significantly
different ($\beta3 = 6.77 \times 10^{-9}$, p = 0.02) from the NH group. The fitted regression lines showed that low α
FFR amplitudes were generally diminished by noise in the HL group. Shaded area=95 % CI of the
regression line.