Male sex hormone and reduced plakoglobin jointly impair atrial conduction and cardiac sodium currents

Laura C. Sommerfeld[1,2,3], Andrew P. Holmes[1,4], Ting Y. Yu[1,5], Christopher O'Shea[1,5],

Deirdre M. Kavanagh[1,6], Jeremy M. Pike[1,6], Thomas Wright[1], Fahima Syeda[1], Areej Aljehani[1],

Tania Kew[1], Victor R. Cardoso[1,7], S. Nashitha Kabir[1], Claire Hepburn[1], Priyanka M. Menon[1],

Sophie Broadway-Stringer[1], Molly O'Reilly[1], Anika Witten[8,9], Lisa Fortmueller[2,3],

Susanne Lutz[11], Alexandra Kulle[12], Georgios V. Gkoutos[7,13,14], Davor Pavlovic[1], Wiebke Arlt[15],

Gareth G. Lavery[15], Richard Steeds[1,16], Katja Gehmlich[1], Monika Stoll[8,9,10], Paulus

Kirchhof[1,3,17], & Larissa Fabritz[1,2,3]

[1]Institute of Cardiovascular Sciences, University of Birmingham, UK

[2]University Center of Cardiovascular Science, University Heart and Vascular Center, UKE Hamburg, Germany

[3]German Center for Cardiovascular Research (DZHK), Standort Hamburg/Kiel/Lübeck, Germany

[4]School of Biomedical Sciences, Institute of Clinical Sciences, University of Birmingham, UK

[5]Research and Training Centre in Physical Sciences for Health, Birmingham, UK

[6]Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham, UK

[7]Institute of Cancer and Genomic Sciences, University of Birmingham, UK

[8]Genetic Epidemiology, Institute for Human Genetics, University of Münster, Germany

[9]Core Facility Genomics of the Medical Faculty, University of Münster, Germany

[10]Cardiovascular Research Institute Maastricht, Department of Biochemistry, Maastricht University, The Netherlands

[11]Institute of Pharmacology and Toxicology, University Medical Center Göttingen, Germany

[12]Division of Paediatric Endocrinology and Diabetes, University Hospital Schleswig-Holstein, Campus Kiel, Germany

[13]Institute of Translational Medicine, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

[14]MRC Health Data Research UK (HDR), Midlands Site, UK

[15]Institute of Metabolism and Systems Research (IMSR) University of Birmingham, UK & Centre for Endocrinology, Diabetes and Metabolism (CEDAM), Birmingham Health Partners

[16]Department of Cardiology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

[17]Department of Cardiology, University Heart & Vascular Center Hamburg, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

Corresponding author:
Professor Larissa Fabritz
University Center of Cardiovascular Science
University Heart and Vascular Center, UKE Hamburg

Hamburg, Germany
+49 40 7410 57980
L.Fabritz@uke.de

Conflicts of interest

The authors have declared no direct conflict of interest in regards to the manuscript.

L.F. has received institutional research grants from governmental and charity funding agencies and several biomedical companies.

P.K. has received research support from several drug and device companies active in atrial fibrillation and has received honoraria from several such companies in the past.

L.F. and P.K. are listed as inventors on two patents held by University of Birmingham (Atrial Fibrillation Therapy WO 015140571, Markers for Atrial Fibrillation WO 2016012783).

Word count: 11408 (all text inclusive of title page, full text, references, figure legends and tables)

Abstract

Androgenic anabolic steroids (AAS) are commonly abused by young men. Male sex associates with earlier manifestation of common and rare cardiac conditions including atrial fibrillation and arrhythmogenic right ventricular cardiomyopathy (ARVC). Clinical data suggest an atrial involvement in ARVC. The disease is caused by desmosomal gene defects such as reduced plakoglobin expression. Analysis of clinical records from 146 ARVC patients identified male preponderance and increased prevalence of atrial arrhythmias in patients with definite ARVC. Definite patients displayed ECG changes suggesting atrial remodelling. To study mechanisms of atrial remodelling due to desmosomal vulnerability and AAS, young adult male mice, heterozygously deficient for plakoglobin (Plako$^{+/-}$) and wildtype (WT) littermates, were chronically exposed to 5α-dihydrotestosterone (DHT) or placebo. DHT increased atrial expression of pro-hypertrophic, fibrotic and inflammatory transcripts. DHT caused atrial conduction slowing, decreased peak sodium current density, reduced action potential amplitude and lowered the peak depolarisation rate in Plako$^{+/-}$ but not WT atria. Super-resolution microscopy revealed a reduction in Na$_v$1.5 clustering in Plako$^{+/-}$ atrial cardiomyocytes following DHT exposure. These data reveal that AAS combined with plakoglobin deficiency cause pathological atrial electrical remodelling in young male hearts. AAS abuse may increase the risk of atrial myopathy in males with desmosomal gene variants.

Key words: arrhythmias, ion channels, sex hormones

Introduction

Several inherited arrhythmia syndromes develop more severe phenotypes in men carrying

pathogenic variants, e.g. Brugada syndrome or Arrhythmogenic Right Ventricular

Cardiomyopathy (ARVC) (1). Male sex also associates with greater incidence of atrial

arrhythmias, both in the general population and in rare conditions (2-7). Effects mediated by

anabolic androgenic steroids (AAS) such as testosterone and the most potent androgen,

5α-dihydrotestosterone (DHT), could contribute. Abuse of AAS is an emerging global health

concern, not restricted to elite athletes but common in the general population, with reports

indicating a 3.3% lifetime prevalence worldwide (8). AAS are abused predominately by men

to increase muscle mass, improve athletic performance and alter appearance (8). However,

AAS can cause cardiac pathology, including hypertrophy and electrophysiological changes

(9-14). Atrial arrhythmias have recently been associated with elevated total plasma

testosterone levels in men (15) and observed in patients known to take AAS devoid of a

clinical indication (16-18). Despite these observations, the mechanisms underpinning cardiac

electrical remodelling in response to higher levels of AAS are largely unknown.

ARVC has recently been reported to show more adverse outcomes in men (19) related to sex

hormone levels (20). Emerging evidence suggests increased incidence of atrial arrhythmias

in cardiomyopathies (21). ARVC is often caused by variants in desmosomal genes including

plakoglobin (22-25). Plakoglobin is located in desmosomal junctional complexes where it

stabilizes cell-cell contacts (26), thereby maintaining mechanical and electrical integrity of the

myocardium.

We hypothesized that a substantial proportion of ARVC patients is suffering from clinically

relevant atrial arrhythmias and that vulnerability of the desmosome, caused by e.g.

plakoglobin reduction, may increase the risk of male sex hormone-induced atrial electrical

remodelling.

To test this hypothesis, we screened patient records from definite and non-definite ARVC

patients seen at a tertiary center inherited cardiac conditions clinic for atrial arrhythmias. We

furthermore employed a murine ARVC model to study the effects of chronically elevated AAS

levels in male mice with heterozygous plakoglobin (gamma-catenin) deficiency (Plako$^{+/-}$) (27)

and their wildtype (WT) littermates.

Results

Atrial arrhythmias and ECG changes in definite ARVC patients

The clinical cohort studied comprised of 146 patients with suspected cardiomyopathy; 97 were identified as "Non-definite" (possible) ARVC cases and 49 as "Definite" ARVC, i.e. presenting with a complete phenotype according to 2010 Task Force Criteria (1). Mean age of the patients at time of ECG analyses was not different between the groups (42 ± 18 years for non-definite vs 43 ± 18 years for definite). 24% of definite ARVC patients experienced atrial fibrillation and/or flutter compared to 3% of the non-definite ARVC patients (Table 1). There was a significant association between sex and ARVC diagnosis type (Table 1, 43% male amongst non-definite vs. 73% male amongst definite patients). Semi-automated analysis of digital ECG lead II recordings focussing on atrial parameters (Figure 1A) showed PR interval prolongation in advanced, definite disease stage (Figure 1B). P wave duration and P wave area were significantly increased in definite compared to non-definite patients. Non-definite ARVC patients exhibited no difference in the analyzed P wave parameters compared to unaffected control subjects. Results of a meta-analysis of clinical observational studies are presented in the discussion.

5α-dihydrotestosterone (DHT) causes general and cardiac growth response in mice

To study increased androgen exposure and desmosomal instability jointly in atria, Plako$^{+/-}$ mice, an established animal model of ARVC (28) and wildtype (WT) littermates were subjected to chronic DHT treatment over 6 weeks (please refer to Figure 2A for an overview scheme). Treatment led to a 3-4-fold increase in serum DHT concentration in both genotypes compared to controls (Ctr) (Figure 2B). Moreover, it increased body weight, seminal vesicle mass/tibia length ratio and atrial weight/tibia length ratio as well as causing left ventricular hypertrophy in Plako$^{+/-}$ animals (Figure 2C, Supplementary Figure 1).

DHT induces ARVC-like atrial ECG changes in plakoglobin-deficient mice

To check whether atrial ECG changes observed in definite ARVC patients are similarly

present in the murine model, ECGs were recorded from mice following DHT treatment. Both

PR interval as well as P wave duration were prolonged in Plako$^{+/-}$ animals exposed to DHT

compared to WT littermates exposed to DHT (Figure 3).

DHT causes atrial unfolded area dilation and atrial conduction slowing in heterozygous

plakoglobin-deficient hearts

Left atrial unfolded area to tibia length ratio was increased in Plako$^{+/-}$ DHT compared to

Plako$^{+/-}$ Ctr (Figure 4A&B), but not in WT left atria after DHT exposure. Chronic DHT

exposure slowed conduction in Plako$^{+/-}$ but not WT atria (Figure 4C&D). The reduction in

conduction velocity in Plako$^{+/-}$ DHT left atria was more pronounced at higher pacing

frequencies. The Plako$^{+/-}$ left atria exposed to DHT exhibited an overall prolongation of 95%

left atrial activation times and increased beat-to-beat activation variability compared to WT Ctr

and Plako$^{+/-}$ Ctr (Figure 5). Connexin expression was not impaired (Supplementary Figure 2).

DHT induces atrial expression of gene profiles implicated in ARVC

Exploratory RNA sequencing analysis confirmed approximately 50% reduction in atrial

plakoglobin (*Jup*) expression in Plako$^{+/-}$ compared to WT animals (Figure 6A). In control

hearts, gene expression patterns did not markedly differ between genotypes (data not

shown). Chronic DHT exposure resulted in significant transcriptional changes in atria of both,

WT and Plako$^{+/-}$ mice (Figure 6B&C). DHT activated expression of genes associated with

muscle growth (e.g. *Igf1*, *Mtpn, Myocd*), but additionally also immune (e.g. *C7*, *Tlr3, Tlr4*) and

pro-fibrotic response genes (e.g. *Col1a1*, *Col3a1, Srf, Lox)* (Figure 6C). Before-mentioned

transcriptional changes were similarly induced in left and right atria (Supplementary Figure 3).

While atrial cardiac myocyte diameter and endomysial collagen deposition was not

significantly different between genotypes after DHT treatment, as quantified in semi-

100 automated histology analysis (Supplementary Figure 5), cell capacitance was increased by

101 DHT treatment in Plako[+/-] DHT (Figure 8C).

DHT reduces action potential amplitude and rate of depolarisation in heterozygous

plakoglobin-deficient left atria

The intracellular microelectrode technique was used to record transmembrane action

potentials (TAPs) from paced, superfused left atria (Figure 7). Paced left atria isolated from

Plako[+/-] DHT mice showed longer activation times compared to all other groups (Figure 7C,

Supplementary Table 1). Action potential amplitude (APA) was reduced in Plako[+/-] DHT left

atria, as well as the peak rate of depolarisation (dV dt^{-1} max) (e.g. 120 ms pacing cycle length

WT Ctr: 118±5 V/s; WT DHT: 120±7 V/s; Plako[+/-] Ctr: 116±4 V/s; Plako[+/-] DHT: 89±5 V/s)

(Figure 7D&E, Supplementary Table 1), both indicative of sodium current impairments. The

Plako[+/-] DHT left atrial cells had a more positive resting membrane potential than WT Ctr

(Supplementary Table 1). Chronic DHT exposure did not modify action potential duration

(APD) in either genotype (Supplementary Table 1). Beat-averaged left atrial optical APD90,

as well as beat-to-beat APD90 variability were similar between all four groups

(Supplementary Table 2).

DHT decreases peak sodium current density in heterozygous plakoglobin-deficient left

atrial cardiac myocytes

To examine mechanisms underlying alterations in action potential morphology and

conduction in the Plako[+/-] DHT left atria, whole-cell patch clamp experiments were performed

monitoring peak sodium current (I_{Na}) amplitude and kinetics. Peak whole-cell I_{Na} density was

decreased by approximately 20% in Plako[+/-] left atrial cells following DHT exposure (Figure

8A&B). Activation kinetics were consistent between all groups (V_{50} activation, WT Ctr: -43±1

mV; WT DHT: -46±1 mV; Plako[+/-] Ctr: -46±1 mV; Plako[+/-] DHT: -46±2 mV). DHT exposure

tended to cause a left shift in steady-state inactivation kinetics in both WT and Plako[+/-], being

statistically significant in Plako[+/-] (Supplementary Figure 5A). The 50% recovery time (P_{50})

from inactivation was significantly longer in the Plako$^{+/-}$ DHT, suggestive of a delayed rate of

sodium channel recovery (Supplementary Figure 5B&C).

Left atrial cell capacitance was elevated by DHT exposure in Plako$^{+/-}$ (Figure 8C). Of note,

there was a considerable spread of individual cell capacitance in DHT-exposed groups,

indicative of variable degrees of hypertrophy. I_{Na} density was negatively correlated against

cell capacitance in Plako$^{+/-}$ left atria cardiac myocytes following DHT treatment (Figure 8D).

This negative correlation was not detected in the WT DHT group. These data suggest that a

hypertrophic response in the Plako$^{+/-}$ DHT left atrial cardiac myocytes was not matched by a

rise in I_{Na}, leading to an overall depletion of I_{Na} density. In contrast, hypertrophy in the WT

DHT left atria was matched by an increase in I_{Na}, so that I_{Na} density was preserved.

Whole left atrial tissue RNA expression of sodium voltage gated channel 5 (*Scn5a*) was

neither significantly affected by genotype, or DHT exposure (Figure 8E).

Na$_v$1.5 cluster depletion in Plako$^{+/-}$ left atrial cardiac myocytes following DHT exposure

To define a molecular cause of I_{Na} density depletion in Plako$^{+/-}$ DHT left atrial cardiac

myocytes, we examined Na$_v$1.5 organisation at super-resolution level using direct Stochastic

Optical Reconstruction Microscopy (dSTORM, workflow see Supplementary Figure 6). Due to

the high variability in T-tubule density between different atrial cells, we focused on Na$_v$1.5

channels located within ca. 200 nm of the contact cell surface membrane by using

TIRF/HILO. Exemplary super-resolution images of Na$_v$1.5 detections, as well as characteristic

cluster maps are shown in Figure 9A.

Characteristics of identified Na$_v$1.5 clusters, i.e. cluster density and cluster area, were similar

in all 4 groups (Supplementary Figure 7A&B). The 95% confidence intervals generated from

our murine WT Ctr left atria experiments suggest a range of $3\text{-}4\times10^{-3}$ Na$_v$1.5 channels per

nm^2 cluster (Supplementary Figure 7C, both mean and median at 3×10^{-3} channels per nm^2

cluster), resulting in an estimated nearest-neighbour distance of 20 nm.

At the Plako$^{+/-}$ DHT left atrial cardiac myocyte membranes, fewer Na$_v$1.5 detections were

observed compared to WT DHT (Figure 9A&B). Of the total detections, the proportion present

in distinct $Na_v1.5$ clusters was significantly lower in the Plako$^{+/-}$ DHT compared to WT DHT

cardiac myocytes (Figure 9C).

Discussion

Main findings

- Atrial arrhythmias and P wave changes are common in patients with ARVC.

- Exposure to the potent androgen 5α-dihydrotestosterone leads to pro-hypertrophic, pro-fibrotic and inflammatory transcriptional signatures in murine atria without overt phenotypic changes.

- Combining chronic 5α-dihydrotestosterone exposure with heterozygous plakoglobin deficiency leads to a profound atrial cardiomyopathy replicating ECG changes in patients with ARVC.

- Mechanistically, increased 5α-dihydrotestosterone concentrations interact with plakoglobin to decrease the number of membrane-localized $Na_v1.5$ clusters, reducing atrial sodium current density and causing atrial conduction slowing.

We report that definite ARVC patients exhibit increased P wave area and prolonged PR interval as well as P wave duration on ECG. A phenocopy of these ECG changes is observed in plakoglobin-deficient mice exposed to supraphysiological 5α-dihydrotestosterone (DHT) concentrations. Heterozygous plakoglobin deficiency predisposes cardiac atrial tissue to hypertrophy and a reduction in sodium current density after six weeks of exposure to elevated concentrations of DHT, resulting in a decrease in the peak upstroke velocity of the atrial action potential ($dV\ dt^{-1}$ max), left atrial conduction slowing and increased electrical beat-to-beat variability. Super-resolution microscopy, dSTORM, identified sarcolemmal depletion of $Na_v1.5$ channels and clusters at left atrial cardiac myocytes isolated from $Plako^{+/-}$ DHT-treated mice. This was accompanied by the functional electrophysiological modifications of reduced atrial $Na_v1.5$ current density and impaired atrial conduction.

The results underpin a role of plakoglobin in regulating $Na_v1.5$ channel cellular localization in

the left atrium and in preserving left atrial electrical integrity in response to stressors such as

pro-hypertrophic DHT exposure.

Patients in the definite ARVC disease stage display a high prevalence of atrial

arrhythmias, P wave changes on ECG and preponderance of male sex

A quarter (24%) of the patients with definite ARVC had atrial arrhythmias in our cohort. Meta-

analysis of published studies (between 1991 and 2021) revealed a weighted mean atrial

arrhythmia prevalence of 15% amongst a total of 1915 ARVC patients (Table 3). Over 99% of

the reported patients were diagnosed with definite ARVC according to accepted criteria. Male

preponderance of atrial phenotypes was reported in other cohorts as well (Table 3).

P wave duration and P wave area showed pathological changes in patients with definite

ARVC, suggesting pathophysiological remodelling of the atria. P wave prolongation and

increased P wave area have been associated with atrial fibrillation (AF) or increased AF

recurrence (29-32). Furthermore, definite ARVC patients displayed longer PR intervals

compared to non-definite patients and controls, suggesting attenuated conduction of the

atrioventricular node. Conduction slowing is a common feature of heart rhythm disorders

including AF, and acts by permitting the development and maintenance of both micro and

macro re-entry (33). In line with our results, Baturova and colleagues recently found evidence

for ARVC disease progression to be paralleled by changes in P wave area (34). Our data

substantiate clinical evidence for progressive atrial conduction disturbances with progression

of ARVC. Our murine data demonstrate that these changes can arise from an interaction

between genetic desmosomal defects and AAS.

DHT exposure interacts with plakoglobin deficiency leading to atrial electrical

dysfunction

Although supraphysiological AAS intake is commonly used by competitive athletes to

enhance performance (doping), reports are unsystematic due to underreporting of AAS

intake. Murine models represent a unique tool to observe early atrial changes arising in

interaction between genetic defects in the desmosome and AAS exposure. We show that

chronic DHT exposure combined with heterozygous deletion of plakoglobin prolongs PR

interval and P wave duration and slows atrial conduction. Affected atria did not show fibrotic

or fatty changes, and connexin expression was not altered. A third important determinant of

conduction velocity is the magnitude of depolarising current, primarily carried by Na^+ through

$Na_v1.5$ channels (35). Our electrophysiological measurements found reduced atrial action

potential amplitude and upstroke velocity (APA and $dV\ dt^{-1}$ max), demonstrating a decreased

sodium current that is sufficient to lower both the magnitude and rate of depolarisation, key

contributors to cardiac conduction velocity. A delay of sodium current recovery times, as

observed in $Plako^{+/-}$ DHT-treated cardiomyocytes, was enhanced at higher pacing

frequencies. Super-resolution microscopy identified reduced availability and altered clustering

of sodium channels as a likely mechanism for these functional defects. The reduced

availability of $Na_v1.5$ channels in cells already operating without a sufficient conduction

reserve (36, 37), is likely to cause more pronounced conduction slowing and beat-to-beat

variability as observed at the higher pacing frequencies in our experiments.

Supraphysiological DHT concentrations increased atrial expression of genes related to

immune response and fibrotic remodelling. Both processes contribute to the pathophysiology

of ARVC as well as AF (38-46) and are likely intertwined. Lysyl oxidase (*Lox*), which we here

report to be upregulated in atria in response to DHT, mediates cross-linking of collagen I and

collagen III fibrils. Both collagen isoforms, amongst others, were also upregulated in DHT-

exposed atria in our study, indicating remodelling of the extracellular matrix (ECM). Lox

overexpression in mice was also found to accelerate the inflammatory response during

angiotensin II (AngII)-induced cardiac hypertrophy, including increased cytokine levels (47).

Among the immune-response genes increased in atrial expression upon DHT treatment were

the Toll-like receptor 3 (*Tlr3*) and the Toll-like receptor 4 (*Tlr4*) in our study. Silencing or

deficiency of these receptors has been demonstrated to improve cardiac function post

myocardial infarction in rodent models by attenuating inflammatory cytokine production and

fibrotic scar formation (48, 49). Toll-like receptors can be activated by endogenous ligands,

including components of the extracellular matrix (50), and have been associated with matrix

turn-over.

Excessive ECM deposition is another pathological driver of conduction defects (51, 52) and

increased perivascular fibrosis has been observed in the ventricular myocardium of male rats

in response to 20 days of testosterone administration (53). Rho associated coiled-coil

containing protein kinase 2 (*Rock2*)-dependent pathways, implicated in cardiac fibroblast

activation, can regulate expression of ECM component levels, including CTGF (*Ccn2*) and α-

smooth muscle actin (*Acta2*) (54). Indeed, the expression of *Rock2, Ccn2,* as well as *Acta2*

transcript, was upregulated in atria after chronic DHT exposure. *Ccn2* expression, in

response to the pro-hypertrophic stimulus AngII, has previously been shown to be mediated

via the transcription factor serum response factor (*Srf*) in cardiac fibroblasts (55), which we

also found to be upregulated in atria subjected to high concentrations of DHT.

Transcriptional changes did not translate to structural changes other than atrial area dilation,

e.g. excessive ECM deposition in atria in our murine study, illustrating the

pathophysiologically relevant interaction of DHT-activated profibrotic signalling with

desmosomal gene defects for the development of atrial conduction slowing.

In the scope of this study, we were able to demonstrate that the combination of a pro-

inflammatory/fibrotic environment, caused by DHT, plus the selective reduction in I_{Na} in

Plako$^{+/-}$ cardiomyocytes is sufficient to induce conduction slowing and increased beat-to-beat

heterogeneity in these vulnerable atria.

While this is the first study showing the molecular effect on *atria*, it has been previously

demonstrated that androgens, including DHT, induce a hypertrophic response in *ventricular*

cardiomyocytes (12). One of the genes consistently elevated in atrial expression after DHT

exposure was insulin growth factor 1 (*Igf1*), a known driver of (cardiac) muscle growth (56,

57). *Igf1* mRNA expression was also found to contribute to atrial fibrotic remodelling and AF

inducibility in a rodent model (58). *IGF1*, containing an androgen response element within its

promoter region (59), is a well-established target of androgen-receptor mediated gene

activation. To our knowledge, we are the first to show that supraphysiological plasma DHT

concentration can induce upregulation of *Igf1* transcript levels in atria. Further evidence of the

pro-hypertrophic atrial gene response to DHT exposure can be based on the downregulation

of the lncRNA Myosin heavy chain associated RNA transcript (*Mhrt*). Repression of *Mhrt*

expression has previously been established as a prerequisite for stress-induced, pathological

cardiac hypertrophy and restoring *Mhrt* expression levels protected murine hearts from

pressure overload-induced hypertrophy (60). Mhrt has been shown to inhibit expression of

the transcription factor myocardin (*Myocd*) (61) and increased *Myocd* expression in peripheral

blood cells of patients has been associated with increased ventricular mass (62). In

accordance with this, down-regulation of *Mhrt* expression upon DHT exposure was paralleled

by upregulation of *Myocd* expression compared to control groups. Since we were not able to

demonstrate a significant increase in atrial cell diameter in histology at the investigated time

point, but were able to demonstrate increased cell capacitance in atrial cardiomyocytes on

single cell level as well as atrial area dilation, we propose that the atrial hypertrophic growth

response to DHT seen is of an eccentric nature.

Reduced plakoglobin decreases left atrial Na$_v$1.5 cluster availability at the membrane in

response to pro-hypertrophic DHT exposure

Super-resolution imaging and electrophysiological techniques suggest that Na$_v$1.5 spatial

sarcolemmal localization is not random, but rather characterized by formation of distinct

clusters (63, 64). To our knowledge, this the first study to evaluate the molecular arrangement

of sarcolemma-localized Na$_v$1.5 channels in atria through the use of dSTORM to allow for

visualisation at nanometer resolution (localisation accuracy estimated to be within 20-30 nm).

This confirmed that the mismatch between cell size and I$_{Na}$ density following DHT exposure in

Plako$^{+/-}$ left atria was due to reduced membrane Na$_v$1.5 channel number and/or defective

cluster availability. Interestingly, the values we estimate for next-neighbour distance,

calculated from cluster density in WT cells, are slightly lower than those reported employing

similar super-resolution methods in ventricular cardiac myocytes (64), suggesting slightly

higher cluster densities in atrial cardiomyocytes.

The number of plasma membrane-localized $Na_v1.5$ detections was reduced in left atrial

cardiomyocytes obtained from $Plako^{+/-}$ DHT and showed a similar trend in $Plako^{+/-}$ control. A

link between plakoglobin and $Na_v1.5$ channel incorporation/trafficking into the cell membrane

at the intercalated disc has been reported in ventricles (65, 66). Interactions of desmosomal

and intercalated disc proteins with sodium channel complex have been demonstrated in

ventricles previously (67) and loss or mutations of the desmosomal components plakophilin-2

and desmoglein-2 were associated with reduced sodium current in different cell and murine

models (67-69). A mismatch in cell size and $Na_v1.5$ cluster availability could have implications

for other cardiomyopathies and pathological hypertrophy (70, 71). Understanding the role of

ion channel cluster properties is in its infancy, but our observations are consistent with other

recent studies reporting a reduced I_{Na} in response to changes in single-molecule $Na_v1.5$

organisation (72, 73).

Implications for patients and athletes

Our results show that atrial arrhythmias are an important clinical feature of ARVC and confirm

male patients are more likely to show a full ARVC phenotype. We demonstrate a previously

unknown interaction between defective desmosomal gene expression and exposure to

androgenic anabolic steroids (AAS) in atria. This may partially explain the occurrence of atrial

conduction slowing and arrhythmias (74-76) in athletes abusing AAS to enhance their

performance. Based on our results, searching for desmosomal gene defects in steroid

abusers with atrial arrhythmias seems warranted. Such analyses may add to a better

understanding of the cardiac damage observed in some of these patients.

Our data, gained from well-controlled murine experiments, demonstrate that reduced

expression of plakoglobin, as commonly observed in cardiac tissue of patients with

pathogenic mutations in a variety of desmosomal genes (77), renders atria susceptible to

AAS-induced pathology. Prevention of atrial arrhythmias in arrhythmogenic cardiomyopathies

is of interest also because they can give rise to inappropriate defibrillator shocks in affected

patients (4, 78) and compromise cardiac function.

We here add exposure to DHT to the list of stimuli aggravating pro-arrhythmic phenotypes in

carriers of desmosomal mutations and demonstrate that this affects atrial electrical function.

Our data also provide an explanation for the stronger phenotypic expression in male gene

carriers with desmosomal mutations and the observed worsened clinical outcome in ARVC

patients with high physiological testosterone levels (20, 22-25).

Methods

See supplement for full methods.

Patient record screening for atrial arrhythmias and semi-automated analysis of digital electrocardiograms (ECGs) from ARVC patients

Adult ARVC patients (>18 years of age) seen at a specialty clinic at a tertiary centre between 2010 and 2021 were classified into two disease severity groups: non-definite and definite cases based on 2010 ARVC Task Force Criteria (TFC) (1).

Clinical records were retrospectively reviewed to obtain information from several modalities including imaging, electrophysiology, histopathology, genetic testing and family history, to cumulatively fulfil a diagnostic classification. Patients exhibiting signs of confirmed disease based on specific combinations of minor or major criteria were classified as the "definite" group. Individuals exhibiting signs on diagnostic investigation congruous with the 2010 TFC as "borderline" or "possible" ARVC were cumulatively considered as the "non-definite" group, as they do not confer a confirmed diagnosis of ARVC. Non-definite cases were included in the analysis to represent individuals in the earlier phases of disease, with a less severe profile of phenotypic expression. Atrial fibrillation and flutter status was extracted from previous electrocardiograms (ECG) and clinical letters.

Digital ECG recordings (10 seconds, sampling frequency 500 Hz) from the most recent follow-up in Inherited Cardiac Conditions Clinic were collated and analyzed using Matlab and BioSigKit (https://doi.org/10.21105/joss.00671). To discern the extent of atrial involvement in definite ARVC patients in comparison with non-definite patients as well as controls, family members of index patients without meeting TFC and/or exclusion of ARVC pathogenic variants attained from targeted gene panel testing, were additionally included ("Control").

Digital ECG analysis was performed by three independent observers in recordings displaying sinus rhythm applying a custom-designed, semi-automated algorithm. ECGs were digitally filtered between 0.5 and 50 Hz and using a Chebyshev type II filter. The R wave was

automatically identified, and all complexes within the recording were averaged to improve

signal quality. The isoelectric line was defined from P wave start to end.

Animal husbandry

Wildtype (WT) and plakoglobin deficient (Plako$^{+/-}$) littermate male mice (28), 129/Sv

background, were housed in individually ventilated cages, (2-7 mice/ cage), monitored daily

under standard conditions: 12 h light/dark cycle, 22±2 °C and 55±10% humidity. Food and

water were available *ad libitum.*

Chronic DHT exposure in the murine model and experimental timeline

Young adult male mice (8-11 weeks) were assigned to either DHT or placebo/control

treatment groups in mixed cages and were fitted with subcutaneous osmotic mini-pumps

(Alzet 2006), containing either DHT (62.5 mg/mL in ethanol), or solvent alone (Control, Ctr),

for 6 weeks (Figure 2). Age and DHT exposure time were matched for all groups. At least 40

minutes before pump implantation, mice were subcutaneously injected with 0.05 mL

of Buprenorphine. Pump implant was performed under anaesthesia with isoflurane inhalation

(max. 4%) in O_2 with a flow rate of 1-2 L/min. Animal handling staff and investigators were

blinded to genotype and treatment. Echocardiography and ECG recording was performed at

6 weeks exposure. Murine hearts were then extracted by thoracotomy under deep terminal

anaesthesia (4-5% isoflurane in O_2, flow rate 1-2 L/min,), and used for *in organ* and *in vitro*

experimental analysis.

Anabolic androgenic steroid measurements

Murine serum DHT concentrations were determined by ultra-performance liquid

chromatography-tandem mass spectrometry (LC-MS/MS) (Waters, Milford, MA, USA) as

described before (79).

Murine awake ECG measurements

Murine ECGs were recorded from awake mice using a tunnel system (ecgTunnel, EMKA Technologies, France) as reported previously (80). Analyses were performed on compound potentials averaged from 20 beats taken at three time-points throughout a 5-min recording with comparable heart rates between groups using the EMKA ECG analysis software. Obtained values were then averaged per animal. For P-wave duration the monophasic part of the P-wave was only taken from ECGs displaying a stable isoelectric line.

Optical mapping of murine left atria

Activation and action potential duration (APD) maps were generated from isolated left atria loaded with the voltage-sensitive dye Di-4-ANEPPS (17.5 µM; Cambridge Bioscience, CA, USA), paced over a range of 120-80 ms cycle length (CL) as described (81-83). To analyze conduction changes in more detail, beat-to-beat variability in whole tissue activation times was evaluated during rapid physiological pacing. To do this, 10 individual activation maps were compared from the final 10 beats of a train of 50 pulses at 80 ms CL (84).

Murine echocardiography

Echocardiography was performed with a dedicated small animal system (Vevo 2100; Visualsonics Fujifilm, Toronto, Ont, Canada) under light anaesthesia (0.5-2% isoflurane in O_2) at a target heart rate of 390-440 bpm (27, 85-87). The investigators were blinded to intervention and genotype. Images were analyzed by a second blinded observer.

Transmembrane action potential recordings

Transmembrane action potentials were recorded from isolated, superfused murine left atria as described previously (81-83, 88).

Cellular electrophysiology

Individual murine left atrial cardiac myocytes were isolated by perfusion with a Tyrode's enzyme solution containing 20 µg/mL Liberase™ (Roche, Indianapolis, IN) or a

collagenase+protease mix, 20 mM taurine and 30 µM $CaCl_2$, via Langendorff over a period of

10-15 min (81, 83). I_{Na} were evoked in voltage-clamp mode using standard protocols and low

Na^+ solution (89). All currents were normalized to cell capacitance.

Murine atrial fiber size and collagen composition

10 µm left atrial transverse sections were stained with FITC-conjugated wheat germ

agglutinin (lectin) and cell diameter as well as endomysial fibrosis were quantified using an

automated analysis tool (90).

Super-resolution microscopy and $Na_v1.5$ cluster analysis

Freshly isolated murine left atrial cardiac myocytes were plated on 10mm diameter laminin-

coated coverslips (Mattek, 35 mm dish, 1.5# coverglass), fixed, permeabilized and blocked.

Cells were stained with primary rabbit anti-$Na_v1.5$ antibody (ASC-005, 1:50, Alomone

Laboratories, Jerusalem, Israel) and after additional washing and blocking they were

incubated in secondary antibody (F(ab')2- Goat anti-Rabbit IgG, Alexa Fluor 647, A212-56,

1:1000, ThermoFisher Scientific, Waltham, MA, USA). Direct stochastic optical reconstruction

microscopy (dSTORM) (91) experiments were performed on a NIKON Eclipse Ti inverted N-

STORM microscope equipped with a NIKON APO 100 x 1.49 NA total internal reflection

fluorescence (TIRF) oil immersion objective. Immunolabelled samples were imaged in 0.5

mg/mL glucose oxidase, 40 µg/mL catalase, 10% wt/vol glucose and 100 mM MEA in PBS,

pH 7.4 to induce Alexa 647 blinking. During dSTORM acquisition, the sample was

continuously illuminated at 640 nm for 20,000 frames. Final rendered images of the localized

molecules were generated using ThunderSTORM (92) and false coloured in FIJI. Final

detections were subjected to a segmentation protocol, using a persistence-based clustering

approach (93).

RNA sequencing of murine left atria

Poly(A) mRNA-enriched libraries were sequenced 75 cycles in a single read mode on the NextSeq-500 System (v2.5 Chemistry, Illumina). Acquired data was trimmed and FASTQ files were aligned using HISAT2 (version 2.1.0; ref Pertea) and the reference genome Ensembl Mus Musculus GRCm38. Differential expression analysis were performed in R (http://www.R-project.org/, R 3.4.1) using the DESeq2 package (94). Differential expression with a false discovery rate <0.05 was deemed significant. Results were visualized using R.

Statistics

All experiments and analyses were performed blinded to genotype and treatment. Data from murine studies was first subjected to outlier analysis, ROUT method, based around a false discovery rate, where α = 0.01 and outliers were removed (Prism v8, GraphPad Software, La Jolla, CA, USA). Associations between categorical variables were analyzed by Fisher´s exact test. Significance between groups for normally distributed data was taken as p<0.05, ordinary one-way or two-way (repeated) measures ANOVA, with Bonferroni's post hoc test, as appropriate. Non-normally distributed data was subjected to Kruskal-Wallis with significance level of p<0.05 and Dunn's post hoc test (Prism v8, GraphPad Software, La Jolla, CA, USA).

Study approval

Ethical approval for analysis of clinical data from Inherited Cardiac Conditions clinic at the University Hospital Queen Elisabeth, Birmingham, was granted by the local department of research ethics at UHB Hospital (Audit number CARMS-16044).

All animal procedures were approved by the UK Home Office (PPL number 30/2967 and PFDAAF77F) and by the institutional review board of University of Birmingham, UK. All animal procedures conformed to the guidelines from Directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes.

Author contributions

LCS performed murine *in vivo* experiments, dSTORM experiments and analysis, gravimetry and histology analysis, RNAseq analysis, P wave analysis in human ECGs and wrote the manuscript. APH performed and analyzed microelectrode, optical mapping, patchclamp and dSTORM studies, performed gravimetry analysis and wrote the manuscript. TYY and COS set up and performed and analyzed optical mapping experiments. DMK designed dSTORM experiments and trained LCS and APH. JMP generated cluster analysis workflows.TW and PMM performed histology. FS codesigned experiments and trained TW, TYY, SNK. AA and TK screened clinical records according to Task Force Criteria. LCS, TK and COS performed semi-automated patient ECG analysis. CH prepared RNA samples and co-supervised PMM. VRC and GVG analyzed RNAseq. MS and AW performed and analyzed RNAseq. SNK and SBS performed *in vivo* experiments and gravimetry. MOR co-supervised TK. LFo analyzed *in vivo* experiments. SL trained LCS and co-supervised PMM. AK performed mass spectrometry of serum DHT concentrations. WA and GGL advised on AAS experimental design, DP co-supervised COS, RS co-supervised AA, KG co-supervised LCS. PK provided input throughout and co-supervised APH, FS, CH, VRC. LFa (co)-supervised LCS, APH, TYY, COS, TW, FS, AA, TK,VRC, SNK, CH, PMM, SBS, MOR, LFo, designed and coordinated the study and wrote the manuscript. All authors reviewed the results, revised the manuscript and approved the final version of it.

Acknowledgements

We thank Clara Apicella, Olivia Grech, Pushpa Patel, Genna Riley, and staff of BMSU Birmingham for expert support. We thankfully acknowledge the Centre of Membrane Proteins and Receptors COMPARE (www.birmingham-nottingham.ac.uk/compare) for both their expertise and infrastructure. We thank the Core Facility Genomics of the Medical Faculty Münster, University of Muenster. We thank all members of the Translational Research in Heart Failure and Arrhythmias Cluster for discussion.

This work was partially supported by European Union (grant agreement No 633196 [CATCH ME] to PK and LF), European Union BigData@Heart (grant agreement EU IMI 116074), British Heart Foundation (FS/13/43/30324 to PK and LF; PG/17/30/32961 to PK and APH, PG/20/22/35093 to PK; AA/18/2/34218 to PK and LF; FS/12/40/29712 to KG), German Centre for Cardiovascular Research supported by the German Ministry of Education and Research (DZHK to PK); Leducq Foundation to PK; UOB research development fund (LF). TYY studentship was supported by PSIBS to LF. Funding from the Wellcome Trust was received by KG (201543/B/16/Z) and WA (WT209492/Z/17/Z). KG is also funded by the MRC (MR/V009540/1). The Institute of Cardiovascular Sciences is a recipient of a BHF Accelerator Award (AA/18/2/34218).

GVG acknowledges support from the NIHR Birmingham ECMC, NIHR Birmingham SRMRC, Nanocommons H2020-EU (731032) and the MRC Heath Data Research UK (HDRUK/CFC/01), an initiative funded by UK Research and Innovation, Department of Health and Social Care (England) and the devolved administrations, and leading medical research charities. GVG and WA receive support from the NIHR Birmingham Biomedical Research Centre. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health and Care Research, the Medical Research Council or the Department of Health.

References

1.   Marcus FI, McKenna WJ, Sherrill D, Basso C, Bauce B, Bluemke DA, et al. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the Task Force Criteria. *European heart journal.* 2010;31(7):806-14.

2.   Morita H, Kusano-Fukushima K, Nagase S, Fujimoto Y, Hisamatsu K, Fujio H, et al. Atrial fibrillation and atrial vulnerability in patients with Brugada syndrome. *J Am Coll Cardiol.* 2002;40(8):1437-44.

3.   Tonet JL, Castro-Miranda R, Iwa T, Poulain F, Frank R, and Fontaine GH. Frequency of supraventricular tachyarrhythmias in arrhythmogenic right ventricular dysplasia. *The American journal of cardiology.* 1991;67(13):1153.

4.   Camm CF, James CA, Tichnell C, Murray B, Bhonsale A, te Riele AS, et al. Prevalence of atrial arrhythmias in arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Heart rhythm.* 2013;10(11):1661-8.

5.   Chu AF, Zado E, and Marchlinski FE. Atrial arrhythmias in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia and ventricular tachycardia. *The American journal of cardiology.* 2010;106(5):720-2.

6.   Baturova MA, Haugaa KH, Jensen HK, Svensson A, Gilljam T, Bundgaard H, et al. Atrial fibrillation as a clinical characteristic of arrhythmogenic right ventricular cardiomyopathy: Experience from the Nordic ARVC Registry. *Int J Cardiol.* 2020;298:39-43.

7.   Priori SG, Blomstrom-Lundqvist C, Mazzanti A, Blom N, Borggrefe M, Camm J, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *European heart journal.* 2015;36(41):2793-867.

8.   Sagoe D, Molde H, Andreassen CS, Torsheim T, and Pallesen S. The global epidemiology of anabolic-androgenic steroid use: a meta-analysis and meta-regression analysis. *Annals of Epidemiology.* 2014;24(5):383-98.

9.   Urhausen A, Albers T, and Kindermann W. Are the cardiac effects of anabolic steroid abuse in strength athletes reversible? *Heart.* 2004;90(5):496-501.

10.   Luijkx T, Velthuis BK, Backx FJG, Buckens CFM, Prakken NHJ, Rienks R, et al. Anabolic androgenic steroid use is associated with ventricular dysfunction on cardiac MRI in strength trained athletes. *International journal of cardiology.* 2013;167(3):664-8.

11.   Alizade E, Avci A, Fidan S, Tabakci M, Bulut M, Zehir R, et al. The Effect of Chronic Anabolic-Androgenic Steroid Use on Tp-E Interval, Tp-E/Qt Ratio, and Tp-E/Qtc Ratio in Male Bodybuilders. *Ann Noninvasive Electrocardiol.* 2015;20(6):592-600.

12.   Marsh JD, Lehmann MH, Ritchie RH, Gwathmey JK, Green GE, and Schiebinger RJ. Androgen receptors mediate hypertrophy in cardiac myocytes. *Circulation.* 1998;98(3):256-61.

13.   Medei E, Marocolo M, Rodrigues DD, Arantes PC, Takiya CM, Silva J, et al. Chronic treatment with anabolic steroids induces ventricular repolarization disturbances: Cellular, ionic and molecular mechanism. *Journal of molecular and cellular cardiology.* 2010;49(2):165-75.

14. Pirompol P, Teekabut V, Weerachatyanukul W, Bupha-Intr T, and
Wattanapermpool J. Supra-physiological dose of testosterone induces
pathological cardiac hypertrophy. *J Endocrinol.* 2016;229(1):13-23.
15. Berger D, Folsom AR, Schreiner PJ, Chen LY, Michos ED, O'Neal WT, et al. Plasma
total testosterone and risk of incident atrial fibrillation: The Atherosclerosis Risk
in Communities (ARIC) study. *Maturitas.* 2019;125:5-10.
16. Sullivan ML, Martinez CM, and Gallagher EJ. Atrial fibrillation and anabolic
steroids. *J Emerg Med.* 1999;17(5):851-7.
17. Tsai WC, Lee TI, Chen YC, Kao YH, Lu YY, Lin YK, et al. Testosterone replacement
increases aged pulmonary vein and left atrium arrhythmogenesis with enhanced
adrenergic activity. *International journal of cardiology.* 2014;176(1):110-8.
18. Lau DH, Stiles MK, John B, Shashidhar, Young GD, and Sanders P. Atrial fibrillation
and anabolic steroid abuse. *Int J Cardiol.* 2007;117(2):e86-7.
19. Rootwelt-Norberg C, Lie OH, Chivulescu M, Castrini AI, Sarvari SI, Lyseggen E, et
al. Sex differences in disease progression and arrhythmic risk in patients with
arrhythmogenic cardiomyopathy. *Europace : European pacing, arrhythmias, and
cardiac electrophysiology : journal of the working groups on cardiac pacing,
arrhythmias, and cardiac cellular electrophysiology of the European Society of
Cardiology.* 2021;23(7):1084-91.
20. Akdis D, Saguner AM, Shah K, Wei C, Medeiros-Domingo A, von Eckardstein A, et
al. Sex hormones affect outcome in arrhythmogenic right ventricular
cardiomyopathy/dysplasia: from a stem cell derived cardiomyocyte-based model
to clinical biomarkers of disease outcome. *European heart journal.* 2017.
21. Wilde AAM, Semsarian C, Marquez MF, Sepehri Shamloo A, Ackerman MJ, Ashley
EA, et al. European Heart Rhythm Association (EHRA)/Heart Rhythm Society
(HRS)/Asia Pacific Heart Rhythm Society (APHRS)/Latin American Heart Rhythm
Society (LAHRS) Expert Consensus Statement on the State of Genetic Testing for
Cardiac Diseases. *Heart rhythm.* 2022.
22. Asimaki A, Syrris P, Wichter T, Matthias P, Saffitz JE, and McKenna WJ. A novel
dominant mutation in plakoglobin causes Arrhythmogenic right ventricular
cardiomyopathy. *American journal of human genetics.* 2007;81(5):964-73.
23. Protonotarios N, Tsatsopoulou A, Anastasakis A, Sevdalis E, McKoy G, Stratos K, et
al. Genotype-phenotype assessment in autosomal recessive arrhythmogenic right
ventricular cardiomyopathy (Naxos disease) caused by a deletion in plakoglobin.
Journal of the American College of Cardiology. 2001;38(5):1477-84.
24. Antoniades L, Tsatsopoulou A, Anastasakis A, Syrris P, Asimaki A, Panagiotakos D,
et al. Arrhythmogenic right ventricular cardiomyopathy caused by deletions in
plakophilin-2 and plakoglobin (Naxos disease) in families from Greece and
Cyprus: genotype-phenotype relations, diagnostic features and prognosis.
European heart journal. 2006;27(18):2208-16.
25. McKoy G, Protonotarios N, Crosby A, Tsatsopoulou A, Anastasakis A, Coonar A, et
al. Identification of a deletion in plakoglobin in arrhythmogenic right ventricular
cardiomyopathy with palmoplantar keratoderma and woolly hair (Naxos
disease). *Lancet (London, England).* 2000;355(9221):2119-24.
26. Li J, Swope D, Raess N, Cheng L, Muller EJ, and Radice GL. Cardiac tissue-restricted
deletion of plakoglobin results in progressive cardiomyopathy and activation of
{beta}-catenin signaling. *Mol Cell Biol.* 2011;31(6):1134-44.
27. Fabritz L, Hoogendijk MG, Scicluna BP, van Amersfoorth SC, Fortmueller L, Wolf S,
et al. Load-reducing therapy prevents development of arrhythmogenic right

ventricular cardiomyopathy in plakoglobin-deficient mice. *J Am Coll Cardiol.*
2011;57(6):740-50.

28.   Kirchhof P, Fabritz L, Zwiener M, Witt H, Schafers M, Zellerhoff S, et al. Age- and
training-dependent development of arrhythmogenic right ventricular
cardiomyopathy in heterozygous plakoglobin-deficient mice. *Circulation.*
2006;114(17):1799-806.

29.   Weinsaft JW, Kochav JD, Kim J, Gurevich S, Volo SC, Afroz A, et al. P wave area for
quantitative electrocardiographic assessment of left atrial remodeling. *PLoS One.*
2014;9(6):e99178.

30.   Magnani JW, Zhu L, Lopez F, Pencina MJ, Agarwal SK, Soliman EZ, et al. P-wave
indices and atrial fibrillation: cross-cohort assessments from the Framingham
Heart Study (FHS) and Atherosclerosis Risk in Communities (ARIC) study. *Am*
Heart J. 2015;169(1):53-61 e1.

31.   Soliman EZ, Prineas RJ, Case LD, Zhang ZM, and Goff DC, Jr. Ethnic distribution of
ECG predictors of atrial fibrillation and its impact on understanding the ethnic
distribution of ischemic stroke in the Atherosclerosis Risk in Communities (ARIC)
study. *Stroke; a journal of cerebral circulation.* 2009;40(4):1204-11.

32.   Gorenek B, Birdane A, Kudaiberdieva G, Goktekin O, Cavusoglu Y, Unalir A, et al. P
wave amplitude and duration may predict immediate recurrence of atrial
fibrillation after internal cardioversion. *Ann Noninvasive Electrocardiol.*
2003;8(3):215-8.

33.   Kirchhof P. The future of atrial fibrillation management: integrated care and
stratified therapy. *Lancet.* 2017;390(10105):1873-87.

34.   Baturova MA, Svensson A, Aneq MA, Svendsen JH, Risum N, Sherina V, et al.
Evolution of P-wave indices during long-term follow-up as markers of atrial
substrate progression in arrhythmogenic right ventricular cardiomyopathy.
Europace : European pacing, arrhythmias, and cardiac electrophysiology : journal
of the working groups on cardiac pacing, arrhythmias, and cardiac cellular
electrophysiology of the European Society of Cardiology. 2021;23(23 Suppl 1):i29-
i37.

35.   Heijman J, Guichard JB, Dobrev D, and Nattel S. Translational Challenges in Atrial
Fibrillation. *Circulation research.* 2018;122(5):752-73.

36.   van Rijen HV, and de Bakker JM. Penetrance of monogenetic cardiac conduction
diseases. A matter of conduction reserve? *Cardiovasc Res.* 2007;76(3):379-80.

37.   van Rijen HV, de Bakker JM, and van Veen TA. Hypoxia, electrical uncoupling, and
conduction slowing: Role of conduction reserve. *Cardiovasc Res.* 2005;66(1):9-11.

38.   Campian ME, Verberne HJ, Hardziyenka M, de Groot EA, van Moerkerken AF, van
Eck-Smit BL, et al. Assessment of inflammation in patients with arrhythmogenic
right ventricular cardiomyopathy/dysplasia. *Eur J Nucl Med Mol Imaging.*
2010;37(11):2079-85.

39.   Chelko SP, Asimaki A, Lowenthal J, Bueno-Beti C, Bedja D, Scalco A, et al.
Therapeutic Modulation of the Immune Response in Arrhythmogenic
Cardiomyopathy. *Circulation.* 2019.

40.   Frustaci A, Chimenti C, Bellocci F, Morgante E, Russo MA, and Maseri A.
Histological substrate of atrial biopsies in patients with lone atrial fibrillation.
Circulation. 1997;96(4):1180-4.

41.   Chung MK, Martin DO, Sprecher D, Wazni O, Kanderian A, Carnes CA, et al. C-
reactive protein elevation in patients with atrial arrhythmias: inflammatory

mechanisms and persistence of atrial fibrillation. *Circulation.* 2001;104(24):2886-
91.

42. Aviles RJ, Martin DO, Apperson-Hansen C, Houghtaling PL, Rautaharju P, Kronmal
632 RA, et al. Inflammation as a risk factor for atrial fibrillation. *Circulation.*
633 2003;108(24):3006-10.

43. Xu J, Cui G, Esmailian F, Plunkett M, Marelli D, Ardehali A, et al. Atrial extracellular
635 matrix remodeling and the maintenance of atrial fibrillation. *Circulation.*
636 2004;109(3):363-8.

44. Boldt A, Wetzel U, Lauschke J, Weigl J, Gummert J, Hindricks G, et al. Fibrosis in
638 left atrial tissue of patients with atrial fibrillation with and without underlying
639 mitral valve disease. *Heart.* 2004;90(4):400-5.

45. Marcus FI, Fontaine GH, Guiraudon G, Frank R, Laurenceau JL, Malergue C, et al.
641 Right ventricular dysplasia: a report of 24 adult cases. *Circulation.*
642 1982;65(2):384-98.

46. Thiene G, Corrado D, Nava A, Rossi L, Poletti A, Boffa GM, et al. Right ventricular
644 cardiomyopathy: is there evidence of an inflammatory aetiology? *European heart*
645 *journal.* 1991;12 Suppl D:22-5.

47. Galan M, Varona S, Guadall A, Orriols M, Navas M, Aguilo S, et al. Lysyl oxidase
647 overexpression accelerates cardiac remodeling and aggravates angiotensin II-
648 induced hypertrophy. *FASEB J.* 2017;31(9):3787-99.

48. Lu C, Ren D, Wang X, Ha T, Liu L, Lee EJ, et al. Toll-like receptor 3 plays a role in
650 myocardial infarction and ischemia/reperfusion injury. *Biochimica et biophysica*
651 *acta.* 2014;1842(1):22-31.

49. Liu L, Wang Y, Cao ZY, Wang MM, Liu XM, Gao T, et al. Up-regulated TLR4 in
653 cardiomyocytes exacerbates heart failure after long-term myocardial infarction. *J*
654 *Cell Mol Med.* 2015;19(12):2728-40.

50. Okamura Y, Watari M, Jerud ES, Young DW, Ishizaka ST, Rose J, et al. The extra
656 domain A of fibronectin activates Toll-like receptor 4. *J Biol Chem.*
657 2001;276(13):10229-33.

51. Verheule S, Sato T, Everett T, Engle SK, Otten D, von der Lohe MR, et al. Increased
659 vulnerability to atrial fibrillation in transgenic mice with selective atrial fibrosis
660 caused by overexpression of TGF-beta 1. *Circulation research.* 2004;94(11):1458-
661 65.

52. Fabritz L, and Kirchhof P. Selective atrial profibrotic signalling in mice and man.
663 *Cardiovasc Res.* 2013;99(4):592-4.

53. Papamitsou T, Barlagiannis D, Papaliagkas V, Kotanidou E, and Dermentzopoulou-
665 Theodoridou M. Testosterone-induced hypertrophy, fibrosis and apoptosis of
666 cardiac cells--an ultrastructural and immunohistochemical study. *Med Sci Monit.*
667 2011;17(9):BR266-73.

54. Akhmetshina A, Dees C, Pileckyte M, Szucs G, Spriewald BM, Zwerina J, et al. Rho-
669 associated kinases are crucial for myofibroblast differentiation and production of
670 extracellular matrix in scleroderma fibroblasts. *Arthritis Rheum.*
671 2008;58(8):2553-64.

55. Ongherth A, Pasch S, Wuertz CM, Nowak K, Kittana N, Weis CA, et al. p63RhoGEF
673 regulates auto- and paracrine signaling in cardiac fibroblasts. *J Mol Cell Cardiol.*
674 2015;88:39-54.

56. Kim J, Wende AR, Sena S, Theobald HA, Soto J, Sloan C, et al. Insulin-like growth
676 factor I receptor signaling is required for exercise-induced cardiac hypertrophy.
677 *Mol Endocrinol.* 2008;22(11):2531-43.

679   57.   Weeks KL, Bernardo BC, Ooi JYY, Patterson NL, and McMullen JR. The IGF1-PI3K-
680   Akt Signaling Pathway in Mediating Exercise-Induced Cardiac Hypertrophy and
681   Protection. *Adv Exp Med Biol.* 2017;1000:187-210.

682   58.   Wang J, Li Z, Du J, Li J, Zhang Y, Liu J, et al. The expression profile analysis of atrial
683   mRNA in rats with atrial fibrillation: the role of IGF1 in atrial fibrosis. *BMC*
684   *Cardiovasc Disord.* 2019;19(1):40.

685   59.   Wu Y, Zhao W, Zhao J, Pan J, Wu Q, Zhang Y, et al. Identification of androgen
686   response elements in the insulin-like growth factor I upstream promoter.
687   *Endocrinology.* 2007;148(6):2984-93.

688   60.   Han P, Li W, Lin CH, Yang J, Shang C, Nuernberg ST, et al. A long noncoding RNA
689   protects the heart from pathological hypertrophy. *Nature.* 2014;514(7520):102-6.

690   61.   Luo Y, Xu Y, Liang C, Xing W, and Zhang T. The mechanism of myocardial
691   hypertrophy regulated by the interaction between mhrt and myocardin. *Cell*
692   *Signal.* 2018;43:11-20.

693   62.   Kontaraki JE, Parthenakis FI, Patrianakos AP, Karalis IK, and Vardas PE. Altered
694   expression of early cardiac marker genes in circulating cells of patients with
695   hypertrophic cardiomyopathy. *Cardiovasc Pathol.* 2007;16(6):329-35.

696   63.   Bhargava A, Lin XM, Novak P, Mehta K, Korchev Y, Delmar M, et al. Super-
697   resolution Scanning Patch Clamp Reveals Clustering of Functional Ion Channels in
698   Adult Ventricular Myocyte. *Circulation research.* 2013;112(8):1112-+.

699   64.   Leo-Macias A, Agullo-Pascual E, Sanchez-Alonso JL, Keegan S, Lin XM, Arcos T, et
700   al. Nanoscale visualization of functional adhesion/excitability nodes at the
701   intercalated disc. *Nature communications.* 2016;7:11.

702   65.   Asimaki A, Kapoor S, Plovie E, Arndt AK, Adams E, Liu ZZ, et al. Identification of a
703   New Modulator of the Intercalated Disc in a Zebrafish Model of Arrhythmogenic
704   Cardiomyopathy. *Sci Transl Med.* 2014;6(240):15.

705   66.   Noorman M, Hakim S, Kessler E, Groeneweg JA, Cox M, Asimaki A, et al.
706   Remodeling of the cardiac sodium channel, connexin43, and plakoglobin at the
707   intercalated disk in patients with arrhythmogenic cardiomyopathy. *Heart*
708   *rhythm : the official journal of the Heart Rhythm Society.* 2013;10(3):412-9.

709   67.   Rizzo S, Lodder EM, Verkerk AO, Wolswinkel R, Beekman L, Pilichou K, et al.
710   Intercalated disc abnormalities, reduced Na(+) current density, and conduction
711   slowing in desmoglein-2 mutant mice prior to cardiomyopathic changes.
712   *Cardiovascular research.* 2012;95(4):409-18.

713   68.   Sato PY, Musa H, Coombs W, Guerrero-Serna G, Patino GA, Taffet SM, et al. Loss of
714   plakophilin-2 expression leads to decreased sodium current and slower
715   conduction velocity in cultured cardiac myocytes. *Circ Res.* 2009;105(6):523-6.

716   69.   Cerrone M, Lin X, Zhang M, Agullo-Pascual E, Pfenniger A, Chkourko Gusky H, et al.
717   Missense mutations in plakophilin-2 cause sodium current deficit and associate
718   with a Brugada syndrome phenotype. *Circulation.* 2014;129(10):1092-103.

719   70.   Hofmann F, Fabritz L, Stieber J, Schmitt J, Kirchhof P, Ludwig A, et al. Ventricular
720   HCN channels decrease the repolarization reserve in the hypertrophic heart.
721   *Cardiovasc Res.* 2012;95(3):317-26.

722   71.   Wagner S, Dybkova N, Rasenack EC, Jacobshagen C, Fabritz L, Kirchhof P, et al.
723   Ca2+/calmodulin-dependent protein kinase II regulates cardiac Na+ channels. *J*
724   *Clin Invest.* 2006;116(12):3127-38.

725   72.   Agullo-Pascual E, Lin XM, Leo-Macias A, Zhang ML, Liang FX, Li Z, et al. Super-
726   resolution imaging reveals that loss of the C-terminus of connexin43 limits

microtubule plus-end capture and Na(V)1.5 localization at the intercalated disc.
Cardiovascular research. 2014;104(2):371-81.

73.   te Riele A, Agullo-Pascual E, James CA, Leo-Macias A, Cerrone M, Zhang ML, et al.
Multilevel analyses of SCN5A mutations in arrhythmogenic right ventricular
dysplasia/cardiomyopathy suggest non-canonical mechanisms for disease
pathogenesis. *Cardiovascular research.* 2017;113(1):102-11.

74.   Furlanello F, Serdoz LV, Cappato R, and De Ambroggi L. Illicit drugs and cardiac
arrhythmias in athletes. *Eur J Cardiovasc Prev Rehabil.* 2007;14(4):487-94.

75.   Akcakoyun M, Alizade E, Gundogdu R, Bulut M, Tabakci MM, Acar G, et al. Long-
term anabolic androgenic steroid use is associated with increased atrial
electromechanical delay in male bodybuilders. *Biomed Res Int.*
2014;2014:451520.

76.   Nieschlag E, and Vorona E. Doping with anabolic androgenic steroids (AAS):
Adverse effects on non-reproductive organs and functions. *Rev Endocr Metab*
Disord. 2015;16(3):199-211.

77.   Asimaki A, Tandri H, Huang H, Halushka MK, Gautam S, Basso C, et al. A new
diagnostic test for arrhythmogenic right ventricular cardiomyopathy. *N Engl J*
Med. 2009;360(11):1075-84.

78.   Takehara N, Makita N, Kawabe J, Sato N, Kawamura Y, Kitabatake A, et al. A
cardiac sodium channel mutation identified in Brugada syndrome associated with
atrial standstill. *J Intern Med.* 2004;255(1):137-42.

79.   Kulle AE, Riepe FG, Melchior D, Hiort O, and Holterhus PM. A novel ultrapressure
liquid chromatography tandem mass spectrometry method for the simultaneous
determination of androstenedione, testosterone, and dihydrotestosterone in
pediatric blood samples: age- and sex-specific reference data. *J Clin Endocrinol*
Metab. 2010;95(5):2399-409.

80.   Silbernagel N, Walecki M, Schafer MK, Kessler M, Zobeiri M, Rinne S, et al. The
VAMP-associated protein VAPB is required for cardiac and neuronal pacemaker
channel function. *FASEB J.* 2018;32(11):6159-73.

81.   Holmes AP, Yu TY, Tull S, Syeda F, Kuhlmann SM, O'Brien S-M, et al. A Regional
Reduction in Ito and IKACh in the Murine Posterior Left Atrial Myocardium Is
Associated with Action Potential Prolongation and Increased Ectopic Activity. *PloS*
one. 2016;11(5):e0154077.

82.   Yu TY, Syeda F, Holmes AP, Osborne B, Dehghani H, Brain KL, et al. An automated
system using spatial oversampling for optical mapping in murine atria.
Development and validation with monophasic and transmembrane action
potentials. *Progress in biophysics and molecular biology.* 2014.

83.   Syeda F, Holmes AP, Yu TY, Tull S, Kuhlmann SM, Pavlovic D, et al. PITX2
Modulates Atrial Membrane Potential and the Antiarrhythmic Effects of Sodium-
Channel Blockers. *J Am Coll Cardiol.* 2016;68(17):1881-94.

84.   O'Shea C, Holmes AP, Yu TY, Winter J, Wells SP, Correia J, et al. ElectroMap: High-
throughput open-source software for analysis and mapping of cardiac
electrophysiology. *Scientific reports.* 2019;9(1):1389.

85.   Kirchhof P, Kahr PC, Kaese S, Piccini I, Vokshi I, Scheld HH, et al. PITX2c is
expressed in the adult left atrium, and reducing Pitx2c expression promotes atrial
fibrillation inducibility and complex changes in gene expression. *Circ Cardiovasc*
Genet. 2011;4(2):123-33.

86.   Blana A, Kaese S, Fortmuller L, Laakmann S, Damke D, van Bragt K, et al. Knock-in
gain-of-function sodium channel mutation prolongs atrial action potentials and
alters atrial vulnerability. *Heart Rhythm.* 2010;7(12):1862-9.

87.   Fabritz L, Kirchhof P, Fortmuller L, Auchampach JA, Baba HA, Breithardt G, et al.
Gene dose-dependent atrial arrhythmias, heart block, and brady-cardiomyopathy
in mice overexpressing A(3) adenosine receptors. *Cardiovasc Res.*
2004;62(3):500-8.

88.   Lemoine MD, Duverger JE, Naud P, Chartier D, Qi XY, Comtois P, et al.
Arrhythmogenic left atrial cellular electrophysiology in a murine genetic long QT
syndrome model. *Cardiovasc Res.* 2011;92(1):67-74.

89.   S OB, Holmes AP, Johnson DM, Kabir SN, C OS, M OR, et al. Increased atrial
effectiveness of flecainide conferred by altered biophysical properties of sodium
channels. *J Mol Cell Cardiol.* 2022;166:23-35.

90.   Winters J, von Braunmuhl ME, Zeemering S, Gilbers M, Brink TT, Scaf B, et al.
JavaCyte, a novel open-source tool for automated quantification of key hallmarks
of cardiac structural remodeling. *Scientific reports.* 2020;10(1):20074.

91.   Kavanagh DM, Smyth AM, Martin KJ, Dun A, Brown ER, Gordon S, et al. A
molecular toggle after exocytosis sequesters the presynaptic syntaxin1a
molecules involved in prior vesicle fusion. *Nature communications.* 2014;5:14.

92.   Ovesny M, Krizek P, Borkovec J, Svindrych ZK, and Hagen GM. ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-
resolution imaging. *Bioinformatics.* 2014;30(16):2389-90.

93.   Pike JA, Khan AO, Pallini C, Thomas SG, Mund M, Ries J, et al. Topological data
analysis quantifies biological nano-structure from single molecule localization
microscopy. *Bioinformatics (Oxford, England).* 2020;36(5):1614-21.

94.   Love MI, Huber W, and Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

95.   Jaoude SA, Leclercq JF, and Coumel P. Progressive ECG changes in arrhythmogenic
right ventricular disease. Evidence for an evolving disease. *European heart*
journal. 1996;17(11):1717-22.

96.   Brembilla-Perrot B, Jacquemin L, Houplon P, Houriez P, Beurrier D, Berder V, et al.
Increased atrial vulnerability in arrhythmogenic right ventricular disease. *Am*
Heart J. 1998;135(5 Pt 1):748-54.

97.   Peters S, Trummel M, and Meyners W. Prevalence of right ventricular dysplasia-
cardiomyopathy in a non-referral hospital. *Int J Cardiol.* 2004;97(3):499-501.

98.   Saguner AM, Ganahl S, Kraus A, Baldinger SH, Medeiros-Domingo A, Saguner AR,
et al. Clinical role of atrial arrhythmias in patients with arrhythmogenic right
ventricular dysplasia. *Circulation journal : official journal of the Japanese*
Circulation Society. 2014;78(12):2854-61.

99.   Wu L, Guo J, Zheng L, Chen G, Ding L, Qiao Y, et al. Atrial Remodeling and Atrial
Tachyarrhythmias in Arrhythmogenic Right Ventricular Cardiomyopathy. *The*
American journal of cardiology. 2016;118(5):750-3.

100.   Bourfiss M, Te Riele AS, Mast TP, Cramer MJ, JF VDH, TA VANV, et al. Influence of
Genotype on Structural Atrial Abnormalities and Atrial Fibrillation or Flutter in
Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy. *J Cardiovasc*
Electrophysiol. 2016;27(12):1420-8.

101.   Mazzanti A, Ng K, Faragli A, Maragna R, Chiodaroli E, Orphanou N, et al.
Arrhythmogenic Right Ventricular Cardiomyopathy: Clinical Course and
Predictors of Arrhythmic Risk. *J Am Coll Cardiol.* 2016;68(23):2540-50.

102.   Gilljam T, Haugaa KH, Jensen HK, Svensson A, Bundgaard H, Hansen J, et al. Heart
824   transplantation in arrhythmogenic right ventricular cardiomyopathy - Experience
825   from the Nordic ARVC Registry. *Int J Cardiol.* 2018;250:201-6.

826   103.   Wu L, Bao J, Liang E, Fan S, Zheng L, Du Z, et al. Atrial involvement in
827   arrhythmogenic right ventricular cardiomyopathy patients referred for
828   ventricular arrhythmias ablation. *J Cardiovasc Electrophysiol.* 2018;29(10):1388-
829   95.

830   104.   Mussigbrodt A, Knopp H, Efimova E, Weber A, Bertagnolli L, Hilbert S, et al.
831   Supraventricular arrhythmias in patients with arrhythmogenic right ventricular
832   dysplasia/cardiomyopathy associate with long-term outcome after catheter
833   ablation of ventricular tachycardias. *Europace : European pacing, arrhythmias, and*
834   *cardiac electrophysiology : journal of the working groups on cardiac pacing,*
835   *arrhythmias, and cardiac cellular electrophysiology of the European Society of*
836   *Cardiology.* 2018;20(7):1182-7.

837   105.   Cardona-Guarache R, Astrom-Aneq M, Oesterle A, Asirvatham R, Svetlichnaya J,
838   Marcus GM, et al. Atrial arrhythmias in patients with arrhythmogenic right
839   ventricular cardiomyopathy: Prevalence, echocardiographic predictors, and
840   treatment. *J Cardiovasc Electrophysiol.* 2019;30(10):1801-10.

841   106.   Kikuchi N, Shiga T, Suzuki A, and Hagiwara N. Atrial tachyarrhythmias and heart
842   failure events in patients with arrhythmogenic right ventricular cardiomyopathy.
843   *Int J Cardiol Heart Vasc.* 2020;31:100669.

Figures and figure legends

Figure 1 - Arrhythmogenic Right Ventricular Cardiomyopathy (ARVC) patients' and control individuals' P wave characteristics derived from digital ECG analysis
(**A**) Lead II ECG recordings from an unaffected (control) as well as a non-definite and definite ARVC patient. Individual cardiac cycles over a duration of 10 sec (grey traces) are overlaid by detected R waves and averaged (black trace). PR interval (blue), P wave duration (orange) and P wave area (green) are marked. (**B**) PR interval and P wave characteristics obtained from semi-automated analysis of the averaged ECG. All parameters are derived from lead II recordings. Mean heart rate ± SEM: Control: 73 ± 3 bpm ; Non-definite ARVC: 75 ± 2 bpm; Definite ARVC: 60 ± 3 bpm. P-values from post hoc tests are reported on the graphs (Kruskal-Wallis ($p<0.05$) with Dunn's post hoc test for PR interval; one-way ANOVA ($p<0.05$) with Bonferroni post hoc test for P wave duration and area). n (number of patients) = Control: 13, Non-definite ARVC: 42, Definite ARVC: 25

A

B

C

Figure 2 - Study design for murine experiments
(**A**) Experimental timeline and methods used in the underlying murine study. All results described were obtained in male WT or Plako+/- mice at 4 months of age, at the end point of the protocol, i.e. after exposure to either 5α-dihydrotestosterone (DHT) or placebo control (Ctr) for 6 weeks. (**B**) Serum DHT concentration (n=11-18 mice/group) and (**C**) body weight (n=48-54 mice/group) are increased following DHT treatment.

Figure 3 - Murine awake ECGs after 6 week chronic control or DHT treatment
(**A**) Exemplary lead II ECG recordings in mice. Shown are compound potentials averaged from
20 subsequent cardiac cycles. PR interval (blue) and monophasic part of P wave (orange) are marked.
(**B**) PR interval (n=30-34 mice/group) and P wave duration (n=20-30 mice/group) are prolonged in
Plako[+/-] DHT compared to WT DHT (2-way ANOVA p<0.05 with post-hoc t-test; p-values are indicated
on graphs).

Figure 4 - Atrial unfolded area and conduction

(**A**) Experimental setup for optical mapping of isolated left atria (LA), scale bar 1 mm. DHT exposure has a significant effect on (**B**) LA unfolded area (p<0.05, 2-way ANOVA) measured from optical mapping raw images and area is significantly increased in Plako$^{+/-}$ left atria subjected to DHT (post hoc unpaired t-test, p-values indicated on graph, n=8-10 LA per group). (**C**) Exemplary isochronal activation maps of LA at 100 ms pacing cycle length (averaged, scale bar 1 mm). Both, heterozygous deletion of plakoglobin and DHT exposure have a significant effect on (**D**) LA conduction velocity (p<0.05, 2-way repeated measures ANOVA), but it is only significantly decreased in LA of Plako$^{+/-}$ DHT animals (Bonferroni-adjusted post hoc test, +p$_{adj}$<0.05 vs Plako$^{+/-}$ Ctr; *p$_{adj}$ <0.05 vs WT Ctr, across all cycle lengths, n=8-9 LA per group). Data plotted as mean ± SEM.

Figure 5 - Atrial activation time and beat-to-beat variation
(**A**) Example individual activation maps taken from the final 5 beats of a train of 50 pulses at 80 ms pacing cycle length (CL). DHT exposure has a significant effect on (**B**) beat averaged 95% left atrial activation times. Both genotype and DHT treatment have a significant effect on (**C**) mean beat-to-beat variation in 95% activation times calculated from the final 10 beats of a train of 50 pulses at 80 ms CL (2-way ANOVA with post-hoc analysis as appropriate, p-values indicated). Individual data points denote single left atria (LA). N = WT Control: 6 LA, WT DHT: 7 LA, Plako+/- Control: 8 LA, Plako+/- DHT: 9 LA. Scale bar in (A) indicates 1 mm.

Figure 6 - Atrial gene expression profiles. Confirmation of Jup deficiency and growth response. (**A**) Normalized read counts of plakoglobin gene Jup from RNA sequencing analysis from all groups, confirming ~50% reduction of expression in heterozygous knock-out animals (Plako^{+/-}) in both left (LA) and right atria (RA) (FDR from DESeq2 WT vs Plako^{+/-} <0.05 for both treatment groups). (**B**) Volcano plot illustrating the effect of DHT treatment, comparing gene expression of WT DHT atria to WT Ctr. Significantly differentially expressed genes at a fold change ≥│0.5│ (adjusted p value from DESeq2 < 0.05) are highlighted in red, top 10 hits are labelled. (**C**) Selected genes significantly regulated (adjusted p value from DESeq2 < 0.05) by DHT exposure in at least one of the genotypes. DHT treatment leads to atrial growth, pro-fibrotic and immune response gene expression.
n= WT Ctr: 3 LA + 3 RA; WT DHT: 6 LA + 6 RA; Plako^{+/-} Ctr: 4 LA + 4 RA; Plako^{+/-} DHT: 6 LA + 6 RA. Ccnd2, cyclin D2; Gdf10, growth differentiation factor; Igf1, insulin-like growth factor 1; Igf2r, insulin-like growth factor 2 receptor; Irs1, insulin receptor substrate 1; Mhrt, myosin heavy chain associated RNA transcript; Mtpn, myotrophin; Psap, prosaposin; Acta2, smooth muscle (α)-2 actin; Adamts9, ADAM metallopeptidase with thrombospondin type 1 motif 9; Ccn2, connective tissue growth factor; Col1a1/3a1/4a1/4a2/5a1, collagen type 1 alpha 1/3 alpha 1/4 alpha 1/4 alpha 2/5 alpha 1; Fbn1, fibrillin 1; Fgf1, fibroblast growth factor 1; Has2, hyaluronan synthase 2; Lox, lysyl oxidase; Myocd, myocardin; Ngfr, nerve growth factor receptor; Rock2, rho associated coiled-coil containing protein

kinase 2; C7, complement C7; Cd, cluster of differentiation 6; Cd9, tetraspanin CD9; Ccr5, C-C motif chemokine receptor 5; Csf1r, colony stimulating factor 1 receptor; Dusp4, dual specificity phosphatase 4; Il34, interleukin 34; Tlr3, toll like receptor 3; Tlr4, toll like receptor 4

Figure 7 - Atrial activation time and action potential characteristics

(**A**) Experimental setup for intracellular microelectrode measurements in paced left atria (LA) to record transmembrane action potentials. Representative examples obtained at 100 ms pacing cycle length are shown in (**B**). Both, heterozygous plakoglobin deletion and DHT exposure have a significant effect on mean (**C**) LA activation time, (**D**) action potential amplitude (APA) and (**E**) maximum upstroke velocity (dV dt^{-1} max) (2-way repeated measures ANOVA, $p < 0.05$). WT Ctr (n=21 cells, N=7 LA), WT DHT (n=13 cells, N=5 LA), Plako$^{+/-}$ Ctr (n=20 cells, N=7 LA) and Plako$^{+/-}$ DHT LA (n=23 cells, N=8 LA). Plako$^{+/-}$ DHT LA have longer activation times, decreased APA and reduced dV dt^{-1} max (Bonferroni-adjusted post hoc analysis: #padj<0.05 vs all other groups, across all cycle lengths). Data averaged per atrium before performing statistical analysis and plotted as mean ± SEM.

Figure 8 - Atrial cardiomyocyte sodium current, cell capacitance and sodium channel expression

(A) Representative left atrial (LA) whole cell peak Na$^+$ current (I_{Na}) traces measured at test potentials -100 to +40 mV. (**B**) Mean current-voltage relationships for WT Ctr, WT DHT, Plako$^{+/-}$ Ctr and Plako$^{+/-}$ DHT. Plako$^{+/-}$ DHT LA cells have reduced I_{Na} density (2-way repeated measures ANOVA with Bonferroni post hoc analysis: #padj<0.05 vs all other groups at the respective test potential). Data plotted as mean ± SEM. Both genotype and DHT treatment show a significant effect on (**C**) individual cell capacitance (2-way ANOVA, p<0.05) as measured in WT Ctr (n=14 cells, N=5 LA), WT DHT (n=16

cells, N=5 LA), Plako$^{+/-}$ Ctr (n=11 cells, N=4 LA) and Plako$^{+/-}$ DHT (n=21 cells, N=5 LA). Data points shown individually and plotted as mean ± SEM. Bonferroni-corrected post hoc analysis shows a significant increase in cell capacity by DHT compared to Ctr in both genotypes. Cell capacitance is highest in LA cardiomyocytes from Plako$^{+/-}$ DHT. (**D**) Scatter plots of individual LA cell capacitance against peak I_{Na} density for WT Ctr (n=14 cells, N=5 LA), WT DHT (n=16 cells, N=5 LA), Plako$^{+/-}$ Ctr (n=11 cells, N=4 LA) and Plako$^{+/-}$ DHT (n=21 cells, N=5 LA). There is a significant negative correlation of peak I_{Na} density against cell capacitance for Plako$^{+/-}$ but not WT LA cells after DHT treatment; linear regression analysis. (**E**) Normalized read counts of sodium channel transcript *Scn5a* in LA obtained from RNA sequencing analysis (n=3-6 LA per group). Data points shown individually and box whiskers denote IQR, max-min and median.

Figure 9 - Sodium channel clustering analysis in atrial cardiomyocyte dSTORM images
(**A**) Brightfield images of fixed left atrial (LA) cardiomyocytes prior to dSTORM recording (top lane),
reconstructed super-resolution images of membrane Na$_v$1.5 and zoom-ins of the indicated area
(middle lanes) as well as corresponding cluster maps generated from binary images (bottom lane).
Detections allocated to a cluster appear in (arbitrary) colour, non-clustered detections remain black.
Scale bar 20 µm and 1.5 µm for zoom-in images, respectively. Genotype has a significant effect on (**B**)
number of Na$_v$1.5 detections at the LA cardiomyocyte membrane and (**C**) fraction of Na$_v$1.5 detections

allocated to a cluster (2-way ANOVA, $p<0.05$). Both number and fraction of clustered membrane $Na_v1.5$ is significantly reduced in cardiomyocytes from Plako[+/-] DHT. Results from post-hoc t-tests are reported on the graphs. For (B) and (C): WT Ctr (n=38 cells, N=7 LA), WT DHT (n=35/36 cells, N=11 LA), Plako[+/-] Ctr (n=20 cells, N=5 LA) and Plako[+/-] DHT (n=28 cells, N=6 LA). Data points were normalized to the mean of the respective WT Ctr group on the same imaging session and are shown individually and plotted as mean ± SEM.

Tables

Table 1 - ARVC patient characteristics

Patient characteristics	Non-definite n=97	Definite n=49	p value
Age at last follow-up, mean ± std (years)	42 ± 18	43 ± 16	0.743
Male, n (%)	42 (43)	36 (73)	**0.0008**
Atrial fibrillation/flutter, n (%)	3 (3)	12 (24)	**0.0001**

Table 2 - Phenotypic characteristics of the murine model

	WT control	WT DHT	Plako[+/-] control	Plako[+/-] DHT
Age at terminal experiment (weeks) (n=59-67)	16.5 ± 0.2	16.7 ± 0.2	16.0 ± 0.2	16.6 ± 0.2
Body weight (g) (n=59-65)	27.9 ± 0.4	29.4 ± 0.4**	27.9 ± 0.4	29.2 ± 0.4[+]
Serum DHT (nM) (n=11-18)	0.25 ± 0.06	0.85 ± 0.21*	0.31 ± 0.11	1.22 ± 0.29[+]
Left atrial weight:tibia length (mg mm^{-1}) (n=29-32)	0.23 ± 0.01	0.24 ± 0.01	0.22 ± 0.01	0.28 ± 0.02[+]
Right atrial weight:tibia length (mg mm^{-1}) (n=28-32)	0.25 ± 0.01	0.29 ± 0.01*	0.26 ± 0.02	0.30 ± 0.03
Seminal vesicle weight:tibia length (mg mm^{-1}) (n=39-45)	6.0 ± 0.4	6.0 ± 0.3	5.5 ± 0.2	6.5 ± 0.3[+]

Murine echocardiography (light anaesthesia, mean heart rate 390-440 bpm)

	WT control	WT DHT	Plako[+/-] control	Plako[+/-] DHT
Heart rate (bpm) (n=18-28)	411 ± 3	415 ± 3	413 ± 3	414 ± 3
Left atrial diameter:tibia length (mm mm^{-1}) (n=15-23)	0.130 ± 0.005	0.140 ± 0.007	0.138 ± 0.006	0.144 ± 0.007
Left ventricular mass:tibia length (mg mm^{-1}) (n=18-28)	5.4 ± 0.3	5.5 ± 0.3	4.9 ± 0.2	6.0 ± 0.3[+]
Left ventricular ejection fraction (%) (n=18-28)	43 ± 3	50 ± 3	47 ± 2	45 ± 3

Data presented as mean ± SEM, n=number of animals per group, *p<0.05 vs WT control, **p<0.01 vs WT control (post-hoc t-test, 2-way ANOVA p<0.05 for DHT treatment), [+]p<0.05 vs Plako[+/-] control (post-hoc t-test, 2-way ANOVA p<0.05 for DHT treatment)

Table 3 - Prevalence of Atrial fibrillation/flutter (Atrial Arrhythmia, AA) in ARVC patients

Study [First author]	Year	Number of patients	Mean age [years, ± SD / (range)]	AA prevalence [%]	Patients with AA; male [%]	Patients without AA; male [%]
Tonet (3)	1991	72	38 (16-73)	24	n.r.	n.r.
Jaoude (95)	1996	74	37.2 ± 13.5 (11-68)	4	n.r.	n.r.
Brembilla-Perot (96)	1998	47	44 ± 18 (17-72)	17	63	n.r.
Peters (97)	2004	80	45.9 (22-91)	30	n.r.	n.r.
Chu (5)	2010	36	47 (17-80)	42	80	76
Camm (4)	2013	248	41.6 ± 14	14	69	50
Saguner (98)	2014	90	49.7 ± 14.6	20	61	64
Wu (99)	2016	294	37.7 ± 14.8	13	62	77
Bourfiss (100)	2016	66	46.4 ± 15.8	21	79	50
Mazzanti (101)	2016	301	38 ± 18	4	n.r.	n.r.
Gilljam (102)	2018	183	Median 46 (14-65)	9	n.r.	n.r.
Wu (103)	2018	100	37.1 ± 12.1	9	n.r.	n.r.
Müssigbrodt (104)	2018	70	53.2 ± 14.0	37	62	73
Cardona-Guarache (105)	2019	117	52 ± 14	22	69	57
Kikuchi (106)	2020	90	44 ± 15	36	78	76
Baturova (34)	2021	100	41 (30-55)	28	n.r.	n.r.
		∑ 1915		Weighted mean: 15%		
Present study: **ARVC diagnosis**						
Non-definite		97	42 ± 18	3	67	43
Definite		49	43 ± 16	24	67	73

SD = standard deviation, n.r. = not reported