Title: Systemic Calcitonin Gene-Related Peptide (CGRP) modifies auditory and vestibular endorgan

electrical potentials, and increases auditory and vestibular hypersensitivities: Implications for vestibular

migraine

Abbreviated Title: CGRP's effect on endorgan and behavior

Authors: Shafaqat M. Rahman[1], Benjamin Liang[1], Stefanie Faucher[1], Elana Fine[1], Raajan Jonnala[1],

Joseph C. Holt[2,3], Choongheon Lee[3], Anne E. Luebke[1,2*]

Affiliations:

[1]University of Rochester, Department of Biomedical Engineering, Rochester, NY 14627

[2]University of Rochester Medical Center, Department of Neuroscience, Del Monte Institute of

Neuroscience, Rochester, NY 14642

[3]University of Rochester Medical Center, Department of Otolaryngology, Rochester, NY 14642

*Corresponding author: Anne E. Luebke, aluebke@ur.rochester.edu

Number of Pages: 21

Number of Figures: 3

Number of Tables: 0

Number of words for Abstract: 225

Number of words for Introduction: 343

Number of words for Discussion: 582

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: This work was fully supported by NIH R01DC017261.

Abstract:

Migraine is a severe and chronic neurological disorder that affects ~18% of people worldwide, the

majority female (3:1). It is characterized by recurrent, debilitating headaches and heightened sensory

sensitivities to stimuli, and migraineurs may also experience a heightened motion sensitivity that leads

to episodes of vertigo and imbalance, termed vestibular migraine (VM). Calcitonin Gene-Related

Peptide (CGRP) is implicated in migraine and is believed to act on brain meninges, midbrain, or

brainstem, and CGRP-based therapeutics have shown efficacy for migraine treatment. However,

CGRP also signals at efferent synapses of the cochlea and vestibular endorgans, and it is unclear if

exogenous CGRP can modulate inner ear function and cause heightened behavioral responses

consistent with VM. In C57BL6/J mice, we tested if intraperitoneally (IP) delivered CGRP can modulate

endorgan potentials to sound [via auditory brainstem responses (ABRs)]and jerk stimulus [via vestibular

sensory evoked potentials (VsEPs)]. We also assessed behavioral measures of phonophobia [acoustic

startle reflex (ASR)] and imbalance [postural sway center of pressure (CoP)] after IP CGRP, and

observed female mice exhibit heightened sensitivities to IP CGRP in all assays, while male mice

showed similar auditory sensitivity, and endorgan effects, but no postural sway changes. In conclusion,

we show that systemically delivered CGRP modulates the nerve activity of the auditory and vestibular

endorgans, and elicits behavioral effects that are consistant with symptoms in a disorder like VM.

Significance statement:

Calcitonin Gene-Related Peptide (CGRP) has been implicated in migraine, and CGRP-based

therapeutics have shown efficacy in the treatment of migraine headaches. Yet, CGRP is also present

in efferent synapses of the inner ear, and it is not clear if increases in systemic CGRP can act directly

on inner ear endorgans, and modify corresponding behavioral responses. In this study, we determined

systemic CGRP changes both cochlear and vestibular endorgan potentials, and produced behaviors

suggestive of phonophobia, and increased postural sway. Peripheral changes in auditory and

vestibular compound action potentials following systemic CGRP, suggest systemic CGRP can act

directly on the inner ear, which opens new therapeutic targets to treat migraine with vertigo.

Introduction:

Migraine is a debilitating chronic neurological disorder that is listed by the World Health Organization (WHO) as the most debilitating neurological disorder worldwide and is estimated to affect 18% of women and 6% of men (Goadsby, Lipton, and Ferrari 2002; Lipton et al. 2007). About 42% of migraineurs have a vestibular component causing balance problems and dizziness (Kuritzky, Toglia, and Thomas 1981; Neuhauser et al. 2001; Neuhauser 2007, 2009), termed vestibular migraine (VM). VM is a major cause of vertigo in dizziness clinics (Kayan and Hood 1984), and patients with migraine, and especially VM, exhibit a heightened sense of sound (Ashkenazi et al. 2009; Lopez-Escamez et al. 2014), yet despite migraine and VM's prevalence, the pathophysiology remains uncertain (Lempert et al. 2022; Lempert et al. 2012).

Calcitonin gene-related peptide (CGRP) is hypothesized to play a role in sensory hypersensitivity in migraine (Russo 2015). In patients, CGRP levels are elevated during migraine, and normalization of CGRP levels is associated with the relief of headache pain which is thought to act at either brain meninges, or the trigeminal system (Goadsby, Lipton, and Ferrari 2002; Juhasz et al. 2005; Lassen et al. 2002). However, CGRP also signals at efferent synapses of the cochlea and vestibular endorgans, so we wondered if the elevated CGRP levels of migraine could affect the inner ear directly. To that end, we modeled the increase in CGRP during migraine, by a systemic intraperitoneal (IP) injection, to determine if systemic CGRP can modulate activity of these endorgans (Dickerson et al. 2016; Luebke et al. 2014; Maison et al. 2003). We were also interested in knowing if increases in systemic CGRP can cause behaviors suggestive of heightened auditory and vestibular sway consistent with VM.

In this study, we tested wildtype C57BL6/J mice to determine if there is a change in endorgan potentials to sound (measured using auditory brainstem responses –ABRs) and jerk stimulus [measured using vestibular stimulus evoked potentials (VsEPs)] after a CGRP injection. We also assessed corresponding behavioral measures of phonophobia (acoustic startle reflex – ASR) and imbalance (postural sway) after systemic CGRP injection.

Methods:

Animals: C57BL6/J mice were obtained from Jackson Laboratories (JAX #664) and were housed under a 12-hour day/night cycle under the care of the University Committee on Animal Resources (UCAR) at the University of Rochester. Mice are housed together in cages (max 5 mice per cage) with *ad libitum* access to food, water, and bedding material. Equal numbers of male and female mice were tested and a total of 82 mice (41 M/ 41F) were used for these studies, which were powered sufficiently to detect male/female differences. Before experimentation on each testing day, mice were equilibrated in the testing room controlled for an ambient temperature between 22-23^{o}C for at least 30 minutes, and remained in this room until completion of experiments. Injections occurred after the equilibration period. Mice used during behavioral experiments were returned to the vivarium, while mice used for ABR/VsEP experiments were euthanized after testing. Mice tested were between 2.3 - 6 months of age. This age range in mice correlates to 20-30 years in humans, which is within the age range that patients with migraine develop symptoms (Lessem 2018). All animal procedures were approved by the University of Rochester's IACUC committee and performed in accordance with the standards set by the NIH.

VsEP and ABR Protocol: C57BL/6 mice were anesthetized with IP urethane/xylazine (1.2 g/kg/20 mg/kg). VsEP and ABR recording methods have been described in detail elsewhere (Lee, Holt, and Jones 2017). Briefly, to record VsEP and ABR, subcutaneous electrodes were placed on the midline over the nuchal crest (G1), behind the right pinna (G2), and on the hip (ground). A noninvasive head clip was used to secure the mouse's head to an electromechanical shaker. During VsEPs, the shaker delivered a transient (jerk) head translation in the naso-occipital axis (± X). To confirm the absence of auditory responses during VsEP testing, a binaural forward masker (intensity: 92 dB SPL; bandwidth: 50~50 kHz) was presented with the use of a free-field speaker (FF1, TDT). For ABR recordings, the same electrodes were used to record ABRs, which were evoked by a single rectangular voltage pulse of 0.03 ms duration (corresponding to 33 kHz cut-off frequency). These click stimuli were presented via a free-field speaker, which was the same driver for acoustic masking during the VsEP recording. A noninvasive rectal thermometer or thermistor was used to monitor the animals' core temperature at 38.0 ± 0.2°C. The standard signal averaging technique described in previous studies (Lee, Holt, and

Jones 2017) was used to record both VsEP and ABR, as shown in Fig. 1A. Recordings of both VsEP

and ABR were made in a soundproof booth. Both VsEPs at maximal intensity +6 dB re: 1 g/ms and

ABRs at maximal intensity 100 dB SPL, were recorded as a stable baseline before a single IP injection

of either vehicle control (PBS) or 0.1 mg/kg CGRP was given.   Control amplitude and latencies are

depicted as the averages of the 2-3 measurements taken during baseline before injection. Following

single IP injection of either vehicle or CGRP, VsEP and ABR recordings were continuously measured

where significant shifts in VsEP and ABR responses were first observed in 10 mins and reaching a

plateau within 15 mins of drug injection (post-CGRP; data not shown). Thus, amplitudes and latencies

depicted as post-saline/post-CGRP are averages of the 2-3 measurements taken after 20-30 min post-

injections.

Drug administration: Injections were performed intraperitoneally (IP) with a fine gauge (30ga) insulin

syringe. Dulbecco PBS served as the vehicle control. IP CGRP was prepared at 0.1 mg/kg (rat α-

CGRP, Sigma). Injection volumes were calibrated so that each animal received 1 injection per testing

day at ~100 ul. For behavioral experiments, mice were tested ~20 minutes after IP delivery of either

vehicle or CGRP.

Behavioral test of phonophobia: Acoustic startle reflex (ASR)

To assess behavioral phonophobia or heightened sound sensitivity, we made use of the acoustic startle

response (ASR).   Experiments were conducted within a sound-attenuating room (IAC, Bronx, NY) with

Sonex foam lined walls. Mice were placed individually in a wire-mesh cage, 5 cm wide, 7 cm long, and

4 cm high, having free sound penetration. The testing cage was oriented so that the mouse's head

faced a Tucker Davis ES2 electrostatic speaker (Masker Speaker) located 46 cm directly in front of the

mouse's head (60dB BBN). The startle speaker was a Radio Shack tweeter that was suspended 15 cm

above the mouse (80-130 dB). The acoustic startle speaker and its supports, the pedestal, acrylic shelf,

and the table on which the apparatus was placed were all covered with echo absorbing foam and

carpeting. Force of the acoustic startle response (ASR) was transduced by an accelerometer and

amplitude scored as integrated root mean square voltage for 100 ms after delivery of the ASR.

Experimental stimuli were controlled, and responses were recorded using custom MATLAB

(MathWorks) software. We tested acoustic startle series in quiet and in noise. There were 12 conditions

in this session with startle stimuli ranging from 80 to 130 dB SPL, delivered in silence or in a continuous

background noise. Background noise, when present, was on for the duration of the inter-trial interval

and broadcast from the masker speaker, as shown schematically in Fig. 2.

<u>Behavioral test of postural sway</u>

Postural sway was measured as a correlate for static balance. Mice were weighed and placed on a

force plate designed to measure forces in the X, Y, Z and its moments in the XY, YZ, and XZ directions.

We used the AMTI Biomechanics Force platform (model HEX6x6) and AMTI automated acquisition

software to obtain forces and moments and then MATLAB code was used to calculate Center of

Pressure (CoP) measures based on a generating a 95% confidence ellipse enclosing 95% of the CoP

trajectory values recorded in a single trial. During a given test, a mouse was first allowed to freely

maneuver on the force plate for 2-5 minutes to acclimate to the novel environment. An accessory

plexiglass guard was placed around the force plate to prevent animals from moving off the test plate.

After acclimation, ~ 10 baseline measurements were taken of the mouse with each measurement

indicating a CoP area (resolution per CoP measurement: 300 samples per second). These

measurements were taken when the mouse showed no movement and its four paws were touching the

force platform. To assess the effect of a vestibular challenge (VC), the mouse was then subjected to a

constant orbital rotation, 125 rpm for 5 minutes, and was placed back onto the force platform. After a

five-minute recovery period, an additional 10 measurements were recorded of the mouse's sway to

determine effect of the vestibular challenge (post VC measures), as schematized in **Fig. 3**. Mice were

first tested after systemic delivery of IP vehicle (20 min after injection), with sway measured both before

and after vestibular challenge. Four days after vehicle control testing, the same mice were injected with

IP CGRP and were tested again 20 minutes after injection (timing correlated with endorgan potential

changes).   While systemic CGRP's effect is believed to dissipate two to three hours post–injection, we

retested after four days to eliminate any residual dizziness due to VC or lingering effects of CGRP. In

some animals we staggered sequence of IP injections, injecting CGRP followed by vehicle testing. We

also injected vehicle 2x to assess test/retest reliability and saw no differences (data not shown). In

certain testing conditions, mice exhibited unreasonably large CoP values (> 15 cm^2) that suggested that

they were either moving during the time of measurement or that they were at the edge of the force

plate, and those values were excluded based on a 10% ROUT analysis (GraphPad Prism 9.0).   Yet,

even with values excluded, a minimum of eight different 95% ellipses were computed per mouse, per

condition tested.

<u>Statistical Analyses</u>

All statistical analyses were conducted in GraphPad Prism 9.0. Repeated measure ANOVA (RM-

ANOVA) and paired t-tests were the primary statistical tools. Bonferroni and Tukey multiple

comparisons test were used to assess for differences between treatment groups. Significance was

set at $p < 0.05$ for all analyses.

<u>Results:</u>

<u>IP CGRP's Effects on ABR:</u>

Representative traces are shown for ABR recordings in female and male C57BL6/J mice **(Fig. 1B-1C)**.

A 2-way RM ANOVA assessing the factors - IP CGRP and sex - showed IP CGRP's effects to be

significant on wave 1 ABR amplitudes (*F (1, 12) = 11.81; p = 0.005*). Female control ABR amplitudes (n

= 6) were computed to be 3.60 ± 0.22 µV but were increased to 4.28 ± 0.25 µV after IP CGRP

(*Bonferroni, p = 0.007*). Likewise, male control amplitudes (n = 7) were computed to be 4.31 ± 0.5 µV

and increased to 4.66 ± 0.44 µV after IP CGRP, but were not deemed statistically significant **(Fig. 1F)**.

Wave 1 ABR amplitudes were normalized so that post-CGRP measurements highlight the change in

amplitude from their respective control measurements (Δ wave1 amplitude*),* and both sexes were

additionally pooled in this depiction **(Fig. 1G)**. When both sexes are pooled (n = 13), a statistically

significant Δ increase of 0.50 ± 0.13 µV was observed in wave 1 amplitudes due to IP CGRP (*paired t-*

test; t, df = 3.9,12; p = 0.002). Wave 2-1 ABR latencies were calculated as the difference between P2

and P1 latencies for control and post-CGRP; however, no significant differences were observed in

latency differences **(Fig. 1H).** A separate group of mice was tested for the effects of PBS as the vehicle

(n = 4), but no significant differences were seen in Δ wave1 amplitude or latency differences between

control and post-vehicle ABR (Δ wave1 amplitude $_{vehicle – control}$ = -0.19 ± 0.21 μV).

<u>IP CGRP's Effects on VsEPs:</u>

Similar to ABR tests, representative traces are shown for VsEP recordings in females and males (**Fig.**

1D-1E). A 2-way RM ANOVA (factors: CGRP, sex) showed CGRP's effects to be significant on P1-N1

VsEP amplitudes (*F (1, 11) = 16.9; p = 0.002*). Female P1-N1 VsEP control amplitudes (n = 6) were

computed to be 0.68 ± 0.06 μV but were reduced to 0.55 ± 0.06 μV after IP CGRP, however, this

reduction was not deemed statistically significant **(Fig. 1I)**. Likewise, male P1-N1 VsEP control

amplitudes (n = 8) were computed to be 0.95 ± 0.10 μV but were reduced significantly to 0.76 ± 0.10 μV

after IP CGRP (*Bonferroni, p = 0.02*). P1-N1 amplitudes were normalized so that post-CGRP

measurements highlight the change in amplitude from their respective control (Δ P1-N1 amplitude*), and*

both sexes were additionally pooled (**Fig. 1J**). When pooling all animals (n = 14), a significant Δ

reduction of -0.16 ± 0.04 uV in P1-N1 amplitude is observed due to IP CGRP (*paired t-test; t, df = 3.63,*

13; p = 0.003).

P1, N1, and P2 latencies were also assessed for VsEPs, and these latencies are depicted as mean ±

SEM for females and males during control and post-CGRP tests **(Fig. 1K).** While no significant

differences were observed due to IP CGRP, modest delays in P1 were observed due to IP CGRP in

either sex. IP CGRP increased female P1 by 26.6 microseconds and male P1 by 22.5 microseconds

compared to P1 control latencies. IP CGRP also increased female N1 by 30 microseconds but had no

effect on Male N1. However, these latency changes were not statistically significant. Comparatively in

mice treated with IP vehicle, no significant P1-N1 amplitude or latency differences were observed

between control and post-vehicle VsEP (Δ P1-N1 amplitude $_{vehicle – control}$ = -0.03 ± 0.08 μV).

<u>CGRP's effect on sound sensitivity</u>

Our sound sensitivity measure in the mouse model makes use of the acoustic startle response (ASR) in

quiet and in background noise. Mice were assessed for the effects of IP CGRP as compared to IP

vehicle on their baseline. The ASR experimental setup is illustrated in **Fig. 2 A** (quiet) and **Fig. 2D**

(background noise 60 dB). In both IP vehicle and IP CGRP injected animals, an increase in ASR was

detected with increasing loudness of the startle stimulus (ES). All animals tested exhibited robust

startle responses.   When ASR was tested after IP CGRP injection and compared to IP vehicle, all

animals exhibited reduced ASR thresholds in both quiet and noise. IP CGRP significantly increased

sound sensitivity to the startle stimulus in quiet (**Fig. 2 B, C**) and in background noise (**Fig. 2 E, F**) in

both female and male mice at all sound levels ($p < 0.0001$).

<u>CGRP's and VC's effects on Postural Sway</u>

Mice were assessed for postural sway by determining center of pressure (CoP) area. An experimental

timeline is shown in **Fig. 3A**. Twenty minutes after delivery of IP vehicle and without a vestibular

challenge (VC), group averages of 95% ellipse areas were computed to be 0.5 ± 0.06 cm^2 for females

and 0.4 ± 0.04 cm^2 for males **(Fig. 3B and C).** For comparison, a deadweight (toy mouse) was also

measured on the force platform, and its average 95% ellipse area was computed to be 0.02 ± 0.0005

cm^2 **(Fig. 3D).** A different cohort of mice (10M/ 10F) was assessed for a test-retest of 95% ellipse areas

after IP vehicle delivery, and testing occurred four days apart. No significant differences were seen in

either sex during the test-retest (mean $\pm$ SE $_{\text{test-retest difference}}$ for females = 0.10 ± 0.3 cm^2 *(Bonferroni, p >*

0.99) and males = $-0.14 + 0.3$ cm^2 *(Bonferroni, p > 0.99)*). A 3-way ANOVA was used to assess the

factors of sex, CGRP, and VC on 95% ellipse areas and is graphically depicted via a violin plot **(Fig.**

3E). Significant differences were seen across the following factors: sex (*$F_{(1.00, 22.00)}$ = 25.26, p <*

0.0001) and CGRP (*$F_{(1.00, 22.00)}$ = 4.67, p = 0.04*). The effects of VC were trending (*$F_{(1.00, 22.00)}$*

= 2.93, p = 0.1). Tukey post-hoc analysis was used to observe differences in specific comparisons

across the three factors. An increase was seen in 95% ellipse due to the VC in female mice treated with

IP vehicle *(p = 0.053)*, as shown in the before/after plot (**Fig. 3F**). However, before the VC (pre-VC), in

females but not males, IP CGRP induced significant increases in CoP ellipse areas by 0.94 ± 0.23 cm^2

(*p = 0.01*) as seen in the before-after plot **(Fig. 3G).** When looking at sex differences, they were not

observed after delivery of IP vehicle. However, during the pre-VC tests after IP CGRP, females had

higher 95% ellipse areas compared to males by 0.96 ± 0.19 cm^2 (*p = 0.007*).

The major and minor axes of 95% ellipse areas were also analyzed with 3-way ANOVA and

post-hoc analysis through Tukey's multiple comparisons test (**Fig. 3H and I**). The major and minor axes

lengths serve as correlates for anteroposterior (AP) and mediolateral (ML) sway magnitudes. In pre-VC

female values, IP CGRP increased AP ellipse lengths by 0.49 + 0.10 cm compared to their IP vehicle

tests (*p = 0.005*). Also, in IP vehicle-treated females, there was an increase in AP ellipse lengths due to

the VC, but this was trending (*t (p = 0.056)*), and IP CGRP-treated females exhibited greater AP

lengths compared to males during pre-VC testing by 0.57 + 0.10 cm, (*p = 0.0002*). When observing ML

sway, like AP sway, IP vehicle-treated females showed increased ML lengths due to VC, but this was

trending (*t (p = 0.07)*). Additionally, IP-CGRP-treated females exhibited increased ML ellipse lengths

compared to their IP vehicle response by 0.12 + 0.03 cm, (*p = 0.02*). No other significant changes were

observed in AP and ML sway.

Discussion:

Using electrophysiological techniques (ABR/VsEP), we observed that systemic, intraperitoneally

delivered CGRP modulates endorgan function in the inner ear with a maximum ~20- 30 minutes after

injection. Exogenous CGRP increases wave 1 amplitudes of the ABR but reduces VsEP P1-N1

amplitudes, suggesting that at least one action of systemic CGRP is at the efferent signaling sites of the

auditory and vestibular periphery. We did not further characterize wave 2 and later peaks of

amplitudes or latencies, as we were primarily interested in wave 1/P1-N1 as an indicator of sensory

processing changes at the periphery.

Furthermore, we observed that systemic CGRP - during this 20–30-minute time - increased

sound sensitivity in both male and female wildtype C57BL6/J mice, whereas it only increased postural

sway in female mice.

A few studies in animal models have provided additional evidence for CGRP's role in vestibular

dysfunction and sound sensitivity. Consistent with current VsEP results where we observed delays in

VsEP latencies due to systemic CGRP (though not statistically significant), the loss of αCGRP In

transgenic mouse lines significantly enhanced activation timing of the macular sensors, through

changes at the endorgan level (Jones et al. 2018). Intracerebral-ventricular (ICV) injection of

nitroglycerin (NTG) caused impaired balance beam walking in rats, and immunohistochemistry

detected more CGRP expression in the vestibular nucleus of the NTG rat model than was observed in

control rats. The administration of a CGRP-receptor antagonist (olcegepant-BIBN4096BS) was shown

to be effective in alleviating impaired balance beam walking (Zhang et al. 2020; Tian et al. 2022).   In

addition, a recent study suggested CGRP has an excitatory role in the auditory nerve response, as

intra-cochlear CGRP infusion was shown to enhance spontaneous and sound-driven auditory nerve

activity in guinea pigs, in comparison to CGRP-null mice where a loss of nerve activity excitation was

observed (Le Prell et al. 2021; Allen and Luebke 2017). Thus, the auditory nerve may be hyper-

sensitized to CGRP's effects during a migraine.

Peripheral effects of systemic CGRP administration on VsEP responses is somewhat surprising,

as the post-CGRP injection VsEP potentials were reduced and it is unclear why. However, this

observation is relevant to VM, which is characterized by sensory misintegration to stimuli, and thus

reduced vestibular sensitivity that occurs concurrently with a heightened auditory sensitivity may

provide a basis for the occurrence of these symptoms.

Additionally, it is unclear why we did not find significant changes in males during postural sway

testing. While more males exhibited significant changes in their VsEP measures, we presume the lack

of CGRP-induced behavioral effects on sway may be due to other postural compensatory mechanisms

present in male mice. Yet, we did discover a robust sex difference when assessing mouse postural

sway, as female mice were much more affected than their male counterparts to injected CGRP.

Similar female-specific sexual dimorphism has been observed in other animal assays assessing

CGRP's effects in migraine but we are the first to show such differences in preclinical migraine

correlates of postural sway (Recober et al. 2009; Paige et al. 2022).

Small molecule blockers of the CGRP receptor 'gepants' have been shown to enter the

periphery, brainstem, and midbrain areas. Future studies will determine if blocking CGRP receptor

signaling by small molecules ('gepants") or by larger monoclonal blocking antibodies [that may or may

not cross blood brain barrier (BBB), but have been shown to be permeable to various peripheral

ganglia] can reduce these sound and imbalance sensitivities present after systemic CGRP injection

(Noseda et al. 2020; Tian et al. 2022; Zhang et al. 2020).

References:
303

Allen, Paul D., and Anne E. Luebke. 2017. 'Reflex Modification Audiometry Reveals Dual Roles for

Olivocochlear Neurotransmission', *Frontiers in Cellular Neuroscience*, 11.

Ashkenazi, A., A. Mushtaq, I. Yang, and M. L. Oshinsky. 2009. 'Ictal and interictal phonophobia in

migraine-a quantitative controlled study', *Cephalalgia*, 29: 1042-8.

Dickerson, I. M., R. Bussey-Gaborski, J. C. Holt, P. M. Jordan, and A. E. Luebke. 2016. 'Maturation of

suprathreshold auditory nerve activity involves cochlear CGRP-receptor complex formation',

Physiol Rep, 4.

Goadsby, P. J., R. B. Lipton, and M. D. Ferrari. 2002. 'Migraine--current understanding and treatment',

N Engl J Med, 346: 257-70.

Jones, S. M., S. Vijayakumar, S. A. Dow, J. C. Holt, P. M. Jordan, and A. E. Luebke. 2018. 'Loss of

alpha-Calcitonin Gene-Related Peptide (alphaCGRP) Reduces Otolith Activation Timing

Dynamics and Impairs Balance', *Front Mol Neurosci*, 11: 289.

Juhasz, G., T. Zsombok, B. Jakab, J. Nemeth, J. Szolcsanyi, and G. Bagdy. 2005. 'Sumatriptan causes

parallel decrease in plasma calcitonin gene-related peptide (CGRP) concentration and migraine

headache during nitroglycerin induced migraine attack', *Cephalalgia*, 25: 179-83.

Kayan, A., and J. D. Hood. 1984. 'Neuro-otological manifestations of migraine', *Brain*, 107 (Pt 4): 1123-

42.

Kuritzky, A., U. J. Toglia, and D. Thomas. 1981. 'Vestibular function in migraine', *Headache*, 21: 110-2.

Lassen, L. H., P. A. Haderslev, V. B. Jacobsen, H. K. Iversen, B. Sperling, and J. Olesen. 2002. 'CGRP

may play a causative role in migraine', *Cephalalgia*, 22: 54-61.

Le Prell, Colleen G., Larry F. Hughes, David F. Dolan, and Sanford C. Bledsoe. 2021. 'Effects of

Calcitonin-Gene-Related-Peptide on Auditory Nerve Activity', *Frontiers in Cell and*

Developmental Biology, 9.

Lee, C., J. C. Holt, and T. A. Jones. 2017. 'Effect of M-current modulation on mammalian vestibular

responses to transient head motion', *J Neurophysiol*, 118: 2991-3006.

Lempert, T., J. Olesen, J. Furman, J. Waterston, B. Seemungal, J. Carey, A. Bisdorff, M. Versino, S.

Evers, A. Kheradmand, and D. Newman-Toker. 2022. 'Vestibular migraine: Diagnostic criteria1',

J Vestib Res, 32: 1-6.

Lempert, T., J. Olesen, J. Furman, J. Waterston, B. Seemungal, J. Carey, A. Bisdorff, M. Versino, S.

Evers, and D. Newman-Toker. 2012. 'Vestibular migraine: diagnostic criteria', *J Vestib Res*, 22:

167-72.

Lessem, Sarah E. 2018. "QuickStats: Percentage of Adults Who Had a Severe Headache or Migraine

in the Past 3 months, by Sex and Age Grou - National Health Interview Survey, United States,

2018." In *Morbidity and Mortality Weekly Report (MMWR)*, edited by National Center for Health

Statistics.

Lipton, R. B., M. E. Bigal, M. Diamond, F. Freitag, M. L. Reed, W. F. Stewart, and Ampp Advisory

Group. 2007. 'Migraine prevalence, disease burden, and the need for preventive therapy',

Neurology, 68: 343-9.

Lopez-Escamez, J. A., J. Dlugaiczyk, J. Jacobs, T. Lempert, R. Teggi, M. von Brevern, and A. Bisdorff.

2014. 'Accompanying Symptoms Overlap during Attacks in Meniere's Disease and Vestibular

Migraine', *Front Neurol*, 5: 265.

Luebke, A. E., J. C. Holt, P. M. Jordan, Y. S. Wong, J. S. Caldwell, and K. E. Cullen. 2014. 'Loss of

alpha-calcitonin gene-related peptide (alphaCGRP) reduces the efficacy of the Vestibulo-ocular

Reflex (VOR)', *J Neurosci*, 34: 10453-8.

Maison, S. F., R. B. Emeson, J. C. Adams, A. E. Luebke, and M. C. Liberman. 2003. 'Loss of alpha

CGRP reduces sound-evoked activity in the cochlear nerve', *J Neurophysiol*, 90: 2941-9.

Neuhauser, H. K. 2007. 'Epidemiology of vertigo', *Curr Opin Neurol*, 20: 40-6.

———. 2009. '[Epidemiology of dizziness and vertigo]', *Nervenarzt*, 80: 887-94.

Neuhauser, H., M. Leopold, M. von Brevern, G. Arnold, and T. Lempert. 2001. 'The interrelations of

migraine, vertigo, and migrainous vertigo', *Neurology*, 56: 436-41.

Noseda, R., A. J. Schain, A. Melo-Carrillo, J. Tien, J. Stratton, F. Mai, A. M. Strassman, and R.

Burstein. 2020. 'Fluorescently-labeled fremanezumab is distributed to sensory and autonomic

ganglia and the dura but not to the brain of rats with uncompromised blood brain barrier',

Cephalalgia, 40: 229-40.

Paige, C., I. Plasencia-Fernandez, M. Kume, M. Papalampropoulou-Tsiridou, L. E. Lorenzo, E. T.

David, L. He, G. L. Mejia, C. Driskill, F. Ferrini, A. L. Feldhaus, L. F. Garcia-Martinez, A. N.

Akopian, Y. De Koninck, G. Dussor, and T. J. Price. 2022. 'A Female-Specific Role for

Calcitonin Gene-Related Peptide (CGRP) in Rodent Pain Models', *J Neurosci*, 42: 1930-44.

Recober, A., A. Kuburas, Z. Zhang, J. A. Wemmie, M. G. Anderson, and A. F. Russo. 2009. 'Role of

calcitonin gene-related peptide in light-aversive behavior: implications for migraine', *J Neurosci*,

29: 8798-804.

Russo, A. F. 2015. 'Calcitonin gene-related peptide (CGRP): a new target for migraine', *Annu Rev*

Pharmacol Toxicol, 55: 533-52.

Tian, R., Y. Zhang, Q. Pan, Y. Wang, Q. Wen, X. Fan, G. Qin, D. Zhang, L. Chen, Y. Zhang, and J.

Zhou. 2022. 'Calcitonin gene-related peptide receptor antagonist BIBN4096BS regulates

synaptic transmission in the vestibular nucleus and improves vestibular function via

PKC/ERK/CREB pathway in an experimental chronic migraine rat model', *J Headache Pain*, 23:

35.

Zhang, Y., Y. Zhang, K. Tian, Y. Wang, X. Fan, Q. Pan, G. Qin, D. Zhang, L. Chen, and J. Zhou. 2020.

'Calcitonin gene-related peptide facilitates sensitization of the vestibular nucleus in a rat model

of chronic migraine', *J Headache Pain*, 21: 72.

Fig. 1:

385   Fig. 2:
386
387

A

Vestibular Challenge (VC)
orbital rotation (5 min)

M_x F_x
M_y M_z
F_y
F_z

$COP_x = M_y/F_z$
$COP_y = -M_x/F_z$

Inject saline (day 1)
or CGRP (day 5)

-20 min   test 10x

1 cm
125 rpm

Post- VC
(5 min)

test 10x

B

COP average = 0.5 cm^2

Center of Pressure$_y$ (COP$_y$- cm)

M-L
A-P

Test toy mouse

Center of Pressure$_X$ (COP$_X$- cm)

C

COP average = 0.4 cm^2

Center of Pressure$_y$ (COP$_y$- cm)

M-L
A-P

Test toy mouse

Center of Pressure$_X$ (COP$_X$- cm)

D

Test toy mouse
95% confidence ellipse
(COP$_y$- cm)
(COP$_X$- cm)

E

CGRP, Post-VC (M)
CGRP, Pre-VC (M)
Vehicle, Post-VC (M)
Vehicle, Pre-VC (M)
CGRP, Post-VC (F)
CGRP, Pre-VC (F)
Vehicle, Post-VC (F)
Vehicle, Pre-VC (F)

ns
ns
ns
ns
*
t (p = 0.053)

Center of Pressure (COP - cm^2)

F

Center of Pressure (CoP - cm^2)

t (p = 0.053)

Vehicle, Pre-VC (F)
Vehicle, Post-VC (F)

G

*

Vehicle, Pre-VC (F)
CGRP, Pre-VC (F)

H

CGRP, Post-VC (M)
CGRP, Pre-VC (M)
Vehicle, Post-VC (M)
Vehicle, Pre-VC (M)
CGRP, Post-VC (F)
CGRP, Pre-VC (F)
Vehicle, Post-VC (F)
Vehicle, Pre-VC (F)

ns
**
t (p = 0.056)

Major Axis- AP (cm)

I

ns
*
t (p = 0.07)

Minor Axis - ML (cm)

 Figure Legends:

Fig. 1: A. Schematic illustrating mouse setup for VsEP/ABR recordings captured during baseline

(control) and 20-30 minutes after intraperitoneal (IP) injections of either PBS vehicle or 0.1 mg/kg

CGRP in male (M) and female (F) C57BL6/J mice. **B-C**. Representative click-evoked ABR waveforms -

in duplicate -were collected for ABR at a maximal intensity of 100 dB SPL, and waves were labeled

(wave1 – wave 4). **D-E.** Likewise, representative VsEP waveforms were collected at +6 dB re: 1 g/ms

having three, labeled response peaks (p1, n1, and p2). Orange and blue colors are used to distinguish

female and male tracings post-CGRP, respectively. Whisker (**F, I**) and before-after (**G, H, J, K**) plots

are used to show changes in amplitude and latencies for ABR and VsEP recordings, and the following

symbols and colors are used: female control (open, orange circle), female post-CGRP (closed, orange

circle), male control (open, blue square), male post-CGRP (closed, blue square). **F-G**. A statistically

significant increase is seen in female wave 1 ABR amplitude post-CGRP (*p = 0.007*), and also when

sexes are pooled (*p = 0.002*). **I-J.** For VsEPs, a statistically significant decrease in P1-N1 VsEP

amplitudes is seen in males (*p = 0.02*) and when sexes are pooled (*p = 0.003*). **H, K.** IP CGRP had no

significant effects on ABR or VsEP latencies.

Fig. 2: Systemic CGRP increases Acoustic Startle Responses (ASR) in both quiet and

background noise, suggesting increased sound sensitivity. A. Schematic of startle stimulus

presented in otherwise quiet background conditions. **B.C.** For female mice (**B**, n=6), and male mice (**C**,

n=6), there were significant differences between vehicle and IP injected CGRP ASR measures in quiet

conditions (p<0.0001). All mice were more sensitive to the same sound level of ASR stimulus after

being injected with CGRP. **D.** Schematic of startle stimulus embedded in continuous 60dB SPL

broadband background noise. **E. F.** Again, ASR embedded in noise was also more sensitive to same

sound level after CGRP injection as shown for female mice (**E**, n=6), male mice (**F**, n=6), (p<0.001). All

mice were more sensitive to the same level of ASR stimulus whether delivered in quiet or in BBN after

being injected with CGRP. Moreover, all mice show ASR increases with sound level from 80 to 130dB

SPL with near maximal startle is elicited by 130 dB SPL stimulus levels. Error bars are SEMs.

Fig. 3: CGRP and VC's effects on postural sway. Figures are graphically depicted in the following manner: vehicle control (open), CGRP (closed), female (orange) and male (blue). (A) The assay timeline involved mice being assessed for effects of IP vehicle control (saline) or IP CGRP on day 1 or day 5, respectively. A 5-minute orbital rotation (125 rpm for 5 minutes, orbital radius = 1 cm) was used as the vestibular challenge. **(B and C)** Group 95% ellipse areas during pre-VC testing were computed to be 0.50 ± 0.06 cm^2 for females and 0.4 ± 0.04 cm^2 for males. **(D)** A toy mouse was measured to be 0.02 ± 0.0005 cm^2 for comparison with live animals. **(E)** Violin plot graphically depicts the distribution of average 95% ellipse areas across sex and treatment and Tukey post-hoc analysis was used to determine significant differences, which were also shown in before/after plots. During IP CGRP pre-VC tests, females had higher 95% ellipse areas compared to males by 0.96 ± 0.19 cm^2 (*p = 0.007*). (**F**) The before-after plot shows a trend (*t(p=0.053)*)) for female mice to exhibit greater sway after vestibular challenge with vehicle injection. **(G)** Before-after plot shows during pre-VC testing in females, IP CGRP caused increased 95% ellipse areas by 0.94 ± 0.23 cm^2 when compared to their IP vehicle response (*p = 0.01*). **(H)** For Pre-VC AP sway in females, IP CGRP increased AP ellipse lengths by $0.49 + 0.10$ cm compared to IP vehicle (*p = 0.005*). A marginal increase in female AP lengths was detected in IP vehicle from pre-VC to post-VC, but this was trending *(t (p = 0.056)*). **(I)** For Pre-VC ML sway in females, IP CGRP increased ML ellipse lengths by $0.12 + 0.03$ cm compared to IP vehicle, (*p = 0.02*). In IP-vehicle treated females, an increase (trend) in ML ellipse lengths is observed from pre-VC to post-VC (*t (p = 0.07)*). IP CGRP-treated females had higher AP lengths compared to males during pre-VC by $0.57 + 0.10$ cm, (*p = 0.0002*), but this sex-specific difference was not seen in ML lengths.