Title: Systemic Calcitonin Gene-Related Peptide (CGRP) modifies auditory and vestibular endorgan

electrical potentials, and increases auditory and vestibular hypersensitivities: Implications for vestibular

migraine

Abbreviated Title: CGRP's effect on inner ear endorgan and behavior

Authors: Shafaqat M. Rahman[1], Benjamin Liang[1], Stefanie Faucher[1], Elana Fine[1], Raajan Jonnala[1],

Joseph C. Holt[2,3], Choongheon Lee[3], Anne E. Luebke[1,2*]

Affiliations:

[1]University of Rochester, Department of Biomedical Engineering, Rochester, NY 14627

[2]University of Rochester Medical Center, Department of Neuroscience, Del Monte Institute of

Neuroscience, Rochester, NY 14642

[3]University of Rochester Medical Center, Department of Otolaryngology, Rochester, NY 14642

*Corresponding author: Anne E. Luebke, aluebke@ur.rochester.edu

Number of Pages: 21

Number of Figures: 3

Number of Tables: 0

Number of words for Abstract: 225

Number of words for Introduction: 343

Number of words for Discussion: 617

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: This work was fully supported by NIH R01DC017261.

Abstract:

Migraine is a severe and chronic neurological disorder that affects ~18% of people worldwide, the majority being female (3:1). It is characterized by recurrent, debilitating headaches and heightened sensory sensitivities. People with migraine may also experience a heightened motion sensitivity that leads to episodes of vertigo and imbalance, termed vestibular migraine (VM). Calcitonin Gene-Related Peptide (CGRP) is implicated in migraine and is believed to act on brain meninges, midbrain, or brainstem, and CGRP-based therapeutics have shown efficacy for migraine treatment. CGRP also signals at efferent synapses of the cochlea and vestibular endorgans, but it is unclear if exogenous CGRP can modulate inner ear function and cause heightened behavioral responses consistent with VM. In C57BL/6 mice, we tested if intraperitoneally (IP) delivered CGRP can modulate endorgan potentials to sound [via auditory brainstem responses (ABRs)] and jerk stimulus [via vestibular sensory evoked potentials (VsEPs)]. We also assessed behavioral measures of phonophobia [acoustic startle reflex (ASR)] and imbalance [postural sway-center of pressure (CoP)] after IP CGRP, and observed female mice exhibit heightened sensitivities to IP CGRP in all assays. Male mice showed similar auditory sensitivity and endorgan effects, but no postural sway changes. In conclusion, we show that systemically delivered CGRP modulates the nerve activity of the auditory and vestibular endorgans and elicits behavioral effects that are consistent with symptoms in a disorder like VM.

Significance statement:

Calcitonin Gene-Related Peptide (CGRP) has been implicated in migraine, and CGRP-based therapeutics have shown efficacy in the treatment of migraine headaches. CGRP is also present in efferent synapses of the inner ear, but we questioned if increases in systemic CGRP can act directly on inner ear endorgans and modify corresponding behavioral responses. In this study, we determined systemic CGRP changes cochlear and vestibular endorgan potentials and produces behaviors suggestive of phonophobia and postural sway. Peripheral changes in auditory and vestibular compound action potentials following systemic CGRP administration suggest CGRP can act directly on the inner ear, which provides a basis for CGRP's involvement in migraine with vertigo and as a therapeutic target.

Introduction:

Migraine is a debilitating chronic neurological disorder that is listed by the World Health Organization (WHO) as the most debilitating neurological disorder worldwide and is estimated to affect 18% of women and 6% of men (Goadsby, Lipton, and Ferrari 2002; Lipton et al. 2007). About 42% of people with migraine have a vestibular component causing balance problems and dizziness (Kuritzky, Toglia, and Thomas 1981; Neuhauser et al. 2001; Neuhauser 2007, 2009), termed vestibular migraine (VM). VM is a major cause of vertigo in dizziness clinics (Kayan and Hood 1984) and patients with migraine, and especially VM, exhibit a heightened sense of sound and motion (Ashkenazi et al. 2009; Lopez-Escamez et al. 2014). Despite migraine and VM's prevalence, the pathophysiology remains uncertain (Lempert et al. 2022; Lempert et al. 2012).

Calcitonin gene-related peptide (CGRP) is intensively studied for its role in migraine's sensory hypersensitivities (Russo 2015). CGRP levels are elevated during migraine, and normalization of CGRP levels is associated with the relief of headache pain which is thought to act at either brain meninges or the trigeminal system (Goadsby, Lipton, and Ferrari 2002; Juhasz et al. 2005; Lassen et al. 2002). CGRP also signals at efferent synapses of the cochlea and vestibular endorgans, so we questioned if elevated CGRP levels in migraine could directly affect the inner ear. To that end, we modeled the increase in CGRP during migraine by a systemic intraperitoneal (IP) injection to determine if systemic CGRP can modulate activity of these endorgans (Dickerson et al. 2016; Luebke et al. 2014; Maison et al. 2003). We also questioned if increases in systemic CGRP can cause behaviors suggestive of heightened auditory and vestibular sway consistent with VM.

In this study, we tested wildtype C57BL/6 mice to determine if there is a change in endorgan potentials to sound (measured using auditory brainstem responses - ABRs) and jerk stimulus (measured using vestibular stimulus evoked potentials - VsEPs) after a CGRP injection. We also assessed corresponding behavioral measures of phonophobia (acoustic startle reflex – ASR) and imbalance (postural sway/center of pressure - CoP) after systemic CGRP injection.

Methods:

Animals: C57BL6/J mice were obtained from Jackson Laboratories (JAX #664) and were housed under

a 12-hour day/night cycle under the care of the University Committee on Animal Resources (UCAR) at

the University of Rochester. Mice were housed together in cages (5 mice per cage) with *ad libitum*

access to food, water, and bedding material. A total of 74 mice (37F/ 37M) were used for these studies,

which were powered sufficiently to detect male/female differences. Before experimentation, mice were

equilibrated in the testing room controlled for an ambient temperature between 22-23^{o}C for at least 30

minutes, and remained in this room until completion of experiments. Injections occurred after the

equilibration period. Mice used during behavioral experiments were returned to the vivarium, while mice

used for ABR/VsEP experiments were euthanized after testing. Mice tested were between 2.3 - 6

months of age. This age range in mice correlates to 20-30 years in humans, which is within the age

range that patients develop migraine symptoms (Lessem 2018). All animal procedures were approved

by the University of Rochester's IACUC committee and performed in accordance with NIH standards.

VsEP and ABR Protocol: C57BL/6 mice were anesthetized with IP urethane/xylazine (1.2 g/kg/20

mg/kg). VsEP and ABR recording methods have been described in detail elsewhere (Lee, Holt, and

Jones 2017). Briefly, to record VsEP and ABR, subcutaneous electrodes were placed on the midline

over the nuchal crest (G1), behind the right pinna (G2), and on the hip (ground). A noninvasive head

clip was used to secure the mouse's head to an electromechanical shaker. During VsEPs, the shaker

delivered a transient (jerk) head translation in the naso-occipital axis (± X). To confirm the absence of

auditory responses during VsEP testing, a binaural forward masker (intensity: 92 dB SPL; bandwidth:

50~50 kHz) was presented with the use of a free-field speaker (FF1, TDT). For ABR recordings, the

same electrodes were used to record ABRs, which were evoked by a single rectangular voltage pulse

of 0.03 ms duration (corresponding to 33 kHz cut-off frequency). These click stimuli were presented via

a free-field speaker, which was the same driver for acoustic masking during the VsEP recording. A

noninvasive rectal thermometer or thermistor was used to monitor the animals' core temperature at

38.0 ± 0.2°C. The standard signal averaging technique described in previous studies (Lee, Holt, and

Jones 2017) was used to record both VsEP and ABR, as shown in **Fig. 1A**. Recordings of both VsEP

and ABR were made in a soundproof booth. Both VsEPs at maximal intensity +6 dB re: 1 g/ms and

ABRs at maximal intensity 100 dB SPL, were recorded as a stable baseline before a single IP injection

of either vehicle control (PBS) or 0.1 mg/kg CGRP was given. Control amplitude and latencies are

depicted as the averages of the 2-3 measurements taken during baseline before injection. Following

single IP injection of either vehicle or CGRP, VsEP and ABR recordings were continuously measured

where significant shifts in VsEP and ABR responses were first observed in 10 mins and reaching a

plateau within 15 mins of drug injection (post-CGRP; data not shown). Thus, amplitudes and latencies

depicted as post-saline/post-CGRP are averages of the 2-3 measurements taken after 20-30 min post-

injections.

Drug administration: Injections were performed intraperitoneally (IP) with a fine gauge (30ga) insulin

syringe. Dulbecco PBS served as the vehicle control. IP CGRP was prepared at 0.1 mg/kg (rat α-

CGRP, Sigma). Injection volumes were calibrated so that each animal received 1 injection per testing

day at ~100 ul. For behavioral experiments, mice were tested ~20 minutes after IP delivery of either

vehicle or CGRP.

Behavioral test of phonophobia: Acoustic startle reflex (ASR)

We used the acoustic startle response (ASR) to assess heightened sound sensitivity as a mouse

surrogate behavior for phonophobia. Experiments were conducted within a sound-attenuating room

(IAC, Bronx, NY) with Sonex foam lined walls. Mice were placed individually in a wire-mesh cage that

is 5 cm wide, 7 cm long, and 4 cm high, and having free sound penetration. The testing cage was

oriented so that the mouse's head faced a Tucker Davis ES2 electrostatic speaker (Masker Speaker)

located 46 cm directly in front of the mouse's head (60dB BBN). The startle speaker was a Radio

Shack tweeter that was suspended 15 cm above the mouse (80-130 dB). The acoustic startle speaker

and its supports, the pedestal, acrylic shelf, and the table on which the apparatus was placed were all

covered with echo absorbing foam and carpeting. Force of the acoustic startle response (ASR) was

transduced by an accelerometer and amplitude scored as integrated root mean square voltage for 100

ms after delivery of the ASR.   Experimental stimuli were controlled, and responses were recorded

using custom MATLAB (MathWorks) software. We tested acoustic startle series in quiet and in noise.

There were 12 conditions in this session, with startle stimuli ranging from 80 to 130 dB SPL, and

delivered either in silence or in a continuous background noise. When present, background noise was

on for the duration of the inter-trial interval and broadcasted from the masker speaker, as shown

schematically in **Fig. 2**.

<u>Behavioral test of postural sway</u>

Postural sway was examined using the center of pressure (CoP) test which acts as a mouse surrogate

behavior for static imbalance. Mice were weighed and placed on a force plate designed to measure

forces in the X, Y, Z and moments in the XY, YZ, and XZ directions. We used the AMTI Biomechanics

Force platform (model HEX6x6) and AMTI automated acquisition software to obtain forces and

moments and then MATLAB code was used to calculate CoP measures based on a confidence ellipse

enclosing 95% of the CoP trajectory values recorded in a single trial. During a given test, mice were

first allowed to freely maneuver on the force plate for 2-5 minutes to acclimate to the novel

environment. An accessory plexiglass guard was placed around the force plate to prevent mice from

moving off the test plate. After acclimation, ~ 10 baseline trials were taken of mice with each trial

indicating a CoP area (resolution per CoP measurement: 300 samples per second). These trials were

taken when the mouse showed no movement and its four paws were touching the force platform. To

assess the effect of a vestibular challenge (VC), mice were then subjected to a constant orbital rotation,

125 rpm for 5 minutes, and were placed back onto the force platform. After a five-minute recovery

period, an additional 10 trials were recorded of the mouse's sway to determine post-VC effects on CoP

as schematized in **Fig. 3**. Mice were first tested after systemic delivery of IP vehicle (20 min after

injection), with sway measured before and after the VC. Four days after vehicle testing, the same mice

were injected with IP CGRP and were tested again 20 minutes after injection (timing correlated with

endorgan potential changes).   While systemic CGRP's effect is believed to dissipate two to three hours

post–injection, we retested after four days to eliminate any residual dizziness due to VC or lingering

CGRP effects. We staggered sequence of IP injections in some animals by injecting CGRP followed by

vehicle testing on the next day. We also injected vehicle 2x to assess test/retest reliability and saw no

differences. Unreasonably large CoP values (> 15 cm^2) suggested mice were either moving during the

trial or that they were at the edge of the force plate, and those values were excluded based on a 10%

ROUT analysis (GraphPad Prism 9.0). Even with values excluded, a minimum of eight different 95%

ellipses were computed per mouse in a given testing condition. Furthermore, we excluded mice that

were outliers by running a second 10% ROUT on averaged CoPs per testing condition.

<u>Statistical Analyses</u>

All statistical analyses were conducted in GraphPad Prism 9.5. Analyses were conducted

separately in females and males. Repeated measure ANOVA (RM-ANOVA) and paired t-tests were

the primary statistical tools. Tukey post hoc analysis was used to assess treatment differences in

females or males, and Bonferroni post hoc analysis was used to assess treatment differences between

sexes. Significance was set at $p < 0.05$ for all analyses.

Results:

<u>IP CGRP's Effects on ABR:</u>

Representative traces are shown for ABR recordings in female and male C57BL6/J mice **(Fig. 1B-1C)**.

Female control ABR amplitudes (n = 6) were computed to be 3.60 ± 0.22 µV but were increased to 4.28

± 0.25 µV after IP CGRP (one-tailed paired t-test; t, df = 3.99, 5; p = 0.005). Likewise, a similar trend

was observed in males as their control amplitudes (n = 7) were computed to be 4.31 ± 0.5 µV and

increased to 4.66 ± 0.44 µV after IP CGRP (one-tailed paired t-test; t, df = 1.94, 6; p = 0.05) **(Fig. 1F)**.

Wave 1 ABR amplitudes were normalized so that post-CGRP measurements highlight the change in

amplitude from their respective control measurements (Δ wave1 amplitude) **(Fig. 1G)**. When sexes are

pooled (n = 13), a significant Δ increase of 0.50 ± 0.13 µV was observed in wave 1 amplitudes due to

IP CGRP (one-tailed paired t-test; t, df = 3.9,12; p = 0.001). Wave 2-1 ABR latencies were calculated

as the difference between P2 and P1 latencies for control and post-CGRP; however, no differences

were observed in latency differences **(Fig. 1H).** A separate group of mice was tested for the effects of

PBS as the vehicle (data not shown, *n* = 4), but no differences were seen in Δ wave1 amplitude or

latency differences between control and post-vehicle (Δ wave1 amplitude $_{vehicle - control}$ = -0.19 ± 0.21 μV).

IP CGRP's Effects on VsEPs:

Similar to ABR tests, representative traces are shown for VsEP recordings in females and males (**Fig.**

1D-1E). Female P1-N1 VsEP control amplitudes (*n* = 6) were computed to be 0.68 ± 0.06 μV and were

reduced to 0.55 ± 0.06 μV after IP CGRP (one-tailed paired t-test; *t, df* = 1.96, 5; *t* (*p* = 0.054)) **(Fig. 1I)**.

Likewise, male P1-N1 VsEP control amplitudes (*n* = 8) were computed to be 0.95 ± 0.10 μV but were

reduced significantly to 0.76 ± 0.10 μV after IP CGRP (one-tailed paired t-test; *t, df* = 3.00, 7; *p* < 0.01).

P1-N1 amplitudes were normalized so that post-CGRP measurements highlight the change in

amplitude from their respective control (Δ P1-N1 amplitude), and both sexes were additionally pooled

(Fig. 1J). When pooling all animals (*n* = 14), a significant Δ reduction of -0.16 ± 0.04 uV in P1-N1

amplitude is observed due to IP CGRP (one-tailed paired t-test**;** *t, df* = 3.65, 13**;** *p* = 0.002).

P1, N1, and P2 latencies were also assessed for VsEPs, and these latencies are depicted as mean ±

SEM for females and males during control and post-CGRP tests **(Fig. 1K).** While no significant

differences were observed due to IP CGRP, modest delays in P1 were observed due to IP CGRP in

either sex. IP CGRP increased female P1 by 26.6 microseconds and male P1 by 22.5 microseconds

compared to P1 control latencies. IP CGRP also increased female N1 by 30 microseconds but had no

effect on Male N1. However, these latency changes were not statistically significant. Comparatively in

mice treated with IP vehicle, no significant P1-N1 amplitude or latency differences were observed

between control and post-vehicle VsEP (Δ P1-N1 amplitude $_{vehicle - control}$ = -0.03 ± 0.08 μV).

CGRP's effect on sound sensitivity

Our sound sensitivity measure in the mouse model makes use of the acoustic startle response (ASR) in

quiet and in background noise. Mice were assessed for the effects of IP CGRP as compared to IP

vehicle on their baseline. The ASR experimental setup is illustrated in **Fig. 2 A** (quiet) and **Fig. 2D**

(background noise 60 dB). In both IP vehicle and IP CGRP injected animals, an increase in ASR was

detected with increasing loudness of the startle stimulus (ES). All animals tested exhibited robust

startle responses (n = 6F/ 6M). When ASR was tested after IP CGRP injection and compared to IP

vehicle, all animals exhibited reduced ASR thresholds in both quiet and noise. IP CGRP significantly

increased sound sensitivity to the startle stimulus in quiet (**Fig. 2 B, C**) and in background noise (**Fig. 2**

E, F) in both female and male mice at all sound levels ($p < 0.0001$ quiet, $p < 0.001$ in noise).

<u>CGRP's and VC's effects on Postural Sway</u>

Mice were assessed for postural sway by determining center of pressure (CoP) ellipse area. An

experimental timeline is shown in **Fig. 3A**. Twenty minutes after delivery of IP vehicle and without a

vestibular challenge (VC), group averages of 95% ellipse areas were computed to be 0.5 ± 0.06 cm^2 for

females and 0.4 ± 0.04 cm^2 for males **(Fig. 3B and C).** For comparison, a deadweight (toy mouse) was

also measured on the force platform, and its average 95% ellipse area was computed to be $0.02 \pm$

0.0005 cm^2 **(Fig. 3D).** A different cohort of mice (n = 10F/ 10M) was assessed for a test-retest of 95%

ellipse areas after IP vehicle treatment, and testing occurred four days apart. A 2-way repeated

measures ANOVA with Bonferroni post-hoc analysis indicated no significant differences in either sex

during the test-retest for females (mean difference = 0.10 ± 0.3 cm^2) or for males (mean difference = -

$0.14 + 0.3$ cm^2) ($p > 0.99$ for either sex). In either sex, 2-way ANOVAs with Tukey post hoc analyses

were used to assess the factors of CGRP and VC on 95% CoP ellipse areas in mice (n = 14F/ 10M)

(Fig. 3E). Differences were seen in females due to IP CGRP (F (1.00, 13.00) = 4.77, p = 0.05) and the

VC (F (1.00, 13.00) = 4.49, p = 0.05). In females after IP vehicle, the VC induced higher CoP ellipse

compared to pre-VC trials (p = 0.02) as shown in **Fig. 3F**. When comparing pre-VC trials, IP CGRP

induced increases in female CoP areas by 0.94 ± 0.23 cm^2 compared to IP vehicle (p = 0.003) as seen

in **Fig. 3G.** Females also had 0.96 ± 0.15 cm^2 higher CoP areas than males based on a separate 2-way

ANOVA assessing sex differences in pre-VC, CGRP and post-VC, CGRP results (p = 0.001).

The major and minor axes of 95% ellipse areas were also analyzed with 2-way ANOVAs across

the factors CGRP and VC (**Fig. 3H and I**). The major and minor axes lengths serve as correlates for

anteroposterior (AP) and mediolateral (ML) sway magnitudes. A 2-way RM ANOVA on major axes

indicated significant effects in females due to IP CGRP (F (1.00, 13.00) = 5.59; p = 0.03) and VC (F

(1.00, 13.00) = 5.41; p = 0.04) whereas these effects were not observed in males. In comparison,

ANOVAs that analyzed minor axes indicated a VC effect in only females (F (1.00, 13.00) = 7.46; p =

0.02). Tukey post-hoc indicated IP CGRP induced larger AP ellipse lengths by 0.49 + 0.10 cm in

females compared to their IP vehicle trials (p = 0.001). Also, in IP vehicle-treated females, there was an

increase in AP ellipse lengths during the post-VC test versus pre-VC (p = 0.02). IP CGRP-treated

females exhibited greater AP lengths compared to males during pre-VC testing by 0.57 + 0.08 cm (p <

0.001). ML lengths increased after the VC in IP vehicle-treated females (p = 0.02), and IP CGRP-

treated females had higher ML ellipse lengths compared to their IP vehicle response (p = 0.007). No

other significant changes were observed in AP and ML sway.

Discussion:

Using electrophysiological techniques (ABR/VsEP), we observed that intraperitoneally delivered

CGRP modulates endorgan function in the inner ear with a maximum ~20- 30 minutes after injection.

Exogenous CGRP increases wave 1 amplitudes of the ABR but reduces VsEP P1-N1 amplitudes,

suggesting at least one action of systemic CGRP is at the efferent signaling sites of the auditory and

vestibular periphery. We did not characterize wave 2 and later peaks, as we were primarily interested in

wave 1/P1-N1 as an indicator of sensory processing changes at the periphery.

Furthermore, we observed that systemic CGRP - during this 20-to-30-minute timeframe -

increased sound sensitivity in male and female C57BL/6J mice, whereas it only increased postural

sway in females.

A few studies in animal models have provided additional evidence for CGRP's role in vestibular

dysfunction and sound sensitivity. Consistent with current VsEP results where we observed delays in

VsEP latencies due to systemic CGRP (though not statistically significant), the loss of αCGRP In

transgenic mouse lines significantly enhanced activation timing of the macular sensors through

changes at the endorgan level (Jones et al. 2018). Intracerebral-ventricular (ICV) injection of

nitroglycerin (NTG) caused impaired balance beam walking in rats, and immunohistochemistry

detected more CGRP expression in the vestibular nucleus of the NTG rat model than was observed in

controls. The administration of a CGRP-receptor antagonist (olcegepant-BIBN4096BS) was shown to

be effective in alleviating impaired balance beam walking (Zhang et al. 2020; Tian et al. 2022).   In

addition, a recent study suggested CGRP has an excitatory role in the auditory nerve response, as

intra-cochlear CGRP infusion was shown to enhance spontaneous and sound-driven auditory nerve

activity in guinea pigs, in comparison to αCGRP-null mice where a loss of nerve activity excitation was

observed (Le Prell et al. 2021; Allen and Luebke 2017). Thus, the auditory nerve may be hyper-

sensitized to CGRP's effects during a migraine.

Peripheral effects of systemic CGRP administration on VsEP responses is somewhat surprising,

as the post-CGRP injection VsEP potentials were reduced and this result requires further examination

in future studies. However, this observation is relevant to VM, which is characterized by sensory

misintegration to stimuli, and thus reduced vestibular sensitivity that occurs concurrently with a

heightened auditory sensitivity may provide a basis for the occurrence of these symptoms.

Additionally, it is unclear why we did not find significant changes in males during postural sway

testing. While more males exhibited significant changes in their VsEP measures, we presume the lack

of CGRP-induced behavioral effects on sway may be due to other postural compensatory mechanisms

present in male mice. Yet, we did discover a robust sex difference when assessing mouse postural

sway, as female mice were much more affected than their male counterparts to injected CGRP.

Similar female-specific sexual dimorphism has been observed in other animal assays assessing

CGRP's effects in migraine but we are the first to show such differences in a preclinical, mouse

surrogate behavior for postural sway (Recober et al. 2009; Paige et al. 2022).

Small molecule blockers of the CGRP receptor 'gepants' have been shown to enter the

periphery, brainstem, and midbrain areas. Ongoing studies will determine if blocking CGRP receptor

signaling by small molecules ('gepants") or by larger monoclonal blocking antibodies (mAbs) - that may

or may not cross blood brain barrier (BBB), but are considered permeable to various peripheral ganglia

- can reduce these sound and imbalance sensitivities due to CGRP (Noseda et al. 2020; Tian et al.

2022; Zhang et al. 2020). The current evidence in favor of anti-CGRP mAbs for treating VM is

compelling, as a prospective observational cohort study over 18 months observed significant reduction

in vertigo and headache frequency in fifty VM patients treated with various anti-CGRP mAbs (Russo et

al. 2023).

References:

Allen, Paul D., and Anne E. Luebke. 2017. 'Reflex Modification Audiometry Reveals Dual Roles for Olivocochlear Neurotransmission', *Frontiers in Cellular Neuroscience*, 11.

Ashkenazi, A., A. Mushtaq, I. Yang, and M. L. Oshinsky. 2009. 'Ictal and interictal phonophobia in migraine-a quantitative controlled study', *Cephalalgia*, 29: 1042-8.

Dickerson, I. M., R. Bussey-Gaborski, J. C. Holt, P. M. Jordan, and A. E. Luebke. 2016. 'Maturation of suprathreshold auditory nerve activity involves cochlear CGRP-receptor complex formation', *Physiol Rep*, 4.

Goadsby, P. J., R. B. Lipton, and M. D. Ferrari. 2002. 'Migraine--current understanding and treatment', *N Engl J Med*, 346: 257-70.

Jones, S. M., S. Vijayakumar, S. A. Dow, J. C. Holt, P. M. Jordan, and A. E. Luebke. 2018. 'Loss of alpha-Calcitonin Gene-Related Peptide (alphaCGRP) Reduces Otolith Activation Timing Dynamics and Impairs Balance', *Front Mol Neurosci*, 11: 289.

Juhasz, G., T. Zsombok, B. Jakab, J. Nemeth, J. Szolcsanyi, and G. Bagdy. 2005. 'Sumatriptan causes parallel decrease in plasma calcitonin gene-related peptide (CGRP) concentration and migraine headache during nitroglycerin induced migraine attack', *Cephalalgia*, 25: 179-83.

Kayan, A., and J. D. Hood. 1984. 'Neuro-otological manifestations of migraine', *Brain*, 107 (Pt 4): 1123-42.

Kuritzky, A., U. J. Toglia, and D. Thomas. 1981. 'Vestibular function in migraine', *Headache*, 21: 110-2.

Lassen, L. H., P. A. Haderslev, V. B. Jacobsen, H. K. Iversen, B. Sperling, and J. Olesen. 2002. 'CGRP may play a causative role in migraine', *Cephalalgia*, 22: 54-61.

Le Prell, Colleen G., Larry F. Hughes, David F. Dolan, and Sanford C. Bledsoe. 2021. 'Effects of

Calcitonin-Gene-Related-Peptide on Auditory Nerve Activity', *Frontiers in Cell and*

Developmental Biology, 9.

Lee, C., J. C. Holt, and T. A. Jones. 2017. 'Effect of M-current modulation on mammalian vestibular

responses to transient head motion', *J Neurophysiol*, 118: 2991-3006.

Lempert, T., J. Olesen, J. Furman, J. Waterston, B. Seemungal, J. Carey, A. Bisdorff, M. Versino, S.

Evers, A. Kheradmand, and D. Newman-Toker. 2022. 'Vestibular migraine: Diagnostic criteria1',

J Vestib Res, 32: 1-6.

Lempert, T., J. Olesen, J. Furman, J. Waterston, B. Seemungal, J. Carey, A. Bisdorff, M. Versino, S.

Evers, and D. Newman-Toker. 2012. 'Vestibular migraine: diagnostic criteria', *J Vestib Res*, 22:

167-72.

Lessem, Sarah E. 2018. "QuickStats: Percentage of Adults Who Had a Severe Headache or Migraine

in the Past 3 months, by Sex and Age Grou - National Health Interview Survey, United States,

2018." In *Morbidity and Mortality Weekly Report (MMWR)*, edited by National Center for Health

Statistics.

Lipton, R. B., M. E. Bigal, M. Diamond, F. Freitag, M. L. Reed, W. F. Stewart, and Ampp Advisory

Group. 2007. 'Migraine prevalence, disease burden, and the need for preventive therapy',

Neurology, 68: 343-9.

Lopez-Escamez, J. A., J. Dlugaiczyk, J. Jacobs, T. Lempert, R. Teggi, M. von Brevern, and A. Bisdorff.

2014. 'Accompanying Symptoms Overlap during Attacks in Meniere's Disease and Vestibular

Migraine', *Front Neurol*, 5: 265.

Luebke, A. E., J. C. Holt, P. M. Jordan, Y. S. Wong, J. S. Caldwell, and K. E. Cullen. 2014. 'Loss of

alpha-calcitonin gene-related peptide (alphaCGRP) reduces the efficacy of the Vestibulo-ocular

Reflex (VOR)', *J Neurosci*, 34: 10453-8.

Maison, S. F., R. B. Emeson, J. C. Adams, A. E. Luebke, and M. C. Liberman. 2003. 'Loss of alpha
CGRP reduces sound-evoked activity in the cochlear nerve', *J Neurophysiol*, 90: 2941-9.

Neuhauser, H. K. 2007. 'Epidemiology of vertigo', *Curr Opin Neurol*, 20: 40-6.

———. 2009. '[Epidemiology of dizziness and vertigo]', *Nervenarzt*, 80: 887-94.

Neuhauser, H., M. Leopold, M. von Brevern, G. Arnold, and T. Lempert. 2001. 'The interrelations of
migraine, vertigo, and migrainous vertigo', *Neurology*, 56: 436-41.

Noseda, R., A. J. Schain, A. Melo-Carrillo, J. Tien, J. Stratton, F. Mai, A. M. Strassman, and R.
Burstein. 2020. 'Fluorescently-labeled fremanezumab is distributed to sensory and autonomic
ganglia and the dura but not to the brain of rats with uncompromised blood brain barrier',
Cephalalgia, 40: 229-40.

Paige, C., I. Plasencia-Fernandez, M. Kume, M. Papalampropoulou-Tsiridou, L. E. Lorenzo, E. T.
David, L. He, G. L. Mejia, C. Driskill, F. Ferrini, A. L. Feldhaus, L. F. Garcia-Martinez, A. N.
Akopian, Y. De Koninck, G. Dussor, and T. J. Price. 2022. 'A Female-Specific Role for
Calcitonin Gene-Related Peptide (CGRP) in Rodent Pain Models', *J Neurosci*, 42: 1930-44.

Recober, A., A. Kuburas, Z. Zhang, J. A. Wemmie, M. G. Anderson, and A. F. Russo. 2009. 'Role of
calcitonin gene-related peptide in light-aversive behavior: implications for migraine', *J Neurosci*,
29: 8798-804.

Russo, A. F. 2015. 'Calcitonin gene-related peptide (CGRP): a new target for migraine', *Annu Rev
Pharmacol Toxicol*, 55: 533-52.

Russo, Cinzia Valeria, Francesco Saccà, Simone Braca, Mattia Sansone, Angelo Miele, Antonio
Stornaiuolo, and Roberto De Simone. 2023. 'Anti-calcitonin gene-related peptide monoclonal
antibodies for the treatment of vestibular migraine: A prospective observational cohort study',
Cephalalgia, 43: 03331024231161809.

Tian, R., Y. Zhang, Q. Pan, Y. Wang, Q. Wen, X. Fan, G. Qin, D. Zhang, L. Chen, Y. Zhang, and J.

Zhou. 2022. 'Calcitonin gene-related peptide receptor antagonist BIBN4096BS regulates

synaptic transmission in the vestibular nucleus and improves vestibular function via

PKC/ERK/CREB pathway in an experimental chronic migraine rat model', *J Headache Pain*, 23:

35.

Zhang, Y., Y. Zhang, K. Tian, Y. Wang, X. Fan, Q. Pan, G. Qin, D. Zhang, L. Chen, and J. Zhou. 2020.

'Calcitonin gene-related peptide facilitates sensitization of the vestibular nucleus in a rat model

of chronic migraine', *J Headache Pain*, 21: 72.

382
383   Fig. 1
384

385
386 Fig. 2
387

388
389 Fig. 3
390

Figure Legends:

Fig. 1. *A,* Schematic illustrating mouse setup for VsEP/ABR recordings captured during baseline

(control) and 20-30 minutes after intraperitoneal (IP) injections of either PBS vehicle or 0.1 mg/kg

CGRP in male (M) and female (F) C57BL6/J mice. *B-C,* Representative click-evoked ABR waveforms -

in duplicate -were collected for ABR at a maximal intensity of 100 dB SPL, and waves were labeled

(wave1 – wave 4). *D-E,* Likewise, representative VsEP waveforms were collected at +6 dB re: 1 g/ms

having three, labeled response peaks (p1, n1, and p2). Orange and blue colors are used to distinguish

female and male tracings post-CGRP, respectively. The following symbols and colors are used: female

- orange, male - square, control - open, CGRP - closed. Sample sizes list - ABR (n = 6F/7M), VsEP (n =

6F/8M). *G & J,* Before/after plots indicate the change in amplitude post-CGRP compared to a control

response normalized to zero. *F-G,* A significant increase is seen in wave 1 ABR amplitudes post CGRP

for female (p = 0.005), male (p = 0.05), and also when sexes are pooled (p = 0.001). *I-J,* For VsEPs, a

statistically significant decrease in P1-N1 VsEP amplitudes was seen in males (p < 0.01), a trend was

seen in females (t (p = 0.054)), and when sexes are pooled (p = 0.002). IP CGRP had no significant

effects on *(H)* ABR or *(K)* VsEP latencies. No amplitude or latency effects were observed in a separate

group of mice (*n* = 4) that were tested for baseline control and IP saline (not shown).

Fig. 2. Systemic CGRP increases Acoustic Startle Responses (ASR) in both quiet and

background noise, suggesting increased sound sensitivity. *A,* Schematic of startle stimulus

presented in otherwise quiet background conditions. *B-C*, In female mice (*B*, n = 6), and male mice (*C*,

n=6), there were significant differences between vehicle and IP CGRP ASR measures in quiet

conditions (p < 0.0001). All mice were more sensitive to the same sound level of ASR stimulus after

being injected with CGRP. *D,* Schematic of startle stimulus embedded in continuous 60dB SPL

broadband background noise. *E-F,* ASR embedded in noise was also more sensitive to same sound

level after CGRP injection as shown for female mice (*E*, n = 6), male mice (*F*, n = 6), (p < 0.001). All

mice were more sensitive to the same level of ASR stimulus whether delivered in quiet or in BBN after

being injected with CGRP. Moreover, all mice showed ASR increases with sound level from 80 to

130dB SPL with near maximal startle is elicited by 130 dB SPL stimulus levels. Error bars are SEMs.

Fig. 3. CGRP and VC's effects on postural sway. Symbols and colors are depicted in a similar scheme as previous figures. *A,* Timeline involved mice being assessed for effects of IP vehicle control (saline) or IP CGRP on day 1 or day 5, respectively. A 5-minute orbital rotation (125 rpm for 5 minutes, orbital radius = 1 cm) was used as the vestibular challenge, and mice were repeatedly tested (n = 14F/10M). *B-C,* Group 95% ellipse areas during pre-VC testing were computed to be 0.50 ± 0.06 cm^2 for females and 0.40 ± 0.04 cm^2 for males. *D,* A toy mouse was measured to be 0.02 ± 0.0005 cm^2 for comparison with live animals. *E,* During IP CGRP pre-VC tests, females had higher 95% ellipse areas compared to males by 0.96 ± 0.19 cm^2 (*p = 0.01*). *F,* Females exhibit greater sway after vestibular challenge with vehicle injection (*p = 0.02*). *G,* Before-after plot shows during pre-VC testing in females, IP CGRP caused increased 95% ellipse areas by 0.94 ± 0.23 cm^2 when compared to their IP vehicle response (*p = 0.003*). *H,* For Pre-VC AP sway in females, IP CGRP increased AP ellipse lengths by 0.49 + 0.10 cm compared to IP vehicle (*p = 0.001*). An increase in female AP lengths was detected in IP vehicle from pre-VC to post-VC, but this was trending *(p = 0.02)*. *I,* For Pre-VC ML sway in females, IP CGRP increased ML ellipse lengths by 0.12 + 0.03 cm compared to IP vehicle, (*p = 0.02*). In IP-vehicle treated females, an increase (trend) in ML ellipse lengths is observed from pre-VC to post-VC (*p = 0.02*), and IP CGRP-treated females had increased ML ellipse lengths compared to their vehicle response by 0.12 + 0.03 cm (*p = 0.007*).