Population genomics of an emergent tri-species hybrid zone

Running title: Genomics of a tri-species hybrid zone

Libby Natola[1,2], Sampath S. Seneviratne[3], and Darren Irwin[1,2]

[1]Department of Zoology, University of British Columbia, Vancouver, BC, Canada;

[2]Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada;

[3]Avian Sciences & Conservation, Department of Zoology, University of Colombo, Colombo, Sri

Lanka

Corresponding Author: Libby Natola, libby.natola@gmail.com

Abstract

Isolating barriers that drive speciation are commonly studied in the context of two-species hybrid zones. There is however evidence that more complex introgressive relationships are common in nature. Here, we use field observations and genomic analysis, including the sequencing and assembly of a novel reference genome, to study an emergent hybrid zone involving two colliding hybrid zones of three woodpecker species: Red-breasted, Red-naped, and Yellow-bellied Sapsuckers (*Sphyrapicus ruber, S. nuchalis,* and *S. varius*). Surveys of the area surrounding Prince George, British Columbia, Canada, show that all three species are sympatric, and Genotyping-by-Sequencing identifies hybrids from each species pair and birds with ancestry from all three species. Observations of mate pair phenotypes and genotypes provide evidence for assortative mating, though there is some heterospecific pairing. Hybridization is more extensive in this tri-species hybrid zone than in two di-species hybrid zones. However, there is no evidence of a hybrid swarm and admixture is constrained to contact zones, so we classify this region as a tension zone and invoke selection against hybrids as a likely mechanism maintaining species boundaries. Analysis of sapsucker age classes does not show disadvantages in hybrid survival to adulthood, so we speculate the selection upholding the tension zone may involve hybrid fecundity. Gene flow among all sapsuckers in di-species hybrid zones suggests introgression likely occurred before the formation of this tri-species hybrid zone, and might result from bridge hybridization, vagrancies, or other three-species interactions.

Keywords: *Sphyrapicus*, sapsucker, hybridization, introgression, Genotyping-by-Sequencing

Introduction

Speciation is the process wherein an ancestral species subdivides into multiple isolated

populations which accumulate differences and eventually become separate species incapable of

interbreeding (Coyne & Orr 2004). Understanding how these subdivisions occur and what allows

differentiation are fundamental topics in evolutionary biology. Biologists refer to the forces that

contribute to differentiation as isolating barriers because they prevent gene flow between the

emergent populations. These barriers can be genetic (Dobzhansky 1936), geographic (Mayr

1947), or behavioural (Wells et al. 1978). Contact zones, where differentiating forms come

together, provide excellent context for identifying isolating barriers and observing the process of

speciation (Barton & Hewitt 1981). The two forms may not interbreed if differentiation is

advanced, but early in speciation they might breed and produce offspring (a process called

hybridization). In this situation, the contact is called a hybrid zone, and if differentiation between

the populations is low, hybrid offspring may breed with each other or backcross with either

parental form (a process termed introgression), opening a bridge of gene flow from one species

to the other.

Hybridization is typically studied in the context of species pairs (Barton and Hewitt 1985,

Moore 1977). However, there are many examples of more complex hybridization relationships in

nature (Ottenburghs 2019). In some cases, one species might act as a conduit or a bridge for gene

flow between two other species. If the "bridge species" introgresses with both additional species

in the "triad", alleles can be shared between two species indirectly. Bridge hybridization has

been observed in suckers (*Catostomus commersoni*, *C. latipinnis*, *C. discobolus*; McDonald et al.

2008), *Heliconius* butterflies (*H. melpomene*, *H. cydno*, and *H. pachinus*; Kronforst 2006) and

Darwin's finches (*Geospiza fortis, G. scandens, G. fulginosa*; Grant and Grant 2020). Similarly,

some species may act as introgressive "hubs", interbreeding with many (otherwise non-

interbreeding) species, such as mallard (*Anas platyrhynchos*) hybridizing with many duck

species, common pheasant (*Phasianus colchicus*) breeding other Pheasant species, and European

herring gull (*Larus argentatus*) which hybridizes with other gulls, potentially facilitating gene

flow between pairs of species that do not directly hybridize (Ottenburghs 2019). In other

systems, multiple species may hybridize among each other creating more direct introgressive

relationships, such as in the *Bos* genus (Wu et al. 2018). Despite there being so many examples

of multispecies hybridization, few studies to date have closely examined the phenomenon in bird

taxa (Ottenburghs 2019).

Red-breasted, Red-naped, and Yellow-bellied sapsuckers (*Sphyrapicus ruber, nuchalis,*

and *varius*) are a recently radiated group of North American woodpeckers. Two well-studied

sapsucker hybrid zones exist in British Columbia, Canada: one in the Mackenzie region between

Red-breasted and Yellow-bellied Sapsuckers; and another involving Red-breasted and Red-

naped Sapsuckers farther south, near the Williams Lake area (Howell 1953, Seneviratne et al.

2016). These zones have historically been separated, but while the Yellow-bellied x Red-

breasted zone has been moving south, the Red-breasted x Red-naped zone has crept north. It was

long predicted that these two hybrid zones would converge into a single three-species contact

zone (Scott et al. 1976, Walters et al. 2002a), but surveys conducted by one of the authors (SS) in

the area from 2009-2011 suggested this interaction had not yet occurred.

By 2018, observations of local ornithologists (Dr. Ken Otter 2018, personal

communication) and birders (as submitted to eBird; https://ebird.org) suggested the convergence

of these hybrid zones had occurred in the Prince George/Quesnel, BC area. Hybrid zones

involving three hybridizing species are rare and allow us the opportunity to examine

evolutionary dynamics of multiple stages of speciation, often referred to as the speciation

continuum (Chhatre et al. 2018, Tarroso et al. 2014). The two colliding zones provide an

opportunity to compare hybrid zone dynamics and genomic patterns in the three-species hybrid

zone as opposed to the two paired species zones. This allows us to directly differentiate patterns

of three species gene flow from two.

In this study, we use field and molecular methods to describe multispecies hybridization

among sapsuckers. We aim to answer the questions: 1) to what degree do Red-naped, Red-

breasted, and Yellow-bellied Sapsuckers hybridize in sympatry? and 2) how do these interactions

vary among different hybrid zones? We address these questions with surveying, mate pair, and

genomic data of sapsuckers in the putative tri-species hybrid zone. We compare patterns of

introgression in this zone with those in each of the two-species hybrid zones.

Materials and Methods:

Surveying

To assess sapsucker density and habitat preferences we performed standardized

presence/absence surveys between 5 and 10 AM during the months of May and June in 2019 and

2020. We did not survey in rainy, cold, or windy weather. We surveyed residential and Forest

Service roads by stopping once every kilometer and playing a standard recording on a Bluetooth

speaker. This recording included two minutes of silence, 3.5 minutes of drumming and call

noises from Red-breasted, Red-naped, and Yellow-bellied sapsuckers, and two more minutes of

silence. During the silences, we listened for sapsucker calls and drumming, and looked for

sapsuckers. We marked each survey point as either having sapsuckers present or absent. If

sapsuckers were present, we identified them to species based on plumage markings. Hybrids

could not be identified to species cross because Red-breasted x Red-naped hybrids and Red-

breasted x Yellow-bellied hybrids cannot be reliably differentiated by plumage. We identified

mated pairs as those seen excavating, feeding, or defending a nest together, or responding to call

playback with paired dry chatter and moth flight display (Walters et al. 2002b).

Sample collection

To reflect the population-wide genotypic composition in the tri-species hybrid zone, we

attempted to catch every sapsucker we found, regardless of phenotype, and made a concerted

effort to fill in any geographical sampling gaps. We trapped 107 birds along two intersecting

transects, one ~115 km N-S on highway 97 from north of Bear Lake, BC to just south of Soda

Creek, BC and another spanning ~140 km E-W between Vanderhoof, BC and McBride, BC on

highway 16. We caught sapsuckers using either dipnets at a nest or canopy nets accompanied by

audio and decoy lures. In hand, we collected ~ 50μL of blood from the brachial vein of each bird

and stored it in Queen's lysis buffer. We banded each individual for later observation and to

prevent resampling. We also collected morphometric data and photographed each bird.

Woodpeckers replace feathers in a reliable sequence as they age, so we determined age classes

based on wear and replacement in the secondary and primary covert feathers (Pyle 1997, pp 163-

181). For analyses, we collapsed age classes to second year (SY, bird hatched previous summer)

or after second year (ASY, bird hatched prior to previous summer). In this study we analyze our

tri-species hybrid zone data along with data from two di-species hybrid zones sampled by

Seneviratne et al. (2016, 2012): one Red-breasted x Yellow-bellied Sapsucker zone and one Red-

breasted x Red-naped Sapsucker zone. For sample collection, DNA extraction, and GBS

sequencing methods of di-species hybrid zones please reference Seneviratne et al. (2016, 2012).

Reference genome assembly

To create a *de novo* reference genome, we submitted tissues of one female Red-breasted

Sapsucker voucher specimen to the University of Delaware Sequencing & Genotyping Center for

Pacific Biosciences Single Molecule Real Time (SMRT) Consensus Long Read (CLR)

sequencing on a Pacific Biosciences Sequel II sequencer. Sequencing yielded 172X coverage.

We assembled the genome in canu v. 1.9 (Koren et al. 2017), resulting in a 1.6 Gb assembly with

an N50 of 4.7 Mb, and an LG50 of 75, with 7596 contigs. To make the assembly haploid for

downstream analysis we kept one haplotig from each of the heterozygous contigs with the Purge

Haplotigs pipeline (Roach et al. 2018). We then masked repetitive elements using Repeat Masker

v. 4.1.1 (Smit et al. 2013), which marked 15.93% of the genome as repeats, the majority of

which (13.71% of the genome) are LINE retroelements. We assessed completeness of our

assembly with the program BUSCO v. 5.1.2 (Manni et al. 2021) and the aves_odb10 lineage

dataset with 8338 BUSCOs. Our assembly is 96% complete with 88% complete single copy

BUSCOs, 8% complete duplicated, 0.9% fragmented, and 3.1% missing. Finally, we used the

Chromosemble tool in Satsuma v. 2.0 (Grabherr et al. 2010) to align our contigs to a Golden-

fronted Woodpecker (*Melanerpes aurifrons*) genome (Wiley & Miller 2020). Our final

assembled genome had 269 scaffolds.

GBS protocol

We extracted DNA using a standard phenol-chloroform extraction and subsequently

followed the genotyping-by-sequencing (GBS) protocol detailed by Elshire et al. (2011) and

Alcaide et al. (2014) using the enzyme *PstI*. We followed lab protocols of Seneviratne et al.

(2016) with the following changes. We incubated the digestion with 1 µL *PstI*, 2 µL 10X buffer,

6µL of common adaptor, 6µL of barcode, and 5 µL of 20ng/µL template DNA at 37˚C for 2

hours. We ran the ligation incubation at 22˚C for 1 hour then 65˚C for 10 minutes. For PCR, we

included 5µL of 5x Phusion buffer, 0.5 µL 10mM dNTPs, 1.25 µL forward and reverse primers,

12.75 µL UltraPure water, and 0.25 µL of PhusionTaq. We ran the reaction protocol as follows:

98 °C for 30 seconds and 18 cycles of: 98°C for 10sec, 65°C for 30sec and 72°C for 30 sec. This

is followed by an extension of 72°C for 5min, followed by 4°C. We used gel extraction to select

for 400-500 bp fragments. We sent 100 ng of DNA from 289 sapsuckers (Supplemental

Materials Table 1) to Genome Quebec for sequencing on either an Illumina HiSeq4000 PE150 or

the NovaSeqSP 6000 PE150 (Table S1) (Illumina, San Diego, CA).

GBS Processing

We demultiplexed all reads using a custom script from Irwin et al. (2018), trimmed reads

with Trimmomatic v 0.38 (Bolger et al. 2014), then aligned them to our Red-breasted Sapsucker

reference genome using BWA v 0.7.17 (Li 2013). We further realigned around indels and called

genotypes using GATK v 3.8 (Van der Auwera & O'Connor 2020). We kept variant sites and

removed all indels and sites that had mapping quality < 20, heterozygosity > 0.6, more than 70%

missing data, GQ < 10, or minor allele frequency ≥ 0.05. We calculated φ in VCFtools v 0.1.16

(Danecek et al. 2011) to estimate relatedness, randomly removed one individual from each pair

of birds with an estimated 4th degree relationship or closer (phi ≥ 0.0224), and filtered out

individuals with more than 60% missing data. After filtering, we retained 251 individuals and

50,639 SNPs.

Data analysis

Using the variants data, we visualized genomic relationships by generating a genomic

PCA in R (R Core Team 2011) using scripts adapted from Irwin et al. (2018). To generate

ADMIXTURE plots, we further pruned data for linkage disequilibrium, removing sites with >

0.6 correlation in 1000 bp windows. We ran ADMIXTURE v 1.3.0 (Alexander & Lange 2011)

for $K = 1$-6 with cross validation and calculated standard errors using 2000 bootstrap replicates.

Cross validation error indicated $K = 3$ had the best predictive accuracy, consistent with the

expectation of this being a three-species hybrid zone. We used the R package Ternary v 1.2.2

(Smith 2017) to create ternary plots of ADMIXTURE data.

Assortative Mating

We evaluated the strength of assortative mating (AM) among the three species in the tri-

species hybrid zone with plumage-based species assignment of birds from 91 sighted pairs. We

classified all birds to pure species or hybrid by plumage, and because different hybrid species

crosses are indistinguishable by eye all hybrids are lumped. First, we calculated the proportion of

conspecific pairs we found in the population (including pairs involving two phenotypic hybrids).

To evaluate whether this proportion falls within expectations under random mating, we

randomized pairings based on observed species and hybrid frequencies 1000 times and

calculated the proportion of conspecific pairs for each randomization. We checked if the

conspecific proportion in our data was within the upper 5% of the simulated distribution. Then,

we calculated plumage AM using the equation:

$$AM = 1 - \frac{2 * \left(\frac{OH}{EH}\right)}{\left(\frac{OC}{EC}\right) + \left(\frac{OH}{EH}\right)}$$

where *OH* signifies observed heterospecific pairings, *EH* is expected heterospecific pairings, *OC*

is observed conspecific pairings, and *EC* is expected conspecific pairings, in accordance with

methods in Scordato et al. (2020). We were able to capture and genotype both mates from 15

pairs. We charted the ADMIXTURE K=3 cluster data of the 15 genotype pairs on a Ternary plot

and measured the Euclidian distances between each pair's Cartesian coordinates on the Ternary

plot. For each individual, we plotted its p coefficient (proportion of genome attributed to Red-

breasted Sapsucker ancestry in ADMIXTURE) by its measured genotypic mate distance. We ran

a linear regression to model predictability of genotypic mate distance based on p coefficient. Our

reasoning with this latter analysis is that the majority of the birds in the hybrid zone are

phenotypic Red-breasted Sapsuckers, such that they may on average mate with more similar

individuals than the other phenotypes are able to.

Hybrid Fitness

To better understand hybrid performance in the tri-species hybrid zone, we asked whether

there is evidence that degree of ancestry admixture is associated with survival. We did this by

examining whether different age classes differed in their average amount of admixture. We were

unable to collect hatching or nestling data in the field, so we first predicted the ADMIXTURE

cluster genotypes of birds in the next generation from our genotypic mate pair data to see if

hybrid sapsuckers were equally represented after one year as we might expect from breeding

data. We averaged ADMIXTURE clusters p, q, and z among mated pairs to predict offspring

ancestry, then calculated a measure *m*, calculated as 1 minus the maximum value across all three

ADMIXTURE clusters, for these "hatch year" birds. Next we distilled admixture within all

sequenced birds to *m* and plotted this measure against sapsucker age class (HY, SY, or ASY) to

understand if hybrids are less likely to survive to older ages, as might be predicted if post-zygotic

isolation causes them to be less fit than pure birds. In addition, we ran a logistic regression on

genomic admixture (*m*) and age class to see if amount of admixture predicted likelihood of

surviving to older age classes.

Results:

The genomic data support the hypothesis that there is a three-species hybrid zone in east-

central British Columbia. In the PCA (Fig. 2), each species formed a distinct group with the

allopatric samples at the distal ends of each cluster. The tri-species hybrid zone samples showed

each species formed a separate genomic cluster. The main axis of variation (PC1, 16.3%)

identifies differences between Yellow-bellied Sapsuckers and Red-breasted/Red-naped

Sapsuckers, and these latter sister species split out across PC2 (3.9%) (Fig. 2). It is important to

note that the relative density of each species varies widely. There is a much higher proportion of

phenotypically Red-breasted birds (70.4% surveyed, 53.5% sampled) than any other species,

followed by phenotypically hybrid birds (20.2% surveyed, 29.1% sampled). We found relatively

few Red-naped (4.0% surveyed, 8.1% sampled) and Yellow-bellied (5.4% surveyed, 9.3%

sampled) birds in the hybrid zone.

There are genomically intermediate birds placed between each pair of species clusters,

with the most being between Red-breasted and Red-naped, and the fewest intermediates between

Red-naped and Yellow-bellied (Fig. 2). Intermediate Red-breasted/Red-naped birds fell along a

wide gradient of PC2 values, but there was a distinct gap separating the two species' clusters.

Intermediates involving Yellow-bellied sapsuckers tended to be more clumped in the very center

of the X-axis, equidistant from each parent cluster. The majority of birds placed between the

major species clusters were collected in the tri-species hybrid zone. Finally, though most birds

were placed in genotypic clusters that matched their phenotypic assignment, there were

genotypic/phenotypic mismatches, shown in the PCA as samples that are the "wrong" color for

their predicted species cluster, pure looking birds with hybrid genotypes, or intermediate looking

birds with pure genotypes. Most mismatched birds were from the tri-species hybrid zone.

ADMIXTURE results similarly showed that many individuals in the study belong to one

of the three distinct species (clusters of ancestry values q, p, or z > 0.995, 14 Red-breasted, 21

Red-naped, 38 Yellow-bellied) (Fig. 3). There were also many birds with both Red-breasted and

Red-naped ancestry (q + p > 0.995, 57 birds). Several birds showed primarily Red-breasted x

Yellow-bellied ancestry (q + z > 0.995, 10 birds), and seven had primarily Red-naped x Yellow-

bellied ancestry (p + z > 0.995). The biggest group of birds had a combination of all three

ancestries (116 birds, hereafter termed "muttsuckers" after McDonald et al. (2008)), many of

which had a low proportion of Yellow-bellied, some Red-naped, and a high proportion of Red-

breasted ancestry. In this way the ADMIXTURE explicitly shows the three-species ancestry

found in many birds in this region.

The ternary plot shows these data more intuitively, with many birds placed on the axis

shared by Red-breasted and Red-naped, especially towards the Red-breasted corner, a few along

the Yellow-bellied axes, and many points within the triangle, signifying three-species ancestry

(Fig. 4). With the tri-species hybrid zone, di-species hybrid zones, and allopatric zones birds split

into separate Ternary plots, it is clearer that the three species are well isolated in allopatry (Fig.

4a), and there is increasingly more admixture as we move from di-species hybridization (Fig. 4b)

to tri-species hybridization (Fig. 4c). It's also more obvious that we caught no "pure" Red-naped

Sapsuckers in the tri-species hybrid zone. Again, there are fewer intermediates showing recent

Yellow-bellied ancestry, and those that do exist tend to be equidistant from either parental group.

The ternary plot also reiterates the "mismatch" pattern in which phenotypic assignment does not

consistently mirror genotypic assignment.

Based on the observed species frequencies, the expected proportion of conspecific

pairings is 0.56. Our pairing randomization showed the observed proportion of conspecific

pairings (0.67) falls in the 99.7th quantile of our randomized pairing distribution, which gives us

high confidence that sapsuckers are mating conspecifically more frequently than would be

expected under random mating ($p <0.01$). The AM calculations show that matings between

phenotypic species are under-represented, but matings between "pure" individuals and plumage

intermediates are more common (Table 1). The genotype mate pairs on the ternary plot have

short genomic distances between mate pairs near Red-breasted Sapsuckers, and much longer

genomic distances connecting pairs with hybrids and lower-density (Fig. 5a). The scatter plot of

p coefficient of randomly chosen mate by pair's genomic distance (Fig. 5b) shows a strong

negative correlation ($R^2 = 0.8$, $F = 52.1$, $p < 0.0001$), meaning that the more Red-breasted

ancestry a bird has, the more closely related it tends to be to its mate.

Looking at age classes as predicted by genomic admixture, we found that hybrids were

not over-represented in the SY or ASY age class (HY $t = 0.11$, $p = 0.91$, SY $t = -0.85$, $p = 0.4$,

Fig. 6), so we have no evidence hybrids suffer lower survival rates than pure species.

Discussion:

Di- vs Tri-species Hybrid Zone Dynamics

Our survey and genomic data conclusively show there is a tri-species hybrid zone in the

Prince George region of BC. Individuals from all three species breed here, often near one or both

of these other sapsucker species. However, we note that the great majority of Red-naped

sapsuckers in the tri-species hybrid zone have some admixture. There is also a demographic

imbalance, with high representation of Red-breasted phenotypes and genotypes. We present

evidence that the sapsuckers breed in all three species pairs in the tri-species hybrid zone, and

there appears to be back-crossing in every direction, indicating direct gene flow and

introgression among all three species. Additionally, we found many "muttsuckers" with ancestry

from all three species, meaning there is likely also indirect gene flow as well, in which some

geneflow could occur between two species that does not involve them directly interbreeding if

hybrids of two species breed with a third species. Backcrossed admixed birds (largest

ADMIXTURE cluster < 0.76) account for 18.9% of birds sampled in the tri-species hybrid zone,

and non-backcrossed hybrids account for 3.3% (largest cluster < 0.51). However, allopatric

individuals from each species form distinct groups (Fig. 4a), showing that the evidence for

substantial introgression is limited to the sympatric region.

Compared to the tri-species hybrid zones, we see less hybridization and introgression in

general within the di-species hybrid zones (as in Grossen et al. 2016, Seneviratne et al. 2016).

There are very few that could be direct hybrids (0.7% had a maximum ADMIXTURE cluster <

0.51), and less than 7% are recently admixed (largest cluster < 0.76), showing that the evidence

for substantial introgression is limited to the sympatric region. As expected by their closer

evolutionary relationship, Red-breasted and Red-naped hybridize the most, with mitochondrial

genetic distance producing estimates of divergence times of 1.090 million years ago (Yellow-

bellied) and 0.32 million years ago (Red-breasted and Red-naped) (Weir & Schluter 2004). Very

little introgression seems to occur between either species and Yellow-bellied, but it is important

to note that there are early generation backcrosses between Yellow-bellied and both Red-naped

and Red-breasted, so at least some hybrids can be viable and fertile. It is intriguing that there is

more admixture in the tri-species hybrid zone than the di-species hybrid zones. Our data don't

have the power to explain this pattern, so it opens some interesting new questions. Does the

addition of a third species reduce assortative mating? Could the distorted demographics in the tri-

species hybrid zone affect mate availability in low-density species, therefore promoting

hybridization? We know habitat influences hybrid zone dynamics in sapsuckers (Natola et al., *in*

preparation). The large stretch of aspen dominated valleys of Prince George area is surrounded

by conifer dominated hills. The latter is less frequented by sapsuckers (Sampath Seneviratne,

personal observation). Perhaps climate or landscape features in this region inherently promote

hybridization.

Interestingly, in the di-species hybrid zones there are several individuals that have

ancestry from the third species that is not one of the two species apparently hybridizing there.

For instance, there are 13 samples from the Red-breasted x Yellow-bellied hybrid zone (bolded

diamonds on Figure 4b) with more Red-naped ancestry than Yellow-bellied ancestry, in addition

to a bird from the same hybrid zone with Red-naped but no Red-breasted ancestry, and a bird

from the Red-breasted x Red-naped zone (bolded square Fig. 4b) with no Red-breasted ancestry

but appears to be a relatively recent Red-naped x Yellow-bellied backcross. Also, one

muttsucker from the Red-breasted x Yellow-bellied hybrid zone (bolded diamond Fig.4b) has

substantial ancestry from all three species. The existence of genotypes in the di-species hybrid

zones in which they don't "belong", suggests greater introgression than previously reported.

These samples were all collected approximately 10 years before local ornithologists noticed the

convergence of the two zones. In addition, we see evidence of many advanced generation

muttsuckers (at least one species' ancestry > 0%, but < 10%). With an estimated generation time

of 1.9 years (Seneviratne et al. 2016), we expect the eight years, or fewer than 5 generations,

since sampling showed no tri-species contact, is insufficient to produce the proportion of late

generation backcrosses with tri-species ancestry we found. It may be possible that the Red-

breasted Sapsuckers, which have existed in allopatry between the two hybrid zones, have acted

as a conduit for introgression of alleles between all three species for generations. Alternatively,

as two di-species hybrid zone birds appear to have very little Red-breasted ancestry at all, and a

Red-breasted x Red-naped hybrid from the Red-breasted x Yellow-bellied hybrid zone has very

little Yellow-bellied ancestry (Fig. 4b), there may be intermittent contact through occasional

long-distance dispersal. In any case, the data showing existence of muttsuckers or birds with

ancestry of species excluded from their hybrid zone is evidence that three-way gene flow among

all three species predates our detection of the tri-species hybrid zone.

Tri-species hybrid zone selection

Though there is a lot of hybridization and introgression in the tri-species hybrid zone, and

to a more limited extent throughout the di-species hybrid zones, there is no evidence of a

collapse into a hybrid swarm. If birds were mating randomly and hybrids were as fit on average

as the parental species, the ternary plot (Fig. 4c) would eventually show most birds in the middle

and few at the corners. Instead, our samples are overwhelmingly concentrated to the corners and

edges of the plot. We propose that the Prince George, BC region is therefore a three-way tension

zone (Barton & Hewitt 1989), maintained by reduced migration in and out of the hybrid zone

and selection against hybrids or hybridization itself. Therefore, we suspect some form of

selection against hybrids, perhaps enhanced by assortative mating, is maintaining species

boundaries and reducing gene flow in this system.

We investigated evidence for assortative mating. In regard to plumage, sapsuckers are not

mating randomly, as evidenced by the higher proportion of conspecific pairings than random,

and the high AM values above the diagonal in Table 1, suggesting some amount of mate

preference for phenotypically similar individuals (note, however, low sample sizes of Yellow-

bellied Sapsuckers in the AM analysis). This provides evidence for a partial mating barrier

between the species. That holds up with the genomic data, where we saw few F1 hybrids (Fig.

4c). However, there appears to be little isolation between phenotypic hybrids and phenotypic

species. Assortative mating was high with Yellow-bellied Sapsuckers, but we had a very small

sample size for this species in these analyses. Assortative mating rates were far lower between

hybrids and pure Red-breasted or hybrids and pure Red-naped Sapsuckers. This is supported by

our evidence of several backcrosses given the number of F1s we found, that hybrids may be

acceptable mates. Note however, the apparent offset of plumage phenotype and genotype in

many birds. Given this offset and acceptance of hybrid birds, it is likely that sapsucker mate

choice is not based solely on plumage. These patterns also suggest plumage may be controlled by

a few genes of large effect, making it an excellent candidate for a genome-wide association

study.

The analysis of the 15 genotyped pairs shows that individuals with Red-breasted

genotypes tend to pair with each other, and hybrids and Red-naped genotypes (no Yellow-bellied

were in this analysis) tend to mate with far more genetically distant sapsuckers (Fig. 5a, b).

These patterns could be explained by a preference for genetically similar individuals. We

speculate this is an effect of relative demography. There are so few Red-naped in the area that

those who are looking for mates "settle" for a mate with an ill matched genome rather than

foregoing breeding entirely. However, with as much outcrossing as we see in the full genomic

dataset it is unlikely assortative mating alone can explain the overall species maintenance (Irwin

2020), so we speculate there must be some selection against hybrids as well. Sources of low

hybrid fitness against hybrids might be related to habitat specialization, as the different sapsucker

species have different preferred climatic conditions and forest compositions and hybrids could be

at a disadvantage (Natola & Burg 2018, Walters et al. 2002b, 2002a). Seasonally migratory

behaviour might also cause low hybrid fitness, as all three of these species have different

migratory strategies (Walters et al. 2002b, 2002a) and intermediate routes could be

disadvantageous (Bensch et al. 1999, Delmore et al. 2016, Scordato et al. 2020). Additionally,

there could be genomic incompatibilities which decrease hybrid fitness.

We investigated one component of hybrid fitness, survival from hatching to adulthood,

by examining age classes and admixture rates. We didn't find any evidence that hybrids are less

likely to survive to adulthood based on our mate pair data, or that second year birds are less

likely to survive to after second year based on our plumage data (Fig. 6). The n = 15 for our mate

pair data is small and may have affected our ability to detect true patterns. While we didn't find

evidence of reduced hybrid survival, it is possible hybrids produce fewer offspring on average

and their fitness is thus diminished. Genetic incompatibilities, caused for instance by copy

number variation and chromosomal inversions, could explain a decrease in hybrid fecundity as

these interactions typically affect the zygotes formed from heterokaryotypic gametes (Rieseberg

& Willis 2007). More research into hybrid fecundity and genomic synteny among sapsucker

species could clarify whether these interactions may be lowering fitness of hybrids.

Conclusions

Our data show that increasing the number of hybridizing species from 2 to 3 changes

hybridization dynamics. Hybridization and introgression are more pervasive in the tri-species

hybrid zone than the di-species hybrid zones. However, that introgression does not extend to

allopatric populations, and hybridization is less common than expected under random mating.

Therefore, this region serves as a tri-species tension zone. As yet, it is unclear what sources of

selection against hybrids maintains the three species in the hybrid zone, yet hybrid survival to

adulthood doesn't seem to explain the pattern. A genomic study on loci under selection and

resistant to introgression may clarify the underlying causes of isolation in sapsuckers.

Furthermore, we show that hybridization is more complex than traditional two-species

interactions in sapsuckers. Indirect gene flow between the three species appears to predate the

tri-species hybrid zone we identified, meaning this history of triple introgression might be the

result of conduits, vagrancies, or other three-species contact. This signifies that sapsucker species

interactions were more complex than we previously suspected, and may indicate a more

pervasive biological pattern present in other taxa.

Acknowledgements:

Kathy Martin, Loren Rieseberg, and Dolph Schluter provided key feedback on all stages

of project development. Afnan Ali, Avery Bartels, Jamie Clarke, Gabriel Pang, Rashika

Ranasinghe, and Kang Wang assisted in field sample collection. This research was funded by the

Natural Sciences and Engineering Research Council of Canada (RGPIN- 2017-03919 and

RGPAS-2017-507830 to DI), the Society of Canadian Ornithologists' Baillie Award, the

American Museum of Natural History's Chapman Award, the Explorer's Club's Mamont Grant,

the Society for the Study of Evolution's R.C. Lewontin Early Award, and the Warner and

Hildegard Hesse Research Award. Thanks to Gavin Hanke (Royal BC Museum) Jocelyn Hudon

(Royal Alberta Museum), Chris Stinson and Ildiko Szabo (Beaty Biodiversity Museum), and

Kevin Winker (University of Alaska Museum) for sharing tissue samples.

References:
431

432
Data Accessibility:
  All scripts, data, and analyses are available on a GitHub repository, and genetic data
will be posted to NCBI. Both will be made public upon manuscript acceptance.

Benefits Generated:
  Benefits from this research accrue from the sharing of our data and results on public
databases as described above.

Author Contributions:
  LN and DI conceived the project. LN assembled reference genome, sampled and
prepared tri-hybrid zone GBS libraries for sequencing, processed reads and conducted analyses,
with advice from DI. SS sampled and prepared GBS libraries from both di-hybrid zones. LN
wrote the manuscript with input from DI and SS.

447　　*Tables:*

449　　*Table 1. Assortative mating matrix based on phenotypic data. Values on and below the diagonal*

450　　*represent the number of pairs found for each cross type. Above the diagonal, italicized, are the*

451　　*calculated AM values indicating the degree of assortative mating between each pair from*

452　　*random mating (0) to complete isolation (1) or completely disassortative mating (-1).*

	Hybrid	*Red-breasted*	*Red-naped*	*Yellow-bellied*
Hybrid	5	*0.41940999*	*0.53861004*	*1*
Red-breasted	21	53	*0.8360373*	*0.66377018*
Red-naped	6	2	2	*1*
Yellow-bellied	0	1	0	*1*

	Hybrid	*Red-breasted*	*Red-naped*	*Yellow-bellied*
Hybrid	5	*0.41940999*	*0.53861004*	*1*
Red-breasted	21	53	*0.8360373*	*0.66377018*
Red-naped	6	2	2	*1*
Yellow-bellied	0	1	0	*1*

457   *Figures:*

458
459   *Figure 1. Map of sample locations for phenotypic hybrids (gray) Red-breasted (red), Red-*
460   *breasted x Yellow-bellied (orange), Red-breasted x Red-naped (purple), Red-naped (blue),*
461   *and Yellow-bellied (yellow) sapsuckers across southeast Canada.*
462

463
464 *Figure 2. Principal Component Analysis of genomic data, each sample is depicted by its*
465 *phenotypic species assignment including tri-species hybrid zone hybrids (grey squares), Red-*
466 *breasted (red circles), Red-naped (blue triangles), Yellow-bellied (yellow diamonds), hybrids*
467 *from Re-breasted x Yellow-bellied di-species hybrid zone (orange x), and hybrids from Red-*
468 *naped x Red-breasted di-species hybrid zone (purple +).*
469

ADMIXTURE

Individual

470
471 *Figure 3. Plot of ADMIXTURE k = 3 data for all samples. Ancestry coefficient was partitioned*
472 *to Red-breasted (q, red), Red-naped (p, blue), and Yellow-bellied (z, yellow) clusters. Plot is*
473 *broken up to show pure birds from each cluster, each species pair cluster, and birds with*
474 *ancestry from all three species.*
475

a.

b

c

Figure 4. Ternary plots for allopatric zone (a), di-species hybrid zones (b), and tri-species

hybrid zone (c) samples. Sample colors depict phenotypic species assignment for Red-breasted

(red), Red-naped (blue), Yellow-bellied (yellow), and hybrid (grey) birds. In panel b, the Red-

breasted x Red-naped di-species hybrid zone samples are shown as squares and the Red-

breasted x Yellow-bellied di-species hybrid zones are shown as diamonds, and birds with

unexpected genotypes given their sampling location are shown in bold. Points are jittered to

show all samples.

a.

b.

Figure 5. Ternary plot of mate pair genotypes from tri-species hybrid zone (a). Colors depict
phenotypic species assignments for Red-breasted (red), Red-naped (blue) and hybrid (grey).
Grey lines connect paired mates. Proportion of Red-breasted Sapsucker ancestry by Euclidian
distance between mate pairs' Cartesian coordinates (b) shows one mate chosen randomly per
pair along with linear regression line, equation, and R^2 value. Shaded areas indicate 95%
confidence level interval.
495

496
497 *Figure 6. Box and whisker plots showing measured admixture (m), for projected hatch year (HY)*
498 *birds, and plumage identified second year (SY) and after second year (ASY) birds.*