Functional connectivity fingerprints of the frontal eye fields and inferior frontal junction in the dorsal vs. ventral prefrontal cortex

Orhan Soyuhos[1] | Daniel Baldauf[1]

[1]Centre for Mind/Brain Sciences (CIMeC), University of Trento, 38068 Rovereto, Italy.

E-mail addresses: orhan.soyuhos@studenti.unitn.it (O. Soyuhos), daniel.baldauf@unitn.it (D. Baldauf).

Abstract

Neuroimaging evidence suggests that the frontal eye field (FEF) and inferior frontal junction (IFJ) govern the encoding of spatial and non-spatial (such as feature- or object-based) representations, respectively, both during visual attention and working memory tasks. However, it is still unclear whether such contrasting functional segregation is also reflected in their underlying functional connectivity patterns. Here, we hypothesized that FEF has predominant functional coupling with spatiotopically organized regions in the dorsal ('where') visual stream, whereas IFJ has predominant functional connectivity with the ventral ('what') visual stream. We applied seed-based functional connectivity analyses to temporally high-resolving resting-state magnetoencephalography (MEG) recordings. We parcellated the brain according to the multimodal Glasser atlas and tested, for various frequency bands, whether the spontaneous activity of each parcel in the ventral and dorsal visual pathway has predominant functional connectivity with FEF or IFJ. The results show that FEF has a robust power correlation with the dorsal visual pathway in beta and gamma bands. In contrast, anterior IFJ (IFJa) has a strong power coupling with the ventral visual stream in delta, beta, and gamma oscillations. Moreover, while FEF is directly phase-coupled with the superior parietal lobe in the beta band, IFJa is directly phase-coupled with the middle and inferior temporal cortex in delta and gamma oscillations. We argue that these intrinsic connectivity fingerprints are congruent with each brain region's function. Therefore, we conclude that FEF and IFJ have dissociable connectivity patterns that fit their respective functional roles in spatial vs. non-spatial top-down attention and working memory control.

KEYWORDS

frontal eye field, inferior frontal junction, functional connectivity, prefrontal cortex, visual pathways, visual attention, working memory.

1 | INTRODUCTION

At any moment, we face an overwhelming stream of rich information from the surrounding environment. Our brain must continuously re-evaluate all the incoming sensory stimulation to prioritize what is relevant for our behavioral goals (Baldauf & Deubel, 2010; Desimone & Duncan, 1995; Fecteau & Munoz, 2006). The prefrontal cortex (PFC), the maestro in orchestrating a range of cognitive functions in the human brain, plays an essential role in this respect (Duncan, 2001; Fuster, 2001; Miller & Cohen, 2001; Rossi et al., 2008). It crucially contributes to the central functions of selective attention and working memory by top-down modulating priority for incoming stimuli according to the context of the ongoing task demands (Jerde et al., 2012; Ptak, 2012). Its distinct control circuits' interareal communications with the rest of the cortex enable the brain to focus its limited perceptual resources on specific locations in space, or on certain features or objects (Baldauf & Desimone,

2014; Bichot et al., 2019; Corbetta & Shulman, 2002; Gregoriou et al., 2009; Gross et al., 2004; Mangun, 1995; Siegel et al., 2008). In this context, human neuroimaging techniques and invasive recordings in primates suggest that the representational content encoded during visual selective attention and working memory can further segregate the functional specialization within PFC (Carrasco, 2011; Cavada & Goldman-Rakic, 1989; Giesbrecht et al., 2003; Goldman-Rakic, 1988; McCarthy et al., 1996; Scalaidhe et al., 1999; Paneri & Gregoriou, 2017; Romanski, 2004; Wilson et al., 1993). In particular, two specific control sites located in the posterior lateral PFC (pIPFC), namely the Frontal Eye Field (FEF) (Kelley et al., 2008; Petit & Pouget, 2019; Vernet et al., 2014) and the Inferior Frontal Junction (IFJ) (Baldauf & Desimone, 2014; Brass et al., 2005; Derrfuss et al., 2005; Muhle-Karbe et al., 2014), show differential neural activity depending on the stimulus and task domain.

Experiments in which attention is covertly directed to a visuospatial cue have led to increased neural activity in FEF, apart from its oculomotor function (Armstrong et al., 2009; Thompson et al., 2005; Thompson & Bichot, 2005). Additionally, muscimol injection and suprathreshold microstimulation techniques revealed FEF's crucial role in spatial attention (Moore & Armstrong, 2003; Moore & Fallah, 2004; Wardak et al., 2006). In this respect, Thompson and Bichot (2005) reported the existence of a visual salience map in primate FEF and noted that its initial neural activities are not selective for non-spatial features such as color or shape, but that their activity evolves over time to select the target location. Along these lines, human neuroimaging and monkey studies indicated that FEF exerts top-down regulatory influences on the extrastriate visual cortex and increases the response gain to retinotopically corresponding targets directly or indirectly through its connections to the intraparietal sulcus (IPS) (Bressler et al., 2008; Gregoriou et al., 2009, 2012; Kelley et al., 2008; Merrikhi et al., 2017). Moreover, transcranial magnetic stimulation (TMS) studies advanced these correlational findings with causal evidence and pointed out FEF's key function in the top-down control of visuospatial attention (Heinen et al., 2017; Marshall et al., 2015; Veniero et al., 2021).

In contrast, experimental designs that require attention to non-spatial features or object representations have consistently invoked neural responses in the inferior frontal cortex around IFJ. This evidence mainly accumulated from human neuroimaging studies due to the lack of corresponding sulcal landmarks in the nonhuman primate brain (Bedini & Baldauf, 2021; Donahue et al., 2018). Nevertheless, the studies on the ventral pre-arcuate of monkey PFC (VPA; regarded as the homolog of IFJ) have provided important insights into the region's functioning. For instance, Bichot and colleagues (2015) reported VPA as the source of feature selection. They showed that the inactivation of VPA reduces the feature-based modulation in ipsilateral V4 and impairs monkeys' performance in detecting target objects in the contralateral visual field (Bichot et al., 2019). Likewise, human neuroimaging and TMS studies reported a top-down modulatory role of IFJ in object- and feature-based attention and working memory (Baldauf & Desimone, 2014; Meyyappan et al., 2021; Zanto et al., 2010, 2011; Zhang et al., 2018). For instance, Zanto and colleagues (2010) found that the functional coupling between the right IFJ and V4 was predictive of the extent of attentional modulation during attention to color. Additionally, they demonstrated that the perturbation of IFJ weakens working memory performance for color features (Zanto et al., 2011). Following this, through directional phase coupling, Baldauf and Desimone (2014) additionally showed that IFJ is the source of attentional modulation of object representations in the inferior temporal (IT) cortex and the driver of non-spatial, object-based attention.

Overall, the evidence from task-based studies suggests that the encoding of to-be-attended or to-be-remembered representational contents functionally segregates PFC and reveals two distinct control circuits in the dorsal and ventral PFC: FEF and IFJ. While FEF is involved in the top-down control of spatial attention and working memory, IFJ contributes to the top-down control of non-spatial (feature- and object-based) attention and working memory (for a review see: Bedini & Baldauf, 2021). Nonetheless, it is still unclear to what extent this functionally specific contrast is reflected in these regions' underlying functional connectivity profiles.

1.1 | Research question and hypothesis

The current study aimed to analyze the intrinsic functional connectivity fingerprints of FEF and IFJ to support their functional segregation within PFC. We argue that FEF and IFJ's interareal communication with regions in the dorsal ('where') and ventral ('what') visual pathways (Goodale & Milner, 1992; Mishkin et al., 1983) would be a prerequisite for their functional involvement in spatial and non-spatial attention and working memory control, respectively; similarly to the structural connectivity fingerprints of areas in IT cortex being predictive of their functional selectivity (Osher et al., 2016; Saygin et al., 2011). In this respect, we hypothesize that FEF has predominant functional connectivity with spatiotopically organized areas in the dorsal visual stream, whereas IFJ has predominant functional connectivity with feature- or object-encoding areas in the ventral visual pathway. To this end, we performed a region of interest (ROI) analysis and explored FEF and IFJ's functional connectivity to the rest of the brain. Additionally, we inquired about the dominant directionality of these interactions between FEF/IFJ

and the visual streams. To this aim, we used various functional connectivity metrics (both phase- and power-based) and epoch lengths to consider their effects on the consistency of results.

2 | MATERIALS AND METHODS

2.1 | Participants

We benefited from the 1200 Subjects Release of Human Connectome Project (HCP) dataset that includes resting-state MEG (rMEG) recordings and structural MRI of 95 healthy participants aged between 22 and 35 (Larson-Prior et al., 2013; Van Essen et al., 2013). The Washington University institutional review board approved data acquisition protocols for HCP. Each subject was asked for verbal consent before the sessions. Thirty-six participants are reported to be monozygotic and dizygotic twins. Out of 95 subjects, we selected 55 independent ones (26 female) after randomly excluding one of the pairs from each twin and checking the data quality. The anonymized dataset is provided in ConnectomeDB (db.humanconnectome.org; Hodge et al., 2016).

2.2 | Procedure

Subjects typically spent 2 or 3 days completing the HCP protocol that includes behavioral tests, resting-state functional magnetic resonance imaging (MRI), task-evoked functional MRI, diffusion-weighted MRI, and task-evoked MEG in addition to structural MRI and rMEG sessions. MEG recordings always took place before MRI sessions to prevent the potential effect of residual magnetization caused by the MRI scanner on the superconducting MEG sensors (Van Essen et al., 2013).

2.3 | MRI data acquisition and preprocessing

Washington University - University of Minnesota Consortium of the Human Connectome Project (WU-Minn HCP Consortium) performed Structural MRI on a customized Siemens 3T "Connectome Skyra" at Washington University (Uğurbil et al., 2013). They acquired two high-resolution T1- (T1w) and T2-weighted (T2w) anatomical images of participants using a standard 32-channel Siemens receive head coil (T1w; 3D MPRAGE; 0.7 mm isotropic; FOV, 224 × 224; 256 sagittal slices; TR=2400 ms; TE=2.14 ms; inversion time (TI), 1000 ms; 8-degree flip angle) (T2w; 3D T2-SPACE; 0.7 mm isotropic; FOV, 224 × 224; 256 sagittal slices; TR=3200 ms; TE=565 ms; variable flip angle). Experts reviewed the quality of scans, considering the tissue contrast, blurriness, and artifacts. If necessary additional scans were arranged for a second time. Also, a radiologist examined the existence of any neuroanatomical anomalies. While the normal variations were noted, subjects with abnormal brain anatomy were excluded from the dataset.

ConnectomeDB provides preprocessed structural data resulting from the HCP structural pipeline (Glasser et al., 2013). The second part of this pipeline is mainly based on a modified version of FreeSurfer's recon-all script to reconstruct a two-dimensional cortical surface from a three-dimensional volume (Dale et al., 1999; Fischl, Sereno, & Dale, 1999). We used the resulting files provided inside the Structural Extended Preprocessed package in the ConnectomeDB for the source reconstruction of MEG data.

2.4 | MEG data acquisition

The rMEG recordings were acquired on a whole head Magnes 3600 scanner (4D Neuroimaging, San Diego, CA) in a magnetically shielded room at Saint Louis University (Larson-Prior et al., 2013). The MEG scanner had 248 magnetometers and 23 reference channels (18 magnetometers and five first-order gradiometers). The data were recorded with a sampling rate of 2034.5101 Hz, delta encoded, and saved in a 4D file format. Electrooculography (EOG), electrocardiography (ECG), and electromyography (EMG) signals were also measured simultaneously to enable offline artifact rejection. Moreover, the experimenters used five Head Position Indicators (HPI) to track the head position in reference to the MEG sensors and marked the anatomical landmarks of nasion and the preauricular points to coregister the SQUID array to each individual's structural MRI. They used a Polhemus Fastrak-III system for the spatial digitization of localizer coils and fiducial points. Additional points defining the subject's head shape (about 2400 points) were also digitized to fine-tune the co-registration.

During the rMEG scanning, subjects lay in a supine position. They were instructed to keep their heads still while fixating a red

crosshair at the center of the projection. Their head positions were recorded before and after every three consecutive sessions. Each run took approximately 6 minutes. Before rMEG, the empty room and participant noise recordings were also acquired to monitor the sensor noise and the noise stemming from the participant due to residual magnetization effects.

2.5 | MEG data preprocessing

The ASCII text files that annotate the indexes of bad channels, bad segments, and non-brain components are provided inside the Preprocessed Resting-State package in the ConnectomeDB. To ensure the reproducibility of our results, we decided to utilize them for preprocessing rMEG data. First, the noisy channels and the bad data segments were marked following a visual inspection. Later, the correlation between signals from adjacent sensors and the ratio of their variance to their neighbors were calculated to mark the outliers as bad channels. Additionally, z-scores for all time points in each channel were computed to identify the segments whose score was higher than a given threshold. The clipping and muscle-related artifacts were detected using ft_artifact_clip.m and ft_artifact_muscle.m functions from the FieldTrip toolbox (Oostenveld et al., 2010). Furthermore, an iterative Independent Component Analysis (ICA) was performed to detect faulty channels and bad segments in the raw data by using the list of bad channels and segments identified beforehand as initial input. In subsequent iterations of this ICA, the input was updated by the bad sensors or segments found in the previous iteration until the ICA algorithm could no longer identify any further bad channels/segments.

In the next step of the MEG preprocessing pipeline, ICA was used this time to remove the artifacts related to eye movements, blinks, and heartbeats from the signal. To this end, first, a bandpass (1.3-150 Hz) and a notch Butterworth (59-61 and 119-121 Kz) filter were applied to the MEG, ECG, and EOG data. The previously identified bad channels and segments were cut out from these three recordings. Afterward, ICA was implemented on the preprocessed MEG data twenty times based on different seed values. Power spectral densities (PSD) and power time courses (PTC) were calculated for the independent components (ICs) of each repetition and corresponding time series of ECG and EOG channels. An automatic classification algorithm detected artifacts based on the correlation between ICs and ECG/EOG recordings. It classified ICs into the brain and non-brain components while also considering pink noise and the noise stemming from the power supply (60 and 120 Hz). The classifier's performance was later confirmed by visual inspection. Ultimately, the best iteration with the highest ratio of brain to artifact components from twenty repetitions was selected. The indices of its brain and non-brain components were reported in ASCII text files. After the removal of the bad channels, bad segments, and artifacts from the rMEG data, we parsed the continuous times series into non-overlapping epochs of either 2 s, 5 s, or 10 s to consider the epoch length's effect on the connectivity measures (Brookes et al., 2011; Fraschini et al., 2016).

2.6 | Source reconstruction of MEG signals

For each epoch length, we utilized the Brainstorm toolbox (Tadel et al., 2011) to reconstruct source-level activation on the native cortex of each subject. We first corrected the DC offset of preprocessed rMEG data and downsampled it to 300 Hz. To estimate sensor noise, we computed the noise covariance matrix from raw empty room recording after removing its DC bias. Additionally, we calculated the lead field matrix for an evenly distributed grid of 15002 cortical sources by fitting a single sphere under each sensor (Huang et al., 1999). Based on these gain and noise covariance matrices, we applied the minimum-norm estimation (MNE) method to inverse-model activations in source space (Baillet et al., 2001; Dale et al., 2000). Here, we chose MNE rather than beamformers because it was shown to perform better for spatially distant sources (Hincapié et al., 2017). Afterward, the source level activity for each subject was projected to an MNI average brain template (FreeSurfer fsaverage; Fischl et al., 1999) and parcellated by the multimodal parcellation atlas (MMP1; Glasser et al., 2016). We calculated the scout activations for 360 regions by taking the maximum absolute value across all vertices per scout and multiplying with its sign. We saved them as Fieldtrip structures to perform functional connectivity analyses in the next step.

The MEG Anatomy package provided in the ConnectomeDB enables the automatic co-registration of MEG sensors with the structural MRI. However, it does not allow projecting sources to a standard brain (such as fsaverage). Since we aimed to interpolate source-level activation of each subject onto the fsaverage template for the group level analysis, we used the pial surface provided in the Structural Extended Preprocessed package to reconstruct sources. In this way, we could also increase the resolution of the cortex from 8004 to 15002 vertices for a better representation of cortical folds. However, we still had to use the MEG Anatomy package to retrieve the coordinates of fiducial points for each subject since the individual MRIs were defaced, and the fiducial points were not stated explicitly.

2.7 | Functional connectivity and directionality analyses

In a second step, we conducted seed-based functional connectivity analyses with seed regions of FEF, anterior IFJ (IFJa), and posterior IFJ (IFJp) in both hemispheres by employing the FieldTrip toolbox (Oostenveld et al., 2010). We first carried out frequency analyses on time series data over all epochs while using the multiple tapers based on the Slepian sequence. The frequencies of interest were the delta (1-4 Hz), theta (4-8 Hz), alpha (8-13 Hz), beta (13-30 Hz), and gamma (30-100 Hz) band with the number of spectral smoothing boxes of 1, 1, 1, 2 and 10, respectively. We computed complex Fourier-spectra between all brain parcels for each frequency band. Based on this Fourier representation of data, we calculated the functional connectivity between each of the six seed regions and the rest of the 359 brain parcels. We decided to use both phase- and amplitude-based functional connectivity measures to take into account the possible distinct roles of phase- and amplitude-coupling (Daffertshofer et al., 2018; Siems & Siegel, 2020). In particular, we chose the imaginary part of the Coherency (iCOH) (Nolte et al., 2004), debiased weighted Phase Lag Index (dwPLI) (Vinck et al., 2011), and orthogonalized Power Envelope Correlation (oPEC) (Hipp et al., 2012) since these metrics are corrected for spatial leakage and have higher group-level repeatability compared to other methods (Bastos & Schoffelen, 2016; Brookes et al., 2011; Colclough et al., 2016; Duan et al., 2021).

We performed specific region-of-interest (ROI) and whole-brain exploratory analyses based on the connectivity metrics mentioned above. For ROI analyses, the target MMP1 parcels in Table 1 were chosen based on a systematic bibliographical search of the relevant literature (Felleman & Van Essen, 1991; Glasser et al., 2016; Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Mishkin et al., 1983) and the list reported in Zimmermann et al. (2018). We first selected the mid-higher visual areas that belong to the dorsal or ventral visual pathway. Additionally, we used the retinotopy atlas by Wang et al. (2015) to identify putative retinotopic parcels of the MMP1 atlas and include them in our definition of the dorsal visual stream, as the topographic organization is generally considered a defining feature of this stream. We excluded parcels near the medial wall, parcels from the superior temporal lobe, and the temporo-parietal junction since they seem to be characterized by distinctive connectional profiles (Kravitz et al., 2011, 2013; Mars et al., 2011). This led to a total of 33 regions with specific visual properties and selectivity to visual features (Table 1). We tested whether the spontaneous activity of each parcel in the ventral and dorsal visual pathway has higher functional connectivity with FEF or IFJ. Following this, we additionally questioned whether the regions that do not show preferential connectivity, neither with FEF nor with IFJ, have a statistically significant intrinsic functional connectivity with any of these seed regions when tested against the mean of whole-brain connectivity values per frequency band.

Finally, we analyzed the dominant directional interactions between the seed regions in PFC and the target areas in the visual cortex during the resting state. To this end, we carried out directionality analyses between FEF, IFJa, IFJp, and all the ROIs in Table 1. We computed partial directed coherence (PDC) (Baccalá & Sameshima, 2001) of each unidirectional interaction between seed regions and visual streams. For each pair, we subtracted the PDC values of one unidirectional connection from the other and tested its significance against zero. We chose PDC from a range of directionality measures because it only considers direct interactions between regions, is generally regarded as insensitive to source leakage, and shows high group-level repeatability, for instance, in comparison to the phase slope index (Colclough et al., 2016).

2.8 | Statistical analyses

We implemented paired Wilcoxon signed-rank tests for each type of analysis because the obtained connectivity values had a non-gaussian distribution. For the same reason, we did not apply z-transformation to the correlation matrix resulting from the oPEC metric. Additionally, we used the false discovery rate (FDR) method to correct for multiple comparisons (Benjamini & Hochberg, 1995; Groppe, 2021). In the hypothesis-driven ROI analyses, we corrected multiple comparisons for all 33 regions in Table 1 (+3 seed regions for PDC measure). Whereas, we constrained the significance test for the whole-brain exploratory analyses to one hemisphere and accordingly corrected multiple comparisons for 180 regions. To visualize FEF and IFJ's predominant functional connectivities across frequency bands or hemispheres, we converted the adjusted p-values to z-scores. We summed the z-score of each parcel and divided them into the number of times they were significant. Importantly, we applied paired significance tests separately for FEF, aIFJ, and pIFJ per region. We did not average the connectivity values of aIFJ and pIFJ before statistical analyses since we suspected them of having different functional connectivity fingerprints based on previous evidence (Baker et al., 2018; Bedini & Baldauf, 2021).

Parcel	Description	Visual Stream	Topographical Organization
Dorsal Visual Pathway			
V6	Sixth Visual Area	Dorsal (Glasser et al., 2016)	✓ (Kravitz et al., 2011)
V6A	Area V6A	Dorsal (Glasser et al., 2016; Zimmermann et al., 2018)	✓ (Kravitz et al., 2011; Wang et al., 2015)
V7	Seventh Visual Area	Dorsal (Glasser et al., 2016)	✓ (Glasser et al., 2016; Wang et al., 2015)
IPS1	IntraParietal Sulcus Area 1	Dorsal (Glasser et al., 2016; Zimmermann et al., 2018)	✓ (Wang et al., 2015)
IP1	Area IntraParietal 1	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
MIP	Medial IntraParietal Area	Dorsal (Zimmermann et al., 2018)	✓ (Kravitz et al., 2011; Wang et al., 2015)
VIP	Ventral IntraParietal Complex	Dorsal (Zimmermann et al., 2018)	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
LIPd	Area Lateral IntraParietal dorsal	-	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
LIPv	Area Lateral IntraParietal ventral	Dorsal (Zimmermann et al., 2018)	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
7AL	Lateral Area 7A	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
7PC	Area 7PC	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
7PL	Lateral Area 7P	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
V3A	Area V3A	Dorsal (Glasser et al., 2016; Goodale & Milner, 1992)	✓ (Glasser et al., 2016; Felleman & Van Essen, 1991; Kravitz et al., 2013; Wang et al., 2015)
V3B	Area V3B	Dorsal (Glasser et al., 2016)	✓ (Wang et al., 2015)
V3CD	Area V3CD	-	✓ (Wang et al., 2015)
LO3	Area Lateral Occipital 3	-	✓ (Wang et al., 2015)
MT	Middle Temporal Area	Dorsal (Goodale & Milner, 1992)	✓ (Glasser et al., 2016; Felleman & Van Essen, 1991; Kravitz et al., 2013)
Ventral Visual Pathway			
V8	Eighth Visual Area	Ventral (Glasser et al., 2016)	-
VMV1	VentroMedial Visual Area 1	Ventral (Glasser et al., 2016)	-
VMV2	VentroMedial Visual Area 2	Ventral (Glasser et al., 2016)	-
VMV3	VentroMedial Visual Area3	Ventral (Glasser et al., 2016)	-
PHT	Area PHT	Ventral (Zimmermann et al., 2018)	-
PH	Area PH	Ventral (Zimmermann et al., 2018)	-
TE1a	Area TE1 anterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	-
TE1m	Area TE1 middle	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	-
TE1p	Area TE1 posterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	-
TE2a	Area TE2 anterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	-
TE2p	Area TE2 posterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	-
TF	Area TF	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011; Zimmermann et al., 2018)	-
TGv	Area TG ventral	Ventral (Kravitz et al., 2013; Zimmermann et al., 2018)	-
FFC	Fusiform Face Complex	Ventral (Glasser et al., 2016)	-
PIT	Posterior Infero Temporal	Ventral (Glasser et al., 2016)	-
VVC	Ventral Visual Complex	Ventral (Glasser et al., 2016)	-

TABLE 1 List of parcels from the MMP1 atlas for ROI analyses. We selected ROIs in the visual cortex as belonging to the ventral or dorsal visual stream by considering previous arguments for their inclusion in the dorsal/ventral pathway, receptive field sizes, and the existence of a topographical organization.

2.9 | Ground truth analysis

Lastly, we performed a ground truth analysis to ensure that our analysis pipeline was implemented correctly and provided valid results. We replicated the results from Hipp et al. (2012) paper for the oPEC metric. In the beta band (13-30 Hz), we calculated the power correlation of left and right MT+ with the rest of the brain. We could successfully replicate the seed-based connectivity of MT+ and reproduce the third figure of their paper by using a different dataset, source-reconstruction method (MNE rather than beamforming), and our own processing pipeline. We did not apply any statistical mask to the figure shared in the supplementary material (Figure S1).

3 | RESULTS

We first report the outcomes for 2 s epoch segmentation and ipsilateral functional connectivity. We detail the results for ROI analyses with respect to phase- and power-based functional connectivity metrics. Additionally, we present PDC in the resting state to expand on the direction of the interaction. The abbreviations for the cortical area names are based on Glasser et al. (2016). The supplementary material provides the findings for 5 and 10 s epoch lengths (Figures S5-6), contralateral connectivity (Figure S9), and whole-brain connectivity analyses (Figures S10-14).

3.1 | Predominant functional connectivities

All three connectivity metrics demonstrated that FEF has robust functional connectivity with the regions in the dorsal visual stream, whereas IFJa has stronger functional connectivity with the areas in the ventral visual stream independently of lateralization and frequency band (Figure 1; two-sided Wilcoxon signed-rank test, $p < 0.05$, FDR-corrected). The direct contrast between the ipsilateral connectivity fingerprints of FEF and IFJa (FEF-IFJa) revealed that FEF has a robust power correlation with the dorsal visual stream in beta and gamma bands. In contrast, IFJa has a strong power coupling with the ventral visual stream in delta, beta, and gamma oscillations. Moreover, both iCOH and dwPLI metrics showed that FEF is phase coupled with superior parietal lobule (SPL) and IPS in the beta band, while IFJa is phase-coupled with the inferior and middle temporal cortex by delta and gamma oscillations (Figure 2). Importantly, the anterior and posterior parts of IFJ have different connectivity patterns. Although the direct contrast between the ipsilateral connectivity fingerprints of FEF and IFJp (FEF-IFJp) also indicates FEF's higher functional connectivity to SPL and IPS, we observed that IFJp has weaker functional connectivity to the middle and ventromedial temporal cortex, particularly in the alpha and beta frequencies. Nevertheless, it still achieves a higher power correlation with the ventral visual areas in the delta frequency compared to FEF (Figures S2 and 2).

The phase-based metrics showed that the significant parcels for FEF-IFJa are lateralized in the left hemisphere. The iCOH and dwPLI metrics did not reveal any predominant coupling for FEF or IFJa with the selected ROIs in the right hemisphere. On the contrary, while the strong power correlation for FEF-IFJa was evident in both hemispheres, the predominant power coupling between the left IFJa and ventral visual stream was weaker than in the right hemisphere (Figure S3A). Additionally, we observed an apparent dissociation within the higher frequencies for the power coupling between visual streams and plPFC in the right hemisphere. The right FEF and IFJa functionally segregate the right visual pathways into dorsal and ventral streams in beta and gamma oscillations. Overall, the results from both phase- and power-based metrics illustrate that FEF and the areas in the dorsal visual stream predominantly communicate in beta frequency. In contrast, IFJa and the regions in the ventral visual stream are functionally connected through delta and gamma oscillations.

For FEF-IFJp, we observed predominant phase coupling only for FEF in the alpha and beta bands, in the left and right hemispheres, respectively (Figure S3). AdditionallyFurthermore, the iCOH and dwPLI measures showed that areas PHT, TE1m, and VMV1 in the ventral stream have a more robust phase relation with FEF than IFJp. Like the contrast between FEF and IFJa, a predominant power correlation exists in both hemispheres also for FEF-IFJp. However, we cannot see any predominant power coupling between IFJp and the ventral visual stream in higher frequencies. Also, the left VMV2 in the ventral visual pathway has a higher power correlation with the left FEF rather than IFJp.

Noticeably, areas V7, V3B, MT, V8, PH, FFC, PIT, and VVC do not show predominant connectivity either with FEF, IFJa, or IFJp (Figures 1 and S2). In this respect, our additional analyses for FEF, IFJa, and IFJp's seed-based intrinsic functional connectivities (one-sided Wilcoxon signed-rank test against the average, $p < 0.05$, FDR-corrected) revealed that V7, V3B, MT, V8, PIT, and VVC do not have statistically

FIGURE 1 Predominant functional connectivity maps of FEF and IFJa across frequency bands and hemispheres for **(A)** oPEC, **(B)** iCOH, and **(C)** dwPLI metrics (two-sided Wilcoxon signed-rank test, p < 0.05, FDR-corrected for 33 ROIs). The results shown here are based on 2 s epoch segmentation.

FIGURE 2 Frequency-specific predominant functional connectivity patterns of FEF, IFJa, and IFJp. The width of a line reflects its relative z-score of the connectivity to other nodes (two-sided Wilcoxon signed-rank test, p < 0.05, FDR-corrected for 33 ROIs). The results shown here are based on 2 s epoch segmentation.

significant power or phase coupling with any of these three seed regions (Figure S4). On the other hand, PH has a significant power correlation both with FEF and IFJa. FFC is significantly power coupled with IFJa and IFJp, but not to FEF. Overall, we observed that all three seed regions have a robust power correlation with the parcels in SPL and IPS. They also have statistically significant spontaneous power coupling with the areas in the ventral visual stream.

3.2 | Influences of chosen epoch length

We observed that 2, 5, and 10 s epoch segmentation affects distinct frequency bands and connectivity metrics differently. In the delta band, IFJa has a consistent robust power correlation with the ventral visual stream. However, its dominance in the ventral pathway decreases in beta and gamma oscillations with increasing segmentation length. In contrast, FEF's power coupling with the dorsal visual stream is consistent in beta and gamma frequencies (Figure S5). Additionally, the results for iCOH are less consistent as the epoch length increases and the number of trials decreases per recording session (Figure S6). Nevertheless, specific predominant functional couplings stand out across different epoch segmentation and connectivity measures: FEF has robust functional connectivity with the areas VIP, LIPd, 7AL, and 7PC in the dorsal visual stream in higher frequencies. Whereas IFJa is predominantly coupled with the areas TE1, TE2, TF, and TGv in the ventral visual stream in lower frequencies.

3.3 | Directional interactions between visual streams and plPFC

Directionality analyses in the frequency domain revealed intrinsic unidirectional information flows between seed regions and the visual streams (Figure 3; two-sided Wilcoxon signed-rank test, $p < 0.001$, FDR-corrected). Most remarkably, PDC showed that directional interactions from SPL and IPS to seed regions are significantly more potent than in the other direction. In contrast, directional influences from seed regions to the middle and inferior temporal cortex are more robust than in the other direction, mainly in delta, theta, and beta frequency bands. In other words, the rMEG time series of the regions in plPFC is predicted by the intrinsic neural activity of the regions in the superior parietal cortex (SPC; SPL and IPS) and themselves predictive of the spontaneous activity of the areas in the ventral visual stream. In the alpha band, we instead observed that the seed regions, particularly IFJa and IFJp, predict the intrinsic neural activity in both visual streams except areas 7AL, 7PL, and VIP in the dorsal stream. The direction of information flow from seed regions to these areas located in the medial part of SPC (mSPC) is not significantly stronger than the other direction for any oscillatory band. Likewise, no region in the ventral stream has a directional influence over any seed region, except left TE1m in the gamma band.

3.4 | Directional Interactions Among FEF, IFJa, and IFJp

Finally, the directionality analyses between FEF, IFJa, and IFJp indicated a direction of information flow from IFJp to the other two seed regions (Figure 4). The PDC values from IFJp to IFJa and IFJp to FEF are statistically stronger than in the other directions in both hemispheres. While left IFJp feeds information to left FEF and left IFJa mainly in the theta band, the same direction of interaction in the right hemisphere is driven through both alpha and beta oscillations. Notably, the directed interaction from right IFJp to right FEF in the alpha band is highly significant ($p < 0.00001$; Figure S7). Besides, we observe that right FEF and IFJa have reciprocal connections in various frequency bands. The direction of spontaneous activity from FEF to IFJa and IFJa to FEF are subserved by delta and alpha oscillations, respectively. In the left hemisphere, we only see unidirectional interactions from FEF to IFJa in the delta and gamma band but not vice versa. Noticeably, the delta frequency band underlies FEF's directional influence over IFJa in both hemispheres ($p < 0.005$).

4 | DISCUSSION

The current study investigated the intrinsic functional connectivity profiles of FEF and IFJ, two specific control circuits in the dorsal and ventral PFC, to provide additional evidence for their respective top-down functions in spatial versus non-spatial (feature and object-based) attention

FIGURE 3 Partial directed coherence between FEF, IFJ, and visual streams. The figure shows the directional influences between seed regions (FEF, IFJa, and FEF) and visual streams. While the green regions receive inputs, the magenta ones send outputs to a seed region (two-sided Wilcoxon signed-rank test, p < 0.001, FDR-corrected for 33+3 ROIs). The results shown here are based on 2 s epoch segmentation.

and working memory control. Our results demonstrated that FEF has a more robust power and phase coupling with the spatiotopically organized regions in the dorsal visual stream. In contrast, IFJa has more substantial power and phase relation with the feature or object encoding regions in the ventral visual pathway (Figure 1). These results suggest that in humans the 'where' and 'what' visual streams (Goodale & Milner, 1992; Mishkin et al., 1983) might extend all the way into pIPFC (Goldman-Rakic, 1996; Macko et al., 1982; Wilson et al., 1993). The spatial versus non-spatial to-be-attended or to-be-remembered stimulus domain might respectively segregate pIPFC into a functionally specific dorsal versus ventral visual pathways.

4.1 | FEF vs. IFJa

FEF has robust functional connectivity with the areas VIP, LIPd, 7AL, and 7PC across different epoch lengths and connectivity metrics in the dorsal stream. In this respect, studies on monkeys reported that area VIP is involved in the analysis of visual motion (Colby et al., 1993; Duhamel et al., 1998), multisensory space representation (Schlack et al., 2005), visuomotor control (Culham et al., 2006), self-motion (Bremmer, 2005; Chen et al., 2011; Schlack et al., 2002) and motion encoding in near-extrapersonal space (Schlack et al., 2003). While area LIP is also to some degree selective to shape (Sereno & Maunsell, 1998), researchers at large showed its role in spatial attention (Bisley & Goldberg, 2003; Thakral & Slotnick, 2009), the choice of saccade target (Andersen et al., 1990; Barash et al., 1991; Blatt et al., 1990; Kusunoki et al., 2000; Kusunoki & Goldberg, 2003) and the creation of priority maps (Gottlieb, 2007; Ipata et al., 2009; Sprague & Serences, 2013). Finally, it is

FIGURE 4 Partial directed coherence between the seed regions within PFC. The figure shows the directional influences among FEF, IFJa, and IFJp in both hemispheres. The width of the arrows reflects z-scores on top of them (two-sided Wilcoxon signed-rank test, FDR-corrected for 33+3 ROIs; see Figure 4). The distribution of subjects' PDC value for each significant directional influence is illustrated in Figures S7 and S8.

evident that SPL (areas 7AL and 7PC) takes part in shifting spatial attention (Behrmann et al., 2004; Caspari et al., 2018; Corbetta et al., 1995; Kelley et al., 2008; Molenberghs et al., 2007; Spadone et al., 2021; Vandenberghe et al., 2001; Yantis et al., 2002) and encoding spatial coordinates for planned reach movements (Baldauf et al., 2008; Connolly et al., 2003). Besides, these results match with FEF's deterministic tractography (Baker et al., 2018). FEF's white matter connections accordingly terminate at the intraparietal sulcus (IP1 and IP2), nearby SPL. Overall, the spontaneous connectivity between FEF and SPC (SPL and IPS) corresponds to their underlying structural connectivity and functional specialization in spatial attention and working memory.

In contrast to FEF, IFJa has a stronger functional coupling with the areas TE1a, TE2a, TF, and TGv in the ventral visual stream. These areas roughly correspond to the anterior middle temporal gyrus (aMTG; area TE1a), inferotemporal cortex (IT; areas TE2a and TF) and temporal pole (TP; area TGv). Notably, studies associated aMTG with social cognition (Xu et al., 2020), language, and semantic processing (Ferreira et al., 2015; Gorno-Tempini & Price, 2001; Rossell et al., 2001; Siuda-Krzywicka et al., 2021; Walker et al., 2011). On this note, this particular functional coupling with the anterior temporal cortex might be due to mind-wandering during the resting-state scan (Chou et al., 2017). In comparison, lesions and neuroimaging studies suggest that TP takes part in face recognition by associating person-specific stored memories to the perceptual representation of people's faces (DeVries & Baldauf, 2019; Kamps et al., 2019; Landi et al., 2021; Olson et al., 2007). Additionally, a relational processing task adapted from Smith et al. (2007) revealed that, in contrast to TGd, TGv is activated vs deactivated in relational primary contrasts when subjects were asked to differentiate objects based on their shapes or textures (Glasser et al., 2016). Lastly, the IT cortex has a direct role in object recognition (Kreiman et al., 2006; Lehky & Tanaka, 2016; Tanaka, 1996; Yamane et al., 2008). Studies on monkeys and humans both show IT cortex's involvement in object-based attention and working memory (Baldauf & Desimone, 2014; Kar et al., 2019; Miller et al., 1996). Furthermore, IFJa, but not FEF, has distinctive white matter connectivity with the temporal cortex, which is in accordance with its predominant functional coupling reported here. IFJa's deterministic tractography demonstrates its structural connectivity to the middle and temporal gyrus (Baker et al., 2018). Also, IFJ's probabilistic DWI tractography is especially well connected with several functional areas in IT cortex, which was not the case for FEF (Baldauf & Desimone, 2014). All in all, in light of previous evidence, we indicate that FEF and IFJa's respective predominant functional coupling with SPC and the temporal cortex in the resting state is congruent with their respective functional segregations in plPFC.

4.2 | Frequency-specific interactions

Different oscillations are associated with different cognitive functions (Başar et al., 2001; Güntekin & Başar, 2016; Harmony et al., 1996). Thanks to the high temporal and spatial resolution of MEG, we could reveal the frequency-specific signatures of the functional coupling between FEF, IFJ, and visual pathways (Figure 2). While these frequency-specific interactions in resting state data hint at general communication patterns between the involved areas, it is important to bear in mind that those interactions can rapidly change either in strength, dominant frequency band, and/or directionality as specific attentional or working memory tasks evolve.

In rest, we found substantial delta oscillatory coupling between IFJ and the areas in the ventral visual stream. In this respect, Harmony and colleagues (1996) observed that attention to internal representations leads to increased delta power during mental processing, such as the Sternberg paradigm that measures working memory performance. They suggested that delta oscillations carry attentional modulations from the frontal cortex to distant neuronal networks (Harmony, 2013). Such that it might have a role in suppressing internal or external distractors that interfere with the ongoing mental state. Noticeably, these statements align with IFJ's role in cognitive control (Brass et al., 2005; Derrfuss et al., 2005) and selective attention (Baldauf & Desimone, 2014; Bedini & Baldauf, 2021). Also, for example, Zarjam et al. (2011) could successfully classify working memory load levels based on the delta activity in the frontal EEG. Moreover, they linked delta synchronization to subjects' internal concentration. Since functional coupling between distant regions requires a slower 'clock' (Jensen & Colgin, 2007), delta oscillations might subserve the intrinsic functional connectivity between IFJ and the areas in the ventral visual stream in the absence of attention to features or objects. This baseline information flow can potentially pave the way for task-evoked activity (Cole et al., 2016) during selective attention and working memory.

In the theta band, we did not find any significant region that predominantly power correlates with FEF or IFJ. Additionally, phase-based measures did not reveal any consistent results. The lack of predominant coupling in the theta band might be because theta oscillations are instead linked to learning and memory through cortico-hippocampal interactions (Başar et al., 2001; Buzsáki, 2002; Goutagny et al., 2009; Herweg et al., 2020; Lega et al., 2012; Lubenov & Siapas, 2009; O'Neill et al., 2013; Siapas et al., 2005). However it is important to note that several task-based studies linked theta oscillation to visual working memory (Liebe et al., 2012; Muthukrishnan et al., 2020; Sauseng et al., 2010) and willed attention (Rajan et al., 2019).

On the other side, we found predominant functional connectivity between FEF and SPC (areas VIP, LIPd, 7AL, and 7PC) in the alpha band. Some of this alpha band activity in the dorsal connections, might be attributed to general arousal (Barry et al., 2020; Brancaccio et al., 2020; Cantero et al., 1999; Pivik & Harman, 1995) or visual mental imagery (Xie et al., 2020) during resting state. However, alpha oscillations are also known as neural signature in task-based attention studies (Babiloni et al., 2006; Bae & Luck, 2019; Bagherzadeh et al., 2020; Capotosto et al., 2009; DeVries et al., 2021, 2019; Geng, 2014; Klimesch et al., 2007; Mazaheri et al., 2011; Noah et al., 2020; Rihs et al., 2007; Sauseng et al., 2005; Thut et al., 2006; Toscani et al., 2010; VanDiepen et al., 2016).

Beta synchronization, which we found more strongly involved in the dorsal connections of FEF, is associated with motor inhibition (Gilbertson et al., 2005; Picazio et al., 2014) and termination (Heinrichs-Graham et al., 2017; Kilavik et al., 2013; Pfurtscheller et al., 1996). In contrast, its desynchronization is linked to movement preparation and execution (Kühn et al., 2004; Pfurtscheller et al., 1997; Pfurtscheller & Lopes da Silva, 1999; Y. Zhang et al., 2008). Even so, the entrainment of beta-band activity through transcranial alternating-current stimulation could slow down voluntary movements in healthy subjects (Pogosyan et al., 2009). In this respect, Engel and Fries (2010) suggested that beta oscillations signal the status quo, i.e., the maintenance of the current sensorimotor set. They claimed that the intention or prediction to hold on to the present state leads to robust beta band coupling. Accordingly, we reason that the robust beta oscillatory coupling between FEF and SPC stems in part from the suppression of eye movements during fixation on the crosshair. It might reflect the inhibition of the oculomotion, sustained attention, and the status quo during resting state. Thereby, movement or visuomovement cells in FEF (Gregoriou et al., 2012) likely drive FEF's intrinsic beta synchronization with SPC. Moreover, the deficits in motor response inhibition are linked to attention deficit hyperactivity disorder (ADHD) (Mostofsky et al., 2001). Compared to control subjects, people with ADHD perform particularly poorly in anti-saccade tasks that require suppressing reflexive eye movements. They also have difficulties maintaining steady fixations for a prolonged time (Munoz et al., 2003; Munoz & Everling, 2004). Consequently, it has been argued that the disruption in FEF and SPC's functional coupling in the beta band might indicate a specific deficit in endogenous visuospatial attention (Caldani et al., 2020).

Engel and Fries (2010) proposed that while beta-band activity signals the maintenance of the status quo, gamma-band oscillations indicate the readiness to change it. In line with this statement, several studies reported task-induced gamma synchrony in object perception, attention, and working memory (Baldauf & Desimone, 2014; Fries, 2005, 2009, 2015; Gregoriou et al., 2009; Howard et al., 2003; Tallon-Baudry et al., 1998; Tallon-Baudry & Bertrand, 1999). Despite its specific role in the task-based activity, all three connectivity measures of our present analysis revealed spontaneous functional connectivity between IFJ and the areas in the ventral visual stream in the gamma band. We suggest that, in particular, IFJ and IT cortex's intrinsic gamma coupling hints at their task-evoked functional connectivity during object-based attention (Baldauf & Desimone, 2014). However, it is also likely that this spontaneous synchronization between IFJ and the ventral stream is driven by mind-wandering or imagery in the scanner (Albers et al., 2013; de Borst et al., 2012; Dentico et al., 2014; Dijkstra et al., 2017, 2018; Pearson, 2019). Finally, the predominant functional coupling between IFJ and the ventral visual stream in the rest might to some extent also be driven by

cross-frequency coupling of delta and gamma rhythms (Bruns & Eckhorn, 2004; Canolty & Knight, 2010).

4.3 | Directionality of interactions

In addition to bidirectional functional coupling, we investigated unidirectional interactions between plPFC and visual streams in the resting state. The majority of our regions of interest in Table 1 exhibited a clear directionality to the respectively connected seed region in the lower frequencies. Remarkably, partial directed coherence revealed directed interactions from mSPC (areas 7AL, 7PL, and VIP) to FEF and IFJ in delta and theta oscillations (Figure 3). Since in the delta and theta range the neural time series of mSPC are statistically more successful in predicting the neural activity in FEF and IFJ than vice versa, we can suggest that SPC drives the spontaneous activity in plPFC during rest. This unidirectional connectivity during rest might reflect similar task-related influences of SPC over plPFC during attention and working memory (Chen et al., 2020; Spadone et al., 2021). Additionally, we observe a frequency-specific direction of interaction in the beta band from parietal areas VIP and 7PL to FEF. A recent study by Spadone et al. (2021) similarly indicated that contrasting stay with shift cues reveals a robust direction of information flow from the right ventral IPS to the right dorsal FEF, specifically in the beta band. Since subjects had to fixate during rMEG, the maintenance of attention and the status quo of beta oscillations could have led to the results we observed in the current study. The feedback influences of IPS onto FEF in the beta band might signal to sustain eye movements on the crosshair. Finding such directionality from the parietal cortex to FEF in the beta band is somewhat surprising and speaks against the general notion that the beta band solely provides top-down signals. For these reasons, it is important to note that the direction of an oscillatory coupling is often task-dependent. Therefore, the direction of interaction between SPC and FEF might change depending on the current task demands. While SPC's highly significant directional influence over plPFC during rest suggests that it might provide some sort of spatial attention signals to FEF, Spadone et al. (2021) also showed that the contrast between shift versus stay cues reveals feedback connections from the right FEF to the right SPL. In this respect, the functional hierarchy is not fixed but dynamic. Additionally, the directionality measures should be taken cautiously in general since unidirectional interactions can be observed also because of the asymmetries in the noise levels (Nolte et al., 2004).

4.4 | IFJa vs. IFJp

The functional connectivity analyses (FEF-IFJa and FEF-IFJp) did not reveal identical predominant intrinsic connectivities for IFJa and IFJp (Figures 1 and S2). In particular, IFJa has generally a stronger functional coupling with the ventral visual stream than IFJp. Since these areas have different intrinsic connectivity fingerprints, we suggest that they are functionally distinct regions (Passingham et al., 2002), as already emphasized in MMP1 (Glasser et al., 2016) and previous studies. For instance, Zanto et al. (2011) pointed out that the right IFJ's dorsal and ventral parts are respectively involved in motion and color processing (see also Schwedhelm et al., 2020). Moreover, Asplund et al. (2010) proposed that IFJ might have a role in both stimulus-driven and goal-directed attention through its participation in the ventral and dorsal attention networks. Furthermore, the deterministic tractography revealed distinct white matter tracts for IFJa and IFJp. IFJa had more robust structural connectivity with the middle and inferior temporal gyrus (areas TE1a, TE1m, and TE2a) compared to IFJp (Baker et al., 2018). Finally, the distinct predominant connectivities of IFJa and IFJp provides strong evidence against the notion that our connectivity results could potentially suffer from a proximity bias (since IFJa is more distant from the visual cortex than IFJp).

We found a highly significant direction of interaction from IFJp to FEF and IFJa in both hemispheres (Figure 4). Within the right hemisphere, alpha and beta oscillations govern these intrinsic directional influences from IFJp to the other seed regions. In the left hemisphere, the same direction of information flow is supported also by theta, in addition to alpha and beta bands. Although the direction of interaction may change depending on task-related activity, it is clear that IFJp feeds information to FEF and IFJa in the resting state. Taken together with the previously mentioned result that IFJp has overall less functional connectivity with the ventral visual cortex, this may hint at its role in providing precursor activity to the output units in FEF and IFJa, which then channel that information to visual cortex. Interestingly, also in non-human primate recordings similar interactions have been observed between VPA (the non-human homologue of IFJ) and FEF in visual search tasks: signals from VPA to FEF helped guiding eye movements to likely targets, and the inactivation of VPA selectively eliminated feature attention signals in FEF (while leaving spatial attention and eye movement control unaffected), such that the animals could no longer find targets during visual search (Bichot et al., 2019).

4.5 | Power- and phase-based connectivity

Power- and phase-coupling potentially reflect partly distinct neuronal mechanisms (Daffertshofer et al., 2018; Siems & Siegel, 2020). While power-based metrics compute functional connectivity between two neuronal oscillations based on their amplitudes' envelope correlation, phased-based measures instead consider the consistent relative phase difference between them (Siegel et al., 2012). In this sense, phase-based measures such as iCOH or dwPLI are more delicate and more conservative while attributing a genuine functional coupling in comparison to power-based metrics to avoid Type I errors (Nolte et al., 2004; Stam et al., 2007; Vinck et al., 2011). However, they both have also been found to be less replicable on the group level and less consistent within and between subjects in resting state data (Colclough et al., 2016). On the other hand, the orthogonalized power envelope correlations (Brookes et al., 2011; Hipp et al., 2012) are shown to be the most replicable methods among several alternative metrics, and, importantly, they are insensitive to spurious coupling stemming from field-spread (Colclough et al., 2016; Duan et al., 2021). In light of these findings, we state that the robustness of power- and phase-based measures in the current study relates to the nature of these metrics' computation. Compared to power correlation, the less number of significant phase-couplings for iCOH and dwPLI might be due to their detailed measurement. Meanwhile, the robust connectivity results for oPEC seem to be more reproducible and generalizable across the population. Our successful replication of the results in Hipp et al. (2012) is an additional proof in this respect.

4 | CONCLUSION

We found that the areas in the dorsal visual stream have a more robust functional coupling with FEF, particularly in the beta oscillations. In contrast, the regions in the ventral visual stream have statistically greater functional connectivity with IFJa, particularly in delta and gamma bands. In light of these and previous findings (Cohen et al., 2008; Cole et al., 2014; Fox et al., 2006; Glasser et al., 2016; Mars et al., 2011; Power et al., 2014; Smith et al., 2009; Tavor et al., 2016; Yeo et al., 2011), we argue that intrinsic connectivity patterns are congruent with each brain region's function. Therefore, we conclude that FEF's functional connectivity with the spatiotopically organized regions in SPL and IPS suggests its role in spatial attention and working memory. In contrast, IFJa's functional coupling with feature- or object-encoding areas in the temporal cortex indicates its role in non-spatial attention and working memory. Our functional connectivity data therefore provides strong evidence that also the prefrontal cortex is organized in a dorsal and ventral stream (Goldman-Rakic, 1996).

The resting-state neuroimaging data is time- and cost-efficient since it eases the acquisition of large datasets and enables analyses of multiple hypotheses over task-free recordings. Nevertheless, it is vital to support the findings from resting-state recordings with task-based settings. To this end, the current study attempted to provide the first direct contrast between the functional connectivity profiles of FEF and IFJ to prompt further research on these regions' functional integration with the rest of the brain. Future studies should directly compare the functional connectivity fingerprints of these regions in task designs as in Meyyappan et al. (2021) to provide a more complete picture. At the same time, it is also crucial to reveal their underlying structural connectivity patterns, ideally by using probabilistic rather than deterministic tractography since the latter underestimates the existing long-range white matter tracts. Overall, the direct contrast between the intrinsic, functional, and anatomical 'connectional fingerprints' (Passingham et al., 2002) of FEF and IFJ would provide further evidence for their functional integration with other cortical regions and a dorsal versus ventral functional segregation within plPFC as an extension of 'where' and 'what' visual pathways.

ACKNOWLEDGMENTS

This research is supported by Ministero degli Affari Esteri e della Cooperazione Internazionale (to O.S.) and Fondazione Cassa Di Risparmio Di Trento E Rovereto (to D.B).

CONFLICT OF INTEREST

The authors report no conflict of interest.

DATA AVAILABILITY STATEMENT

The data used in the current study provided and publicly made available by the 1200 Subjects Release of Human Connectome Project (Larson-Prior et al., 2013; Van Essen et al., 2013) in ConnectomeDB (db.humanconnectome.org; Hodge et al., 2016).

ORCID

Orhan Soyuhos https://orcid.org/0000-0002-2858-3979

Daniel Baldauf https://orcid.org/0000-0001-5764-506X

REFERENCES

Albers, A. M., Kok, P., Toni, I., Dijkerman, H. C., & de Lange, F. P. (2013). Shared Representations for Working Memory and Mental Imagery in Early Visual Cortex. *Current Biology*, *23*(15), 1427–1431. https://doi.org/10.1016/j.cub.2013.05.065

Andersen, R. A., Bracewell, R. M., Barash, S., Gnadt, J. W., & Fogassi, L. (1990). Eye position effects on visual, memory, and saccade-related activity in areas LIP and 7a of macaque. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *10*(4), 1176–1196.

Armstrong, K. M., Chang, M. H., & Moore, T. (2009). Selection and Maintenance of Spatial Information by Frontal Eye Field Neurons. *Journal of Neuroscience*, *29*(50), 15621–15629. https://doi.org/10.1523/JNEUROSCI.4465-09.2009

Asplund, C. L., Todd, J. J., Snyder, A. P., & Marois, R. (2010). A central role for the lateral prefrontal cortex in goal-directed and stimulus-driven attention. *Nature Neuroscience*, *13*(4), 507–512. https://doi.org/10.1038/nn.2509

Babiloni, C., Vecchio, F., Bultrini, A., Luca Romani, G., & Rossini, P. M. (2006). Pre- and Poststimulus Alpha Rhythms Are Related to Conscious Visual Perception: A High-Resolution EEG Study. *Cerebral Cortex*, *16*(12), 1690–1700. https://doi.org/10.1093/cercor/bhj104

Baccalá, L. A., & Sameshima, K. (2001). Partial directed coherence: A new concept in neural structure determination. *Biological Cybernetics*, *84*(6), 463–474. https://doi.org/10.1007/PL00007990

Bae, G.-Y., & Luck, S. J. (2019). Decoding motion direction using the topography of sustained ERPs and alpha oscillations. *NeuroImage*, *184*, 242–255. https://doi.org/10.1016/j.neuroimage.2018.09.029

Bagherzadeh, Y., Baldauf, D., Pantazis, D., & Desimone, R. (2020). Alpha Synchrony and the Neurofeedback Control of Spatial Attention. *Neuron*, *105*(3), 577-587.e5. https://doi.org/10.1016/j.neuron.2019.11.001

Baillet, S., Mosher, J. C., & Leahy, R. M. (2001). Electromagnetic brain mapping. *IEEE Signal Processing Magazine*, *18*(6), 14–30. https://doi.org/10.1109/79.962275

Baker, C. M., Burks, J. D., Briggs, R. G., Conner, A. K., Glenn, C. A., Sali, G., McCoy, T. M., Battiste, J. D., O'Donoghue, D. L., & Sughrue, M. E. (2018). A Connectomic Atlas of the Human Cerebrum-Chapter 1: Introduction, Methods, and Significance. *Operative Neurosurgery (Hagerstown, Md.)*, *15*(suppl_1), S1–S9. https://doi.org/10.1093/ons/opy253

Baldauf, D., Cui, H., & Andersen, R. A. (2008). The Posterior Parietal Cortex Encodes in Parallel Both Goals for Double-Reach Sequences. *Journal of Neuroscience*, *28*(40), 10081–10089. https://doi.org/10.1523/JNEUROSCI.3423-08.2008

Baldauf, D., & Desimone, R. (2014). Neural mechanisms of object-based attention. *Science (New York, N.Y.)*, *344*(6182), 424–427. https://doi.org/10.1126/science.1247003

Baldauf, D., & Deubel, H. (2010). Attentional landscapes in reaching and grasping. *Vision Research*, *50*(11), 999–1013. https://doi.org/10.1016/j.visres.2010.02.008

Barash, S., Bracewell, R. M., Fogassi, L., Gnadt, J. W., & Andersen, R. A. (1991). Saccade-related activity in the lateral intraparietal area. I. Temporal properties; comparison with area 7a. *Journal of Neurophysiology*, *66*(3), 1095–1108. https://doi.org/10.1152/jn.1991.66.3.1095

Barry, R. J., De Blasio, F. M., Fogarty, J. S., & Clarke, A. R. (2020). Natural alpha frequency components in resting EEG and their relation to arousal. *Clinical Neurophysiology*, *131*(1), 205–212. https://doi.org/10.1016/j.clinph.2019.10.018

Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and theta oscillations govern cognitive processes. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, *39*(2–3), 241–248. https://doi.org/10.1016/s0167-8760(00)00145-8

Bastos, A. M., & Schoffelen, J.-M. (2016). A Tutorial Review of Functional Connectivity Analysis Methods and Their Interpretational Pitfalls. *Frontiers in Systems Neuroscience*, *9*. https://doi.org/10.3389/fnsys.2015.00175

Bastos, A. M., Vezoli, J., Bosman, C. A., Schoffelen, J.-M., Oostenveld, R., Dowdall, J. R., De Weerd, P., Kennedy, H., & Fries, P. (2015). Visual Areas Exert Feedforward and Feedback Influences through Distinct Frequency Channels. *Neuron*, *85*(2), 390–401. https://doi.org/10.1016/j.neuron.2014.12.018

Bechtold, B. Violin plots for Matlab, Github Project. https://doi.org/10.5281/zenodo.4559847 (2016).

Bedini, M., & Baldauf, D. (2021). Structure, function and connectivity fingerprints of the frontal eye field versus the inferior frontal junction: A comprehensive comparison. *European Journal of Neuroscience*, *54*(4), 5462–5506. https://doi.org/10.1111/ejn.15393

Bedini, M., Olivetti, E., Avesani, P., & Baldauf, D. (2021). The anatomical pathways underlying spatial versus non-spatial attention. *Cognitive Processing, 22 (1),* 35-36.

Behrmann, M., Geng, J. J., & Shomstein, S. (2004). Parietal cortex and attention. *Current Opinion in Neurobiology*, *14*(2), 212–217. https://doi.org/10.1016/j.conb.2004.03.012

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, *57*(1), 289–300.

Bichot, N. P., Rossi, A. F., & Desimone, R. (2005). Parallel and Serial Neural Mechanisms for Visual Search in Macaque Area V4. *Science*. https://doi.org/10.1126/science.1109676

Bichot, N. P., Xu, R., Ghadooshahy, A., Williams, M. L., & Desimone, R. (2019). The role of prefrontal cortex in the control of feature attention in area V4. *Nature Communications*, *10*(1), 5727. https://doi.org/10.1038/s41467-019-13761-7

Bisley, J. W., & Goldberg, M. E. (2003). Neuronal Activity in the Lateral Intraparietal Area and Spatial Attention. *Science*, *299*(5603), 81–86. https://doi.org/10.1126/science.1077395

Blatt, G. J., Andersen, R. A., & Stoner, G. R. (1990). Visual receptive field organization and cortico-cortical connections of the lateral intraparietal area (area LIP) in the macaque. *The Journal of Comparative Neurology*, *299*(4), 421–445. https://doi.org/10.1002/cne.902990404

Brancaccio, A., Tabarelli, D., Bigica, M., & Baldauf, D. (2020). Cortical source localization of sleep-stage specific oscillatory activity. *Scientific Reports*, *10*(1), 6976. https://doi.org/10.1038/s41598-020-63933-5

Brass, M., Derrfuss, J., Forstmann, B., & Cramon, D. Y. von. (2005). The role of the inferior frontal junction area in cognitive control. *Trends in Cognitive Sciences*, *9*(7), 314–316. https://doi.org/10.1016/j.tics.2005.05.001

Bremmer, F. (2005). Navigation in space – the role of the macaque ventral intraparietal area. *The Journal of Physiology*, *566*(1), 29–35. https://doi.org/10.1113/jphysiol.2005.082552

Bressler, S. L., Tang, W., Sylvester, C. M., Shulman, G. L., & Corbetta, M. (2008). Top-Down Control of Human Visual Cortex by Frontal and Parietal Cortex in Anticipatory Visual Spatial Attention. *Journal of Neuroscience*, *28*(40), 10056–10061. https://doi.org/10.1523/JNEUROSCI.1776-08.2008

Brookes, M. J., Hale, J. R., Zumer, J. M., Stevenson, C. M., Francis, S. T., Barnes, G. R., Owen, J. P., Morris, P. G., & Nagarajan, S. S. (2011). Measuring functional connectivity using MEG: Methodology and comparison with fcMRI. *NeuroImage*, *56*(3), 1082–1104. https://doi.org/10.1016/j.neuroimage.2011.02.054

Bruns, A., & Eckhorn, R. (2004). Task-related coupling from high- to low-frequency signals among visual cortical areas in human subdural recordings. *International Journal of Psychophysiology*, *51*(2), 97–116. https://doi.org/10.1016/j.ijpsycho.2003.07.001

Buzsáki, G. (2002). Theta Oscillations in the Hippocampus. *Neuron*, *33*(3), 325–340. https://doi.org/10.1016/S0896-6273(02)00586-X

Caldani, S., Isel, F., Septier, M., Acquaviva, E., Delorme, R., & Bucci, M. P. (2020). Impairment in Attention Focus During the Posner Cognitive Task in Children With ADHD: An Eye Tracker Study. *Frontiers in Pediatrics*, *8*. https://doi.org/10.3389/fped.2020.00484

Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in Cognitive Sciences*, *14*(11), 506–515. https://doi.org/10.1016/j.tics.2010.09.001

Cantero, J. L., Atienza, M., Salas, R. M., & Gómez, C. M. (1999). Alpha EEG coherence in different brain states: An electrophysiological index of the arousal level in human subjects. *Neuroscience Letters*, *271*(3), 167–170. https://doi.org/10.1016/S0304-3940(99)00565-0

Capotosto, P., Babiloni, C., Romani, G. L., & Corbetta, M. (2009). Frontoparietal Cortex Controls Spatial Attention through Modulation of Anticipatory Alpha Rhythms. Journal of Neuroscience, 29(18), 5863–5872. https://doi.org/10.1523/JNEUROSCI.0539-09.2009

Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, *51*(13), 1484–1525. https://doi.org/10.1016/j.visres.2011.04.012

Caspari, N., Arsenault, J. T., Vandenberghe, R., & Vanduffel, W. (2018). Functional Similarity of Medial Superior Parietal Areas for Shift-Selective Attention Signals in Humans and Monkeys. *Cerebral Cortex*, *28*(6), 2085–2099. https://doi.org/10.1093/cercor/bhx114

Cavada, C., & Goldman-Rakic, P. S. (1989). Posterior parietal cortex in rhesus monkey: II. Evidence for segregated corticocortical networks linking sensory and limbic areas with the frontal lobe. *Journal of Comparative Neurology*, *287*(4), 422–445. https://doi.org/10.1002/cne.902870403

Chen, A., DeAngelis, G. C., & Angelaki, D. E. (2011). Representation of Vestibular and Visual Cues to Self-Motion in Ventral Intraparietal Cortex. *Journal of Neuroscience*, *31*(33), 12036–12052. https://doi.org/10.1523/JNEUROSCI.0395-11.2011

Chen, X., Zirnsak, M., Vega, G. M., Govil, E., Lomber, S. G., & Moore, T. (2020). Parietal Cortex Regulates Visual Salience and Salience-Driven Behavior. *Neuron*, *106*(1), 177-187.e4. https://doi.org/10.1016/j.neuron.2020.01.016

Chou, Y., Sundman, M., Whitson, H. E., Gaur, P., Chu, M.-L., Weingarten, C. P., Madden, D. J., Wang, L., Kirste, I., Joliot, M., Diaz, M. T., Li, Y.-J., Song, A. W., & Chen, N. (2017). Maintenance and Representation of Mind Wandering during Resting-State fMRI. *Scientific Reports*, *7*(1), 40722. https://doi.org/10.1038/srep40722

Cohen, A. L., Fair, D. A., Dosenbach, N. U. F., Miezin, F. M., Dierker, D., Van Essen, D. C., Schlaggar, B. L., & Petersen, S. E. (2008). Defining functional areas in individual human brains using resting functional connectivity MRI. *NeuroImage*, *41*(1), 45–57. https://doi.org/10.1016/j.neuroimage.2008.01.066

Colby, C. L., Duhamel, J. R., & Goldberg, M. E. (1993). Ventral intraparietal area of the macaque: Anatomic location and visual response properties. *Journal of Neurophysiology*, *69*(3), 902–914. https://doi.org/10.1152/jn.1993.69.3.902

Colclough, G. L., Woolrich, M. W., Tewarie, P. K., Brookes, M. J., Quinn, A. J., & Smith, S. M. (2016). How reliable are MEG resting-state connectivity metrics? *NeuroImage*, *138*, 284–293. https://doi.org/10.1016/j.neuroimage.2016.05.070

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and Task-Evoked Network Architectures of the Human Brain. *Neuron*, *83*(1), 238–251. https://doi.org/10.1016/j.neuron.2014.05.014

Cole, M. W., Ito, T., Bassett, D. S., & Schultz, D. H. (2016). Activity flow over resting-state networks shapes cognitive task activations. *Nature Neuroscience*, *19*(12), 1718–1726. https://doi.org/10.1038/nn.4406

Connolly, J. D., Andersen, R. A., & Goodale, M. A. (2003). FMRI evidence for a "parietal reach region" in the human brain. *Experimental Brain Research*, *153*(2), 140–145. https://doi.org/10.1007/s00221-003-1587-1

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215. https://doi.org/10.1038/nrn755

Corbetta, M., Shulman, G. L., Miezin, F. M., & Petersen, S. E. (1995). Superior Parietal Cortex Activation During Spatial Attention Shifts and Visual Feature Conjunction. *Science*, *270*(5237), 802–805. https://doi.org/10.1126/science.270.5237.802

Culham, J. C., Cavina-Pratesi, C., & Singhal, A. (2006). The role of parietal cortex in visuomotor control: What have we learned from neuroimaging? *Neuropsychologia*, *44*(13), 2668–2684. https://doi.org/10.1016/j.neuropsychologia.2005.11.003

Daffertshofer, A., Ton, R., Kringelbach, M. L., Woolrich, M., & Deco, G. (2018). Distinct criticality of phase and amplitude dynamics in the resting brain. *NeuroImage*, *180*, 442–447. https://doi.org/10.1016/j.neuroimage.2018.03.002

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage*, *9*(2), 179–194. https://doi.org/10.1006/nimg.1998.0395

Dale, A. M., Liu, A. K., Fischl, B. R., Buckner, R. L., Belliveau, J. W., Lewine, J. D., & Halgren, E. (2000). Dynamic statistical parametric mapping: Combining fMRI and MEG for high-resolution imaging of cortical activity. *Neuron*, *26*(1), 55–67. https://doi.org/10.1016/s0896-6273(00)81138-1

de Borst, A. W., Sack, A. T., Jansma, B. M., Esposito, F., de Martino, F., Valente, G., Roebroeck, A., di Salle, F., Goebel, R., & Formisano, E. (2012). Integration of "what" and "where" in frontal cortex during visual imagery of scenes. *NeuroImage*, *60*(1), 47–58. https://doi.org/10.1016/j.neuroimage.2011.12.005

de Vries, I. E. J. de, Marinato, G., & Baldauf, D. (2021). Decoding Object-Based Auditory Attention from Source-Reconstructed MEG Alpha Oscillations. *Journal of Neuroscience*, *41*(41), 8603–8617. https://doi.org/10.1523/JNEUROSCI.0583-21.2021

de Vries, I. E. J., Savran, E., van Driel, J., & Olivers, C. N. L. (2019). Oscillatory Mechanisms of Preparing for Visual Distraction. *Journal of Cognitive Neuroscience*, *31*(12), 1873–1894. https://doi.org/10.1162/jocn_a_01460

de Vries, E., & Baldauf, D. (2019). Attentional weighting in the face processing network: a magnetic response image-guided magnetoencephalography study using multiple cyclic entrainments. *Journal of cognitive neuroscience, 31(10),* 1573-1588. https://doi.org/10.1162/jocn_a_01428

Dentico, D., Cheung, B. L., Chang, J.-Y., Guokas, J., Boly, M., Tononi, G., & Van Veen, B. (2014). Reversal of cortical information flow during visual imagery as compared to visual perception. *NeuroImage*, *100*, 237–243. https://doi.org/10.1016/j.neuroimage.2014.05.081

Derrfuss, J., Brass, M., Neumann, J., & von Cramon, D. Y. (2005). Involvement of the inferior frontal junction in cognitive control: Meta-analyses of switching and Stroop studies. *Human Brain Mapping*, *25*(1), 22–34. https://doi.org/10.1002/hbm.20127

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review of Neuroscience*, *18*, 193–222. https://doi.org/10.1146/annurev.ne.18.030195.001205

Dijkstra, N., Bosch, S. E., & Gerven, M. A. J. van. (2017). Vividness of Visual Imagery Depends on the Neural Overlap with Perception in Visual Areas. *Journal of Neuroscience*, *37*(5), 1367–1373. https://doi.org/10.1523/JNEUROSCI.3022-16.2016

Dijkstra, N., Mostert, P., Lange, F. P. de, Bosch, S., & van Gerven, M. A. (2018). Differential temporal dynamics during visual imagery and perception. *ELife*, *7*, e33904. https://doi.org/10.7554/eLife.33904

Donahue, C. J., Glasser, M. F., Preuss, T. M., Rilling, J. K., & Van Essen, D. C. (2018). Quantitative assessment of prefrontal cortex in humans relative to nonhuman primates. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(22), E5183–E5192. https://doi.org/10.1073/pnas.1721653115

Duan, W., Chen, X., Wang, Y.-J., Zhao, W., Yuan, H., & Lei, X. (2021). Reproducibility of power spectrum, functional connectivity and network construction in resting-state EEG. *Journal of Neuroscience Methods*, *348*, 108985. https://doi.org/10.1016/j.jneumeth.2020.108985

Duhamel, J.-R., Colby, C. L., & Goldberg, M. E. (1998). Ventral Intraparietal Area of the Macaque: Congruent Visual and Somatic Response Properties. *Journal of Neurophysiology*, *79*(1), 126–136. https://doi.org/10.1152/jn.1998.79.1.126

Duncan, J. (2001). An adaptive coding model of neural function in prefrontal cortex. *Nature Reviews Neuroscience*, *2*(11), 820–829. https://doi.org/10.1038/35097575

Engel, A. K., & Fries, P. (2010). Beta-band oscillations—Signalling the status quo? *Current Opinion in Neurobiology*, *20*(2), 156–165. https://doi.org/10.1016/j.conb.2010.02.015

Fecteau, J. H., & Munoz, D. P. (2006). Salience, relevance, and firing: A priority map for target selection. *Trends in Cognitive Sciences*, *10*(8), 382–390. https://doi.org/10.1016/j.tics.2006.06.011

Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate cerebral cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, *1*(1), 1–47. https://doi.org/10.1093/cercor/1.1.1-a

Ferreira, R. A., Göbel, S. M., Hymers, M., & Ellis, A. W. (2015). The neural correlates of semantic richness: Evidence from an fMRI study of word learning. *Brain and Language*, *143*, 69–80. https://doi.org/10.1016/j.bandl.2015.02.005

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *NeuroImage*, *9*(2), 195–207. https://doi.org/10.1006/nimg.1998.0396

Fischl, B., Sereno, M. I., Tootell, R. B., & Dale, A. M. (1999). High-resolution intersubject averaging and a coordinate system for the cortical surface. *Human Brain Mapping*, *8*(4), 272–284. https://doi.org/10.1002/(sici)1097-0193(1999)8:4<272::aid-hbm10>3.0.co;2-4

Fox, M. D., Corbetta, M., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2006). Spontaneous neuronal activity distinguishes human dorsal and ventral attention systems. *Proceedings of the National Academy of Sciences*, *103*(26), 10046–10051. https://doi.org/10.1073/pnas.0604187103

Fraschini, M., Demuru, M., Crobe, A., Marrosu, F., Stam, C. J., & Hillebrand, A. (2016). *The effect of epoch length on estimated EEG functional*

connectivity and brain network organisation. *Journal of neural engineering*, *13*(3), 036015. https://doi.org/10.1088/1741-2560/13/3/036015

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480. https://doi.org/10.1016/j.tics.2005.08.011

Fries, P. (2009). Neuronal Gamma-Band Synchronization as a Fundamental Process in Cortical Computation. *Annual Review of Neuroscience*, *32*(1), 209–224. https://doi.org/10.1146/annurev.neuro.051508.135603

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, *88*(1), 220–235. https://doi.org/10.1016/j.neuron.2015.09.034

Fuster, J. M. (2001). The prefrontal cortex--an update: Time is of the essence. *Neuron*, *30*(2), 319–333. https://doi.org/10.1016/s0896-6273(01)00285-9

Geng, J. J. (2014). Attentional mechanisms of distractor suppression. Current Directions in Psychological Science, 23(2), 147–153. https://doi.org/10.1177/0963721414525780

Giesbrecht, B., Woldorff, M. G., Song, A. W., & Mangun, G. R. (2003). Neural mechanisms of top-down control during spatial and feature attention. NeuroImage, 19(3), 496–512. https://doi.org/10.1016/S1053-8119(03)00162-9

Gilbertson, T., Lalo, E., Doyle, L., Lazzaro, V. D., Cioni, B., & Brown, P. (2005). Existing Motor State Is Favored at the Expense of New Movement during 13-35 Hz Oscillatory Synchrony in the Human Corticospinal System. *Journal of Neuroscience*, *25*(34), 7771–7779. https://doi.org/10.1523/JNEUROSCI.1762-05.2005

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178. https://doi.org/10.1038/nature18933

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, *80*, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Goldman-Rakic, P. S. (1988). Topography of cognition: Parallel distributed networks in primate association cortex. *Annual Review of Neuroscience*, *11*, 137–156. https://doi.org/10.1146/annurev.ne.11.030188.001033

Goldman-Rakic, P. S. (1996). The prefrontal landscape: Implications of functional architecture for understanding human mentation and the central executive. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *351*(1346), 1445–1453. https://doi.org/10.1098/rstb.1996.0129

Goodale, M. A., & Milner, A. D. (1992). Separate visual pathways for perception and action. *Trends in Neurosciences*, *15*(1), 20–25. https://doi.org/10.1016/0166-2236(92)90344-8

Gorno-Tempini, M. L., & Price, C. J. (2001). Identification of famous faces and buildings: A functional neuroimaging study of semantically unique items. *Brain*, *124*(10), 2087–2097. https://doi.org/10.1093/brain/124.10.2087

Gottlieb, J. (2007). From Thought to Action: The Parietal Cortex as a Bridge between Perception, Action, and Cognition. *Neuron*, *53*(1), 9–16. https://doi.org/10.1016/j.neuron.2006.12.009

Goutagny, R., Jackson, J., & Williams, S. (2009). Self-generated theta oscillations in the hippocampus. *Nature Neuroscience*, *12*(12), 1491–1493. https://doi.org/10.1038/nn.2440

Gregoriou, G. G., Gotts, S. J., & Desimone, R. (2012). Cell-type-specific synchronization of neural activity in FEF with V4 during attention. *Neuron*, *73*(3), 581–594. https://doi.org/10.1016/j.neuron.2011.12.019

Gregoriou, G. G., Gotts, S. J., Zhou, H., & Desimone, R. (2009). High-Frequency, Long-Range Coupling Between Prefrontal and Visual Cortex During Attention. *Science*. https://doi.org/10.1126/science.1171402

Groppe, D.M. (2021). fdr_bh, MATLAB Central File Exchange. Retrieved from https://www.mathworks.com/matlabcentral/fileexchange/27418-fdr_bh

Gross, J., Schmitz, F., Schnitzler, I., Kessler, K., Shapiro, K., Hommel, B., & Schnitzler, A. (2004). Modulation of long-range neural synchrony reflects temporal limitations of visual attention in humans. *Proceedings of the National Academy of Sciences*, *101*(35), 13050–13055. https://doi.org/10.1073/pnas.0404944101

Güntekin, B., & Başar, E. (2016). Review of evoked and event-related delta responses in the human brain. *International Journal of Psychophysiology*, *103*, 43–52. https://doi.org/10.1016/j.ijpsycho.2015.02.001

Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing. *Frontiers in Integrative Neuroscience*, *7*. https://doi.org/10.3389/fnint.2013.00083

Harmony, T., Fernández, T., Silva, J., Bernal, J., Díaz-Comas, L., Reyes, A., Marosi, E., Rodríguez, M., & Rodríguez, M. (1996). EEG delta activity: An indicator of attention to internal processing during performance of mental tasks. *International Journal of Psychophysiology*, *24*(1), 161–171. https://doi.org/10.1016/S0167-8760(96)00053-0

Heinen, K., Feredoes, E., Ruff, C. C., & Driver, J. (2017). Functional connectivity between prefrontal and parietal cortex drives visuo-spatial attention shifts. *Neuropsychologia*, *99*, 81–91. https://doi.org/10.1016/j.neuropsychologia.2017.02.024

Heinrichs-Graham, E., Kurz, M. J., Gehringer, J. E., & Wilson, T. W. (2017). The functional role of post-movement beta oscillations in motor termination. *Brain Structure and Function*, *222*(7), 3075–3086. https://doi.org/10.1007/s00429-017-1387-1

Herweg, N. A., Solomon, E. A., & Kahana, M. J. (2020). Theta Oscillations in Human Memory. *Trends in Cognitive Sciences*, *24*(3), 208–227. https://doi.org/10.1016/j.tics.2019.12.006

Hincapié, A.-S., Kujala, J., Mattout, J., Pascarella, A., Daligault, S., Delpuech, C., Mery, D., Cosmelli, D., & Jerbi, K. (2017). The impact of MEG source reconstruction method on source-space connectivity estimation: A comparison between minimum-norm solution and

beamforming. *NeuroImage*, *156*, 29–42. https://doi.org/10.1016/j.neuroimage.2017.04.038

Hipp, J. F., Hawellek, D. J., Corbetta, M., Siegel, M., & Engel, A. K. (2012). Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nature Neuroscience*, *15*(6), 884–890. https://doi.org/10.1038/nn.3101

Hodge, M. R., Horton, W., Brown, T., Herrick, R., Olsen, T., Hileman, M. E., McKay, M., Archie, K. A., Cler, E., Harms, M. P., Burgess, G. C., Glasser, M. F., Elam, J. S., Curtiss, S. W., Barch, D. M., Oostenveld, R., Larson-Prior, L. J., Ugurbil, K., Van Essen, D. C., & Marcus, D. S. (2016). ConnectomeDB—Sharing human brain connectivity data. *NeuroImage*, *124*, 1102–1107. https://doi.org/10.1016/j.neuroimage.2015.04.046

Howard, M. W., Rizzuto, D. S., Caplan, J. B., Madsen, J. R., Lisman, J., Aschenbrenner-Scheibe, R., Schulze-Bonhage, A., & Kahana, M. J. (2003). Gamma Oscillations Correlate with Working Memory Load in Humans. *Cerebral Cortex*, *13*(12), 1369–1374. https://doi.org/10.1093/cercor/bhg084

Huang, M. X., Mosher, J. C., & Leahy, R. M. (1999). A sensor-weighted overlapping-sphere head model and exhaustive head model comparison for MEG. *Physics in Medicine and Biology*, *44*(2), 423–440. https://doi.org/10.1088/0031-9155/44/2/010

Ipata, A. E., Gee, A. L., Bisley, J. W., & Goldberg, M. E. (2009). Neurons in the lateral intraparietal area create a priority map by the combination of disparate signals. *Experimental Brain Research*, *192*(3), 479–488. https://doi.org/10.1007/s00221-008-1557-8

Jensen, O., & Colgin, L. L. (2007). Cross-frequency coupling between neuronal oscillations. *Trends in Cognitive Sciences*, *11*(7), 267–269. https://doi.org/10.1016/j.tics.2007.05.003

Jerde, T. A., Merriam, E. P., Riggall, A. C., Hedges, J. H., & Curtis, C. E. (2012). Prioritized Maps of Space in Human Frontoparietal Cortex. *Journal of Neuroscience*, *32*(48), 17382–17390. https://doi.org/10.1523/JNEUROSCI.3810-12.2012

Ji, J. L., Spronk, M., Kulkarni, K., Repovš, G., Anticevic, A., & Cole, M. W. (2019). Mapping the human brain's cortical-subcortical functional network organization. *NeuroImage*, *185*, 35–57. https://doi.org/10.1016/j.neuroimage.2018.10.006

Kamps, F. S., Morris, E. J., & Dilks, D. D. (2019). A face is more than just the eyes, nose, and mouth: fMRI evidence that face-selective cortex represents external features. *Neuroimage, 184*, 90-100. https://doi.org/10.1016/j.neuroimage.2018.09.027

Kar, K., Kubilius, J., Schmidt, K., Issa, E. B., & DiCarlo, J. J. (2019). Evidence that recurrent circuits are critical to the ventral stream's execution of core object recognition behavior. *Nature neuroscience, 22(6)*, 974-983. https://doi.org/10.1038/s41593-019-0392-5

Kassebaum, P. (2021). circularGraph, GitHub repository. Retrieved from https://github.com/paul-kassebaum-mathworks/circularGraph

Kelley, T. A., Serences, J. T., Giesbrecht, B., & Yantis, S. (2008). Cortical Mechanisms for Shifting and Holding Visuospatial Attention. *Cerebral Cortex*, *18*(1), 114–125. https://doi.org/10.1093/cercor/bhm036

Kilavik, B. E., Zaepffel, M., Brovelli, A., MacKay, W. A., & Riehle, A. (2013). The ups and downs of beta oscillations in sensorimotor cortex. *Experimental Neurology*, *245*, 15–26. https://doi.org/10.1016/j.expneurol.2012.09.014

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88. https://doi.org/10.1016/j.brainresrev.2006.06.003

Kravitz, D. J., Saleem, K. S., Baker, C. I., & Mishkin, M. (2011). A new neural framework for visuospatial processing. *Nature Reviews Neuroscience*, *12*(4), 217–230. https://doi.org/10.1038/nrn3008

Kravitz, D. J., Saleem, K. S., Baker, C. I., Ungerleider, L. G., & Mishkin, M. (2013). The ventral visual pathway: An expanded neural framework for the processing of object quality. *Trends in Cognitive Sciences*, *17*(1), 26–49. https://doi.org/10.1016/j.tics.2012.10.011

Kühn, A. A., Williams, D., Kupsch, A., Limousin, P., Hariz, M., Schneider, G.-H., Yarrow, K., & Brown, P. (2004). Event-related beta desynchronization in human subthalamic nucleus correlates with motor performance. *Brain*, *127*(4), 735–746. https://doi.org/10.1093/brain/awh106

Kusunoki, M., & Goldberg, M. E. (2003). The Time Course of Perisaccadic Receptive Field Shifts in the Lateral Intraparietal Area of the Monkey. *Journal of Neurophysiology*, *89*(3), 1519–1527. https://doi.org/10.1152/jn.00519.2002

Kusunoki, M., Gottlieb, J., & Goldberg, M. E. (2000). The lateral intraparietal area as a salience map: The representation of abrupt onset, stimulus motion, and task relevance. *Vision Research*, *40*(10), 1459–1468. https://doi.org/10.1016/S0042-6989(99)00212-6

Kreiman, G., Hung, C. P., Kraskov, A., Quiroga, R. Q., Poggio, T., & DiCarlo, J. J. (2006). Object selectivity of local field potentials and spikes in the macaque inferior temporal cortex. *Neuron, 49(3)*, 433-445. https://doi.org/10.1016/j.neuron.2005.12.019

Landi, S. M., Viswanathan, P., Serene, S., & Freiwald, W. A. (2021). A fast link between face perception and memory in the temporal pole. *Science*, *373*(6554), 581–585. https://doi.org/10.1126/science.abi6671

Larson-Prior, L. J., Oostenveld, R., Della Penna, S., Michalareas, G., Prior, F., Babajani-Feremi, A., Schoffelen, J.-M., Marzetti, L., de Pasquale, F., Di Pompeo, F., Stout, J., Woolrich, M., Luo, Q., Bucholz, R., Fries, P., Pizzella, V., Romani, G. L., Corbetta, M., & Snyder, A. Z. (2013). Adding dynamics to the Human Connectome Project with MEG. *NeuroImage*, *80*, 190–201. https://doi.org/10.1016/j.neuroimage.2013.05.056

Lega, B. C., Jacobs, J., & Kahana, M. (2012). Human hippocampal theta oscillations and the formation of episodic memories. *Hippocampus*, *22*(4), 748–761. https://doi.org/10.1002/hipo.20937

Lehky, S. R., & Tanaka, K. (2016). Neural representation for object recognition in inferotemporal cortex. *Current Opinion in Neurobiology*, *37*, 23–35. https://doi.org/10.1016/j.conb.2015.12.001

Liebe, S., Hoerzer, G. M., Logothetis, N. K., & Rainer, G. (2012). Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nature Neuroscience*, *15*(3), 456–462. https://doi.org/10.1038/nn.3038

Lubenov, E. V., & Siapas, A. G. (2009). Hippocampal theta oscillations are travelling waves. *Nature*, *459*(7246), 534–539. https://doi.org/10.1038/nature08010

Macko, K. A., Jarvis, C. D., Kennedy, C., Miyaoka, M., Shinohara, M., Sololoff, L., & Mishkin, M. (1982). Mapping the Primate Visual System with [2-14C]Deoxyglucose. *Science*, *218*(4570), 394–397. https://doi.org/10.1126/science.7123241

Mangun, G. R. (1995). Neural mechanisms of visual selective attention. Psychophysiology, 32(1), 4–18. https://doi.org/10.1111/j.1469-8986.1995.tb03400.x

Mars, R. B., Jbabdi, S., Sallet, J., O'Reilly, J. X., Croxson, P. L., Olivier, E., Noonan, M. P., Bergmann, C., Mitchell, A. S., Baxter, M. G., Behrens, T. E. J., Johansen-Berg, H., Tomassini, V., Miller, K. L., & Rushworth, M. F. S. (2011). Diffusion-Weighted Imaging Tractography-Based Parcellation of the Human Parietal Cortex and Comparison with Human and Macaque Resting-State Functional Connectivity. *Journal of Neuroscience*, 31(11), 4087–4100. https://doi.org/10.1523/JNEUROSCI.5102-10.2011

Marshall, T. R., O'Shea, J., Jensen, O., & Bergmann, T. O. (2015). Frontal eye fields control attentional modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 35(4), 1638–1647. https://doi.org/10.1523/JNEUROSCI.3116-14.2015

Mazaheri, A., DiQuattro, N. E., Bengson, J., & Geng, J. J. (2011). Pre-Stimulus Activity Predicts the Winner of Top-Down vs. Bottom-Up Attentional Selection. PLOS ONE, 6(2), e16243. https://doi.org/10.1371/journal.pone.0016243

McCarthy, G., Puce, A., Constable, T., Krystal, J. H., Gore, J. C., & Goldman-Rakic, P. (1996). Activation of Human Prefrontal Cortex during Spatial and Nonspatial Working Memory Tasks Measured by Functional MRI. *Cerebral Cortex*, 6(4), 600–611. https://doi.org/10.1093/cercor/6.4.600

Merrikhi, Y., Clark, K., Albarran, E., Parsa, M., Zirnsak, M., Moore, T., & Noudoost, B. (2017). Spatial working memory alters the efficacy of input to visual cortex. *Nature Communications*, 8(1), 15041. https://doi.org/10.1038/ncomms15041

Meyyappan, S., Rajan, A., Mangun, G. R., & Ding, M. (2021). Role of Inferior Frontal Junction (IFJ) in the Control of Feature versus Spatial Attention. *Journal of Neuroscience*, 41(38), 8065–8074. https://doi.org/10.1523/JNEUROSCI.2883-20.2021

Miller, E. K., & Cohen, J. D. (2001). An Integrative Theory of Prefrontal Cortex Function. *Annual Review of Neuroscience*, 24(1), 167–202. https://doi.org/10.1146/annurev.neuro.24.1.167

Miller, E. K., Erickson, C. A., & Desimone, R. (1996). Neural Mechanisms of Visual Working Memory in Prefrontal Cortex of the Macaque. *Journal of Neuroscience*, 16(16), 5154–5167. https://doi.org/10.1523/JNEUROSCI.16-16-05154.1996

Mishkin, M., Ungerleider, L. G., & Macko, K. A. (1983). Object vision and spatial vision: Two cortical pathways. *Trends in Neurosciences*, 6, 414–417. https://doi.org/10.1016/0166-2236(83)90190-X

Molenberghs, P., Mesulam, M. M., Peeters, R., & Vandenberghe, R. R. C. (2007). Remapping Attentional Priorities: Differential Contribution of Superior Parietal Lobule and Intraparietal Sulcus. *Cerebral Cortex*, 17(11), 2703–2712. https://doi.org/10.1093/cercor/bhl179

Moore, T., & Armstrong, K. M. (2003). Selective gating of visual signals by microstimulation of frontal cortex. *Nature*, 421(6921), 370–373. https://doi.org/10.1038/nature01341

Moore, T., & Fallah, M. (2004). Microstimulation of the Frontal Eye Field and Its Effects on Covert Spatial Attention. *Journal of Neurophysiology*, 91(1), 152–162. https://doi.org/10.1152/jn.00741.2002

Mostofsky, S. H., Lasker, A. G., Cutting, L. E., Denckla, M. B., & Zee, D. S. (2001). Oculomotor abnormalities in attention deficit hyperactivity disorder: A preliminary study. *Neurology*, 57(3), 423–430. https://doi.org/10.1212/WNL.57.3.423

Muhle-Karbe, P. S., Andres, M., & Brass, M. (2014). Transcranial Magnetic Stimulation Dissociates Prefrontal and Parietal Contributions to Task Preparation. *Journal of Neuroscience*, 34(37), 12481–12489. https://doi.org/10.1523/JNEUROSCI.4931-13.2014

Munoz, D. P., Armstrong, I. T., Hampton, K. A., & Moore, K. D. (2003). Altered Control of Visual Fixation and Saccadic Eye Movements in Attention-Deficit Hyperactivity Disorder. *Journal of Neurophysiology*, 90(1), 503–514. https://doi.org/10.1152/jn.00192.2003

Munoz, D. P., & Everling, S. (2004). Look away: The anti-saccade task and the voluntary control of eye movement. *Nature Reviews Neuroscience*, 5(3), 218–228. https://doi.org/10.1038/nrn1345

Muthukrishnan, S. P., Soni, S., & Sharma, R. (2020). Brain Networks Communicate Through Theta Oscillations to Encode High Load in a Visuospatial Working Memory Task: An EEG Connectivity Study. *Brain Topography*, 33(1), 75–85. https://doi.org/10.1007/s10548-019-00739-3

Noah, S., Powell, T., Khodayari, N., Olivan, D., Ding, M., & Mangun, G. R. (2020). Neural Mechanisms of Attentional Control for Objects: Decoding EEG Alpha When Anticipating Faces, Scenes, and Tools. Journal of Neuroscience, 40(25), 4913–4924. https://doi.org/10.1523/JNEUROSCI.2685-19.2020

Nolte, G., Bai, O., Wheaton, L., Mari, Z., Vorbach, S., & Hallett, M. (2004). Identifying true brain interaction from EEG data using the imaginary part of coherency. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, 115(10), 2292–2307. https://doi.org/10.1016/j.clinph.2004.04.029

Ó Scalaidhe, S. P., Wilson, F. A. W., & Goldman-Rakic, P. S. (1999). Face-selective Neurons During Passive Viewing and Working Memory Performance of Rhesus Monkeys: Evidence for Intrinsic Specialization of Neuronal Coding. *Cerebral Cortex*, 9(5), 459–475. https://doi.org/10.1093/cercor/9.5.459

Olson, I. R., Plotzker, A., & Ezzyat, Y. (2007). The Enigmatic temporal pole: A review of findings on social and emotional processing. *Brain: A Journal of Neurology*, 130(Pt 7), 1718–1731. https://doi.org/10.1093/brain/awm052

O'Neill, P.-K., Gordon, J. A., & Sigurdsson, T. (2013). Theta Oscillations in the Medial Prefrontal Cortex Are Modulated by Spatial Working Memory and Synchronize with the Hippocampus through Its Ventral Subregion. *Journal of Neuroscience*, 33(35), 14211–14224. https://doi.org/10.1523/JNEUROSCI.2378-13.2013

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010, December 23). *FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data* [Research Article]. Computational Intelligence and Neuroscience; Hindawi. https://doi.org/10.1155/2011/156869

Osher, D. E., Saxe, R. R., Koldewyn, K., Gabrieli, J. D. E., Kanwisher, N., & Saygin, Z. M. (2016). Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple Visual Categories across Cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, 26(4), 1668–1683.

https://doi.org/10.1093/cercor/bhu303

Paneri, S., & Gregoriou, G. G. (2017). Top-Down Control of Visual Attention by the Prefrontal Cortex. Functional Specialization and Long-Range Interactions. *Frontiers in Neuroscience*, *11*, 545. https://doi.org/10.3389/fnins.2017.00545

Passingham, R. E., Stephan, K. E., & Kötter, R. (2002). The anatomical basis of functional localization in the cortex. *Nature Reviews Neuroscience*, *3*(8), 606–616. https://doi.org/10.1038/nrn893

Pearson, J. (2019). The human imagination: The cognitive neuroscience of visual mental imagery. *Nature Reviews Neuroscience*, *20*(10), 624–634. https://doi.org/10.1038/s41583-019-0202-9

Petit, L., & Pouget, P. (2019). The comparative anatomy of frontal eye fields in primates. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *118*, 51–64. https://doi.org/10.1016/j.cortex.2019.02.023

Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG synchronization and desynchronization: Basic principles. *Clinical Neurophysiology*, *110*(11), 1842–1857. https://doi.org/10.1016/S1388-2457(99)00141-8

Pfurtscheller, G., Neuper, C., Andrew, C., & Edlinger, G. (1997). Foot and hand area mu rhythms. *International Journal of Psychophysiology*, *26*(1), 121–135. https://doi.org/10.1016/S0167-8760(97)00760-5

Pfurtscheller, G., Stancák, A., & Neuper, C. (1996). Post-movement beta synchronization. A correlate of an idling motor area? *Electroencephalography and Clinical Neurophysiology*, *98*(4), 281–293. https://doi.org/10.1016/0013-4694(95)00258-8

Picazio, S., Veniero, D., Ponzo, V., Caltagirone, C., Gross, J., Thut, G., & Koch, G. (2014). Prefrontal Control over Motor Cortex Cycles at Beta Frequency during Movement Inhibition. *Current Biology*, *24*(24), 2940–2945. https://doi.org/10.1016/j.cub.2014.10.043

Pivik, R. T., & Harman, K. (1995). A reconceptualization of EEG alpha activity as an index of arousal during sleep: All alpha activity is not equal. *Journal of Sleep Research*, *4*(3), 131–137. https://doi.org/10.1111/j.1365-2869.1995.tb00161.x

Pogosyan, A., Gaynor, L. D., Eusebio, A., & Brown, P. (2009). Boosting Cortical Activity at Beta-Band Frequencies Slows Movement in Humans. *Current Biology*, *19*(19), 1637–1641. https://doi.org/10.1016/j.cub.2009.07.074

Power, J. D., Schlaggar, B. L., & Petersen, S. E. (2014). Studying Brain Organization via Spontaneous fMRI Signal. *Neuron*, *84*(4), 681–696. https://doi.org/10.1016/j.neuron.2014.09.007

Ptak, R. (2012). The Frontoparietal Attention Network of the Human Brain: Action, Saliency, and a Priority Map of the Environment. *The Neuroscientist*, *18*(5), 502–515. https://doi.org/10.1177/1073858411409051

Rajan, A., Siegel, S. N., Liu, Y., Bengson, J., Mangun, G. R., & Ding, M. (2019). Theta Oscillations Index Frontal Decision-Making and Mediate Reciprocal Frontal–Parietal Interactions in Willed Attention. Cerebral Cortex, 29(7), 2832–2843. https://doi.org/10.1093/cercor/bhy149

Rihs, T. A., Michel, C. M., & Thut, G. (2007). Mechanisms of selective inhibition in visual spatial attention are indexed by α-band EEG synchronization. *European Journal of Neuroscience*, *25*(2), 603–610. https://doi.org/10.1111/j.1460-9568.2007.05278.x

Romanski, L. M. (2004). Domain specificity in the primate prefrontal cortex. *Cognitive, Affective, & Behavioral Neuroscience*, *4*(4), 421–429. https://doi.org/10.3758/CABN.4.4.421

Rossell, S. L., Bullmore, E. T., Williams, S. C. R., & David, A. S. (2001). Brain activation during automatic and controlled processing of semantic relations: A priming experiment using lexical-decision. *Neuropsychologia*, *39*(11), 1167–1176. https://doi.org/10.1016/S0028-3932(01)00049-5

Rossi, A. F., Pessoa, L., Desimone, R., & Ungerleider, L. G. (2008). The prefrontal cortex and the executive control of attention. *Experimental Brain Research*, *192*(3), 489. https://doi.org/10.1007/s00221-008-1642-z

Sauseng, P., Griesmayr, B., Freunberger, R., & Klimesch, W. (2010). Control mechanisms in working memory: A possible function of EEG theta oscillations. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1015–1022. https://doi.org/10.1016/j.neubiorev.2009.12.006

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., Gruber, W. R., & Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with human EEG alpha activity. *European Journal of Neuroscience*, *22*(11), 2917–2926. https://doi.org/10.1111/j.1460-9568.2005.04482.x

Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D. E., & Saxe, R. R. (2011). Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nature Neuroscience*, *15*(2), 321–327. https://doi.org/10.1038/nn.3001

Schlack, A., Hoffmann, K.-P., & Bremmer, F. (2002). Interaction of linear vestibular and visual stimulation in the macaque ventral intraparietal area (VIP). *European Journal of Neuroscience*, *16*(10), 1877–1886. https://doi.org/10.1046/j.1460-9568.2002.02251.x

Schlack, A., Hoffmann, K.-P., & Bremmer, F. (2003). Selectivity of macaque ventral intraparietal area (area VIP) for smooth pursuit eye movements. *The Journal of Physiology*, *551*(2), 551–561. https://doi.org/10.1113/jphysiol.2003.042994

Schlack, A., Sterbing-D'Angelo, S. J., Hartung, K., Hoffmann, K.-P., & Bremmer, F. (2005). Multisensory Space Representations in the Macaque Ventral Intraparietal Area. *Journal of Neuroscience*, *25*(18), 4616–4625. https://doi.org/10.1523/JNEUROSCI.0455-05.2005

Schwedhelm, P., Baldauf, D., & Treue, S. (2020). The lateral prefrontal cortex of primates encodes stimulus colors and their behavioral relevance during a match-to-sample task. *Scientific reports, 10(1)*, 1-12. https://doi.org/10.1038/s41598-020-61171-3

Sereno, A. B., & Maunsell, J. H. R. (1998). Shape selectivity in primate lateral intraparietal cortex. *Nature*, *395*(6701), 500–503. https://doi.org/10.1038/26752

Siapas, A. G., Lubenov, E. V., & Wilson, M. A. (2005). Prefrontal Phase Locking to Hippocampal Theta Oscillations. *Neuron*, *46*(1), 141–151. https://doi.org/10.1016/j.neuron.2005.02.028

Siegel, M., Donner, T. H., & Engel, A. K. (2012). Spectral fingerprints of large-scale neuronal interactions. *Nature Reviews Neuroscience*, *13*(2), 121–134. https://doi.org/10.1038/nrn3137

Siegel, M., Donner, T. H., Oostenveld, R., Fries, P., & Engel, A. K. (2008). Neuronal Synchronization along the Dorsal Visual Pathway Reflects the Focus of Spatial Attention. *Neuron*, *60*(4), 709–719. https://doi.org/10.1016/j.neuron.2008.09.010

Siems, M., & Siegel, M. (2020). Dissociated neuronal phase- and amplitude-coupling patterns in the human brain. *NeuroImage*, *209*, 116538. https://doi.org/10.1016/j.neuroimage.2020.116538

Siuda-Krzywicka, K., Witzel, C., Bartolomeo, P., & Cohen, L. (2021). Color Naming and Categorization Depend on Distinct Functional Brain Networks. *Cerebral Cortex*, *31*(2), 1106–1115. https://doi.org/10.1093/cercor/bhaa278

Smith, R., Keramatian, K., & Christoff, K. (2007). Localizing the rostrolateral prefrontal cortex at the individual level. *NeuroImage*, *36*(4), 1387–1396. https://doi.org/10.1016/j.neuroimage.2007.04.032

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N., Watkins, K. E., Toro, R., Laird, A. R., & Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during activation and rest. *Proceedings of the National Academy of Sciences*, *106*(31), 13040–13045. https://doi.org/10.1073/pnas.0905267106

Spadone, S., Wyczesany, M., Della Penna, S., Corbetta, M., & Capotosto, P. (2021). Directed Flow of Beta Band Communication During Reorienting of Attention Within the Dorsal Attention Network. *Brain Connectivity*, *11*(9), 717–724. https://doi.org/10.1089/brain.2020.0885

Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in the human occipital, parietal and frontal cortices. *Nature Neuroscience*, *16*(12), 1879–1887. https://doi.org/10.1038/nn.3574

Stam, C. J., Nolte, G., & Daffertshofer, A. (2007). Phase lag index: Assessment of functional connectivity from multi channel EEG and MEG with diminished bias from common sources. *Human Brain Mapping*, *28*(11), 1178–1193. https://doi.org/10.1002/hbm.20346

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A user-friendly application for MEG/EEG analysis. *Computational Intelligence and Neuroscience*, *2011*, 879716. https://doi.org/10.1155/2011/879716

Tallon-Baudry, C., & Bertrand, O. (1999). Oscillatory gamma activity in humans and its role in object representation. *Trends in Cognitive Sciences*, *3*(4), 151–162. https://doi.org/10.1016/S1364-6613(99)01299-1

Tallon-Baudry, C., Bertrand, O., Peronnet, F., & Pernier, J. (1998). Induced γ-Band Activity during the Delay of a Visual Short-Term Memory Task in Humans. *Journal of Neuroscience*, *18*(11), 4244–4254. https://doi.org/10.1523/JNEUROSCI.18-11-04244.1998

Tanaka, K. (1996). Inferotemporal Cortex and Object Vision. *Annual Review of Neuroscience*, *19*(1), 109–139. https://doi.org/10.1146/annurev.ne.19.030196.000545

Tavor, I., Jones, O. P., Mars, R. B., Smith, S. M., Behrens, T. E., & Jbabdi, S. (2016). Task-free MRI predicts individual differences in brain activity during task performance. *Science*, *352*(6282), 216–220. https://doi.org/10.1126/science.aad8127

Thakral, P. P., & Slotnick, S. D. (2009). The role of parietal cortex during sustained visual spatial attention. *Brain Research*, *1302*, 157–166. https://doi.org/10.1016/j.brainres.2009.09.031

Thompson, K. G., & Bichot, N. P. (2005). A visual salience map in the primate frontal eye field. In *Progress in Brain Research* (Vol. 147, pp. 249–262). Elsevier. https://doi.org/10.1016/S0079-6123(04)47019-8

Thompson, K. G., Biscoe, K. L., & Sato, T. R. (2005). Neuronal Basis of Covert Spatial Attention in the Frontal Eye Field. *Journal of Neuroscience*, *25*(41), 9479–9487. https://doi.org/10.1523/JNEUROSCI.0741-05.2005

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). α-Band Electroencephalographic Activity over Occipital Cortex Indexes Visuospatial Attention Bias and Predicts Visual Target Detection. *Journal of Neuroscience*, *26*(37), 9494–9502. https://doi.org/10.1523/JNEUROSCI.0875-06.2006

Toscani, M., Marzi, T., Righi, S., Viggiano, M. P., & Baldassi, S. (2010). Alpha waves: A neural signature of visual suppression. *Experimental Brain Research*, *207*(3), 213–219. https://doi.org/10.1007/s00221-010-2444-7

Uğurbil, K., Xu, J., Auerbach, E. J., Moeller, S., Vu, A. T., Duarte-Carvajalino, J. M., Lenglet, C., Wu, X., Schmitter, S., Van de Moortele, P. F., Strupp, J., Sapiro, G., De Martino, F., Wang, D., Harel, N., Garwood, M., Chen, L., Feinberg, D. A., Smith, S. M., … Yacoub, E. (2013). Pushing spatial and temporal resolution for functional and diffusion MRI in the Human Connectome Project. *NeuroImage*, *80*, 80–104. https://doi.org/10.1016/j.neuroimage.2013.05.012

van Diepen, R. M., Miller, L. M., Mazaheri, A., & Geng, J. J. (2016). The Role of Alpha Activity in Spatial and Feature-Based Attention. ENeuro, *3*(5), ENEURO.0204-16.2016. https://doi.org/10.1523/ENEURO.0204-16.2016

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., & Ugurbil, K. (2013). The WU-Minn Human Connectome Project: An overview. *NeuroImage*, *80*, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Vandenberghe, R., Gitelman, D. R., Parrish, T. B., & Mesulam, M. M. (2001). Functional Specificity of Superior Parietal Mediation of Spatial Shifting. *NeuroImage*, *14*(3), 661–673. https://doi.org/10.1006/nimg.2001.0860

Veniero, D., Gross, J., Morand, S., Duecker, F., Sack, A. T., & Thut, G. (2021). Top-down control of visual cortex by the frontal eye fields through oscillatory realignment. *Nature Communications*, *12*(1), 1757. https://doi.org/10.1038/s41467-021-21979-7

Vernet, M., Quentin, R., Chanes, L., Mitsumasu, A., & Valero-Cabré, A. (2014). Frontal eye field, where art thou? Anatomy, function, and non-invasive manipulation of frontal regions involved in eye movements and associated cognitive operations. *Frontiers in Integrative Neuroscience*, *8*, 66. https://doi.org/10.3389/fnint.2014.00066

Vinck, M., Oostenveld, R., van Wingerden, M., Battaglia, F., & Pennartz, C. M. A. (2011). An improved index of phase-synchronization for electrophysiological data in the presence of volume-conduction, noise and sample-size bias. *NeuroImage*, *55*(4), 1548–1565. https://doi.org/10.1016/j.neuroimage.2011.01.055

Walker, G. M., Schwartz, M. F., Kimberg, D. Y., Faseyitan, O., Brecher, A., Dell, G. S., & Coslett, H. B. (2011). Support for anterior temporal involvement in semantic error production in aphasia: New evidence from VLSM. *Brain and Language*, *117*(3), 110–122. https://doi.org/10.1016/j.bandl.2010.09.008

Wang, L., Mruczek, R. E. B., Arcaro, M. J., & Kastner, S. (2015). Probabilistic Maps of Visual Topography in Human Cortex. *Cerebral Cortex*, *25*(10), 3911–3931. https://doi.org/10.1093/cercor/bhu277

Wardak, C., Ibos, G., Duhamel, J.-R., & Olivier, E. (2006). Contribution of the Monkey Frontal Eye Field to Covert Visual Attention. *Journal of Neuroscience*, *26*(16), 4228–4235. https://doi.org/10.1523/JNEUROSCI.3336-05.2006

Wilson, F. A. W., Scalaidhe, S. P. Ó., & Goldman-Rakic, P. S. (1993). Dissociation of Object and Spatial Processing Domains in Primate Prefrontal Cortex. *Science*, *260*(5116), 1955–1958. https://doi.org/10.1126/science.8316836

Xie, S., Kaiser, D., & Cichy, R. M. (2020). Visual Imagery and Perception Share Neural Representations in the Alpha Frequency Band. *Current Biology*, *30*(13), 2621-2627.e5. https://doi.org/10.1016/j.cub.2020.04.074

Xu, J., Wang, C., Xu, Z., Li, T., Chen, F., Chen, K., Gao, J., Wang, J., & Hu, Q. (2020). Specific Functional Connectivity Patterns of Middle Temporal Gyrus Subregions in Children and Adults with Autism Spectrum Disorder. *Autism Research*, *13*(3), 410–422. https://doi.org/10.1002/aur.2239

Yamane, Y., Carlson, E. T., Bowman, K. C., Wang, Z., & Connor, C. E. (2008). A neural code for three-dimensional object shape in macaque inferotemporal cortex. *Nature Neuroscience*, *11*(11), 1352–1360. https://doi.org/10.1038/nn.2202

Yantis, S., Schwarzbach, J., Serences, J. T., Carlson, R. L., Steinmetz, M. A., Pekar, J. J., & Courtney, S. M. (2002). Transient neural activity in human parietal cortex during spatial attention shifts. *Nature Neuroscience*, *5*(10), 995–1002. https://doi.org/10.1038/nn921

Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165. https://doi.org/10.1152/jn.00338.2011

Zanto, T. P., Rubens, M. T., Bollinger, J., & Gazzaley, A. (2010). Top-down modulation of visual feature processing: The role of the inferior frontal junction. *NeuroImage*, *53*(2), 736–745. https://doi.org/10.1016/j.neuroimage.2010.06.012

Zanto, T. P., Rubens, M. T., Thangavel, A., & Gazzaley, A. (2011). Causal role of the prefrontal cortex in top-down modulation of visual processing and working memory. *Nature Neuroscience*, *14*(5), 656–661. https://doi.org/10.1038/nn.2773

Zarjam, P., Epps, J., & Chen, F. (2011). Characterizing working memory load using EEG delta activity. *2011 19th European Signal Processing Conference*, 1554–1558.

Zhang, X., Mlynaryk, N., Ahmed, S., Japee, S., & Ungerleider, L. G. (2018). The role of inferior frontal junction in controlling the spatially global effect of feature-based attention in human visual areas. *PLOS Biology*, *16*(6), e2005399. https://doi.org/10.1371/journal.pbio.2005399

Zhang, Y., Chen, Y., Bressler, S. L., & Ding, M. (2008). Response preparation and inhibition: The role of the cortical sensorimotor beta rhythm. *Neuroscience*, *156*(1), 238–246. https://doi.org/10.1016/j.neuroscience.2008.06.061

Zimmermann, M., Mars, R. B., de Lange, F. P., Toni, I., & Verhagen, L. (2018). Is the extrastriate body area part of the dorsal visuomotor stream? *Brain Structure & Function*, *223*(1), 31–46. https://doi.org/10.1007/s00429-017-1469-0