ASAR RNAs control replication through interactions with dozens of RNA binding proteins.

Mathew J. Thayer[1,*], Michael B. Heskett[2,4], Leslie G. Smith[1], Paul Spellman[2,3,], and Phillip A. Yates[1].

[1]Department of Chemical Physiology and Biochemistry
[2] Department of Molecular and Medical Genetics
[3]Cancer Early Detection Advanced Research Center, Knight Cancer Institute
Oregon Health & Science University
3181 S. W. Sam Jackson Park Road
Portland, Oregon 97239, USA

[4]Stanford Cancer Institute
291 Campus Drive, Stanford CA, 94305.

*Corresponding author.

ORCID: 0000-0001-6483-1661

TEL: (503) 494-2447

FAX: (503) 494-7368

Email:thayerm@ohsu.edu

Abstract

ASARs are a family of very-long noncoding RNAs that control replication timing on

individual human autosomes, and are essential for chromosome integrity. The eight

known ASAR genes express RNAs that remain localized to their parent chromosomes.

Analysis of eCLIP data revealed >30 nuclear matrix/RNA binding proteins (RBPs) that

interact with multiple ASAR RNAs. An ~7kb domain within the ASAR6-141 RNA shows a

striking density of RBP interaction sites. Genetic deletion and ectopic integration assays

indicate that the ~7kb RNA binding protein domain contains functional sequences for

controlling replication timing of entire chromosomes *in cis*. Depletion of 10 different

ASAR-associated RBPs results in loss of the chromosome territory localization of ASAR

RNAs, and alters the normal synchronous replication timing program on all human

autosomes, recapitulating the effect of individual ASAR gene knockouts on a genome-

wide scale. Our results indicate that ASAR RNAs serve as an essential RNA scaffold for

the assembly of RBP complexes that function in normal chromosome replication and help

maintain the structural integrity of each mammalian chromosome.

Introduction:

The vast majority of mammalian DNA replicates in homologous regions of chromosome pairs in a highly synchronized manner (Dileep and Gilbert, 2018; Mukhopadhyay et al., 2014; Rivera-Mulia et al., 2018). However, genetic disruption of non-protein coding ASAR ("ASynchronous replication and Autosomal RNA") genes causes a delay in replication timing on individual human chromosomes in *cis*, resulting in highly asynchronous replication between pairs of autosomes (Donley et al., 2015; Heskett et al., 2020; Heskett et al., 2022; Stoffregen et al., 2011). The first two ASAR genes were identified at the breakpoints of human chromosome translocations that displayed a chromosome-wide delay in replication timing that is associated with a chromosome-wide delay in mitotic chromosome condensation [DRT/DMC; (Donley *et al.*, 2015; Stoffregen *et al.*, 2011). ASAR RNAs are extremely long (>185 kb), and remain associated with the chromosome territories where they are transcribed (Chang et al., 2007; Donley *et al.*, 2015; Donley et al., 2013; Platt et al., 2018; Stoffregen *et al.*, 2011). One unusual characteristic of ASAR genes is that they are subject to an epigenetic program that results in random Allelic Expression Imbalance [AEI; (Donley *et al.*, 2015; Heskett *et al.*, 2020; Heskett et al., 2021; Heskett *et al.*, 2022; Stoffregen *et al.*, 2011)]. In addition, as the name implies ASAR genes display epigenetically programmed ASynchronous Replication Timing (ASRT) between alleles (Donley *et al.*, 2015; Heskett *et al.*, 2020; Heskett *et al.*, 2022; Stoffregen *et al.*, 2011). Recently, we used these notable ASAR characteristics to identify 69 additional ASAR gene candidates located on every human autosome (Heskett *et al.*, 2020; Heskett *et al.*, 2022). The eight genetically validated ASAR genes are located on human chromosomes 1, 6, 8, 9 and 15 (Donley *et al.*, 2015;

Heskett *et al.*, 2020; Heskett *et al.*, 2022; Stoffregen *et al.*, 2011). Altogether, these

findings suggest that ASARs represent ubiquitous essential *cis*-acting elements of

mammalian chromosome replication timing and structural integrity.

Chromosome-associated long non-protein coding RNAs are known to be important for

numerous aspects of chromosome dynamics in diverse eukaryotes from yeast to

mammals (Creamer et al., 2021; Ding et al., 2019; Faber and Vo, 2022; Menon and

Meller, 2015; Nozawa and Gilbert, 2019). The mammalian nucleus is rich in heterogenous

nuclear RNA (hnRNA), representing >95% of the total RNA Polymerase II output (Nozawa

and Gilbert, 2019; St Laurent et al., 2012). The vast majority of this hnRNA is poorly

defined, has been referred to as the "Dark Matter" of the genome (St Laurent *et al.*, 2012),

and includes spliced and unspliced intronic sequences, circular RNAs (Kristensen et al.,

2019), very long intergenic non-coding RNAs (vlincRNAs) (St Laurent et al., 2016), as

well as poorly defined RNAs that include chromatin associated RNAs [caRNAs; (Nozawa

et al., 2017)] and repeat rich RNAs that can be detected by C0t1 DNA [which is comprised

primarily of LINE and SINE sequences; reviewed in (Creamer *et al.*, 2021; Frank and

Rippe, 2020; Nozawa and Gilbert, 2019; St Laurent *et al.*, 2016)]. Some of these nuclear

RNAs have been studied extensively and comprise abundant species with defined

functions, while others are thought to be the by-products of other nuclear processes with

no known function and have been collectively called "RNA debris" (Nozawa and Gilbert,

2019).

Heterogeneous nuclear ribonucleoproteins (hnRNPs) represent a large family of

abundant RNA-binding proteins (RBPs) that contribute to multiple aspects of nucleic acid

metabolism. Many of the proteins that are in hnRNP complexes share general features,

but differ in domain composition and functional properties [reviewed in (Geuens et al.,

2016)]. The functions of hnRNPs vary according to their subcellular localization. Most of

the hnRNPs possess a conventional nuclear localization signal and are predominantly

present in the nucleus during steady state, where they function in various RNA metabolic

processes including transcription, splicing, and mRNA export [reviewed in (Geuens *et al.*,

2016)].

Here, we used publicly available eCLIP data from ENCODE (Moore et al., 2020) to

identify RBPs that interact with multiple ASAR RNAs, and found >30 different RBPs,

including HNRNPA1, HNRNPU, HNRNPUL1, HNRNPC, and HNRNPM, that interact with

three known ASARs. We also found that an ~7kb region within the ~185 kb ASAR6-141

RNA has a surprisingly high density of interacting RBPs, and that deletion of this ~7kb

"RBP domain" from the endogenous *ASAR6-141* gene results in a delayed replication

timing phenotype that is comparable to deletion of the entire ~185 kb *ASAR6-141* gene

on human chromosome 6. In addition, ectopic integration of transgenes expressing the

~7kb RBP domain into autosomes or into the inactive X chromosome resulted in retention

of the RNA within the chromosome territory of the integrated chromosomes, and resulted

in delayed replication timing, delayed mitotic chromosome condensation, and

chromosome structure instability in *cis*. Taken together, these results indicate that the

~7kb RBP domain represents a critical region within the *ASAR6-141* gene that controls

replication timing of human chromosome 6.

Previous studies found that siRNA mediated depletion, or forced expression of

dominant interfering mutants of HNRNPU resulted in dissociation of C0t1 and Xist RNAs

from their chromosome territories (Creamer *et al.*, 2021; Hall et al., 2014; Hasegawa et

al., 2010; Kolpa et al., 2021). Here we found that shRNA mediated depletion of 10

different ASAR-associated RBPs (HNRNPU, HNRNPC, HNRNPL, HNRNPUL1,

HNRNPM, HLTF, KHSRP, KHDRBS1, UCHL5, or TIA1/TIAL1) results in loss of the

chromosome territory localization of multiple ASAR RNAs. In addition, knockdown of

HNRNPU, HNRNPUL1, UCHL5, or TIA/TIAL1 dramatically altered the normal

synchronous replication timing program of all human autosome pairs, recapitulating the

effect of individual ASAR gene knockouts on a genome-wide scale.

Many recent studies have revealed a surprisingly dynamic and stochastic nature of

chromatin in eukaryotic nuclei, which is reminiscent of a multiphase fluid behavior

(Laghmach et al., 2021a; Laghmach et al., 2021b; Rippe, 2022). Disruption of this liquid-

liquid phase using brief exposure to 1,6-hexanediol results in alterations of higher order

chromatin organization in living cells (Liu et al., 2021; Ulianov et al., 2021). Here, we found

that a brief exposure to 1,6-hexanediol interfered with the chromosome territory

localization of multiple ASAR and C0T1 RNAs. In addition, we found that new RNA

synthesis is required for recovery of the chromosome territory association of C0t1 RNA

following 1,6-hexanediol exposure. In summary, these results indicate that ASAR RNAs

serve as an essential RNA scaffold for the assembly of RBP complexes that function in

normal chromosome replication timing, and in the maintenance of structural integrity of

each mammalian chromosome.

Results

We first sought to identify RBPs that interact with the known ASAR RNAs using

publicly available eCLIP data [see ENCODE; (Moore *et al.*, 2020)]. Using a strategy of

mapping the eCLIP RNA sequence reads to the human genome for >120 RBPs from two

different human cell lines, HEPG2 and K562, we identified dozens of RBPs that interact

with known ASAR RNAs (Table S1; also see Table S2). An example of the alignment of

the eCLIP RNAseq reads to the genomic region containing *ASAR6-141* is shown in Figure

1A, and also illustrates the genomic location of RNAseq reads from total RNA isolated

from both cell lines (also from ENCODE). Given the wide distribution of RNAseq reads

from total RNA that map to *ASAR6-141* in both cell lines, we were surprised to detect an

~7 kb region that contained virtually all of the eCLIP sequences from both cell lines. In

total, we detected 693 independent eCLIP reads from 67 different RBPs in HEPG2 and

217 independent eCLIP reads from 55 RBPs in K562, with sequences from 33 different

RBPs being shared between the two cell lines, and all reads mapping to the ~7kb region

(see Tables S1 and S2). The zoomed in region in Figure 1A shows alignment of the eCLIP

sequences to the ~7kb <u>R</u>NA <u>B</u>inding <u>P</u>rotein <u>D</u>omain (RBPD; see Table S1). In addition,

because K562 cells are female we could directly compare the RBPs that are shared

between the ~7kb RBPD and XIST RNA within the same eCLIP dataset. We found that

51 out of the 55 RBPs that are bound to the ~7kb RBPD in K562 cells are also bound to

XIST RNA (Tables S1 and S2).

Disruption of the ~7kb RBPD within ASAR6-141 results in delayed replication.

To determine if the ~7kb RBPD within the *ASAR6-141* gene contains replication timing

activity we used CRISPR/Cas9 mediated engineering to delete the ~7 kb sequence from

the endogenous locus. For this analysis we designed single guide RNAs (sgRNAs) to

unique sequences on either side of the ~7kb RBPD (see Fig. 1A). We expressed these

sgRNAs in combination with Cas9 in human HTD114 cells and screened clones for

deletion of the ~7kb RBPD (Fig. S1). We chose HTD114 cells for this analysis because

they maintain AEI and ASRT of both imprinted and random monoallelic genes, they have

a stable karyotype that does not change significantly following transfection, drug selection

and sub-cloning, and knockouts of the eight known ASAR genes, including ASAR6-141,

result in delayed replication timing, delayed mitotic condensation and chromosome

structure instability in *cis* (Breger et al., 2005; Chang *et al.*, 2007; Donley *et al.*, 2015;

Donley *et al.*, 2013; Heskett *et al.*, 2020; Heskett *et al.*, 2022; Platt *et al.*, 2018; Stoffregen

et al., 2011). Because *ASAR6-141* expression is mono-allelic in the HTD114 cells

(Heskett *et al.*, 2020), we isolated clones that had heterozygous or homozygous deletions

of the ~7kb RBPD. We determined which alleles were deleted based on retention of the

different base pairs of heterozygous SNPs located within the ~7kb RBPD (see Fig. S1).

To assay replication timing of individual chromosomes we quantified DNA synthesis

in mitotic cells using a BrdU terminal label assay [Fig. 1B; and (Smith et al., 2001; Smith

and Thayer, 2012). In addition, the HTD114 cells contain a centromeric polymorphism on

chromosome 6, which allows for an unambiguous distinction between the two

chromosome 6 homologs [(Donley *et al.*, 2013; Platt *et al.*, 2018; Stoffregen *et al.*, 2011);

also see Fig. 1D and E]. From our previous studies we knew that the chromosome 6 with

the smaller centromere is linked to the expressed allele of *ASAR6-141* (Heskett *et al.*,

2020). For simplicity, we refer to the chromosome 6 with the larger centromere as 6A and

to the chromosome 6 with the smaller centromere as 6B. As expected, prior to disruption

of the ~7kb RBPD chromosomes 6A and 6B, as well as all other chromosome pairs

assayed, display synchronous replication [Fig. 1H; and see (Breger *et al.*, 2005; Donley

et al., 2015; Donley *et al.*, 2013; Stoffregen *et al.*, 2011)]. Figure 1C-1D shows this

replication timing analysis on a mitotic cell where the ~7kb RBPD was deleted from the

expressed allele (CHR6B). Note that CHR6B contains more BrdU incorporation than

CHR6A (Fig. 1E-1G). Quantification of the BrdU incorporation in CHR6B and CHR6A in

multiple cells indicated that deletion of the expressed allele of the ~7kb RBPD resulted in

a significant delay in replication timing (Fig. 1H). This is in contrast to cells containing a

deletion of the ~7kb RBPD from the silent allele (CHR6A) where the BrdU incorporation

is comparable between CHR6B and CHR6A (Fig. 1H). A similar asynchronous replication

pattern between the chromosome 6 homologs was present in cells with homozygous

deletions, with CHR6B representing the later replicating allele (Fig. 1H). In addition, we

found that the delayed replication associated with deletion of the ~7kb RBPD on CHR6B

is comparable to the delayed replication associated with deletion of the entire *ASAR6-*

141 gene from the same chromosome (Fig. 1H). These results indicate that the ~7kb

RBPD contains functional sequences for controlling replication timing, and that

expression of the ~7kb RBPD is important for controlling replication timing of chromosome

6.

Ectopic integration of transgenes expressing the ~7kb RBPD into autosomes.

To determine if the ~7 kb RBPD is sufficient to induce delayed replication in an

ectopic integration assay, we generated two transgenes containing the ~7kb RBPD, one

in the sense orientation and one in the antisense orientation with respect to the promoter

(Fig. 2A). These transgenes were introduced into mouse cells and individual clones

isolated. We initially screened individual clones for expression of the transgenes using

RNA FISH with the ~7kb RBPD as probe. Because the mouse cells that we used are

female, we included an RNA FISH probe for Xist to serve as positive control for the RNA

FISH assay. This two-color RNA FISH assay allowed us to identify clones with single sites

of transgene expression, and allowed us to directly compare the RNA FISH signals from

the transgenes to that of the endogenous Xist RNA hybridization signals expressed from

the inactive X chromosomes. Figure 2B shows examples of the RNA FISH signals in cells

expressing either the sense or antisense transgenes. Large clouds of RNA FISH

hybridization signals that are comparable to Xist RNA hybridization signals were detected

in 4 out of 12 clones expressing the sense transgene. In contrast, 0 out of 12 clones

expressing the antisense transgene expressed RNA FISH hybridization signals

comparable to Xist RNA signals, but instead only small pinpoint sites of RNA FISH

hybridization were detected (Fig. 2B). Note that some of the mouse cells are tetraploid

and therefore contain two Xist RNA hybridization signals.

To determine which mouse chromosomes contain the transgenes, we used DNA

FISH using the ~7kb RBPD DNA as probe in combination with the inverted DAPI banding

pattern to identify the mouse chromosomes in metaphase spreads. We then confirmed

the identity of the mouse chromosomes using DNA FISH on metaphase spreads using

the ~7kb RBPD in combination with a BAC probe from the chromosome of interest. For

example, clone ~7kb(+)A5, which contains the sense-strand transgene integrated in

mouse chromosome 5 is shown in Figure S2A. Next, to determine if the sense strand

RNA expressed from the ~7kb RBPD transgene is retained within the chromosome 5

territory, we used RNA-DNA FISH with the ~7kb RBPD as RNA probe plus a chromosome

paint to detect mouse chromosome 5 DNA in clone ~7kb(+)A5. Figure 2C shows that the

~7kb RBPD RNA is localized to one of the chromosome 5s, indicating that the ~7kb RBPD

RNA is retained within the chromosome 5 territory.

Next to determine if the ~7kb RBPD transgene alters replication timing, we visualized

DNA synthesis in mitotic cells using a BrdU terminal label assay [see Fig. 1B; and (Smith

et al., 2001; Smith and Thayer, 2012)]. In this assay, cells were exposed to BrdU for five

hours, harvested for mitotic cells, and processed for BrdU incorporation and for FISH

using the ~7kb RBPD transgene to identify the integrated chromosome. Figure 2D-E

shows an example of this replication-timing assay in a mitotic cell containing the ~7kb

RBPD sense transgene integrated into mouse chromosome 5. Note that the chromosome

containing the ~7kb RBPD transgene is delayed in mitotic condensation and contains

more BrdU incorporation than any other chromosome within the same cell. In contrast,

integration of transgenes with the ~7kb RBPD in the antisense orientation did not result

in delayed condensation nor in delayed replication of mouse chromosomes (see Fig. 2F

and 2G).

Ectopic integration of the ~7kb RBPD transgene into the inactive X chromosome.

Ectopic integration of *Xist* transgenes into autosomes has been an instrumental tool in

characterization of *Xist* functions, including the ability to delay replication timing and

induce gene silencing (Lee et al., 1996; Minks and Brown, 2009; Pintacuda et al., 2017).

We next sought to determine the consequences of ectopic integration of the ~7kb RBPD

into the inactive X chromosome. The ectopic integration assays described above involved

an RNA FISH screen for clones expressing the ~7kb RBPD, and included an RNA FISH

probe for Xist RNA as positive control. Therefore, to identify transgene integrations into

the inactive X chromosome, we simply screened an additional 50 clones for colocalization

of RNA FISH hybridization signals for both Xist and the ~7kb RBPD sense transgene.

Figure 3A shows this RNA FISH assay on cells where the Xist and ~7kb RBPD

hybridization signals are colocalized. We note that the intensity of the Xist RNA FISH

signal was enhanced in the area where the ~7kb RBPD hybridization signal was detected,

suggesting that the presence of the ~7kb RBPD RNA promoted preferential localization

of Xist RNA in the same region of the X chromosome territory that contains the ~7kb

RBPD RNA.

To confirm that the mouse X chromosome contains the transgene, we used DNA FISH

using the ~7kb RBPD DNA as probe in combination with an X chromosome paint probe

on metaphase spreads. An example of this analysis is shown in Figure S2B-2C, and

indicates that the ~7kb RBPD sense transgene is integrated into the X chromosome.

During this analysis we detected delayed mitotic condensation on X chromosomes, and

the X chromosomes with delayed mitotic condensation contain the ~7kb RBPD

transgene, an example of this analysis is shown in Figure 3C-3D.

Next to determine if the ~7kb RBPD transgene alters replication timing of the inactive

X chromosome, we visualized DNA synthesis in mitotic cells using our BrdU terminal label

assay (Smith *et al.*, 2001; Smith and Thayer, 2012). Figure 3E-3H shows an example of

this assay in a mitotic cell containing the ~7kb RBPD sense transgene integrated into the

inactive X chromosome. In this assay, cells were exposed to BrdU for five hours,

harvested for mitotic cells, and processed for BrdU incorporation and for FISH using the

~7kb RBPD transgene plus an X chromosome paint probe to identify the integrated

chromosome. In summary, integration of the 7kb RBPD transgene was sufficient to cause

DRT/DMC on the inactive X chromosome.

Depletion of RBPs disrupts the chromosome territory localization of ASAR RNAs.

The results described above identified >2 dozen RBPs expressed in two different cell

lines that interact with an ~7kb domain within ASAR6-141 RNA, and that this ~7kb RBPD

is critical for controlling chromosome-wide DNA replication timing. We next used shRNA

knockdown of 13 RBPs that bind within the ~7kb RBPD plus 5 additional RBPs that

interact with two other ASAR RNAs and/or Xist RNA (see Table S1) to determine if these

RBPs function in the chromosome territory localization of ASAR RNAs. For this analysis

we used shRNA knockdown of the RBPs in HTD114 cells. Figure 4A shows the

quantification of HNRNPU by Western blot and by quantitative immunofluoresence

assays, and indicates an ~70% reduction in HNRNPU in both assays. Figure 4B shows

the RNA-DNA FISH for ASAR6-141 RNA in cells expressing empty vector and shows the

expected chromosome territory localized RNA FISH signal on one chromosome 6. In

contrast, depletion of HNRNPU resulted in a dramatic disruption of the ASAR6-141 RNA

hybridization signal into multiple punctate foci that are no longer associated with the

chromosome 6 DNA probe.

Next, to determine if the chromosomal localization of other ASARs was also affected

by HNRNPU knockdown we carried out RNA-DNA FISH for localization of ASAR1-187

and ASAR6 RNAs simultaneously. For this analysis we used Fosmid probes to detect the

ASAR RNAs plus BAC probes to detect the DNA for the ASAR genes located on

chromosomes 1 and 6. Figure 4D shows the RNA-DNA FISH assay on cells expressing

empty vector, and indicates that the RNA FISH signals for both ASARs are closely

associated with their DNA FISH signals on the two different chromosomes. In contrast,

cells with HNRNPU knockdown showed a dispersed punctate nuclear pattern for both

ASAR RNAs, and that the punctate foci show overlapping hybridization signals for both

ASAR RNAs. In addition to the punctate nuclear pattern of RNA localization following

HNRNPU depletion, we also detected cells with cytoplasmic localization of ASAR1-187

and ASAR6 RNAs (Fig. 4F-4G). This cytoplasmic localization was also detected for

ASAR6-141 RNA (not shown). Similar results were obtained following shRNA depletion

of HNRNPC, HNRNPL, HNRNPUL1, HNRNPM, HLTF, KHSRP, KHDRBS1, UCHL5,

TIA1/TIAL1 (Fig. S3). In contrast, shRNA depletion of SAFB, SAFB2, SAFB plus SAFB2,

HNRNPA1, or CTCF did not affect ASAR RNA localization (Fig. S4).

Depletion of RBPs causes asynchronous replication on all autosomes.

Next to determine if ASAR associated RBPs function during the normally synchronous

replication timing program between pairs of homologous chromosomes, we analyzed

DNA synthesis in mitotic cells depleted for HNRNPU, HNRNPUL1, UCHL5 or TIA1/TIAL1

using the BrdU terminal label assay [Fig. 1B; and (Smith *et al.*, 2001; Smith and Thayer,

2012)]. Figure 5 shows this analysis in cells that have been depleted for HNRNPU.

Analysis of cells prior to RBP depletion indicated that each autosome pair displays a

stereotypical and synchronous BrdU incorporation pattern for each autosome pair (Fig.

5A). In contrast, following HNRNPU depletion individual mitotic spreads displayed

dramatic differential incorporation of BrdU into pairs of homologous chromosomes. Figure

5B shows examples of asynchronous replication timing between every pairs of autosome.

Note that each chromosome pair contains dramatic differential BrdU incorporation

between homologs that represents considerable asynchronous replication timing. Figure

5C-5E shows three examples of the chromosome pairs 1, 4 and 18. Quantification of the

BrdU incorporation in chromosomes 1 and 6 in multiple cells depleted for HNRNPU or

TIA1/TIAL1 indicated that each pair of chromosomes displays significant asynchrony (Fig.

5F). In addition, ~1/3 of all mitotic spreads contained one or a few chromosomes that

displayed the full delay in replication timing/delay in mitotic chromosome condensation

(DRT/DMC) phenotype (two examples are shown in Fig. 5G-5H).

A brief exposure to 1,6 hexanediol disrupts ASAR and C0t1 RNA localization.

Recent studies have found that a brief exposure to 1,6-hexanediol results in alterations

of higher order chromatin organization in living cells, which is consistent with disruption

of a liquid-liquid phase separation [LLPS; see (Liu *et al.*, 2021; Ulianov *et al.*, 2021)]. The

shRNA depletion assays described above resulted in mislocalization of multiple ASAR

RNAs into punctate foci, which is suggestive of liquid-liquid phase separation. To

determine if 1,6-hexanediol can disrupt the chromosome territory localization of ASAR

RNAs we exposed HTD114 cells to 4% 1,6-hexanediol for 20 minutes, and then

processed the cells for RNA-DNA FISH. Figure 6 shows this analysis and indicates that

the chromosome territory localization of ASAR6 and ASAR6-141 RNA is disrupted in cells

exposed to 1,6-hexanediol, with both RNAs being detected as a diffuse hybridization

pattern in the nucleus and cytoplasm. In addition, washing out the 1,6-hexanediol and

allowing the cells to recover for 4 hours revealed that the ASAR RNAs regained their

normal chromosome territory localization pattern (Fig. 6D and 6G).

Next to determine if exposure to 1,6-hexanediol can also disrupt the chromosome

territory localization of C0t1 RNA we exposed HA(4)A cells, which are mouse A9 hybrid

cells containing human chromosome 4, to 4% 1,6-hexanediol for 20 minutes and then

processed the cells for RNA-DNA FISH. Figure 7 shows this analysis and indicates that

the chromosome territory localization of hCot1 RNA is disrupted in cells exposed to 1,6-

hexanediol, and that the hC0t1 RNA is detected as a diffuse hybridization pattern in the

nucleus and cytoplasm (Fig. 7B). In addition, washing out the 1,6-hexanediol and allowing

the cells to recover for 4 hours revealed that the hC0t1 RNA regained its normal

chromosome territory association pattern (Fig. 7C). To determine if new RNA synthesis

is required to reestablish the chromosome territory localization of hC0t1 RNA, we treated

cells with 1,6-hexanediol for 20 minutes, washed out the 1,6-hexanediol and allowed the

cells to recover for 4 hours either in the presence or absence of Actinomycin D. Figure

7D-7E shows this analysis and indicates that hC0t RNA was not detected on the human

chromosome territory in the presence of Actinomycin D, indicating that new RNA

synthesis is required for the generation of chromosome territory localization of C0t1 RNA.

Discussion:

We recently identified 69 ASAR candidate genes, and genetically validated 6 out of 6 as

controlling replication timing of individual chromosomes, bringing the total of genetically

validated ASAR genes to eight (Heskett *et al.*, 2020; Heskett *et al.*, 2022). The eight ASAR

genes (*ASAR1-187, ASAR6, ASAR6-141, ASAR8-2.7, ASAR9-23, ASAR9-24, ASAR9-*

30 and ASAR15) display striking epigenetically controlled AEI, ASRT and their RNAs

remain localized to their parent chromosome territories (Donley *et al.*, 2015; Heskett *et*

al., 2020; Heskett *et al.*, 2022; Stoffregen *et al.*, 2011). Evidence that ASAR RNA

mediates the chromosome-wide effects on replication timing came from oligonucleotide

mediated RNA degradation of ASAR6 RNA that was expressed from a BAC transgene

integrated into a mouse chromosome, thus reversing the chromosome-wide delayed

replication timing induced by the *ASAR6* transgene (Platt *et al.*, 2018). In this report, we

used publicly available RNA-protein interaction data from ENCODE in combination with

a strategy of mapping the eCLIP sequences to the human genome to identify >30 nuclear

matrix/RBPs that interact with multiple ASAR RNAs. One unanticipated observation that

came from this analysis was that an ~7kb domain within the ~185kb ASAR6-141 RNA

contained virtually all of the RBP interaction sites for >30 different RBPs in two different

human cell lines. Using genetic deletion and ectopic integration assays we found that the

~7kb RBPD contains functional sequences for controlling replication timing of entire

chromosomes in *cis*. We also found that shRNA-mediated depletion of 10 different ASAR-

associated RBPs results in loss of the chromosome territory association of multiple ASAR

RNAs. In addition, we found that depletion of HNRNPU, HNRNPUL1, UCHL5, and

TIA/TIAL1 alters the normally synchronous replication timing that occurs between pairs

of homologous chromosomes, recapitulating the effect of individual ASAR gene

knockouts on a genome-wide scale. Furthermore, we found that a brief exposure of cells

to 1,6-hexanediol disrupts the chromosome territory localization of multiple ASAR RNAs

as well as C0t1 RNA, and that new RNA synthesis is required to restore the chromosome

territory localization of C0t1 RNA.

Previous studies have suggested that long nascent transcripts that are detected by

C0t1 DNA play a dynamic structural role that promotes the open architecture of active

chromosome territories (Creamer *et al.*, 2021; Hall *et al.*, 2014). The RNA species

detected by C0t1 DNA are predominantly L1 sequences, and these L1 RNAs remain

associated with the chromosome territories where they are made (Hall *et al.*, 2014).

Additional studies have described <u>rep</u>etitive RNA (repRNA) as important regulators of the

dynamic organization of genomic loci into membrane-less subcompartments with distinct

nuclear functions [reviewed in (Frank and Rippe, 2020)]. Recent studies have also

proposed that abundant nuclear proteins such as HNRNPU nonspecifically interact with

"RNA debris" that creates a dynamic nuclear mesh that regulates interphase chromatin

structure (Nozawa *et al.*, 2017; Nozawa and Gilbert, 2019). One important distinction that

can be made between ASARs and these other hnRNAs [i.e. C0t1 RNA, repRNA, caRNA,

and "RNA debris" (Creamer *et al.*, 2021; Frank and Rippe, 2020; Hall *et al.*, 2014; Kolpa

et al., 2021; Nozawa *et al.*, 2017; Nozawa and Gilbert, 2019)] is that ASARs are defined

genetically, and these other RNAs are not. Thus, we have used both loss of function and

gain of function assays to define the roles that ASARs play in chromosome dynamics,

which includes control of chromosome-wide replication timing and mitotic chromosome

condensation (Donley *et al.*, 2015; Donley *et al.*, 2013; Heskett *et al.*, 2020; Heskett *et*

al., 2022; Platt *et al.*, 2018; Stoffregen *et al.*, 2011). We also note that C0t1 DNA, when

used as an RNA FISH probe, can detect ASAR6 RNA that is expressed and localized to

an individual chromosome territory (Figure S4), indicating that at least some of the RNA

FISH signal detected by C0T1 DNA represents ASARs. In addition, we found that a brief

exposure to 1,6-hexanediol disrupts the chromosome territory localization of multiple

ASAR RNAs, suggesting that ASAR RNAs participate in the nuclear LLPS processes that

organize chromosome territories. Given our recent discovery of ASAR candidate genes

on every autosome, representing ~3% of the human genome (Heskett *et al.*, 2022), we

propose that the functional "chromosome associated RNA" species that control

chromosome dynamics within mammalian nuclei are encoded by ASAR genes. Whether

or not there are nuclear functions of "RNA debris" that are independent of ASARs will

require genetic and/or functional validation.

Here we identified several hnRNPs (HNRNPU, HNRNPUL1, HNRNPC, and HNRNPM)

as important interaction partners for the chromosome territory localization of ASAR RNAs.

In addition, we found that the KH domain (HNRNP<u>K</u> <u>H</u>omology domain) containing

proteins KHSRP and KHDRBS1 also contribute to the chromosome territory localization

of ASAR RNAs. We also found that depletion of HNRNPU, HNRNPUL1, UCHL5, and

TIA1/TIAL1 proteins resulted in a dramatic genome-wide disruption of the normally

synchronous replication timing program on all autosome pairs. Taken together our results

indicate that the association of ASAR RNAs with hnRNPs and nuclear matrix associated

proteins is an essential step in the regulation of the replication timing program on

mammalian chromosomes.

References:
Breger, K.S., Smith, L., and Thayer, M.J. (2005). Engineering translocations with
delayed replication: evidence for cis control of chromosome replication timing. Hum Mol
Genet *14*, 2813-2827.
Chang, B.H., Smith, L., Huang, J., and Thayer, M. (2007). Chromosomes with delayed
replication timing lead to checkpoint activation, delayed recruitment of Aurora B and
chromosome instability. Oncogene *26*, 1852-1861.
Creamer, K.M., Kolpa, H.J., and Lawrence, J.B. (2021). Nascent RNA scaffolds
contribute to chromosome territory architecture and counter chromatin compaction. Mol
Cell. 10.1016/j.molcel.2021.07.004.
Dileep, V., and Gilbert, D.M. (2018). Single-cell replication profiling to measure
stochastic variation in mammalian replication timing. Nature communications *9*, 427.
10.1038/s41467-017-02800-w.
Ding, D.Q., Okamasa, K., Katou, Y., Oya, E., Nakayama, J.I., Chikashige, Y., Shirahige,
K., Haraguchi, T., and Hiraoka, Y. (2019). Chromosome-associated RNA-protein
complexes promote pairing of homologous chromosomes during meiosis in
Schizosaccharomyces pombe. Nature communications *10*, 5598. 10.1038/s41467-019-
13609-0.
Donley, N., Smith, L., and Thayer, M.J. (2015). ASAR15, A cis-Acting Locus that
Controls Chromosome-Wide Replication Timing and Stability of Human Chromosome
15. PLoS Genet *11*, e1004923. 10.1371/journal.pgen.1004923.
Donley, N., Stoffregen, E.P., Smith, L., Montagna, C., and Thayer, M.J. (2013).
Asynchronous Replication, Mono-Allelic Expression, and Long Range Cis-Effects of
ASAR6. PLoS Genet *9*, e1003423.
Faber, M.W., and Vo, T.V. (2022). Long RNA-Mediated Chromatin Regulation in Fission
Yeast and Mammals. Int J Mol Sci *23*. 10.3390/ijms23020968.
Frank, L., and Rippe, K. (2020). Repetitive RNAs as Regulators of Chromatin-
Associated Subcompartment Formation by Phase Separation. J Mol Biol *432*, 4270-
4286. 10.1016/j.jmb.2020.04.015.
Geuens, T., Bouhy, D., and Timmerman, V. (2016). The hnRNP family: insights into
their role in health and disease. Hum Genet *135*, 851-867. 10.1007/s00439-016-1683-5.
Hall, L.L., Carone, D.M., Gomez, A.V., Kolpa, H.J., Byron, M., Mehta, N., Fackelmayer,
F.O., and Lawrence, J.B. (2014). Stable C0T-1 repeat RNA is abundant and is
associated with euchromatic interphase chromosomes. Cell *156*, 907-919.
10.1016/j.cell.2014.01.042.
Hasegawa, Y., Brockdorff, N., Kawano, S., Tsutui, K., Tsutui, K., and Nakagawa, S.
(2010). The matrix protein hnRNP U is required for chromosomal localization of Xist
RNA. Dev Cell *19*, 469-476. 10.1016/j.devcel.2010.08.006.
Heskett, M.B., Smith, L.G., Spellman, P., and Thayer, M.J. (2020). Reciprocal
monoallelic expression of ASAR lncRNA genes controls replication timing of human
chromosome 6. RNA (New York, N.Y.) *26*, 724-738. 10.1261/rna.073114.119.
Heskett, M.B., Spellman, P.T., and Thayer, M.J. (2021). Differential Allelic Expression
among Long Non-Coding RNAs. Non-coding RNA *7*. 10.3390/ncrna7040066.
Heskett, M.B., Vouzas, A.E., Smith, L.G., Yates, P.A., Boniface, C., Bouhassira, E.E.,
Spellman, P.T., Gilbert, D.M., and Thayer, M.J. (2022). Epigenetic Control of Hundreds

of Chromosome-Associated lncRNA Genes Essential for Replication and Stability.
bioRxiv, 2022.2004.2025.489405. 10.1101/2022.04.25.489405.
Kolpa, H.J., Creamer, K.M., Hall, L.L., and Lawrence, J.B. (2021). SAF-A mutants
disrupt chromatin structure through dominant negative effects on RNAs associated with
chromatin. Mamm Genome. 10.1007/s00335-021-09935-8.
Kristensen, L.S., Andersen, M.S., Stagsted, L.V.W., Ebbesen, K.K., Hansen, T.B., and
Kjems, J. (2019). The biogenesis, biology and characterization of circular RNAs. Nat
Rev Genet *20*, 675-691. 10.1038/s41576-019-0158-7.
Kruskal, J.B. (1964). Multidimensional scaling by optimizing goodness of fit to a
nonmetric hypothesis. Psychometrika *29*, 1–27.
Laghmach, R., Di Pierro, M., and Potoyan, D. (2021a). A Liquid State Perspective on
Dynamics of Chromatin Compartments. Front Mol Biosci *8*, 781981.
10.3389/fmolb.2021.781981.
Laghmach, R., Di Pierro, M., and Potoyan, D.A. (2021b). The interplay of chromatin
phase separation and lamina interactions in nuclear organization. Biophys J *120*, 5005-
5017. 10.1016/j.bpj.2021.10.012.
Lee, J.T., Strauss, W.M., Dausman, J.A., and Jaenisch, R. (1996). A 450 kb transgene
displays properties of the mammalian X-inactivation center. Cell *86*, 83-94.
Liu, X., Jiang, S., Ma, L., Qu, J., Zhao, L., Zhu, X., and Ding, J. (2021). Time-dependent
effect of 1,6-hexanediol on biomolecular condensates and 3D chromatin organization.
Genome Biol *22*, 230. 10.1186/s13059-021-02455-3.
Menon, D.U., and Meller, V.H. (2015). Identification of the Drosophila X chromosome:
The long and short of it. RNA biology *12*, 1088-1093. 10.1080/15476286.2015.1086864.
Minks, J., and Brown, C.J. (2009). Getting to the center of X-chromosome inactivation:
the role of transgenes. Biochem Cell Biol *87*, 759-766.
Moore, J.E., Purcaro, M.J., Pratt, H.E., Epstein, C.B., Shoresh, N., Adrian, J., Kawli, T.,
Davis, C.A., Dobin, A., Kaul, R., et al. (2020). Expanded encyclopaedias of DNA
elements in the human and mouse genomes. Nature *583*, 699-710. 10.1038/s41586-
020-2493-4.
Mukhopadhyay, R., Lajugie, J., Fourel, N., Selzer, A., Schizas, M., Bartholdy, B., Mar,
J., Lin, C.M., Martin, M.M., Ryan, M., et al. (2014). Allele-specific genome-wide profiling
in human primary erythroblasts reveal replication program organization. PLoS Genet *10*,
e1004319. 10.1371/journal.pgen.1004319.
Nozawa, R.S., Boteva, L., Soares, D.C., Naughton, C., Dun, A.R., Buckle, A.,
Ramsahoye, B., Bruton, P.C., Saleeb, R.S., Arnedo, M., et al. (2017). SAF-A Regulates
Interphase Chromosome Structure through Oligomerization with Chromatin-Associated
RNAs. Cell *169*, 1214-1227.e1218. 10.1016/j.cell.2017.05.029.
Nozawa, R.S., and Gilbert, N. (2019). RNA: Nuclear Glue for Folding the Genome.
Trends Cell Biol *29*, 201-211. 10.1016/j.tcb.2018.12.003.
Pintacuda, G., Wei, G., Roustan, C., Kirmizitas, B.A., Solcan, N., Cerase, A., Castello,
A., Mohammed, S., Moindrot, B., Nesterova, T.B., and Brockdorff, N. (2017). hnRNPK
Recruits PCGF3/5-PRC1 to the Xist RNA B-Repeat to Establish Polycomb-Mediated
Chromosomal Silencing. Mol Cell *68*, 955-969.e910. 10.1016/j.molcel.2017.11.013.
Platt, E.J., Smith, L., and Thayer, M.J. (2018). L1 retrotransposon antisense RNA within
ASAR lncRNAs controls chromosome-wide replication timing. J Cell Biol *217*, 541-553.
10.1083/jcb.201707082.

Rippe, K. (2022). Liquid-Liquid Phase Separation in Chromatin. Cold Spring Harb
Perspect Biol *14*. 10.1101/cshperspect.a040683.
Rivera-Mulia, J.C., Dimond, A., Vera, D., Trevilla-Garcia, C., Sasaki, T., Zimmerman, J.,
Dupont, C., Gribnau, J., Fraser, P., and Gilbert, D.M. (2018). Allele-specific control of
replication timing and genome organization during development. Genome Res *28*, 800-
811. 10.1101/gr.232561.117.
Smith, L., Plug, A., and Thayer, M. (2001). Delayed Replication Timing Leads to
Delayed Mitotic Chromosome Condensation and Chromosomal Instability of
Chromosome Translocations. Proc Natl Acad Sci U S A *98*, 13300-13305.
Smith, L., and Thayer, M. (2012). Chromosome replicating timing combined with
fluorescent in situ hybridization. J Vis Exp *10*, e4400.
St Laurent, G., Savva, Y.A., and Kapranov, P. (2012). Dark matter RNA: an intelligent
scaffold for the dynamic regulation of the nuclear information landscape. Frontiers in
genetics *3*, 57. 10.3389/fgene.2012.00057.
St Laurent, G., Vyatkin, Y., Antonets, D., Ri, M., Qi, Y., Saik, O., Shtokalo, D., de Hoon,
M.J., Kawaji, H., Itoh, M., et al. (2016). Functional annotation of the vlinc class of non-
coding RNAs using systems biology approach. Nucleic Acids Research. 2 *44*, 3233-
3252. 10.1093/nar/gkw162.
Stoffregen, E.P., Donley, N., Stauffer, D., Smith, L., and Thayer, M.J. (2011). An
autosomal locus that controls chromosome-wide replication timing and mono-allelic
expression. Hum Mol Genet *20*, 2366-2378.
Ulianov, S.V., Velichko, A.K., Magnitov, M.D., Luzhin, A.V., Golov, A.K., Ovsyannikova,
N., Kireev, II, Gavrikov, A.S., Mishin, A.S., Garaev, A.K., et al. (2021). Suppression of
liquid-liquid phase separation by 1,6-hexanediol partially compromises the 3D genome
organization in living cells. Nucleic Acids Research. 2 *49*, 10524-10541.
10.1093/nar/gkab249.
Zhu, Y., Bye, S., Stambrook, P.J., and Tischfield, J.A. (1994). Single-base deletion
induced by benzo[a]pyrene diol epoxide at the adenine phosphoribosyltransferase locus
in human fibrosarcoma cell lines. Mutat Res *321*, 73-79.
543
544

545   **Figure legends**
546
Figure 1. RNA binding proteins interact with an~7kb domain within ASAR6-141
RNA. A) Integrative Genomics Viewer view showing the genomic location of *ASAR6-141*.
The genomic locations of RNAseq and eCLIP reads from HEPG2 and K562 (from
ENCODE) are shown. The zoomed in view shows the ~7kb RBPD with the location of all
of the eCLIP reads that map within the region (see Table S2). The location of sgRNAs
are shown, the asterisks mark the sgRNAs from (Heskett *et al.*, 2020). B) Schematic
illustration of the BrdU terminal label protocol(Smith and Thayer, 2012). Cells were
treated with BrdU (green) for 5.5 hours and then harvested for mitotic cells. C-H) BrdU
incorporation in HTD114 cells containing a heterozygous deletion of the the ~7kb RBPD.
Cells containing a deletion of the ~7kb RBPD from the expressed allele of *ASAR6-141*
were exposed to BrdU, harvested for mitotic cells, and subjected to DNA FISH using
chromosome 6 centromeric probe (red; D-E). The larger centromere resides on the
chromosome 6 with the silent *ASAR6-141* allele (6A), and the smaller centromere resides
on the chromosome 6 with the expressed *ASAR6-141* allele (6B). DNA was stained with
DAPI (blue). Bar is 10 uM. F and G) DAPI staining and BrdU incorporation were quantified
by calculating the number of pixels (area x intensity) and displayed as a ratio of BrdU
incorporation in chromosome 6B divided by the BrdU incorporation in chromosome 6A.
H) Quantification of BrdU incorporation in multiple cells with heterozygous (Δ7kb-6B,
expressed allele; or Δ7kb-6A, silent allele) or homozygous (Δ7kb-6B/6A) deletions of the
~7kb RBPD. Also included is the quantification of BrdU incorporation in heterozygous
deletions of the entire ~185kb ASAR6-141 gene from chromosome 6B (Δ185-6B) or 6A
(Δ185-6A) (Heskett *et al.*, 2020). **P = 0.00027 and *P = 0.03.
569
570
Figure 2. Replication timing in mouse cells containing *ASAR6-141 ~7kb RBPD*
transgenes. A) Schematic view of the ~7kb RBPD in both the sense and antisense
orientation. The promoter used to drive expression is from the ASAR6 gene (Platt *et al.*,
2018). B) Two color RNA FISH assay for expression of the ~7kb RBPD transgenes.
Individual clones were screened for expression of the ASAR6-141 (~7kb RBPD)
transgenes (red). RNA hybridization using an Xist probe (green) was used as positive
control. The left and right panels represent examples of clones containing the sense and
antisense transgenes, respectively. The arrows mark the ASAR6-141 RNA. C) RNA from
the ~7kb RBPD remains localized to mouse chromosomes. RNA-DNA FISH using the
~7kb RBPD as RNA FISH probe (red), plus a mouse chromosome 5 paint to detect
chromosome 5 DNA (green). D and E) Cells containing the sense ~7kb RBPD transgene
integrated into mouse chromosome 5 were exposed to BrdU (green), harvested for mitotic
cells, and subjected to DNA FISH using the ~7kb RBPD (red). Note that the chromosome
that contains the transgene shows delayed mitotic condensation and more BrdU
incorporation than any other chromosome within the same cell. F-G) Cells containing the
antisense ~7kb RBPD transgene integrated into mouse chromosomes were exposed to
BrdU (green), harvested for mitotic cells, and subjected to DNA FISH using the ~7kb
RBPD (red). Note that the chromosome that contains the transgene shows normal mitotic
condensation and only a small amount of BrdU incorporation.
590

Figure 3. Replication timing in mouse cells containing *ASAR6-141 ~7kb RBPD* transgene integrated into the inactive X chromosome. A) Two color RNA FISH assay for expression of the sense ~7kb RBPD transgene plus Xist RNA. Individual clones were screened for expression of the sense strand ~7kb RBPD transgene (red) in combination with RNA FISH for Xist (green). Examples of three different cells showing colocalization of the ASAR RNA and Xist RNA. The white lines show the path for the pixel intensity profiles in panel B. B) Pixel intensity profiles across the Xist RNA hybridization domain showing enhanced signal intensity over the ~7kb RBPD RNA hybridization domain. C and D) Delayed mitotic chromosome condensation of the X chromosome that contains the ~7kb RBPD transgene. An example of a mitotic spread processed for DNA FISH using the ~7kb RBPD as probe (magenta) plus a mouse X chromosome paint (red). The arrows mark the location of the transgene hybridization signal. E-H) Cells containing the sense ~7kb RBPD transgene integrated into the mouse X chromosome were exposed to BrdU (green), harvested for mitotic cells, and subjected to DNA FISH using the ~7kb RBPD (magenta) plus a mouse X chromosome paint. Note that the chromosome that contains the transgene (arrows) displays delayed mitotic condensation and more BrdU incorporation than any other chromosome within the same cell.

Figure 4. Depletion of HNRNPU results in disruption of the chromosome territory localization of ASAR RNAs. A) Quantitation of HNRNPU by Western blot (top) and by quantitative immunofluorescence (bottom). B and C) RNA-DNA FISH for ASAR6-141 RNA. HTD114 cells were transfected with empty vector (B) or vector expressing an shRNA against HNRNPU (C). The white arrows mark the RNA FISH signals and the arrowheads mark the DNA FISH signals. B) Examples of three different nuclei showing the RNA hybridization pattern for ASAR6-141 RNA in cells transfected with empty vector. C) The top and bottom panels show examples of two different nuclei showing a punctate RNA hybridization pattern for ASAR6-141 RNA in cells that were depleted for HNRNPU. D-H) RNA-DNA FISH for ASAR1-187 and ASAR6 in cells depleted for HNRNPU. D) Examples of nuclei from cells transfected with empty vector. The white arrows mark the RNA FISH signals for ASAR6 (red) and ASAR1-187 (green). The DNA FISH signals for *ASAR6* DNA (orange) and *ASAR1-187* (magenta) are marked with colored arrows as shown. The merged image shows that the RNA and DNA FISH signals are adjacent to each other. E) RNA FISH image showing a punctate nuclear pattern for ASAR6 RNA (red) and ASAR1-187 RNA (green). The merged image shows that the FISH signal for both ASARs are colocalized. F-H) RNA-DNA FISH showing cytoplasmic localization of ASAR6 and ASAR1-187 RNAs. The white arrows mark the sites of colocalization of ASAR6 and ASAR1-187 RNA in punctate foci in the cytoplasm. The FISH signals for *ASAR6* DNA (orange) and *ASAR1-187* DNA (magenta) are marked with colored arrows as shown. DNA was stained with DAPI, and Bars are 10*u*m.

Figure 5. Depletion of HNRNPU results in asynchronous replication on autosome pairs. A) BrdU incorporation in parental HTD114 cells. Cells were exposed to BrdU for five hours, processed for BrdU incorporation using an antibody against BrdU. This panel represents chromosomes from multiple mitotic spreads showing representative BrdU incorporation in chromosome. Each autosome pair were captured from the same mitotic cell, and each pair displays a stereotypical BrdU incorporation pattern that is consistent

with synchronous replication timing. B) HTD114 cells were transfected with the HNRNPU
shRNA expression vector, exposed to BrdU for five hours, and processed for BrdU
incorporation. This panel represents chromosomes from multiple mitotic spreads showing
representative BrdU incorporation in pairs of autosomes. Each autosome pair were
captured from the same mitotic cell, and each pair displays a differential BrdU
incorporation pattern that is consistent with asynchronous replication timing. C) Shows
two chromosome 1 homologs side by side. For each CHR1 homolog (A or B) we show
the inverted DAPI staining (black and white), chromosome 1 centromeric probe (red),
BrdU incorporation (green), and DAPI staining (blue). Panels D and E show the BrdU
incorporation in chromosomes 4 (CHR4A and CHR4B) and 18 (CHR18A and CHR18B),
respectively. For each chromosome pair the inverted DAPI staining (black and white),
BrdU incorporation (green), and DAPI staining (blue) are shown. Quantification of BrdU
incorporation in multiple cells indicated that HNRNPU or TIA/TIAL1 depletion resulted in
significant asynchrony (P < 1 x 10^{-5}) between chromosome 1 and chromosome 6
homologs. The P values are form the Kruskal-Wallis test (Kruskal, 1964). G) DRT/DMC
on chromosome 6 following HNRNPU depletion. HTD114 cells were transfected with the
HNRNPU shRNA expression vector, exposed to BrdU for 5 hours, harvested for mitotic
cells, and processed for BrdU incorporation (green) and DNA FISH using a chromosome
6 centromeric probe (red). H) DRT/DMC on chromosome 1 following UCHL5 depletion.
HTD114 cells were transfected with the UCHL5 shRNA expression vector, exposed to
BrdU for 5 hours, harvested for mitotic cells, and processed for BrdU incorporation (green)
and DNA FISH using a chromosome 1 paint as probe (red). The arrow marks the
chromosome 1 with DRT/DMC, and the inset shows only the DAPI staining of the
chromosome 1 highlighting the DMC.

661
Figure 6. 1,6 hexanediol disrupts the chromosome territory association of ASAR
RNAs. HTD114 cells were exposed to 4% 1,6 hexanediol (HD) for 20 minutes, washed
extensively and allowed to recover for 4 hours in complete media, and processed for
RNA-DNA FISH. A-C) RNA FISH using an ASAR6-141 probe plus a BAC probe from the
ASAR6 locus to detect DNA. D-F) RNA FISH using an ASAR6 probe plus a BAC probe
from the ASAR6 locus to detect DNA. Panels A and D are HTD114 cells prior to exposure.
B and E) HTD114 cells exposed to HD for 20 minutes. C and F) HTD114 cells exposed
to HD for 20 minutes followed by 4 hour incubation in the absence of HD (4hr washout).

670
Figure 7. 1,6 hexanediol disrupts the chromosome territory association of C0t1
RNA. HA(4)A cells were exposed to 4% 1,6 hexanediol (HD) for 20 minutes, washed
extensively and allowed to recover for 4 hours in complete media, and processed for
RNA-DNA FISH using C0t1 as probe from both RNA and DNA. A) Control HA(4)A cells
prior to HD exposure. B) HA(4)A cells exposed to HD for 20 minutes. C) HA(4)A cells
exposed to HD for 20 minutes, washed extensively and allowed to recover for 4 hours in
complete media. D and E) HA(4)A cells exposed to HD for 20 minutes, washed
extensively and allowed to recover for 4 hours in complete media containing Actinomycin
D (ActD). F) Control HA(4)A cells exposed to Actinomycin D for 4 hours.

680
Figure S1. CRISPR/Cas9 deletion of the ~7kb RBPD. A) Integrative Genomics Viewer
view showing the genomic location of *ASAR6-141*. The genomic locations of RNAseq

and eCLIP reads from HEPG2 and K562 (from ENCODE) are shown. The zoomed in
view shows the ~7kb RBPD with the location of all of the eCLIP reads that map within the
region (see Table S2). The location of sgRNAs that flank the ~7kb RBPD are shown. The
location of PCR primers (1-6) and the heterozygous SNP rs17070386 are also shown. B)
Sanger sequencing traces on PCR products generated from genomic DNA using primers
5 and 6 (see Table S3). In previous studies we generated mouse monochromosomal
hybrid cells containing the two chromosome 6 homologs, and sequencing traces from
these two hybrids, containing either chromosome 6A (CHR6A) or chromosome 6B
(CHR6B) are shown below the HTD114 panel. Sequencing traces from PCR products
generated from 4 heterozygous deletions, two with CHR6A deletions {D6A(7-82-29) and
D6A(7-82-13)} and two with CHR6B deletions {D6B(7-18-1) and D6B(7-63-1)} are shown.
C) Junction PCR (primers 1 and 4) on genomic DNA isolated from heterozygous {D6A(7-
82-29), D6A(7-82-13), D6B(7-18-1) and D6B(7-63-1)} or homozygous deletions {D6B/6A-
1 and D6B/6A-2} are shown. Parental HTD114 DNA was used as control. D) Internal PCR
products (primers 5 and 6) on genomic DNA isolated from heterozygous {D6A(7-82-29),
D6A(7-82-13), D6B(7-18-1) and D6B(7-63-1)} or homozygous deletions {D6B/6A-1 and
D6B/6A-2} are shown. Parental HTD114 DNA was used as control.
700
Figure S2. Identification of mouse chromosomes containing the ~7kb RBPD
transgene. A) DNA FISH on mouse chromosomes from clone ~7kb(+)A5. Mitotic spread
showing DNA FISH using the ~7kb RBPD (magenta; marked with arrows) and a mouse
chromosome 5 BAC (RP24-107D17; red; marked with arrowheads). B and C) DNA FISH
on mouse chromosomes from clone ~7kb(+)X1. Mitotic spread showing DNA FISH using
the ~7kb RBPD (magenta; marked with arrows) and a mouse X chromosome paint (red).
707
Figure S3. Disruption of the chromosome territory localization of ASAR1-187 and
ASAR6 RNAs following shRNA depletion of RBPs. HTD114 cells were transfected
with empty vector (A) or shRNA expression vectors for KHSRP (B), HNRNPL (C),
HNRNPUL1 (D), UCHL5 (E), KHDRBS1 (F), HTLF (G), HNRNPC (H), or TIA/TIAL1 (note
that the single knockdowns of TIA or TIAL1 did not disrupt ASAR RNA localization, not
shown). Cells were processed for RNA FISH using Fosmid probes (see Table S3) for
ASAR1-187 (green) and ASAR6 (red).
715
716 **Figure S4. RNA FISH on cells depleted for RBPs that did not disrupt ASAR**
717 **localization.** HTD114 cells were transfected with shRNA expression vectors for SAFB
718 (A), SAFB2 (B), SAFB and SAFB2 (C), HNRNPA1 (D), CTCF (E), or empty vector (F).
719 Cells were processed for RNA FISH using Fosmid probes (see Table S3) for ASAR1-187
720 (green) and ASAR6 (red).
721
Figure S5. Human C0t1 DNA detects ASAR6 RNA. Mouse cells containing an *ASAR6*
BAC (RP11-767E7) transgene integrated into mouse chromosome 3 were previously
described (Donley *et al.*, 2013). A-C) RNA-DNA FISH using and ASAR6 fosmid probe to
ASAR6 RNA (green; arrows) plus a mouse chromosome 3 paint probe to detect DNA
(red; arrowheads). D-F) RNA-DNA FISH using human C0t1 DNA to detect RNA (green;
arrows) plus the *ASAR6* BAC (red arrowheads) to detect DNA.
728

STAR Methods

Cell culture

HTD114 cells are a human *APRT* deficient cell line derived from HT1080 cells (Zhu et al., 1994). Mouse C2C12 cells were used for the transgene integration assays, transfected using Lipofectamine 2000 (Invitrogen) according to the manufacturer's recommendations, and clones containing transgene integrations were isolated following selection in media containing G418. All cells were grown in DMEM (Gibco) supplemented with 10% fetal bovine serum (Hyclone), and were grown in a humidified incubator at 37°C in a 5% carbon dioxide atmosphere.

shRNA depletion of RBPs

Using Lipofectamine 2000 (Invitrogen), according to the manufacturer's recommendations, we co-transfected HTD114 cells with plasmids encoding puromycin resistance (PGK-PURO) and shRNAs against 18 different RBPs (Millipore/Sigma). 24h after transfection, cells were plated in media containing 10 ug/ml puromycin (Millipore/Sigma), and 24h-48h following puromycin selection cells were processed for Western blot, RNA-DNA FISH or for replication timing assays (see below).

Western blot analysis

CRISPR/Cas9 engineering

Using Lipofectamine 2000 (Invitrogen), according to the manufacturer's recommendations, we co-transfected HTD114 cells with plasmids encoding GFP,

sgRNAs and Cas9 endonuclease (Origene). Each plasmid encoded sgRNA was designed

to bind at the indicated locations (Figure S1). 48h after transfection, cells were plated at

clonal density and allowed to expand for 2-3 weeks. The presence of deletions was

identified by PCR using the primers described in Figure S1. The single cell colonies that

grew were analyzed for heterozygous deletions by PCR (Figure S1). We used retention

of a heterozygous SNP to identify the disrupted allele, and homozygosity at the SNP

confirmed that cell clones were homogenous.

PCR analysis

Genomic DNA was isolated from tissue culture cells using TRIZOL Reagent

(Invitrogen). PCR was performed in a 12.5μL volume using 50-100ng of genomic DNA,

1x Standard Taq Buffer (New England Biolabs, Inc.), 200μM each deoxynucleotide

triphosphates, 0.2μM of each primer, and 3 units of Taq DNA Polymerase (New England

Biolabs, Inc.) under the following reaction conditions: 95°C for 2 minutes, followed by 30-

40 cycles of 95°C for 30 seconds, 55-62°C for 45 seconds, and 72°C for 1 minute, with a

final extension time of 10 minutes at 72°C. PCR products were separated on 1% agarose

gels, stained with ethidium bromide, and photographed under ultraviolet light illumination.

Sequencing of PCR products was carried out at the Vollum Institute DNA Sequencing

Core facility.

DNA FISH

Mitotic chromosome spreads were prepared as described previously (Smith *et al.*,

2001). After RNase (100μg/ml) treatment for 1h at 37°C, slides were washed in 2XSSC

and dehydrated in an ethanol series and allowed to air dry. Chromosomal DNA on the

slides was denatured at 75°C for 3 minutes in 70% formamide/2XSSC, followed by

dehydration in an ice-cold ethanol series and allowed to air dry. BAC and Fosmid DNAs

were labeled using nick translation (Vysis, Abbott Laboratories) with Spectrum Orange-

dUTP, Spectrum Aqua-dUTP or Spectrum Green-dUTP (Vysis). Final probe

concentrations varied from 40-60 ng/µl. Centromeric probe cocktails (Vysis) and/or whole

chromosome paint probes (Metasystems) plus BAC or Fosmid DNAs were denatured at

75°C for 10 minutes and prehybridized at 37°C for 10 minutes. Probes were applied to

denatured slides and incubated overnight at 37°C. Post-hybridization washes consisted of

one 3-minute wash in 50% formamide/2XSSC at 40°C followed by one 2-minute rinse in

PN (0.1M Na_2HPO_4, pH 8.0/2.5% Nonidet NP-40) buffer at RT. Coverslips were mounted

with Prolong Gold antifade plus DAPI (Invitrogen) and viewed under UV fluorescence

(Olympus).

RNA-DNA FISH

Cells were plated on Poly-L-Lysine coated (Millipore Singa) glass microscope slides

at ~50% confluence and incubated for 4 hours in complete media in a 37°C humidified

CO_2 incubator. Slides were rinsed 1X with sterile RNase free PBS. Cell Extraction was

carried out using ice cold solutions as follows: Slides were incubated for 30 seconds in

CSK buffer (100mM NaCl/300mM sucrose/3mM $MgCl_2$/10mM PIPES, pH 6.8), 10 minutes

in CSK buffer/0.1% Triton X-100, followed by 30 seconds in CSK buffer. Cells were then

fixed in 4% paraformaldehyde in PBS for 10 minutes and stored in 70% EtOH at -20°C

until use. Just prior to RNA FISH, slides were dehydrated through an EtOH series and

allowed to air dry. Denatured probes were prehybridized at 37°C for 10 min, applied to

non-denatured slides and hybridized at 37°C for 14-16 hours. Post-hybridization washes

consisted of one 3-minute wash in 50% formamide/2XSSC at 40°C followed by one 2-

minute rinse in 2XSSC/0.1% TX-100 for 1 minute at RT. Slides were then fixed in 4%

paraformaldehyde in PBS for 5 minutes at RT, and briefly rinsed in 2XSSC/0.1% TX-100

at RT. Coverslips were mounted with Prolong Gold antifade plus DAPI (Invitrogen) and

slides were viewed under UV fluorescence (Olympus). Z-stack images were generated

using a Cytovision workstation. After capturing RNA FISH signals, the coverslips were

removed, the slides were dehydrated in an ethanol series, and then processed for DNA

FISH, beginning with the RNase treatment step, as described above.

Chromosome replication timing assay

The BrdU replication timing assay was performed as described previously on

exponentially dividing cultures and asynchronously growing cells (Smith and Thayer,

2012). Mitotic chromosome spreads were prepared and DNA FISH was performed as

described above. The incorporated BrdU was then detected using a FITC-labeled anti-

BrdU antibody (Roche). Coverslips were mounted with Prolong Gold antifade plus DAPI

(Invitrogen), and viewed under UV fluorescence. All images were captured with an

Olympus BX Fluorescent Microscope using a 100X objective, automatic filter-wheel and

Cytovision workstation. Individual chromosomes were identified with either chromosome-

specific paints, centromeric probes, BACs or by inverted DAPI staining. Utilizing the

Cytovision workstation, each chromosome was isolated from the metaphase spread and

a line drawn along the middle of the entire length of the chromosome. The Cytovision

software was used to calculate the pixel area and intensity along each chromosome for

each fluorochrome occupied by the DAPI and BrdU (FITC) signals. The total amount of

fluorescent signal in each chromosome was calculated by multiplying the average pixel

intensity by the area occupied by those pixels. The BrdU incorporation into human

chromosomes containing CRISPR/Cas9 modifications was calculated by dividing the total

incorporation into the chromosome with the deleted chromosome divided by the BrdU

incorporation into the non-deleted chromosome within the same cell. Boxplots were

generated from data collected from 8-12 cells per clone or treatment group. Differences

in measurements were tested across categorical groupings by using the Kruskal-Wallis

test (Kruskal, 1964) and listed as P-values for the corresponding plots.

RESOURCE AVAILABILITY

Further information and requests for resources and reagents should be directed to and
will be fulfilled by the lead contact, Mathew Thayer (thayerm@ohsu.edu)

Materials availability statement

All reagents generated in this study will be made freely available upon request.

Acknowledgments:

M.J.T. was supported by NIH NIGMS (R01GM114162 and R01GM130703).

M.B.H was supported by NIH NCI 4K00CA245677-03

Author contributions:

Declaration of interests

The authors declare no competing interests.

FIGURE 1

FIGURE 2

A 7 kb RBP domain

promoter

sense
or
antisense

B
XIST RNA
ASAR6-141 RNA (sense)

XIST RNA
ASAR6-141 RNA (antisense)

C
Cell#1 Cell#2 Cell#3 Cell#4

ASAR6-141 RNA

Chr5

merge

D ASAR6-141 DNA

E ASAR6-141 DNA BrdU

F ASAR6-141 DNA

G ASAR6-141 DNA BrdU

Figure 3

FIGURE 4

A

CHR1 CHR2 CHR3 CHR4 CHR5

CHR6 CHR7 CHR8 CHR9 CHR10 CHR11
t(3;11)

CHR12 CHR13 CHR14 CHR15 CHR16 CHR17

CHR18 CHR19 CHR20 CHR21 CHR22 CHRX CHRY

B

CHR1 CHR2 CHR3 CHR4 CHR5

CHR6 CHR7 CHR8 CHR9 CHR10 CHR11
t(3;11)

CHR12 CHR13 CHR14 CHR15 CHR16 CHR17

CHR18 CHR19 CHR20 CHR21 CHR22 CHRX CHRY

C

CHR1A   CHR1B

D

CHR4A   CHR4B

F

Chromosome 1   Chromosome 6

Ratio of BrdU incorporation

— HNRNPU TIA/TIAL1 — HNRNPU TIA/TIAL1
shRNA

E

CHR18A   CHR18B

G

G2 DNA
synthesis

CHR6 cen
BrdU
DAPI

H

CHR1 paint
BrdU
DAPI

A
ASAR6 BAC DNA
ASAR6-141 RNA

B
ASAR6-141 RNA
20 minutes HD

C
ASAR6 BAC DNA
ASAR6-141 RNA
20 minutes HD
4hr washout

D
ASAR6 BAC DNA
ASAR6 RNA

E
ASAR6 RNA
20 minutes HD

F
ASAR6 BAC DNA
ASAR6 RNA
20 minutes HD
4hr washout

FIGURE 6

FIGURE 7

A	hC0T1 DNA / hC0T1 RNA / merge	**Untreated**
B	hC0T1 DNA / hC0T1 RNA / merge	**20 min 4% HD**
C	hC0T1 DNA / hC0T1 RNA / merge	**20 min 4% HD/ 4hr wash**
D	hC0T1 DNA / hC0T1 RNA / merge	
E	hC0T1 DNA / hC0T1 RNA / merge	**20 min 4% HD/ 4hr wash +ActD**
F	hC0T1 DNA / hC0T1 RNA / merge	**4hr ActD**

A

FIGURE S1

sgRNA 1 2

rs17070386 5 6

sgRNA 3 4

B SNP analysis of rs17070386 within deleted region (primers 5 plus 6)

HTD114

D6A(7-82-29)

D6B(7-18-1)

CHR6A

D6A(7-82-13)

D6B(7-63-1)

CHR6B

C

HTD114
D6B/6A-1
D6B(7-18-1)
D6B(7-63-1)
D6A(7-82-29)
D6B/6A-2
D6A(7-82-13)
M

Junction PCR
Primers 1 and 4

D

HTD114
D6B/6A-1
D6B(7-18-1)
D6B(7-63-1)
D6A(7-82-29)
D6B/6A-2
D6A(7-82-13)
M

Internal PCR
Primers 5 and 6

FIGURE S2

Chr5 BAC
~7kb RBPD

~7kb RBPD

ChrX paint
~7kb RBPD

FIGURE S4

FIGURE S5

A — ASAR6 RNA / DAPI

B — mChr3 DNA / DAPI

C — merge

D — hCot1 RNA / DAPI

E — ASAR6 DNA / DAPI

F — merge