Colour vision is aligned with natural scene statistics at 4-months of age

Alice E. Skelton[1], Anna Franklin[1], & Jenny M. Bosten[2]*

1 - The Sussex Colour Group, University of Sussex, UK ; 2 – The Sussex Vision Lab, University of Sussex, UK.

* **Correspondence** and requests for materials should be addressed to Jenny Bosten
Email: j.bosten@sussex.ac.uk

Author Contributions:

Conceptualization: AES, AF, JMB

Methodology: AES, AF, JMB

Investigation: AES, AF, JMB

Project administration: AES, AF, JMB

Writing – original draft: AES, AF, JMB

Writing – review & editing: AES, AF, JMB

Competing interests: Authors declare that they have no competing interests.
Data Availability: The data that support the findings of this study are openly available in OSF
[https://osf.io/wx5bt/?view_only=1ac006c69cc449a58329b96be0c7e14e]

Research Highlights

1.   We find infants' hue sensitivity is aligned with the distribution of colours in the natural

world, as it is in adults

2.   At just 4-months, infants' visual system is tailored to extract and represent the

statistical regularities of the natural world

3.   This points to the fundamental drive for the human brain to attempt to optimally

encode by representing statistical regularities even at a young age.

Abstract

Visual perception in adult humans is thought to be optimised to represent the statistical

regularities of natural scenes (e.g. Parraga et al., 2000; Simoncelli, 2003). In infancy and early

childhood, visual input is required to drive visual development (Berardi et al., 2000;

Blakemore & Cooper, 1970; Jandó et al., 2012; Knudsen, 2004), but whether or not scene

statistics are extracted and represented colour is currently unknown. In adults, visual

sensitivity to different hues shows an asymmetry, which relates to the statistical regularity of

colour in the natural world (Bosten et al., 2015). We measured colour discrimination in

infants to investigate whether or not the visual system can in very early life represent scene

statistics. Our results reveal the earliest association between vision and natural scene

statistics that has yet been found: even as young as four months of age, colour vision is

aligned with the distributions of colours in natural scenes.

Introduction

Human perception is optimised   in many ways to extract information from the

environments that we find ourselves in. Process to calibrate perception to the contents of

natural environments occur over different time scales, both over evolutionary history

(Changizi et al., 2006; Osorio & Vorobyev, 1996; Regan et al., 2001), and within the lifetime of

the individual (Schefrin & Werner, 1990). Some calibrative processes undoubtedly occur

during infancy: for example, face (Kelly et al., 2007), speech (Kuhl et al., 2006) and music

(Trehub & Hannon, 2006) perception are refined during infancy to show biases for the stimuli

encountered in infants' individual environments. In mature perceptual systems, there is

tuning to the statistical regularities of visual scenes: visual channels are thought to be tuned

to extract the regularities of spatial frequency (Hansen & Hess, 2006; Knill et al., 1990;

Tadmor & Tolhurst, 1994), orientation (Hansen & Essock, 2004), and attributes of colour

(Long et al., 2006) in natural scenes. Relatively little is known about the developmental

trajectory of visual representations of the statistical regularities of natural scenes. Ellemberg

et al., 2012 have found that representations of the spatial frequency structure of natural

scenes are present in 10 year-olds but undeveloped in 6- and 8- year-olds. The only existing

work on infants finds that at 9-10 months, but not 3-6 months, infants' looking preference

and their Event Related Potential components (P100 and P400) appear to distinguish

between natural textures and one type of synthetic manipulation of those textures (Balas &

Conlin, 2014; Balas & Woods, 2014). One interpretation of this work is that infants tune to

natural texture statistics by 9-months. However, the lack of a consistent effect of another

type of natural texture manipulation (contrast polarity), and the potential role of other low-

level visual differences independent of naturalness suggests that further work is needed to

clarify the nature of these effects.

Colour vision is a good example of a perceptual system that is optimised at several different

levels to extract information from natural scenes. Human colour vision relies on three classes

of retinal receptors, long- (L), medium- (M) and short- (S) wavelength sensitive cones. The

particular sensitivities of the human cones are thought to have evolved to optimally

represent colour information relevant to the diets or skin colours of our primate ancestors

(Changizi et al., 2006; Osorio & Vorobyev, 1996; Regan et al., 2001). Postreceptorally, colour

is processed predominantly in two chromatically opponent channels, which themselves seem

to be optimised to minimise the redundancy of information acquired about natural scenes

(Ruderman et al., 1998). There has been recent interest in a third level of alignment with

natural scene statistics: colour discrimination is poorest along a blue-yellow colour axis

(Honjyo & Nonaka, 1970; Krauskopf & Karl, 1992; Pearce et al., 2014; Álvaro et al., 2017)

which is intermediate between the colour axes along which the two chromatically opponent

retinogeniculate colour mechanisms are tuned (the vertical and horizontal axes in Figure 1a).

It is the blue-yellow axis that contains most variance in natural scenes (Bosten et al., 2015;

Webster & Mollon, 1997). See figure 1b & c for an example of the variation found in natural
scenes.

Figure 1 a) Our version of the MacLeod Boynton chromaticity diagram (Macleod & Boynton,
1979), using the Stockman, MacLeod and Johnson cone fundamentals (Stockman et
al., 1993) representing the signals carried by the chromatically opponent
retinogeniculate colour mechanisms, L/(L+M) and S/(L+M). The blue-yellow axis is
intermediate to the cardinal axes, indicated by the dashed line. b) An example of a
hyperspectrally imaged natural scene (Nascimento et al., 2002), and c) A histogram
showing the distribution of chromaticities of the natural scene in (b) plotted in the
MacLeod Boynton chromaticity diagram. The largest variance in chromaticities is
along the negative diagonal. d) Infant and e) adult target chromaticities along the 8
individual axes presented in the MacLeod Boynton chromaticity diagram. The hue
angles of the cardinal axes are indicated in (e). Towards the centre of the figure,
colours are desaturated, increasing in saturation radially along the axes. f) An

example target (upper panel) displayed within 6.9° of the participant's current fixation

point. If the target was subsequently fixated for a minimum of 160ms, it was replaced

by a reward smiling face (lower), a sound was played, and a hit was recorded.

In the present study we investigated the development of representations of the distribution

of colours in natural scenes by measuring colour discrimination in infants using a novel task

(see methods). We chose infants aged 4-6 months, because it is thought that the S-cone

reliant colour-opponent mechanism develops relatively late at 2-3 months (Bornstein, 1975;

Brown & Teller, 1989; Knoblauch et al., 2001; Suttle et al., 2002), and it is only after this point

that infants have trichromatic colour vision. For 60 infants and 19 adults we measured eye

movements to chromatic targets varying across 8 levels of saturation in 8 hue directions

(Figure 1(d) and (e)).

For each hue, when rate of detection is plotted against target saturation, a colour

discrimination threshold can be defined at a certain performance criterion (**alpha K**) along

the resulting psychometric function, which was fit using a local linear 'model-free' algorithm

(Zychaluk & Foster, 2009) (Figure 2a). Through thresholds for each of the eight colours we fit

colour discrimination ellipses, where the distance from the centre of the ellipse in each hue

direction is the discrimination threshold in a version of the MacLeod-Boynton chromaticity

diagram (Macleod & Boynton, 1979) based on the Stockman, MacLeod and Johnson cone

fundamentals (Stockman et al., 1993). We chose the MacLeod-Boynton diagram as a colour

metric because it represents along its cardinal axes activity in the two dominant post-

receptoral retinogeniculate colour mechanisms (Webster & Mollon, 1991). If colour

discrimination is limited by noise in either of these cardinal colour mechanisms, the resulting

discrimination ellipse would be oriented either horizontally or vertically. An off-axis bias in

colour discrimination, such as a blue-yellow bias, means that discrimination must be

determined by noise in non-cardinal colour mechanisms(Bosten et al., 2015).

Figure 2. (a) Example psychometric function, and psychometric functions for infants and

adults for the 8 colour axes tested, all showing the probability of target detection as a

function of stimulus intensity (saturation). The example psychometric function

demonstrates how different alpha criteria result in different threshold measurements

along the same psychometric function. For the infant and adult data, each individual

graph shows group average psychometric functions for a specific hue axis, which is a

radial line in the MacLeod Boynton chromaticity diagram (Figure 1 (d) and (e)), noted

by the angle of the hue axis in the bottom left of each figure. (b) For each of a range

of alphas, 8 colour discrimination thresholds were extracted (one for each hue), and

then an ellipse was fit to each set of 8 thresholds plotted in the MacLeod-Boynton

chromaticity diagram. Separate discrimination ellipses were plotted for the 3 groups:

two infant groups (younger and older than the infant sample median age) and adults.

The figures show ellipses for 3 example alphas. (c) In order to quantify the extent of

the blue-yellow bias of each colour discrimination ellipse, an 'axis ratio' was

calculated. This is the ratio of the length of the negative diagonal of the ellipse to the

length of the positive diagonal, when the ellipse is plotted in a normalised version of

the MacLeod Boynton chromaticity diagram, scaled to equate variances in the x and y

directions. In both 3(b) and 3(c), ellipses fitted from thresholds extracted at lower

alphas are shown lighter in colour, and those from higher alphas are shown darker in

colour. (d) Axis ratios are plotted as a function of alpha with bootstrapped 95%

confidence intervals for each of the 3 groups.

Results

For our primary analysis we fit discrimination ellipses for three groups, two infant samples

split above and below the median age (younger infants: N=30, M_{age}= 4.55 months, SD=0.27,

and older infants: N=30, M_{age}= 5.82 months, SD=0.51), and adults. From the fitted ellipses we

quantified the asymmetry in colour discrimination by taking an *axis ratio*. This is the ratio of

the length of the ellipse along the negative diagonal to the length along the positive diagonal,

in a normalised colour space in which the variances along the x and y axes are equated

(Bosten et al., 2015). If the axis ratio is greater than 1, then colour discrimination is relatively

poor in a blue-yellow colour direction, aligned with the axis of maximum variability of the

colours in natural scenes.

For each of the three age groups and for each hue, we achieved very clear group

psychometric functions (Figure 2a). One obvious difference between the groups is in the

positions of asymptotes of the psychometric functions. For adults, performance quickly

asymptotes close to 100% detection. For younger infants the asymptotes are at about 30-

40% and for older infants they are at about 50-60%. This large difference means that any

particular performance criterion (alpha) would fall on very different parts of the

psychometric function for the different groups. This prompted us to extract thresholds at

different alphas, and to fit different discrimination ellipses to each set of thresholds at

different alphas (Jones et al., 2015; Trehub et al., 1980). Figure 2(d) shows the results of this

analysis: for all three age groups, axis ratio is large at low alphas and then decreases as a

function of alpha.

The large axis ratios show that for all three age groups colour discrimination is selectively

poor in the blue-yellow colour direction, as would be expected if colour vision is aligned with

the statistics of natural scenes. Figure 2(d) shows axis ratio as a function of threshold alpha.

For infants, axis ratios are large at low alphas (1.89 for younger infants at alpha=0.08; 2.36 for

older infants at alpha=0.08), but as alpha increases they tend towards 1. Adults show the

same decrease in axis ratio with alpha as infants, but maintain an axis ratio larger than one at

all alphas. We conducted similar analyses at the level of individual participants for all infants

as one group and all adults as another group (for details see SI *Individual participant*

analysis). Here, we found that there is a significant effect of alpha on axis ratio for both

infants (using a Skillings-Mack test (Chatfield & Mander, 2009), a non-parametric Friedman

type statistic which can be used when there are random data missing as is the case in the

current study where the number and type of trials completed by each infant varies,

T(18)=102.57, *p*=<.001) and adults (Skillings-Mack, T(18)=59.79, *p*=<0.001).

Discussion

The strong dependence of axis ratio on alpha that we observe for infants implies that the

psychometric function describing performance as a function of saturation has both a higher

mean and steeper slope for blue and yellow targets than for red and green targets. This does

not fit the usual expectation, where the noise that limits detection tends to increase the

mean (determined by signal:noise ratio) and *reduce* the slope(Arazi et al., 2017; Legge et al.,

1987). One possible explanation for the dependency of axis ratio on alpha is that non-visual

processes contribute differentially to visual performance at different alphas. Specifically, it is

generally assumed that the slope of the psychometric function is determined by visual signals

and sensory noise, while attentional 'lapses' affect only its upper asymptote(Swanson &

Birch, 1992; Wichmann & Hill, 2001). However, especially for infants, we propose that this is

an unsafe assumption. Instead, there may be an interaction between visual and non-visual

contributions to psychophysical decisions so that both the positively sloping region and the

upper asymptote are influenced by non-visual factors such as attention and motivation.

Importantly, the non-visual factors may have their greatest influence at high signal strengths.

In the current study, an infant may be more motivated (e.g. by attentional salience) to make

an eye movement towards a coloured stimulus with greater saturation than one with lower

saturation. Thus, the lower reaches of the psychometric function at lower stimulus strengths

may be more singularly determined by visual sensitivity than the upper reaches, where the

presence or absence of attention and motivation may have a relatively greater influence on

the probability of detection. This may explain why axis ratio is largest at low stimulus

strengths, where visual sensitivity (tuned to natural scene statistics) is having its purest

influence on detection performance.

Our results show that like adults, infants as young as 4 months of age show a bias in colour

discrimination in a direction predicted from the distributions of colours in natural scenes.

Though there is a significant improvement in colour discrimination with age (see SI

Development of Colour Sensitivity), the discrimination ellipses show no evidence for

development in the degree of bias (Figure 2b). Our results are compatible with two schemes:

either the blue-yellow bias in colour discrimination is innate, perhaps calibrated to natural

scene statistics neurogenetically, or it develops very early by calibration to the statistics of

natural scenes, between the ages of 2-3 months (when trichromacy first develops) and 4

months (the age of the youngest infants in our sample).

Our finding that bias in colour discrimination is a function of threshold criterion (alpha) for all

three age groups has important implications for psychophysics. The choice of alpha in

psychophysical analysis is arbitrary, but there are conventions that are often applied. For

example, alpha is often defined half way between the guess rate and the asymptote of the

psychometric function (Watson & Pelli, 1983). Our finding implies that the choice of alpha

may have a substantial impact on the conclusions of psychophysical studies, and that actually

a lower alpha than is typically chosen may better isolate the sensory processes (at least in

infants) that psychophysicists are interested in.

The relationship between axis ratio and alpha that we have observed would not have been

apparent had we not measured full psychometric functions. Typically, in infant studies,

discrimination performance is measured using a variant of the 'preferential looking'

technique, where gaze durations in the presence and absence of stimulus targets are

compared statistically. This often results in one number to decide whether a given target is

above or below threshold (seen versus not seen). Discrimination thresholds measured using

preferential looking for infants of different ages (Knoblauch et al., 2001) but owing to the

time-consuming nature of preferential looking, are available for only a limited number of

colour axes (see SI *Development of Colour Sensitivity*). Our method of using eye movements

(see also (Jones et al., 2014)) is more efficient, allowing measurement of performance at a

greater number of points and full psychometric functions, as is good practice for

psychophysics in adults. Our efficient measure enables infants' colour discrimination ellipses

to be revealed for the first time.

Our results provide the first evidence that post-receptoral vision is aligned to natural scene

statistics in early infancy. A major finding in colour vision science is that the L-, M- and S-

cones are optimally spectrally positioned to extract colour information from the natural

world. If the alignment that we see in infants is achieved by neurogenetic calibration, then

there must be genetic optimisation post-receptorally as well. If the aligning process is

ontogenetic, then the infant brain must be able to extract and calibrate to statistical

information within a matter of months. This revises previous research that has demonstrated

statistical optimisation of perception only later in infancy for social or linguistic stimuli. To

distinguish these theoretically important accounts, future research will need to apply our

methods to genetically similar infants living in radically different visual environments.

Methods

Participants

Sixty four infants (age range= 4.01-6.80 months, mean age= 5.18 months, SD= 0.76 months) took part in the experiment. Four infants were excluded from the study due to fussiness preventing them from proceeding past the initial calibration phase. All participants were born at 40 +/- 2 weeks, met a minimum birth weight of 2500g, had no known visual deficits, and no history of epilepsy, and reported no family history of colour vision deficiency.

Nineteen adult participants were recruited by experimenters (13 female, mean age= 26.42 years). Participants had normal colour vision as assessed by the Ishihara pseudoisochromatic plates (38 plate edition, (Ishihara, 1936)) and the Lanthony tritan album (Lanthony, 1987).

Stimuli and apparatus

Target stimuli were coloured circles that subtended a visual angle of 6.9° on a surround metameric with equal energy white that subtended a visual angle of 41°. The chromatic difference between target and surround was isoluminant for the standard observer (Stockman et al., 1993). In order to prevent individual participants detecting targets via residual luminance differences with the surround, low-pass luminance noise was added over the whole display. The mean luminance of both target and surround was 28 cdm^{-2}, and luminance noise was linearly distributed to range between ±10% of the mean luminance. The chromaticities of the target stimuli varied in saturation along 8 axes within a version the MacLeod-Boynton (1979) chromaticity diagram (Macleod & Boynton, 1979) based on the Stockman, MacLeod and Johnson (1993) cone fundamentals (Stockman et al., 1993). The centre of our stimulus space was a point metameric with equal energy white (the background chromaticity), and there were 8 points per axis (64 points in total) increasing in saturation away from the centre. Four of the axes were the cardinal directions in the MacLeod-Boynton chromaticity diagram; two axes were in the blue-yellow direction (the cerulean line defined from 476 to 576 nm(Mollon, 2006)), and two axes were a reflection of this line through the vertical axis of the MacLeod-Boynton chromaticity diagram.

For adults, stimuli were spaced logarithmically along these axes, to focus the collection of data closest to where adult threshold levels were expected to be. For infants stimuli were sampled in equal steps along each axis (see main text Figure 1). Due to infants' higher

discrimination thresholds (Knoblauch et al., 2001) stimuli were sampled from a wider range

of chromaticities than adults.

Stimuli were presented on a Diamond Pro 2070SB CRT Monitor (Mitsubishi, Tokyo, Japan) via

a PC-driven ViSaGe MKII Stimulus Generator (Cambridge Research Systems Ltd, Rochester,

UK). Eye-movement data was recorded via an Eyelink 1000 system (SR Research, Ontario,

Canada), and the experimental programme was run in MATLAB (R2016a, The MathWorks

Inc., Natick, MA). The system was gamma-corrected using a LS-100 luminance meter (Konica-

Minolta, Tokyo, Japan), and the colour calibration achieved using a spectrascan PR655

spectroradiometer (Photo Research Inc., Chatsworth, CA).

Procedure

Infant participants were seated in a car seat mounted in a chair 60cm from the

screen, and watched a cartoon on screen during camera set up, and then completed a 9-

point spatial calibration. If necessary (e.g. if a participant moved a lot), participant eye

movements were recalibrated during testing.

Before each trial, participants were presented with a blank screen of luminance noise for

250ms. Targets appeared within 6.9° of the participants' current fixation point on the display.

If participants fixated the target (or a 1° area around the target) for 160ms, this was recorded

as a successful fixation (a 'hit'), the target was replaced with an image of a smiley face, and a

short musical tune was played. If the target was not fixated within 2250ms, no auditory or

visual reward was played, and a miss was recorded (see figure *Methods.1*).

Figure Methods. 1) Example trial procedure. Participants were presented with a blank screen
of luminance noise for 250ms before a target was presented. If fixated for a minimum
of 160ms, the target was replaced by a smiley face, and a was noise presented, and a
hit was recorded for the trial. After another blank display of luminance noise, the next
target was displayed and was not fixated within 2250ms, no reward was presented,
and the next trial started.

If participants' gaze left the screen, a looming black and white spiral accompanied by a loud

noise was played to refocus infants' visual attention to the screen. Once the participant was

centrally fixated, the experimenter pressed a key to allow trials to start again.

For adult participants, trials were presented as 10 blocks of 64 trials (total 640 trials), the

testing duration was approximately 30 minutes. During each block of trials, each chromaticity

was presented once, in a random order, resulting in each chromaticity being presented 10

times to each participant. Participants were told that their task was to continually search for

and look at targets on the screen.

In order to encourage maximum engagement with the task for infants, informed by pilot

testing (N=4), 3/10 trials presented a maximally saturated stimulus. Stimuli were presented in

blocks of 64, and testing was continued until the infant became disengaged with the task.

References

Arazi, A., Censor, N., & Dinstein, I. (2017). Neural variability quenching predicts individual perceptual
abilities. *Journal of Neuroscience*, *37*(1), 97-109.
Balas, B., & Conlin, C. (2014). Early visual ERP components are sensitive to natural texture
appearance. *Journal of Vision*, *14*(10), 1431-1431.
Balas, B., & Woods, R. (2014). Infant preference for natural texture statistics is modulated by
contrast polarity. *Infancy*, *19*(3), 262-280.
Berardi, N., Pizzorusso, T., & Maffei, L. (2000). Critical periods during sensory development. *Current
opinion in neurobiology*, *10*(1), 138-145.
Blakemore, C., & Cooper, G. F. (1970). Development of the brain depends on the visual environment.
Nature, *228*(5270), 477.
Bornstein, M. H. (1975). Qualities of color vision in infancy. *Journal of experimental child psychology*,
19(3), 401-419.
Bosten, J., Beer, R., & MacLeod, D. (2015). What is white? *Journal of Vision*, *15*(16), 5-5.
Brown, A. M., & Teller, D. Y. (1989). Chromatic opponency in 3-month-old human infants. *Vision
Research*, *29*(1), 37-45.
Changizi, M. A., Zhang, Q., & Shimojo, S. (2006). Bare skin, blood and the evolution of primate colour
vision. *Biology letters*, *2*(2), 217-221.

Chatfield, M., & Mander, A. (2009). The Skillings–Mack test (Friedman test when there are missing
data). *The Stata Journal*, *9*(2), 299.
Ellemberg, D., Hansen, B. C., & Johnson, A. (2012). The developing visual system is not optimally
sensitive to the spatial statistics of natural images. *Vision Research*, *67*, 1-7.
Hansen, B. C., & Essock, E. A. (2004). A horizontal bias in human visual processing of orientation and
its correspondence to the structural components of natural scenes. *Journal of vision*, *4*(12),
5-5.
Hansen, B. C., & Hess, R. F. (2006). Discrimination of amplitude spectrum slope in the fovea and
parafovea and the local amplitude distributions of natural scene imagery. *Journal of Vision*,
6(7), 3-3.
Honjyo, K., & Nonaka, M. (1970). Perception of white in a 10° field. *JOSA*, *60*(12), 1690-1694.
Ishihara, S. (1936). Series of plates designed as tests for colour-blindness.
Jandó, G., Mikó-Baráth, E., Markó, K., Hollódy, K., Török, B., & Kovacs, I. (2012). Early-onset
binocularity in preterm infants reveals experience-dependent visual development in
humans. *Proceedings of the National Academy of Sciences*, *109*(27), 11049-11052.
Jones, P., Kalwarowsky, S., Atkinson, J., Braddick, O., & Nardini, M. (2014). Automated Measurement
of Resolution Acuity in Infants Using Remote Eye-Tracking [Article]. *Investigative
Ophthalmology & Visual Science*, *55*(12), 8102-8110. https://doi.org/10.1167/iovs.14-15108
Jones, P. R., Kalwarowsky, S., Braddick, O. J., Atkinson, J., & Nardini, M. (2015). Optimizing the rapid
measurement of detection thresholds in infants. *Journal of Vision*, *15*(11), 2-2.
Kelly, D. J., Quinn, P. C., Slater, A. M., Lee, K., Ge, L., & Pascalis, O. (2007). The other-race effect
develops during infancy: Evidence of perceptual narrowing. *Psychological Science*, *18*(12),
1084-1089.
Knill, D. C., Field, D., & Kerstent, D. (1990). Human discrimination of fractal images. *JOSA A*, *7*(6),
1113-1123.
Knoblauch, K., Vital-Durand, F., & Barbur, J. (2001). Variation of chromatic sensitivity across the life
span [Article]. *Vision Research*, *41*(1), 23-36. https://doi.org/10.1016/S0042-6989(00)00205-
4
Knudsen, E. I. (2004). Sensitive periods in the development of the brain and behavior. *Journal of
cognitive neuroscience*, *16*(8), 1412-1425.
Krauskopf, J., & Karl, G. (1992). Color discrimination and adaptation. *Vision research*, *32*(11), 2165-
2175.
Kuhl, P. K., Stevens, E., Hayashi, A., Deguchi, T., Kiritani, S., & Iverson, P. (2006). Infants show a
facilitation effect for native language phonetic perception between 6 and 12 months.
Developmental science, *9*(2), F13-F21.
Lanthony, P. (1987). A tritan album. In *Colour Vision Deficiencies VIII* (pp. 141-143). Springer.
Legge, G. E., Kersten, D., & Burgess, A. E. (1987). Contrast discrimination in noise. *JOSA A*, *4*(2), 391-
404.
Long, F., Yang, Z., & Purves, D. (2006). Spectral statistics in natural scenes predict hue, saturation,
and brightness [Article]. *Proceedings of the National Academy of Sciences of the United
States of America*, *103*(15), 6013-6018. https://doi.org/10.1073/pnas.0600890103
Macleod, D., & Boynton, R. (1979). Chromaticity diagram showing cone excitation by stimuli of equal
luminance [Letter]. *Journal of the Optical Society of America*, *69*(8), 1183-1186.
https://doi.org/10.1364/JOSA.69.001183
Mollon, J. (2006). Monge: the verriest lecture, Lyon, July 2005. *Visual Neuroscience*, *23*(3-4), 297-
309.
Nascimento, S. M., Ferreira, F. P., & Foster, D. H. (2002). Statistics of spatial cone-excitation ratios in
natural scenes. *JOSA A*, *19*(8), 1484-1490.
Osorio, D., & Vorobyev, M. (1996). Colour vision as an adaptation to frugivory in primates. *Proc. R.
Soc. Lond. B*, *263*(1370), 593-599.

Parraga, C., Troscianko, T., & Tolhurst, D. (2000). The human visual system is optimised for
processing the spatial information in natural visual images [Article]. *Current Biology*, *10*(1),
35-38. https://doi.org/10.1016/S0960-9822(99)00262-6

Pearce, B., Crichton, S., Mackiewicz, M., Finlayson, G. D., & Hurlbert, A. (2014). Chromatic
illumination discrimination ability reveals that human colour constancy is optimised for blue
daylight illuminations. *PloS one*, *9*(2), e87989.

Regan, B. C., Julliot, C., Simmen, B., Vienot, F. o., Charles–Dominique, P., & Mollon, J. D. (2001).
Fruits, foliage and the evolution of primate colour vision. *Philosophical Transactions of the
Royal Society of London B: Biological Sciences*, *356*(1407), 229-283.

Ruderman, D. L., Cronin, T. W., & Chiao, C.-C. (1998). Statistics of cone responses to natural images:
implications for visual coding. *JOSA A*, *15*(8), 2036-2045.

Schefrin, B. E., & Werner, J. S. (1990). Loci of spectral unique hues throughout the life span. *Journal
of the Optical Society of America A*, *7*(2), 305-311. https://doi.org/10.1364/JOSAA.7.000305

Simoncelli, E. (2003). Vision and the statistics of the visual environment [Review]. *Current Opinion in
Neurobiology*, *13*(2), 144-149. https://doi.org/10.1016/S0959-4388(03)00047-3

Stockman, A., Macleod, D., & Johnson, N. (1993). Spectral sensitivities of the human cones [Review].
Journal of the Optical Society of America a-Optics Image Science and Vision, *10*(12), 2491-
2521. https://doi.org/10.1364/JOSAA.10.002491

Suttle, C. M., Banks, M. S., & Graf, E. W. (2002). FPL and sweep VEP to tritan stimuli in young human
infants. *Vision research*, *42*(26), 2879-2891.

Swanson, W. H., & Birch, E. E. (1992). Extracting thresholds from noisy psychophysical data.
Perception & psychophysics, *51*(5), 409-422.

Tadmor, Y., & Tolhurst, D. (1994). Discrimination of changes in the second-order statistics of natural
and synthetic images. *Vision research*, *34*(4), 541-554.

Trehub, S. E., & Hannon, E. E. (2006). Infant music perception: Domain-general or domain-specific
mechanisms? *Cognition*, *100*(1), 73-99.

Trehub, S. E., Schneider, B. A., & Endman, M. (1980). Developmental changes in infants' sensitivity to
octave-band noises. *Journal of Experimental Child Psychology*, *29*(2), 282-293.

Watson, A., & Pelli, D. (1983). QUEST - A Baysian adaptive psychometric method [Article]. *Perception
& Psychophysics*, *33*(2), 113-120. https://doi.org/10.3758/BF03202828

Webster, M., & Mollon, J. (1991). Changes in color appearance following post-receptoral adaptation
[Article]. *Nature*, *349*(6306), 235-238. https://doi.org/10.1038/349235a0

Webster, M. A., & Mollon, J. (1997). Adaptation and the color statistics of natural images. *Vision
research*, *37*(23), 3283-3298.

Wichmann, F. A., & Hill, N. J. (2001). The psychometric function: I. Fitting, sampling, and goodness of
fit. *Perception & psychophysics*, *63*(8), 1293-1313.

Zychaluk, K., & Foster, D. (2009). Model-free estimation of the psychometric function [Article].
Attention Perception & Psychophysics, *71*(6), 1414-1425.
https://doi.org/10.3758/APP.71.6.1414

Álvaro, L., Linhares, J. M., Moreira, H., Lillo, J., & Nascimento, S. M. (2017). Robust colour constancy
in red-green dichromats. *PloS one*, *12*(6), e0180310.

Acknowledgements: We thank the infant participants and their parents, Ken Knoblauch, Daniel Osario, and Don

MacLeod for helpful discussion. This research was funded by a European Research Council Grant project,

"COLOURMIND," 772193 (to A.F).

Competing interests: The authors report no conflict of interest.

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Jenny Bosten, j.bosten@sussex.ac.uk

437