Is the production of reactive oxygen and nitrogen species by macrophages associated with better infectious control in the experimental disseminated and pulmonary mucormycosis?

Amanda Ribeiro dos Santos [1], Thais Fernanda Fraga-Silva [2], Débora de Fátima Almeida Donanzam [1], Angela Carolina Finatto [3], Camila Marchetti [3], Maria Izilda Andrade [4], Olavo Speranza de Arruda [3], Maria Sueli Parreira de Arruda [3], James Venturini [1] *.

[1] Universidade Federal de Mato Grosso do Sul (UFMS), Faculdade de Medicina, Campo Grande, MS, 79070-900, Brazil.

[2] Universidade de São Paulo, Departamento de Bioquímica e Imunologia, Escola de Medicina de Ribeirão Preto, SP, 14049-900, Brazil.

[3] Universidade Estadual Paulista (Unesp), Faculdade de Ciências, Bauru, SP, 17033-360, Brazil.

[4] Lauro de Souza Lima Institute, Bauru, SP, 17034-971, Brazil.

*Corresponding author: Prof. Dr. James Venturini, Cidade Universitária, Unit 9, 79070-900, Campo Grande, MS, Brazil, Phone/Fax: +55 (67) 3345-7370, james.venturini@ufms.br

Abstract

Different levels of resistance against *Rhizopus oryzae* infection have been observed between inbred (BALB/c) and outbred (Swiss) mice, with is associated with the genetic background of each mouse strain. Considering that macrophages play an important role in host resistance to *Rhizopus* species, we use the different infectious outcomes observed in experimental mucormycosis to identify the most efficient macrophages responses pattern against *R. oryzae in vitro* and *in vivo*. For this, we compared BALB/c and Swiss macrophage activity pre-and-post intravenous or intratracheal *R. oryzae* infections. Production of hydrogen peroxide (H_2O_2) and nitric oxide (NO) was determined in cultures of peritoneal (PMΦ) or alveolar macrophages (AMΦ) challenged, or not, with heat-killed spores of *R. oryzae*. Levels of TNF-α and IL-10 were also measured to enhance our findings. Naïve PMΦ from BALB/c increased the production of H_2O_2, TNF-α, and IL-10 in the presence of heat-killed spores of *R. oryzae,* while naïve PMΦ from Swiss mice was less responsive. Naïve AMΦ from two strains of mice were less reactive to heat-killed spores of *R. oryzae* than PMΦ. On 30 days of *R. oryzae* intravenous infection, lower fungal load in BALB/c strain of mice was accompanied by higher production of H_2O_2 by PMΦ when compared with Swiss mice. Differently, AMΦ from BALB/c mice showed higher production of NO, TNF-α, and IL-10 after 7 days of intratracheal infection and after 30 days, lower fungal load, when compared with Swiss mice. According to the set of experiments performed, our findings reveal that independently of mice strain, PMΦ is more reactive against *R. oryzae* in the first contact than AMΦ. In addition, increased PMΦ production of H_2O_2 at the end of disseminated infection is related to efficient fungal clearance observed in resistant (BALB/c). Our findings provide new evidence to understand the parasite-hosts relationship in mucormycosis.

1. Introduction

Mucormycosis is a devastating invasive fungal infection predominantly caused by *Rhizopus* species (1). It has emerged globally as a public health threat during the COVID-19 pandemic (2–4). The infection occurs via diverse routes such as inhalation, percutaneous, or ingestion of spores. Mucormycosis may have a wide variety of disease manifestations, with rhinocerebral and pulmonary diseases being the most common (5). The mortality rate varies from 46%-70% and up to 90% upon dissemination (6–8).

Risk factors for mucormycosis include diabetes mellitus, neutropenia, sustained immunosuppressive therapy, chronic prednisone use, and iron chelation therapy (7). During the COVID-19 pandemic, the use of immunomodulatory drugs (e.g., systemic corticosteroids and tocilizumab) and COVID-19-induced immune dysregulation increased the risk of mucormycosis (9,10). Cases of COVID-19–associated mucormycosis (CAM) have been reported worldwide (11–20), and have been mostly associated with underlying uncontrolled diabetes mellitus diseases. Nevertheless, cutaneous and soft-tissue mucormycosis has been observed in immunocompetent individuals (21,22), such as those induced by traumatic implantation of contaminated soil and water during tornadoes (23).

Considering the high rates of mortality and morbidity associated with this life-threatening disease (14), it has a prognosis and outcome that have not significantly improved over the last decades, mainly due to the difficulty in forming an early diagnosis and the limited activity of current antifungal agents against Mucorales (24,25). In addition, mucormycosis has an incompletely understood pathogenesis, particularly related to the parasite-hosts relationship.

Experimental models of mucormycosis are widely used for the evaluation of antifungal therapy and, in these studies, immunosuppression is used to induce fungal dissemination (26–29). Nevertheless, the effect of immunosuppressive drugs restricts the evaluation of the immunological mechanisms involved in fungal resistance. Recently, our group has focused on studies exploring *Rhyzopus*-host interplay using immunocompetent models, inducing both disseminated and pulmonary mucormycosis (30,31). Using BALB/c and Swiss mice, we determined a resistant and less resistant model of mucormycosis, respectively. While BALB/c mice has been shown better capacity in decreasing fungal load until 30 days of intravenous and intratracheal infection, Swiss mice has been shown be less responsive to *R. oryzae* infectious and consequently more prolonged viable fungal presence in internal organs (30,31). We and others (32,33) have highlighting the importance of evaluating immune response in immunocompetent mice to understand the mechanisms involved in the resistance to Mucorales agents, as well as, the mucormycosis pathogenesis in immunocompetent individuals (21,22).

Andrianaki and collaborators (2018) revealed the essential role of *Rhizopus*–macrophage interplay for pulmonary mucormycosis. Using an immunocompetent model, it was shown that the development of mucormycosis is related to the prolonged intracellular survival of the fungus inside of the macrophages (33). Considering the poor knowledge of macrophage activity during mucormycosis in the context of natural resistance; in the present study, we employed resistant and less resistant models to determine the reactive oxygen and nitrogen species production by *R. oryzae*-infected macrophages

2. Material and methods

2.1 Mice. Two-month-old female inbred BALB/c and outbred Swiss mice from the

Animal House at the Laboratório de Imunopatologia Experimental (LIPE) of UNESP

(Univ Estadual Paulista, Bauru, SP, Brazil) were divided into groups randomly. We

provide food and water sterile *ad libitum*. This study was carried out in strict accordance

with the recommendations in the Guide for the Care and Use of Laboratory Animals of

the National Institutes of Health and Brazilian College of Animal Experimentation. The

study was approved by an Institutional Animal Care and Use Committee (IACUC)

(Protocol Number: 1608/46/01/2013-CEUA-FC) by Animal Experimentation Ethics

Committee of the School of Sciences of Bauru, UNESP. All surgery was performed under

ketamine/xylazine anesthesia, and all efforts were made to minimize suffering.

2.2 Fungal strains. *R. oryzae* (IAL 3796) was previously obtained from the fungal

collection of Instituto Lauro de Souza Lima (ILSL), and submitted to species

identification by Adolfo Lutz Institute (São Paulo, SP, Brazil). The fungi were maintained

by monthly subculturing on Sabouraud dextrose agar (SDA) slants (Difco Laboratories,

Detroit, Michigan, USA).

2.3 Experimental design. Mice were randomly separated into two main groups

according to the route of inoculation: intravenous groups (*Rhi*-IV*)* and intratracheal

groups (*Rhi*- IT). For the *Rhi*-IV groups, Swiss and BALB/c mice were inoculated within

3.0×10^4 viable spores of *R. oryzae* in the caudal vein. For the *Rhi*- IT groups, Swiss and

BALB/c mice were inoculated within 2.0×10^6 viable spores of *R. oryzae* in the trachea.

Next, groups of six *R. oryzae*-infected mice were evaluated on days 7 and 30 post-

infection (p.i.). Non-infected groups were composed of BALB/c and Swiss mice

inoculated with sterile saline solution (SSS) by intravenous or intratracheal route.

2.4 Fungal infection. Fungi were washed carefully with SSS, and the suspension was

mixed twice for 10 s on a vortex-mixer and decanted off for 5 min. The supernatants were

collected and washed twice. We determined fungal viability by cotton blue staining. In

Rhi-IVgroups, a volume of 100 μL of fungal suspension (3×10^4 spores of *R. oryzae*) was

inoculated into the lateral tail vein. In *Rhi*-IT groups, mice were previously anesthetized

ketamine and xylazine (80 and 10 mg/kg body weight, respectively) by intraperitoneal

route. After the tracheal exposition, each mouse received a volume of 40 μL (2×10^6

spores of *R. oryzae*) of the suspension. The incision was sutured with surgical thread, and

the mice were kept in a warm place and observed for their recovery process.

2.5 Collection of the biological material. Mice were anesthetized with isoflurane then

euthanized via CO_2 asphyxiation. Peritoneal lavage (PL) and bronchoalveolar lavage

(BAL) was performed in *Rhi*- IV and *Rhi*-IT groups, respectively, using cold and sterile

phosphate-buffered saline (PBS). Fragments of the brain, liver, lung, spleen, and kidneys

were collected and submitted to microbiological evaluation.

2.6 Recovery of viable fungi. Quantitative colony culture is a wide method for organ

fungal burden determination in per gram tissue in which whole tissue is grinding into

suspension before inoculating onto culture plates. However this methods causes non-

viability in Mucorales (21), and to avoid damage to fragile hyphal and consequent false-

negative results in our quantitative analysis (34), the fragment method was employed in

our set of experiments, as previously reported (30,31,35). For this, ten fragments (2 x 2

mm) of the brain, liver, lung, spleen, and kidneys were cultured on Sabouraud agar plates

at 25ºC for a maximum of seven days. We counted fragments of fungal growths and

expressed the results as the frequency of *R. oryzae*-positive fragments per total of

cultivated.

2.7 Macrophages cell culture. BAL and PL suspensions were centrifuged for 10 minutes

at 410 *g*, and cells were resuspended in 1.0 mL of supplemented RPMI-1640 (Nutricell,

Campinas, SP, Brazil). The supplemented RPMI-1640 contained: 10% heat-inactivated

fetal calf serum (Nutricell), penicillin (100 UI mL^{-1}), streptomycin (100 mg mL^{-1})

(Sigma-Aldrich, St. Louis, MO, USA), and amphotericin B (0.25 μg mL^{-1}) (Sigma-

Aldrich). Cell concentration was adjusted to 1.0×10^5 mononuclear phagocytes mL^{-1} as

judged by the uptake of 0.02% neutral red (Sigma) and confirmed by expression of F4/80

by Fluorescence-Activated Cell Sorting (FACS). Cells were plated into 96-well flat-

bottomed microtiter plates (Greiner BioOne, Frickenhausen, Germany), and incubated for

two hours at 37°C and 5% CO_2 in a humidified chamber to allow cells to adhere and

spread. Non-adherent cells were removed by washing the wells three times with RPMI,

and we used the remaining adherent cells (>95% mononuclear phagocytes as assessed by

morphological examination) for experiments. The adherent cells were cultured at 37°C

and 5% CO_2 in supplemented RPMI-1640 with or without heat-killed spores of *R. oryzae*

(*R. oryzae*-Ag) (1 spore: 1 cell). As an internal control for macrophage activity, cells were

cultured with 10 μg ml-1 lipopolysaccharide (Sigma-Aldrich). After 24 hours, cell-free

supernatants were harvested and stored at -80°C for cytokine analysis.

2.8 Production of hydrogen peroxide (H_2O_2). The production of H_2O_2 was estimated

according Russo et al. (1989). Briefly, adhered mononuclear phagocytes obtained as

described before were maintained in RPMI-1640 culture medium at 37°C and 5% CO_2

for 24 h. At the end of the cell culture period, we removed the supernatants from the wells,

added the phenol red solution containing dextrose (Sigma), phenol red (Sigma), and

horseradish peroxidase type II (Sigma), and incubated the plate at 37°C in 5% CO_2 for

one hour. Reaction was stopped by adding 1 N NaOH. The H_2O_2 concentration was

determined using a colorimetric microreader (ELx 800; BioTek Instruments Inc.,

Winooski, VE, USA)(36).

2.9 Detection of levels of nitric oxide (NO). The production of NO was estimated

following the Griess' method (Green et al., 1982). Production of nitrite, a stable end

product of NO, was measured in the cell-free supernatants of adhered mononuclear

phagocytes cultured. Briefly, a volume of 0.1 mL of cell-free supernatant with an equal

volume of Griess reagent was incubated for 10 minutes at room temperature. Griess

reagent was prepared using: 1% sulfanilamide (Synth, Diadema, SP, Brazil), 0.1%

naphthalene diamine dihydrochloride (Sigma), and 2.5% H3PO4. The nitrite

accumulation was quantified using a colorimetric microreader (ELx 800; BioTek

Instruments Inc., Winooski, VE, USA). Nitrite concentration was determined using

sodium nitrite (Sigma) diluted in RPMI-1640 medium as a standard (36).

2.10 Dosage of TNF-α and IL-10. IL-10 and TNF-α levels were measured in the cell-

free supernatants of cell cultures using a cytokine Duo-Set Kit (R&D Systems,

Minneapolis, MI, USA), according the manufacturer's instructions.

2.11 Statistical analyses. Normality tests of data was determined by Shapiro–Wilk's test.

For comparing two independent samples *t*-test was performed. For multiple comparisons,

ANOVA with the Tukey post-test was used. Pearson's correlation coefficient was

determined to measure the statistical association between two continuous variables. All

statistical tests were performed using GraphPad Prism version 5.0 for Windows

(GraphPad Software, San Diego, CA). *P*-value of 5% or lower was considered

statistically significant (Zar, 2010).

3. Results

3.1 Production of reactive oxygen reactive species and IL-10 by heat-killed *R. oryzae*-

infected peritoneal macrophages is dependent of mouse genetic background.

First, we evaluated the *in vitro* response of peritoneal (PMΦ) and alveolar (AMΦ)

macrophages obtained from non-infected BALB/c and Swiss mice challenged with heat-

killed *R. oryzae* (*R.oryzae*-Ag). To evaluate the PMΦ and AMΦ response we measured

reactive oxygen and nitrogen species (H_2O_2 and NO) in the cultures. We also measured

the levels of TNF-α and IL-10 to validate our findings.

In the presence of heat-killed spores of *R. oryzae*, PMΦ from the two mouse

strains showed decreased production of the NO (Fig 1A) and increased the TNF-α

production (Fig 1C). Interestedly, *R. oryzae*-infected PMΦ from BALB/c showed

increased production of H_2O_2 and IL-10. (Fig 1B, Fig 1D).

In general, AMΦ from both Swiss and BALB/c mice showed similar production

of NO and H_2O_2 (Fig 1E, Fig 1F). A slight increased production of TNF-α by *R. oryzae*-

infected AMΦ obtained from BALB/c was observed. In this set of experiments, no

detectable production of IL-10 was observed.

Fig 1. Peritoneal and alveolar macrophages from naive Swiss and BALB/c mice
react differently in front of heat-killed *R. oryzae*. (A) NO, (B) H2O2, (C) TNF-α, (D)
IL-10 levels in cell-free supernatant of peritoneal macrophages, and (E) NO (F) H2O2,
(G) TNF-α levels in cell-free supernatant of alveolar macrophages of non-infected Swiss,
and BALB/c mice co-cultured or not with heat-killed spores of R. oryzae. Student's t-test;
n=5-7; *p < 0.05, ** p< 0.01, ***p< 0.001). NO: nitric oxide; H2O2: hydrogen peroxide;
TNF-α: tumor necrosis factor-alpha; IL-10: interleukin 10; PMΦ: peritoneal
macrophages; AlvMΦ: alveolar macrophages.

3.2 Enhanced production of H_2O_2 by peritoneal macrophages was associated with better clearance of *R. oryzae* in a model of disseminated mucormycosis.

In the present study, BALB/c and Swiss mice strains showed more and less resistance, respectively, against *in vivo R. oryzae* infection, as previously reported (30,31). The course of infection was characterized by similar fungal load on day 7 for both mouse strain, followed by a decreased fungal load on day 30, that was more expressive in BALB/c mice (Fig 2A).

In order to explore the results of *in vitro R. oryzae*-infected macrophages experiments, we hypothesized that the production of reactive species of oxygen and nitrogen by macrophages could be involved in the differential response observed between these two mice strains.

To assess our hypothesis, we first evaluated the specific activity of PMΦ derived from BALB/c and Swiss mice that were intravenously infected with *R. oryzae*. For this, on days 7 and 30 p.i., PMΦ were recovered from the peritoneal cavity and challenged with heat-killed spore of *R. oryzae*.

On day 7, PMΦ from Swiss mice showed higher production of NO (Fig 2B) and TNF-α (Fig 2D) than PMΦ from BALB/c mice. On the other hand, on day 30, more resistant BALB/c mice showed higher levels of H_2O_2 (Fig 2C). No differences in the production of NO, TNF-α, and IL-10 were observed between both mouse strains (Fig 2B, Fig 2D, Fig 2E).

Next, we performed a correlation test between production of H_2O_2 and fungal load. An inverse correlation was observed between the production of H_2O_2 and total fungal load in the BALB/c mice (Fig 2G), but not in the Swiss mice (Fig 2F).

Fig 2. Increased PMΦ production of H_2O_2 at the end of disseminated infection is related to efficient fungal clearance in BALB/c mice. (A) Total fungal load (B) NO,

(C) H_2O_2, (D) TNF-α, and (E) IL-10 levels in cell-free supernatant of peritoneal macrophages from Swiss or BALB/c mice co-cultured with heat-killed spores of *R. oryzae*. Linear regression analysis between H_2O_2 levels and total fungal load in BALB/c (F) and Swiss (G) mice. The infected group was composed of mice inoculated intravenously with 3 x 10^4 spores of *R. oryzae* and evaluated after 7 and 30 days. Any significant differences relative to infected samples compared to different times post infection (letters) and to different strains (*) are indicated. (Student's t-test; n=5-7; *p < 0.05, ** p< 0.01, ***p< 0.001). NO: nitric oxide; H_2O_2: hydrogen peroxide; TNF-α: tumor necrosis factor alpha; IL-10: interleukin 10.

3.3 Enhanced initial pro-inflammatory response and NO production by alveolar macrophages of BALB/c mice seems to better control the pulmonary mucormycosis.

BALB/c and Swiss mice were intratracheally infected with *R. oryzae* to evaluate the tissue-specific response. As observed in intravenous infection, both BALB/c and Swiss mice showed a decrease in viable fungal load after 30 days p.i., and BALB/c mice showed better fungal clearance than Swiss mice (Fig 3A).

On day 7, AMΦ from BALB/c mice showed higher production of NO (Fig. 3B), TNF-α (Fig 3D), and IL-10 (Fig 3E) than AMΦ from Swiss mice. Lower production of H_2O_2 by AMΦ from BALB/c mice was observed (Fig 3C).

On day 30, no differences in the production of NO, H_2O_2, TNF-α, and IL-10 by AlvMΦ were observed (Fig 3B, Fig 3C, Fig 3D, Fig 3E).

Fig 3. AMΦ from BALB/c mice showed higher production of NO, TNF-α, and IL-10 than AMΦ from Swiss mice after 7 days of *R. oryzae* pulmonary infection. (A) Total fungal load, (B) NO, (C) H_2O_2, (D) TNF-α, and (E) IL-10, levels in cell-free supernatant of alveolar macrophages from Swiss or BALB/c mice co-cultured with heat-killed spores of *R. oryzae*. The infected group was composed of mice inoculated intratracheally with 2 ×10^6 spores of *R. oryzae* and evaluated after 7 and 30 days. Any significant differences relative to infected samples compared to different times post infection (letters) and to different strains (*) are indicated. (Student´s t-test; n=5-7; *p < 0.05, ** p< 0.01, ***p< 0.001). NO: nitric oxide; H_2O_2: hydrogen peroxide; TNF-α: tumor necrosis factor alpha; IL-10: interleukin 10.

4. Discussion

Here we use the different outcomes observed in experimental mucormycosis between the immunocompetent BALB/c and Swiss mice to highlight the protective pattern of macrophages responses against *R. oryzae*. We found that macrophages from different microenvironments showed different responses against *R. oryzae* not only *in vitro* but also during *in vivo* infection. Also, the better outcome was associated with higher H_2O_2 produced by PM, and higher initial NO, TNF-α, and IL-10 produced by AMΦ during experimental disseminated and pulmonary mucormycosis, respectively. In addition, the genetic background interferes in the macrophage response, so, regardless of the microenvironment, the BALB/c's macrophage is more effective in eliminating the fungus.

Inbred mouse strains vary widely in their degree of innate susceptibility to systemic fungal infections (37–41). BALB/c and Swiss mice are resistant and susceptible strains, respectively, in experimental infections caused by *Cryptococcus neoformans* (Rhodes et al., 1980), *Candida albicans* (38,42), and *Paracoccidioides brasiliensis* (41). It was identified that the genetic target C5-deficiency (Hc0 allele, hemolytic complement) modulates the host's initial response and causes susceptibility and ineffective inflammatory response against fungal infections (38,39). Active as an anaphylatoxin, the C5 molecule is a power chemotactic factor for polymorphonuclear leukocytes. Lack of C5 would block complement, thereby decreasing the opsonization and recruitment of phagocytic cells (39). Although the higher susceptibility of Swiss mice to fungal infections may be explained by a deficiency in the Hc0 allele (41–43), our results suggest that other genetic targets may be involved in the less responsiveness of macrophages from non-infected Swiss mice to heat-killed *R. oryzae*.

According to clinical and experimental data, individuals who lack phagocytes or present impaired phagocytic functions show a higher risk of developing mucormycosis

(6,44,45). We observed in the *in vitro* part of this study that peritoneal macrophages from

the most resistant strain (BALB/c) showed a potent inflammatory response mediated by

H_2O_2 and TNF-α. (46) in the first contact with *R. oryzae* antigen. A previous *in vitro* study

showed that the inactivated Mucorales cells can stimulate high levels of TNF-α, IL-1β,

IL-6, I-L8, GM-CSF, and MCP-1 production by humans cells of different immune

subsets, with paralleled upregulation of transcriptional activity of IL-1β and TNF-α

(47,48). Since the oxidative microbicidal production by neutrophils and monocytes is

related to *R. oryzae* hyphae damage and killing (49,50), our results indicate an initial

proinflammatory response against *R. oryzae* can lead to a better outcome in experimental

mucormycosis.

Unexpectedly, in the first contact with *R. oryzae* antigen peritoneal macrophages,

in general, decreased the production of NO. Similar to our results, it was demonstrated

that *Aspergillus nidulans* melanin inhibited the NO production by lipopolysaccharide

(LPS)-stimulated peritoneal macrophages, accompanied by a slight stimulatory effect on

TNF-α production (51). Melanin is a type of pigment in the fungal cell wall of black

molds and has been shown to block the effects of hydrolytic enzymes on the cell wall

(52). Considering the importance of this finding, we suggest that the immunomodulatory

effects induced by the *R. oryzae* cell wall on peritoneal macrophages need to be better

investigated.

Differently from peritoneal macrophages, alveolar macrophages from the BALB/c

strain produced higher NO and TNF-α levels than less resistant strain, during the *in vivo*

model of pulmonary mucormycosis. It is known that pulmonary activated macrophages

are the major defense against fungal invasion (53). In contact with macrophages

receptors, TNF-α provides signals that lead to an induction of antimicrobial activity. This

activity depends on NO synthase activation and NO production (54). Some studies are

demonstrating that this process is essential to fungal death. In paracoccidioidomycosis (PCM), TNF-a induces *P. brasiliensis* killing by H_2O_2 and NO release (55). In cryptococcosis, TNF-α significantly promoted macrophage NO production and anticryptococcal activity(56). In addition, TNF-α enhances pulmonary alveolar macrophage phagocytosis and oxygen production during initial contact with *A. fumigatus* (57). It is known that TNF-α contributes to the influx and activation of neutrophils and mononuclear cells in the lungs during the filamentous fungal challenge (58). In mucormycosis, the TNFα signaling may have a protective response since patients treated with a tumor necrosis factor inhibitor have a higher risk to developed disseminated mucormycosis(59,60).

Although the association of NO with the killing of yeast cells supports the importance of this metabolite in host protection (61), other studies have been demonstrating that the overproduction of this metabolite is associated with susceptibility in experimental PCM (62,63). Nascimento, et. al (2002) observed that high levels of NO induce T cell immunosuppression during *P. brasiliensis* infection (Nascimento et al., 2002). These reports suggest that the protective or deleterious NO role *in vivo* depends on the balance between its fungicidal and immunosuppressive properties. Our data demonstrate that after intravenous *R. oryzae* infection, stronger and early production of NO and TNF-a by peritoneal macrophages from Swiss mice does not result in late better fungal clearance. On the other hand, after a pulmonary *R. oryzae* infection, higher and early production of NO and TNF-a by alveolar macrophages from BALB/c mice was accompanied by higher levels of IL-10, resulting in later better control of pulmonary mucormycosis. Suggesting that in BALB/c mice, an initial balanced immune response is essential for better infection control.

IL-10 is an anti-inflammatory cytokine that can impede pathogen clearance. But

it also can ameliorate immunopathology (64). The balance between pro-and anti-

inflammatory cytokines is crucial for the host´s defense against *A. fumigatus*. An *in vitro*

study with mononuclear cells (MNC) stimulated with *A. fumigatus* showed that hyphae

of this fungi induced the release of IL-10 by MNC, and this process was dependent on

endogenous IL-1 (65). Whereas inflammatory cytokines such as IFN-γ, TNF-α, and IL-

18 activate monocytes and neutrophils to ingest and kill *A. fumigatus* conidia and hyphae,

the subsequent release of the anti-inflammatory cytokine IL-10, are responsible for down-

regulating the potential deleterious overstimulation induced by the inflammatory

mediators (66).

In general, one of the biological effects of IL-1 is an increased synthesis of IL-10

(67). Netea et al., (2004) showed that *Candida* stimulated IL-1 release by cells, and

secondarily IL-10 release (68). In Mucormycosis, although high production of IL-10 by

Mucorales-specific T cells from patients was related to susceptibility in the late phase of

the disease (69), our results showed a correlation between higher IL-10 production by

alveolar macrophages in the initial moments of *R. oryzae* infection and better outcome in

experimental pulmonary mucormycosis. We suggest that high levels of IL-10 were a

biological effect to balance the pro-inflammatory response mediated by high levels of

TNF-α and NO. A fundamental step for an efficient immune response.

While alveolar macrophages from a more resistant strain showed an initial higher

TNF-α, NO, and IL-10 response to a pulmonary infection, the peritoneal macrophages

from the same strain showed a late and strong response mediated by H_2O_2 after the

intravenous *R. oryzae* infection. It is important to note that after infection by the

intravenous route, the decrease in viable *R. oryzae* was inversely proportional to the

release of H_2O_2 by peritoneal macrophages from a more resistant strain of mice. It

indicates the relation of this mediator with protection in Mucormycosis. This result agrees with the study done by Andrianaki et al. (33), where they prove the susceptibility of *A. fumigatus* and *R. oryzae* conidia to oxidative damage induced by H_2O_2. In this context, an important detail was that the BALB/c strain only showed a more efficient fungal clearance after 30 days of infection. We suggest that the adaptative immune response potentiated the activity of the macrophages evaluated in this study.

Macrophages are the first immune cells to interact with invasive pathogens. Depending on macrophage's activity, they can prime the adaptive immune response, which is far more aggressive and specific to pathogens (70). Generally, high levels of ROS production by macrophages are linked to intracellular events, including the activation of transcription factors such as NFκB (71), and induction of mitogenesis (72). In addition, *in vitro*, and *in vivo* studies already showed that *R. oryzae* could trigger a Th-17 response mediated by high amounts of IL-23 by dendritic cells (73), and also the association of high levels of IL-17 and IFN-γ with better *R. oryzae* elimination by immunocompetent BALB/c and C57BL/6 mice (30,31). Considering the studies above and the evidence found here, we hypothesized that adaptive immune responses developed by BALB/c mice were essential to enhance the fungicidal activity of peritoneal macrophages and efficiently kill *R. oryzae*.

The absence of additional experiments to explore the intracellular mechanisms mediating the differences in ROS production and to confirm the adaptive immune response related to a more efficient macrophages response front of *R. oryzae*-Ag was the main limitation of the present study, and we suggest that more studies should be done to clarify the results observed here.

In summary, our findings reveal that independently of mice strain, PMΦ is more reactive against *R. oryzae* in the first contact than AMΦ. In addition, increased PMΦ

production of H_2O_2 at the end of disseminated infection is related to efficient fungal

clearance observed in resistant (BALB/c). Our findings provide new evidence to

understand the parasite-hosts relationship in mucormycosis.

References

1. Severo LC, Oliveira FDM, Dreher R, Teixeira PZ, Porto NDS, Londero AT. Zygomycosis: A
report of eleven cases and a review of the Brazilian literature. Rev Iberoam Micol. março
de 2002;19(1):52–6.

2. Bitar D, Van Cauteren D, Lanternier F, Dannaoui E, Che D, Dromer F, et al. Increasing
incidence of zygomycosis (mucormycosis), France, 1997-2006. Emerging Infect Dis.
setembro de 2009;15(9):1395–401.

3. Kume H, Yamazaki T, Abe M, Tanuma H, Okudaira M, Okayasu I. Increase in aspergillosis
and severe mycotic infection in patients with leukemia and MDS: comparison of the data
from the Annual of the Pathological Autopsy Cases in Japan in 1989, 1993 and 1997.
Pathol Int. novembro de 2003;53(11):744–50.

4. Neofytos D, Horn D, Anaissie E, Steinbach W, Olyaei A, Fishman J, et al. Epidemiology and
outcome of invasive fungal infection in adult hematopoietic stem cell transplant
recipients: analysis of Multicenter Prospective Antifungal Therapy (PATH) Alliance
registry. Clin Infect Dis. 1º de fevereiro de 2009;48(3):265–73.

5. Ribes JA, Vanover-Sams CL, Baker DJ. Zygomycetes in human disease. Clin Microbiol Rev.
abril de 2000;13(2):236–301.

6. Ibrahim AS, Spellberg B, Walsh TJ, Kontoyiannis DP. Pathogenesis of Mucormycosis. Clin
Infect Dis. 1º de fevereiro de 2012;54(Suppl 1):S16–22.

7. Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele RL, et al.
Epidemiology and outcome of zygomycosis: a review of 929 reported cases. Clin Infect
Dis. 1º de setembro de 2005;41(5):634–53.

8. Spellberg B, Edwards J, Ibrahim A. Novel perspectives on mucormycosis:
pathophysiology, presentation, and management. Clin Microbiol Rev. julho de
2005;18(3):556–69.

9. Baddley JW, Thompson GR III, Chen SCA, White PL, Johnson MD, Nguyen MH, et al.
Coronavirus Disease 2019–Associated Invasive Fungal Infection. Open Forum Infectious
Diseases. 1º de dezembro de 2021;8(12):ofab510.

10. Narayanan S, Chua JV, Baddley JW. COVID-19 associated Mucormycosis (CAM): risk
factors and mechanisms of disease. Clin Infect Dis. 22 de agosto de 2021;ciab726.

11. Hussain S, Baxi H, Riad A, Klugarová J, Pokorná A, Slezáková S, et al. COVID-19-Associated
Mucormycosis (CAM): An Updated Evidence Mapping. Int J Environ Res Public Health. 30
de setembro de 2021;18(19):10340.

12.   Rabagliati R, Rodríguez N, Núñez C, Huete A, Bravo S, Garcia P. COVID-19-Associated
445   Mold Infection in Critically Ill Patients, Chile. Emerg Infect Dis. maio de 2021;27(5):1454–
446   6.

447   13.   Patel A, Agarwal R, Rudramurthy SM, Shevkani M, Xess I, Sharma R, et al. Multicenter
448   Epidemiologic Study of Coronavirus Disease–Associated Mucormycosis, India - Volume
449   27, Number 9—September 2021 - Emerging Infectious Diseases journal - CDC. [citado 1º
450   de fevereiro de 2022]; Disponível em: https://wwwnc.cdc.gov/eid/article/27/9/21-
451   0934_article

452   14.   Hoenigl M, Seidel D, Carvalho A, Rudramurthy SM, Arastehfar A, Gangneux JP, et al. The
453   emergence of COVID-19 associated mucormycosis: a review of cases from 18 countries.
454   The Lancet Microbe [Internet]. 25 de janeiro de 2022 [citado 1º de fevereiro de
455   2022];0(0). Disponível em: https://www.thelancet.com/journals/lanmic/article/PIIS2666-
456   5247(21)00237-8/fulltext

457   15.   Bonates P, João GAP, Cruz KS, Ferreira M de S, Baía-da-Silva DC, de Farias MEL, et al. Fatal
458   rhino-orbito-cerebral mucormycosis infection associated with diabetic ketoacidosis post-
459   COVID-19. Rev Soc Bras Med Trop. 54:e0358-2021.

460   16.   Monte Junior ES do, Santos MELD, Ribeiro IB, Luz G de O, Baba ER, Hirsch BS, et al. Rare
461   and Fatal Gastrointestinal Mucormycosis (Zygomycosis) in a COVID-19 Patient: A Case
462   Report. Clin Endosc. novembro de 2020;53(6):746–9.

463   17.   Pauli MA, Pereira L de M, Monteiro ML, de Camargo AR, Rabelo GD. Painful palatal lesion
464   in a patient with COVID-19. Oral Surg Oral Med Oral Pathol Oral Radiol. junho de
465   2021;131(6):620–5.

466   18.   Nasir N, Farooqi J, Mahmood SF, Jabeen K. COVID-19 associated mucormycosis: a life-
467   threatening complication in patients admitted with severe to critical COVID-19 from
468   Pakistan. Clin Microbiol Infect. novembro de 2021;27(11):1704–7.

469   19.   Dulski TM. Notes from the Field: COVID-19–Associated Mucormycosis — Arkansas, July–
470   September 2021. MMWR Morb Mortal Wkly Rep [Internet]. 2021 [citado 1º de fevereiro
471   de 2022];70. Disponível em:
472   https://www.cdc.gov/mmwr/volumes/70/wr/mm7050a3.htm

473   20.   Mejía-Santos H. Notes from the Field: Mucormycosis Cases During the COVID-19
474   Pandemic — Honduras, May–September 2021. MMWR Morb Mortal Wkly Rep [Internet].
475   2021 [citado 1º de fevereiro de 2022];70. Disponível em:
476   https://www.cdc.gov/mmwr/volumes/70/wr/mm7050a2.htm

477   21.   Cornely OA, Alastruey-Izquierdo A, Arenz D, Chen SCA, Dannaoui E, Hochhegger B, et al.
478   Global guideline for the diagnosis and management of mucormycosis: an initiative of the
479   European Confederation of Medical Mycology in cooperation with the Mycoses Study
480   Group Education and Research Consortium. The Lancet Infectious Diseases. 1º de
481   dezembro de 2019;19(12):e405–21.

482   22.   Lelievre L, Garcia-Hermoso D, Abdoul H, Hivelin M, Chouaki T, Toubas D, et al.
483   Posttraumatic mucormycosis: a nationwide study in France and review of the literature.
484   Medicine (Baltimore). novembro de 2014;93(24):395–404.

23. Neblett Fanfair R, Benedict K, Bos J, Bennett SD, Lo YC, Adebanjo T, et al. Necrotizing
cutaneous mucormycosis after a tornado in Joplin, Missouri, in 2011. N Engl J Med. 6 de
dezembro de 2012;367(23):2214–25.

24. Kontoyiannis DP, Lewis RE, Lortholary O, Lotholary O, Spellberg B, Petrikkos G, et al.
Future directions in mucormycosis research. Clin Infect Dis. fevereiro de 2012;54 Suppl
1:S79-85.

25. Skiada A, Lanternier F, Groll AH, Pagano L, Zimmerli S, Herbrecht R, et al. Diagnosis and
treatment of mucormycosis in patients with hematological malignancies: guidelines from
the 3rd European Conference on Infections in Leukemia (ECIL 3). Haematologica. 1º de
abril de 2013;98(4):492–504.

26. Rodríguez MM, Pastor FJ, Calvo E, Salas V, Sutton DA, Guarro J. Correlation of in vitro
activity, serum levels, and in vivo efficacy of posaconazole against Rhizopus microsporus
in a murine disseminated infection. Antimicrob Agents Chemother. dezembro de
2009;53(12):5022–5.

27. Lewis RE, Albert ND, Liao G, Hou J, Prince RA, Kontoyiannis DP. Comparative
pharmacodynamics of amphotericin B lipid complex and liposomal amphotericin B in a
murine model of pulmonary mucormycosis. Antimicrob Agents Chemother. março de
2010;54(3):1298–304.

28. Lewis RE, Liao G, Wang W, Prince RA, Kontoyiannis DP. Voriconazole pre-exposure selects
for breakthrough mucormycosis in a mixed model of Aspergillus fumigatus-Rhizopus
oryzae pulmonary infection. Virulence. agosto de 2011;2(4):348–55.

29. Lewis RE, Ben-Ami R, Best L, Albert N, Walsh TJ, Kontoyiannis DP. Tacrolimus enhances
the potency of posaconazole against Rhizopus oryzae in vitro and in an experimental
model of mucormycosis. J Infect Dis. 1º de março de 2013;207(5):834–41.

30. Santos AR dos, Fraga-Silva TF, F Almeida D de, dos Santos RF, Finato AC, Amorim BC, et
al. Rhizopus-host interplay of disseminated mucormycosis in immunocompetent mice.
Future Microbiology [Internet]. 20 de julho de 2020 [citado 24 de julho de 2020];
Disponível em: https://www.futuremedicine.com/doi/10.2217/fmb-2019-0246

31. Dos Santos AR, Fraga-Silva TF, de Fátima Almeida-Donanzam D, Dos Santos RF, Finato AC,
Soares CT, et al. IFN-γ Mediated Signaling Improves Fungal Clearance in Experimental
Pulmonary Mucormycosis. Mycopathologia. 29 de outubro de 2021;

32. Bao W, Jin L, Fu H jing, Shen Y nian, Lu G xia, Mei H, et al. Interleukin-22 Mediates Early
Host Defense against Rhizomucor pusilluscan Pathogens. PLoS One [Internet]. 17 de
junho de 2013 [citado 18 de julho de 2019];8(6). Disponível em:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3684593/

33. Andrianaki AM, Kyrmizi I, Thanopoulou K, Baldin C, Drakos E, Soliman SSM, et al. Iron
restriction inside macrophages regulates pulmonary host defense against Rhizopus
species. Nature Communications. 20 de agosto de 2018;9(1):3333.

34. Zhang SX, Babady NE, Hanson KE, Harrington AT, Larkin PMK, Leal SM, et al. Recognition
of Diagnostic Gaps for Laboratory Diagnosis of Fungal Diseases: Expert Opinion from the
Fungal Diagnostics Laboratories Consortium (FDLC). J Clin Microbiol. 18 de junho de
2021;59(7):e0178420.

35. Venturini J, Golim MA, Alvares AM, Locachevic GA, Arruda OS, Arruda MSP.
Morphofunctional evaluation of thymus in hyperglycemic-hypoinsulinemic mice during
dermatophytic infection. FEMS Immunol Med Microbiol. junho de 2011;62(1):32–40.

36. Fraga-Silva TFC, Marchetti CM, Mimura LAN, Locachevic GA, Golim MA, Venturini J, et al.
Relationship among Short and Long Term of Hypoinsulinemia-Hyperglycemia,
Dermatophytosis, and Immunobiology of Mononuclear Phagocytes. Mediators Inflamm.
2015;2015:342345.

37. Calich VL, Singer-Vermes LM, Siqueira AM, Burger E. Susceptibility and resistance of
inbred mice to Paracoccidioides brasiliensis. Br J Exp Pathol. outubro de 1985;66(5):585–
94.

38. Radovanovic I, Mullick A, Gros P. Genetic control of susceptibility to infection with
Candida albicans in mice. PLoS One. 20 de abril de 2011;6(4):e18957.

39. Rhodes JC, Wicker LS, Urba WJ. Genetic Control of Susceptibility to Cryptococcus
neoformans in Mice. Infect Immun. agosto de 1980;29(2):494–9.

40. Svirshchevskaya EV, Shevchenko MA, Huet D, Femenia F, Latgé JP, Boireau P, et al.
Susceptibility of mice to invasive aspergillosis correlates with delayed cell influx into the
lungs. Int J Immunogenet. outubro de 2009;36(5):289–99.

41. Vilani-Moreno F, Fecchio D, de Mattos MC, Moscardi-Bacchi M, Defaveri J, Franco M.
Study of pulmonary experimental paracoccidioidomycosis by analysis of bronchoalveolar
lavage cells: resistant vs. susceptible mice. Mycopathologia. 1998;141(2):79–91.

42. Hasenclever HF. Comparative pathogenicity of Candida albicans for mice and rabbits. J
Bacteriol. julho de 1959;78(1):105–9.

43. Maheshwari RK, Tandon RN, Feuillette AR, Mahouy G, Badillet G, Friedman RM.
Interferon inhibits Aspergillus fumigatus growth in mice: an activity against an
extracellular infection. J Interferon Res. fevereiro de 1988;8(1):35–44.

44. Ribes JA, Vanover-Sams CL, Baker DJ. Zygomycetes in human disease. Clin Microbiol Rev.
abril de 2000;13(2):236–301.

45. Waldorf AR, Ruderman N, Diamond RD. Specific susceptibility to mucormycosis in murine
diabetes and bronchoalveolar macrophage defense against Rhizopus. J Clin Invest. julho
de 1984;74(1):150–60.

46. Bradley JR. TNF-mediated inflammatory disease. J Pathol. janeiro de 2008;214(2):149–60.

47. Belic S, Page L, Lazariotou M, Waaga-Gasser AM, Dragan M, Springer J, et al. Comparative
Analysis of Inflammatory Cytokine Release and Alveolar Epithelial Barrier Invasion in a
Transwell® Bilayer Model of Mucormycosis. Front Microbiol. 2018;9:3204.

48. Wurster S, Thielen V, Weis P, Walther P, Elias J, Waaga-Gasser AM, et al. Mucorales
spores induce a proinflammatory cytokine response in human mononuclear phagocytes
and harbor no rodlet hydrophobins. Virulence. 17 de 2017;8(8):1708–18.

49.   Diamond RD, Clark RA. Damage to Aspergillus fumigatus and Rhizopus oryzae hyphae by
oxidative and nonoxidative microbicidal products of human neutrophils in vitro. Infect
Immun. novembro de 1982;38(2):487–95.

50.   Diamond RD, Haudenschild CC, Erickson NF. Monocyte-mediated damage to Rhizopus
oryzae hyphae in vitro. Infect Immun. outubro de 1982;38(1):292–7.

51.   Gonçalves R de CR, Kitagawa RR, Raddi MSG, Carlos IZ, Pombeiro-Sponchiado SR.
Inhibition of nitric oxide and tumour necrosis factor-α production in peritoneal
macrophages by Aspergillus nidulans melanin. Biol Pharm Bull. 2013;36(12):1915–20.

52.   Heinekamp T, Thywissen A, Macheleidt J, Keller S, Valiante V, Brakhage AA. Aspergillus
fumigatus melanins: interference with the host endocytosis pathway and impact on
virulence. Front Microbiol [Internet]. 2013 [citado 23 de agosto de 2018];3. Disponível
em: https://www.frontiersin.org/articles/10.3389/fmicb.2012.00440/full

53.   Leopold Wager CM, Wormley FL. Classical versus alternative macrophage activation: the
Ying and the Yang in host defense against pulmonary fungal infections. Mucosal
Immunol. setembro de 2014;7(5):1023–35.

54.   Salim T, Sershen CL, May EE. Investigating the Role of TNF-α and IFN-γ Activation on the
Dynamics of iNOS Gene Expression in LPS Stimulated Macrophages. PLoS One. 8 de junho
de 2016;11(6):e0153289.

55.   Moreira AP, Dias-Melicio LA, Peraçoli MTS, Calvi SA, Victoriano de Campos Soares AM.
Killing of Paracoccidioides brasiliensis yeast cells by IFN-gamma and TNF-alpha activated
murine peritoneal macrophages: evidence of H(2)O (2) and NO effector mechanisms.
Mycopathologia. julho de 2008;166(1):17–23.

56.   Kawakami K, Qureshi MH, Koguchi Y, Zhang T, Okamura H, Kurimoto M, et al. Role of
TNF-alpha in the induction of fungicidal activity of mouse peritoneal exudate cells against
Cryptococcus neoformans by IL-12 and IL-18. Cell Immunol. 10 de abril de 1999;193(1):9–
16.

57.   Roilides E, Dimitriadou-Georgiadou A, Sein T, Kadiltsoglou I, Walsh TJ. Tumor necrosis
factor alpha enhances antifungal activities of polymorphonuclear and mononuclear
phagocytes against Aspergillus fumigatus. Infect Immun. dezembro de
1998;66(12):5999–6003.

58.   Mehrad B, Strieter RM, Standiford TJ. Role of TNF-alpha in pulmonary host defense in
murine invasive aspergillosis. J Immunol. 1º de fevereiro de 1999;162(3):1633–40.

59.   Keijzer A, Valk P van der, Ossenkoppele GJ, Loosdrecht A van de. Mucormycosis in a
patient with low risk myelodysplasia treated with anti-TNF-alpha. Haematologica. 1º de
janeiro de 2006;91(12 Suppl):ECR51–ECR51.

60.   Singh P, Taylor SF, Murali R, Gomes LJ, Kanthan GL, Maloof AJ. Disseminated
mucormycosis and orbital ischaemia in combination immunosuppression with a tumour
necrosis factor alpha inhibitor. Clinical & Experimental Ophthalmology. 2007;35(3):275–
80.

61.   Gonzalez A, de Gregori W, Velez D, Restrepo A, Cano LE. Nitric oxide participation in the
604   fungicidal mechanism of gamma interferon-activated murine macrophages against
605   Paracoccidioides brasiliensis conidia. Infect Immun. maio de 2000;68(5):2546–52.

606   62.   Bocca AL, Hayashi EE, Pinheiro AG, Furlanetto AB, Campanelli AP, Cunha FQ, et al.
607   Treatment of Paracoccidioides brasiliensis-infected mice with a nitric oxide inhibitor
608   prevents the failure of cell-mediated immune response. J Immunol. 15 de setembro de
609   1998;161(6):3056–63.

610   63.   Nascimento FRF, Calich VLG, Rodríguez D, Russo M. Dual role for nitric oxide in
611   paracoccidioidomycosis: essential for resistance, but overproduction associated with
612   susceptibility. J Immunol. 1º de maio de 2002;168(9):4593–600.

613   64.   Couper KN, Blount DG, Riley EM. IL-10: The Master Regulator of Immunity to Infection.
614   The Journal of Immunology. 1º de maio de 2008;180(9):5771–7.

615   65.   Roilides E, Dimitriadou A, Kadiltsoglou I, Sein T, Karpouzas J, Pizzo PA, et al. IL-10 exerts
616   suppressive and enhancing effects on antifungal activity of mononuclear phagocytes
617   against Aspergillus fumigatus. The Journal of Immunology. 1º de janeiro de
618   1997;158(1):322–9.

619   66.   Grünig G, Corry DB, Leach MW, Seymour BW, Kurup VP, Rennick DM. Interleukin-10 is a
620   natural suppressor of cytokine production and inflammation in a murine model of allergic
621   bronchopulmonary aspergillosis. J Exp Med. 17 de março de 1997;185(6):1089–99.

622   67.   Dinarello CA. Biologic basis for interleukin-1 in disease. Blood. 15 de março de
623   1996;87(6):2095–147.

624   68.   Netea MG, Sutmuller R, Hermann C, Graaf CAAV der, Meer JWMV der, Krieken JH van, et
625   al. Toll-Like Receptor 2 Suppresses Immunity against Candida albicans through Induction
626   of IL-10 and Regulatory T Cells. The Journal of Immunology. 15 de março de
627   2004;172(6):3712–8.

628   69.   Potenza L, Vallerini D, Barozzi P, Riva G, Forghieri F, Zanetti E, et al. Mucorales-specific T
629   cells emerge in the course of invasive mucormycosis and may be used as a surrogate
630   diagnostic marker in high-risk patients. Blood. 17 de novembro de 2011;118(20):5416–9.

631   70.   Ghuman H, Voelz K. Innate and Adaptive Immunity to Mucorales. J Fungi (Basel). 5 de
632   setembro de 2017;3(3).

633   71.   Sun Y, Oberley LW. Redox regulation of transcriptional activators. Free Radical Biology
634   and Medicine. 1º de janeiro de 1996;21(3):335–48.

635   72.   Joneson T, Bar-Sagi D. A Rac1 Effector Site Controlling Mitogenesis through Superoxide
636   Production. J Biol Chem. 17 de julho de 1998;273(29):17991–4.

637   73.   Chamilos G, Ganguly D, Lande R, Gregorio J, Meller S, Goldman WE, et al. Generation of
638   IL-23 producing dendritic cells (DCs) by airborne fungi regulates fungal pathogenicity via
639   the induction of T(H)-17 responses. PLoS ONE. 23 de setembro de 2010;5(9):e12955.

Striking image. Overview of *in vitro* and *in vivo* macrophages response in BALB/c and Swiss mice infected with *R. oryzae* via different routes of infection. Macrophage activity of BALB/c and Swiss mice were evaluated pre-and-post intravenous or intratracheal *R. oryzae* infection. NO, H_2O_2, TNF-α, and IL-10 concentrations were evaluated in macrophages co-cultured or not with heat-killed spores of *R. oryzae*. Infected mice were intravenously or intratracheally inoculated with 3×10^4 and 2×10^6 spores of *R. oryzae* respectively, and evaluated after 7 and 30 days. Summarized results of the fungal load on day 30 are also shown. NO: nitric oxide; H_2O_2: hydrogen peroxide; TNF-α: tumor necrosis factor alpha; IL-10: interleukin 10; PMΦ: peritoneal macrophages; AlvMΦ: alveolar macrophages. Image by Débora Almeida-Donanzam via Inkscape

Figure 1

Figure 2

Figure 3