1
18 **Aging and sperm signals alter DNA break formation and repair in the *C. elegans* germline**
19
20
21
22 Erik Toraason[1][¶], Victoria L. Adler[1][¶], and Diana E. Libuda[1],[*]
23
24 1 University of Oregon, Department of Biology, Institute of Molecular Biology, Eugene, OR,
25 97403
26
27 [¶] These authors contributed equally to this work
28
29 [*] Corresponding Author (dlibuda@uoregon.edu)
48 Short title: DNA break formation and repair in aging oocytes
49
50

Abstract

Female reproductive aging is associated with decreased oocyte quality and fertility. The nematode *Caenorhabditis elegans* is a powerful system for understanding the biology of aging and exhibits age-related reproductive defects that are analogous to those observed in many mammals, including dysregulation of DNA repair. *C. elegans* germline function is influenced simultaneously by both reproductive aging and signals triggered by limited supplies of sperm, which are depleted over chronological time. To delineate the causes of DNA repair defects in aged *C. elegans* germlines, we assessed both DNA double strand break (DSB) induction and repair during meiotic prophase progression in aged germlines which were depleted of self-sperm, mated, or never exposed to sperm. We find that germline DSB induction is dramatically reduced only in hermaphrodites which have exhausted their endogenous sperm, suggesting that a signal due specifically to sperm depletion downregulates DSB formation. We also find that DSB repair is delayed in aged germlines regardless of whether hermaphrodites had either a reduction in sperm supply or an inability to endogenously produce sperm. These results demonstrate that in contrast to DSB induction, DSB repair defects are a feature of *C. elegans* reproductive aging independent of sperm presence. Finally, we demonstrate that the ubiquitin E2 ligase variant UEV-2 is required for efficient DSB repair specifically in young germlines, implicating UEV-2 in the regulation of DNA repair during reproductive aging. In summary, our study demonstrates that DNA repair defects are a feature of *C. elegans* reproductive aging and uncovers parallel mechanisms regulating efficient DSB formation in the germline.

Author Summary

Aging leads to a decline in the quality of the female reproductive cells, known as oocytes.

Oocytes subjected to reproductive aging experience an increase in both infertility and

aneuploidies that cause miscarriages and birth defects. The nematode *Caenorhabditis elegans*

is a classic model system used to determine the mechanisms of aging. Old *C. elegans* oocytes

accrue many defects which may contribute to their reduced quality, including dysregulation of

DNA repair. *C. elegans* fertility and germline function is also regulated oocyte-independently by

sperm-dependent signals. To determine how aging and sperm may independently impact DNA

repair in aging *C. elegans* oocytes, we control oocyte aging and sperm presence independently

to evaluate their effects on DNA break formation and repair. We find that running out of sperm

reduces the levels of DNA breaks which are produced, but the efficiency of DNA repair declines

during aging independent of sperm effects. We also identify a protein which specifically

promotes DNA repair in the oocytes of young animals, suggesting that this protein may regulate

DNA repair in the germline during aging. Taken together, our research defines aging-specific

and aging-independent mechanisms which regulate the genome integrity of oocytes.

Introduction

Genome integrity must be preserved during gamete development, as any genetic

defects incurred may have detrimental effects on progeny or fertility. Meiosis, the specialized

cell division that generates haploid gametes such as eggs and sperm, utilizes specific DNA

repair pathways to both ensure accurate chromosome segregation and preserve genomic

integrity. During early meiotic prophase I, DNA double-strand breaks (DSBs) are intentionally

induced across the genome by the conserved topoisomerase-like protein Spo11 (Keeney *et al.*

1997; Dernburg *et al.* 1998). A specific subset of these breaks must be repaired by

recombination as crossovers, creating the physical connections between homologous

chromosomes required for accurate chromosome segregation. Failure to repair meiotic DSBs

accurately and efficiently can contribute to infertility or risk the formation of *de novo* germline

mutations.

Gamete quality is negatively impacted in organisms of advanced chronological age

(Broekmans *et al.* 2007). In many organisms, oocyte quality in particular declines starkly with

maternal age (Luo *et al.* 2009, 2010; Moghadam *et al.* 2022). Oocyte aging is associated with

conserved phenotypic changes, including loss of sister chromatid cohesion, dysregulation of

DNA repair gene expression, and derepression of heterochromatin and retroviral elements (Luo

et al. 2010; Achache *et al.* 2021; Raices *et al.* 2021; Chatzidaki *et al.* 2021; Wasserzug-Pash *et*

al. 2022).

The nematode *Caenorhabditis elegans* is a key model system for the study of aging

biology, including age-related infertility (Mack *et al.* 2018). *C. elegans* hermaphrodites (which

produce oocytes as adults) undergo reproductive senescence due to declining oocyte quality

and incur many of the defects observed in the aging mammalian ovary (Andux and Ellis 2008;

Luo *et al.* 2009, 2010; Achache *et al.* 2021). Unlike many mammalian systems, however, which

generate oocytes *in utero* and hold them in dictyate arrest until ovulation, *C. elegans*

hermaphrodites continuously produce new oocytes during their adult reproductive period (Albert

Hubbard and Greenstein 2000). Mitotic proliferation and ovulation of oocytes is dependent upon

signals from sperm, which are stored at the end of the germline in a specialized compartment

called the 'spermatheca' (Mccarter *et al.* 1999; Cinquin *et al.* 2016). "Obligate female" mutants,

which do not produce sperm, therefore exhibit dramatically slowed germline proliferation and

progression (Doniach and Hodgkin 1984; Schedl and Kimble 1988; Mccarter *et al.* 1999;

Cinquin *et al.* 2016). The *C. elegans* germline is organized in a spatial temporal gradient

wherein oocytes mitotically proliferate at the distal tip and move proximally through the germline

as they progress through meiotic prophase I (Albert Hubbard and Greenstein 2000).Thus,

oocyte nuclei at all stages of meiotic prophase I are simultaneously present in the adult germline

and enable assessment of meiotic events which are dynamic across prophase, such as the

induction and repair of DSBs.

Multiple lines of evidence suggest that preservation of genome integrity is important for

the maintenance of oocyte quality during reproductive aging. Human females carrying DNA

repair protein variants exhibit extended fertility (Ruth *et al.* 2021). *C. elegans* mutants with

extended reproductive periods are also resilient to exogenous DNA damage and upregulate

genes associated with DNA repair (Luo *et al.* 2010). Further, recent evidence demonstrated that

DNA damage and repair are altered in aged *C. elegans* germlines (Achache *et al.* 2021; Raices

et al. 2021). By the fourth day of adulthood, *C. elegans* oocyte nuclei exhibit fewer programmed

DSBs, delayed loading of recombination proteins, and increased engagement of error-prone

repair mechanisms (Achache *et al.* 2021; Raices *et al.* 2021).

Sperm also regulate *C. elegans* germline physiology and reproduction. *C. elegans*

hermaphrodites produce sperm only during a late stage in larval development (L'Hernault 2006).

By the third to fourth day of adulthood, these sperm are depleted, which leads to a premature

cessation of reproduction in *C. elegans* hermaphrodites (Luo *et al.* 2010). Sperm depletion also

induces broad transcriptional remodeling independent of aging processes, resulting in a 'female-

like' transcriptional profile (Angeles-Albores *et al.* 2017). Mating extends the hermaphrodite

reproductive span on average to the sixth day of adulthood, after which declining oocyte quality

limits fertility (Luo *et al.* 2010). Mating and even exposure to males, however, also induces

deleterious responses in hermaphrodites leading to premature demise (Maures *et al.* 2014; Shi

and Murphy 2014). It remains unknown how reproductive aging, signaling induced by the

presence or depletion of sperm, and mating intersect to regulate meiotic processes in aged *C.*

elegans germlines.

To define DNA repair defects which are specific to reproductive aging, we assayed

levels of DSB formation and repair in the meiotic oocytes of aged mated and unmated *C.*

elegans hermaphrodites, as well as feminized germline mutants that do not produce sperm (*fog-*

2 mutants). We demonstrate that while the depletion of sperm downregulates DSB induction in

aged germlines, delayed DSB repair is a shared feature of aging germlines independent of

sperm presence. Finally, we identify the ubiquitin E2 ligase variant protein UEV-2 as a putative

regulator of DNA repair during germline aging. Taken together, our work distinguishes DNA

repair defects specific to reproductive aging and identifies parallel mechanisms regulating

gamete quality in the immortal germline.

Methods

***Caenorhabditis elegans* strains and maintenance**

Caenorhabditis elegans strains were maintained at 20°C on nematode growth medium (NGM)

plates seeded with OP50 *Escherichia coli* bacteria. All experiments were performed in the N2

genetic background. Strains used in this experiment include AV761 (*GFP::cosa-1* II; *spo-*

11(me44) IV/ nT1[qIs51]), AV676 (*GFP::cosa-1* II; *fog-2(q71)* V), N2 (wildtype), CB4108 (*fog-*

2(q71) V*)*, DLW135 (*uev-2(gk960600gk429008gk429009); rgr-1(gk429013)* III), DLW199

(libIs4[*pie-1p::uev-2::unc-54 3'UTR*] III:7007600), N2 (wild type), VC30168 (Million Mutation

Project strain carrying *uev-2(gk960600)*), and WBM1119 (wbmIs60 [*pie-1p::3XFLAG::dpy-10*

crRNA::unc-54 3'UTR] (III:7007600)).

In experiments with aged animals, L4 hermaphrodites were isolated and maintained on NGM

plates seeded with OP50 in the absence of males. Strains which produced self progeny were

transferred to new NGM plates seeded with OP50 2 days post-L4 to prevent overconsumption

of food from F1 progeny. At this transfer, if the experimental cohort was to be mated, young

adult male N2 worms were additionally added to these plates at a ratio of ~1.5-2 males per

hermaphrodite. Mated hermaphrodites were again transferred to new NGM plates with OP50

~20-26 hours after males were added and male animals were discarded.

Strain DLW135 was generated by backcrossing VC30168 to N2 10 times. VC30168 was

created by the Million Mutations Project (Thompson *et al.* 2013) and carried many mutations in

addition to the *uev-2(gk960600)* allele of interest. Following backcrossing, mutations on

Chromosomes I, II, IV, V, and X were assumed to have been eliminated. To determine the

success of backcrossing on removing undesired mutations *in cis* with *uev-2* on Chromosome III,

we assessed the presence of known flanking mutations to *uev-2(gk960600gk429008gk429009)*.

Presence of the upstream most proximal genic mutation to *uev-2*, *pho-9(gk429005)*, was

assessed via PCR amplification using OneTaq 2x Master Mix (forward primer DLO1142 5'-

ACCCATTTCCCATTCAATCA-3' reverse primer DLO1143 5'-TTGTAATCTGCCCCAAAAGG-3')

and subsequent HpaII restriction digest (New England Biolabs). DLW135 carried a wild type

allele of *pho-9*, indicating that the region of Chromosome III upstream of *uev-2* was successfully

reverted to wild-type sequence by recombination. However, the closely linked (~1 cM)

downstream allele *rgr-1(gk429013)* was preserved in DLW135, as confirmed by Sanger

sequencing (Sequetech) of a PCR amplified region of the *rgr-1* locus using OneTaq 2x Master

Mix (forward primer DLO1140 5'-TGGAATGGGACTTCCTCTTG-3' reverse primer DLO1141 5'-

TTTCCAAAAGCCAGGACATC-3') isolated using a GeneJET PCR Purification kit

(ThermoFisher). The *rgr-1(gk429013)* allele is a single base pair substitution resulting in a

S360N missense mutation. RGR-1 is a Mediator complex subunit involved in transcriptional

activation that is required for embryonic viability (Shim *et al.* 2002). S360N does not disrupt a

predicted functional domain, and mutants carrying *rgr-1(gk429013)* survive embryogenesis and

are fertile, indicating that this mutation does not severely disrupt function of the RGR-1 protein.

As RGR-1 is not known to play a role in DNA damage repair, and *uev-2* has been previously

demonstrated to modulate germline sensitivity to DNA damage (Luo *et al.* 2010), the

phenotypes we observed using DLW135 in this manuscript are not best explained by the

presence of the *rgr-1(gk429103)* mutation. For simplicity, DLW135 mutants are referred to as

'*uev-2* mutants' in the text of this manuscript.

CRISPR/Cas9 genome editing

Strain DLW199 was generated using the SKILODGE transgenic system (Silva-García *et al.*

2019). WBM1119 was injected with 40ng/μL pRF4 purified plasmid, 40ng/μL purified PCR

amplicon of the full *uev-2* coding sequence with 35bp homology arms to the wbmIs60 landing

site (Phusion polymerase, forward primer DLO1144 5'-

tcccaaacaattaaaaatcaaattttcttttccagATGCGAAGACGTAGCAACAG-3' reverse primer DLO1154

5'-taattggacttagaagtcagaggcacgggcgcgagatgTTAGTTTTCGATGTCAATTGGT-3'), 0.25 μg/μL

Cas9 enzyme (IDT), 100ng/μL tracrRNA (IDT), and 56ng/μL crRNA DLR002 (5'-

GCUACCAUAGGCACCACGAG-3'). Dpy F1 progeny were isolated and screened for insertion

at the wbmIs60 locus by PCR following the SKILODGE recommended protocols (primers

CGSG130, CGSG117 (Silva-García *et al.* 2019)).

The candidate insertion identified among progeny from the above injected hermaphrodites

contained an undesired additional 43bp of sequence between the 5' 3xFLAG tag of the edited

wbmIs60 landing site and the start codon of the *uev-2* coding sequence. The strain carrying this

insertion allele was backcrossed 3x to N2 and was CRISPR/Cas9 edited again to remove the

undesired 5' sequence. Worms were injected with 0.25 μg/μL Cas9 (IDT), 100ng/μL tracrRNA

(IDT), 28ng/μL gRNA DLR022 (5'-GAUCUUUAUAAUCACCGUCA-3'), 28ng/μL gRNA DLR023

(5'-UGUUGCUACGUCUUCGCAUC-3'), 25ng/μL ssODN donor DLO1173 (5'-

AACAATTAAAAATCAAATTTTCTTTTCCAGATGCGGAGGCGAAGTAATAGACAATATGTTGA

TCTCTCATATTTTCGCGAAAC-3'), and 40ng/μL purified pRF4 plasmid. Successful removal of

the 3xFLAG sequence and undesired 43bp inserted sequence were confirmed by PCR and

Sanger sequencing (Sequetech).

Nematode irradiation

C. elegans worms were maintained at 20°C on NGM plates seeded with OP50 prior to and

following irradiation. Irradiation was performed using a Cs^{137} source (University of Oregon).

Immunofluorescence sample preparation and microscopy

Immunofluorescence samples were prepared as in (Libuda *et al.* 2013). Nematodes were

dissected in 1x Egg Buffer (118 mM NaCl, 48 mM KCl_2, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 25 mM

HEPES pH7.4, 0.1% Tween20) and were fixed in 1x Egg Buffer with 1% paraformaldehyde for 5

min on a SuperFrost Plus slide (VWR). Slides were then flash frozen in liquid nitrogen and the

cover slip was removed before the slides were placed in ice cold methanol for 1 minute. Slides

were washed in 1xPBST (1x PBS, 0.1% Tween20) 3x for 10 minutes before they were placed in

Block (1xPBST with 0.7% bovine serum albumin) for a minimum of one hour. 50μL of primary

antibody diluted in PBST (see below for specific antibody dilutions) was then placed on each

slide and samples were incubated for 16-18hrs in a dark humidifying chamber with parafilm

coverslips. Slides were then washed 3x in 1xPBST for 10 minutes. 50μL of secondary antibody

diluted 1:200 in PBST were then added to each sample and slides were incubated for 2hr in a

dark humidifying chamber with parafilm coverslips. Slides were washed 3x in 1xPBST for 10

minutes, and then 50μL of 2μg/mL DAPI was applied to each slide. Samples were incubated in

a dark humidifying chamber with parafilm coverslips for 5 minutes, then were washed 1x in

PBST for 5 minutes. Slides were mounted with a No 1.5 coverslip (VWR) and sealed with nail

polish. All slides were maintained at 4°C until imaging.

Immunofluorescence images were acquired at 512x512 or 1024x1024 pixel dimensions on an

Applied Precision DeltaVision microscope with a 63x lens and a 1.5x optivar. All images were

acquired in 3 dimensions with Z-stacks at 0.2μm intervals. In a minority of aged unirradiated

germlines, we noted that most nuclei in mid-late pachytene exhibited high levels of RAD-51 and

condensed DNA morphology characteristic of apoptosis. These aberrant gonads were not

included in our analyses. Images were deconvolved with Applied Precision softWoRx software

and individual image tiles were stitched into a single image for analysis using the Grid/Collection

Stitching module in Fiji with regression threshold 0.7 (Preibisch *et al.* 2009) or using Imaris

Stitcher software (Bitplane).

Specific antibodies used and their dilution factors are: Rabbit αRAD-51 (1:500), Chicken αRAD-

51 (1:1000), Rabbit αDSB-2 (1:5000) (Rosu *et al.* 2013), Rabbit αGFP (1:1500), Chicken

αGFP(1:1000), Alexa Fluor 488 Goat αChicken (1:200), Alexa Fluor 555 Goat αChicken (1:200),

Alexa Fluor 555 Goat αRabbit (1:200), and Alexa Fluor 488 Goat αRabbit (1:200).

Image analysis and quantification

Images were analyzed as described in (Toraason *et al.* 2021). Image quantification was

performed using Imaris software (Bitplane). Individual nuclei within stitched gonads were

identified as Surface objects (Smooth 0.1-0.15, Background 3-4, Seed Point Diameter 3-4)

based on DAPI staining intensity. Manual thresholding of specific values were used per gonad

to generate surfaces which represented the nuclei observed. Defined surfaces were then split to

designate individual nuclei using the Imaris Surfaces Split module. Nuclei which were partially

imaged or overlapped with other nuclei were eliminated from the analysis. RAD-51 foci were

defined as Spot objects (Estimated XY Diameter 0.1, Model PSF-elongation 1.37, Background

Subtraction enabled). To determine the number of RAD-51 foci per nucleus, we either utilized

the "Find Spots Close to Surface" MATLAB module (Threshold value 0.1) or utilized the "Closest

Distance to Surface" statistic calculated by Imaris to find the number of Spots ≤0.1μm distant

from nuclei. The length of each germline was defined using the Imaris Measurements tool.

Measurement points were specifically placed at the beginning of the premeiotic tip and the end

of pachytene. For germlines which had a defined transition zone by DAPI morphology, points

were also placed at the start and end of the transition zone.

Nuclei positions were transformed from 3D coordinates to a linear order using the Gonad

Linearization Algorithm implemented in R (Toraason *et al.* 2021). Gonad length in germlines

which lacked a defined transition zone (e.g. *fog-2* mutants, Supplemental Figure 2) was

normalized to the distance from the premeiotic tip to the end of pachytene, where the premeiotic

tip begins at position 0 and the end of pachytene is at position 1. In all other germlines, the

gonad length was normalized to pachytene, where the beginning of pachytene is position 0 and

the end of pachytene is position 1.

Germline DSB-2 staining was analyzed in Imaris using germlines stitched in Fiji as described

above. The length of the germline was defined using the Imaris Measurements tool. Specific

points were placed at the beginning of the transition zone, end of the transition zone, beginning

of the DSB-2 zone (defined as the row of nuclei in which most nuclei had DSB-2 staining), the

end of the DSB-2 zone, the final position of one or more nuclei which had DSB-2 staining, and

the end of pachytene. The measured distances were then normalized to pachytene, where the

beginning of pachytene is position 0 and the end of pachytene is position 1.

GFP::COSA-1 foci were quantified manually from late pachytene nuclei from 3D z-stacks using

Fiji. Nuclei that were completely contained within the image stack were quantified in the last few

rows of the late pachytene region in which all nuclei displayed bright GFP::COSA-1 foci (last ~3-

6 rows of late pachytene in old *fog-2* germlines, last ~6-8 rows of late pachytene in young *fog-2*

germlines).

fog-2 **Brood Viability Assay**

C. elegans worms were maintained at 20°C during fertility assays. Feminized *fog-2* mutants

were synchronized in age by placing gravid mated CB4108 females onto an NGM plate seeded

with OP50 for one hour. Hatched female progeny were isolated as L4s from these plates and

were kept in isolation from males to prevent mating. At adult day 1, 2, 3, 4, or 5, these isolated

fog-2 females were then placed on individual plates with n=2 young adult N2 males each. Mated

fog-2 females were then subsequently transferred to new NGM plates seeded with OP50 with

young adult N2 males at either 6hr, 12hr, 18hr, 24hr, and 48hr after the first mating, or at 24hr

and 48hr after the first mating. 72hr after the first mating, adult females were discarded. Plates

were scored ~24hr after the parent female was removed for hatched progeny, dead eggs, and

unfertilized oocytes. Brood viability was calculated as (hatched progeny) / (hatched progeny +

dead eggs). Fertility assays were replicated twice with n=5 females of each age group assayed

per replicate.

During the course of the brood viability assays, some mated *fog-2* females exhibited matricidal

hatching. This phenotype was more pronounced in aged worms, consistent with previous work

which showed that matricidal hatching is exacerbated with maternal age (Pickett and Kornfeld

2013). Only eggs which were successfully ovulated were scored in the assay.

Statistics

All statistics were calculated in R (v4.0.3). Data wrangling was performed using the Tidyverse

package (v1.3.0) (Wickham *et al.* 2019). Specific statistical tests used are denoted in the figure

legends and text. P values were adjusted for multiple comparisons when appropriate. If 3

pairwise comparisons were being performed, Bonferroni correction was applied. If >3 pairwise

comparisons were performed, Holm-Bonferroni correction was instead applied to reduce the risk

of type II statistical errors.

Data and Code Availability

The gonad linearization algorithm is available on the Libuda Lab GitHub

<github.com/libudalab/Gonad-Analysis-Pipeline> and on the Libuda Lab website

<libudalab.org>

<u>Results</u>

Meiotic DNA break levels are influenced by both aging and sperm depletion

To determine the relative contributions of reproductive aging and sperm depletion to

DNA break repair dynamics in the *C. elegans* germline, we examined DNA break levels in the

oocytes of aged hermaphrodites which were mated (to prevent sperm depletion) or unmated (to

permit sperm depletion) (Figure 1A). DNA breaks were quantified using immunofluorescence to

visualize the recombinase RAD-51, which marks DSBs designated for repair by recombination

(Gartner and Engebrecht 2022). While the cytological appearance of RAD-51 foci indicates the

occurrence of DSBs, disappearance of RAD-51 foci indicates progression of a DSB event

further through a DSB repair pathway. "Young" germlines were isolated from N2 hermaphrodites

on the first day of adulthood (1 day post-L4, Figure 1A), while "aged" germlines were isolated

from N2 hermaphrodites on their fourth day of adulthood (4 days post-L4, Figure 1A). Aged

hermaphrodites were maintained either unmated to males, or mated with males from their

second to third day post-L4 larval stage (Figure 1A, see Methods).

To quantify the profile of DSB induction and repair across prophase I, we counted the

number of RAD-51 foci per nucleus in oocytes from young and aged animals throughout the

germline (see Methods). Under normal conditions, RAD-51 foci accumulate within nuclei

following DSB induction by the conserved endonuclease SPO-11 in early pachytene (Dernburg

et al. 1998; Colaiácovo *et al.* 2003). Then, as nuclei progress through mid and late pachytene,

these RAD-51 foci decline in number as DSBs are repaired (Colaiácovo *et al.* 2003). During

early pachytene, the amount of RAD-51 foci per nucleus was similar between aged mated

germlines and young germlines (Figure 1B-C, Bin 2 Mann-Whitney U test p=0.258). Young

germlines, however, accumulated a higher total number of RAD-51 foci per nucleus (Figure 1B-

C, Bin 3 Mann-Whitney U test p=0.005), suggesting that DSB induction or RAD-51 loading is

slightly compromised in aged mated germlines. We further noted that RAD-51 foci in aged

unmated germlines were greatly decreased throughout early pachytene as compared to both

young and aged mated germlines (Figure 1B-C, Bins 2-3 Mann-Whitney U test p<0.001),

indicating that sperm depletion in aged germlines may affect meiotic DSB induction and/or RAD-

51 loading.

In contrast to early pachytene, nuclei throughout mid pachytene from aged mated

germlines maintained higher levels of RAD-51 than young germlines (Figure 1B-C, Bins 4-5

Mann-Whitney U test p<0.05). We observed a similar effect in the aged unmated germlines,

which also displayed elevated numbers of RAD-51 foci relative to young germlines throughout

mid pachytene (Figure 1B-C, Bins 4-5 Mann-Whitney U test p<0.001). Thus, DSB repair at mid-

pachytene may be delayed in aging germlines regardless of mating or sperm depletion. Notably,

by late pachytene the number of RAD-51 foci per nucleus converged between young, aged

mated, and aged unmated germlines (Figure 1B-C, Bin 6 Mann-Whitney U test p>0.05),

indicating that ultimately all DSBs can be repaired or minimally offload RAD-51 in aged

germlines. Taken together, our results suggest that parallel mechanisms may regulate DNA

break levels in aged *C. elegans* germlines: 1) depletion of sperm downregulates DSB induction

and/or RAD-51 loading; and, 2) reproductive aging delays RAD-51 foci unloading at mid

pachytene.

To determine if the persistent RAD-51 foci in aged mated and unmated germlines were

derived from the programmed meiotic DSBs, we also examined RAD-51 foci in *spo-11(me44)*

null mutants, which do not form meiotic DSBs (Supplemental Figure 1) (Colaiácovo *et al.* 2003).

We did not observe a notable increase in nuclei with RAD-51 foci in aged *spo-11* germlines,

indicating that the persistent RAD-51 foci present at mid pachytene in aged wildtype gonads are

likely derived from normal meiotic functions and processes, such as SPO-11 activity.

Nuclei which are competent for DSB induction in the *C. elegans* germline have their

chromatin marked with the protein DSB-2 (Rosu *et al.* 2013). To assess if the altered

accumulation of DSBs which we observed in aged unmated germlines coincided with a change

in competency for DSB induction, we quantified the extent of young and aged germlines in

which ≥50% of nuclei exhibited DSB-2 staining (the "DSB-2 zone", Supplemental Figure 2A-B).

DSB-2 accumulates on meiotic chromatin beginning in the transition zone (leptotene/zygotene)

and is offloaded from the majority of nuclei by mid pachytene (Rosu *et al.* 2013; Toraason *et al.*

2021). Mutants which incur errors in crossover formation, however, maintain DSB-2 on meiotic

chromatin later into pachytene (Rosu *et al.* 2013). While the length of the DSB-2 zone was only

subtly altered in aged mated germlines (Supplemental Figure 2B-C, Mann-Whitney U test

p=0.027), the DSB-2 zone persisted later into pachytene in aged unmated germlines relative to

young germlines (Supplemental Figure 2B-C, Mann-Whitney U test p=0.008). Thus, our data

indicate that the extent of DSB-2 marked pachytene nuclei is influenced both by aging and by

the absence of sperm.

Meiotic DNA breaks are elevated in aged feminized germlines

To uncouple the relationship between sperm depletion and reproductive aging in

regulating DSB induction and repair, we examined RAD-51 levels in germlines which have

never been impacted by sperm or mating. Hermaphrodites carrying the *fog-2(q71)* mutation do

not produce sperm during larval development but proliferate a full adult complement of oocytes

(Schedl and Kimble 1988), rendering them "obligate females." Due to the absence of signaling

from sperm in *fog-2* mutants, both germline stem cell proliferation and meiotic progression are

halted, such that meiotic oocytes are held within the gonad (Mccarter *et al.* 1999; Cinquin *et al.*

2016). Nonetheless, feminized mutants undergo reproductive senescence and exhibit reduced

oocyte quality with age (Supplemental Figure 3; (Andux and Ellis 2008; Luo *et al.* 2009))

We analyzed the levels of RAD-51 foci in oocyte nuclei from young (1 day post-L4),

aged (4 days post-L4), and old (6 days post-L4) *fog-2* germlines (Figure 2A-B). During our

experiments we noted that the cytologically distinctive transition zone, which demarcates

meiotic entry and is composed of nuclei undergoing active chromosome movement to facilitate

pairing, was dramatically reduced in aged *fog-2* germlines (Supplemental Figure 4). Previous

work has shown that mitotic germ cell proliferation is reduced in feminized and sperm-depleted

germlines (Cinquin *et al.* 2016). Thus, the absence of a transition zone in aged *fog-2* germlines

may be the product of two parallel effects: 1) nuclei in the transition zone completing the pairing

process and therefore exhibiting the classic "cage-like" morphology of paired chromosomes

found in pachytene nuclei; and, 2) decreased proliferation also limiting the number of new nuclei

which enter meiosis. This lack of the transition zone in aged *fog-2* germlines presented a

challenge for staging meiotic nuclei to make comparisons between young and aged gonads. To

quantify RAD-51 levels in *fog-2* germlines independent of meiotic stages, we normalized the

germline length with position 0 at the premeiotic tip and position 1 at the end of pachytene and

used a sliding window to assay RAD-51 foci within the germline (Figure 2D, see Methods)

(Toraason *et al.* 2021). To describe the RAD-51 profile of aging *fog-2* germlines, we calculated

two metrics: 1) the "RAD-51 zone" indicating the extent of the germline which contained nuclei

with RAD-51 foci; and, 2) the "peak RAD-51 window" indicating the maximum levels of RAD-51

within the germlines.

To assess whether the proportion of germline nuclei with RAD-51 foci was altered in

aging *fog-2* germlines, we calculated the "RAD-51 zone" of each age group, which was defined

as the germline distance extending from the most distal (near the premeiotic tip) to the most

proximal (near the end of pachytene) windows in which at least 50% of nuclei had one or more

RAD-51 foci (Figure 2B, 2D). We found that the RAD-51 zone extended more distally in the

germline in aged and old *fog-2* animals as compared to young germlines (Figure 2B, 2D). This

distal expansion of the RAD-51 zone can likely be explained by transition zone nuclei in young

germlines completing the pairing process and entering pachytene as the germline ages. In

contrast, the proximal end of the RAD-51 zone only subtly shifted distally in aged and old

germlines (Figure 2B, 2D). This result indicates that later prophase I nuclei within aged *fog-2*

germlines continue to either maintain or induce RAD-51 marked DSBs.

To determine if the number of RAD-51 marked DSBs in *fog-2* germline nuclei were

altered with age, we identified the "peak RAD-51 windows" in each age group, defined as the

window in which the mean RAD-51 foci per nucleus was highest (Figure 2C, 2D arrowheads).

We noted that the position of the peak RAD-51 window moved distally in aged and old *fog-2*

germlines (Figure 2D arrowheads), suggesting that the spatial regulation of DSB induction and

repair may change as feminized germlines age. The number of RAD-51 foci per nucleus within

the peak RAD-51 window was not significantly different in aged germlines as compared to

young gonads (Figure 2C-D, Mann-Whitney U test p=1.000). Old germlines, however, exhibited

a significant ~1.5 fold increase in RAD-51 foci per nucleus as compared to young and aged

germlines within the peak RAD-51 window, indicating that *fog-2* mutant germ cells accumulate

RAD-51 foci during aging (Figure 2C-D, Mann-Whitney U test p <0.001). This result notably

differs from aged unmated wildtype germlines, which exhibit reduced DSBs with age (Figure 1B-

C) (Achache *et al.* 2021; Raices *et al.* 2021). Our data therefore support a model in which sperm

depletion, rather than absence of sperm, downregulates meiotic DSB induction.

DSB repair is altered in aged feminized germlines

The accumulation of RAD-51 foci observed in aging *fog-2* germlines may be the product

of: 1) increased induction of DSBs; 2) defects in DSB repair; or, 3) a combination of these

effects. To assess the efficiency of DSB repair during *fog-2* germline aging, we exposed young

(1 day post-L4) and aged (4 days post-L4) *fog-2* mutant females to 5000 Rads of ionizing

radiation (Figure 3A, Supplemental Figure 5), inducing ~118 DSBs per nucleus throughout the

germline (Yokoo *et al.* 2012). We then allowed the animals to age for 2 days to resolve this DNA

damage before assessing germlines for persistent unrepaired DSBs as marked by RAD-51 foci

(Figure 3A). As *fog-2* germlines accumulate DSBs during aging (Figure 2C-D), we established

baseline levels of DNA damage based on comparing RAD-51 foci in animals of equivalent ages

that were never exposed to radiation to the irradiated cohorts (Figure 3A). We noted

considerable inter-nucleus variance in the RAD-51 foci, which persisted following irradiation in

both young and aged germlines (Supplemental Figure 5). This effect was particularly prominent

in the distal germlines of both groups (Supplemental Figure 5C-D), suggesting that a

subpopulation of nuclei in the mitotic germline or early stages of meiosis are uniquely

susceptible to exogenous DNA damage regardless of parental age.

To estimate the residual DSBs derived from irradiation which were not yet repaired two

days post irradiation, we calculated the median number of RAD-51 foci in a sliding window

across the germline (Figure 3B, see Methods) and subtracted the unirradiated median RAD-51

foci from the irradiated median RAD-51 foci in each window (Figure 3C). Both young and aged

germlines maintain high levels of damage in the distal germline following irradiation (germline

position 0.0-0.5, Figure 3C, Supplemental Figure 5C). Nuclei in the proximal region of young

irradiated germlines did not consistently maintain median DNA break levels higher than

baseline; whereas aged irradiated germlines maintained a median elevation of ~6-10 RAD-51

foci per nucleus (germline position 0.5-1.0, Figure 3C). This result indicates that aged *fog-2*

germlines exhibit DNA repair defects specifically in nuclei at later stages of meiotic prophase I.

Taken together, our experiments in "feminized" germlines demonstrate that DNA repair

efficiency is altered in aging germlines independent of any signals from sperm.

Defects in meiotic DSB repair may disrupt the formation of interhomolog crossovers

(Gartner and Engebrecht 2022). To determine if crossover recombination is impeded in aged

fog-2 germlines, we quantified bright GFP::COSA-1 foci, which mark joint molecules designated

to become crossovers at late pachytene (Yokoo *et al.* 2012). Under normal conditions, each of

the six pairs of *C. elegans* homologous chromosomes form one crossover and therefore most

nuclei exhibit only six COSA-1 foci (Yokoo *et al.* 2012). We observed that the number of COSA-

1 foci in late pachytene nuclei of old *fog-2* gonads were altered relative to their young

counterparts (Figure 4A, Chi square test p=0.033), with a subtle but significant increase

specifically in the fraction of nuclei with 7 COSA-1 foci (Figure 4A, Fisher's Exact test p=0.038).

Thus, our data suggests that DSB repair in old *fog-2* germlines may be altered in a manner

which impacts the number of designated crossovers. Further, the late pachytene region of the

germline with bright COSA-1 foci was reduced in old *fog-2* germlines (Figure 4B). This

phenotype is reminiscent of synapsis deficient mutants, which form COSA-1 foci only in a very

small region of late pachytene (Cahoon *et al.* 2019). Previous work has shown that aged

unmated wildtype germlines incur defects in crossover formation which reduce the number of

crossovers formed (Luo *et al.* 2010; Achache *et al.* 2021; Raices *et al.* 2021). Our results in *fog-*

2 mutants therefore demonstrate that these defects are not shared in aged feminized *C.*

elegans germlines, suggesting that sperm presence influences the germ cell capacity for

crossover formation during reproductive aging.

UEV-2 is required for 'youthful' germline DSB repair

To identify proteins which may regulate DSB repair in the aging *C. elegans* germline, we

looked to candidate genes upregulated in long-reproductive *sma-2* mutant oocytes, which

exhibit DNA damage resilience in addition to delayed reproductive senescence (Luo *et al.*

2010). The *sma-2* DNA damage resilience phenotype requires upregulation of the ubiquitin E2

ligase variant UEV-2, suggesting that this protein may promote efficient germline DNA repair

(Luo *et al.* 2010). UEV proteins lack a catalytic cysteine residue conserved in E2 ubiquitin

ligases (Sancho *et al.* 1998) but have been shown to form heterodimeric complexes with other

E2 ubiquitin ligases to influence their function, implying regulatory roles for this protein class

(Vandemark *et al.* 2001; Wijk and Timmers 2010).

To assess the influence of UEV-2 on DSB repair during germline aging, we utilized a

strain carrying the putative null allele *uev-2(gk960600)*, which ablates the translation initiation

site and second exon boundary of the gene (Supplemental Figure 6; see Methods). With the

uev-2 mutant strain, we examined the number of RAD-51 foci in germline nuclei derived from

young (1 day post-L4) or aged (4 days post-L4) animals (Figure 5A). Aged *uev-2* mutants were

also mated to avoid the DSB induction defects associated with sperm depletion (Figure 5A). If

UEV-2 functions to promote efficient DSB repair in young gonads but becomes dysregulated or

loses function during aging, then we would expect *uev-2* mutants to exhibit defects in DSB

repair in young germlines but minimal additional defects in aged germlines. Indeed, when we

compared the levels of RAD-51 observed in young and aged mated wildtype and *uev-2*

germlines, we observed DSB repair defects that did not accumulate with age. In early

pachytene, young and aged *uev-2* mutants exhibited similar levels of RAD-51 to young wildtype

germlines (Figure 5B-C, Bins 2-3 Mann-Whitney U test p>0.05), indicating that UEV-2 is not

required for meiotic DSB induction nor RAD-51 loading. In contrast, at mid pachytene, young

uev-2 mutant germlines maintained elevated RAD-51 foci relative to young wildtype germlines

(Figure 5B-C, Bins 4-5 Mann-Whitney U test $p<0.05$). The specific levels of DSBs at mid

pachytene in young *uev-2* mutants were also indistinguishable from aged wildtype germlines

(Figure 5B-C, Bins 4-5 Mann-Whitney U test $p>0.05$). These results at mid pachytene indicate

that DSB repair is delayed in young *uev-2* mutants to an extent which recapitulates the effect we

observe during wildtype aging. Aged *uev-2* germline RAD-51 levels at mid pachytene were

statistically indistinguishable from either young or aged mated wildtype germlines (Figure 5B-C,

Bins 4-5 Mann-Whitney U test $p>0.05$), suggesting that the *uev-2* mutation does not grossly

exacerbate DSB repair defects with age.

In late pachytene, the specific rates of DSB resolution diverged slightly between young

and aged *uev-2* and wildtype germlines (Figure 5B-C, Bins 6-7 Mann-Whitney U test $p<0.05$),

suggesting that UEV-2-independent and age-specific effects may contribute to DSB resolution

at this meiotic stage. Taken together, our results indicate that loss of *uev-2* in young germlines

is sufficient to phenocopy the mid-pachytene patterns of DSB repair observed in an aged

wildtype context. This observation supports a model in which UEV-2 functions in young

germlines specifically to promote efficient DSB repair.

Overexpression of *uev-2* alters RAD-51 foci levels in an aged oocytes

As loss of *uev-2* in young germlines appeared to "prematurely age" RAD-51 foci

patterns, we hypothesized that overexpression of *uev-2* in aged germlines could ameliorate

persistent RAD-51 foci at mid pachytene. To test this hypothesis, we used CRISPR/Cas9

genome editing to generate a germline-specific overexpression construct of *uev-2* driven by the

pie-1 promoter (*pie-1p::uev-2*, see Methods). We then assessed for the presence of DSBs as

marked by RAD-51 in the germlines of young (1 day post-L4) or aged (4 days post-L4) mated

animals overexpressing UEV-2 and compared those levels to young and aged mated wildtype

germlines (Figure 6A).

At the beginning of early pachytene, both young and aged mated *pie-1p::uev-2* mutants

initially accumulated DSBs at levels similar to young and aged mated wildtype gonads (Figure

6B-C, Bin 2 Mann-Whitney U test p>0.05). However, aged mated *pie-1p::uev-2* mutants

accumulated fewer total DSBs than young wildtype, aged mated wildtype, and young *pie-*

1p::uev-2 germlines (Figure 6B-C, Bin 3 Mann-Whitney U test p<0.05). At mid pachytene, young

pie-1p::uev-2 germlines maintained elevated RAD-51 foci over young wildtype germlines,

suggesting that overexpression of *uev-2* deleteriously impacted DSB repair in this context

(Figure 6B-C, Bin 4-5 Mann Whitney U test p≤0.001). This effect was not preserved in aged

mated *pie-1p::uev-2* germlines, which exhibited similar DSB levels as young wildtype germlines

at the beginning of mid pachytene and slightly elevated foci at the end of mid pachytene (Figure

6B-C, Mann-Whitney U test Bin 4 p=0.156 Bin 5 p=0.023).

Throughout late pachytene, young *pie-1p::uev-2* germlines maintained subtle but

significantly elevated DSBs relative to young and aged wildtype germlines (Figure 6B-C, Bin 6-7

Mann-Whitney U test p<0.05). Conversely, aged mated *pie-1p::uev-2* germlines maintained

significantly fewer RAD-51 foci throughout late pachytene than young wildtype or young *pie-*

1p::uev-2 germlines (Figure 6B-C, Bin 6-7 Mann-Whitney U test p<0.05). Taken together, these

data suggest that UEV-2 is not the sole regulator of DSB repair efficiency during *C. elegans*

germline aging and appears to have age-dependent functions in regulating meiotic DSB

accumulation and repair.

Discussion

C. elegans germline function is impacted both by reproductive aging and sperm signals.

Our study demonstrates that aged *C. elegans* germlines exhibit delayed DSB repair in mid-

pachytene regardless of mated status, suggesting that deficiencies in germline DNA repair are a

product of reproductive aging. We further find that sperm depletion, but not absence of sperm,

reduces RAD-51 marked DSBs at early pachytene, suggesting that loss of signals from sperm

downregulate DSB induction. Taken together, our study supports a model in which signals due

to sperm depletion and reproductive aging operate in parallel to influence meiotic DSB induction

and repair (Figure 7).

Sperm depletion and DSB induction

Our data indicate that aged unmated germlines exhibit dramatically reduced RAD-51 foci

in early pachytene (Figure 7). Previous work has similarly reported that unmated

hermaphrodites induce fewer DSBs with age (Achache *et al.* 2021; Raices *et al.* 2021). We find,

however, that mating is sufficient to rescue RAD-51 foci accumulation at early pachytene in

aged germlines. This effect in mated hermaphrodites may be due to cues from sperm-specific

signals, seminal fluid components, or male pheromones, all of which impact hermaphrodite

physiology (Shi and Murphy 2014; Aprison *et al.* 2022). While our study cannot distinguish

between these male- and mating-dependent effects, it is notable that aged feminized *fog-2*

germlines, which have never been exposed to sperm or males, do not exhibit reduced RAD-51

marked DSBs during aging. Thus, we propose that DSB induction in aged germlines is primarily

repressed by signals caused specifically from sperm depletion rather than reproductive aging,

mating-induced signaling, or exposure to males. We further found that *spo-11(me44)* mutants

do not exhibit increased DSBs with age, suggesting that the DSBs observed in aged gonads

come from the endogenous meiotic machinery. This result contrasts with previous work done

using *spo-11(ok79)* mutants, which incur SPO-11 independent DSBs with age (Raices *et al.*

2021). Thus, our data raises the possibility that the specific nature of *spo-11* mutation or

background strain-specific effects influence the occurrence of exogenous DSBs in aged

germlines.

Why might sperm depleted hermaphrodites downregulate germline DSB induction?

Recent evidence has unveiled a potential transition in hermaphrodite gonad function following

sperm depletion (Kern *et al.* 2021). After all sperm have been utilized from the spermatheca,

hermaphrodites continue to lay unfertilized oocytes and secrete a nutrient-rich yolk in what has

been suggested to be a form of 'primitive lactation' (Kern *et al.* 2021). Thus, the reduction in

DSBs induced in germ cells following sperm depletion may be a product of the hermaphrodite

germline functionally changing from a reproductive organ to a system which produces food for

offspring. Reduced DSB formation, then, may be indicative of the metabolic resources of the

hermaphrodite being reallocated in favor of providing nutritional supplement for progeny.

Multiple mutants in *C. elegans* have been reported to exhibit age-dependent decline in

meiotic DSB induction (Tang *et al.* 2010; Rosu *et al.* 2013). Our research raises the possibility

that these proteins may mediate or respond to signals from sperm. Also, both reproductive

aging and *C. elegans* yolk secretion are regulated by insulin/insulin-like growth factor signaling

(Luo *et al.* 2010; Kern *et al.* 2021). How reproductive aging and sperm depletion signals are

integrated through this pathway to enact distinct phenotypes that impact germline function

remains unknown but opens an avenue for future investigation. In summary, we have

illuminated a regulatory mechanism specifically associated with sperm depletion which

downregulates DSB induction in the *C. elegans* germline.

DSB repair and *C. elegans* reproductive aging

Aged *C. elegans* germlines exhibit multiple DNA repair defects, including delays in

recombination protein loading and increased engagement of error-prone repair mechanisms

(Raices *et al.* 2021). Both RAD-51 loading and error prone pathway engagement are regulated

by DSB end resection (Gartner and Engebrecht 2022), suggesting that differences in DSB

repair during aging may be derived from defects at this DNA processing step. We demonstrate

that the E2 ligase variant UEV-2 is required for 'youthful' patterns of RAD-51 foci resolution

during mid pachytene, indicating that a loss of UEV-2 or an age-related change in its function

may underly the DNA repair defects in aged germlines. However, overexpression of UEV-2 is

not sufficient to rescue persistent RAD-51 foci at mid pachytene in aged germlines and instead

introduces DSB repair defects in young germlines. These results suggest that the specific

levels of *uev-2* expression or the co-expression of other proteins may be important for the

function of UEV-2 in DNA repair processes.

While the specific molecular functions of UEV-2 remain unknown, previous yeast two-

hybrid assays have evidenced that UEV-2 may interact with BRC-1, the *C. elegans* BRCA1

homolog (Gudgen *et al.* 2004). BRCA1 is an E3 ubiquitin ligase thought to regulate many DNA

repair steps, including DSB resection (Cruz-García *et al.* 2014). Recent studies have

demonstrated that BRC-1 is vital for preventing error prone repair in the *C. elegans* germline (Li

et al. 2020; Kamp *et al.* 2020). Given these results, we propose that UEV-2 may modulate BRC-

1 activity in the germline to regulate resection of DSBs and promote efficient recombination.

Under this model, overexpression of *uev-2* or loss of its function may cause hyper- or hypo-DSB

resection respectively, and thus have a deleterious impact on the efficiency of recombination.

Taken together, our work demonstrates that UEV-2 is involved in regulating efficient and

'youthful' meiotic DSB repair, thereby opening avenues to future work uncovering the specific

roles this protein plays in meiosis.

<u>Acknowledgements</u>

We thank the CGC (funded by National Institutes of Health (NIH) P40 OD010440) for strains.

We thank C. Cahoon, A. Naftaly, and N. Kurhanewicz for thoughtful comments on this

manuscript. This work was supported by the National Institutes of Health T32GM007413 and

Advancing Science in America (ARCS) Foundation Award to E.T., and National Institutes of

Health R00HD076165 and R35GM128890 to D.E.L. D.E.L. is also a recipient of a March of

Dimes Basil O'Connor Starter Scholar award and Searle Scholar Award.

<u>Competing Interests</u>

The authors declare no competing interests.

Figures

Figure 1

Figure 1. DNA double strand break levels are altered during *C. elegans* germline aging. A)

Schemes used to isolate young (1 day post-L4) and aged (4 days post L4) worms for

experiments. Days count ~18-24 hour periods after hermaphrodites were isolated as L4 larvae

and are separated by alternating grey shaded boxes. B) Representative images of RAD-51 foci

meiotic nuclei in young and aged germlines. Scale bars represent 5µm. C) RAD-51 foci per

nucleus in oocytes. Line plots represent the mean RAD-51 foci per nucleus along the length of

the germline in a sliding window encompassing 0.1 units of normalized germline distance with a

step size of 0.01 germline distance units. Mean RAD-51 foci were calculated from nuclei

analyzed in n=9 total germlines derived from ≥3 experimental replicates within each age group.

Shaded areas around each line represent ± SEM. Total nuclei analyzed (Bins 1/2/3/4/5/6/7)

Young: 185/117/146/107/117/97/83; Aged Mated: 234/205/192/173/154/177/96; Aged Unmated:

268/147/161/175/191/152/110. Germlines distances were normalized to the start (0) and end (1)

of pachytene based on DAPI morphology (see Methods). For analysis, the germline was divided

into 7 bins encompassing the transition zone (Bin 1), early pachytene (Bins 2-3), mid pachytene

(Bins 4-5), and late pachytene (Bins 6-7). The germline positions at which each bin start and

end are marked on the X axis as vertical grey lines. Heat maps below each bin display the p

values of pairwise comparisons of RAD-51 foci per nucleus counts within that bin. P values

were calculated using Mann-Whitney U tests with Bonferroni correction for multiple

comparisons. Blue tiles indicate significant differences (adjusted p value <0.05) and grey tiles

indicate nonsignificant effects (adjusted p value >0.05).

Figure 2

Figure 2. DNA double strand break levels increase with age in *fog-2* feminized germlines.

A) Schemes used to isolate young (1 day post-L4), aged (4 days post L4), and old (6 days post

L4) *fog-2(q71)* worms for experiments. Days count ~18-24 hour periods after hermaphrodites

were isolated as L4 larvae and are separated by alternating grey shaded boxes. B)

Representative whole germline images of young, aged, and old *fog-2* germlines. The RAD-51

zone, defined as the region of the germline in which the majority of nuclei have one or more

RAD-51 foci, is indicated with a green dashed line. All germlines are oriented with the distal

mitotic tip on the left and the end of pachytene on the right. Scale bars represent 20µm. Grey

numbered boxes indicate the positions of the images presented in panel C. C) Representative

images of the peak levels of RAD-51 foci observed in meiotic nuclei of young, aged, and old

fog-2 germlines. Scale bars represent 5µm. Each panel is numbered to indicate the position in

the germlines displayed in panel B that each inset was taken from. D) RAD-51 foci per nucleus

in *fog-2(q71)* oocytes. Line plots represents the mean RAD-51 foci per nucleus along the length

of the germline in a sliding window encompassing 0.1 units of normalized germline distance with

a step size of 0.01 germline distance units. Mean RAD-51 foci were calculated from nuclei

analyzed in n=9 total germlines derived from ≥3 experimental replicates within each age group.

Shaded areas around each line represent ± SEM. Average nuclei quantified in each bin ±

standard deviation: Young *fog-2* 173±21.3, Aged *fog-2* 143±32.0, Old *fog-2* 148.5±29.6.

Germline distance was normalized to the premeiotic tip (0) and end of pachytene (1) based on

DAPI morphology (see Methods). Arrowheads indicate the "peak RAD-51" windows, defined as

the windows along the length of the germline of each age group with the highest RAD-51 foci

per nucleus. P values were calculated by Mann-Whitney U test comparisons of RAD-51 counts

within these peak windows with Bonferroni correction for multiple comparisons (n.s. = p>0.05,

*** = p<0.001). Vertical dotted lines indicate the distal and proximal bounds of the RAD-51 zone

for each age group, defined as windows in which the median RAD-51 foci per nucleus count

was ≥1.

Figure 3. DNA double strand break repair is disrupted in aged *fog-2* feminized germlines.

A) Schemes used to isolate young and aged *fog-2(q71)* worms for experiments. Days count

~18-24 hour periods after hermaphrodites were isolated as L4 larvae and are separated by

alternating grey shaded boxes. Irradiated (IR) germlines were exposed to ionizing radiation at

the first or fourth day of adulthood and were allowed to recover for two days before analysis

(see Methods). Unirradiated germlines (No IR) were never exposed to radiation. B) RAD-51 foci

per nucleus in irradiated (IR) and unirradiated (no IR) oocytes. Line plots represents the mean

RAD-51 foci per nucleus along the length of the germline in a sliding window encompassing 0.1

units of normalized germline distance with a step size of 0.01 germline distance units. Plots in

panel B share an X axis with the plot in panel C. Mean RAD-51 foci were calculated from nuclei

analyzed in n=9 total germlines derived from ≥3 experimental replicates within each age group.

Shaded areas around each line represent ± SEM. Average nuclei quantified in each bin ±

standard deviation: Young *fog-2* No IR 141.6±26.6, Young *fog-2* IR 134.4±27.9, Old *fog-2* No IR

148.5±29.6, Old *fog-2* IR 142.4±30. Germline distance was normalized to the premeiotic tip (0)

and end of pachytene (1) based on DAPI morphology (see Methods). Representative images of

young and aged IR and No IR germlines are displayed in Supplemental Figure 5. C) Median

RAD-51 foci per nucleus in irradiated germlines above median levels in unirradiated germlines

of the same age (calculated as median RAD-51 foci in IR gonads – median RAD-51 foci in non-

IR gonads within each window along the length of the germline).

Figure 4. Crossover designation is altered in aged *fog-2* germlines. A) Stacked bar plot of the percent of nuclei in young (1 day post L4) and old (6 days post L4) *fog-2* mutants with the given number of GFP::COSA-1 foci. The p value displayed was calculated by Chi Square test. N values indicate the number of nuclei scored. B) Representative images of GFP::COSA-1 localization in young and old *fog-2* mutant germlines. Scale bars represent 20μm. Germlines are oriented with the proximal end on the left and distal end on the right. Vertical low opacity white lines designate the end of pachytene.

Figure 5

Figure 5. UEV-2 is required for 'youthful' DNA repair. A) Schemes used to isolate young (1 day post-L4) and aged (4 days post L4) *uev-2* mutant worms for experiments. Days count ~18-24 hour periods after hermaphrodites were isolated as L4 larvae and are separated by alternating grey shaded boxes. B) Representative images of RAD-51 foci in meiotic nuclei of

young *uev-2*, aged mated *uev-2*, young wildtype, and aged mated wildtype germlines. Scale

bars represent 5µm. C) RAD-51 foci per nucleus in oocytes. Line plots represent the mean

RAD-51 foci per nucleus along the length of the germline in a sliding window encompassing 0.1

units of normalized germline distance with a step size of 0.01 germline distance units. Mean

RAD-51 foci were calculated from nuclei analyzed in n=9 total germlines derived from ≥3

experimental replicates within each age and genotype group. Shaded areas around each line

represent ± SEM. Total nuclei analyzed (Bins 1/2/3/4/5/6/7) Young Wildtype:

185/117/146/107/117/97/83; Aged Wildtype Mated: 234/205/192/173/154/177/96; Young *uev-2*:

186/135/134/113/112/132/95; Aged *uev-2* Mated: 129/161/155/158/167/142/120. Germlines

distances were normalized to the start (0) and end (1) of pachytene based on DAPI morphology

(see Methods). For analysis, the germline was divided into 7 bins encompassing the transition

zone (Bin 1), early pachytene (Bins 2-3), mid pachytene (Bins 4-5), and late pachytene (Bins 6-

7). The germline positions at which each bin start and end are marked on the X axis as vertical

grey lines. Heat maps below each bin display the p values of pairwise comparisons of RAD-51

foci per nucleus counts within that bin. P values were calculated using Mann-Whitney U tests

with Holm-Bonferroni correction for multiple comparisons. Blue tiles indicate significant

differences (adjusted p value <0.05) and grey tiles indicate nonsignificant effects (adjusted p

value >0.05).

Figure 6

Figure 6. Germline *uev-2* overexpression differentially impacts DSB levels in young and aged germlines. A) Schemes used to isolate young (1 day post-L4) and aged (4 days post L4) worms for experiments. Days count ~18-24 hour periods after hermaphrodites were isolated as

L4 larvae and are separated by alternating grey shaded boxes. B) Representative images of

RAD-51 foci in meiotic nuclei of young *pie-1p::uev-2*, aged *pie-1::uev-2*, and young wildtype

germlines. Scale bars represent 5µm. C) RAD-51 foci per nucleus in oocytes. Line plots

represent the mean RAD-51 foci per nucleus along the length of the germline in a sliding

window encompassing 0.1 units of normalized germline distance with a step size of 0.01

germline distance units. Mean RAD-51 foci were calculated from nuclei analyzed in n=9 total

germlines derived from ≥3 experimental replicates within each age group. Shaded areas around

each line represent ± SEM. Total nuclei analyzed (Bins 1/2/3/4/5/6/7) Young Wildtype:

185/117/146/107/117/97/83; Aged Wildtype Mated: 234/205/192/173/154/177/96; Young *pie-*

1p::uev-2: 182/192/162/125/116/97; Aged *pie-1p::uev-2* Mated: 140/149/136/127/126/95/102.

Germlines distances were normalized to the start (0) and end (1) of pachytene based on DAPI

morphology (see Methods). For analysis, the germline was divided into 7 bins encompassing

the transition zone (Bin 1), early pachytene (Bins 2-3), mid pachytene (Bins 4-5), and late

pachytene (Bins 6-7). The germline positions at which each bin start and end are marked on the

X axis as vertical grey lines. Heat maps below each bin display the p values of pairwise

comparisons of RAD-51 foci per nucleus counts within that bin. P values were calculated using

Mann-Whitney U tests with Holm-Bonferroni correction for multiple comparisons. Blue tiles

indicate significant differences (adjusted p value <0.05) and grey tiles indicate nonsignificant

effects (adjusted p value >0.05).

Figure 7

Figure 7. Model of aging and sperm effects on DSB levels during germline aging.

References

Achache H., R. Falk, N. Lerner, T. Beatus, and Y. B. Tzur, 2021 Oocyte aging is controlled by

mitogen-activated protein kinase signaling. Aging Cell 20.

Albert Hubbard E. J., and D. Greenstein, 2000 The Caenorhabditis elegans gonad: A test tube

for cell and developmental biology. Dev. Dyn. 218: 2–22.

Andux S., and R. E. Ellis, 2008 Apoptosis maintains oocyte quality in aging Caenorhabditis

elegans females. PLoS Genet. 4.

Angeles-Albores D., D. H. W. Leighton, T. Tsou, T. H. Khaw, I. Antoshechkin, *et al.*, 2017 The

Caenorhabditis elegans female-like state: Decoupling the transcriptomic effects of aging

and sperm status. G3 Genes, Genomes, Genet. 7: 2969–2977.

Aprison E. Z., S. Dzitoyeva, and I. Ruvinsky, 2022 Serotonin coordinates reproductive functions

in Caenorhabditis. BioRXiv.

Broekmans F. J., E. A. H. Knauff, E. R. te Velde, N. S. Macklon, and B. C. Fauser, 2007 Female

reproductive ageing: current knowledge and future trends. Trends Endocrinol. Metab. 18:

58–65.

Cahoon C. K., J. M. Helm, and D. E. Libuda, 2019 Synaptonemal Complex Central Region

Proteins Promote Localization of Pro-crossover Factors to Recombination Events During

Caenorhabditis elegans Meiosis. Genetics 213: 395–409.

Chatzidaki E. E., S. Powell, B. J. H. Dequeker, J. Gassler, M. C. C. Silva, *et al.*, 2021 Ovulation

suppression protects against chromosomal abnormalities in mouse eggs at advanced

maternal age. Curr. Biol. 31: 4038-4051.e7.

Cinquin A., M. Chiang, A. Paz, S. Hallman, O. Yuan, *et al.*, 2016 Intermittent Stem Cell Cycling

Balances Self-Renewal and Senescence of the C. elegans Germ Line. PLoS Genet. 12.

Colaiácovo M. P., A. J. MacQueen, E. Martinez-Perez, K. McDonald, A. Adamo, *et al.*, 2003

Synaptonemal complex assembly in C. elegans is dispensable for loading strand-exchange

proteins but critical for proper completion of recombination. Dev. Cell 5: 463–474.

Cruz-García A., A. López-Saavedra, and P. Huertas, 2014 BRCA1 accelerates CtIP-ediated

DNA-end resection. Cell Rep. 9: 451–459.

Dernburg A. F., K. McDonald, G. Moulder, R. Barstead, and M. Dresser, 1998 Meiotic

Recombination in C. elegans Initiates by a Conserved Mechanism and Is Dispensable for

Homologous Chromosome Synapsis. Cell 94: 387–389.

Doniach T., and J. Hodgkin, 1984 A Sex-Determining Gene, fem-1, Required for Both Male and

Hermaphrodite Development in Caenorhabditis elegans. Dev. Biol. 106: 223–235.

Gartner A., and J. A. Engebrecht, 2022 DNA repair, recombination, and damage signaling.

Genetics 220.

Gudgen M., A. Chandrasekaran, T. Frazier, and L. Boyd, 2004 Interactions within the ubiquitin

pathway of Caenorhabditis elegans. Biochem. Biophys. Res. Commun. 325: 479–486.

Kamp J. A., R. van Schendel, I. W. Dilweg, and M. Tijsterman, 2020 BRCA1-associated

structural variations are a consequence of polymerase theta-mediated end-joining. Nat.

Commun. 11: 1–10.

Keeney S., C. N. Giroux, and N. Kleckner, 1997 Meiosis-Specific DNA Double-Strand Breaks

Are Catalyzed by Spo11, a Member of a Widely Conserved Protein Family. Cell 88: 375–

384.

Kern C. C., S. J. Townsend, A. Salzmann, N. B. Rendell, G. W. Taylor, *et al.*, 2021 C. elegans

feed yolk to their young in a form of primitive lactation. Nat. Commun. 12.

L'Hernault S. W., 2006 Spermatogenesis. WormBook 1–14.

Li Q., S. Hariri, and J. Engebrecht, 2020 Meiotic double-strand break processing and crossover

patterning are regulated in a sex-specific manner by BRCA1-BARD1 in C. elegans.

Genetics 216: 359–379.

Libuda D. E., S. Uzawa, B. J. Meyer, and A. M. Villeneuve, 2013 Meiotic chromosome

structures constrain and respond to designation of crossover sites. Nature 502: 703–706.

Luo S., W. M. Shaw, J. Ashraf, and C. T. Murphy, 2009 TGF-ß Sma/Mab signaling mutations

uncouple reproductive aging from somatic aging. PLoS Genet. 5.

Luo S., G. A. Kleemann, J. M. Ashraf, W. M. Shaw, and C. T. Murphy, 2010 TGF-β and Insulin

Signaling Regulate Reproductive Aging via Oocyte and Germline Quality Maintenance. Cell

143: 299–312.

Mack H. I. D., T. Heimbucher, and C. T. Murphy, 2018 The nematode Caenorhabditis elegans

as a model for aging research. Drug Discov. Today Dis. Model. 27: 3–13.

Maures T. J., L. N. Booth, B. A. Benayoun, Y. Izrayelit, F. C. Schroeder, *et al.*, 2014 Males

shorten the life span of C. elegans hermaphrodites via secreted compounds. Science (80-.

). 343: 541–544.

Mccarter J., B. Bartlett, T. Dang, and T. Schedl, 1999 On the Control of Oocyte Meiotic

Maturation and Ovulation in Caenorhabditis elegans. Dev. Biol. 205: 111–128.

Moghadam A. R. E., M. T. Moghadam, M. Hemadi, and G. Saki, 2022 Oocyte quality and aging.

J. Bras. Reprod. Assist. 26: 105–122.

Pickett C. L., and K. Kornfeld, 2013 Age-related degeneration of the egg-laying system

promotes matricidal hatching in Caenorhabditis elegans. Aging Cell 12: 544–553.

Preibisch S., S. Saalfeld, and P. Tomancak, 2009 Globally optimal stitching of tiled 3D

microscopic image acquisitions. Bioinforma. Appl. 25: 1463–1465.

Raices M., R. Bowman, S. Smolikove, and J. L. Yanowitz, 2021 Aging Negatively Impacts DNA

Repair and Bivalent Formation in the C. elegans Germ Line. Front. Cell Dev. Biol. 9.

Rosu S., K. A. Zawadzki, E. L. Stamper, D. E. Libuda, and A. L. Reese, 2013 The C. elegans

DSB-2 Protein Reveals a Regulatory Network that Controls Competence for Meiotic DSB

Formation and Promotes Crossover Assurance. PLoS Genet 9: 1–23.

Ruth K. S., F. R. Day, J. Hussain, A. Martínez-Marchal, C. E. Aiken, *et al.*, 2021 Genetic

insights into biological mechanisms governing human ovarian ageing. Nature 596: 393–

397.

Sancho E., M. R. Vila, L. Sanchez-Pulido, J. J. Lozano, R. Paciucci, *et al.*, 1998 Role of UEV-1,

an Inactive Variant of the E2 Ubiquitin-Conjugating Enzymes, in In Vitro Differentiation and

Cell Cycle Behavior of HT-29-M6 Intestinal Mucosecretory Cells. Mol. Cell. Biol. 18: 576–

589.

Schedl T., and J. Kimble, 1988 fog-2, a Germ-Line-Specific Sex Determination Gene Required

for Hermaphrodite Spermatogenesis in Caenorhabditis elegans. Genetics 119: 46–61.

Shi C., and C. T. Murphy, 2014 Mating Induces Shrinking and Death in *Caenorhabditis* Mothers.

Science (80-.). 343: 536–540.

Shim E. Y., A. K. Walker, and T. Keith Blackwell, 2002 Broad requirement for the mediator

subunit RGR-1 for transcription in the Caenorhabditis elegans embryo. J. Biol. Chem. 277:

30413–30416.

Silva-García C. G., A. Lanjuin, C. Heintz, S. Dutta, N. M. Clark, *et al.*, 2019 Single-copy knock-in

loci for defined gene expression in caenorhabditis elegans. G3 Genes, Genomes, Genet.

9: 2195–2198.

Tang L., T. Machacek, Y. M. Mamnun, A. Penkner, J. Gloggnitzer, *et al.*, 2010 Mutations in

Caenorhabditis elegans him-19 Show Meiotic Defects That Worsen with Age. Mol. Biol.

857  Cell 21: 885–896.

858 Thompson O., M. Edgley, P. Strasbourger, S. Flibotte, B. Ewing, *et al.*, 2013 The million

859  mutation project: A new approach to genetics in Caenorhabditis elegans. Genome Res. 23:

860  1749–1762.

861 Toraason E., V. L. Adler, N. A. Kurhanewicz, A. DiNardo, A. M. Saunders, *et al.*, 2021

862  Automated and customizable quantitative image analysis of whole Caenorhabditis elegans

863  germlines. Genetics 217.

864 Vandemark A. P., R. M. Hofmann, C. Tsui, C. M. Pickart, and C. Wolberger, 2001 Molecular

865  Insights into Polyubiquitin Chain Assembly: Crystal Structure of the Mms2/Ubc13

866  Heterodimer. Cell 105: 711–720.

867 Wasserzug-Pash P., R. Rothman, E. Reich, L. Zecharyahu, O. Schonberger, *et al.*, 2022 Loss

868  of heterochromatin and retrotransposon silencing as determinants in oocyte aging. Aging

869  Cell 21.

870 Wickham H., M. Averick, J. Bryan, W. Chang, L. McGowan, *et al.*, 2019 Welcome to the

871  Tidyverse. J. Open Source Softw. 4: 1686.

872 Wijk S. J. L., and H. T. M. Timmers, 2010 The family of ubiquitin-conjugating enzymes (E2s):

873  deciding between life and death of proteins. FASEB J. 24: 981–993.

874 Yokoo R., K. A. Zawadzki, K. Nabeshima, M. Drake, S. Arur, *et al.*, 2012 COSA-1 Reveals

875  Robust Homeostasis and Separable Licensing and Reinforcement Steps Governing

876  Meiotic Crossovers. Cell 149: 75–87.