The DNA Damage response and cell competition are p53- and Xrp1-dependent processes that suppress hyperplastic aneuploidy

Chaitali Khan[1] and Nicholas E. Baker[2]

Department of Genetics
Albert Einstein College of Medicine
1300 Morris Park Avenue
Bronx
NY 10461
USA

1. Present address: Cell and Developmental Biology Center, National Heart Lung and Blood Institute, National Institutes of Health, Bethesda, MD 20892, USA

2. Author for correspondence: Nicholas.baker@einsteinmed.edu

ABSTRACT

It is important to understand how p53 suppresses tumorigenesis. P53 activity contributes to many instances of cell competition in mammals. This has not been seen for *Drosophila* p53, where the transcription factor Xrp1 is an effector of cell competition. Xrp1 is induced in a p53-dependent manner by DNA damage, and we report that Xrp1 mediates multiple functions of p53 in the DNA damage response, contributing to p53-dependent gene transcription and DNA damage-induced apoptosis. Differences in either Xrp1 or p53 activity, occurring between wild type and mutant cells experiencing mild genotoxic stress, both resulted in cell competition. Unexpectedly, cell competition due to differential p53 activity did not require Xrp1 but instead was restrained by Xrp1. We show that Xrp1 has a p53-independent role in removing genomically-altered cells. Both Xrp1 and p53 limit the accumulation of abnormal cells that results from genotoxicity, and we propose that genomic alterations enhance cellular growth of p53 mutant cells and promote cell competition, potentially contributing to the tumorigenesis in p53 mutants.

INTRODUCTION

Mutations in the tumor suppressor gene p53 occur frequently in many cancer types (Muller and Vousden 2014). This comes as no surprise since p53 plays an essential role in ensuring genomic integrity by serving as the key effector of DNA damage response (DDR) (Levine, Hu, and Feng 2006; Kastenhuber and Lowe 2017). However, the emphasis on p53's role in DDR as its tumor suppression mechanism has recently been questioned. Point mutations that block p53 function in the acute DDR have little effect in blocking tumor growth, and conditional knock-out studies point to a critical window for tumor suppressor function that is subsequent to the resolution of DNA damage (Christophorou et al. 2005; Christophorou et al. 2006; Li et al. 2012; Kaiser and Attardi 2018). Besides responding to DNA damage, p53 has other functions ranging from metabolic regulation, cell-cell recognition and cell competition (Baker 2020; Lacroix et al. 2020; Pilley, Rodriguez, and Vousden 2021). These other functions might also be important for tumor suppression by p53. Here we focus on the role of p53 in cell competition. Cell competition is a phenomenon whereby interaction between cells leads to the elimination of cells with lower fitness, and in mammals higher p53 activity is often associated with lower fitness (Merino et al. 2015; Nagata and Igaki 2018; Baker 2020; Morata 2021; Pilley, Rodriguez, and Vousden 2021). Cell competition was first described in the case of *Drosophila* cells heterozygous for mutations in ribosomal protein genes ($Rp^{+/-}$ cells) (Morata and Ripoll 1975). Cell competition is not limited to $Rp^{+/-}$ mutants but occurs in response to a variety of genetic differences, in mammals as well as in *Drosophila* (Maruyama and Fujita 2017; Baker 2017).

One function of cell competition is to suppress tumorigenesis by eliminating abnormal cells prior to transformation (Vishwakarma and Piddini 2020; Gogna and Moreno 2020). Indeed, cell competition removes *Drosophila* cells mutated for certain genes defining apical-basal polarity of epithelial cells, which otherwise form neoplastic tumors (Brumby and Richardson 2003; Igaki, Pagliarini, and Xu 2006; Kanda and Igaki 2020). In addition, our recent work demonstrated that ribosomal protein (Rp) genes participate in the recognition and removal of certain aneuploid cells through cell competition. Almost all tumors are aneuploid, and genotypes that promote aneuploidy are cancer prone in humans and in mouse models, therefore it is expected that mechanisms that reduce aneuploidy are tumor suppressive (Hanks et al. 2004; Matsuura et al. 2006; Foijer, Draviam, and Sorger 2008; García-Castillo et al. 2008; Baker et al. 2009; Snape et al. 2011;

Mukherjee et al. 2014). Because the *Rp* genes are distributed around the genome, in

Drosophila as in humans, aneuploidy and other large-scale genetic alterations generally

change *Rp* gene dose, and such changes could serve as a sensor for genomic alterations

(Uechi, Tanaka, and Kenmochi 2001; Marygold et al. 2007; McNamee and Brodsky 2009).

Accordingly, clones of cells with segmental chromosome deletions are only tolerated

when they do not affect *Rp* gene dose, but eliminated by cell competition when they

reduce dose of *Rp* genes (Ji et al. 2021).

Aneuploidy is also known to cause a stress response due to the altered expression

of many proteins, not just Rp, leading to cell-autonomous deficits in growth and survival

(Torres et al. 2007; Dekanty et al. 2012; Brennan et al. 2019; Joy et al. 2021). In *Drosophila*,

blocking apoptosis of the aneuploid cells that result from chromosome instability, and

which experience cell-autonomous stress, leads to invasive tumors (Dekanty et al. 2012;

Joy et al. 2021).

Differences in p53 activity between cells are a key feature of many instances of

cell competition in mammals, and by themselves are sufficient to lead to competition

between mammalian cells (Baker, Kiparaki, and Khan 2019; Pilley, Rodriguez, and

Vousden 2021). For example, in mouse hematopoietic stem cell chimeras, p53 activation

by low dose ionizing radiation leads to elimination of cells with higher relative levels of

p53 (Bondar and Medzhitov 2010; Marusyk et al. 2010). Similarly, in mouse embryo

chimeras, aneuploid cells are eliminated by the presence of euploid cells in a p53-

dependent manner (Singla et al. 2020). In cultured mammalian cells, direct

pharmacological activation of p53, in the absence of other stressors, is sufficient to

promote competitive elimination (Wagstaff et al. 2016).

Heterozygous mutation of human Rp genes activates p53 through inhibition of

MDM2 (Donati et al. 2011; Bursac et al. 2014). In *Drosophila*, there is no evidence that p53

is activated by Rp mutations, p53 is not required for the cell competition of $Rp^{+/-}$ cells, and

the MDM2 proteins, whose inhibition in mammalian p53 mutant cells is the basis for p53

activation, are absent from *Drosophila* (Chakraborty et al. 2015; Kale et al. 2015; Lee et al.

2018). There is, however, a role for p53 in protective metabolic changes that permit

Drosophila cells expressing elevated *myc* to eliminate nearby wild type cells. This is an

example of super-competition where it is the wild-type cells that are eliminated (de La

Cova et al. 2014).

Instead of p53, the transcription factor that is important in the competition of $Rp^{+/-}$ cells is the bZip domain protein Xrp1. Xrp1 protein is barely expressed in wild type imaginal discs, but induced by Rp mutations (Lee et al. 2016; Baillon et al. 2018; Lee et al. 2018). Xrp1 expression is induced in $Rp^{+/-}$ cells not by p53 but through an activity of the RpS12 protein; this activity seems to be enhanced in $Rp^{+/-}$ cells (Kale et al. 2018; Lee et al. 2018; Ji et al. 2019). Aneuploid cells that arise as a consequence of irradiation, or telomere loss, are eliminated from *Drosophila* tissues in a p53-independent manner (Wichmann, Jaklevic, and Su 2006; McNamee and Brodsky 2009; Titen and Golic 2008). The majority of such aneuploid cell elimination depends on RpS12, suggesting that it represents a cell competition process similar to that affecting $Rp^{+/-}$ cells (Ji et al. 2021). Xrp1 binds to DNA as a heterodimer with another bZip protein, Irbp18, as originally discovered though binding of Xrp1 and Irbp18 to terminal repeats of P element transposons (Francis et al. 2016). The Xrp1/Irbp18 heterodimer acts as a sequence-specific transcription factor (Kiparaki et al. 2022). In $Rp^{+/-}$ mutant cells, *Xrp1* and *irbp18* transcriptionally autoregulate, which is needed for their expression and is essential for cell competition (Baillon et al. 2018; Blanco, Cooper, and Baker 2020). Xrp1 is a rapidly-evolving gene without any clear mammalian homolog, although Irbp18 is the *Drosophila* homolog of C/EBP proteins (Blanco, Cooper, and Baker 2020). Xrp1 may be functionally related to DDIT3/CHOP, a mammalian bZip protein with roles in the ER stress response, because DDIT3/CHOP heterodimerizes with C/EBP, like Xrp1 does with Irbp18, because Xrp1 expression is also induced by ER stress, and because overexpression of human DDIT3/CHOP in *Drosophila* produces a phenotype resembling that of Xrp1 expression (Blanco, Cooper, and Baker 2020; Brown et al. 2021; Ochi et al. 2021).

Xrp1 was initially found as one of the most abundantly expressed transcriptional targets of p53 following irradiation (Brodsky et al. 2004), and Xrp1 has also been proposed to contribute genome integrity, because irradiation of Xrp1 mutant flies potentiates loss of heterozygosity (Akdemir et al. 2007). Xrp1 may also contribute to DNA recombination, since it promotes transposition of the *Drosophila* P elements, to which it also binds directly (Francis et al. 2016).

We hypothesized that Xrp1 is an effector of p53 function in the *Drosophila* DDR, and that by activating Xrp1 expression directly, ie by-passing p53, *Rp* mutant cells activate a portion of what might otherwise be considered p53 responses, and that this includes cell competition (Baker, Kiparaki, and Khan 2019). If this was correct, then Xrp1

should be required for aspects of the p53-dependent DDR, and p53 activity differences

should be able, through Xrp1, to trigger competition between cells, by-passing the

requirement for an *Rp* mutation. Here, we test these hypotheses. We confirm that Xrp1 is

upregulated by p53 and show that Xrp1 is needed for expression of multiple p53 target

genes, contributing to p53-dependent apoptosis in response to irradiation. We show that

differential activation of p53 does indeed cause cell competition in *Drosophila*, as it does

in mammals. Unexpectedly, however, the mechanism of p53-driven cell competition was

not Xrp1-dependent. On the contrary, Xrp1 suppressed the expansion of *p53$^{-/-}$* mutant cell

clones experiencing genotoxic stress. Our work then revealed roles of both Xrp1 and p53

in limiting the accumulation of damaged and aneuploid cells. This suggests that it is cells

with genomic alterations resulting from DNA damage that are behind the super-

competitive behavior of *p53* mutant cells, which is enhanced in the absence of Xrp1

because of the p53-independent role of Xrp1 in eliminating aneuploid cells. Our work

indicates that the connection between p53 and cell competition seen in mammals is

shared in *Drosophila*. Additionally, it highlights the importance of cell competition as a

mechanism containing genomic instability by eliminating aneuploid cells, and suggests

that under some conditions, aneuploid cells that evade cell competition themselves

become super-competitors.

RESULTS

DNA damage induced expression of Xrp1

Xrp1 mRNA is upregulated by p53 within 30 mins of irradiation of *Drosophila*

embryos (Brodsky et al. 2004; Akdemir et al. 2007), perhaps by direct binding of the p53

within Xrp1 locus (Kurtz et al. 2019). To confirm that Xrp1 is a p53 target in imaginal

discs, we explored Xrp1 expression in third instar larval wing discs using a fly line

expressing Xrp1 protein with a C-terminal HA tag, encoded at the endogenous locus

(Blanco, Cooper, and Baker 2020). Wild type wing discs exhibited few Xrp1-HA positive

cells; these might reflect spontaneous DNA damage or another cellular stress (Figure 1A).

Exposure to 1500 Rads of γ-rays (IR) induced strong Xrp1-HA positive cells by 2h after

IR, which further increased by 4h after IR (Figure 1B-C). DNA damage induced Xrp1-HA

expression was non-uniform, with some cells showing higher expression levels than

others, which was different from the more uniform Xrp1-HA expression seen in *Rp$^{+/-}$*

mutant wing discs (Figure 1D). A similar, salt-and-pepper pattern of Xrp1 was observed
with Xrp1 antibody (Figure 1E and F). The specificity of Xrp1-HA expression was further
confirmed by its reduction after RNAi for Xrp1(Figure 1G).

Next, we tested the requirement of p53, using a dominant-negative form of p53
encoded by UAS-$p53^{H159N}$. This was expressed in posterior wing compartments, so that
Xrp1-HA expression levels could be compared to anterior compartments of the same
wing discs that lacked dominant-negative p53 expression. Dominant-negative p53
significantly reduced IR-induced Xrp1-HA (Figure 1H), as previously reported for Xrp1
transcription in irradiated embryos (Brodsky et al. 2004; Akdemir et al. 2007). Imaginal
disc Xrp1 was also p53-dependent at the transcript level, with a ~3x reduction in mRNA
level seen after qRT-PCR from irradiated $p53^{-/-}$ wing discs (Figure 1- figure supplement
1A).

Interestingly, Xrp1-HA protein expression was only slightly affected by Irbp18
knock-down (Figure 1I), arguing that autoregulation contributes little to IR-induced $Xrp1$
transcription. Consistent with this, 50% of the normal $Xrp1$ mRNA level remained in the
$Xrp1^{m2-73}$/Df mutant background, in accordance with continued transcription of the
functionally null $Xrp1^{m2-73}$ point mutant allele (Figure 1- figure supplement 1A). This was
different from Xrp1 expression in $Rp^{+/-}$ mutants, where Xrp1 autoregulation is important
(Baillon et al. 2018; Blanco, Cooper, and Baker 2020). The expression of the Xrp1-LacZ
enhancer trap, a reporter upregulated in $Rp^{+/-}$ mutants (Baillon et al. 2018; Lee et al. 2018),
was also moderately elevated after irradiation (Figure 1J, K). Xrp1-LacZ expression,
however, depended on Xrp1 and Irbp18 function, both after irradiation and in the
unirradiated control (Figure 1- figure supplement 1B-E). Unexpectedly, reduced p53
function slightly elevated the expression of Xrp1-LacZ expression after irradiation,
instead of reducing it (Figure 1L). The discordance between regulation of Xrp1-LacZ
expression with that of Xrp1 mRNA and Xrp1-HA protein suggests that Xrp1-LacZ
differs from endogenous Xrp1 regulation, either at the transcriptional or post-
transcriptional levels.

Regulation of p53-dependent DDR target genes by Xrp1
It is known from mammalian studies that p53 regulates a complex transcriptional
program of hundreds or even thousands of genes (Levine, Hu, and Feng 2006). However,
depending on the study, only 122-346 p53-dependent genes have a p53 consensus motif

in the promoter or upstream region, suggesting that they could be direct p53 targets (Wei et al. 2006; Fischer 2017). It is thought that p53 directly controls context-specific, core transcription programs that in turn regulate further secondary, indirect targets (Andrysik et al. 2017). Since Xrp1 is a transcription factor and a p53 target in *Drosophila*, Xrp1 is a candidate to bring secondary target genes under the indirect control of p53. Accordingly, when we compared p53-dependent genes in the DDR (Brodsky et al. 2004; Akdemir et al. 2007) with Xrp1-dependent genes in $Rp^{+/-}$ wing imaginal discs (Lee et al. 2018) and Xrp1-dependent genes in wing imaginal discs over-expressing Xrp1 (Baillon et al. 2018). We found 21 of 42 Rapidly Induced p53-target genes (RIPD) were Xrp1-regulated in wing discs (Figure 2A). These genes were associated with Gene Ontology (GO) terms transcription factors, cell death, DNA repair, and redox response (Figure 2B).

To test if Xrp1 was needed for regulating the DNA damage induced targets of p53, we employed available reporter lines and also confirmed gene expression by qRT-PCR. *GstD-lacZ* is a redox response reporter that is induced in response to oxidative stress (Sykiotis and Bohmann 2008), and is elevated in $Rp^{+/-}$ wing discs downstream to Xrp1 (Ji et al. 2021; Kiparaki et al. 2022). *GstD-lacZ* was expressed in response to IR, dependent on p53 (Figure2C-D). We found that Xrp1 and Irbp18 were needed for p53-dependent expression of *GstD-lacZ* after IR (Figure 2F-G). This was also in accordance with Xrp1 binding sites being present in the *GstD*-lacZ reporter (Brown et al. 2021). To confirm that antioxidant genes themselves were affected like the *GstD-lacZ* reporter was, we measured *GstD1* mRNA levels in wing discs. Induction of *GstD1* mRNA by IR indeed depended on both p53 and Xrp1 (Figure 2M).

Another stress pathway activated in DDR is the Jun Kinase pathway (Picco and Pages 2013). JunK activity is also Xrp1-dependent in $Rp^{+/-}$ wing discs (Lee et al. 2018; Ji et al. 2019). We found the JNK reporter *TRE-dsRed* was induced by IR in a p53-dependent manner (Figure 2H-J), and was also Xrp1/Irbp18 dependent (Figure 2K & L). The TNF-alpha ligand *eiger (egr)* has been proposed to activate JNK signaling in the DDR (Sanchez et al. 2019). Surprisingly, however, *egr* transcription was p53-dependent but independent of Xrp1, which questions whether it is solely responsible for JNK signaling in the DDR (Fig. 2N).

All except one of the p53-dependent DNA repair genes induced in DDR were also potential Xrp1 targets (Figure 2B). We looked at wing disc mRNA levels of representative genes from the major DNA repair pathways. These were the homologous recombination

pathway gene *rad50*, the translesion polymerase DNA polymerase zeta subunit 1, *mus205*,

and Non-Homologous End Joining (NHEJ) pathway genes DNA ligase 4 (*lig4*) and *Ku80*.

We found that IR-dependent expression of all these genes depended on Xrp1 at least as

strongly as upon p53 (Figure 2O-R). A well-known aspect of DDR is apoptotic death of

severely damaged cells, dependent in *Drosophila* on p53-dependent induction of the pro-

apoptotic genes *rpr* and *hid* (Brodsky et al. 2004; Dekanty, Barrio, and Milán 2015). To

explore the potential role of Xrp1, we examined reporter lines for *rpr* and *hid*, which are

expressed in response to IR and are widely used as p53 reporters (Brodsky et al. 2000;

Tanaka-Matakatsu et al. 2009; Fan et al. 2010). Two reporter lines for *hid* are the *hid5′F-*

wt-EGFP reporter, which contains ~2 kb spanning the *hid* promoter from +16 to −2210bp

relative to the transcription start site, and the *hid$^{20-10}$-lacZ* reporter, which contains a

distinct 10 kb sequence from the *hid* upstream region (Tanaka-Matakatsu et al. 2009; Fan

et al. 2010) (Figure 3A). As expected, *hid5′F-wt-EGFP* was highly induced after IR (Figure

3B-C), and was p53-dependent (Figure 3D). Both *Xrp1* and *Irbp18* RNAi also reduced the

hid5′F-wt-EGFP expression to the same extent as p53 dominant-negative did (Figure 3E-

F). The *hid$^{20-10}$-lacZ* reporter was also induced by IR and was dependent upon p53 (Figure

3G-I). Both *Xrp1* and *Irbp18* RNAi reduced IR-dependent *hid$^{20-10}$-lacZ* expression, although

not so much as p53 dominant-negative (Figure 3J-K). We assessed *rpr* expression using

the *rpr^{150}-lacZ* reporter, containing a 150bp sequence including a putative p53 binding site

~4.8kb upstream of the *rpr* gene (Brodsky et al. 2000). In controls, *rpr^{150}-lacZ* reporter

exhibits uniform expression in most of the wing disc, on which is superimposed a region

of elevated expression bordering the posterior wing pouch and notum region (Alpar,

Bergantiños, and Johnston 2018; Li et al. 2020) and (Figure 3L). Because IR enhanced all

aspects of *rpr^{150}-lacZ* expression (Figure 3L,M and Figure 3- Supplemental 1A), we

measured the average *rpr^{150}-lacZ* expression level in anterior and posterior compartments

and computed the ratio of posterior-to-anterior expression level as a readout of genetic

perturbations applied to the posterior compartments (Figure 3- Supplemental 1B). The

increase in *rpr^{150}-lacZ* following IR was p53-dependent, as expected, but unaffected by

RNAi for either Xrp1 or Irbp18 (Figure 3N-P- Supplemental 1B).

To confirm that these studies of *hid* and *rpr* reporters were indeed reflective of the

expression of the genes themselves, mRNA levels were measured in wing discs after IR.

Consistent with the reporter studies, induction of *hid* mRNA by irradiation was

completely dependent on p53, and partially dependent on Xrp1 (Figure 3Q). Induction of

rpr mRNA expression was completely dependent on p53 but was unaffected by the loss
of Xrp1(Figure 3R).

281   Taken together, these results strongly support the notion that p53 target genes
identified through the *Drosophila* DDR include secondary targets that are more directly
induced by the p53-dependent expression of Xrp1. Only two p53 target genes, *egr* and
rpr, were induced independently of Xrp1 out of the nine examined, although *hid*
induction has both Xrp1-dependent and -independent components. Analysis of p53 and
Xrp1 binding motifs supported this conclusion. Sequences matching the p53 binding
consensus has been located within both the *rpr^{150}-lacZ* reporter and the *hid5'F-wt-EGFP*
reporter (Brodsky et al. 2000; Wichmann, Uyetake, and Su 2010). Consensus binding sites
for the Xrp1/Irbp18 heterodimer have been determined by bacterial one-hybrid studies
(Fly Factor Survey) (Zhu et al. 2010) and from ChIP-seq after Xrp1 over-expression in
larval wing disc (Baillon et al. 2018). Such sequences are footprinted by Xrp1/Irbp18 in
the P element terminal repeat, (Francis et al. 2016), and sufficient to confer Xrp1/Irbp18-
dependent transcription on luciferase reporters (Kiparaki et al. 2022). Using an *in-silico*
motif finder (INSECT) (Rohr et al. 2013), we found three Xrp1/Irbp18 binding motifs in
the *hid5'F-wt-EGFP* reporter sequence and two in the *hid$^{20-10}$-lacZ* reporter sequence
(Figure 3T), but we found none in the *rpr^{150}-lacZ* reporter that was expressed
independently of Xrp1 and Irbp18 (Figure 3T). We conclude that, while *rpr* could be a
direct target of p53 in the DDR, *hid* expression is partially dependent on Xrp1 induction
downstream of p53, and the *hid5'F-wt-EGFP* reporter entirely so.

Xrp1 is sufficient for the expression of p53- dependent DDR target genes
302   It is possible that Xrp1 might activate p53 to create positive feedback in the DDR.
We used the *rpr^{150}-lacZ* reporter, which is activated by p53 but not Xrp1, to test how p53
activity was affected by Xrp1 over-expression. As expected, Xrp1 overexpression in the
wing pouch resulted in huge amount of cell death (Figure 4- figure supplement 1A-B),
but no discernable increase in *rpr^{150}-lacZ* levels (Figure 4- figure supplement 1C-D).
Similar results were obtained by expressing Xrp1 in the differentiating eye discs (Fig. 4A-
B). Thus, Xrp1 does not seem to cause p53 activation, but to act downstream of p53.

309   Next, we tested if Xrp1 was sufficient for the expression of target genes. We found
that Xrp1 overexpression in the eye disc led to a robust upregulation of *hid$^{20-10}$-lacZ*
(Figure 4 C and D) and *hid5'F-wt-EGFP* (Figure 4E and F). Xrp1 also regulates JNK and

oxidative stress responses after DNA damage, and Xrp1 overexpression was sufficient to
activate reporters for these cellular pathways. Levels of both *JNK-dsRED* and *GstD-lacZ*
were highly upregulated by Xrp1 (Figure 4- figure supplement 1E and F & G and H).
These results support the idea that Xrp1 functions downstream of p53, and is sufficient
for the expression of a subset of p53- dependent genes.
317

Xrp1 contributes to DNA damage induced cell death

To see whether the regulation of cell death genes by Xrp1 was functionally
significant, the *p53*-dependent apoptosis that occurs in response to irradiation was
examined. Cell death begins within 30 minutes after IR treatment and is completely
dependent on p53 until 4-5 hours later (Wichmann, Jaklevic, and Su 2006; McNamee and
Brodsky 2009). Accordingly, p53 dominant-negative expression in posterior
compartments substantially reduced the IR-induced cell death, leaving of only a few cells
positive for the activated Dcp-1 caspase (Figure 5B & D). Xrp1 RNAi (two independent
lines) and Irbp18 RNAi both reduced cell death in the posterior compartment (Figure 5C
& D), to a lesser extent than p53 dominant-negative did.

We also examined the cell-autonomy of p53 and Xrp1 function using clones of
mutant cells. Wild type and *p53^{+/-}* heterozygous cells exhibited similar cell death levels
after irradiation, but cell death was absent from *p53^{-/-}* null clones (Figure 5F,H). There
was little cell death observed in *Xrp1^{+/-}* (heterozygous) and *Xrp1^{-/-}* (null) tissue as opposed
to high amount of cell death observed in *Xrp1^{+/+}* (wild type twin spot) (Figure 5G,H)
(effects of Xrp1 mutation on *Rp* mutant cells are also dominant through
haploinsufficiency) (Lee et al. 2018). Control clones had no effect on IR-induced cell death
(Figure 5E,H).

Taken together, these results show that Xrp1 contributes to p53-mediated cell
death in response to irradiation, although this is partial because p53 also regulates cell
death genes directly such as *rpr* (Figure 3R). We also tested the contribution of Xrp1 to
effects of p53 over-expression. Over-expression of p53 in the developing eye causes a
rough eye phenotype in adults (Figure 5- Supplemental figure 1A (Ollmann et al. 2000;
Jin et al. 2000). Xrp1 knock-down very slightly reduced the rough eye phenotype, unlike
dominant-negative p53, which rescued completely (Figure 5- figure supplement 1B,C).
p53 over-expression in wing discs resulted in massive cell death that was reduced only
weekly by Xrp1 knock-down or mutant genotypes (Figure 5- figure supplement 1D -H).

This partial dependence on Xrp1 may reflect the observation that p53 over-expression led to rather little Xrp1 protein expression (Figure 5 figure supplement 1I), which is a difference from endogenous p53 activity in the DDR, suggesting that IR induced (Figure 1H) and over-expressed p53 mechanisms may not be identical.

Differential activation of p53 triggers cell competition in *Drosophila*

Differences in p53 activity between mammalian cells often result in cell competition (Bondar and Medzhitov 2010; Wagstaff et al. 2016; Zhang et al. 2017). In *Drosophila*, p53 is dispensable for the elimination of $Rp^{+/-}$ cells (Kale et al. 2015; Lee et al. 2018), and cells with and without the p53 gene do not compete with one another (de La Cova et al. 2014). If Xrp1 is a downstream effector of *Drosophila* p53, however, and differences in Xrp1 activity lead to cell competition, then we hypothesized that differences in p53 activity between cells might lead to competition, in *Drosophila* as in mammals.

To test this hypothesis, we developed a strategy for chronic mild activation of p53 by feeding the genotoxic drug Cisplatin (Cis-Pt). Cis-Pt causes p53 activation through double-strand breaks and other DNA damage (Dasari and Tchounwou 2014). Since feeding with 0.5-1mM Cis-Pt produce high levels of DNA damage and cell death even in non-mosaics (Figure 6 Supplement 1A) (Khan et al. 2017), we modified this protocol by reducing Cis-Pt dosage. Third instar larvae fed with 0.01 mM or 0.02 mM Cis-Pt (starting at 48h after egg laying) exhibited negligible cell death or DNA damage, as measured by anti-active Dcp1 or anti γH2Av, and only a few Dcp-1 positive cells were seen after feeding 0.05mM Cis-Pt (Figure 6 figure supplement 1A). Anti-γH2Av detects histone H2Av phosphorylated at ser-237 and is a marker for double-strand breaks (Madigan, Chotkowski, and Glaser 2002). We used 0.05 mM Cis-Pt feeding for subsequent experiments, although some results were also corroborated using 0.02 mM Cis-Pt. Neither concentration caused adult defects such as notched wing margins that are seen after more severe DNA damage (Wells and Johnston 2012), although there was a pupariation delay of ~12h or ~ 24h associated with 0.02 mM and 0.05 mM Cis-Pt respectively, and 0.05mM Cis-Pt led to some arrest of development at the pharate pupa stage. We then used FRT mediated somatic recombination in *p53* heterozygotes to generate reciprocal clones of *p53^-/-^* and *p53^+/+^* cells, with Cis-Pt fed from the time of clone

induction, leading to clones of cells with and without p53 activity that had been exposed

to mild genotoxic stress (Figure. 6A & see methods).

The p53 reporter *rpr^{150}*-lacZ was elevated by 0.05 mM Cis-Pt diet (Figure. 6B, C).

In the absence of Cis-Pt, the ratio of *p53$^{-/-}$* and *p53$^{+/+}$* clone sizes was similar to controls, as

reported previously (Figure. 6D, E, H) (de La Cova et al. 2014). By contrast, in the presence

of Cis-Pt the ratio of *p53$^{-/-}$* clone size to sister *p53$^{+/+}$*clone sizes was ~4 by 72 hours after

clone induction and ~ 6 at 96h after clonal induction, although Cis-Pt had little effect on

control clone sizes (Figure. 6F-H). Similar results were seen using 0.02mM Cis-Pt (Figure

6- figure supplement 1B-F). We confirmed these results in eye development, where

0.02mM Cis-Pt enhanced the contribution of *p53$^{-/-}$* mutant clones to adult eyes (Figure 6

figure supplement 1G-J). We also compared absolute *p53$^{-/-}$* clone sizes in the presence and

absence of Cis-Pt. This analysis revealed that *p53$^{-/-}$* mutant clones grew ~ 2.5-fold bigger

in the presence of Cis-Pt (Figure 6I), whereas the reciprocal *p53$^{+/+}$* sister clones grew ~2

fold smaller (Figure 6J).

These results were consistent with competition between clones of *p53$^{+/+}$* cells

experiencing p53 activity and clones of *p53$^{-/-}$* cells, but they also could be explained by a

cell-autonomous p53-dependent growth retardation in the presence of Cis-Pt. To

distinguish these possibilities, we performed a 'neutral-clone assay' that compared the

growth of *p53$^{+/+}$* clones adjacent to *p53$^{-/-}$* sister clones with the growth of independent,

spatially uncorrelated clones within the same wing discs, derived from simultaneous

mitotic recombination of a distinct chromosome. In the absence of Cis-Pt, the two

independent categories of clone, one the sisters of *p53$^{-/-}$* clones and one the sisters of *p53$^{+/+}$*

clones, were both indistinguishable in size. By contrast, in the presence of 0.05 mM Cis-

Pt, the clones with *p53$^{-/-}$* sisters were ~4-fold smaller than the clones with *p53$^{+/+}$* sisters

(Figure 6K-M). These results show that the growth disadvantage of clones of wild type

cells with *p53$^{-/-}$* sisters after Cis-Pt exposure is not a cell-autonomous effect of p53 activity,

but is a property only of wild type clones with nearby *p53$^{-/-}$* sister clones. This indicates

that differences in p53 activity lead to cell competition in *Drosophila* imaginal discs, as

they do in many mammalian tissues, and that *p53$^{-/-}$* cells out-compete nearby cells in

which *p53* is active, even when the p53 activity is mild compared to that usually studied

as representative of the DDR.

We asked if the growth disadvantage *p53$^{+/+}$* clones with *p53$^{-/-}$* sister clones was due

to more robust DNA damage signaling than the *p53$^{-/-}$* mutant cells. We saw no difference

in γH2Av staining between *p53^{-/-}* mutant clones and their *p53^{+/+}* sisters in the absence of

Cis-Pt (Figure 6- figure supplement 2A-B). There was no significant change after Cis-Pt

feeding, indicating that Cis-Pt causes only low levels of DNA damage that do not

accumulate substantially in *p53^{-/-}* mutant cells (Figure 6- figure supplement 2E-H). By

contrast, irradiation with 1500 Rads of γ-rays gave rise to significantly higher γH2Av

levels, that were still further elevated in *p53^{-/-}* mutant clones compared to their *p53^{+/+}* sister

clones (Figure 6- figure supplement 2D-C). This accumulation of *p53^{-/-}* mutant cells with

DNA damage might be due to lack of apoptosis of such cells in the wing disc any such

effect of 0.05mM Cis-Pt must be significantly smaller.

p53-mediated cell competition independent of Xrp1

Since Xrp1 is downstream of p53 in the DDR, we hypothesized that it could be

higher Xrp1 that results in cell competition of wild type cells growing next to *p53^{-/-}* mutant

cells. To test this, we first examined the effect of Cis-Pt on Xrp1 mutant cells. Although

Xrp1^{-/-} mutant clones grew like their *Xrp1^{+/+}* sister clones in the absence of Cis-Pt, as

reported previously Figure 7(A, E) (Baillon et al. 2018; Lee et al. 2018), Cis-Pt exposure

led to larger *Xrp1^{-/-}* mutant clones and smaller *Xrp1^{+/+}* twin-spots (Figure 7B, D, E). We

also observed that *irbp18^{-/-}* mutant clones grew larger than their *irbp18^{+/+}* sister clones in

the presence of Cis-Pt, suggesting that the Xrp1/Irbp18 heterodimer was activated by

Cis-Pt (Figure 7G- I). Thus, Cis-Pt has similar effects on clones of *Xrp1* and *p53* mutant

cells, consistent with both genes functioning in the same pathway.

To test whether Xrp1 was the downstream effector of p53 in cell competition, we

explored whether different levels of p53 activity could lead to cell competition in the

absence of Xrp1. We depleted posterior compartments of Xrp1 by RNAi and examined

competition between *p53^{-/-}* mutant clones and their *p53^{+/+}* twin-spots. In the absence of

Cis-Pt, the ratio of *p53^{-/-}* and *p53^{+/+}* clone sizes was comparable in the anterior and the

posterior compartments, indicating that it was not affected by Xrp1 (Figure 8A-C). If Xrp1

was required for competition between cells with different p53 activities, we expected that

p53^{-/-} and *p53^{+/+}* clone sizes would remain similar in Xrp1-depleted posterior

compartments in the presence of Cis-Pt. Contrary to this expectation, *p53^{-/-}* clones were

larger than *p53^{+/+}* twin spots when Xrp1 was depleted in the presence of Cis-Pt, just as

they had been without Xrp1 depletion (Figure 8D-F). To analyze this in more detail, we

compared the clone sizes of all the genotypes. We found that, in the absence of Cis-Pt, growth of all the $p53^{-/-}$ and $p53^{+/+}$ clones was equivalent, regardless of *Xrp1* knockdown, confirming earlier results (Figure 6E,H). In the presence of Cis-Pt, all the $p53^{-/-}$ clones grew larger and all the $p53^{+/+}$ clones grew less, also confirming earlier experiments (Figure 6G,H). Neither of these Cis-Pt effects was reversed by Xrp1 knockdown (Figure 8D-F). In fact, in the presence of Cis-Pt, the $p53^{-/-}$ clones depleted for Xrp1 grew significantly larger than plain $p53^{-/-}$ clones in the presence of Cis-Pt (Figure 8G). In addition, the adjacent $p53^{+/+}$ sister clones depleted for Xrp1 grew less than $p53^{+/+}$ sister clones able to express Xrp1, although this difference was not statistically significant after correction for multiple testing (Figure 8H).

These results were not consistent with our original hypothesis that Xrp1 mediates the competitive elimination of cells with higher p53 activity, because the p53-dependent growth reduction of wild type clones under genotoxic stress and near to $p53^{-/-}$ clones occurred independently of Xrp1. In addition, $p53^{-/-}$ clones exposed to Cis-Pt grew even more when Xrp1 was depleted, indicating an unexpected, p53-independent function of Xrp1 under genotoxic stress.

Xrp1 function in mediating competitive cell death to limit genomic instability

If Xrp1 was activated through p53, we would not expect Xrp1 expression or function in $p53^{-/-}$ cells exposed to Cis-Pt. Instead, we observed a salt-and pepper Xrp1 expression pattern after Cis-Pt exposure that included $p53^{-/-}$, $p53^{+/-}$, and $p53^{+/+}$ cells (Figure 8- figure supplement 1A,B). This expression was much lower than seen after 1500 Rads of IR. By contrast, Cis-Pt-treated wing discs containing *Xrp1* clones showed Xrp1 expression only in $Xrp1^{+/+}$ cells, not $Xrp1^{+/-}$ cells or $Xrp1^{-/-}$ cells (Figure 8- figure supplement 1C). These results were unexpected, because previously we had found that p53 was required for DNA damage to induce Xrp1 expression (Figure 1H and Figure 1-figure supplement 1A). Using Xrp1-HA to monitor Xrp1 expression, and enGal4 to express dominant-negative p53 in posterior compartments, we confirmed the salt-and-pepper pattern of Xrp1 expression after Cis-Pt treatment, and, surprisingly, found Xrp1-HA expression was increased where p53 was inhibited, not reduced (Figure 8 supplement 1D-G). We also noticed that Xrp1 expression typically occurred in small cell clusters, possibly clonal.

Notably, we have previously described p53-independent induction of Xrp1 in wing discs in response to IR (Ji et al. 2021). Unlike p53-dependent Xrp1 expression in the acute DDR, which occurs within 2-4h, p53-independent induction peaks 16-24h after irradiation. This Xrp1 expression is thought to occur in aneuploid and segmental-aneuploid cells that arise following DNA damage (McNamee and Brodsky 2009; Ji et al. 2021). More than half of this expression depends on RpS12, which is also required for Xrp1 expression in $Rp^{+/-}$ cells (Lee et al. 2018; Ji et al. 2019; 2021). Here, we verified that Xrp1-HA was expressed in scattered cells 24h following irradiation (Figure 8 supplement 1H,I). Xrp1-HA expression was completely eliminated by knock-down of Xrp1, but not by knock-down of Irbp18 (Figure 8 supplement 1J,K). Like the results with Cis-Pt, however, 24h post-irradiation Xrp1-HA expression was enhanced several-fold by p53 inhibition (Figure 8 supplement 1 L, N). Like Xrp1 expression in aneuploid cells, p53-independent Xrp1 expression was significantly reduced in the RpS12 mutant background (Figure 8M,N). These results are consistent with the notion that, following irradiation or Cis-Pt feeding, the p53-dependent DDR suppresses the acquisition of aneuploidy, and that if p53 is inhibited then an increased frequency of aneuploid cells results, and these cells express Xrp1 in an RpS12-dependent manner. Xrp1 activity would then be expected to remove these cells by cell competition.

To explore the impact of p53 on genome integrity and Xrp1 activity further, we looked at IR-induced DNA damage using anti-γH2Av. Consistent with previous results, 24h after irradiation most acute DNA damage has been repaired although a limited amount remains, along with some continuing cell death (Figure 9A) (Wichmann, Jaklevic, and Su 2006; McNamee and Brodsky 2009). A portion of the cell death continues in the absence of p53 (Wichmann, Jaklevic, and Su 2006; McNamee and Brodsky 2009). Consistent with the conclusion that this includes competitive death of aneuploid cells (McNamee and Brodsky 2009; Ji et al. 2021), we found that the p53-independent cell death was substantially reduced by Xrp1 knockdown (Figure 9C,D). Anti-γH2Av labeling 24h after irradiation was greatly increased in the absence of p53, indicating persistence of DNA damage (Figure 9B). Anti-γH2Av labeling was even further increased after Xrp1 knockdown in the absence of p53, showing that Xrp1 contributes to the elimination of cells with DNA damage even in the absence of p53, presumably through cell competition that removes aneuploid cells. Consistent with this hypothesis, Xrp1 expression was

induced in a subset of the damaged cells that persisted when p53 was inhibited (Figure 9E,F).

To assess the effect of aneuploidy, the effect of Cis-Pt on clones of *rpS12* mutant cells was examined. Whereas the cell competition-defective $rpS12^{G97D}$ allele has no effect on clone growth in the wing disc in the absence of DNA damage (Figure 9 supplement 1 A-C) (Kale et al. 2018), $rpS12^{G97D/G97D}$ clones overgrew their $rpS12^{+/+}$ twin-spots in the presence of Cis-Pt (Figure 9 supplement 1 A-C), although the effect was smaller than that seen for p53 or Xrp1 mutants. Importantly, although RpS12 is implicated in the competitive removal of *Rp* mutant and aneuploid cells, it did not affect cell death in the acute DDR 2h after IR, suggesting that Xrp1 function in acute DDR depends on p53 not RpS12 (Figure 9 supplement 1D-F). These results are consistent with previous conclusions that Xrp1 expression can be induced by at least two independent pathways, by p53 in the acute DDR, and by RpS12 in the competitive elimination of $Rp^{+/-}$ cells and of aneuploid cells monosomic for one or more *Rp* gene loci, that latter typically occurring ~24h after acute DNA damage (Lee et al. 2018; Ji et al. 2021). Both mechanisms need to be considered to fully account for Xrp1 requirements in the presence of Cis-Pt.

DISCUSSION

Drosophila p53 shows sequence similarity and conserved overall structure with human p53, and is the key effector of the DDR in *Drosophila* as well as in mammals (Brodsky et al. 2000; Ollmann et al. 2000). There are some differences: in mammals, for example, DNA damage results in cell cycle (G1 and G2/M) arrest due to p53-dependent transcriptional activation of p21 and other targets 14-3-3σ and mir34a (Chen 2016), whereas cell cycle arrest in the *Drosophila* DDR is mediated by *grps/Chk1* and *mnk/Chk2*, independently of p53 (Brodsky et al. 2004). So far, *Drosophila* p53 has not been shown to be important for eliminating cells by competition, which is clearly a function of p53 in mammals (Baker 2020). Cell competition of multiple *Drosophila* genotypes, including the $Rp^{+/-}$ genotype where cell competition was first described, depends on the transcription factor Xrp1 (Lee et al. 2016; 2018; Baillon et al. 2018; Langton et al. 2021; Ochi et al. 2021; Kiparaki et al. 2022). Xrp1 is also a target of p53 in the acute DDR, suggesting that Xrp1 could mediate a relationship between p53 and cell competition in *Drosophila* (Baker, Kiparaki, and Khan 2019). In this work, we establish that Xrp1 is a major effector of p53

in the *Drosophila* DDR, and we show that differences in p53 activity cause cell competition

in *Drosophila*, as in mammals. Xrp1 retards the competitive expansion of $p53^{-/-}$ mutant

cells, however, through its p53-independent role in limiting accumulation of aneuploid

cells. This correlates with a role of Xrp1 in preventing accumulation of damaged cells in

$p53^{-/-}$ mutant clones under genotoxic stress, suggesting that it is these damaged cells that

are responsible for cell competition. Our results and conclusions are summarized in

Figure 10.

Xrp1 is an effector of p53 functions in the DDR

We show that, in imaginal discs, Xrp1 is induced downstream of p53 in the acute

phase of DDR at both RNA and protein levels (Figure 1). Unlike Xrp1 induction in $Rp^{+/-}$

cells, Xrp1 expression in the DDR occurs independently of autoregulatory transcription,

which may be why the Xrp1-LacZ enhancer trap shows less upregulation than the

endogenous transcript and protein. Multiple p53-dependent DDR genes, the majority in

our study, were Xrp1-dependent, including the pro-apoptotic gene *hid*. Even though *hid*-

reporters have been used as readouts of p53 activity (Fan et al. 2010; Zhang et al. 2014;

Ruiz-Losada et al. 2021; Chakravarti et al. 2022), they may be more accurately described

as targets of Xrp1. Other Xrp1 targets include DNA repair genes, GstD1, and Jun-kinase

signaling. Accordingly, Xrp1 is functionally required in the DDR, and in the absence of

Xrp1 DNA damage-induced cell death is reduced, although not eliminated, perhaps

because the pro-apoptotic gene *rpr* seems to be a direct target of p53 that does not require

Xrp1.

So far, integrating data from gene expression studies, binding motif and ChIP-Seq

searches, and global run-on sequencing of nascent mRNA has not yielded a consensus

on the core set of direct p53 target genes in mammals (Riley et al. 2008; Allen et al. 2014;

Fischer 2017). It is plausible that, as in *Drosophila*, mammalian p53 acts as a master

transcription factor that regulates a network of gene expression responses both directly

and also through intermediate transcription factors.

Cell competition due to differential p53 activity

If Xrp1 is activated and functions downstream of p53 in the DDR, we hypothesized

that differences in p53 activity between cells might lead to competition, as occurs between

cells with different Xrp1 levels in many examples of cell competition (Baker, Kiparaki,

and Khan 2019). To test this, we generated cells with chronic p53 activity by feeding the
genotoxic chemical Cis-Pt at concentrations that provoked p53 activity detectable with
the *rpr-LacZ* reporter but relatively little sign of acute DNA damage. Mutating p53 in
clones of cells then produced a situation where p53 activity was elevated in the wild type
cells compared to their p53-mutant neighbors. This clearly led to a competitive
disadvantage for wild type cells (Figure 6D-J). Clones of such cells showed diminished
growth in comparison to neutral clones of similar genotypes that did not have *p53*-mutant
sisters nearby (Figure 6K-M). Interestingly, the *p53*-mutant clones showed enhanced
growth, but our experiments did not permit us to distinguish whether this reflected the
absence of a cell-autonomous growth retardation that would normally result from p53
activity, or a non-autonomous stimulation due to out-competing nearby p53-wild type
cells.

These findings show that imaginal disc cells with higher p53 activity than their
neighbors suffer from cell competition with them, in *Drosophila* as in mammals. This has
not been recognized previously because *p53* mutant and wild type cells do not compete
in the absence of genotoxic stress, and because p53 is dispensable for the competitive
elimination of $Rp^{+/-}$ cells (Kale et al. 2015; de La Cova et al. 2004). $Rp^{+/-}$ cell competition is
independent of p53 because Xrp1 is induced in this genotype by RpS12, not p53 (Lee et
al. 2018). Given our present results, failure of wild type and *p53* mutant cells to compete
indicates that under normal conditions the baseline *p53* activity of imaginal disc cells is
below the threshold necessary for cell competition. Clonal expansion of *p53* mutant cells
in mammals indicates that this is not the case in mammalian tissues (Zhang et al. 2017).
The higher baseline p53 activity in mammals could reflect differences in p53 regulation
and turnover, but also could be caused simply by the much larger mammalian genome
providing a bigger target for spontaneous DNA damage. Many examples of mammalian
cell competition involving p53 activity also relate to genotoxic or mutant situations that
are expected to activate p53 (Nijhof et al. 2007; Bondar and Medzhitov 2010; Fernandez-
Antoran et al. 2019).

p53 depleted cells accumulate damage and compete with their neighbors
Our initial hypothesis was that, under chronic genotoxic stress, p53 would induce
Xrp1 expression through the DDR and that differences in Xrp1 expression would mediate
cell competition. Xrp1 expression was indeed induced by chronic Cis-Pt exposure.

Unexpectedly, however, Xrp1 expression also occurred independently of p53, and was
actually increased by p53 inhibition. Additionally, Xrp1 knock-down did not protect wild
type cells from competition with *p53* mutant cells under genotoxic stress, and may even
exacerbate it, while the overgrowth of *p53* mutant cells was potentiated by Xrp1-
depletion. These results were inconsistent with the idea of cell competition due to Xrp1
induction as part of the DDR, and indicate that Xrp1 in fact blunts an independent
pathway of cell competition in response to p53 differences. Below we suggest a plausible
explanation for these findings that is consistent with the available data.

One p53-independent route of Xrp1 expression is through RpS12. This occurs both
in $Rp^{+/-}$ cells and in cells with IR-induced genomic damage thought to change *Rp* gene
dose (Lee et al. 2018; Ji et al. 2021). Other mechanisms of Xrp1-induction, for example
following direct inhibition of translation factors, appear less dependent on RpS12
(Kiparaki et al. 2022). We showed here that Xrp1 expression in IR-damaged genomes was
also enhanced by p53 inhibition, and that this enhancement depended at least partially
on RpS12, the sensor of altered Rp gene dose. This suggests that, after irradiation, p53
function reduces the accumulation of cells with aneuploidy and other changes in genome
content that would induce Xrp1 expression through RpS12. We also found that p53
promoted the resolution of DNA damage and clearance of badly damaged cells following
irradiation, so that a great increase in cells with persistent unresolved DNA damage
resulted from irradiation when p53 was inhibited. The number of cells with unresolved
DNA damage is detectably increased further by inhibition of Xrp1 along with p53. Since
this represents a p53-independent role of Xrp1, it is more likely to represent failure to
eliminate cells with genome alterations by cell competition, rather than the role of Xrp1
in the DDR, because the latter is p53-dependent.

These studies of the response to IR now suggest a new hypothesis. Xrp1 expression
in imaginal discs exposed to Cis-Pt is not only due to a continuous DDR, mediated by
p53, but also occurs in sporadic cells where genotoxic stress has led to genome
rearrangements or aneuploidy, and hence to RpS12-dependent Xrp1 expression. This
could explain the salt-and-pepper Xrp1 expression pattern, and why Xrp1 expression in
the presence of Cis-Pt is enhanced by p53 inhibition. This notion is supported by the
finding that clones of cells homozygous for the cell competition-defective allele of *rpS12*
also have a growth advantage in the presence of Cis-Pt, like clones of cells mutant for *p53*
or *Xrp1* do.

Importantly, however, wild type cells exposed to Cis-Pt suffer a competitive growth disadvantage in the vicinity of clones of *p53* mutant cells that is independent of Xrp1. We suggest that this is related to the accumulation of DNA damage and genomic alterations that we demonstrated occurs in cells exposed to genotoxic stress that are unable to activate p53. Because this occurs even in the presence of wild type Xrp1, which we show removes many cells damaged by irradiation, we suggest that *p53* mutant clones contain cells with damaged genomes that cannot be recognized by the RpS12, Xrp1-dependent cell competition pathway that predominantly recognizes *Rp* gene haploinsufficiency. These could include cells with trisomies, or other increases in gene content that are not characterized by *Rp* haploinsufficiency, or cells with genetic alterations so substantial that they generate stress due to the mismatched levels of many proteins, not only ribosomal proteins. Evidence for just such a stress was originally observed in yeast strains with extra chromosomes, and has been demonstrated in *Drosophila* imaginal discs by inhibition of spindle assembly checkpoint genes, which leads to aneuploidy typically characterized by increased DNA content (Torres et al. 2007; Dekanty et al. 2012; Joy et al. 2021). It is already known that such genotypes can be growth-promoting and tumorigenic in *Drosophila* when cell death is prevented (Clemente-Ruiz and Milán 2015). Our model is that such cells accumulate in p53-deficient clones after genotoxic stress and exhibit enhanced growth and competitively inhibit nearby wild type cells, an effect that is further enhanced by the absence of Xrp1, which results in the accumulation of further damaged cells. In this model, differences in p53 activity do not lead to cell competition in *Drosophila* directly, but as a consequence of different levels of genomic damage that arise in cells with and without p53 function.

It would be interesting to examine whether genome damage underlies examples of p53-dependent cell competition observed in mammals. Our results, and model, are consistent with the idea that p53 suppresses tumorigenesis in part by promoting cell competition to remove abnormal cells, and that absence of p53 both abolishes this protection and facilitates the accumulation of hyperplastic aneuploid cells that may promote tumorigenesis.

ACKNOWLEDGEMENTS

We thank Drs. J. Secombe, M. Kiparaki, S. Nair, A. Kumar and S. Muliyil for comments on the manuscript. Fly stocks were obtained from the Bloomington Drosophila

Stock Center (supported by NIH P40OD018537) and the Vienna Drosophila Resource
Center. Confocal microscopy was performed in the Analytical Imaging Facility of the
Albert Einstein College of Medicine (supported by the NCI P30CA013330) using the Leica
SP8 microscope acquired through NIH SIG 1S10 OD023591. The monoclonal antibody
mAb40-1a developed by J.R.Sanes was obtained from the Developmental Studies
Hybridoma Bank, created by the NICHD and maintained at the University of Iowa,
Department of Biology, Iowa City, IA 52242. This project was supported by grant
GM104213 from the National Institutes of Health.

678
679

MATERIAL AND METHODS

1. Fly husbandry:

Flies were raised at 25^0C on standard cornmeal and molasses medium, containing
the following ingredients per 1L: 18g yeast; 22g molasses; 80g malt extract; 9g agar; 65g
cornmeal; 2.3g methyl para-benzoic acid; 6.35ml propionic acid. Larvae were not
distinguished for their sex prior to dissection.

2. DNA damage and genotoxic drug treatment:

γ-radiation: Flies were grown in the standard cornmeal medium at 25^0C. After egg
laying was done for 24 hours, the flies were transferred to a fresh vial. Following this, the
larvae were allowed to grow until the wandering late III instar stage and DNA damage
was induced by treating them with γ-radiation from cesium-137 using a shepherd
irradiator. For most of the experiments, 1500 Rads were administered, and dissection was
performed at a specified time point after the irradiation.

Cisplatin treatment: Cisplatin dosage was calibrated to sub-lethal concentration
as explained in the results section. To avoid using large amount of cisplatin, we used a
smaller volume (2.5ml) of fly food. Fresh solution was made every day from powdered
Cisplatin by dissolving it in sterilized water at a concentration of 4mM (3mg in 250ml),
and it was serially diluted such that the final concentration of Cisplatin in the 2.5ml fly
food was 0.2mM or 0.05mM. The Cisplatin solution was added to the fly food by
micropipette, drop by drop, making sure it disperses throughout the food.

3. Cell death and clone size quantification:

      Cell death: Cell death was quantified by counting number of Dcp-1 positive cells with the help of ImageJ cell counter plugin. For UAS-Gal4 experiments Dcp-1 positive cells in Gal4 to the control compartment was counted and compared as the ratio across different genetic perturbations. In case of mutant wing discs, the number of Dcp-1 positive cell was counted in the entire disc and absolute number were compared.

      Clone size quantification: Clone size was quantified from the wing pouch region, by manually drawing boundaries using ImageJ. The ratio of total clone area to total area of twin spot was computed and compared across different treatment or genotypes.

4. Immunohistochemistry:

      Imaginal discs were fixed using 4% formaldehyde in 1× PBS, followed by 2 h blocking with 0.5% bovine serum albumin (BSA) dissolved in 0.1% PBST (1×PBS with 0.1% Triton X-100). Primary antibody incubations were performed at 4°C overnight, followed by 2 h secondary incubation at room temperature. All washes following antibody incubation were performed in 0.1% PBST. Primary antibodies were against, rabbit anti-cleaved Dcp-1 (Cell Signaling 9578, 1:200), mouse anti-γH2Av (DSHB UNC93-5.2.1, 1:50), mouse anti-βGalactosidase (DSHB 40a-1, 1:50), mouse anti-p53 (DSHB 25F4, 1:50) and mouse anti-GFP (DSHB 12A6, 1:50), anti-Xrp1 (1:250; gift from Rio lab). Alexa-Fluor (Thermo Fisher-Invitrogen) and Cyanine dye (Jackson labs) conjugated secondary antibodies were used at 1:200 dilution. Image acquisition was performed on Leica-Sp8 microscope, and intensity adjustments were made using Photoshop CC 2018 and Image J was used for image processing.

      Previous experiments have supported that significant results could be obtained from 5 replicates, in most of the cases more than 5 sample were analyzed. Experiments were repeated at least 2 times or more for most of the results. No exclusion of the data points was made while performing statistical significance. For the quantitative data N values are indicted in the figure legends. Statistical significance was performed using student t-test for independent or dependent means as indicated in figure legends. For certain data sets one-way ANOVA with Bonferroni-Holm multiple comparison correction was performed as mentioned in the figure legends. Exact p values are indicated in the figures.

5. q-RT PCR:

Total RNA was extracted from wing imaginal discs of late third instar larvae treated with 2000 Rads of IR, untreated larvae were taken as control for normalization. Wing discs were collected in TRIzol reagents (Ambion) and RNA was isolated according to the manufacturer's instructions. RNA samples were DNase treated according to manufacturer's instructions (Thermo Fisher; AM1907) and reverse transcribed using Verso cDNA Synthesis Kit (Thermo Fisher; AB1453). The cDNA was used to perform the q-RT PCR reactions following the manufacturer's instructions for the TaqMan™ Fast Advanced Master Mix (Thermo Fisher; 4444557). Custom design FAM-MGB TaqMan probes were utilized for q-RT PCR. The gene expression values were normalized to housekeeping gene RpL32 and were also confirmed with actin as a second control (data not included). Following calculations were used for estimating fold change in gene expression $\Delta Ct = Ct$ (gene of interest) – Ct (housekeeping gene), $\Delta\Delta Ct = \Delta Ct$ (3-4 hr IR) – ΔCt (untreated average) and Fold gene expression $= 2\text{^} -(\Delta\Delta Ct)$. 3 biological replicates were analysed for each condition and genotype. The gene expression analysis was performed using technical triplicates for each sample. It has been previously established that significant results can obtained 3 biological and 3 technical triplicates.

The following TaqMan probes were used in this study:

1. p53 Dm02154335_g1
2. Xrp1 Dm02142088_g1
3. rpr Dm01823054_s1
4. hid Dm01823031_m1
5. GstD1 Dm02362792_s1
6. eiger Dm01794373_m1
7. rad50 Dm01817359_g1
8. PolZ1 Dm01843262_g1
9. DNAlig4 Dm01838632_g1
10. Ku80 Dm01791676_g1
11. *Drosophila* RpL32 Dm02151827_g1
12. *Drosophila* actin Dm02361909_s1

765
766

767 FIGURES AND FIGURE LEGENDS:

Khan and Baker, Figure. 1

Figure 1: DNA damage induced Xrp1 expression

Xrp1 protein levels in the wild type III instar wing discs stained for HA, in 0 Rads (control) (A), 2 hours after 1500 Rads (B), 4 hours after 1500 Rads (C) and in $Rp18^{+/-}$ mutant (D). Larval wing discs stained with Xrp1 antibody, 0 Rads (control) (E) and 4 hours after 1500 Rads (F) (note: Xrp1 antibody show high noise to signal ratio compared to Xrp1-HA). Xrp1-HA expression at 4 hours post 1500 Rads in the posterior compartment compared relative to the anterior compartment taken as an internal control, *en-Gal4* driving *UAS-Xrp1* RNAi (G), p53 dominant negative (*UAS-p53^{H159N}*) (H), and *UAS-Irbp18* RNAi (I). III instar wing discs assayed for *Xrp1^{02515}*-lacZ enhancer trap expression detected by anti-βgal antibody in 0 Rads (J), 4 hours after 1500 Rads (K) and in the wing disc

expressing *UAS-p53[H159N]* under control of *en-Gal4*, 4 hours after 1500 Rads (L). N= 7-8 discs

were analyzed for every genotype, each experiment was repeated at least 2-3 times.

Genotypes: (A-C) *Xrp1-HA*, (D) *p{hs:FLP}; FRT42 RpS18 p{ubi:GFP}*, (E-F) *w[1118]* (G) *en-*

Gal4::UAS-RFP, UAS-Xrp1 RNAi (VDRC 107860); *Xrp1-HA*, (H) *en-Gal4::UAS-RFP, UAS-*

p53DN[H159N] (Bl. 8420); *Xrp1-HA*, (I) *en-Gal4::UAS-RFP, UAS-Irbp18 RNAi; Xrp1-HA*, (J-K)

P{PZ}Xrp1[02515], (L) *en-Gal4::UAS-RFP,UAS-p53DN[H159N]* (Bl. 8420); *P{PZ}Xrp1[02515]*

Figure 1- Supplemental 1: Xrp1 RNA expression in response to DNA damage

qRT-PCR analysis for Xrp1 mRNA expression levels from III instar wing discs from the

genotypes *w[1118]*, *p53[5A-1-4]* and *Xrp1[m2-73]/Df1* collected 3-4 hours after irradiation (1500

RADs) and normalized to untreated (0 Rads) sample (A). III instar wing discs assayed for

Xrp1[02515]-lacZ enhancer trap expression by anti-βgal antibody in 0 Rads for *en>Xrp1 RNAi*

(B), *en>Irbp18 RNAi* (C) and 4 hours after 1500 Rads treatment in genotypes *en>Xrp1*

RNAi (D), *en>Irbp18 RNAi* (E). Number of wing discs analyzed for genotypes were N= 3

for B, N=2 for C, N=3 for D and N=5 for E, Data is plotted as mean± SEM, two-tailed

student t-test for independent means was performed for statistical significance.

Genotype: (A) w^{1118}, *yw; p53$^{[5A-1-4]}$*, and *Xrp1^{m2-73}*/ *Df 1*, (B & C) *en-Gal4::UAS-RFP, UAS-Xrp1*

RNAi (VDRC 107860); *P{PZ}Xrp1^{02515}* (C & E) *en-Gal4::UAS-RFP, UAS-Irbp18 RNAi;*

P{PZ}Xrp1^{02515}

Figure 2: Xrp1 serves as the effector of p53 in DNA damage response

Vein diagram showing the overlap of p53 (Akdemir et al. 2007; Brodsky et al. 2004) and Xrp1 (Lee et al. 2018; Baillon et al. 2018) target genes (A). Table showing the genes organized according to GO categories from radiation-induced p53-dependent (RIPD) and genes common between Xrp1- and p53- dependent transcription program (highlighted in bold) (B). Expression of *GstD*-lacZ in III instar wing discs measured by staining for anti-βgal antibody, in the wild type (no IR) (C), 4 hours after 2000 Rads in wild type disc (D), *UAS-p53^{H159N}* (E), *UAS-Xrp1 RNAi* (F) and *UAS-Irbp18* RNAi (G). Expression of *TRE-dsRed* in III instar wing discs measured by staining for anti-RFP antibody, in the wild type (no IR) (H), 4 hours after 1500 Rads in wild type disc (I), *UAS-p53^{H159N}* (J). *UAS-Xrp1 RNAi* (K) and *UAS-Irbp18 RNAi* (L). qRT-PCR analysis for mRNA expression levels from III instar wing discs of *w^{1118}*, *p53^{[5A-1-4]}* and *Xrp1^{m2-73}/Df1* collected 3-4 hours after irradiation (1500 RADs) and normalized to untreated (0 Rads) sample, for the following genes *GstD1* (M), *egr* (N), *rad50* (O), *mus205* (P), *lig4* (Q) and *Ku80* (R). N= 7-8 discs were analyzed for every genotype, each experiment was repeated at least 2-3 times. Data is plotted as mean± SEM, two-tailed student t-test for independent means was performed for statistical significance.

Genotype: (C & D) *GstD-LacZ*, (E) *GstD-LacZ, UAS-p53^{H159N}* (Bl. 8420); *hh-Gal4::UAS-GFP*, (F) *GstD-LacZ, UAS-Xrp1 RNAi* (VDRC 107860); *hh-Gal4::UAS-GFP*, (G) *GstD-LacZ, UAS-Irbp18 RNAi; hh-Gal4::UAS-GFP*, (H and I) *TRE-dsRed*, (J) *TRE-dsRed::en-Gal4, UAS-p53^{H159N}* (Bl. 8420), (K) *TRE-dsRed::en-Gal4, UAS-Xrp1 RNAi* (VDRC 107860), (L) *TRE-dsRed::en-Gal4, UAS-Irbp18 RNAi*, (M-R) *w^{1118}*, *yw; p53^{[5A-1-4]}*, and *Xrp1^{m2-73}/Df1*

Figure 3: Xrp1 controls expression of the regulatory sequences of pro-apoptotic gene

hid

Schematic of H99 locus with approximate positions of the *rpr*, *hid*, and *grm* and the

reporter lines used for the analysis (A). Expression of *hid5'F-wt-EGFP* measured by

staining for anti-GFP antibody, in the wild type (no IR) (B) and 4 hours after 2000 Rads

in wild type wing disc (C), *UAS-p53^{H159N}* (D), *UAS-Xrp1 RNAi* (E) and *UAS-Irbp18* RNAi

(F). *hid^{20-10}-lacZ* expression measured by anti-βgal staining, in the wild type (no IR) (G)

and 4 hours after 2000 Rads in wild type wing disc (H), *UAS-p53^{H159N}* (I), *UAS-Xrp1* (J)

and *UAS-Irbp18* RNAi (K). *rpr^{150}-lacZ* expression measured by anti-βgal staining, in the

wild type (no IR) (L) and 4 hours after 2000 Rads in wild type wing disc (M). *UAS-p53^{H159N}*

(N), *UAS-Xrp1* (O) and *UAS-Irbp18* RNAi (P). qRT-PCR analysis for mRNA expression

levels from III instar wing discs of *w^{1118}*, *p53^{[5A-1-4]}* and *Xrp1^{m2-73}/Df1* collected 3-4 hours after

irradiation (1500 RADs) and normalized to untreated (0 Rads) sample, for the following

genes, *rpr* (Q), and *hid* (R). Putative DNA binding motif of Xrp1/Irbp18 heterodimer,

adapted from Fly Factor Survey (S). Position of Xrp1/Irbp18 DNA motif in the upstream

regulatory region of *hid* relative to the TSS (T). N= 5-8 discs was analyzed for every

genotype, each experiment was repeated at least 2-3 times. Data is plotted as mean± SEM,

two-tailed student t-test for independent means was performed for statistical

significance.

Genotype: (B & C) *hid5'F-wt-EGFP*, (D) *hid5'F-wt-EGFP; en-Gal4::UAS-RFP, UAS-p53^{H159N}*

(Bl. 8420), (E) *hid5'F-wt-EGFP; en-Gal4::UAS-RFP, UAS-Xrp1 RNAi* (VDRC 107860), (F)

hid5'F-wt-EGFP; en-Gal4::UAS-RFP, UAS-Irbp18 RNAi, (G & H) *yw; hid^{20-10}-lacZ*, (I) *en-*

Gal4::UAS-RFP, UAS-p53^{H159N} (Bl. 8420); *hid^{20-10}-lacZ*, (J) *en-Gal4::UAS-RFP, UAS-Xrp1*

RNAi (VDRC 107860); *hid^{20-10}-lacZ*, (K) *en-Gal4::UAS-RFP, UAS-Irbp18 RNAi; hid^{20-10}-lacZ*,

(L & M) *rpr^{150}-lacZ*, (N) *rpr^{150}-lacZ, UAS-p53^{H159N}* (Bl. 8420); *hh-Gal4::UAS-GFP*, (O) *rpr^{150}-*

lacZ, UAS-Xrp1 RNAi (VDRC 107860); *hh-Gal4::UAS-GFP,* (P) *rpr^{150}-lacZ, UAS-Irbp18*

RNAi; hh-Gal4::UAS-GFP, (Q-R) *w^{1118}*, *yw; p53^{[5A-1-4]},* and *Xrp1^{m2-73}/Df1*

Figure 3- Supplemental 1: *rpr¹⁵⁰lacZ* **intensity quantification**

Average intensity quantification for *rpr¹⁵⁰lacZ* from the wing the anterior compartment of the wing pouch with and without IR, for the genotypes shown in Figure 3 L-P. *rpr¹⁵⁰lacZ* average intensity in the anterior compartments is elevated by IR, regardless of gene expression or depletion in the P compartments (A). The ratio of mean posterior to mean anterior LacZ expression for each. Only p53-DN reduced the relative *rpr-LacZ* expression level (B). Statistics: Data is plotted as mean± SEM; n=5 for *irbp18* RNAi, n=7 for other genotypes; significance assessed by one-way Anova with Bonferroni-Holm multiple comparison correction. For panel B, genotypes were compared only to the unirradiated control.

Figure 3 - source data file 1: intensity quantification for *rpr¹⁵⁰lacZ*

Genotype: (A & B) *rpr¹⁵⁰-lacZ*, *rpr¹⁵⁰-lacZ, UAS-p53^H159N* (Bl. 8420); *hh-Gal4::UAS-GFP*, *rpr¹⁵⁰-lacZ, UAS-Xrp1 RNAi* (VDRC 107860); *hh-Gal4::UAS-GFP, rpr¹⁵⁰-lacZ, UAS-Irbp18 RNAi; hh-Gal4::UAS-GFP*

Figure 4: Xrp1 is sufficient for the expression of p53- dependent DDR target genes

rpr^{150}-lacZ reporter expression measured by anti-βgal (red) and cell death by anti-Dcp-1 (green) in III instar eye imaginal disc expressing UAS-RFP (A) and UAS-Xrp1 long isoform (B), under the control of *GMR-Gal4* (blue). *hid$^{20-10}$-lacZ* reporter expression measured by anti-βgal (red) and cell death by anti-Dcp-1 (green) in III instar eye imaginal disc expressing UAS-RFP (C) and UAS-Xrp1 long isoform (D) under the control of *GMR-Gal4* (blue). *hid5'F-wt-EGFP* reporter expression measured by anti-GFP (red) and cell death by anti-Dcp-1 (green) in III instar eye imaginal disc expressing UAS-RFP (E) and UAS-Xrp1 long isoform (F) under the control of *GMR-Gal4* (blue). N= 7-8 discs were analyzed for every genotype; each experiment was repeated at least 2 times. All the control and experimental genotypes shown in this figure were kept at 18⁰C in order to lessen the effect of Xrp1 overexpression.

Genotype: (A-A') *GMR-Gal4::UAS-RFP, rpr^{150}-lacZ*, (B-B') *GMR-Gal4::UAS-RFP::UAS-Xrp1, rpr^{150}-lacZ*, (C-C') *GMR-Gal4::UAS-RFP; hid$^{20-10}$-lacZ*, (D-D') *GMR-Gal4::UAS-RFP::UAS-Xrp1; hid$^{20-10}$-lacZ*, (E-E') *hid5'F-wt-EGFP; GMR-Gal4::UAS-RFP*, (F-F') *hid5'F-wt-EGFP; GMR-Gal4::UAS-RFP, UAS-Xrp1*

Figure 4- Supplemental 1: Xrp1 is sufficient for the expression of p53- dependent DDR target genes

Xrp1-HA protein measured by anti-HA (red) and cell death by Dcp-1 (green) in the III instar wing disc expressing UAS-RFP control (A) and UAS-Xrp1 long isoform (B) under the control of *nub-Gal4* (blue). *rpr¹⁵⁰-lacZ* reporter expression measured by anti-βgal (red) and cell death by anti-Dcp-1 (green) in III instar wing imaginal disc expressing UAS-RFP (C) and UAS-Xrp1 (D) under the control of *nub-Gal4*. *TRE-EGFP* reporter expression measured by anti-GFP (red) and cell death by anti-Dcp-1 (green) in III instar wing imaginal disc expressing UAS-RFP control (E) and UAS-Xrp1 (F) under the control of *nub-Gal4* (blue). *GstD-lacZ* reporter expression measured by anti-GFP (red) and cell death by anti-Dcp-1 (green) in III instar wing imaginal disc expressing UAS-RFP (G) and UAS-Xrp1 (H) under the control of *nub-Gal4* (blue). N= 7-8 discs were analyzed for every genotype, each experiment was repeated at least 2 times. All the control and experimental

genotypes shown in this figure were kept at 18⁰C in order to lessen the effect of Xrp1

overexpression.

Genotype: (A) *nub-Gal4::UAS-RFP*, (B) *nub-Gal4::UAS-RFP::UAS-Xrp1-HA*, (C) *nub-*

Gal4::UAS-RFP, rpr¹⁵⁰-lacZ, (D) *nub-Gal4::UAS-RFP::UAS-Xrp1-HA, rpr¹⁵⁰-lacZ*, (E) *nub-*

Gal4::UAS-RFP, TRE-GFP, (F) *nub-Gal4::UAS-RFP::UAS-Xrp1-HA, TRE-GFP*, (G) *nub-*

Gal4::UAS-RFP, GstD-lacZ, (H) *nub-Gal4::UAS-RFP::UAS-Xrp1-HA, GstD-lacZ*

Figure 5: Xrp1 is partly required for DNA damage induced cell death

Drosophila III instar wing imaginal discs dissected at 2 hours after 1500 Rads, cell death is

measured by Dcp-1 (green and gray scale) in different genetic perturbations using en-

Gal4 (posterior compartment (red)), UAS-RFP (A), *UAS-p53^{H159N}* (B) and *UAS-Xrp1* RNAi

(C). Dcp-1 positive cells were quantified and plotted as the ratio of the number of Dcp-1
positive cells in anterior and posterior compartment (D). Wing discs dissected 4 hours
after 1500 RADs and stained for Dcp-1 positive cells in *FRT82* (E), *FRT82p53[5A-1-4]* (F), and
FRT82Xrp1m2-73 (G). (H) Dcp-1 positive cells quantified inside the clone (RFP-/-) and
inside the sister clone (RFP+/+). 4 hours timepoint was taken to have uniform
distribution of Dcp-1 positive cells. Statistics: (A-D) N= 15 each genotype. (E-H) N= 8
each genotype. Data is plotted as mean± SEM, two-tailed student t-test for independent
means for (A-D) and for dependent mean for (E-H) was performed for statistical
significance.

Figure 5 - source data file 1: intensity quantification for Dcp1 (Figure 5A-D)

Figure 5 - source data file 2: intensity quantification for Dcp1 (Figure 5E-H)

Genotype: (A) *en-Gal4::UAS-RFP,* (B) *en-Gal4::UAS-RFP, UAS-p53DNH159N* (Bl. 8420), (C)
en-Gal4::UAS-RFP, UAS-Xrp1 RNAi (Bl. 34521), (E) *p{hs:FLP}; FRT82-RFP/FRT82* (30 min
hs, 48 hr A.E.L), (F) *p{hs:FLP}; FRT82-RFP/FRT82-p53[5A-1-4]* (30 min hs, 48 hr A.E.L), (G)
p{hs:FLP}; FRT82-RFP/FRT82-Xrp1m2-73 (30 min hs, 48 hr A.E.L)

Figure 5- Supplemental 1: Xrp1 do not rescue p53 O/E phenotype

(A) Bright field photograph of *Drosophila* eye, rough eye phenotype due to constitutive expression of p53 under the control of GMR-promoter (*GMR-p53*). (B) Overexpression of dominant-negative p53 (*UAS-p53^H159N*) rescue the rough eye phenotype. (C) Xrp1 knock down by RNAi have slight effect on rough eye phenotype. (D) Control *nub-Gal4* exhibit few Dcp-1 positive cells and no p53 protein was detected by anti-p53 antibody. (E) *nub-Gal4* driving p53 overexpression (*GUS-p53*) in the wing pouch exhibit elevated cell death measured by Dcp-1 as well as p53 protein measured by anti-p53 antibody. (F) Co-

expressing *UAS-Xrp1* RNAi and *GUS-p53* under *nub-Gal4* lead to a decrease in cell death

by approximately 50%. (G) Removing one copy by $Xrp1^{m2-73}$/+ had no effect on cell death.

(H) Removing both copies of Xrp1 ($Xrp1^{m2-73}$/$Xrp1^{attp\,flox}$) lead to a decrease in cell death by

approximately 50%. Xrp1-HA expression assayed by anti-HA antibody in wing disc

expressing p53 under control of *nub-Gal4* (I & I'). (J) Dcp1 intensity was quantified using

ImageJ, N=6 for each genotype. Data is plotted as mean± SEM, two-tailed student t-test

for independent means. All the control and experimental genotypes shown in this figure

were kept at 18^0C to lessen the effect of p53 overexpression.

Figure 5 supplemental 1- source data file 1: Dcp1 quantification D-H and J

Genotype: (A) *GMR-p53,* (B) *GUS-p53, GMR-Gal4; UAS-p53^{H159N}* (Bl. 4021), (C) *GUS-p53,*

GMR-Gal4; UAS-Xrp1 RNAi (Bl. 4521), (D) *nub-Gal4; w^{1118}*, (E) *nub-Gal4, GUS-p53,* (F) *nub-*

Gal4, GUS-p53; UAS-Xrp1 RNAi (Bl. 4521), (G) *nub-Gal4, GUS-p53; Xrp1^{m2-73}/+,* (H) *nub-*

Gal4, GUS-p53; Xrp1^{m2-73}/Xrp1$^{attp\,flox}$, (I) *nub-Gal4, GUS-p53; Xrp1-HA*

Figure 6: Chronic exposure to sub-lethal genotoxic drug gives competitive advantage to *p53* mutant over wild type cells

(A) Schematic describing the timing for Cis-Pt treatment, clonal induction and dissection time point (made using BioRender software). (B-C) *rpr[150]-lacZ* reporter expression in III instar wing imaginal disc measured by anti-βgal in control no Cis-Pt and 96 hours after 0.05mM Cis-Pt treatment. (D-G) III instar wing imaginal disc stained for RFP, heterozygous cells one copy RFP (intermediate grey), sister clone with two copy of RFP (dark grey) and clone marked by absence of RFP (black), 76 hours after clonal induction (a.c.i) untreated discs (no Cis-Pt) *FRT82* (D) & *FRT82p53[5A-1-4]* (E) and 96 hours a.c.i + 0.05mM Cis-Pt treatment in *FRT82* (F) & *FRT82p53[5A-1-4]* (G). (H) Total clone (*FRT82/82*) to sister clone (*FRT82-RFP/82-RFP*) area measurements in individual discs plotted as ratio at 72 hours a.c.i in no Cis-Pt, 72 hours a.c.i + 0.05mM Cis-Pt and 96 hours a.c.i + 0.05mM Cis-Pt. (I) Untreated (no Cis-Pt) clone to treated (0.05mM Cis-Pt) clone area ratio (calculated per disc) for *FRT82* clone and for *p53[-/-]* mutant clone after 72 hours a.c.i. (J) Ratio of *FRT82* sister clone (*FRT82-RFP/82-RFP*) to *p53[+/+]* sister clone (*FRT82-RFP/82-RFP*) 72 hours a.c.i in untreated discs (no Cis-Pt) and 0.05mM Cis-Pt. (K-L) Neutral clone assay for confirming competitive disadvantage of sister clone associated with *p53[-/-]* mutant clones, images are shown for 0.05mM Cis-Pt condition. Control disc (*FRT42-alz/FRT42; Frt82-RFP*) with *FRT42* clone and its sister clone (*FRT42-alz/42-alz*) (K) and the same disc with *FRT82* clone and its sister clone (*FRT82-RFP/82-RFP*) (K'). Experimental disc (*FRT42-alz/FRT42; FRT82p53[5A-1-4]*) carrying *FRT42* clone and its sister clone (*FRT42-alz/42-alz*) (L), the same disc with *p53[-/-]* mutant clone and sister clone (*FRT82-RFP/82-RFP*) (L'). Measurements of *FRT42-alz/42-alz* sister clone to *FRT82-RFP/82-RFP* sister clone area under no Cis-Pt 72 hours a.c.i and 0.05mM Cis-Pt at 96 hours a.c.i in control discs and the *FRT42-alz/42-alz* sister clone to *FRT82-RFP/82-RFP* sister clone ratio in experimental discs in no Cis-Pt at 72 hours a.c.i and 0.05mM Cis-Pt 96 hours a.c.i (M). Note: for neutral clone assay the discs with non-overlapping *FRT42-alz* and *FRT82-RFP* were analyzed. This was done to avoid any effect of lack of p53 mutant clone on *FRT42-alz* clone within the same discs. Statistics: N= 10 discs each genotype (for D-J), data is plotted as mean± SEM, two-tailed student t-test for independent means was performed for statistical significance. N = 8 for (K-M), data is plotted as mean± SEM, two-tailed student t-test for dependent means was performed for statistical significance.

Figure 6 - source data file 1: Clone and sister clone area quantification D-J

Figure 6 - source data file 2: Clone and sister clone area measurement for neutral clone

assay K-M

Genotype: (B & C) rpr^{150}-lacZ, (D) *p{hs:FLP}; FRT82-RFP/ FRT82* (30 min hs, 48 hr A.E.L),

(E) *p{hs:FLP}; FRT82-RFP/ FRT82-p53$^{[5A-1-4]}$* (30 min hs, 48 hr A.E.L), (F) *p{hs:FLP}; FRT82-*

RFP/ FRT82 (30 min hs, 48 hr A.E.L), (G) *p{hs:FLP}; FRT82-RFP/ FRT82-p53$^{[5A-1-4]}$* (30 min

hs, 48 hr A.E.L), (K & K') *p{hs:FLP}; FRT42-alZ/ FRT42; FRT82-RFP/ FRT82* (20 min hs, 36

hr A.E.L), (L & L') *p{hs:FLP}; FRT42-alZ/ FRT42; FRT82-RFP/ FRT82-p53$^{[5A-1-4]}$* (20 min hs,

36 hr A.E.L)

Khan and Baker, Figure. 6- supplemental 1

Figure 6- Supplemental 1: Sub-cell lethal dosage of Cis-Pt do not cause heightened cell

death but give competitive advantage to *p53$^{-/-}$* mutant cells

(A) III instar wing imaginal discs assayed for cell death by anti-Dcp-1 staining and DNA

damage by anti-γH2Av antibody at different concentrations (0.01mM, 0.02mM, 0.05mM

and 0.5mM) of Cis-Pt at two different time points 72 or 96 hours after the treatment, as

marked in the figure. 0.5mM Cisplatin samples were analyzed at 15 hours' time point due

difficulty in recovering viable larvae at later time point 72 and 96 hours. (B-E) III instar

wing imaginal disc stained for RFP, heterozygous cells one copy RFP (intermediate grey),

sister clone with two copy of RFP (dark grey) and clone is marked by absence of RFP

(black) 72 hours after clonal induction in *FRT82* (B), *FRT82p53*[5A-1-4] (C) and 72 hours after

clonal induction (a.c.i) + 0.02mM Cis-Pt treatment in *FRT82* (D) & *FRT82p53*[5A-1-4] (E). (F)

Total clone to sister clone (*FRT82-RFP/82-RFP*) area measurements in individual discs,

plotted as ratio at 72 hours a.c.i in no Cis-Pt, and 72 hours a.c.i + 0.02mM Cis-Pt treatment.

(G-J) Bright field photograph of adult eye in untreated larvae (no Cis-Pt treatment) in

FRT82 (G) & *FRT82p53*[5A-1-4] (H), and in treated larvae (0.02mM Cis-Pt treatment) *FRT82*

(I) & *FRT82p53*[5A-1-4] (J). N= 10 discs each genotype in (B-F), data is plotted as mean± SEM,

two-tailed student t-test for independent means was performed for statistical

significance.

Figure 6 – Supplemental 1- source data file 1: Clone size quantification B-F

1018

1019 Genotype: (A) *w*[1118], (B-D) *p{hs:FLP}; FRT82-RFP/FRT82* (30 min hs, 48 hr A.E.L), (C-E)

1020 *p{hs:FLP}; FRT82-RFP/FRT82-p53*[5A-1-4] (30 min hs, 48 hr A.E.L), (G-I) *eyflp; FRT82-*

1021 *RFP/FRT82*, (H-J) *eyflp; FRT82-RFP/FRT82-p53*[5A-1-4]

1022

1023

Figure 6- Supplemental 2: *p53*$^{-/-}$ **mutant and wild clone do not show any significant differences in DDR upon 0.05mM Cis-Pt**

III instar wing imaginal discs stained for γH2Av (green and grey scale) and marked for clone (Black), sister clone (bright red) and heterozygous tissue (dull red). (A-B) γH2Av levels in *FRT82* (A), and *FRT82p53*[5A-1-4] (B) under no Cis-Pt condition after 72 hours after clonal induction (a.c.i). (C-D) 1500 Rads γ-radiation showed significantly higher levels of γH2Av, no significant difference was observed between clone and sister clone of *FRT82*

1031 (C), *p53^{-/-}* mutant clone showing higher number of γH2Av positive cells compared to sister

(C), *p53⁻ᐟ⁻* mutant clone showing higher number of γH2Av positive cells compared to sister

clone (D). (E-H) Slight overall increase in γ-H2Av level upon 0.05mM Cisplatin treatment

in *FRT82* (E), and *FRT82p53[5A-1-4]* (F) at 72 hours a.c.i and *FRT82* (G), and *FRT82p53[5A-1-4]*

(H) at 96 hours a.c.i. N = 15 discs observed and compared for γ-H2Av staining for each

genotype, experiment was repeated at least 2-3 times.

Genotype: (A, C, E & G) *p{hs:FLP}; FRT82-RFP/FRT82* (30 min hs, 48 hr A.E.L), (B, D, F &

H) *p{hs:FLP}; FRT82-RFP/ FRT82-p53[5A-1-4]* (30 min hs, 48 hr A.E.L)

1038

1039

Figure 7: Lack of Xrp1/irbp18 facilitate growth of cells over wild type cells

(A-D) III instar wing imaginal disc stained for RFP, heterozygous cells one copy RFP

(intermediate grey), sister clone with two copy of RFP (dark grey) and clone is marked

by absence of RFP (black) under control condition (no Cis-Pt) 72 hours after clonal

induction (a.c.i) in *FRT82* (A), *FRT82Xrp1^{m2-73}* (B) and 96 hours a.c.i + 0.05mM Cis-Pt

treatment in *FRT82* (C) & *FRT82Xrp1^{m2-73}* (D). (E) Total clone to total sister clone (*FRT82-*

RFP/82-RFP) area measurements in individual discs, plotted as ratio at 72 hours a.c.i + no

Cis-Pt, 72 hours and 96 hours after clonal induction + 0.05mM Cis-Pt treatment. (F-G) III

instar wing imaginal disc stained for GFP, heterozygous cells one copy GFP (intermediate

grey), sister clone with two copy of GFP (dark grey) and clone is marked by absence of

GFP (black) 96 hours a.c.i + 0.05mM Cis-Pt treatment in *FRT80* (F) & *irbp18^{[f05006]} FRT80*

(G). (H) Total clone to total sister clone (*GFP-FRT80/80-GFP*) area measurements in

individual discs, plotted as ratio at 72 hours a.c.i + no Cis-Pt, 72 hours and 96 a.c.i +

0.05mM Cis-Pt treatment. N= 10 discs each genotype, data is plotted as mean± SEM, two-

tailed student t-test for independent sample was performed for statistical significance.

Figure 7 - source data file 1: Clone size quantification A-E

Figure 7 - source data file 2: Clone size quantification F-H

1057

1058 Genotype: (A & C) *p{hs:FLP}; FRT82-RFP/ FRT82* (30 min hs, 48 hr A.E.L), (B & D)

1059 *p{hs:FLP}; FRT82-RFP/ FRT82-Xrp1^{m2-73}* (30 min hs, 48 hr A.E.L), (F) *p{hs:FLP}; GFP-FRT80/*

1060 *irbp18^{[f05006]}-FRT80* (30 min hs, 48 hr A.E.L), (G) *p{hs:FLP}; GFP-FRT80/irbp18^{[f05006]}-FRT80*

1061 (30 min hs, 48 hr A.E.L)

1062

Figure 8: Lack of Xrp1 co-operates with *p53*[-/-] mutant cell expansion

(A-B) III instar wing imaginal disc stained for RFP, heterozygous cells one copy RFP (intermediate grey, dull red), sister clone with two copy of RFP (dark grey, bright red) and clone is marked by absence of RFP (black) and the engrailed compartment (blue). Image showing *p53*[-/-] mutant and *FRT82-RFP/82-RFP* clone 72 hours after clonal induction (a.c.i), comparing the total clone to sister clone area in anterior versus posterior compartment in control (*en-Gal4::UAS-GFP; FRT82-RFP/FRT82-p53*[5A-1-4]) (A) and experimental (*en-Gal4::UAS-GFP, UAS-Xrp1 RNAi; FRT82-RFP/FRT82-p53*[5A-1-4]) (B). (C) Quantification of the *p53*[-/-] mutant clone to sister clone (*FRT82-RFP/82-RFP*) area in anterior and posterior compartment from the data shown in (A) and (B). (D-E) Images showing *p53*[-/-] mutant and sister clone (*Frt82-RFP/82-RFP*) 96 hours a.c.i + 0.05mM Cis-Pt treatment, comparing the total clone to sister clone area in anterior versus posterior

compartment in control (*en-Gal4::UAS-GFP;FRT82-RFP/FRT82-p53*[5A-1-4]) (D) and

experimental (*en-Gal4::UAS-GFP, UAS-Xrp1 RNAi; FRT82-RFP/FRT82-p53*[5A-1-4]) (E). (F)

Quantification of the *p53*[-/-] mutant clone to sister clone (*FRT82-RFP/82-RFP*) area in

anterior and posterior compartment from the data shown in (D) and (E). (G) Total *p53*[-/-]

mutant clone area quantified in anterior and posterior compartment from Cis-Pt treated

disc in control (*en-Gal4::UAS-GFP;FRT82-RFP/FRT82-p53*[5A-1-4]) and experimental (*en-*

Gal4::UAS-GFP, UAS-Xrp1 RNAi;FRT82-RFP/FRT82-p53[5A-1-4]). (H) Total *p53*[-/-] mutant

sister clone area quantified in anterior and posterior compartment from Cis-Pt treated

disc in control (*en-Gal4::UAS-GFP;FRT82-RFP/FRT82-p53*[5A-1-4]) and experimental (*en-*

Gal4::UAS-GFP, UAS-Xrp1 RNAi; FRT82-RFP/FRT82-p53[5A-1-4]). Statistics: (A-B, C) N= 12

for each genotype, (D-E, F-G) N= 16 for each genotype, data is plotted as mean± SEM,

One-way Anova with Bonferroni-Holm multiple comparison correction was performed

for panel C and G-H, Holm p values are displayed on the graphs.

Figure 8- source data file 1- clone area quantification A-H

1090

1091 Genotype: (A & D) *p{hs:FLP}; en-Gal4::UAS-GFP; FRT82-RFP/FRT82-p53*[5A-1-4] (30 min hs,

1092 48 hr A.E.L), (B & E) *p{hs:FLP}; en-Gal4::UAS-GFP, UAS-Xrp1 RNAi* (VDRC 107860);

1093 *FRT82-RFP/FRT82-p53*[5A-1-4] (30 min hs, 48 hr A.E.L)

1094

Figure 8- Supplemental 1: Xrp1 induction independent of p53

(A-C) III instar wing imaginal disc stained for RFP, heterozygous cells one copy RFP (intermediate grey), sister clone with two copy of RFP (dark grey) and clone is marked by absence of RFP (black) and stained for anti-Xrp1 (long isoform) in *FRT82* (A), *FRT82p53[5A-1-4]* (B) and *FRT82Xrp1^{m2-73}* (C). (D-G) Xrp1-HA expression in control disc (D), 72 hours (E), 96 hours (F) and 96 hours in p53 dominant negative (*UAS-p53^{H159N}*) after 0.05mM Cis-Pt (G). (I-M) Xrp1-HA expression in control disc (H), 24 hours after 1500 rads (I), *UAS-Xrp1* RNAi (J), *UAS-Irbp1* RNAi (K), *UAS-p53^{H159N}* (L) and *UAS-p53^{H159N}* in *rpS12^{G97D/G97D}* mutant background (M). (N) Ratio of Xrp1 positive cells in anterior to posterior compartment in *UAS-p53^{H159N}* and *UAS-p53^{H159N}* in *rpS12^{G97D/G97D}* mutant

1106 background. N= 8 for (L-N), data is plotted as mean± SEM, student t-test two-tailed for
1107 independent sample was performed for statistical significance.

1108 Figure 8 – Supplemental 1- source data file 1: quantification for Xrp1-HA positive cells L-
1109 N

1110

Genotype: (A) *p{hs:FLP}; FRT82-RFP/FRT82* (30 min hs, 48 hr A.E.L), (B) *p{hs:FLP}; FRT82-*
RFP/FRT82-p53[5A-1-4] (30 min hs, 48 hr A.E.L), (C) *p{hs:FLP}; FRT82-RFP/FRT82-Xrp1[m2-*
73](30 min hs, 48 hr A.E.L), (D-G) *Xrp1-HA*, (H & I) *Xrp1-HA*, (J) *en-Gal4::UAS-RFP; UAS-*
Xrp1 RNAi (VDRC 107860); *Xrp1-HA*, (K) *en-Gal4::UAS-RFP; UAS-Irbp18 RNAi; Xrp1-HA*,
(L) *en-Gal4::UAS-RFP; UAS-p53DN[H159N]* (Bl. 8420); *Xrp1-HA*, (M) *en-Gal4::UAS-RFP; UAS-*
p53DN[H159N] (Bl. 8420); *Xrp1-HA-rpS12[G97D/G97D]*

1117

1118

Figure 9: Xrp1 limits genomic instabilities by removing aneuploid cells in an p53-
independent manner

(A-C) III instar wing disc from larvae dissected 24 hours after 1500 Rads of γ-rays
treatment, stained for cell death (Dcp-1, green), DNA damage signaling (γH2Av, red) in

control (*en-Gal4, UAS-GFP*) (A), *p53*[-/-] mutant (B) and Xrp1 k/d in *p53*[-/-] mutant

background (C). (D) Quantification of Dcp-1 positive cells anterior versus posterior

compartment in *p53*[-/-] mutant and Xrp1 k/d in *p53*[-/-] mutant background. (E) γH2Av and

(F) Xrp1-HA levels 24 hours after irradiation in the posterior compartment of wing disc

expressing *p53*[H159N] under the control of engrailed Gal4. N= 15 for (A-D), data is plotted

as mean± SEM, student t-test two-tailed for independent sample was performed for

statistical significance. N= 8, for (E & F).

Figure 9- source data file 1- Dcp-1 positive cell number quantification A-D

1131

1132 Genotype: Figure 9 (A) *en-Gal4::UAS-GFP; FRT82-RFP/FRT82-p53*[5A-1-4], (B) *en-Gal4::UAS-*

1133 *GFP; FRT82-RFP/FRT82-p53*[5A-1-4], (C) *en-Gal4::UAS-GFP, UAS-Xrp1 RNAi* (VDRC

1134 107860)*; FRT82-RFP/FRT82-p53*[5A-1-4]

1135

Figure 9- Supplemental 1: RpS12 remove aneuploid cells by cell competition created by chronic 0.05mM Cis-Pt treatment

(A-B) III instar wing imaginal disc stained for GFP, heterozygous cells one copy GFP (intermediate grey), sister clone with two copy of GFP (dark grey) and clone is marked by absence of GFP (black) 96 hours after clonal induction (a.c.i) + 0.05mM Cis-Pt

treatment in *GFP-FRT80* (A) & *rpS12^{G97D}-FRT80* (B). (C) Total clone to total sister clone (*GFP-FRT80/GFP-FRT80*) area measurements in individual discs, plotted as ratio at 72 hours a.c.i + no Cis-Pt, 72 hours and 96 hours a.c.i + 0.05mM Cis-Pt treatment. (D-F) Cell death measured by anti-Dcp-1 (green and gray scale) in wing imaginal discs dissected at 2 hours after 1500 Rads, measured in en-Gal4 (posterior compartment (red)) versus anterior compartment *UAS-RFP; rpS12^{G97D}* (D), and *UAS-Xrp1 RNAi; rpS12^{G97D}* (E). Dcp-1 positive cells were quantified in anterior and posterior compartment and plotted as the ratio of the number of Dcp-1 positive cells (F). N= 12 discs for (A-C), N= 6 discs for (D-F), data is plotted as mean± SEM, student t-test two-tailed for independent sample was performed for statistical significance.

Figure 9 – Supplemental 1- source data file 1: Clone and sister area quantification A-C

Figure 9 – Supplemental 1- source data file 2: Dcp-1 positive cell number quantification D-F

Genotype: (A) *p{hs:FLP}; GFP-FRT80/ FRT82* (30 min hs, 48 hr A.E.L), (B) *p{hs:FLP}; GFP-FRT80/rpS12^{G97D/G97D}-FRT80* (30 min hs, 48 hr A.E.L), (D) *en-Gal4::UAS-RFP; rpS12^{G97D/G97D}*, (E) *en-Gal4::UAS-RFP, UAS-Xrp1 RNAi* (VDRC 107860); *rpS12^{G97D/G97D}*

1161

Figure 10: Model

(A) Wild type cells activate p53 in response to IR, p53 activate Xrp1 which in turn mediate many functions of p53 in DNA damage response. p53 via Xrp1 promote DNA repair, cell death to keep a check on accumulation of cells with abnormal genomic content. p53 strongly regulate cell death in comparison to Xrp1, which is important to remove damaged cells from the tissue, failure of which results in the accumulation of damaged cells that leads to compromised genomic integrity. Cells escaping p53-Xrp1 DDR are further removed by RpS12-Xrp1 dependent pathway that remove aneuploid cells via cell competition.

(B) Absence of p53, abrogates DDR (both DNA repair and cell death). Lack of DDR leads to build up of cells harboring abnormal genomic content (aneuploidy, chromosomal rearrangement, deletions). Persistence of such cells gives competitive growth advantage to $p53^{-/-}$ mutant cells. Part of this growth is opposed by p53-independent role of Xrp1 in removing aneuploid cells by cell competition. Simultaneous loss of both p53 and Xrp1 causes an additive by promoting persistence of damaged and aneuploid cells which in turn impart aggressive growth phenotype to the cells (hyperplastic aneuploidy).

1179

(C) $p53^{-/-}$ mutant (dark grey cells) and $p53^{+/+}$ (light grey cells) cells grow to the same extent

in unchallenged condition, $p53^{-/-}$ mutant cells gain competitive advantage over $p53^{+/+}$ cells

upon treatment with genotoxin, the competitive growth advantage is further enhanced

by simultaneous lack of both Xrp1 and p53 (black cells). We propose that the growth of

p53 and Xrp1 double mutant cells is further promoted due to greater accumulation of

cells with hyperplastic aneuploidies (black cells).

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

 Supplementary Table 1: Key Resources Table

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Drosophila melanogaster*)	Xrp1		FLYBASE: FBgn0261113	
gene (*Drosophila melanogaster*)	irbp18		FLYBASE: FBgn0036126	
genetic reagent (*D. melanogaster*)	p53		FLYBASE: FBgn0039044	
genetic reagent (*D. melanogaster*)	rpS12		FLYBASE: FBgn0286213	
genetic reagent (*D. melanogaster*)	*Xrp1^{m2-73}*	(Lee et al. 2016)	FLYBASE: FBal0346068	Bloomington Drosophila Stock Center #81270
genetic reagent (*D. melanogaster*)	*Xrp1^{02515}*	(Spradling et al. 1999)	FLYBASE: FBal0009448	Bloomington Drosophila Stock Center #11569
genetic reagent (*D. melanogaster*)	*Xrp1$^{attP\ flox}$*	(Blanco, Cooper, and Baker 2020)		Strain maintained in Dr. Nicholas Baker's lab

genetic reagent (*D. melanogaster*)	Xrp1[HA]	(Blanco, Cooper, and Baker 2020)		Strain maintained in Dr. Nicholas Baker's lab
genetic reagent (*D. melanogaster*)	UAS-Xrp1 (long isoform)	(Tsurui-Nishimura et al. 2013)	FLYBASE: FBal0284387	
genetic reagent (*D. melanogaster*)	*irbp18[05006]*	(Francis et al. 2016)	FLYBASE: FBal0185260	Exelisis Drosophila Collection #f05006
genetic reagent (*D. melanogaster*)	*rpS12[G97D]*		FLYBASE: FBgn0286213	Strain maintained in Dr. Nicholas Baker's lab
genetic reagent (*D. melanogaster*)	*p53[5A-1-4]*		FLYBASE: FBgn0039044	Bloomington Drosophila Stock Center #6815
genetic reagent (*D. melanogaster*)	UAS-p53.H159N.Ex-2 (Chr. II)		FLYBASE: FBti0040583	Bloomington Drosophila Stock Center #8420
genetic reagent (*D. melanogaster*)	UAS-p53.H159N.Ex -3 (Chr. III)		FLYBASE: FBti0040584	Bloomington Drosophila Stock Center #8421
genetic reagent (*D. melanogaster*)	UAS-dsRNA[Xrp1]	(Ni et al. 2011)	FLYBASE: FBti0144696	Bloomington Drosophila Stock Center #34521
genetic reagent (*D. melanogaster*)	UAS-dsRNA[Xrp1]		FLYBASE: FBst0479673	Vienna Drosophila Resource

				Center #107860
genetic reagent (*D. melanogaster*)	UAS-dsRNA^{irbp18}	(Ni et al. 2011)	FLYBASE: FBti0140125	Bloomington Drosophila Stock Center #33652
genetic reagent (*D. melanogaster*)	UAS-dsRNA^{w}	(Ni et al. 2011)	FLYBASE: FBtp0064645	Bloomington Drosophila Stock Center #33623
genetic reagent (*D. melanogaster*)	arm-LacZ	(Vincent, Girdham, and O'Farrell 1994)	FLYBASE: FBal0040819	
genetic reagent (*D. melanogaster*)	Ubi-GFP	(Davis, O'Farrell, and Girdham 1995)	FLYBASE: FBal0047085	
genetic reagent (*D. melanogaster*)	M(2)56F (mutating RpS18)	Laboratory of Y. Hiromi	FLYBASE: FBal0011916	
genetic reagent (*D. melanogaster*)	hs-FLP	(Struhl and Basler 1993)	FLYBASE: FBtp0001101	
genetic reagent (*D. melanogaster*)	ey-FLP	(Newsome, Åsling, and Dickson 2000)	FLYBASE: FBal0098303	
genetic reagent (*D. melanogaster*)	GMR-gal4	(Freeman 1996)	FLYBASE: FBtp0001315	
genetic reagent (*D. melanogaster*)	P{neoFRT}42D	(Xu and Rubin 1993)	FLYBASE: FBti0141188	Bloomington Drosophila Stock Center #1802

genetic reagent (*D. melanogaster*)	P{neoFRT}80 B	(Xu and Rubin 1993)	FLYBASE: FBti0002073	Bloomington Drosophila Stock Center #1988
genetic reagent (*D. melanogaster*)	P{neoFRT}82 B	(Xu and Rubin 1993)	FLYBASE: FBti0002074	Bloomington Drosophila Stock Center #2050, 2051
Antibody	Polyclonal Rabbit anti-XRP1 (short isoform)	(Francis et al. 2016)		1:200 dilution
Antibody	Mouse anti-HA Tag monoclonal	Cell Signalling Technology	Cat #2367 RRID:AB_10 691311	1:100 dilution
Antibody	Mouse anti-beta galactosidase (mAb40-1a) monoclonal	DSHB	RRID: AB_23 14509	1:100 dilution
Antibody	Rabbit anti-active-Dcp1 polyclonal	Cell Signalling Technology	Cat #9578 RRID:AB_27 21060	1:50 dilution
Antibody	Mouse anti-p53 (25F4) monoclonal	DSHB	RRID: AB_57 9787	1:20 dilution

1209
1210
1211

1212

1213

1214

1215

1216

1217

1218

1219

REFERENCES

Akdemir, F., A. Christich, N. Sogame, J. Chapo, and J. M. Abrams. 2007. "P53 Directs Focused Genomic Responses in Drosophila." *Oncogene* 26 (36): 5184–93. https://doi.org/10.1038/sj.onc.1210328.

Allen, Mary Ann, Zdenek Andrysik, Veronica L Dengler, Hestia S Mellert, Anna Guarnieri, Justin A Freeman, Kelly D Sullivan, et al. 2014. "Global Analysis of P53-Regulated Transcription Identifies Its Direct Targets and Unexpected Regulatory Mechanisms." *ELife* 3 (May). https://doi.org/10.7554/elife.02200.

Alpar, Lale, Cora Bergantiños, and Laura A. Johnston. 2018. "Spatially Restricted Regulation of Spätzle/Toll Signaling during Cell Competition." *Developmental Cell* 46 (6): 706-719.e5. https://doi.org/10.1016/j.devcel.2018.08.001.

Andrysik, Zdenek, Matthew D. Galbraith, Anna L. Guarnieri, Sara Zaccara, Kelly D. Sullivan, Ahwan Pandey, Morgan MacBeth, Alberto Inga, and Joaquín M. Espinosa. 2017. "Identification of a Core TP53 Transcriptional Program with Highly Distributed Tumor Suppressive Activity." *Genome Research* 27 (10): 1645–57. https://doi.org/10.1101/gr.220533.117.

Baillon, Ludovic, Federico Germani, R Claudia, Jochen Hilchenbach, and Konrad Basler. 2018. "Xrp1 Is a Transcription Factor Required for Cell Competition-Driven Elimination of Loser Cells." *Scientific Report*, no. July: 1–10. https://doi.org/10.1038/s41598-018-36277-4.

Baker, N E. 2017. "Mechanisms of Cell Competition Emerging from Drosophila Studies." *Current Opinion in Cell Biology* 48: 40–46. https://doi.org/10.1016/j.ceb.2017.05.002.

Baker, Fang Jin, Karthik B. Jeganathan, and Jan M. van Deursen. 2009. "Whole Chromosome Instability Caused by Bub1 Insufficiency Drives Tumorigenesis through Tumor Suppressor Gene Loss of Heterozygosity." *Cancer Cell* 16 (6): 475–86. https://doi.org/10.1016/j.ccr.2009.10.023.

Baker, N E. 2020. "Emerging Mechanisms of Cell Competition." *Nature Reviews Genetics* 21 (11): 683–97. https://doi.org/10.1038/s41576-020-0262-8.

Baker, N.E., M. Kiparaki, and C. Khan. 2019. "A Potential Link between P53, Cell Competition and Ribosomopathy in Mammals and in Drosophila." *Developmental Biology* 446 (1). https://doi.org/10.1016/j.ydbio.2018.11.018.

Blanco, Jorge, Jacob C. Cooper, and Nicholas E. Baker. 2020. "Roles of C/EBP Class BZip Proteins in the Growth and Cell Competition of Rp ('Minute') Mutants in Drosophila." *ELife* 9: 1–27. https://doi.org/10.7554/eLife.50535.

Bondar, Tanya, and Ruslan Medzhitov. 2010. "P53-Mediated Hematopoietic Stem and Progenitor Cell Competition." *Cell Stem Cell* 6 (4): 309–22. https://doi.org/10.1016/j.stem.2010.03.002.

Brennan, Christopher M., Laura Pontano Vaites, Jonathan N. Wells, Stefano Santaguida, Joao A. Paulo, Zuzana Storchova, J. Wade Harper, Joseph A. Marsh, and Angelika Amon. 2019. "Protein Aggregation Mediates Stoichiometry of Protein Complexes in Aneuploid Cells." *Genes and Development* 33 (15–16): 1031–47. https://doi.org/10.1101/gad.327494.119.

Brodsky, M H, W Nordstrom, G Tsang, E Kwan, G M Rubin, and J M Abrams. 2000. "Drosophila P53 Binds a Damage Response Element at the Reaper Locus." *Cell* 101: 103–13. https://doi.org/10.1016/S0092-8674(00)80627-3.

Brodsky, M H, Brian T. Weinert, Garson Tsang, Yikang S. Rong, Nadine M. McGinnis, Kent G.
Golic, Donald C. Rio, and Gerald M. Rubin. 2004. "Drosophila Melanogaster MNK/Chk2 and
P53 Regulate Multiple DNA Repair and Apoptotic Pathways Following DNA Damage."
Molecular and Cellular Biology 24 (3): 1219–31. https://doi.org/10.1128/mcb.24.3.1219-
1231.2004.
Brown, Brian, Sahana Mitra, Finnegan D Roach, Deepika Vasudevan, and Hyung Don Ryoo.
2021. "The Transcription Factor Xrp1 Is Required for PERK-Mediated Antioxidant Gene
Induction in Drosophila." *ELife* 10: 74047. https://doi.org/10.7554/eLife.
Brumby, A.M., and H.E. Richardson. 2003. "Scribblemutants Cooperate with Oncogenic Ras or
Notch to Cause Neoplastic Overgrowth in Drosophila." *The EMBO Journal* 22: 5769–79.
Bursac, Sladana, Maja Cokaric, Giulio Donati, and Sinisa Volarevic. 2014. "Activation of the
Tumor Suppressor P53 upon Impairment of Ribosome Biogenesis." *BBA - Molecular Basis
of Disease* 1842 (6): 817–30. https://doi.org/10.1016/j.bbadis.2013.08.014.
Chakraborty, Riddhita, Ying Li, Lei Zhou, and Kent G. Golic. 2015. "Corp Regulates P53 in
Drosophila Melanogaster via a Negative Feedback Loop." *PLoS Genetics* 11 (7): 1–25.
https://doi.org/10.1371/journal.pgen.1005400.
Chakravarti, Ananya, Heshani N. Thirimanne, Savanna Brown, and Brian R. Calvi. 2022.
"Drosophila P53 Isoforms Have Overlapping and Distinct Functions in Germline Genome
Integrity and Oocyte Quality Control." *ELife* 11 (January).
https://doi.org/10.7554/eLife.61389.
Chen, Jiandong. 2016. "The Cell-Cycle Arrest and Apoptotic Functions of P53 in Tumor Initiation
and Progression." *Cold Spring Harbor Perspectives in Medicine* 6 (3).
https://doi.org/10.1101/cshperspect.a026104.
Christophorou, M. A., I. Ringshausen, A. J. Finch, L. Brown Swigart, and G. I. Evan. 2006. "The
Pathological Response to DNA Damage Does Not Contribute to P53-Mediated Tumour
Suppression." *Nature* 443 (7108): 214–17. https://doi.org/10.1038/nature05077.
Christophorou, Maria A., Dionisio Martin-Zanca, Laura Soucek, Elizabeth R. Lawlor, Lamorna
Brown-Swigart, Emmy W. Verschuren, and Gerard I. Evan. 2005. "Temporal Dissection of
P53 Function in Vitro and in Vivo." *Nature Genetics* 37 (7): 718–26.
https://doi.org/10.1038/ng1572.
Clemente-Ruiz, Marta, and Marco Milán. 2015. "Epithelial Cell Division: Keeping Aneuploidy
Levels in Check." *Current Biology*. Cell Press. https://doi.org/10.1016/j.cub.2015.02.008.
Dasari, Shaloam, and Paul Bernard Tchounwou. 2014. "Cisplatin in Cancer Therapy: Molecular
Mechanisms of Action." *European Journal of Pharmacology* 740: 364–78.
https://doi.org/https://doi.org/10.1016/j.ejphar.2014.07.025.
Davis, Ilan, Patrick H. O'Farrell, and Charles, H. Girdham. 1995. "A Nuclear GFP That Marks
Nuclei in Living Drosophila Embryos; Maternal Supply Overcomes a Delay in the
Appearance of Zygotic Fluorescence." *Developmental Biology* 170 (2): 726–29.
Dekanty, A., L. Barrio, and M. Milán. 2015. "Contributions of DNA Repair, Cell Cycle Checkpoints
and Cell Death to Suppressing the DNA Damage-Induced Tumorigenic Behavior of
Drosophila Epithelial Cells." *Oncogene* 34 (8): 978–85.
https://doi.org/10.1038/onc.2014.42.
Dekanty, A., Lara Barrio, Mariana Muzzopappa, Herbert Auer, and Marco Milán. 2012.
"Aneuploidy-Induced Delaminating Cells Drive Tumorigenesis in Drosophila Epithelia."

Proceedings of the National Academy of Sciences of the United States of America 109 (50): 20549–54. https://doi.org/10.1073/pnas.1206675109.

Donati, G, S Bertoni, E Brighenti, M Vici, Trere´1 Trere´1, S Volarevic, L Montanaro, and M Derenzini. 2011. "The Balance between RRNA and Ribosomal Protein Synthesis Up- and Downregulates the Tumour Suppressor P53 in Mammalian Cells." *Oncogene* 30: 3274–88. https://doi.org/10.1038/onc.2011.48.

Fan, Y, T v Lee, D Xu, Z Chen, a-F Lamblin, H Steller, and a Bergmann. 2010. "Dual Roles of Drosophila P53 in Cell Death and Cell Differentiation." *Cell Death and Differentiation* 17 (6): 912–21. https://doi.org/10.1038/cdd.2009.182.

Fernandez-Antoran, David, Gabriel Piedrafita, Kasumi Murai, Swee Hoe Ong, Albert Herms, Christian Frezza, and Philip H. Jones. 2019. "Outcompeting P53-Mutant Cells in the Normal Esophagus by Redox Manipulation." *Cell Stem Cell* 25 (3): 329-341.e6. https://doi.org/10.1016/j.stem.2019.06.011.

Fischer, M. 2017. "Census and Evaluation of P53 Target Genes." *Oncogene* 36 (28): 3943–56. https://doi.org/10.1038/onc.2016.502.

Foijer, Floris, Viji M. Draviam, and Peter K. Sorger. 2008. "Studying Chromosome Instability in the Mouse." *Biochimica et Biophysica Acta - Reviews on Cancer*. https://doi.org/10.1016/j.bbcan.2008.07.004.

Francis, Malik Joseph, Siobhan Roche, Michael Jeffrey Cho, Eileen Beall, Bosun Min, Ronaldo Paolo Panganiban, and Donald C. Rio. 2016. "Drosophila IRBP BZIP Heterodimer Binds P-Element DNA and Affects Hybrid Dysgenesis." *Proceedings of the National Academy of Sciences of the United States of America* 113 (46): 13003–8. https://doi.org/10.1073/pnas.1613508113.

Freeman, Matthew. 1996. "Reiterative Use of the EGF Receptor Triggers Differentiation of All Cell Types in the Drosophila Eye." *Cell*. Vol. 87.

García-Castillo, Herbert, Ana Isabel Vásquez-Velásquez, Horacio Rivera, and Patricio Barros-Núñez. 2008. "Clinical and Genetic Heterogeneity in Patients with Mosaic Variegated Aneuploidy: Delineation of Clinical Subtypes." *American Journal of Medical Genetics, Part A* 146 (13): 1687–95. https://doi.org/10.1002/ajmg.a.32315.

Gogna, Rajan, and Eduardo Moreno. 2020. "Emerging Role of Cell Competition in Cancer." *Seminars in Cancer Biology*. Academic Press. https://doi.org/10.1016/j.semcancer.2020.04.005.

Hanks, Sandra, Kim Coleman, Sarah Reid, Alberto Plaja, Helen Firth, David FitzPatrick, Alexa Kidd, et al. 2004. "Constitutional Aneuploidy and Cancer Predisposition Caused by Biallelic Mutations in BUB1B." *Nature Genetics* 36 (11): 1159–61. https://doi.org/10.1038/ng1449.

Igaki, Tatsushi, Raymond A. Pagliarini, and Tian Xu. 2006. "Loss of Cell Polarity Drives Tumor Growth and Invasion through JNK Activation in Drosophila." *Current Biology* 16 (11): 1139–46. https://doi.org/10.1016/j.cub.2006.04.042.

Jean-Paul Vincent, Chaeles H. Girdham, and Patrick H. O'Farrell. 1994. "A Cell-Autonomous, Ubiquitous Marker for the Analysis of Drosophila Genetic Mosaics." *Developmental Biology* 164: 328–31.

Ji, Zhejun, Jacky Chuen, Marianthi Kiparaki, and Nicholas Baker. 2021. "Cell Competition Removes Segmental Aneuploid Cells from Drosophila Imaginal Disc-Derived Tissues Based on Ribosomal Protein Gene Dose." *ELife* 10 (April). https://doi.org/10.7554/eLife.61172.

Ji, Zhejun, Marianthi Kiparaki, Virginia Folgado, Amit Kumar, Jorge Blanco, Gerard Rimesso, Jacky Chuen, Yang Liu, Deyou Zheng, and Nicholas E. Baker. 2019. "Drosophila RpS12 Controls Translation, Growth, and Cell Competition through Xrp1." *PLoS Genetics* 15 (12). https://doi.org/10.1371/journal.pgen.1008513.

Jin, Shengkan, Sebastian Martinek, Woo S Joo, Jennifer R Wortman, Nebojsa Mirkovic, Andrej Sali, Mark D Yandell, Nikola P Pavletich, Michael W Young, and Arnold J Levine. 2000. "Identification and Characterization of a P53 Homologue in Drosophila Melanogaster" 97 (13).

Joy, Jery, Lara Barrio, Celia Santos-Tapia, Daniela Romão, Nikolaos Nikiforos Giakoumakis, Marta Clemente-Ruiz, and Marco Milán. 2021. "Proteostasis Failure and Mitochondrial Dysfunction Leads to Aneuploidy-Induced Senescence." *Developmental Cell* 56 (14): 2043-2058.e7. https://doi.org/10.1016/j.devcel.2021.06.009.

Kaiser, Alyssa M., and Laura D. Attardi. 2018. "Deconstructing Networks of P53-Mediated Tumor Suppression in Vivo." *Cell Death and Differentiation* 25 (1): 93–103. https://doi.org/10.1038/cdd.2017.171.

Kale, Zhejun Ji, Marianthi Kiparaki, Jorge Blanco, Gerard Rimesso, Stephane Flibotte, and Nicholas E Baker. 2018. "Ribosomal Protein S12e Has a Distinct Function in Cell Competition." *Developmental Cell* 44 (1): 42–55. https://doi.org/10.1016/j.devcel.2017.12.007.

Kale, W Li, C. H. Lee, and N E Baker. 2015. "Apoptotic Mechanisms during Competition of Ribosomal Protein Mutant Cells: Roles of the Initiator Caspases Dronc and Dream/Strica." *Cell Death and Differentiation* 22 (8): 1300–1312. https://doi.org/10.1038/cdd.2014.218.

Kanda, Hiroshi, and Tatsushi Igaki. 2020. "Mechanism of Tumor-Suppressive Cell Competition in Flies." *Cancer Science* 111 (10): 3409–15. https://doi.org/10.1111/cas.14575.

Kastenhuber, Edward R., and Scott W. Lowe. 2017. "Putting P53 in Context." *Cell* 170 (6): 1062–78. https://doi.org/10.1016/j.cell.2017.08.028.

Khan, C., S. Muliyil, C. Ayyub, and B.J. Rao. 2017. "The Initiator Caspase Dronc Plays a Non-Apoptotic Role in Promoting DNA Damage Signalling in D. Melanogaster." *Journal of Cell Science* 130 (18). https://doi.org/10.1242/jcs.200782.

Kiparaki, Marianthi, Chaitali Khan, Virginia Folgado-Marco, Jacky Chuen, Panagiotis Moulos, and Nicholas E Baker. 2022. "The Transcription Factor Xrp1 Orchestrates Both Reduced Translation and Cell Competition upon Defective Ribosome Assembly or Function." *ELife* 11 (February). https://doi.org/10.7554/elife.71705.

Kurtz, Paula, Amanda E. Jones, Bhavana Tiwari, Nichole Link, Annika Wylie, Charles Tracy, Helmut Krämer, and John M. Abrams. 2019. "Drosophila P53 Directs Nonapoptotic Programs in Postmitotic Tissue." *Molecular Biology of the Cell* 30 (11): 1339–51. https://doi.org/10.1091/mbc.E18-12-0791.

La Cova, Claire de, Mauricio Abril, Paola Bellosta, Peter Gallant, and Laura A. Johnston. 2004. "Drosophila Myc Regulates Organ Size by Inducing Cell Competition." *Cell* 117 (1): 107–16. https://doi.org/10.1016/S0092-8674(04)00214-4.

La Cova, Claire de, Nanami Senoo-Matsuda, Marcello Ziosi, D. Christine Wu, Paola Bellosta, Catarina M. Quinzii, and Laura A. Johnston. 2014. "Supercompetitor Status of Drosophila Myc Cells Requires P53 as a Fitness Sensor to Reprogram Metabolism and Promote Viability." *Cell Metabolism* 19 (3): 470–83. https://doi.org/10.1016/j.cmet.2014.01.012.

Lacroix, Matthieu, Romain Riscal, Giuseppe Arena, Laetitia Karine Linares, and Laurent le Cam. 2020. "Metabolic Functions of the Tumor Suppressor P53: Implications in Normal Physiology, Metabolic Disorders, and Cancer." *Molecular Metabolism*. Elsevier GmbH. https://doi.org/10.1016/j.molmet.2019.10.002.

Langton, Paul F., Michael E. Baumgartner, Remi Logeay, and Eugenia Piddini. 2021. "Xrp1 and Irbp18 Trigger a Feed-Forward Loop of Proteotoxic Stress to Induce the Loser Status." Edited by Norbert Perrimon. *PLOS Genetics* 17 (12): e1009946. https://doi.org/10.1371/journal.pgen.1009946.

Lee, C.-H., G. Rimesso, D. M. Reynolds, J. Cai, and N. E. Baker. 2016. "Whole-Genome Sequencing and IPLEX MassARRAY Genotyping Map an EMS-Induced Mutation Affecting Cell Competition in Drosophila Melanogaster." *G3:Genes|Genomes|Genetics* 6 (October): 3207–17. https://doi.org/10.1534/g3.116.029421.

Lee, Marianthi Kiparaki, Jorge Blanco, Virginia Folgado, Zhejun Ji, Amit Kumar, Gerard Rimesso, and Nicholas E. Baker. 2018. "A Regulatory Response to Ribosomal Protein Mutations Controls Translation, Growth, and Cell Competition." *Developmental Cell*, 1–14. https://doi.org/10.1016/j.devcel.2018.07.003.

Levine, a J, W Hu, and Z Feng. 2006. "The P53 Pathway: What Questions Remain to Be Explored?" *Cell Death and Differentiation* 13 (6): 1027–36. https://doi.org/10.1038/sj.cdd.4401910.

Li, Tongyuan, Ning Kon, Le Jiang, Minjia Tan, Thomas Ludwig, Yingming Zhao, Richard Baer, and Wei Gu. 2012. "Tumor Suppression in the Absence of P53-Mediated Cell-Cycle Arrest, Apoptosis, and Senescence." *Cell* 149 (6): 1269–83. https://doi.org/10.1016/j.cell.2012.04.026.

Li, Chenxi Wu, Xiang Ding, Wenzhe Li, and Lei Xue. 2020. "Toll Signaling Promotes JNK-Dependent Apoptosis in Drosophila." *Cell Division* 15 (1). https://doi.org/10.1186/s13008-020-00062-5.

Madigan, James P, Heather L Chotkowski, and Robert L Glaser. 2002. "DNA Double-Strand Break-Induced Phosphorylation of Drosophila Histone Variant H2Av Helps Prevent Radiation-Induced Apoptosis." *Nucleic Acids Research* 30 (17): 3698–3705. https://doi.org/10.1093/nar/gkf496.

Marusyk, Andriy, Christopher C. Porter, Vadym Zaberezhnyy, and James DeGregori. 2010. "Irradiation Selects for P53-Deficient Hematopoietic Progenitors." *PLoS Biology* 8 (3). https://doi.org/10.1371/journal.pbio.1000324.

Maruyama, Takeshi, and Yasuyuki Fujita. 2017. "Cell Competition in Mammals — Novel Homeostatic Machinery for Embryonic Development and Cancer Prevention." *Current Opinion in Cell Biology* 48: 106–12. https://doi.org/https://doi.org/10.1016/j.ceb.2017.06.007.

Marygold, Steven J., John Roote, Gunter Reuter, Andrew Lambertsson, Michael Ashburner, Gillian H. Millburn, Paul M. Harrison, et al. 2007. "The Ribosomal Protein Genes and Minute Loci of Drosophila Melanogaster." *Genome Biology* 8 (10). https://doi.org/10.1186/gb-2007-8-10-r216.

Matsuura, Shinya, Yoshiyuki Matsumoto, Ken Ichi Morishima, Hideki Izumi, Hiroshi Matsumoto, Emi Ito, Keisuke Tsutsui, et al. 2006. "Monoallelic BUB1B Mutations and Defective Mitotic-Spindle Checkpoint in Seven Families with Premature Chromatid Separation (PCS)

Syndrome." *American Journal of Medical Genetics* 140 A (4): 358–67.
https://doi.org/10.1002/ajmg.a.31069.

McNamee, Laura M., and Michael H. Brodsky. 2009. "P53-Independent Apoptosis Limits DNA
Damage-Induced Aneuploidy." *Genetics* 182 (2): 423–35.
https://doi.org/10.1534/genetics.109.102327.

Merino, Marisa M., Christa Rhiner, Jesus M. Lopez-Gay, David Buechel, Barbara Hauert, and
Eduardo Moreno. 2015. "Elimination of Unfit Cells Maintains Tissue Health and Prolongs
Lifespan." *Cell* 160 (3): 461–76. https://doi.org/10.1016/j.cell.2014.12.017.

Morata, Ginés. 2021. "Cell Competition: A Historical Perspective." *Developmental Biology*.
Elsevier Inc. https://doi.org/10.1016/j.ydbio.2021.02.012.

Morata, Ginés, and Pedro Ripoll. 1975. "Minutes: Mutants of Drosophila Autonomously
Affecting Cell Division Rate." *Developmental Biology* 42 (2): 211–21.
https://doi.org/https://doi.org/10.1016/0012-1606(75)90330-9.

Mukherjee, M., G. Ge, N. Zhang, D. G. Edwards, P. Sumazin, S. K. Sharan, P. H. Rao, D. Medina,
and D. Pati. 2014. "MMTV-Espl1 Transgenic Mice Develop Aneuploid, Estrogen Receptor
Alpha (ER)-Positive Mammary Adenocarcinomas." *Oncogene* 33 (48): 5511–22.
https://doi.org/10.1038/onc.2013.493.

Muller, Patricia A J, and Karen H. Vousden. 2014. "Mutant P53 in Cancer: New Functions and
Therapeutic Opportunities." *Cancer Cell* 25 (3): 304–17.
https://doi.org/10.1016/j.ccr.2014.01.021.

Nagata, Rina, and Tatsushi Igaki. 2018. "Cell Competition: Emerging Mechanisms to Eliminate
Neighbors." *Development Growth and Differentiation*. Blackwell Publishing.
https://doi.org/10.1111/dgd.12575.

Newsome, Timothy P., Bengt Åsling, and Barry J. Dickson. 2000. "Analysis of Drosophila
Photoreceptor Axon Guidance in Eye-Specific Mosaics." *Development* 127: 851–60.

Ni, Jian Quan, Rui Zhou, Benjamin Czech, Lu Ping Liu, Laura Holderbaum, Donghui Yang-Zhou,
Hye Seok Shim, et al. 2011. "A Genome-Scale ShRNA Resource for Transgenic RNAi in
Drosophila." *Nature Methods* 8 (5): 405–7. https://doi.org/10.1038/nmeth.1592.

Nijhof, Joanne G W, Aat M Mulder, Ewoud N Speksnijder, Esther M Hoogervorst, Leon H F
Mullenders, and Frank R de Gruijl. 2007. "Growth Stimulation of UV-Induced DNA Damage
Retaining Epidermal Basal Cells Gives Rise to Clusters of P53 Overexpressing Cells." *DNA
Repair* 6 (11): 1642–50. https://doi.org/https://doi.org/10.1016/j.dnarep.2007.05.008.

Ochi, Naotaka, Mai Nakamura, Rina Nagata, Naoki Wakasa, Ryosuke Nakano, and Tatsushi Igaki.
2021. "Cell Competition Is Driven by Xrp1-Mediated Phosphorylation of Eukaryotic
Initiation Factor 2α." *PLoS Genetics* 17 (12).
https://doi.org/10.1371/JOURNAL.PGEN.1009958.

Ollmann, Michael, Lynn M Young, Charles J di Como, Felix Karim, Marcia Belvin, Stephanie
Robertson, Kellie Whittaker, et al. 2000. "Drosophila P53 Is a Structural and Functional
Homolog of the Tumor Suppressor P53." *Cell* 101: 91–101.

Picco, V., and G. Pages. 2013. "Linking JNK Activity to the DNA Damage Response." *Genes &
Cancer* 4 (9–10): 360–68. https://doi.org/10.1177/1947601913486347.

Pilley, Steven, Tristan A Rodriguez, and Karen H Vousden. 2021. "Mutant P53 in Cell-Cell
Interactions." *Gene & Dev* 35: 433–48. https://doi.org/10.1101/gad.347542.

Riley, Todd, Eduardo Sontag, Patricia Chen, and Arnold Levine. 2008. "Transcriptional Control of Human P53-Regulated Genes." *Nature Reviews Molecular Cell Biology* 9 (5): 402–12. https://doi.org/10.1038/nrm2395.

Rohr, Cristian O, R Gonzalo Parra, Patricio Yankilevich, and Carolina Perez-Castro. 2013. "INSECT: IN-Silico SEarch for Co-Occurring Transcription Factors." *Bioinformatics* 29 (22): 2852–58. https://doi.org/10.1093/bioinformatics/btt506.

Ruiz-Losada, Mireya, Raul González, Ana Peropadre, Alejandro Gil-Gálvez, Juan J. Tena, Antonio Baonza, and Carlos Estella. 2021. "Coordination between Cell Proliferation and Apoptosis after DNA Damage in Drosophila." *Cell Death and Differentiation*. https://doi.org/10.1038/s41418-021-00898-6.

Sanchez, Juan A., Duarte Mesquita, Marýa C. Ingaramo, Federico Ariel, Marco Milan, and Andres Dekanty. 2019. "Eiger/TNFα-Mediated Dilp8 and ROS Production Coordinate Intra-Organ Growth in Drosophila." *PLoS Genetics* 15 (8). https://doi.org/10.1371/journal.pgen.1008133.

Singla, Shruti, Lisa K. Iwamoto-Stohl, Meng Zhu, and Magdalena Zernicka-Goetz. 2020. "Autophagy-Mediated Apoptosis Eliminates Aneuploid Cells in a Mouse Model of Chromosome Mosaicism." *Nature Communications* 11 (1). https://doi.org/10.1038/s41467-020-16796-3.

Snape, Katie, Sandra Hanks, Elise Ruark, Patricio Barros-Núñez, Anna Elliott, Anne Murray, Andrew H. Lane, et al. 2011. "Mutations in CEP57 Cause Mosaic Variegated Aneuploidy Syndrome." *Nature Genetics* 43 (6): 527–29. https://doi.org/10.1038/ng.822.

Spradling, Allan C, Dianne Stern, Amy Beaton, E Jay Rhem, Todd Laverty, Nicole Mozden, Sima Misra, and Gerald M Rubin. 1999. "The Berkeley Drosophila Genome Project Gene Disruption Project: Single P-Element Insertions Mutating 25% of Vital Drosophila Genes." *Genetics* 153: 135–77. https://academic.oup.com/genetics/article/153/1/135/6047835.

Struhl, Gary, and Konrad Basler. 1993. "Organizing Activity of Wingless Protein in Drosophila." *Cell*. Vol. 72.

Sykiotis, Gerasimos P., and Dirk Bohmann. 2008. "Keap1/Nrf2 Signaling Regulates Oxidative Stress Tolerance and Lifespan in Drosophila." *Developmental Cell* 14 (1): 76–85. https://doi.org/10.1016/j.devcel.2007.12.002.

Tanaka-Matakatsu, Miho, Jinhua Xu, Leping Cheng, and Wei Du. 2009. "Regulation of Apoptosis of Rbf Mutant Cells during Drosophila Development." *Developmental Biology* 326 (2): 347–56. https://doi.org/10.1016/j.ydbio.2008.11.035.

Tian Xu, and Gerald M. Rubin. 1993. "Analysis of Genetic Mosaics in Developing and Adult Drosophila Tissues." *Development* 117: 1223–37.

Titen, Simon W A, and Kent G. Golic. 2008. "Telomere Loss Provokes Multiple Pathways to Apoptosis and Produces Genomic Instability in Drosophila Melanogaster." *Genetics* 180 (4): 1821–32. https://doi.org/10.1534/genetics.108.093625.

Torres, Eduardo M, Tanya Sokolsky, Cheryl M Tucker, Leon Y Chan, Monica Boselli, Maitreya J Dunham, and Angelika Amon. 2007. "Effects of Aneuploidy on Cellular Physiology and Cell Division in Haploid Yeast." *Science* 317 (5840): 916–24. https://www.science.org.

Tsurui-Nishimura, Noriko, Thanh Quang Nguyen, Tomonori Katsuyama, Tatsurou Minami, Hirofumi Furuhashi, Yoshiteru Oshima, and Shoichiro Kurata. 2013. "Ectopic Antenna Induction by Overexpression of CG17836/Xrp1 Encoding an AT-Hook DNA Binding Motif

Protein in Drosophila." *Bioscience, Biotechnology, and Biochemistry* 77 (2): 339–44.
https://doi.org/10.1271/bbb.120756.
Uechi, Tamayo, Tatsuo Tanaka, and Naoya Kenmochi. 2001. "A Complete Map of the Human
Ribosomal Protein Genes: Assignment of 80 Genes to the Cytogenetic Map and
Implications for Human Disorders." *Genomics* 72 (3): 223–30.
https://doi.org/10.1006/geno.2000.6470.
Vishwakarma, Medhavi, and Eugenia Piddini. 2020. "Outcompeting Cancer." *Nature Reviews*
Cancer 20 (3): 187–98. https://doi.org/10.1038/s41568-019-0231-8.
Wagstaff, Laura, Maja Goschorska, Kasia Kozyrska, Guillaume Duclos, Iwo Kucinski, Anatole
Chessel, Lea Hampton-O'Neil, et al. 2016. "Mechanical Cell Competition Kills Cells via
Induction of Lethal P53 Levels." *Nature Communications* 7 (April).
https://doi.org/10.1038/ncomms11373.
Wei, Chia-Lin, Qiang Wu, Vinsensius B Vega, Kuo Ping Chiu, Patrick Ng, Tao Zhang, Atif Shahab,
et al. 2006. "A Global Map of P53 Transcription-Factor Binding Sites in the Human
Genome." *Cell* 124 (1): 207–19. https://doi.org/10.1016/j.cell.2005.10.043.
Wells, Brent S., and Laura A. Johnston. 2012. "Maintenance of Imaginal Disc Plasticity and
Regenerative Potential in Drosophila by P53." *Developmental Biology* 361 (2): 263–76.
https://doi.org/10.1016/j.ydbio.2011.10.012.
Wichmann, Anita, Burnley Jaklevic, and Tin Tin Su. 2006. "Ionizing Radiation Induces Caspase-
Dependent but Chk2- and P53-Independent Cell Death in Drosophila Melanogaster."
Proceedings of the National Academy of Sciences 103 (26): 9952–57.
https://doi.org/10.1073/pnas.0510528103.
Wichmann, Anita, Lyle Uyetake, and Tin Tin Su. 2010. "E2F1 and E2F2 Have Opposite Effects on
Radiation-Induced P53-Independent Apoptosis in Drosophila." *Developmental Biology* 346
(1): 80–89. https://doi.org/10.1016/j.ydbio.2010.07.023.
Zhang, Bingqing, Sonam Mehrotra, Wei Lun Ng, and Brian R. Calvi. 2014. "Low Levels of P53
Protein and Chromatin Silencing of P53 Target Genes Repress Apoptosis in Drosophila
Endocycling Cells." *PLoS Genetics* 10 (9). https://doi.org/10.1371/journal.pgen.1004581.
Zhang, Yinyin Xie, Ying Zhou, Cong Xiang, Lai Chen, Chenxi Zhang, Xiaoshuang Hou, Jiong Chen,
Hui Zong, and Geng Liu. 2017. "P53 Pathway Is Involved in Cell Competition During Mouse
Embryogenesis." *Proceedings of the National Academy of Sciences* 114 (3): 498–503.
https://doi.org/10.1073/pnas.1617414114.
Zhu, Lihua Julie, Ryan G Christensen, Majid Kazemian, Christopher J Hull, Metewo Selase
Enuameh, Matthew D Basciotta, Jessie A Brasefield, et al. 2010. "FlyFactorSurvey: A
Database of Drosophila Transcription Factor Binding Specificities Determined Using the
Bacterial One-Hybrid System ." *Nucleic Acids Research* 39 (suppl_1): D111–17.
https://doi.org/10.1093/nar/gkq858.

1563