Title: Social effects of rabies infection in male vampire bats (*Desmodus rotundus*)

Elsa M. Cárdenas Canales[*1]
Sebastian Stockmaier[*2]
Eleanor Cronin[2]
Tonie Rocke[3]
Jorge E. Osorio[1]
Gerald G. Carter[2,4]

*co-first authors

Corresponding authors: EMCC (crdenascanal@wisc.edu), SS (sebastian.stockmaier24@gmail.com)

Affiliations
1. Department of Pathobiological Sciences, School of Veterinary Medicine, University of Wisconsin-Madison, Madison, WI 53706, USA
2. Department of Evolution, Ecology and Organismal Biology, The Ohio State University
3. U.S. Geological Survey, National Wildlife Health Center, Madison, WI 53711, USA
4. Smithsonian Tropical Research Institute, Balboa Ancón, Panama

Word counts: Title (11); Abstract (179); Main text + figure/table legends + acknowledgements/funding (2497)

Abstract (resumen en español, S1)

Rabies virus (RABV) transmitted by the common vampire bat (*Desmodus rotundus*) poses a threat to agricultural development and public health throughout the Neotropics. The ecology and evolution of rabies host-pathogen dynamics are influenced by two infection-induced behavioral changes. RABV-infected hosts often exhibit increased aggression which facilitates transmission, and rabies also leads to reduced activity and paralysis prior to death. Although several studies document rabies-induced behavioral changes in rodents and other dead-end hosts, surprisingly few studies have measured these changes in vampire bats, the key natural reservoir throughout Latin America. Here, we take advantage of an experiment designed to test the safety and efficacy of an oral rabies vaccine in captive male vampire bats to quantify for the first time how rabies affects allogrooming and aggressive behaviors in the vampire bat. Compared to non-rabid vampire bats, rabid individuals reduced their allogrooming prior to death, but we did not detect increases in aggression among bats. To put our results in context, we review what is known and what remains unclear about behavioral changes of rabid vampire bats.

Introduction

Rabies virus (RABV) transmitted by the blood-feeding common vampire bat (*Desmodus rotundus*) creates a substantial burden for agricultural development and public health throughout Latin America, with deadly rabies outbreaks occurring in livestock [1–3] and humans [4,5]. RABV is transmitted by direct contact between the virus-laden saliva of the infected bat and the other animal's broken skin, eyes, or mucous membranes. The common vampire bat is a key vector because RABV transmissions can occur both among vampire bats and when they

bite livestock, wildlife, or less frequently, humans, leading to cross-species transmission [2,6].
Recent studies combining mathematical modeling, RABV phylodynamics, and the ecology,
demography, and dispersal of vampire bats have shown great potential to predict and mitigate
these pathogen spillover events [6–9].
Surprisingly few studies, however, have directly investigated how RABV infection affects
vampire bat behavior. In mustelids [10,11], canines [12,13], rodents [14], and humans [15],
RABV can lead to paralysis without obvious increases in aggression before death ("paralytic"
rabies), but it can also induce aggression and biting ("furious" rabies) which is likely to increase
transmission to other hosts (pathogen manipulation) [16]. Given that aggressive interactions are
commonly observed in vampire bats, especially among males [17–19], and that RABV is
detectable in the saliva at the end of infection [20,21], increases in aggression in rabid vampire
bats should enhance transmission. In a previous study [21], seven confirmed naturally RABV-
exposed vampire bats showed no obvious symptoms and survived, while seven others
presented two distinct disease outcomes. Three bats showed *furious* rabies presentation with
hypersalivation, excess vocalizations, teeth chattering, aggression towards handlers and other
bats, and irritability to light and sound. Four bats showed *paralytic* rabies presentation with
social isolation, lethargy, and apparent respiratory distress. Although these anecdotal
observations demonstrate both presentations are possible, they appear in some studies but not
others (Table 1), and the relative probability of paralytic versus furious symptoms in rabid
vampire bats remains unclear.
Besides biting, another possible transmission pathway is allogrooming, i.e., the licking
and chewing of a conspecific's fur and skin [22]. Allogrooming takes up about 3-5% of a bat's
active time [23]. It is sometimes targeted to wounds on the skin and can reopen minor wounds
(GGC, personal observation) creating potential for transfer of saliva into open wounds.
Allogrooming of the face and mouth is sometimes followed by regurgitations of ingested blood
(e.g., [24]), which could also lead to RABV transmission [25]. No study has yet quantified
changes in allogrooming in rabid bats.
During a study to evaluate a recombinant rabies vaccine candidate for vampire bats, we
opportunistically measured rates of aggression and allogrooming in 40 captive male vampire
bats that were experimentally infected with RABV. We then compared aggression and
allogrooming in non-rabid bats to bats confirmed positive for rabies at death or the end of study.
82

Material and Methods

84
Capture and care
We collected behavior data from 40 male common vampire bats that were part of a larger
sample of bats used to test a viral-vectored recombinant mosaic glycoprotein rabies vaccine
candidate. The bats were captured in the State of San Luis Potosí, México, July-Aug 2018, and
transported to the U.S. Geological Survey National Wildlife Health Center in Madison,
Wisconsin, USA (for details see 21). Bats were individually marked by combinations of 0-4 bat
bands (Porzana Limited) on the right or left forearm.
92
Experimental procedure
For the rabies vaccine study, bats were caged according to three treatments: (1) oral
vaccination, (2) topical vaccination, or (3) placebo control. Groups remained caged together for
~120 days before being challenged with RABV. One week prior to the challenge, we reassigned
the bats into new cages so individuals that received different treatments would be mixed and
acclimated to each other (3 cages with 13, 13, and 14 bats). All bats were challenged with a
heterologous RABV variant (of coyote origin) at a dose of $10^{3.3}$ tissue culture infective dose
($TCID_{50}$/mL), injected intramuscularly into each masseter muscle (50 μL on each side) in April

2019 (127 days post-vaccination). We began quantifying behaviors one day after the challenge
(day 1 post challenge). To confirm death by rabies, we performed a direct fluorescent antibody
test for RABV in brain smear impressions of bats following standard procedures [26]. To detect
RABV shedding in the saliva, we collected oral swabs periodically from all individuals, daily if
clinical signs were observed, and upon death. Swabs were tested using methods described
elsewhere [27,28].
107
Behavioral data collection
After the bats were challenged with RABV, they were recorded using an infrared surveillance
system (Amcrest 960 H/+) with a different camera pointed to the interior of each cage through a
clear acrylic viewing window. In each of the three cages, we sampled behaviors three hours per
night (at hours 0100, 0300, and 0500) during the most active period [29]. At every new-minute
mark, an observer that was blind to the infection status of the bats, stopped the video and
recorded the presence or absence of either allogrooming or aggression within a 5-second time
window, and the identities of the actor and receiver. Allogrooming involves licking or chewing
another bat's fur or skin and often occurs with two bats allogrooming each other simultaneously
(video S1). Aggressive events included biting and fighting (video S2), and a behavior we call
"clinging" where a bat bites into another bat's neck and clings onto it for a prolonged time period
while the target is actively trying to shake off the aggressor (video S3).
120
Statistical analysis
For each night we collected 180 presence/absence samples per group except for two nights
when 53 samples and 29 samples were missing due to camera outages (resulting in a total of
18,818 behavioral samples). We counted the number of observed allogrooming and aggression
events for each bat and divided it by the three sampled hours to estimate behavioral rates per
hour. We estimated the 95% confidence intervals around the mean rate for each of the two
treatment groups, rabid and non-rabid, using bootstrapping (percentile method, 5000 iterations,
boot R package, [30]). We also plotted the mean difference in behavior rates of rabid bats
versus non-rabid bats over each day of the study.
Because the exact timeline of infection for any rabid bat was unclear, we plotted for
every rabid bat an effect size (standardized mean difference between observed and expected)
during increasingly large (nested) time periods, from 1 day before the rabid bat's death to 15
days before death (excluding one bat that survived until the end of the experiment 50 days after
the challenge). To do this, we calculated the mean behavior count for each focal rabid bat for a
given period prior to death (e.g., 4 days = days 1-4 prior to death), then compared that observed
mean count to the expected count (i.e., the mean count of all non-rabid bats within the same
group and time period). To get an effect size for each time period, we standardized this mean
difference:
139

$$standardized\ mean\ difference = \frac{mean\ count\ of\ rabid\ bat - mean\ count\ of\ non-rabid\ bats}{pooled\ standard\ deviation}$$

140
141
142 **Results**
143
Fourteen bats died after the experimental RABV challenge; all were confirmed RABV positive,
and deaths occurred in all three cages and in all three treatment groups (5 controls, 4 oral
vaccinates, and 5 topically vaccinated). We detected no difference between the vaccination
treatments on behavioral rates in rabid bats (Fig. S5, Table S6). The time of death ranged from
9-29 days post-challenge. One other RABV-challenged bat that had been topically vaccinated
was alive by the end of the experiment (after 50 days) but confirmed rabid after it was

euthanized. None of the 10 vaccinated bats that became rabid shed virus. In the unvaccinated
bats, we detected RABV shedding in the saliva of 3 bats on the day of death, and one of these
was also shedding RABV the day prior to death.
In rabid bats, we did not detect a clear increase in aggression given (Fig. 1) or received
(Fig. S2). Rabid vampire bats both gave and received less allogrooming than their healthy
cagemates on average about 12 days after inoculation, and this group difference increased as
we considered time periods closer to their death (Fig. 2, S3). The decrease in allogrooming and
low levels of aggression are consistent with paralytic rather than 'furious' rabies.
158
159

160
161
Figure 1: No evidence of increased aggression in rabid male vampire bats prior to death. Panel (a)
shows timeline of aggression event counts for each rabid (orange points) and non-rabid (blue point)
vampire bat across three cages. Point size reflects the rate of observed events per hour. Red circles
show RABV positive saliva sample. Panel (b) shows mean rate of aggression events per hour ± 95% CIs
for rabid and non-rabid bats starting one day after inoculation with RABV. Panel (c) shows the
standardized mean difference ± 95% CIs between rabid bats and healthy cagemates during shrinking
time intervals before death. Outliers in panel c are caused by one rabid bat (group 2-i) that showed

aggression 16 days post-challenge (2 days prior to death). See table S4 for confidence intervals.
170

171
172
Figure 2: Reduced allogrooming prior to death in rabid male vampire bats. Panel (a) shows timeline
of allogrooming events counts for each rabid (orange points) and non-rabid (blue point) vampire bat
across three cages. Point size reflects the rate of observed events per hour. Red circles show RABV
positive saliva sample. Panel (b) shows mean rate of allogrooming events per hour ± 95% CIs for rabid
and non-rabid bats starting one day after inoculation with RABV. Panel (c) shows the standardized mean
difference ± 95% CIs between rabid bats and healthy cagemates during shrinking time intervals before
death. See table S4 for confidence intervals.
180
181

Table 1: Studies anecdotally describing rabies-induced changes in social behavior of vampire bats after injection or natural exposure.

Reference	Result	Method	Type of virus used	Comments
Present study	No increase in aggressive behavior observed, reduced social grooming	Behavioral sampling	Coyote variant	40 male vampire bats. Some previously vaccinated. 15 bats rabies confirmed.
[25]	No aggression, grooming unobserved	Anecdotal observation	T-9/95 vampire bat field isolate from before 2001 (no detailed information). Study published in 2009. Isolate location unknown.	10 vampire bats (7 males, 3 females). 4 bats confirmed positive. Signs of paralytic rabies in all 4.
[31]	No aggression, grooming unobserved	Anecdotal observation	CASS-88 vampire bat origin, isolated from infected cow brain in 1988. Study published in 1998.	24 vampire bats (no information on sex). Some bats previously vaccinated with oral vaccine. 16 confirmed positive. Signs of paralytic rabies in 10/16. Others showed no obvious clinical signs.
[20]	No aggression, grooming unobserved	Anecdotal observation	CASS-88 vampire bat origin, isolated from infected cow brain in 1988. Study published in 2005.	14 bats (six males, eight females). 11 died. Paralysis of wings and hind-legs prior to death in 3/11. Others showed depression, hypoactivity and anorexia.
[32]	No aggression, grooming unobserved	Anecdotal observation	CASS-88 vampire bat origin, isolated from infected cow brain in 1988. Study published in 2002.	Test of rabies vaccine. 9/10 control bats (vaccinated with saline) died of rabies (no information on sex). Altered reflexes, tremor, and paralysis were observed 72–24 hours before death in rabid bats.
[33]	No aggression, grooming unobserved	Anecdotal observation	Brldr2918 vampire bat field isolate from 1997. Study published in 2005. Study bat location and location of virus isolate are <100 km apart.	10 bats died of RABV (no detailed information on sex/deaths). 8 showed signs of paralytic rabies, 2 showed no clinical signs
[34]	No aggression, grooming unobserved	Anecdotal observation	Brldr2918 vampire bat field isolate from 1997. Study published in 2008. No information on bat capture location	10 bats died of RABV (no detailed information on sex/deaths) some were previously vaccinated (orally, fur-transmissible). All showed signs of paralytic rabies.
[21]	Aggression, grooming unobserved	Anecdotal observation	Natural exposure	Total of 14 confirmed rabid male bats. 7 showed no clinical signs, 3 the furious form and 4 the paralytic form.
[35]	Potential aggression, grooming unobserved	Anecdotal observation	Natural exposure	Introduction of wild bats to an existing captive colony (no information on sex). After 2 months, fighting started. Several bats from original colony were mutilated and tested for rabies. Suggested aggression of introduced bats.
[36]	Aggression, grooming unobserved	Anecdotal observation	Natural exposure	Aggression observed in naturally infected rabid bats. 14 of 24 bats observed showed clinical signs including hyperexcitability, aggressiveness, and paralysis before death.

Discussion

Rabid male vampire bats reduced their allogrooming, and probably as a consequence, also received less allogrooming. This change could be due to either rabies-induced paralysis or generalized sickness behaviors, such as lethargy, causing passive self-isolation [37–40]. All bats showed low rates of aggression, and we saw no clear increase in aggression in the rabid bats, regardless of vaccine treatment group.

Several other studies did not observe heightened aggression in rabid vampire bats (Table 1). One possible reason for this lack of observations is reduced selection on RABV to increase aggression in vampire bats because they are highly social, frequently aggressive, and bite other hosts [41]. Another possibility is that distinct RABV strains differ in pathogenicity and clinical forms of the disease (e.g., presence of aggression) across species [42–45]. Studies describing natural infections often report some aggression, but experimental RABV challenges that failed to find evidence of aggression used viral strains that were not currently circulating or, as in our case, used a strain derived from a different species (Table 1). It would be interesting to determine if infection with endemic vampire bat RABV strains may induce a higher proportion of furious versus paralytic disease in vampire bats.

Similar to some field observations of rabid vampire bats [17,36], some rabid bats in our study seemed disproportionally targeted, receiving more aggression prior to death (Fig. S2, e.g., bat group 1-j, 2-i, 2-d, 3-b). Acute infections can have a social cost, such as increased antagonism by conspecifics [46], and it would be interesting to examine evidence for avoidance of or aggression towards rabid individuals more closely in this and other species [40].

Times until death in rabid vampire bats varied from 9 to 29 days, but one of the 15 rabid males remained alive until the end of the experiment, 50 days after infection. The bat was previously vaccinated, but its neurologic function declined over the final weeks losing coordination and mobility. We did not detect RABV in its saliva. The causes of this prolonged survival remain unclear.

In the late stage of infection, RABV spreads to the salivary glands and is excreted in saliva [42]. Evidence of RABV shedding in vampire bats prior to or at the time of death has been demonstrated before [20,33,47]. Here, we detected RABV shedding in saliva of 3 not vaccinated bats of the 15 rabies-infected bats, which allowed us to overlay behavioral measures with pathogen shedding (Fig. 1, 2, S2, S3). These 3 vampire bats did not groom others much when the virus was detectable in their saliva. Similarly, we did not observe heightened social aggression in these periods before death. Future work should more closely quantify the relationship between rabies shedding and behavioral changes.

Some caveats to the interpretations of our experimental results and those of others (Table 1) should be considered. First, given that social aggression can be rare and brief, absence of evidence of social aggression is not evidence of absence. Occasional aggression towards handlers and other bats by rabid bats was observed during daily checkups. Second, the administered RABV challenge dose, route, and site of inoculation are not standardized across experiments and often differ (Table 1). As in our study, the RABV challenge strains used by researchers do not typically represent currently endemic viruses, are derived from different species, or are adapted to other species before their use in experimental infections. More standardized experimental infections are needed to disentangle the role of administered dose and temporal overlap of circulating strains on rabies-induced behavioral changes in natural reservoirs such as vampire bats.

In conclusion, we observed reductions in allogrooming and low levels of aggression that indicated paralytic but not furious rabies presentation in 15 rabid male vampire bats relative to 25 non-rabid male bats. In comparison to other previous reports involving natural rabies exposures that report elevated aggression (Table 1), our results suggest variation in behavioral effects by RABV strain.

Ethics: Field work was carried out under permit *SGPA/DGVS/003242/18* from the Mexican Secretariat of Environment and Natural Resources. All animal husbandry practices and experiments were approved by the USGS-National Wildlife Health Center Institutional Animal Care and Use Committee (*Protocol EP180418*).

Data Availability: All data and R code to repeat the analysis is publicly available on Figshare: https://doi.org/10.6084/m9.figshare.19991204.v2

Competing interests: We have no competing interests.

Acknowledgements: We thank Osric Wong, Natalie Sebunia, Jaelynn Butler, Michael Shechtman, Olivia Kalczynski, and Raven Hartman for scoring behavioral videos. Daniel Streicker, Horacio Delpietro, Alvaro Aguilar-Setién, and Charles Rupprecht provided relevant literature.

Funding: SS and GGC are supported by the National Science Foundation (grant IOS-2015928).

Authors contributions: EMCC, SS, and GGC designed the study. EMCC carried out experiments. SS and EC analyzed the data. TR and JEO coordinated the study and provided resources and lab space. GGC conceived of the idea and supervised the data analysis and writing. All authors contributed to draft of manuscript and gave final approval for publication.

References

1. Benavides JA, Rojas Paniagua E, Hampson K, Valderrama W, Streicker DG. 2017 Quantifying the burden of vampire bat rabies in Peruvian livestock. *PLOS Negl Trop Dis* **11**, e0006105. (doi:10.1371/journal.pntd.0006105)

2. Johnson N, Arechiga-Ceballos N, Aguilar-Setien A. 2014 Vampire bat rabies: ecology, epidemiology and control. *Viruses* **6**, 1911–28. (doi:10.3390/v6051911)

3. Martínez-Burnes J, López A, Medellín J, Haines D, Loza E, Martínez M. 1997 An outbreak of vampire bat-transmitted rabies in cattle in northeastern Mexico. *Can Vet J* **38**, 175–177.

4. Lopez A, Miranda P, Tejada E, Fishbein DB. 1992 Outbreak of human rabies in the Peruvian jungle. *Lancet* **339**, 408–411. (doi:10.1016/0140-6736(92)90088-k)

5. Schneider MC, Romijn PC, Uieda W, Tamayo H, da Silva DF, Belotto A, da Silva JB, Leanes LF. 2009 Rabies transmitted by vampire bats to humans: an emerging zoonotic disease in Latin America? *Rev Panam Salud Publica* **25**, 260–269. (doi:10.1590/s1020-49892009000300010)

6. Streicker DG *et al.* 2012 Ecological and anthropogenic drivers of rabies exposure in vampire bats: implications for transmission and control. *Proc Biol Sci* **279**, 3384–92. (doi:10.1098/rspb.2012.0538)

7. Blackwood JC, Streicker DG, Altizer S, Rohani P. 2013 Resolving the roles of immunity, pathogenesis, and immigration for rabies persistence in vampire bats. *Proc Natl Acad Sci USA* **110**, 20837–20842. (doi:10.1073/pnas.1308817110)

8.   Streicker DG *et al.* 2016 Host–pathogen evolutionary signatures reveal dynamics and future
invasions of vampire bat rabies. *Proc Natl Acad Sci USA* **113**, 10926–10931.
(doi:10.1073/pnas.1606587113)

9.   Streicker DG, Allgeier JE. 2016 Foraging choices of vampire bats in diverse landscapes:
potential implications for land-use change and disease transmission. *J Appl Ecol* **53**, 1280–
1288. (doi:10.1111/1365-2664.12690)

10. Moegle H, Knorpp F. 1978 Zur Epidemiologie der Wildtiertollwut: 2. Mitteilung:
Beobachtungen über den Dachs 1. *Zentralblatt für Veterinärmedizin Reihe B* **25**, 406–415.

11. Smith GC. 2002 The role of the badger (Meles meles) in rabies epizootiology and the
implications for Great Britain. *Mammal Rev* **32**, 12–25.

12. George JP. 1980 Description clinique de la rage du renard.

13. Shuangshoti S, Thorner PS, Teerapakpinyo C, Thepa N, Phukpattaranont P, Intarut N,
Lumlertdacha B, Tepsumethanon V, Hemachudha T. 2016 Intracellular spread of rabies
virus is reduced in the paralytic form of canine rabies compared to the furious form. *PLOS*
Negl Trop Dis **10**, e0004748. (doi:10.1371/journal.pntd.0004748)

14. Winkler WG, Schneider NJ, Jennings WL. 1972 Experimental rabies infection in wild
rodents. *J Wildl Dis* **8**, 99–103.

15. Hemachudha T, Wacharapluesadee S, Mitrabhakdi E, Wilde H, Morimoto K, Lewis RA.
2005 Pathophysiology of human paralytic rabies. *J Neurovirol* **11**, 93–100.
(doi:10.1080/13550280590900409)

16. Klein SL. 2003 Parasite manipulation of the proximate mechanisms that mediate social
behavior in vertebrates. *Physiol Behav* **79**, 441–449. (doi:10.1016/S0031-9384(03)00163-X)

17. Delpietro HA, Russo RG, Carter GG, Lord RD, Delpietro GL. 2017 Reproductive
seasonality, sex ratio and philopatry in Argentina's common vampire bats. *R Soc Open Sci*
4, 160959.

18. Wilkinson GS. 1985 The social organization of the common vampire bat. I. Pattern and
cause of association. *Behav Ecol Sociobiol* **17**, 111–121.

19. Wilkinson GS. 1985 The social organization of the common vampire bat: II. Mating system,
genetic structure, and relatedness. *Behav Ecol Sociobiol*, 123–134.

20. Aguilar-Setien A, Loza-Rubio E, Salas-Rojas M, Brisseau N, Cliquet F, Pastoret PP, Rojas-
Dotor S, Tesoro E, Kretschmer R. 2005 Salivary excretion of rabies virus by healthy vampire
bats. *Epidemiol infect* **133**, 517–522. (doi:10.1017/s0950268805003705)

21. Cárdenas-Canales EM *et al.* 2020 Clinical presentation and serologic response during a
rabies epizootic in captive common vampire bats (*Desmodus rotundus*). *Trop Med Infect Dis*
5. (doi:10.3390/tropicalmed5010034)

22. Wilkinson GS. 1986 Social grooming in the common vampire bat, *Desmodus rotundus.*
Anim Behav **34**, 1880–1889. (doi:http://dx.doi.org/10.1016/S0003-3472(86)80274-3)

23. Carter G, Leffer L. 2015 Social grooming in bats: are vampire bats exceptional? *PLoS One*
10, e0138430. (doi:10.1371/journal.pone.0138430)

24. Carter GG, Wilkinson GS. 2015 Social benefits of non-kin food sharing by female vampire
bats. *Proc Biol Sci* **282**. (doi:10.1098/rspb.2015.2524)

25. Agronegócios P, Aparecida MC, Souza M, Nassar C, Cortez A, Sakai T, Itou T, Sequetin
EM, José L. 2009 Experimental infection of vampire bats Desmodus rotundus (E. Geoffroy)
maintained in captivity by feeding defibrinated blood added with rabies virus. *Braz J Vet Res*
Anim Sci **46**, 92–100.

26. Rupprecht CE, Fooks AR, Abela-Ridder B. 2018 *Laboratory techniques in rabies*. World
Health Organization.

27. Gigante CM *et al.* 2018 Multi-site evaluation of the LN34 pan-lyssavirus real-time RT-PCR
assay for post-mortem rabies diagnostics. *PLoS One* **13**, e0197074.
(doi:10.1371/journal.pone.0197074)

28. Wadhwa A *et al.* 2017 A Pan-Lyssavirus Taqman real-time RT-PCR assay for the detection
of highly variable rabies virus and other lyssaviruses. *PLoS Negl Trop Dis* **11**, e0005258.
(doi:10.1371/journal.pntd.0005258)

29. Ripperger SP, Carter GG. 2021 Social foraging in vampire bats is predicted by long-term
cooperative relationships. *PLOS Biol* **19**, e3001366. (doi:10.1371/journal.pbio.3001366)

30. Canty A, Ripley B. 2017 boot: Bootstrap R (S-Plus) functions. *R package version* **1**, 3–20.

31. Sétien AA, Brochier B, Tordo N, De Paz O, Desmettre P, Péharpré D, Pastoret P-P. 1998
Experimental rabies infection and oral vaccination in vampire bats (*Desmodus rotundus*).
Vaccine **16**, 1122–1126.

32. Aguilar-Setién A, Leon YC, Tesoro EC, Kretschmer R, Brochier B, Pastoret P-P. 2002
Vaccination of vampire bats using recombinant vaccinia-rabies virus. *J Wildl Dis* **38**, 539–
544. (doi:10.7589/0090-3558-38.3.539)

33. Almeida MF, Martorelli LFA, Aires CC, Sallum PC, Durigon EL, Massad E. 2005
Experimental rabies infection in haematophagous bats *Desmodus rotundus*. *Epidemiol*
Infect **133**, 523–527. (doi:10.1017/s0950268804003656)

34. Almeida MF, Martorelli LFA, Aires CC, Barros RF, Massad E. 2008 Vaccinating the vampire
bat *Desmodus rotundus* against rabies. *Virus Res* **137**, 275–277.
(doi:10.1016/j.virusres.2008.07.024)

35. Horst R, Langworthy M. 1972 Rabies in a colony of vampire bats. *J Mammal* **53**, 903–905.

36. Delpietro HA, de Díaz AMO, Larghi OP. 1985 Comportamiento en cautividad de vampiros
rabiosos infectados naturalmente. *Veterinaria Argentina* **2**, 748–756.

37. Ripperger SP, Stockmaier S, Carter GG. 2020 Tracking sickness effects on social
encounters via continuous proximity sensing in wild vampire bats. *Behav Ecol* 31, 1296-
1302. (doi:10.1093/beheco/araa111)

38. Stockmaier S, Bolnick DI, Page RA, Carter GG. 2020 Sickness effects on social interactions
depend on the type of behaviour and relationship. *J Anim Ecol* 89, 1387-1394.
(doi:10.1111/1365-2656.13193)

39. Stockmaier S, Bolnick DI, Page RA, Carter GG. 2018 An immune challenge reduces social
grooming in vampire bats. *Anim Behav* **140**, 141–149. (doi:10.1016/j.anbehav.2018.04.021)

40. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Ancel Meyers L, Bolnick DI. 2021
Infectious diseases and social distancing in nature. *Science* **371**, eabc8881.
(doi:10.1126/science.abc8881)

41. Hart BL. 1990 Behavioral adaptations to pathogens and parasites: five strategies. *Neurosci*
Biobehav Rev **14**, 273–294.

42. Banyard AC, Davis A, Gilbert AT, Markotter W. 2020 Bat rabies. In Rabies Elsevier, 231–
276. (doi:10.1016/B978-0-12-818705-0.00007-8)

43. Davis AD, Gordy PA, Bowen RA. 2013 Unique characteristics of bat rabies viruses in big
brown bats (*Eptesicus fuscus*). *Arch Virol* **158**, 809–820.

44. Obregón-Morales C, Aguilar-Setién Á, Martínez LP, Galvez-Romero G, Martínez-Martínez
FO, Aréchiga-Ceballos N. 2017 Experimental infection of *Artibeus intermedius* with a
vampire bat rabies virus. *Comp Immunol Microbiol Infect Dis* **52**, 43–47.

45. Morimoto K, Hooper DC, Carbaugh H, Fu ZF, Koprowski H, Dietzschold B. 1998 Rabies
virus quasispecies: implications for pathogenesis. *Proc Natl Acad Sci USA* **95**, 3152–3156.

46. McFarland R, Henzi SP, Barrett L, Bonnell T, Fuller A, Young C, Hetem RS. 2021 Fevers
and the social costs of acute infection in wild vervet monkeys. *Proc Natl Acad Sci USA* **118**.
(doi: 10.1073/pnas.2107881118)

47. Moreno JA, Baer GM. 1980 Experimental rabies in the vampire bat. *Am J Trop Med Hyg* **29**,
254–259.