A class IV adenylate cyclase CyaB is required for capsule polysaccharide production and biofilm formation in *Vibrio parahaemolyticus*

Regmi, A., J.G. Tague, K. Boas Lichty, and E.F. Boyd*

Department of Biological Sciences, University of Delaware, Newark, DE 19716

Corresponding author*

E. Fidelma Boyd

Department of Biological Sciences

341 Wolf Hall, University of Delaware

Newark, DE 19716

Phone: (302) 831-1088. Fax: (302) 831-2281 Email: fboyd@udel.edu

ABSTRACT

CRP (cyclic AMP receptor protein), encoded by *crp*, is a global regulator that is

activated by cAMP (cyclic AMP), a second messenger synthesized by a class I adenylate

cyclase (AC-I) encoded by *cyaA* in *Escherichia coli*. cAMP-CRP is required for growth on

non-preferred carbon sources and is a global regulator. We constructed in-frame non-

polar deletions of the *crp* and *cyaA* homologs in *Vibrio parahaemolyticus* and found that

the Δ*crp* mutant did not grow in minimal media supplemented with non-preferred

carbon sources, but the Δ*cyaA* mutant grew similar to wild type. Bioinformatics analysis

of the *V. parahaemolyticus* genome identified a 181 amino acid protein annotated as a

class IV adenylate cyclase (AC-IV) named CyaB, a member of the CYTH protein

superfamily. AC-IV phylogeny showed CyaB was present in Gamma- and Alpha-

Proteobacteria as well as Planctomycete and Archaea. Only the bacterial CyaB proteins

contained an N-terminal motif HFxxxExExK indicative of adenylyl cyclase activity.

Both *V. parahaemolyticus cyaA* and *cyaB* genes functionally complemented an *E. coli*

Δ*cyaA* mutant. The Δ*crp* and Δ*cyaB*/Δ*cyaA* mutants showed defects in growth on non-

preferred carbon sources, and in swimming and swarming motility, indicating cAMP-

CRP is an activator. The Δ*cyaA* and Δ*cyaB* single mutants had no defects in these

phenotypes indicating AC-IV complements AC-I. Capsule polysaccharide and biofilm

production assays showed significant defects in Δ*crp*, Δ*cyaB*/Δ*cyaA*, and the Δ*cyaB*

mutant, whereas Δ*cyaA* behaved similar to wild type. This is consistent with a role of

cAMP-CRP as an activator of these phenotypes and establishes a cellular role for AC-IV

in capsule and biofilm formation, which to date has been unestablished.

IMPORTANCE

Here, we characterized the roles of CRP and CyaA in *V. parahaemolyticus,* showing

cAMP-CRP was an activator of metabolism, motility, capsule and biofilm formation.

These results are in contrast to cAMP-CRP in *V. cholerae,* which represses capsule and

biofilm formation. Previously, only an AC-I CyaA had been identified in *Vibrio* species.

Our data showed that an AC-IV CyaB homolog is present in *V. parahaemolyticus* and

was required for optimal growth. The data demonstrated that CyaB was essential for

capsule production and biofilm formation uncovering a physiological role of AC-IV in

bacteria. The data showed that the *cyaB* gene was widespread among *Vibrionaceae*

species and several other Gamma-Proteobacteria, but in general, its phylogenetic

distribution was limited. Our phylogenetic analysis also demonstrated that in some

species the *cyaB* gene was acquired by horizontal gene transfer.

INTRODUCTION

Carbon catabolite repression (CCR) is a global regulatory mechanism that represses

functions required for secondary carbon metabolism when a preferred carbon source is

present (1, 2). In *Escherichia coli*, CCR is mediated by several mechanisms such as the

regulation of the biosynthesis of catabolic enzymes, the inhibition of carbon transport,

and metabolic signaling (3, 4). Some of the major players involved in CCR are the global

regulator CRP (**c**AMP **R**eceptor **P**rotein), adenylate cyclase (AC), the signaling molecule

cAMP (3′,5′-cyclic AMP), and components of phosphoenolpyruvate:sugar

phosphotransferase system (PTS) transporters (1, 3-7). CRP functions as a homodimer,

activated by the second messenger cAMP and in *E. coli* cAMP-CRP is a global

transcriptional activator of non-preferred carbon genes in the absence of glucose (8-10).

In *E. coli*, CRP has an N-terminal cAMP binding domain (CBD; residues 1–136) and a C-

terminal helix turn helix DNA binding domain (DBD; residues 139–209) (11, 12). When

bound by cAMP, CRP undergoes a conformation change that allows the protein to

interact with target promoter regions at consensus sequences (13, 14).

When glucose is present, it is imported by the PTS transporter, resulting in

dephosphorylation of the EIIA component of PTS. EIIA^{glc} inhibits non-preferred carbon

transporters, a mechanism known as inducer exclusion, independent of cAMP signaling

(2, 4, 15-17). When glucose is absent, EIIA^{glc} is phosphorylated and P~EIIA^{glc} activates

adenylate cyclase to produce cAMP (4, 18). However, growth on non-PTS carbon

sources, such as lactose, and in limited nitrogen and phosphorous conditions also have

reduced cAMP levels, indicating the presence of a catabolite modulating factor, which

Monod suggested in his last paper (19). Recent research showed that cAMP signals

were modulated by metabolic precursors such as α-ketoacids to control the allocation of

proteomic resources by a feedback system rather than carbon metabolism (20, 21).

There are six classes (I-VI) of adenylate cyclases (AC) based on primary amino

acid sequence, all with different catalytic mechanisms and structures, indicating

different evolutionary origins (22, 23). *Escherichia coli* and many other *Proteobacteria*

contain a single AC gene *cyaA*, which encodes a class I AC (AC-I). The 810 amino acid

CyaA protein has two functionally distinct domains, the N-terminal catalytic cyclase

domain and the C-terminal regulatory domain (11, 12, 24). Class II ACs (AC-II) are

comprised of bacterial toxins such as the exotoxin from *Bordetella pertussis* (25). The class

III ACs (AC-III) are the most diverse and widespread, and encompass adenylate

cyclases present in eukaryotes and prokaryotes (26). A structurally diverse AC was

identified in *Prevotella ruminicola* D31d and forms the sole representative of AC-V (27).

A class VI AC was described in *Rhizobium elti*, a 345 amino acid protein that was shown

to have cyclase activity by complementation of an *E. coli cyaA* deletion mutant (28).

Class IV AC (AC-IV) along with mammalian thiamine triphosphatases and

inorganic triphosphatases belong to the CYTH-like domain superfamily (29). CYTH

proteins are present in all three domains, Bacteria, Archaea and Eukaryota, and perform

a variety of functions (29). CYTH proteins contain a triphosphate tunnel metaloenzyme

(TTM) fold and act on triphosphorylated substrates with at least one divalent metal

cation required for catalysis. A hallmark of CYTH proteins is an N-terminal ExExK

motif (30). In bacteria, an AC-IV CyaB was initially described in *Aeromonas hydrophila*, a

small 181 amino acid protein with a single domain (31). The foundational work in *A.*

hydrophila indicated that *cyaB* was not expressed under standard growth conditions, but

cyaB did functionally complement an *E. coli cyaA* deletion mutant (31). A CyaB homolog

was also identified from *Yersinia pestis*, its crystal structure (2FJT) was determined and

showed to be very different from AC-I and AC-III enzymes. AC-IV formed a dimer like

AC-III, but with a distinct antiparallel eight-stranded barrel active site and could cyclize

both ATP and GTP (32-34). The physiological role of bacterial AC-IV was not

established. Archaea contain numerous proteins annotated as AC-IV, but cAMP

biosynthesis has not been shown for this domain (35). A recent study examining a

CYTH-like protein annotated as an AC-IV from the Crenarchaeote *Sulfolobus*

acidocaldarius showed that it had triphosphatase activity, but no adenylate cyclase

activity (35). Vogt and colleagues differentiated CYTH-like proteins (with phosphatase

activity) from AC-IV (with adenylyl cyclase activity) using sequence similarity network

analysis and functional studies (35). They suggested that a true AC-IV has an extended

N-terminal motif HFxxxxExExK and the absence of HF indicates a CYTH protein with

either thiamine triphosphatase (ThTPase) or inorganic triphosphatase activity (35).

In the present study, the function and activation of CRP in *V. parahaemolyticus*

was investigated for the first time. In-frame non-polar deletions of *crp* (VP2793) and

cyaA (VP2987) in *V. parahaemolyticus* RIMD2201633 were constructed and examined for

growth on various carbon sources. The Δ*crp* mutant grew on minimal medium M9 3%

NaCl supplemented with glucose (M9G) and was unable to grow on non-PTS carbon

sources, but the Δ*cyaA* mutant grew similar to wild type. This suggested the presence of

an additional uncharacterized AC in *V. parahaemolyticus*. Bioinformatics analysis

identified ORF VP1760 that encoded a putative AC-IV named CyaB. The phylogeny of

AC-IV among Bacteria and their closest orthologs in Archaea was examined to

determine its distribution and evolutionary history. To examine whether *cyaA* and *cyaB*

were both functional, an *E. coli* Δ*cyaA* mutant was functionally complemented with both

genes. Growth pattern analysis of the Δ*cyaB*, Δ*cyaA* and a double Δ*cyaB*/Δ*cyaA* mutants

on different carbon sources was also examined. Swimming and swarming motility

assays, capsule polysaccharide (CPS) production and biofilm formation assays were

performed to determine whether cAMP-CRP played a role in these important

phenotypes.

RESULTS

CRP is an essential regulator in *V. parahaemolyticus*.

cAMP-CRP is a global regulator of nutrient transport and metabolism in many bacterial

species (13, 36-40). Homologs of CRP and CyaA are present in *V. parahaemolyticus*, ORF

VP2793 and VP2987, respectively (**Fig. S1**). The CRP and CyaA proteins of *V.*

parahaemolyticus shared 98% and 82% amino acid identity respectively to homologs in *V.*

cholerae. The genome context of both genes is conserved across *V. parahaemolyticus*, *V.*

cholerae, and *E. coli* (**Fig. S1**). To begin to investigate the physiological role of CRP in *V.*

parahaemolyticus, we first constructed an in-frame non-polar deletion of *crp* (VP2793)

and *cyaA* (VP2987) in *V. parahaemolyticus* RIMD2210633. The Δ*crp* mutant displayed a

small colony morphology on LB supplemented with 3% NaCl (LBS) plates **(Fig. S2A)**.

Growth pattern analysis in LBS broth and M9S D-glucose broth demonstrated that the

Δ*crp* strain had a growth defect **(Fig. 1A**-**B**). Next, we examined the Δ*cyaA* mutant,

expecting a phenotype similar to the Δ*crp* mutant, as shown in *E. coli* and *V. cholerae*

Δ*cyaA* mutants. However, growth of Δ*cyaA* on LBS plates gave identical colony

morphology to wild type (**Fig. S2A)** and similarly in LBS and M9 supplemented with

3% NaCl (M9S) and D-glucose broth, Δ*cyaA* presented a similar pattern to wild type

(**Fig. 1A**-**B**). In addition, growth in alternative carbon sources, M9S supplemented with

L-arabinose, L- ribose, or D-gluconate, showed the Δ*crp* strain was defective, but the

Δ*cyaA* mutant grew similar to wild type (**Fig 2A-C**). These data demonstrate that CRP is

required for catabolism of alternative carbon sources similar to *V. cholerae*, but that *V.*

parahaemolyticus contained an additional uncharacterized adenylate cyclase that can

synthesize cAMP in the absence of *cyaA* to activate CRP. Alternatively, CRP could be

acting independently of cAMP, but given its high homology to CRP from *V. cholerae* this

is less likely.

Identification of a class IV adenylate cyclase (AC-IV) CyaB in *V. parahaemolyticus*.

BLASTP analysis using CyaA (VP2987) as a seed to search the *V. parahaemolyticus*

RIMD2210633 genome, only hit to itself, indicating no other homolog of AC-I was

present. Then, a search of the *V. parahaemolyticus* genome for adenylate cyclase classes

AC-II using locus WP_080378568 from *Bordetella pertussis*, AC-III using locus PWU33926

from *Pseudomonas aeruginosa*, and AC-IV using locus UNB56521 from *A. hydrophila,* was

performed. In addition, ten AC proteins (loci Rv0386, Rv1264, Rv1318c, Rv1319c,

Rv1320c, Rv1625c, Rv1647, Rv1900c, Rv2212, Rv3645) encompassing AC-II and AC-III

from *Mycobacterium tuberculosis* that were shown to have activity were also used as

seeds. These searches identified a homolog of AC-IV from *A. hydrophila* named CyaB.

No additional adenylate cyclases were identified in *V. parahaemolyticus* RIMD2210633.

The 181 amino acid CyaB protein lacked transmembrane domains and signal peptide

cleavage sites. CyaB contained only one domain and unlike AC-I, AC-II and AC-III does

not contain any regulatory or signaling domains. CyaB shared 56% amino acid identity

with CyaB of *A. hydrophila* and 45% amino acid identity with CyaB from *Y. pestis*. CyaB

from both *A. hydrophila* and *Y. pestis* were shown to have AC activity generating cAMP,

however a physiological role was not established (31-34). In *V. parahaemolyticus, cyaB* is

encoded on the negative strand upstream of VP1761, which encoded a DUF962-domain

178 hypothetical protein separate by an intergenic region of 88-bp. The AC-IV CyaB protein

179 has been structurally characterized in *Y. pestis* (2FJT and 3NOY) and a crystal structure

180 is also available for VP1760 from *V. parahaemolyticus* (2ACA) (32-34). Comparison of *Y.*

181 *pestis* and *V. parahaemolyticus* primary sequence and crystal structures shows they are

182 highly similar with active sites conserved between the proteins (**Fig. S3**). The primary

183 sequence of all three bacterial sequences contained the N-terminal conserved

184 HFxxxxExExK motif and the archaea sequences contained the shorter ExExK motif (**Fig.**

185 **S3**).

186 **Distribution and phylogenetic analysis of AC-IV among Bacteria**.

187 Using *cyaB* (VP1760) as a seed using BLAST analysis, we examined the distribution

188 among nearly 18,000 members of the family *Vibrionaceae* in the NCBI genome database.

189 The *cyaB* gene was present in all characterized *V. parahaemolyticus* strains and in many

190 members of the Harveyi clade, to which *V. parahaemolyticus* belongs, including *V.*

191 *campbellii, V. harveyi, V. jasicida, V. owensii,* and *V. rotiferanus* (**Fig. 3**). The CyaB proteins

192 in these species showed ~70% amino acid identity with CyaB from *V. parahaemolyticus*,

193 as compared to ~95% amino acid identity among the CyaA proteins. On the *cyaB* gene

194 tree, within the Harveyi clade, species formed two divergent branches; one branch

195 representing *V. parahaemolyticus* strains and the second branch containing the remaining

196 species. This suggests that *cyaB* in *V. parahaemolyticus* has diverged significantly from

197 *cyaB* in the other Harveyi clade species. Interestingly, CyaB was absent from most *V.*

alginolyticus strains, a species closely related to *V. parahaemolyticus*. Our analysis did

identify a *V. alginolyticus* strain that contained a *cyaB* gene on a plasmid that clustered

within the same group as *cyaB* present on a plasmid in *V. coralliilyticus* indicating that

cyaB was acquired by horizontal gene transfer in these species (**Fig. 3**). The *cyaB* gene

was present in ~35 *V. cholerae* strains (from a total of 1,700 genomes in the NCBI

database) and two strains of its sister species *V. metoecus* from a total of 30 genomes,

which clustered together on the tree. The *cyaB* gene from species within the genera

Aliivibrio and most *Photobacterium* formed separate divergent branches from *cyaB* in

Vibrio species (**Fig. 3**). However, *cyaB* from five *Photobacterium* species clustered within

Vibrio clades suggesting that these *Photobacterium* species acquired a copy of the *cyaB*

gene from *Vibrio*, indicating horizontal gene transfer (**Fig. 3**). Comparison of CyaA

phylogeny with the CyaB phylogeny from the same set of species also demonstrated

additional instances of horizontal transfer among species shown by the non-congruency

of the branching patterns for several CyaB proteins (**Fig. S4**).

Next, we examined the distribution and phylogeny of AC-IV among Bacteria and

Archaea genomes in the NCBI genome database (**Fig. 4**). This analysis showed that AC-

IV were present predominantly among Gamma-Proteobacteria. CyaB from some species

belonging to the genera *Shewanella, Marinomonas, Psychromonas,* and *Morganella*

clustered closely with CyaB from *Vibrio* species. Species belonging to the genera

Aeromonas, Yersinia, Brennia, Dickeya, Hafnia, Proteus Providencia, and *Pectobacterium*

contained a copy of CyaB that branched along phylogeny lines indicating that CyaB is

ancestral to these lineages (**Fig. 4**). A handful of Alpha-Proteobacteria, *Cohaesibacter*,

Hoeflea, Kangsaoukella, Maritalea, and Pseudovibrio, also contained a CyaB homolog that

was annotated as AC-IV and these clustered together and branched distantly from

CyaB in Gamma-Proteobacteria (**Fig. 4**). Proteins showing greater than 25% amino acid

identity with greater than 90% query coverage to CyaB from *V. parahaemolyticus* were

present in Archaea and branched divergently from AC-IV homologs in Bacteria (**Fig. 4**).

Amino acid sequence comparisons followed by WebLogo analysis of Proteobacteria or

Archaea CyaB sequences was performed and generated sequence logos for the entire

protein in both groups to identify sequence conservation patterns (**Fig. S5 and S6**). In

this analysis, the previously identified AC-IV defining N-terminal motif HFxxxExExK

was present in all Gamma-Proteobacteria and Alpha-Proteobacteria sequences

examined in this study (**Fig. S5**). In contrast, CyaB representatives from Planctomycete

and Archaea contained the N-terminal motif ExExK, which was proposed to be present

in CYTH proteins with only phosphatase activity (**Fig. S6**) (35).

AC-IV CyaB functionally complements an *E. coli cyaA* deletion mutant.

To begin to determine whether CyaB has adenylate cyclase activity, we functionally

complemented an *E. coli* ∆*cyaA* mutant with the *cyaA* or *cyaB* gene from *V.*

parahaemolyticus. *E. coli* ∆*cyaA*, *E. coli* ∆*cyaA* pBBR*cyaB,* and *E. coli* ∆*cyaA* pBBR*cyaA* were

grown in M9S supplemented with glucose as a sole carbon source and all strains grew

similarly (**Fig. 5A**). Growth was examined in M9S supplemented with gluconate as a

sole carbon source. In this analysis, the growth defect of *E. coli* Δ*cyaA* was rescued in *E.*

coli Δ*cyaA* complemented with either pBBR*cyaA* or pBBR*cyaB* (**Fig. 5B**).

To investigate the role of *cyaB* in *V. parahaemolyticus* physiology further, we

constructed a *cyaB* deletion mutant strain. We examined growth of wild type and the

Δ*cyaB* mutant on solid and in liquid media. The Δ*cyaB* mutant had similar colony

morphology to wild type on LBS plates (**Fig. S2A**) and showed identical growth

patterns in LBS broth to wild type (**Fig. 1A**). However, when the Δ*cyaB* mutant was

grown overnight in M9S D-glucose and used to inoculate M9 supplemented with D-

glucose, a longer lag phase than wild type was observed, but a similar biomass was

reached (**Fig. 1B**).

Next, we constructed a Δ*cyaB*/Δ*cyaA* double mutant strain by deleting *cyaA* from the

Δ*cyaB* background. The double mutant demonstrated small translucent colony

morphology on LBS similar to the colonies observed for Δ*crp* (**Fig. S2A and C**).

Addition of cAMP to the media restored the large colony morphology for the double

mutant indicating that the defect was due to the absence of cAMP (**Fig. S2D**).

Interestingly, a glycerol stock of the double mutant when streaked on LBS plates

produced small translucent colonies as well as regular opaque colonies, suggesting

suppressor mutations were occurring (**Fig. S2B**). We did not observed this phenotype in

either of the single Δ*cyaA or* Δ*cyaB* mutants. In *V. cholerae*, which contains only an AC-I

CyaA, the ΔcyaA mutant reversion to wild-type phenotypes was also observed (41). For

our studies, small colonies were always used for phenotypic assays. The ΔcyaB/ΔcyaA

double mutant showed a growth defect on M9S D-glucose (**Fig. 1B**) and showed no

growth in M9 supplemented with L-arabinose, L-ribose, or D-gluconate **(Fig. 2A-C)**.

These growth assays were performed in M9S L-arabinose, L-ribose, or D-gluconate with

inoculum obtained after overnight growth in M9S D-glucose. These growth assays

showed that the ΔcyaA mutant grew similar to wild type, but the ΔcyaB mutant growth

curve showed a different slope (**Fig. 2**). In contrast, when mutants were grown

overnight in LBS and then inoculated into alternative carbon sources, ΔcyaB growth

curves showed a shorter lag phase and a significantly greater biomass than ΔcyaA,

which grew similar to wild type (**Fig. S7**). Analysis of the regulatory regions of both AC

genes identified putative CRP binding sites in the regulatory region of *cyaA*, but not in

the *cyaB* regulatory region (**Fig. S8**). These data suggest that *cyaA* is under CRP control

and that *cyaB* is not, and that CyaA and CyaB can complement each other.

cAMP-CRP regulation of swimming and swarming motility.

Unlike *V. cholerae* which only produces a polar flagellum, *V. parahaemolyticus* produces

both polar and lateral flagella required for swimming and swarming motility,

respectively, which are both activated by the sigma factor RpoN (42). To investigate the

role of CRP in *V. parahaemolyticus* motility, we conducted swimming assays and found

that the ΔcyaA and ΔcyaB mutants grew similar to wild type (**Fig. 6A**). The Δcrp and

Δ*cyaB*/Δ*cyaA* double mutants were defective in swimming compared to wild type, but

did not show a complete defect similar to the Δ*rpoN* mutant (**Fig. 6A**). Also, Δ*crp* and

Δ*cyaB*/Δ*cyaA* mutant showed a swarming defect as indicated by the absence of the

budding phenotype (**Fig. 6B**). The Δ*cyaA* and Δ*cyaB* mutants grew similar to wild type

producing typical budding swarming cells as previously shown for this species (**Fig.**

6B) (43-45). The WT, Δ*cyaA* and Δ*cyaB* swarming colonies were examined further to

show that the swarming colony structural differences were not due to mutants forming.

To accomplish this, different parts of the swarming colony were re-streaked onto

swarming plates and examined after 48 h at 30°C, which again showed similar colony

morphologies to the original colonies (**Fig. S9**). Since Δ*crp* and Δ*cyaB*/*cyaA* both gave a

small colony morphology on swarming plates after 48 h of growth at 30°C, we extended

the incubation time to 64 h at 30°C to ensure no swarming motility occurred **(Fig. S10).**

Overall, the data demonstrate that cAMP-CRP is an activator of both swimming and

swarming motility in *V. parahaemolyticus*.

cAMP-CRP and AC-IV CyaB are required for capsule and biofilm production.

Capsule polysaccharide (CPS) is an important component of biofilm and is positively

controlled by the quorum sensing master regulator OpaR in *V. parahaemolyticus* (43, 44,

46-49). To determine whether CRP plays a role in CPS formation, we examined the Δ*crp*,

Δ*cyaA*, Δ*cyaB*, and Δ*cyaB*/*cyaA* mutants on HI medium containing Congo red dye and

CaCl$_2$. After incubation for 48 h at 30°C, the wild type strain produced a rugose opaque

colony as did the ΔcyaA mutant, both indicatingf CPS production (**Fig. 7**). However, the

ΔcyaB mutant produced a smooth translucent colony as did the Δcrp and ΔcyaB/ΔcyaA

mutants indicating absence of CPS (**Fig. 7**). In biofilm assays, wild type and ΔcyaA

produced significant biofilm, whereas Δcrp, ΔcyaB, and ΔcyaB/ΔcyaA produced

significantly less biofilm compared to wild type (**Fig. 8**). Overall, these data

demonstrate that both CRP and CyaB, but not CyaA, are required for CPS and biofilm

production in *V. parahaemolyticus*. We functionally complemented a *V. parahaemolyticus*

ΔcyaB mutant with the *cyaB* gene on an expression vector and a biofilm assay was

performed. In this assay the wild type pBBREV (empty vector), ΔcyaB pBBREV, and

ΔcyaB pBBRcyaB were examined and the ΔcyaB pBBRcyaB complement showed

restoration of biofilm to wild type levels (**Fig. S11**).

Both Δcrp and ΔopaR deletion mutants have significant defects in capsule formation and

biofilm production. It was previously shown that OpaR is an activator of CPS binding

to the regulatory regions of the CpsA-K biosynthesis cluster and CPS regulators CspQ,

CpsR, and CpsS in *V. parahaemolyticus* BB22OP (50). We examined whether CRP and

OpaR interact to control CPS production. To accomplish this, functional complement

experiments of Δcrp and ΔopaR with either *crp* or *opaR* were performed. As controls,

both Δcrp and ΔopaR were functionally complemented with their respective gene. The

ΔopaR pBADopaR showed restoration of CPS and Δcrp pBADcrp produced a large non-

translucent colony (**Fig. 9A**). The ΔopaR pBADcrp strain remained translucent,

suggesting that *crp* cannot rescue the Δ*opaR* CPS defect (**Fig. 9A**). Also, the Δ*crp*

pABAD*opaR* mutant did not appear to restore CPS (**Fig. 9A**), however in biofilm assays,

Δ*crp* pBAD*crp* produced significantly more biofilm than Δ*crp* pBADEV (**Fig. 9B, Fig.**

9C). In the Δ*crp* pBAD*opaR* strain, significantly more biofilm was produced compared to

Δ*crp* pBADEV, but not to the same level as Δ*crp* pBAD*crp* **(Fig. 9D)**. These data suggest

that CRP is epistatic to OpaR in the biofilm regulatory network, but additional factors

are involved in biofilm regulation, which needs to be examined further.

DISCUSSION

In this study, the physiological roles of CRP and adenylate cyclase were examined in *V.*

parahaemolyticus, which led to the identification of an uncharacterized AC-IV named

CyaB. The data showed that in *V. parahaemolyticus*, CRP was an important regulator of

metabolism and motility, acting as a positive regulator of both swimming and

swarming motility. In addition, the data showed that CRP was required for capsule and

biofilm formation. In *E. coli* and other enterobacteria deleting *cyaA*, which encodes the

canonical AC-I, results in CRP remaining inactive. Thus in these species, the Δ*cyaA* and

Δ*crp* mutants have similar phenotypes (51, 52). In *V. cholerae* and *V. vulnificus*, deletion

of *cyaA* also gave phenotypes similar to Δ*crp*, showing defects in metabolism and

motility (38, 53, 54). In *V. parahaemolyticus*, the Δ*cyaA* mutant showed phenotypes

similar to wild type. This unexpected finding led us to search for proteins with putative

AC function in the *V. parahaemolyticus* genome leading to the identification of the *cyaB*

gene (VP1760). CyaB belongs to the CYTH protein superfamily (Pfam PF01928) and was

annotated as a class AC-IV, which contained one domain and was homologous to AC-

IV proteins previously described in *A. hydrophila*, *Y. pestis,* and thermophilic

archaebacteria (31-34). Our phylogenetic analysis showed that CyaB is widespread

throughout the *Vibrionaceae* family generally present in all strains of each species.

Several instances of horizontal gene transfer of *cyaB* were noted. Overall, AC-IV

homologs were present predominantly among Gamma-Proteobacteria in species

belonging to the genera *Aeromonas, Shewanella, Yersinia, Proteus, Providencia, Dickeya,*

and *Pectobacterium,* amongst others. In these genera, the CyaB phylogeny usually

followed evolutionary relationships, but some *Shewanella* species for example nested

within *Vibrio* CyaB clades suggesting horizontal gene transfer. Homologs were also

present in Alpha-Proteobacteria, but only in a handful of genera; *Cohaesibacter, Hoeflea,*

Kangsaoukella, Maritalea, and Pseudovibrio. Outside of Bacteria, proteins annotated as AC-

IV were present in Planctomycete, Crenarchaetote and Euryarchaeote species. All

bacterial homologs of AC-IV described in this study contained the N-terminal motif

HFxxxxExExK. The amino acid residues HF are an important distinguishing feature

present in proteins with nucleotidyl cyclase activity, whereas among Planctomycete and

Archaea species only the N-terminal motif ExExK is present. Vogt and colleagues recent

work showed that archaeal CYTH-like proteins are functionally diverge from bacterial

AC-IV CyaB-like proteins (35). They proposed that a true AC-IV requires the extended

N-terminal signature HFxxxxExExK, with the HF motif absent in other CYTH proteins

that function as either TTM or ThTPase (35).

The seminal studies of AC-IV in *A. hydrophila* showed that the protein was only active at

high temperatures and alkaline pH conditions (31). Furthermore, their studies showed

that deleting *cyaB* did not have any cellular effects under the conditions examined (31).

A study in *Shewanella oneidensis* characterized three AC genes, *cyaA, cyaB*, and *cyaC*

members of AC-I, AC-IV, and AC-III present in that species. They showed each *cya* gene

functionally complemented an *E. coli* Δ*cyaA*, but did not determine any other role for

cyaB (55). Indeed, to date, no physiological role for AC-IV has been shown. In our study,

we show that deleting *cyaB* results in significant defects in metabolism, capsule

production, and biofilm formation. The defects in CPS production and biofilm

formation in both the Δ*crp* and Δ*cyaB* mutants indicate that cAMP-CRP is a positive

regulator of these phenotypes and CyaB is essential for this function. The data may also

suggest that cAMP, per se, is not directly involved, but that a function specific to the

CYTH superfamily could be.

Studies in *V. cholerae* identified cAMP-CRP as a negative regulator of capsule

production and biofilm formation (38, 56). It was showed that cAMP-CRP negatively

regulated CPS (encoded by *vsp* loci) and biofilm matrix proteins by regulating genes

encoding diguanylate cyclases and phosphodiesterases. In *V. cholerae*, the *vsp*

biosynthetic operon is negatively controlled by HapR, the quorum sensing master

regulator, and it was shown that CRP was epistatic to HapR in the regulation of the

VspT and VspR regulators of capsule production (38, 56-59). In these studies both Δ*crp*

and Δ*cyaA* showed decreased expression of *hapR* suggesting that cAMP-CRP was a

positive regulator of HapR (38, 56, 60). In *V. parahaemolyticus*, the quorum sensing

master regulator OpaR is a positive regulator of capsule and biofilm production,

however, whether cAMP-CRP controls OpaR is unknown (43, 44, 46-50). Our

preliminary bioinformatics analysis identified two putative CRP binding sites in the

regulator region of OpaR suggesting CRP may be a direct regulator of *opaR* in *V.*

parahaemolyticus (**Fig. S12**). Putative CRP binding sites were also identified in *cspA-K*

regulator promoter regions, which will need to be investigated further (**Fig. S12**). It is of

interest to note that although both *V. cholerae* and *V. parahaemolyticus* use conserved

regulatory pathways, QS and CCR, to modulate capsule and biofilm production, the

effects are the direct opposite in each species. It will be of interest to determine how this

diversity in the control of important phenotypes in these related species has evolved

and how CyaB plays a role.

MATERIALS AND METHODS

Bacterial strains, media, and culture conditions.

All strains and plasmids used in this study are listed in **Table 1**. A streptomycin-

resistant clinical isolate *V. parahaemolyticus* RIMD2210633 was used throughout the

study. Unless stated otherwise, all *V. parahaemolyticus* strains were grown in Luria

Bertani (LB) medium (Fischer Scientific, Pittsburgh, PA) containing 3% NaCl (LBS) at

37°C with aeration or M9 medium media (Sigma Aldrich, St. Louis, MO) containing 3%

NaCl (M9S) supplement with 10mM of various carbon sources. Antibiotics were added

to growth media at the following concentrations: Ampicillin (Amp), 100 μg/ml,

streptomycin (Sm), 200 μg/ml, chloramphenicol (Cm), 10 μg/ml, and kanamycin (Km),

50 μg/ml when required.

Construction of mutant strains in *V. parahaemolyticus* RIMD2210633.

Splice by overlapping extension (SOE) PCR and an allelic exchange (61) was used to

construct in-frame, non-polar deletion mutants of *crp* (VP2793), *cyaA* (VP2987), *cyaB*

(VP1760) and a double deletion *cyaB/cyaA* in *V. parahaemolyticus* RIMD2210633. Primer

pairs were designed for *crp* (VP2793), *cyaA* (VP2987), and *cyaB* (VP1760) PCR

amplification using *V. parahaemolyticus* RIMD2210633 genomic DNA as a template. All

primers used in this study are listed in **Table S1**. Using the PCR amplified fragments of

each gene, SOE PCR was conducted to create truncated version of *crp* (leaving 81-bps of

633-bp VP2793), *cyaA* (153-bp of 2,529-bp VP2987) and *cyaB* (84-bp of 546-bp VP1760).

The truncated PCR fragments were cloned into the suicide vector pDS132 and named

pDSΔ*crp*, pDSΔ*cyaA*, and pDSΔ*cyaB*. pDSΔ*crp*, pDSΔ*cyaA*, and pDSΔ*cyaB* were

transformed into *E. coli* strain β2155 λ*pir*, which were conjugated separately into *V.*

parahaemolyticus. Conjugation was conducted by spread plating both strains onto LBS

plate containing 0.3 mM diaminopimelic acid (DAP). Bacterial growth from this plate

was scraped, suspended in LBS, serially diluted, and plated on LBS plates containing

streptomycin and chloramphenicol. After overnight growth, colonies were examined to

identify single crossovers of the suicide plasmids into the target site via PCR. The

colonies with single crossover were grown over night in LBS with no antibiotic and

plated onto LBS plate containing 10% sucrose to select for double crossover deletion

mutants. The double deletion mutant Δ*cyaB*/Δ*cyaA* was constructed by deleting *cyaA*

from Δ*cyaB* background following the above protocol. The gene deletions were

confirmed by PCR and DNA sequencing.

Bioinformatics analysis.

The presence of CRP binding sites (BS) in the regulatory regions of *crp, cyaA* and *cyaB* in

V. parahaemolyticus were examined using PRODORIC software using both the *E. coli* and

V. cholerae consensus sequences for Crp (62). WebLogo was used to generate graphical

representations of the multiple sequence alignments used to generate phylogenetic trees

for CyaB (63). Each single letter symbol is the single letter designation of an amino acid

at each position in the protein. The height of the each stack for each position indicates

the sequence conservation at that position and the height of each symbol within a stack

represent the frequency of each amino acid at that position.

Functional complementation of *E. coli* Δ*cyaA* and *V. parahaemolyticus* mutants.

Primer pairs were designed to amplify *cyaA* (VP2987) and *cyaB* (VP1760) using wild

type genomic DNA as template (**Table S1**). Purified PCR products of *cyaA* or *cyaB*, and

the vector pBBR1MCS were digested using BamHI and XbaI. Both the insert and the

vector were gel purified. The genes were each cloned into the vector pBBR1MCS

creating pBBR*cyaA* or pBBR*cyaB*, which were then transformed into *E. coli* B2155 using

the calcium chloride transformation method. *E. coli* β2155 containing pBBR*cyaA* and

pBBR*cyaB* were transformed into *E. coli* Δ*cyaA* and grown overnight in media

supplemented with 100 mM isopropyl-β-d-thiogalactopyranoside (IPTG) (64). For

functional complementation of *V. parahaemolyticus* Δ*crp*, Δ*cyaB* and Δ*opaR* a similar

approach was used, but *crp* and *opaR* as well as *cyaB* were cloned into pBAD33.

Complement strains were verified using PCR and sequencing.

Growth pattern analysis.

Bacterial cultures were grown overnight in LBS or M9S supplemented with 10mM

glucose at 37°C with aeration. An overnight growth culture was spun down,

resuspended, and washed with PBS and a 1: 40 dilution was then used to conduct

growth pattern analysis in M9S supplemented with 10 mM of each individual carbon

sources. Growth curves were performed in a 96 well plate at 37°C with aeration. Optical

density (OD595) was measured each hour for 24 hours. Each experiment was performed

in triplicate with two biological replications.

Swimming and swarming motility assay.

Swimming assays were performed in semi-solid media, LB 2% NaCl with 0.3% agar and

swarming assays were performed in heart infusion (HI) medium (Remel, Lenexa, KS)

containing 2% NaCl and 1.5 % agar. To examine swimming behavior, a single colony of

the test bacterium was picked and stabbed in the center of the LB plate and incubated at

37°C for 24 h. For the swarming assay, a single colony was picked and gently inoculated

on the surface of the HI plate, and the plates were incubated at 30°C for 48 h. Each

experiment was performed in triplicate with three biological replications.

Capsule polysaccharide and biofilm production assay.

Capsule polysaccharide (CPS) formation assays were conducted as previously

described (43-45). Wild type and mutant strains were grown on HI media containing

1.5% agar, 2% NaCl, 2.5 mM $CaCl_2$, and 0.25% Congo red dye for 48 h at 30°C. Each

image is an example from at least three biological replicates performed in triplicate.

Biofilm assays were performed as previously described (43-45). Briefly, overnight

cultures were used to inoculate strip wells in a 1:40 dilution in LBS. Cultures were

incubated for 24 h at 37°C prior to analysis. Biofilm was stained with crystal violet and

solubilized with dimethyl sulfoxide (DMSO) (ThermoFisher Scientific, USA). Optical

density was measured at 595nm to quantify biofilm.

Phylogenetic analysis of AC-IV.

pBLAST searches with VP1760 from *V. parahaemolyticus* used as a seed were performed

on all completed and draft genomes published on the National Center for

Biotechnology Institute (NCBI) website. Sequences were aligned using CLUSTALW

(65). The evolutionary history was inferred by using the Maximum Likelihood method

and Kimura 2-parameter model (66). The *cyaB* gene tree with the highest log likelihood

(-16932.68) is shown. The percentage of trees in which the associated taxa clustered

together is shown next to the branches. A discrete Gamma distribution was used to

model evolutionary rate differences among sites (3 categories (+*G*, parameter = 0.5711)).

This analysis involved 61 nucleotide sequences. Codon positions included were

1st+2nd+3rd. There were a total of 543 positions in the final dataset. The CyaB tree was

inferred by using the Maximum Likelihood method and Le_Gascuel_2008 model. The

tree with the highest log likelihood (-21377.64) is shown. A discrete Gamma distribution

was used to model evolutionary rate differences among sites (3 categories (+*G*,

parameter = 0.7213)). This analysis involved 177 amino acid sequences and a total of 145

positions in the final dataset. Initial tree(s) for the heuristic search were obtained

automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise

distances estimated, and then selecting the topology with superior log likelihood value.

Trees are drawn to scale, with branch lengths measured in the number of substitutions

per site. All positions with less than 95% site coverage were eliminated, i.e., fewer than

5% alignment gaps, missing data, and ambiguous bases were allowed at any position

(partial deletion option). Phylogenetic analyses were conducted with MEGA X (67). For

the *Vibrio* species that contained CyaB, a second tree was constructed for the same set of

species based on CyaA protein sequences and compared with the CyaA tree to examine

differences in branching patterns amongst species.

ACKNOWLEDGMENTS

This research was supported in part by a National Science Foundation grant (award

IOS-1656688) to E.F.B. J.G.T was funded by a University of Delaware graduate

fellowship award.

REFERENCES:

1.   Botsford JL, Harman JG. 1992. Cyclic AMP in prokaryotes. Microbiol Rev 56:100-

22.

2.   Görke B, Stülke J. 2008. Carbon catabolite repression in bacteria: many ways to

make the most out of nutrients. Nature Reviews Microbiology 6:613-624.

3.   Deutscher J. 2008. The mechanisms of carbon catabolite repression in bacteria.

Curr Opin Microbiol 11:87-93.

4.   Deutscher J, Francke C, Postma PW. 2006. How phosphotransferase system-

related protein phosphorylation regulates carbohydrate metabolism in bacteria.

Microbiol Mol Biol Rev 70:939-1031.

5.   Danchin A, Guiso N, Roy A, Ullmann A. 1984. Identification of the *Escherichia coli*

cya gene product as authentic adenylate cyclase. J Mol Biol 175:403-8.

6.   Dumay V, Danchin A, Crasnier M. 1996. Regulation of *Escherichia coli* adenylate

cyclase activity during hexose phosphate transport. Microbiology (Reading) 142 (

Pt 3):575-583.

7.   Makman RS, Sutherland EW. 1965. Adenosine 3',5'-Phosphate in *Escherichia coli*. J

Biol Chem 240:1309-14.

8.   Stülke J, Hillen W. 1999. Carbon catabolite repression in bacteria. Curr Opin

Microbiol 2:195-201.

9.   Kolb A, Busby S, Buc H, Garges S, Adhya S. 1993. Transcriptional regulation by

cAMP and its receptor protein. Annu Rev Biochem 62:749-95.

10.   Zheng D, Constantinidou C, Hobman JL, Minchin SD. 2004. Identification of the

CRP regulon using in vitro and in vivo transcriptional profiling. Nucleic Acids

Res 32:5874-93.

11.   Roy A, Danchin A. 1982. The cya locus of *Escherichia coli* K12: organization and

gene products. Mol Gen Genet 188:465-71.

12.   Roy A, Danchin A, Joseph E, Ullmann A. 1983. Two functional domains in

adenylate cyclase of *Escherichia coli*. J Mol Biol 165:197-202.

13.   Berg OG, von Hippel PH. 1988. Selection of DNA binding sites by regulatory

proteins. II. The binding specificity of cyclic AMP receptor protein to recognition

sites. J Mol Biol 200:709-23.

14.   Benoff B, Yang H, Lawson CL, Parkinson G, Liu J, Blatter E, Ebright YW, Berman

HM, Ebright RH. 2002. Structural basis of transcription activation: the CAP-alpha

CTD-DNA complex. Science 297:1562-6.

15.   Nelson SO, Wright JK, Postma PW. 1983. The mechanism of inducer exclusion.

Direct interaction between purified III of the phosphoenolpyruvate:sugar

phosphotransferase system and the lactose carrier of *Escherichia coli*. EMBO J

2:715-20.

16.   Inada T, Kimata K, Aiba H. 1996. Mechanism responsible for glucose-lactose

diauxie in *Escherichia coli*: challenge to the cAMP model. Genes Cells 1:293-301.

17.   Hogema BM, Arents JC, Bader R, Eijkemans K, Yoshida H, Takahashi H, Aiba H,

Postma PW. 1998. Inducer exclusion in *Escherichia coli* by non-PTS substrates: the

role of the PEP to pyruvate ratio in determining the phosphorylation state of

enzyme IIAGlc. Mol Microbiol 30:487-98.

18.   Saier MH, Jr., Feucht BU, Hofstadter LJ. 1976. Regulation of carbohydrate uptake

and adenylate cyclase activity mediated by the enzymes II of the

phosphoenolpyruvate: sugar phosphotransferase system in *Escherichia coli*. J Biol

Chem 251:883-92.

19.   Ullmann A, Tillier F, Monod J. 1976. Catabolite modulator factor: a possible

mediator of catabolite repression in bacteria. Proc Natl Acad Sci U S A 73:3476-9.

20.   Danchin A, Dondon L, Daniel J. 1984. Metabolic alterations mediated by 2-

ketobutyrate in *Escherichia coli* K12. Mol Gen Genetics 193:473-478.

21.   You C, Okano H, Hui S, Zhang Z, Kim M, Gunderson CW, Wang YP, Lenz P,

Yan D, Hwa T. 2013. Coordination of bacterial proteome with metabolism by

cyclic AMP signalling. Nature 500:301-6.

22.   Bârzu O, Danchin A. 1994. Adenylyl cyclases: a heterogeneous class of ATP-

utilizing enzymes. Prog Nucleic Acid Res Mol Biol 49:241-83.

23. Ramakrishnan G, Jain A, Chandra N, Srinivasan N. 2016. Computational

recognition and analysis of hitherto uncharacterized nucleotide cyclase-like

proteins in bacteria. Biol Direct 11:27.

24. Crasnier M, Dumay V, Danchin A. 1994. The catalytic domain of *Escherichia coli*

K-12 adenylate cyclase as revealed by deletion analysis of the cya gene. Mol Gen

Genet 243:409-16.

25. Ladant D, Ullmann A. 1999. Bordatella pertussis adenylate cyclase: a toxin with

multiple talents. Trends Microbiol 7:172-6.

26. Sinha SC, Sprang SR. 2006. Structures, mechanism, regulation and evolution of

class III nucleotidyl cyclases. Rev Physiol Biochem Pharmacol 157:105-40.

27. Cotta MA, Whitehead TR, Wheeler MB. 1998. Identification of a novel adenylate

cyclase in the ruminal anaerobe, Prevotella ruminicola D31d. FEMS Microbiol

Lett 164:257-60.

28. Téllez-Sosa J, Soberón N, Vega-Segura A, Torres-Márquez ME, Cevallos MA.

2002. The Rhizobium etli cyaC product: characterization of a novel adenylate

cyclase class. J Bacteriol 184:3560-8.

29. Iyer LM, Aravind L. 2002. The catalytic domains of thiamine triphosphatase and

CyaB-like adenylyl cyclase define a novel superfamily of domains that bind

organic phosphates. BMC Genomics 3:33.

30.   Martinez J, Truffault V, Hothorn M. 2015. Structural Determinants for Substrate

Binding and Catalysis in Triphosphate Tunnel Metalloenzymes. J Biol Chem

290:23348-60.

31.   Sismeiro O, Trotot P, Biville F, Vivares C, Danchin A. 1998. Aeromonas

hydrophila adenylyl cyclase 2: a new class of adenylyl cyclases with thermophilic

properties and sequence similarities to proteins from hyperthermophilic

archaebacteria. J Bacteriol 180:3339-44.

32.   Gallagher DT, Kim SK, Robinson H, Reddy PT. 2011. Active-site structure of

class IV adenylyl cyclase and transphyletic mechanism. J Mol Biol 405:787-803.

33.   Gallagher DT, Smith NN, Kim SK, Heroux A, Robinson H, Reddy PT. 2006.

Structure of the class IV adenylyl cyclase reveals a novel fold. J Mol Biol 362:114-

22.

34.   Smith N, Kim SK, Reddy PT, Gallagher DT. 2006. Crystallization of the class IV

adenylyl cyclase from Yersinia pestis. Acta Crystallogr Sect F Struct Biol Cryst

Commun 62:200-4.

35.   Vogt MS, Ngouoko Nguepbeu RR, Mohr MKF, Albers SV, Essen LO, Banerjee A.

2021. The archaeal triphosphate tunnel metalloenzyme SaTTM defines structural

determinants for the diverse activities in the CYTH protein family. J Biol Chem

297:100820.

36.   Kovacikova G, Skorupski K. 2001. Overlapping binding sites for the virulence

599   gene regulators AphA, AphB and cAMP-CRP at the *Vibrio cholerae* tcpPH

600   promoter. Mol Microbiol 41:393-407.

601   37.   Li CC, Merrell DS, Camilli A, Kaper JB. 2002. ToxR interferes with CRP-

602   dependent transcriptional activation of ompT in *Vibrio cholerae*. Mol Microbiol

603   43:1577-89.

604   38.   Liang W, Pascual-Montano A, Silva AJ, Benitez JA. 2007. The cyclic AMP

605   receptor protein modulates quorum sensing, motility and multiple genes that

606   affect intestinal colonization in *Vibrio cholerae*. Microbiology (Reading) 153:2964-

607   2975.

608   39.   Popovych N, Tzeng SR, Tonelli M, Ebright RH, Kalodimos CG. 2009. Structural

609   basis for cAMP-mediated allosteric control of the catabolite activator protein.

610   Proc Natl Acad Sci U S A 106:6927-32.

611   40.   Shimada T, Fujita N, Yamamoto K, Ishihama A. 2011. Novel roles of cAMP

612   receptor protein (CRP) in regulation of transport and metabolism of carbon

613   sources. PLoS One 6:e20081.

614   41.   Skorupski K, Taylor RK. 1997. Cyclic AMP and its receptor protein negatively

615   regulate the coordinate expression of cholera toxin and toxin-coregulated pilus in

616   *Vibrio cholerae*. Proc Natl Acad Sci U S A 94:265-70.

42.   Whitaker WB, Richards GP, Boyd EF. 2014. Loss of sigma factor RpoN increases

intestinal colonization of *Vibrio parahaemolyticus* in an adult mouse model. Infect

Immun 82:544-56.

43.   Kalburge SS, Carpenter MR, Rozovsky S, Boyd EF. 2017. Quorum Sensing

Regulators Are Required for Metabolic Fitness in *Vibrio parahaemolyticus*. Infect

Immun 85.

44.   Tague JG, Hong J, Kalburge SS, Boyd EF. 2022. Regulatory Small RNA Qrr2 Is

Expressed Independently of Sigma Factor-54 and Can Function as the Sole Qrr

Small RNA To Control Quorum Sensing in *Vibrio parahaemolyticus*. J Bacteriol

204:e0035021.

45.   Tague JG, Regmi A, Gregory GJ, Boyd EF. 2021. Fis Connects Two Sensory

Pathways, Quorum Sensing and Surface Sensing, to Control Motility in *Vibrio*

parahaemolyticus. Front Microbiol 12:669447.

46.   McCarter LL. 1998. OpaR, a homolog of Vibrio harveyi LuxR, controls opacity of

Vibrio parahaemolyticus. J Bacteriol 180:3166-73.

47.   Guvener ZT, McCarter LL. 2003. Multiple regulators control capsular

polysaccharide production in *Vibrio parahaemolyticus*. J Bacteriol 185:5431-41.

48.   Enos-Berlage JL, Guvener ZT, Keenan CE, McCarter LL. 2005. Genetic

determinants of biofilm development of opaque and translucent *Vibrio*

parahaemolyticus. Mol Microbiol 55:1160-82.

49.   Kernell Burke A, Guthrie LT, Modise T, Cormier G, Jensen RV, McCarter LL,

638   Stevens AM. 2015. OpaR controls a network of downstream transcription factors

639   in *Vibrio parahaemolyticus* BB22OP. PLoS One 10:e0121863.

640   50.   Gode-Potratz CJ, McCarter LL. 2011. Quorum sensing and silencing in *Vibrio*

641   *parahaemolyticus*. J Bacteriol 193:4224-37.

642   51.   Hufnagel DA, Evans ML, Greene SE, Pinkner JS, Hultgren SJ, Chapman MR.

643   2016. The Catabolite Repressor Protein-Cyclic AMP Complex Regulates csgD and

644   Biofilm Formation in Uropathogenic *Escherichia coli*. J Bacteriol 198:3329-3334.

645   52.   Kalivoda EJ, Stella NA, O'Dee DM, Nau GJ, Shanks RM. 2008. The cyclic AMP-

646   dependent catabolite repression system of Serratia marcescens mediates biofilm

647   formation through regulation of type 1 fimbriae. Appl Environ Microbiol

648   74:3461-70.

649   53.   Kim YR, Kim SY, Kim CM, Lee SE, Rhee JH. 2005. Essential role of an adenylate

650   cyclase in regulating *Vibrio vulnificus* virulence. FEMS Microbiol Lett 243:497-503.

651   54.   Kim YR, Lee SE, Kim B, Choy H, Rhee JH. 2013. A dual regulatory role of cyclic

652   adenosine monophosphate receptor protein in various virulence traits of *Vibrio*

653   *vulnificus*. Microbiol Immunol 57:273-80.

654   55.   Charania MA, Brockman KL, Zhang Y, Banerjee A, Pinchuk GE, Fredrickson JK,

655   Beliaev AS, Saffarini DA. 2009. Involvement of a membrane-bound class III

adenylate cyclase in regulation of anaerobic respiration in *Shewanella oneidensis*

MR-1. J Bacteriol 191:4298-306.

56.   Fong JCN, Yildiz FH. 2008. Interplay between Cyclic AMP-Cyclic AMP Receptor

Protein and Cyclic di-GMP Signaling in *Vibrio cholerae* Biofilm Formation. J

Bacteriol 190:6646-6659.

57.   Hammer BK, Bassler BL. 2003. Quorum sensing controls biofilm formation in

Vibrio cholerae. Mol Microbiol 50:101-4.

58.   Zhu J, Mekalanos JJ. 2003. Quorum sensing-dependent biofilms enhance

colonization in *Vibrio cholerae*. Dev Cell 5:647-56.

59.   Yildiz FH, Liu XS, Heydorn A, Schoolnik GK. 2004. Molecular analysis of

rugosity in a *Vibrio cholerae* O1 El Tor phase variant. Mol Microbiol 53:497-515.

60.   Silva AJ, Pham K, Benitez JA. 2003. Haemagglutinin/protease expression and

mucin gel penetration in El Tor biotype *Vibrio cholerae*. Microbiology 149:1883-

1891.

61.   Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. 1989. Site-directed

mutagenesis by overlap extension using the polymerase chain reaction. Gene

77:51-9.

62.   Grote A, Klein J, Retter I, Haddad I, Behling S, Bunk B, Biegler I, Yarmolinetz S,

Jahn D, Münch R. 2009. PRODORIC (release 2009): a database and tool platform

for the analysis of gene regulation in prokaryotes. Nucleic Acids Res 37:D61-5.

63.   Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo

generator. Genome Res 14:1188-90.

64.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita

M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame,

single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

65.   Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the

sensitivity of progressive multiple sequence alignment through sequence

weighting, position-specific gap penalties and weight matrix choice. Nucleic

Acids Res 22:4673-80.

66.   Kimura M. 1980. A simple method for estimating evolutionary rates of base

substitutions through comparative studies of nucleotide sequences. J Mol Evol

16:111-20.

67.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol

35:1547-1549.

68.   Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, Tagomori K, Iijima Y,

Najima M, Nakano M, Yamashita A, Kubota Y, Kimura S, Yasunaga T, Honda T,

Shinagawa H, Hattori M, Iida T. 2003. Genome sequence of *Vibrio*

parahaemolyticus: a pathogenic mechanism distinct from that of *V. cholerae*. Lancet

361:743-9.

69.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita

697   M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame,

698   single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006 0008.

699   70.   Dehio C, Meyer M. 1997. Maintenance of broad-host-range incompatibility group

700   P and group Q plasmids and transposition of Tn5 in Bartonella henselae

701   following conjugal plasmid transfer from *Escherichia coli*. J Bacteriol 179:538-40.

702   71.   Philippe N, Alcaraz J, Coursange E, Geiselmann J, Schneider D. 1994.

703   Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria.

704   Plasmid 51:246-255.

705   72.   Kovach ME, Phillips RW, Elzer PH, Roop RM, Peterson KM. 1994. pBBR1MCS: a

706   broad-host-range cloning vector. Biotechniques 16:800-802.

707   73.   Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation, modulation,

708   and high-level expression by vectors containing the arabinose PBAD promoter. J

709   Bacteriol 177:4121-30.

Figure Legends

Fig. 1. Growth pattern analysis in LBS and M9S D-Glucose. Growth pattern analysis

of *V. parahaemolyticus* RIMD2210633 (WT), Δ*cyaA*, Δ*cyaB*, Δ*cyaB*/Δ*cyaA*, and Δ*crp* in **(A)**

LBS **(B)** M9S + 10mM D-glucose grown at 37°C with aeration for 24 h. For growth in M9,

the strains were grown overnight in M9 3%NaCl (M9S) + 10mM glucose (M9GS) at 37°C

with aeration for 24 h. Optical density (OD$_{595}$) was measured every hour for 24 h; means

and standard errors of at least two biological replicates are displayed. For all growth

curves the area under curve (AUC) was calculated for each strain and a Student's *t*-test

was performed, *P<0.05; *** *P*<0.001.

Fig. 2. Growth pattern analysis in alternative carbon sources. Growth pattern analysis

of *V. parahaemolyticus* RIMD2210633 (WT), Δ*cyaA*, Δ*cyaB*, Δ*cyaB*/Δ*cyaA*, and Δ*crp* in **(A)**

M9S + 10mM L-arabinose, **(B)** M9S + 10mM L-ribose **(C)** M9S + 10mM D-gluconate at

37°C with aeration for 24 h. Inoculums were grown overnight in M9S + 10mM glucose

at 37°C with aeration. Optical density (OD$_{595}$) was measured every hour for 24 h; means

and standard errors of at least two biological replicates are displayed. For all growth

curves the area under curve (AUC) was calculated for each strain and a Student's *t*-test

was performed, ** P<0.01; *** *P*<0.001.

Fig. 3. Evolutionary analysis by of *cyaB* among the family *Vibrionaceae*. Phylogenetic

analysis was conducted in MEGA X. The evolutionary history was inferred by using the

Maximum Likelihood method and Kimura 2-parameter model. The *cyaB* gene tree with

the highest log likelihood (-16932.68) is shown. The percentage of trees in which the

associated taxa clustered together is shown next to the branches. This analysis involved

61 nucleotide sequences and a total of 543 positions in the final dataset.

Fig. 4. Evolutionary analysis by of CyaB among Gamma- Proteobacteria, Alpha-

Proteobacteria, Planctomycete and Archaea species. The CyaB tree was inferred by

using the Maximum Likelihood method and Le_Gascuel_2008 model. The tree with the

highest log likelihood (-21377.64) is shown. This analysis involved 177 amino acid

sequences and a total of 145 positions in the final dataset.

Fig. 5: Complementation analysis of *E. coli* Δ*cyaA*. The *E. coli* Δ*cyaA* strain was

complemented with a functional copy of *cyaA* or *cyaB* from *V. parahaemolyticus* and

growth curves were conducted in **(A)** M9 + 10 mM glucose **(B)** M9 + 10mM gluconate

for 48 h at 37°C with aeration. Optical density (OD_{595}) was measured every hour for

24 h; means and standard errors of three biological replicates are displayed. For all

growth curves the area under curve (AUC) was calculated for each strain and a

Student's *t*-test was performed, ** $P<0.01$.

Fig. 6. Characterization of motility in *V. parahaemolyticus* mutants. (A) Swimming

assays were conducted using wild type, Δ*crp*, Δ*cyaA*, Δ*cyaB*, and Δ*cyaB*/Δ*cyaA* mutants

grown at 37°C for 24 h in LB 2% NaCl with 0.3% agar. The diameter of the region

covered by the bacterial strains were measured and represented as bar graph. The error

bars indicate standard error of mean and an Unpaired Student t-test was conducted.

The significant difference is denoted by asterisks ***, P<0.001 and NS indicates not

significant **(B)** Swarming assays. Colonies of wild type and mutant strains Δ*crp*, Δ*cyaA*,

Δ*cyaB*, and Δ*cyaB*/Δ*cyaA* were assessed for swarming motility in HI containing 2% NaCl

and 1.5% agar at 30°C for 48 h.

Fig. 7. Capsule polysaccharide production. Colonies of *V. parahaemolyticus*

RIMD2210633 and mutant strains were inoculated on HI plate containing Congo red

dye and $CaCl_2$ grown at 37°C for 24 h.

Fig. 8. Biofilm formation. *V. parahaemolyticus* strains were grown in HI broth for 24 h.

A. The biofilms were visualized using crystal violet and then dissolved using DMSO,

and the optical density of the biofilms were measured at OD and **B.** represented as bar

graphs. Error bars indicate standard error of mean (SEM). Unpaired Student *t* test was

conducted to obtain the *P* values. The significant difference in biofilm production is

denoted by asterisks. *, P<0.05, **, P<0.01, ***, P<0.001. NS, not significant.

Fig. 9. A. Capsule polysaccharide (CPS) assays. The pBAD*opaR* and pBAD*crp*

complementation plasmids were conjugated into *V. parahaemolyticus* mutant strains.

Cells were grown on HI congo red $CaCl_2$ media with 0.1% L-arabinose and 12.5 μg/mL

chloramphenicol at 30ºC for 48 h prior to imaging. **B.** Biofilm production assays. *V.*

parahaemolyticus Δ*crp* mutant containing pBAD33 (pBADEV), pBAD*crp*, or pBAD*opaR*

expression plasmids were grown in 3% LB with 12.5 μg/mL chloramphenicol and 0.1%

L-arabinose for 24 hours at 37°C. **C.** Biofilm solubilized in DMSO. **D**. Biofilm measured

for absorbance at OD_{595}. Error bars indicate standard error of mean (SEM). An Unpaired

Student *t* test was conducted to obtain ***, P<0.001.

Table 1. Bacterial strains and plasmids used in this study

Bacterial strains	Genotype or description	References or sources
Vibrio parahaemolyticus		
RIMD2210633	O3:K6 clinical isolate, Sm^r	(68)
Δ*crp*	RIMD2210633 Δ*crp* (VP2793), Sm^r	This study
Δ*cyaA*	RIMD2210633 Δ*cyaA* (VP2987), Sm^r	This study
Δ*cyaB*	RIMD2210633 Δ*cyaB* (VP1760), Sm^r	This study
Δ*cyaB*/Δ*cyaA*	RIMD2210633 Δ*cyaB* Δ*cyaA* (VP2987), Sm^r	This study
Δ*rpoN*	RIMD2210633 Δ*rpoN* (VP2670), Sm^r	(42)
Δ*opaR*	RIMD2210633 Δ*opaR* (VP2516), Sm^r	(43)
Escherichia coli		
BW25113	Wild-Type *E. coli* K12 strain	(69)
Δ*cyaA*	BW25113, F-, *Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ⁻, rph-1, ΔcyaA751::kan, Δ(rhaD-rhaB)568, hsdR514*	(69)
DH5αλpir	*Δlac pir*	Laboratory collection
B2155λpir	*ΔdapA::emr pir*, for bacterial conjugation	(70)
B2155pDS*crp*	B2155 strain harboring pDS*crp*	This study
B2155pDS*cyaA*	B2155 strain harboring pDS*cyaA*	This study
B2155pDS*cyaB*	B2155 strain harboring pDS*cyaB*	This study
Complement strains		This study
E. coli Δ*cyaA*pBBR*cyaA*	*E. coli* Δ*cyaA* harboring pBBR*cyaA*	This study
E. coli Δ*cyaA*pBBR*cyaB*	*E. coli* Δ*cyaA* harboring pBBR*cyaB*	This study
E. coli Δ*cyaA*pBBR*cyaB*	*E. coli* Δ*cyaA* harboring pBBR*cyaB*	This study
V. parahaemolyticus Δ*cyaB*pBBR*cyaB*	*V. parahaemolyticus* Δ*cyaB* harboring pBBR*cyaB*	This study
V. parahaemolyticus Δ*crp*pBAD*crp*	*V. parahaemolyticus* Δ*crp* harboring pBAD*crp*	This study
V. parahaemolyticus Δ*opaR*pBAD*opaR*	*V. parahaemolyticus* Δ*opaR* harboring pBAD*opaR*	This study
V. parahaemolyticus Δ*crp*pBAD*opaR*	*V. parahaemolyticus* Δ*crp* harboring pBAD*opaR*	This study
Plasmids		
pDS132	Suicide plasmid, Cm^r, SacB	(71)
pDSΔ*crp*	pDS132 harboring truncated *crp*, Cm^r	This study

pDSΔ*cyaA*	pDS132 harboring truncated *cyaA*, Cm^r	This study
pDSΔ*cyaB*	pDS132 harboring truncated *cyaB*, Cm^r	This study
pBBR1MCS	Expression vector, P_LAC promoter, Cm^r	(72)
pBAD33	Expression vector, P_BAD promoter, Cm^r	(73)
pBAD*crp*	pBAD harboring full-length *crp* (VP2793), Cm^r	This study
pBAD*opaR*	pBAD harboring full-length *opaR*, Cm^r	This study
pBBR*cyaA*	pBBR1MCS harboring full-length *cyaA* (VP2987), Cm^r	This study
pBBR*cyaB*	pBBR1MCS harboring full-length *cyaB*(VP1760), Cm^r	This study

A. LB 3% NaCl (LBS)

B. M9 3% NaCl (M9S)

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

A. Swimming motility

B. Swarming motility

Fig. 6

Capsule polysaccharide (CPS) production

Rugose OP	Rugose OP	Smooth TR	Smooth TR	Smooth TR	Smooth TR
WT	$\triangle cyaA$	$\triangle cyaB$	$\triangle cyaB/$ $\triangle cyaA$	$\triangle crp$	$\triangle rpoN$

Fig. 7

A. Biofilm formation assay

WT   Δ*cyaA*   Δ*cyaB*   Δ*cyaB/*   Δ*crp*   Δ*rpoN*
                                       Δ*cyaA*

B. Biofilm quantification

Fig. 8

A.

$\Delta opaR$
pBAD$opaR$

$\Delta opaR$
pBADcrp

Δcrp
pBADEV

Δcrp
pBADcrp

Δcrp
pBAD$opaR$

B.

C.

Δcrp
pBADEV

Δcrp
pBADcrp

Δcrp
pBAD$opaR$

(-)
control

D.

Fig. 9.