**A class IV adenylate cyclase CyaB is required for capsule**

**polysaccharide production and biofilm formation in *Vibrio***

***parahaemolyticus***

**Regmi, A., J.G. Tague, and E.F. Boyd***

Department of Biological Sciences, University of Delaware, Newark, DE

19716

Corresponding author*

E. Fidelma Boyd

Department of Biological Sciences

341 Wolf Hall, University of Delaware

Newark, DE 19716

Phone: (302) 831-1088.  Fax: (302) 831-2281 Email: fboyd@udel.edu

**ABSTRACT**

CRP (cyclic AMP receptor protein), encoded by *crp*, is a global transcription factor that is activated by cyclic AMP (cAMP), a second messenger synthesized from ATP by adenylate cyclase, *cyaA* in *Escherichia coli*. In this species, growth on the preferred carbon source glucose, results in low levels of cAMP, and inactive CRP. When glucose is absent, cAMP levels are high which activates CRP to regulate 100s of genes. To investigate the role of CRP in *Vibrio parahaemolyticus*, we constructed non-polar deletions of the *crp* and *cyaA* homologs. The Δ*crp* mutant did not grow in minimal media supplemented with non-glucose carbon sources, however the Δ*cyaA* mutant grew similar to wild type suggesting the presence of an additional adenylate cyclase. Bioinformatics analysis identified a homolog of a class IV adenylate cyclase, CyaB present in *V. parahaemolyticus*. Both *V. parahaemolyticus cyaA* and *cyaB* functionally complemented an *E. coli* Δ*cyaA* mutant indicating adenylate cyclase activity. Growth pattern analysis showed that *cyaA* is controlled by CRP, but *cyaB* is not. cAMP-CRP was required for motility with the Δ*crp* and Δ*cyaB*/Δ*cyaA* mutants having a defect in swimming and swarming motility. The Δ*cyaA* mutant produces capsule polysaccharide (CPS) and biofilm similar to wild type. However, the Δ*cyaB* mutant was defective in CPS production and biofilm formation, similar to the Δ*crp* and Δ*cyaB*/Δ*cyaA* mutants, indicating that CyaB, but not CyaA, is required for these phenotypes. Phylogenetic analysis of *cyaB* showed broad distribution within the *Vibrionaceae* family and was acquired by horizontal gene transfer in several species.

## IMPORTANCE

The physiological role of class IV adenylate cyclases remains unknown and previously only a class I adenylate cyclase was identified within *Vibrio* species. Here we identify and characterize a class IV AC in *V. parahaemolyticus*. Our analysis shows that CyaB is essential for optimal growth and is required for capsule polysaccharide production and biofilm formation. The *cyaB* gene is widespread among *Vibrionaceae* species and phylogenetic analysis demonstrates that in some species the *cyaB* gene was acquired by horizontal gene transfer.

## INTRODUCTION

Many bacterial species repress alternative carbon catabolism gene expression when glucose is present, a mechanism known as carbon catabolite repression (CCR) (1). Two major players involved in CCR are adenylate cyclase (AC) encoded by *cyaA*, which produces cyclic AMP (cAMP) from ATP, and the global regulator CRP (**c**AMP **R**eceptor **P**rotein) encoded by *crp*. cAMP is a second messenger that activates CRP, which functions as a homodimer, and in *Escherichia coli* is a global regulator of metabolism (2-4). The CRP regulon includes genes for the transport and catabolism of non-glucose carbon sources. In *E. coli*, the 209 amino acid CRP protein has an N-terminal cAMP binding domain (CBD; residues 1–136) and a C-terminal helix turn helix DNA binding domain (DBD; residues 139–209) (5, 6). When bound by cAMP, CRP undergoes a conformation change that allows the protein to interact with target promoter regions at consensus sequences, a CRP-box consisting of a palindromic 16-bp sequence TGT**G**ANNNNNNT**C**ACA (5, 7). Analysis of 323 CRP-binding sites showed that G at position 4 and C at position 13 of the CRP-box are highly conserved (8). Also, in *E. coli* it was shown that CRP can bind cGMP, but this does not result in transcriptional changes or the conformational change required for DNA binding (9).

There are six classes (I-VI) of adenylate cyclases (AC) based on primary amino acid sequence, with different catalytic mechanisms and structures indicating different evolutionary origins (10, 11). *Escherichia coli* and many other *Gamma-Proteobacteria*

contain a single adenylate cyclase gene *cyaA*, which encodes a class I AC (AC-I). The 810

amino acid CyaA protein has two functionally distinct domains; the N-terminal

catalytic cyclase domain and the C-terminal regulatory domain. Studies in *Vibrio* species

have shown an expanded physiological role of CyaA in bacteria, specifically in

virulence, biofilm formation, motility and type III secretion regulation (12-17). In

addition, the regulatory role of CyaA is amplified by its regulation of other global

regulators such as the stress response sigma factor RpoS and quorum sensing LuxR

homologs. Class II ACs (AC-II) are comprised of bacterial exotoxins such as the

calmodulin-activated exotoxin from *Bordetella pertussis*, with an N-terminal cyclase

catalytic domain and a C-terminal hemolytic domain (18). The class III ACs are the most

diverse and widespread and encompass adenylate cyclases and guanylate cyclases

(GCs) from eukaryotic and prokaryotic species (19). A well-characterized AC-III

enzyme from bacteria is CyaC from *Pseudomonas aeruginosa* that has receptor binding

activity as well as cyclase activity. This species also contains a class I and a class II AC

(20). A class IV AC was first described in *Aeromonas hydrophila* and named CyaB, and is

a small 181 amino acid protein with unusual alkalinophilic and thermophilic activity (21).

An AC-IV was also described in *Yersinia pestis* and its structure determined (22). In

*Shewanella oneidensis* MR-1, an AC-IV was identified and shown to have adenylate

cyclase activity. This species also contains AC-I encode by *cyaA* and an AC-III encoded

by *cyaC* (23). The physiological role of AC-IVs among bacteria is unknown. A

structurally diverse AC was identified in *Prevotella ruminicola* D31d and forms the sole

representative of AC-V (24). A class VI AC was described in *Rhizobium elti* and was

shown to have cyclase activity by complementation of an *E. coli cyaA* mutant (25).

Similar to *E. coli*, *V. cholerae*, a human gastrointestinal pathogen, has a single AC,

a CyaA homolog that synthesizes cAMP, which binds to CRP to activate transcription

(17). cAMP-CRP was shown to play a crucial role in controlling virulence gene

expression, biofilm formation and quorum sensing in *V. cholerae* through positive

regulation of *hapR (luxR* homolog*)*, the quorum sensing master regulator (12, 17, 26).

HapR is a negative regulator of cholera toxin encoded *ctxAB* and the toxin co-regulated

pilus encoded by *tcp* as well as biofilm formation in this species. HapR is a repressor of

AphA, which is a positive regulator of the *ctxAB* and *tcp* genes, and AphA positively

regulates VpsR, a positive regulator of the *vsp* loci required for capsule polysaccharide

(CPS) formation and biofilm (12, 27, 28). Thus, HapR and cAMP-CRP are negative

regulators of CT and TCP expression and biofilm formation (17, 26). In *V. cholerae*,

transcriptomic analysis demonstrated that CRP is a global regulator controlling 176

genes (17). A recent study using ChIP-Seq demonstrated 119 CRP binding sites in the

genome of *V. cholerae*, most overlapping with previously identified cAMP-CRP binding

sites (29).

*Vibrio parahaemolyticus* is a halophile present in the marine environment, where it

exists free living or associated with marine flora and fauna, forming biofilms (30-32). V.

*parahaemolyticus* is also an important human pathogen associated with pandemic

infectious disease. Consumption of raw or undercooked shellfish infected with *V.*

*parahaemolyticus* causes gastroenteritis in human and is the leading cause of bacterial

seafood borne gastroenteritis worldwide (32, 33). In this species, the QS master

regulator OpaR (LuxR homolog) positively regulates biofilm by directly, positively

regulating *cpsQ* and the *cspA-K* loci required for CPS production, a major component of

biofilm (34-40). The role of cAMP-CRP in *V. parahaemolyticus* physiology has not been

previously examined.

In the present study, the function and activation of CRP in *V. parahaemolyticus*

was investigated. We constructed in-frame non-polar deletions of *crp* (VP2793) and *cyaA*

(VP2987) homologs in *V. parahaemolyticus* RIMD2201633 and examined their growth

patterns in various carbon sources. The Δ*crp* mutant grew on minimal medium M9 3%

NaCl (M9) supplemented with glucose (M9G), but was unable to grow on L-arabinose,

L- ribose, or D-gluconate as sole carbon sources. The Δ*crp* mutant also showed defects

in both swimming and swarming motility. Interestingly, the Δ*cyaA* mutant grew on all

carbon sources similar to wild type and showed no motility defects. This suggested the

presence of an additional uncharacterized adenylate cyclase gene in *V. parahaemolyticus*.

Bioinformatics analysis identified ORF VP1760 that encoded a putative class IV AC,

which we named *cyaB*, present in all *V. parahaemolyticus* strains, but absent from most *V.*

*cholerae* strains. Both *cyaA* and *cyaB* functionally complemented an *E. coli* Δ*cyaA* mutant.

In growth pattern analysis, the Δ*cyaB* and Δ*cyaA* mutants showed significantly different

growth curves from each other on all media examined. In addition, the Δ*cyaB* mutant

showed significantly different growth patterns depending on the growth media used

for overnight cultures used for inoculations. In motility assays, both Δ*cyaA* and Δ*cyaB*

mutants behaved similar to wild type, but the double Δ*cyaB*/Δ*cyaA* mutant showed

growth and motility defects similar to the Δ*crp* mutant. Next, we examined whether

cAMP-CRP played a role in CPS and biofilm formation. The Δ*crp* mutant showed

smooth translucent colony morphology demonstrating a lack of CPS, indicating  that

cAMP-CRP is a positive regulator of CPS. The Δ*cyaA* mutant produced CPS similar to

wild type with a rough rugose morphology, however the Δ*cyaB* showed a smooth

translucent colony morphology, indicating a lack of CPS. Similarly, in biofilm assays,

the Δ*cyaA* mutant produced a biofilm similar to wild type whereas the Δ*crp*, Δ*cyaB*, and

Δ*cyaB*/Δ*cyaA* mutants produce significantly less biofilm. Thus, our data show that

cAMP-CRP modulates CPS and biofilm and CyaB is essential for this function.

Phylogenetic analysis of the *cyaB* gene showed it is phylogenetically widespread among

the family *Vibrionaceae* and has been acquired by horizontal gene transfer in many

species.

**RESULTS**

**CRP is an essential regulator in *V. parahaemolyticus.***

cAMP-CRP is a global regulator of nutrient transport in many bacterial species (7-9, 13,

15, 17). Homologs of CRP and CyaA are present in *V. parahaemolyticus*, ORF VP2793 and

VP2987, respectively (**Fig. S1**). The CRP protein of *V. parahaemolyticus* shared 98% amino

acid identity to CRP in *V. cholerae*. The CyaA shared 82% amino acid identity with that

of *V. cholerae*. The genome context of both genes was conserved across *V.*

*parahaemolyticus*, *V. cholerae* and *E. coli* (**Fig. S1**). To begin to investigate the

physiological role of CRP in *V. parahaemolyticus,* we first constructed an in-frame non-

polar deletion of *crp* (VP2793) and *cyaA* (VP2987) in *V. parahaemolyticus* RIMD2210633.

The Δ*crp* mutant displayed a small colony morphology on LBS plates **(Fig. S2A)**.

Growth pattern analysis in LBS broth and M9S D-glucose broth demonstrated that the

Δ*crp* strain had a growth defect **(Fig. 1A**-**B**). We examined the effects of deleting *cyaA*,

predicting it should behave similar to the Δ*crp* deletion mutant, as seen in *E. coli* and *V.*

*cholerae* Δ*cyaA* mutants. Growth of Δ*cyaA* on LBS plates showed identical colony

morphology to wild type (**Fig. S2A)** and the growth pattern of Δ*cyaA* in LBS and M9S

D-glucose broth showed a similar pattern to wild type (**Fig. 1A**-**B**).

Next, we examined growth in alternative carbon sources and found that in M9S

supplemented with L-arabinose, L- ribose, or D-gluconate, the Δ*crp* strain was unable to

grow (**Fig 2A-C**). These data demonstrate that CRP is required for catabolism of

alternative carbon sources similar to many other species. Growth analysis of the Δ*cyaA*

mutant in M9S supplemented with L-arabinose, L- ribose, or D-gluconate, showed that

it grew similar to wild type (**Fig 2A-C**). These data suggested that *V. parahaemolyticus*

contained an additional uncharacterized adenylate cyclase that could synthesize cAMP

in the absence of *cyaA*. Alternatively, though less likely, CRP could act independently of

cAMP.

**Identification of a class IV adenylate cyclase CyaB in *V. parahaemolyticus*.**

BLAST analysis using CyaA as a seed to search the RIMD2210633 genome, only hit to

itself, indicating no other homolog of the class I adenylate cyclase was present. We then

searched the *V. parahaemolyticus* genome for class II, III and class IV adenylate cyclases

and identified a homolog of a class IV adenylate cyclase, which we named CyaB. No

additional adenylate cyclases were identified in *V. parahaemolyticus*. The CyaB protein

shared 45% amino acid identity with CyaB of *Y. pestis* and 56% amino acid identity with

CyaB of *A. hydrophila* (**Fig. S3**). In both of these species, the CyaB protein has been

structurally characterized and shown to have AC activity generating cAMP, however its

physiological role is unknown (21, 22, 41, 42). CyaB lacks transmembrane domains and

signal peptide cleavage sites indicating it is a cytosolic protein. To determine whether *V.*

*parahaemolyticus* CyaB produces cAMP, we functionally complemented an *E. coli* Δ*cyaA*

mutant with either *V. parahaemolyticus cyaA* or *cyaB* and examined growth in D-glucose

and D-gluconate. The *E. coli* Δ*cyaA* did not demonstrate growth defect in D-glucose, but

was defective in growth in D-gluconate **(Fig. 3A-B)**. The growth defect of E. *coli* Δ*cyaA*

in D-gluconate was rescued by E. *coli* Δ*cyaA* strain complemented by either *V.*

*parahaemolyticus cyaA* (VP2987) or *cyaB* (VP1760), indicating both, CyaA and CyaB, have

adenylate cyclase activity (**Fig. 3A-B**).

We reasoned that our lack of Δ*cyaA* mutant phenotypes was due to *cyaB*

complementing the Δ*cyaA* mutant. To investigate this further, we constructed a Δ*cyaB*

strain and examined growth. The Δ*cyaB* mutant had similar colony morphology to wild

type (**Fig. S2A**) and showed identical growth patterns in LBS to wild type (**Fig. 1**).

However, when the Δ*cyaB* mutant was grown overnight in M9S D-glucose and used to

inoculate M9 supplemented with D-glucose, L-arabinose, L- ribose, or D-gluconate, a

longer lag phase was observed, but a similar biomass to wild type was reached (**Fig.**

**2A-C)**.

Next, we constructed a Δ*cyaB*/Δ*cyaA* double mutant strain by deleting *cyaA* from

the Δ*cyaB* background. The double mutant demonstrated small translucent colony

morphology on LBS similar to the colonies observed for Δ*crp* (**Fig. S2A)**. Interestingly, a

plate streaked on LBS from the glycerol stock gave small translucent colonies and

regular opaque colonies, suggesting suppressor mutations were occurring. We did not

see this phenotype in either of the single *cya* mutants (**Fig. S2B**). Addition of cAMP to

the media restored the large colony morphology for the double mutant (**Fig. S2B-D**). In

*V. cholerae,* which contains only the AC-I CyaA, the Δ*cyaA* mutant reversion to wild-

type phenotypes was also observed (12). For our analysis, small colonies were always

used for characterization studies. The Δ*cyaB*/Δ*cyaA* double mutant showed a growth

defect on M9S D-glucose (**Fig. 1B**) and showed no growth in M9 supplemented with L-

arabinose, L-ribose, or D-gluconate **(Fig. 2A-C)**. The growth assays were performed in

M9S L-arabinose, L-ribose, or D-gluconate with inoculum obtained after overnight

growth in M9S D-glucose. These growth assays showed that the Δ*cyaA* mutant grew

similar to wild type, but the Δ*cyaB* mutant had a significantly longer lag phase.

However, when strains were grown overnight in LBS and then inoculated into

alternative carbon sources, the Δ*cyaB* mutant showed a shorter lag phase and a

significantly greater biomass than the Δ*cyaA* mutant, which grew similar to wild type

(**Fig. 4**). This suggest that *cyaA* is under CRP control and that *cyaB* is not under

catabolite repression. Analysis of the regulatory regions of both genes identified

putative CRP binding sites in the regulatory region of *cyaA* but not *cyaB* (data not

shown). The growth assays also demonstrate that CyaB functions as an adenylate

cyclase and CyaA and CyaB can complement each other.

**CRP regulation of swimming and swarming motility**.

*Vibrio parahaemolyticus* produces both polar and lateral flagella required for swimming

and swarming motility, respectively (43). To investigate the role of CRP in *V.*

*parahaemolyticus* motility, we conducted swimming assays and found that the Δ*crp*

mutant was defective in swimming compared to wild type but did not show a complete

defect similar to the Δ*rpoN* mutant (**Fig. 5A**). Similarly, the Δ*crp* mutant also showed a

defect in swarming as indicated by the absence of the budding phenotype compared to

wild type (**Fig. 5B**). Neither of the single Δ*cyaA* nor Δ*cyaB* mutants showed a motility

defect, however the Δ*cyaB*/Δ*cyaA* double mutant showed defects similar to Δ*crp* (**Fig. 5A**

**and 5B**). These data demonstrate that the cAMP-CRP is required for motility of *V.*

*parahaemolyticus*.

**CRP and CyaB required for CPS and biofilm production.**

Capsule polysaccharide (CPS) production is a key component of biofilm and is

positively controlled by the quorum sensing master regulator OpaR in *V.*

*parahaemolyticus* (34, 36, 38-40, 44). To determine whether CRP plays a role in CPS

formation, we examined the Δ*crp* and *cya* mutants on HI medium supplemented with

congo red and $CaCl_2$, incubated for 48 hrs at 30°C. The wild type strain produced a

rugose opaque colony as did the Δ*cyaA* mutant, indicative of CPS production (**Fig. 6**).

However, the Δ*cyaB* mutant produced a smooth translucent colony indicative of the

absence of CPS. The Δ*crp and* Δ*cyaB*/Δ*cyaA* mutants also did not produce CPS,

presenting a smooth translucent colony morphology (**Fig. 6**). These data show that CRP

is a positive regulator of CPS production and *cyaB* is also essential. In biofilm assays, a

similar pattern emerged as seen with CPS, where the Δ*cyaA* mutant produce a biofilm

similar to wild type, however both the Δ*crp* and Δ*cyaB* mutants produced significantly

less biofilm (**Fig. 7**). Overall, these data demonstrate that both CRP and CyaB, but not

CyaA, are required for CPS and biofilm production in *V. parahaemolyticus*.

**Distribution and phylogenetic analysis of *cyaB*.**

CyaB was first identified in a strain of *A. hydrophila, Y. pestis*, and a number of

hyperthermophilic archaebacterial (21). We used *cyaB* (VP1760) as a seed and examined

the distribution among members of the family *Vibrionaceae* in the NCBI genome

database. The *cyaB* gene was present in all *V. parahaemolyticus* strains and was highly

conserved. CyaB was absent from most *V. alginolyticus* strains, a closely related species

to *V. parahaemolyticus*, but was present in other members of the Harveyi clade to which

*V. parahaemolyticus* belongs, such as *V. campbellii, V. harveyi, V. jasicida, V. owensii, and V.*

*rotiferanus* (**Fig. 8**). These species showed ~70% amino acid identity with CyaB, but ~95%

amino acid identity to CyaA from *V. parahaemolyticus*. On the *cyaB* gene tree, within the

Harveyi clade, species formed two divergent branches, one branch representing *V.*

*parahaemolyticus* strains and the second branching pattern representing the other

species. This suggests that *cyaB* in *V. parahaemolyticus* has diverged significantly from

*cyaB* in the other Harveyi clade species. Of note is a *cyaB* gene from a strain of *V.*

*alginolyticus*, which is present on a plasmid that clusters within the same group as *cyaB*

from *V. coralliilyticus*, which is also present on a plasmid (**Fig. 8**). This suggest that in

these strains and species, *cyaB* was acquired by horizontal gene transfer. In general, the

*cyaB* gene from the genera *Aliivibrio* and *Photobacterium* cluster closely together and are

divergent from *Vibrio* species, as one would expect if the gene is ancestral to the groups

(**Fig. 8**). However, *cyaB* from five *Photobacterium* species clustered within *Vibrio* clades

indicating that these *Photobacterium* species acquired a copy of the *cyaB* gene from

*Vibrio*, another example of horizontal gene transfer of *cyaB* in these groups (**Fig. 8**).

Comparison of CyaA phylogeny with the CyaB phylogeny from the same set of species

also demonstrated additional instances of horizontal transfer of *cyaB* among species

shown by the non-congruency of the branching patterns for several CyaB proteins (**Fig.**

**S4**).

**DISCUSSION**

Here we describe the physiological role of a class IV adenylate cyclase (AC). In this

study, we demonstrated that in *V. parahaemolyticus*, CRP is an important regulator of

metabolism and motility, acting as a positive regulator of both swimming and

swarming motility. In addition, we showed that CRP is a positive regulator of CPS and

biofilm formation. Multiple studies in *E. coli* and enterobacteria, in general, have

demonstrated that deleting *cyaA* impairs the cell's cAMP production thus deactivating

CRP, which led to Δ*cyaA* and Δ*crp* mutants demonstrating identical phenotypes (45, 46).

In *V. cholerae* and *V. vulnificus,* deletion of *cyaA* gave phenotypes identical to Δ*crp,*

showing defects in metabolism and motility (17, 47, 48). In our study in *V.*

*parahaemolyticus,* the Δ*cyaA* mutant showed phenotypes similar to wild type. This

unexpected finding prompted us to search for proteins with putative AC function in the

*V. parahaemolyticus* genome and led us to identify *cyaB* (VP1760). CyaB belongs to CYTH

superfamily of proteins, a class IV AC, which contains one domain. The canonical AC

CyaA, which is a class I type, contains two domains: a cyclase domain and a regulatory

domain. CyaB contains a single domain and a homolog was previously identified in *A*.

*hydrophila*, *Y. pestis* and hyperthermophilic archaebacteria (21, 22, 41, 42). In *A.*

*hydrophila*, CyaB was only active at high temperatures and alkaline pH conditions and

not active under normal laboratory condition (21). In that study, it was demonstrated

that deleting *cyaB* did not show any cAMP-CRP phenotypic defects. However, when

*cyaB* was expressed from a plasmid in the Δ*cyaA* and Δ*cyaA*/Δ*cyaB* mutant strains, it was

able to restore the phenotypes back to wild type (21). No physiological role for CyaB

has been uncovered. In contrast, in our study, deleting *cyaB* showed defects in

metabolism, CPS production, and biofilm formation similar to the Δ*crp* mutant, but with

less severity. This indicates that CyaB is important for these phenotypes and that CyaA

could be compensating for the loss of CyaB. The double mutant Δ*cyaB*/Δ*cyaA*

demonstrated phenotypes identical to Δ*crp* indicating that only after removing both

CyaA and CyaB, the cells became devoid of cAMP.

Under optimal growth conditions in M9S D-glucose, deleting *cyaA* did not have

any impact on CRP function, thus we do not see any phenotypic changes. However,

deleting *cyaB* showed growth differences absent in the Δ*cyaA* mutant. When the Δ*cyaB*

mutant was grown overnight in M9S D-glucose, cell growth was impaired on non-PTS

sugars, presenting as a longer lag phase, suggesting in these cells *cyaA* was repressed

and needs time to adjust to non-PTS sugars. However, when the Δ*cyaB* mutant is grown

overnight in LBS, cells are primed for growth on non-PTS sugars, as *cyaA* is expressed

in LBS and these cells show significantly better growth and shorter lag phases than

either wild type or the Δ*cyaA* mutant. To investigate this further, bioinformatics analysis

was used to identify the putative CRP binding sites in the regulatory region of *cyaA* and

*cyaB*. We identified putative CRP binding sites in the regulatory region of *cyaA*, one of

which overlaps the RNAP binding site, characteristic of a repressor (data not shown).

No CRP binding sites were identified in the regulatory region of *cyaB*, suggesting that

this gene is regulated differently than *cyaA*. The data also suggest that the presence of

*cyaB* may have a negative impact on growth since the Δ*cyaB* mutant grows better than

wild type and the Δ*cyaA* mutant. The differences between the Δ*cyaA* and Δ*cyaB* mutants

will likely be explained in terms of differences in expression and activation of these

enzymes in response to specific environmental signals.

The defects in CPS production and biofilm formation in both the Δ*crp* and Δ*cyaB*

mutants indicate that cAMP-CRP is a positive regulator of these phenotypes and CyaB

is essential for this function. Studies in *V. cholerae* identified cAMP-CRP as a negative

regulator of capsule production and biofilm formation (17, 49). These studies showed

that cAMP-CRP negatively regulated vibrio polysaccharide (VPS) genes and biofilm

matrix proteins. The quorum sensing regulator HapR was also shown to negatively

regulate these phenotypes in *V. cholerae* (50-52). In *V. cholerae*, *vsp* gene regulation is

negatively controlled by HapR and it was shown that *crp* was epistatic to *hapR* in the

regulation of *vspT* and *vspR* regulators of VSP production (17, 49-52). It is of interest to

note that both a Δ*crp* and Δ*cyaA* mutants showed decreased expression of *hapR*

indicating that CRP is a positive regulator of HapR (16, 17, 49). In *V. parahaemolyticus*,

the quorum sensing regulator OpaR is a positive regulator of CPS and biofilm,

however, whether cAMP-CRP controls OpaR is unknown. Our preliminary

bioinformatics analysis idenitifed putative CRP binding sites in the regulator region of

OpaR suggesting CRP is may also regulated opaR in *V. parahaemolyticus*, which will

need to be investigated further.

**MATERIALS AND METHODS**

**Bacterial strains, media, and culture conditions.**

All strains and plasmids used in this study are listed in **Table 1**. A streptomycin-

resistant clinical isolate *V. parahaemolyticus* RIMD2210633 was used throughout the

study. Unless stated otherwise, all *V. parahaemolyticus* strains were grown in Luria

Bertani (LB) medium (Fischer Scientific, Pittsburgh, PA) containing 3% NaCl (LBS) at

37°C with aeration or M9 medium media (Sigma Aldrich, St. Louis, MO) containing 3%

NaCl (M9S) supplement with 10mM of various carbon sources. Antibiotics were added

to growth media at the following concentrations: Ampicillin (Amp), 100 μg/ml,

streptomycin (Sm), 200 μg/ml, chloramphenicol (Cm), 10 μg/ml, and kanamycin (Km),

50 μg/ml when required.

**Construction of mutant strains in *V. parahaemolyticus* RIMD2210633.**

Splice by overlapping extension (SOE) PCR and an allelic exchange (53) was used to

construct in-frame, non-polar deletion mutants of *crp* (VP2793), *cyaA* (VP2987), *cyaB*

(VP1760) and double deletion *cyaB/cyaA* in *V. parahaemolyticus* RIMD2210633. Primer

pairs were designed for *crp* (VP2793), *cyaA* (VP2987), and *cyaB* (VP1760) PCR

amplification using *V. parahaemolyticus* RIMD2210633 genomic DNA as a template. All

primers used in this study are listed in **Table S1**. Using the PCR amplified fragments of

each gene, SOE PCR was conducted to obtain truncated version of *crp* (81-bps of 633-bp

VP2793), *cyaA* (153-bp of 2,529-bp VP2987) and *cyaB* (84-bp of 546-bp VP1760). The

truncated PCR fragments were cloned into the suicide vector pDS132 and named

pDS*Δcrp*, pDS*ΔcyaA,* and pDS*ΔcyaB*. pDS*Δcrp*, pDS*ΔcyaA,* and pDS*ΔcyaB* were

transformed into *E. coli* strain β2155 *λpir*, which were conjugated separately into *V.*

*parahaemolyticus*. Conjugation was conducted by spread plating both strains onto LBS

plate containing 0.3 mM diaminopimelic acid (DAP). Bacterial growth from this plate

was scrapped, suspended in LBS, serially diluted, and plated on LBS plates containing

streptomycin and chloramphenicol. After overnight growth, colonies were examined to

identify single crossovers of the suicide plasmids into the target site via PCR. The

colonies with single crossover were grown over night in LBS with no antibiotic and

plated onto LBS plate containing 10% sucrose to select for double crossover deletion

mutant. The double deletion mutant Δ*cyaB*/Δ*cyaA* was constructed by deleting *cyaA*

from Δ*cyaB* background following the above protocol. The gene deletions were

confirmed by PCR and DNA sequencing.

**Functional complementation of *E. coli* Δ*cyaA*.**

Primer pairs were designed to amplify *cyaA* (VP2987) and *cyaB* (VP1760) using wild

type genomic DNA as template (**Table S1**). Purified PCR products of *cyaA* and *cyaB*,

and the vector pBBR1MCS were digested using BamHI and XbaI. Both the insert and

the vector were gel purified. The genes were each cloned into the vector pBBR1MCS

creating pBB*cyaA* or pBB*cyaB*, which were then transformed into *E. coli* B2155 using the

calcium chloride transformation method. *E. coli* β2155 containing pBB*cyaA* and pBB*cyaB*

were transformed into *E. coli* Δ*cyaA* (54). The transformed complementation strain was

selected for using LB agar supplemented with Km and Cm. Complement strains were

verified using PCR and sequencing. IPTG (0.5mM) was used to induce the transcription

of the respective genes from the vector.

**Growth pattern analysis.**

Bacterial cultures were grown overnight in LBS or M9S supplemented with 10mM

glucose at 37°C with aeration. An overnight growth culture was spun down,

resuspended, and washed with PBS and a 1: 40 dilution was then used to conduct

growth pattern analysis in M9S supplemented with 10 mM of each individual carbon

sources. Growth curves were performed in a 96 well plate at 37°C with aeration. Optical

density (OD595) was measured each hour for 24 hours. The data was plotted using

GraphPad Prism 7.00, GraphPad software, La Jolla, California USA,

www.graphpad.com.

**Swimming and Swarming motility assay.**

Swimming assays were performed in semi-solid media, LB 2% NaCl with 0.3% agar and

swarming assays were performed in heart infusion (HI) medium (Remel, Lenexa, KS)

containing 2% NaCl and 1.5 % agar. To examine swimming behavior, a single colony of

the test bacterium was picked and stabbed in the center of the LB plate and incubated at

37°C for 24 hrs. For the swarming assay, a single colony was picked and gently

inoculated on the surface of the HI plate, and the plates were incubated at 30°C for 48

hrs.

**Capsule polysaccharide production assay.**

Capsule polysaccharide (CPS) formation assays were conducted as previously

described (55, 56). Wild type and mutants of interest were grown on HI media

containing 1.5% agar, 2% NaCl, 2.5 mM $CaCl_2$, and 0.25% Congo red dye for 48 hrs at

30°C. Each image is an example from at least three biological replicates.

**Biofilm formation assay.**

Biofilm assays were performed as previously described (39, 40, 57). Briefly, overnight

cultures were used to inoculate strip wells in a 1:40 dilution in LBS. Cultures were

incubated for 24 hrs at 37°C prior to analysis. Biofilm was stained with crystal violet and

solubilized with dimethyl sulfoxide (DMSO) (ThermoFisher Scientific, USA). Optical

density was measured at 595nm to quantify biofilm.

**Phylogenetic analysis of *cyaB*.**

pBLAST searches with VP1760 from *V. parahaemolyticus* used as a seed were performed

on all completed and draft *Vibrionaceae* genomes published on the National Center for

Biotechnology Institute (NCBI) website. Sequences were aligned using CLUSTALW

(58). Phylogenetic analysis was conducted with MEGA X(59). The evolutionary history

was inferred by using the Maximum Likelihood method and Kimura 2-parameter

model (60). The tree with the highest log likelihood (-16932.68) is shown. The

percentage of trees in which the associated taxa clustered together is shown next to the

branches. Initial tree(s) for the heuristic search were obtained automatically by applying

Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using

the Maximum Composite Likelihood (MCL) approach, and then selecting the topology

with superior log likelihood value. A discrete Gamma distribution was used to model

evolutionary rate differences among sites (3 categories (+$G$, parameter = 0.5711)). The

tree is drawn to scale, with branch lengths measured in the number of substitutions per

site. This analysis involved 61 nucleotide sequences. Codon positions included were

1st+2nd+3rd. All positions with less than 95% site coverage were eliminated, that is

fewer than 5% alignment gaps, missing data, and ambiguous bases were allowed at any

position (partial deletion option). There were a total of 543 positions in the final dataset.

For the species that contained CyaB, a second tree was constructed for the same set of

species based on CyaA protein sequences and compared with the CyaA tree for

differences in branching patterns amongst species.

**ACKNOWLEDGMENTS**

This research was supported in part by a National Science Foundation grant (award

IOS-1656688) to E.F.B. J.G.T was funded by a University of Delaware graduate

fellowship award.

**REFERENCES:**

1.      Botsford JL, Harman JG. 1992. Cyclic AMP in prokaryotes. Microbiol Rev 56:100-

22.

2.      Stülke J, Hillen W. 1999. Carbon catabolite repression in bacteria. Curr Opin

Microbiol 2:195-201.

3.      Kolb A, Busby S, Buc H, Garges S, Adhya S. 1993. Transcriptional regulation by

cAMP and its receptor protein. Annu Rev Biochem 62:749-95.

4.      Zheng D, Constantinidou C, Hobman JL, Minchin SD. 2004. Identification of the

CRP regulon using in vitro and in vivo transcriptional profiling. Nucleic Acids

Res 32:5874-93.

5.      Benoff B, Yang H, Lawson CL, Parkinson G, Liu J, Blatter E, Ebright YW, Berman

HM, Ebright RH. 2002. Structural basis of transcription activation: the CAP-alpha

CTD-DNA complex. Science 297:1562-6.

6.      Lawson CL, Swigon D, Murakami KS, Darst SA, Berman HM, Ebright RH. 2004.

Catabolite activator protein: DNA binding and transcription activation. Curr

Opin Struct Biol 14:10-20.

7.      Berg OG, von Hippel PH. 1988. Selection of DNA binding sites by regulatory

proteins. II. The binding specificity of cyclic AMP receptor protein to recognition

sites. J Mol Biol 200:709-23.

8.     Shimada T, Fujita N, Yamamoto K, Ishihama A. 2011. Novel roles of cAMP

454            receptor protein (CRP) in regulation of transport and metabolism of carbon

455            sources. PLoS One 6:e20081.

456    9.     Popovych N, Tzeng SR, Tonelli M, Ebright RH, Kalodimos CG. 2009. Structural

457            basis for cAMP-mediated allosteric control of the catabolite activator protein.

458            Proc Natl Acad Sci U S A 106:6927-32.

459    10.    Bârzu O, Danchin A. 1994. Adenylyl cyclases: a heterogeneous class of ATP-

460            utilizing enzymes. Prog Nucleic Acid Res Mol Biol 49:241-83.

461    11.    Ramakrishnan G, Jain A, Chandra N, Srinivasan N. 2016. Computational

462            recognition and analysis of hitherto uncharacterized nucleotide cyclase-like

463            proteins in bacteria. Biol Direct 11:27.

464    12.    Skorupski K, Taylor RK. 1997. Control of the ToxR virulence regulon in *Vibrio*

465            *cholerae* by environmental stimuli. Mol Microbiol 25:1003-1009. .

466    13.    Kovacikova G, Skorupski K. 2001. Overlapping binding sites for the virulence

467            gene regulators AphA, AphB and cAMP-CRP at the Vibrio cholerae tcpPH

468            promoter. Mol Microbiol 41:393-407.

469    14.    Li CC, Crawford JA, DiRita VJ, Kaper JB. 2000. Molecular cloning and

470            transcriptional regulation of ompT, a ToxR-repressed gene in Vibrio cholerae.

471            Mol Microbiol 35:189-203.

15.  Li CC, Merrell DS, Camilli A, Kaper JB. 2002. ToxR interferes with CRP-

dependent transcriptional activation of ompT in Vibrio cholerae. Mol Microbiol

43:1577-89.

16.  Silva AJ, Pham K, Benitez JA. 2003. Haemagglutinin/protease expression and

mucin gel penetration in El Tor biotype Vibrio cholerae. Microbiology 149:1883-

1891.

17.  Liang W, Pascual-Montano A, Silva AJ, Benitez JA. 2007. The cyclic AMP

receptor protein modulates quorum sensing, motility and multiple genes that

affect intestinal colonization in Vibrio cholerae. Microbiology (Reading) 153:2964-

2975.

18.  Ladant D, Ullmann A. 1999. Bordatella pertussis adenylate cyclase: a toxin with

multiple talents. Trends Microbiol 7:172-6.

19.  Sinha SC, Sprang SR. 2006. Structures, mechanism, regulation and evolution of

class III nucleotidyl cyclases. Rev Physiol Biochem Pharmacol 157:105-40.

20.  Smith RS, Wolfgang MC, Lory S. 2004. An adenylate cyclase-controlled signaling

network regulates Pseudomonas aeruginosa virulence in a mouse model of acute

pneumonia. Infect Immun 72:1677-84.

21.  Sismeiro O, Trotot P, Biville F, Vivares C, Danchin A. 1998. Aeromonas

hydrophila adenylyl cyclase 2: a new class of adenylyl cyclases with thermophilic

properties and sequence similarities to proteins from hyperthermophilic

archaebacteria. J Bacteriol 180:3339-44.

22.   Smith N, Kim SK, Reddy PT, Gallagher DT. 2006. Crystallization of the class IV

adenylyl cyclase from Yersinia pestis. Acta Crystallogr Sect F Struct Biol Cryst

Commun 62:200-4.

23.   Charania MA, Brockman KL, Zhang Y, Banerjee A, Pinchuk GE, Fredrickson JK,

Beliaev AS, Saffarini DA. 2009. Involvement of a membrane-bound class III

adenylate cyclase in regulation of anaerobic respiration in Shewanella oneidensis

MR-1. J Bacteriol 191:4298-306.

24.   Cotta MA, Whitehead TR, Wheeler MB. 1998. Identification of a novel adenylate

cyclase in the ruminal anaerobe, Prevotella ruminicola D31d. FEMS Microbiol

Lett 164:257-60.

25.   Téllez-Sosa J, Soberón N, Vega-Segura A, Torres-Márquez ME, Cevallos MA.

2002. The Rhizobium etli cyaC product: characterization of a novel adenylate

cyclase class. J Bacteriol 184:3560-8.

26.   Skorupski K, Taylor RK. 1997. Cyclic AMP and its receptor protein negatively

regulate the coordinate expression of cholera toxin and toxin-coregulated pilus in

Vibrio cholerae. Proc Natl Acad Sci U S A 94:265-70.

27.   Matson JS, Withey JH, DiRita VJ. 2007. Regulatory networks controlling Vibrio

cholerae virulence gene expression. Infect Immun 75:5542-9.

28.    Beyhan S, Bilecen K, Salama SR, Casper-Lindley C, Yildiz FH. 2007. Regulation of

512           rugosity and biofilm formation in Vibrio cholerae: comparison of VpsT and VpsR

513           regulons and epistasis analysis of vpsT, vpsR, and hapR. J Bacteriol 189:388-402.

514    29.    Manneh-Roussel J, Haycocks JRJ, Magán A, Perez-Soto N, Voelz K, Camilli A,

515           Krachler AM, Grainger DC. 2018. cAMP Receptor Protein Controls Vibrio

516           cholerae Gene Expression in Response to Host Colonization. mBio 9.

517    30.    Colwell RR, Kaper JB, Joseph A. 1977. Vibrio cholerae, Vibrio parahaemolyticus,

518           and other vibrios: occurrence and distribution in Chesapeake. Science 198:394-

519           396.

520    31.    Joseph SW, Colwell RR, Kaper JB. 1982. *Vibrio parahaemolyticus* and related

521           halophilic *Vibrios.* Crit Rev Microbiol 10:77-124.

522    32.    DePaola A, Kaysner CA, Bowers J, Cook DW. 2000. Environmental investigations

523           of Vibrio parahaemolyticus in oysters after outbreaks in Washington, Texas, and

524           New York (1997 and 1998). Appl Environ Microbiol 66:4649-54.

525    33.    Nair G, Ramamurthy T, Bhattacharya S, Dutta B, Takeda Y, Sack D. 2007. Global

526           dissemination of *Vibrio parahaemolyticus* serotype O3:K6 and its serovariants. Clin

527           Microbiol Rev 20:39-48

528    34.    McCarter LL. 1998. OpaR, a homolog of Vibrio harveyi LuxR, controls opacity of

529           Vibrio parahaemolyticus. J Bacteriol 180:3166-73.

35.  Enos-Berlage JL, McCarter LL. 2000. Relation of capsular polysaccharide

production and colonial cell organization to colony morphology in Vibrio

parahaemolyticus. J Bacteriol 182:5513-20.

36.  Kernell Burke A, Guthrie LT, Modise T, Cormier G, Jensen RV, McCarter LL,

Stevens AM. 2015. OpaR controls a network of downstream transcription factors

in Vibrio parahaemolyticus BB22OP. PLoS One 10:e0121863.

37.  Gode-Potratz CJ, McCarter LL. 2011. Quorum sensing and silencing in Vibrio

parahaemolyticus. J Bacteriol 193:4224-37.

38.  Guvener ZT, McCarter LL. 2003. Multiple regulators control capsular

polysaccharide production in Vibrio parahaemolyticus. J Bacteriol 185:5431-41.

39.  Kalburge SS, Carpenter MR, Rozovsky S, Boyd EF. 2017. Quorum Sensing

Regulators Are Required for Metabolic Fitness in Vibrio parahaemolyticus. Infect

Immun 85.

40.  Tague JG, Hong J, Kalburge SS, Boyd EF. 2022. Regulatory Small RNA Qrr2 Is

Expressed Independently of Sigma Factor-54 and Can Function as the Sole Qrr

Small RNA To Control Quorum Sensing in Vibrio parahaemolyticus. J Bacteriol

204:e0035021.

41.  Gallagher DT, Kim SK, Robinson H, Reddy PT. 2011. Active-site structure of

class IV adenylyl cyclase and transphyletic mechanism. J Mol Biol 405:787-803.

42.    Gallagher DT, Smith NN, Kim SK, Heroux A, Robinson H, Reddy PT. 2006.

Structure of the class IV adenylyl cyclase reveals a novel fold. J Mol Biol 362:114-

22.

43.    Whitaker WB, Richards GP, Boyd EF. 2014. Loss of sigma factor RpoN increases

intestinal colonization of *Vibrio parahaemolyticus* in an adult mouse model. Infect

Immun 82:544-56.

44.    Enos-Berlage JL, Guvener ZT, Keenan CE, McCarter LL. 2005. Genetic

determinants of biofilm development of opaque and translucent Vibrio

parahaemolyticus. Mol Microbiol 55:1160-82.

45.    Hufnagel DA, Evans ML, Greene SE, Pinkner JS, Hultgren SJ, Chapman MR.

2016. The Catabolite Repressor Protein-Cyclic AMP Complex Regulates csgD and

Biofilm Formation in Uropathogenic Escherichia coli. J Bacteriol 198:3329-3334.

46.    Kalivoda EJ, Stella NA, O'Dee DM, Nau GJ, Shanks RM. 2008. The cyclic AMP-

dependent catabolite repression system of Serratia marcescens mediates biofilm

formation through regulation of type 1 fimbriae. Appl Environ Microbiol

74:3461-70.

47.    Kim YR, Kim SY, Kim CM, Lee SE, Rhee JH. 2005. Essential role of an adenylate

cyclase in regulating Vibrio vulnificus virulence. FEMS Microbiol Lett 243:497-

503.

48.     Kim YR, Lee SE, Kim B, Choy H, Rhee JH. 2013. A dual regulatory role of cyclic

569            adenosine monophosphate receptor protein in various virulence traits of Vibrio

570            vulnificus. Microbiol Immunol 57:273-80.

571     49.     Fong JCN, Yildiz FH. 2008. Interplay between Cyclic AMP-Cyclic AMP Receptor

572            Protein and Cyclic di-GMP Signaling in Vibrio cholerae Biofilm Formation.

573            Journal of Bacteriology 190:6646-6659.

574     50.     Hammer BK, Bassler BL. 2003. Quorum sensing controls biofilm formation in

575            Vibrio cholerae. Mol Microbiol 50:101-4.

576     51.     Zhu J, Mekalanos JJ. 2003. Quorum sensing-dependent biofilms enhance

577            colonization in Vibrio cholerae. Dev Cell 5:647-56.

578     52.     Yildiz FH, Liu XS, Heydorn A, Schoolnik GK. 2004. Molecular analysis of

579            rugosity in a Vibrio cholerae O1 El Tor phase variant. Mol Microbiol 53:497-515.

580     53.     Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. 1989. Site-directed

581            mutagenesis by overlap extension using the polymerase chain reaction. Gene

582            77:51-9.

583     54.     Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita

584            M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame,

585            single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

586     55.     Güvener Z, McCarter L. 2003. Multiple regulators control capsular

587            polysaccharide production in *Vibrio parahaemolyticus*. J Bacteriol 185:5431-41.

588    56.    Kalburge SS, Carpenter MR, Rozovsky S, Boyd EF. 2017. Quorum sensing

589           regulators are required for metabolic fitness in *Vibrio parahaemolyticus*. Infect

590           Immun 85:930-16.

591    57.    Tague JG, Regmi A, Gregory GJ, Boyd EF. 2021. Fis Connects Two Sensory

592           Pathways, Quorum Sensing and Surface Sensing, to Control Motility in Vibrio

593           parahaemolyticus. Front Microbiol 12:669447.

594    58.    Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the

595           sensitivity of progressive multiple sequence alignment through sequence

596           weighting, position-specific gap penalties and weight matrix choice. Nucleic

597           Acids Res 22:4673-80.

598    59.    Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

599           Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol

600           35:1547-1549.

601    60.    Kimura M. 1980. A simple method for estimating evolutionary rates of base

602           substitutions through comparative studies of nucleotide sequences. J Mol Evol

603           16:111-20.

604    61.    Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, Tagomori K, Iijima Y,

605           Najima M, Nakano M, Yamashita A, Kubota Y, Kimura S, Yasunaga T, Honda T,

606           Shinagawa H, Hattori M, Iida T. 2003. Genome sequence of *Vibrio*

607    *parahaemolyticus*: a pathogenic mechanism distinct from that of *V. cholerae*. Lancet

608        361:743-9.

609    62.    Grenier F, Matteau D, Baby V, Rodrigue S. 2014. Complete Genome Sequence of

610        Escherichia coli BW25113, Genome Announc, vol 2.

615    **Table 1. Bacterial strains and plasmids used in this study**

| Bacterial strains | Genotype or description | References or sources |
|---|---|---|
| *Vibrio parahaemolyticus* | | |
| RIMD2210633 | O3:K6 clinical isolate, Sm$^r$ | (61) |
| Δ*crp* | RIMD2210633 Δ*crp* (VP2793), Sm$^r$ | This study |
| Δ*cyaA* | RIMD2210633 Δ*cyaA* (VP2987), Sm$^r$ | This study |
| Δ*cyaB* | RIMD2210633 Δ*cyaB* (VP1760), Sm$^r$ | This study |
| Δ*cyaB*/Δ*cyaA* | RIMD2210633 Δ*cyaB* Δ*cyaA* (VP2987), Sm$^r$ | This study |
| Δ*rpoN* | RIMD2210633 Δ*rpoN* (VP2670), Sm$^r$ | (43) |
| *Escherichia coli* | | |
| BW25113 | Wild-Type *E. coli* K12 strain | (62) |
| Δ*cyaA* | BW25113, F-, *Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ-, rph-1, ΔcyaA751::kan, Δ(rhaD-rhaB)568, hsdR514* | |
| DH5α λpir | | Laboratory collection |
| B2155λpir | *ΔdapA::emr pir*, for bacterial conjugation | Laboratory collection |
| B2155pDS*crp* | B2155 strain harboring pDS*crp* | This study |
| B2155pDS*cyaA* | B2155 strain harboring pDS*cyaA* | This study |
| B2155pDS*cyaB* | B2155 strain harboring pDS*cyaB* | This study |
| **Complement strains** | | This study |
| *E. coli* Δ*cyaA*pCyaA | *E. coli* Δ*cyaA* harboring pBBR*cyaA* | This study |
| *E. coli* Δ*cyaA*pCyaB | *E. coli* Δ*cyaA* harboring pBBRcyaA | This study |
| **Plasmids** | | |
| pDS132 | Suicide plasmid, Cm$^r$, SacB | |
| pDSΔ*crp* | pDS132 harboring truncated *crp*, Cm$^r$ | This study |
| pDSΔ*cyaA* | pDS132 harboring truncated *cyaA*, Cm$^r$ | This study |
| pDSΔ*cyaB* | pDS132 harboring truncated *cyaB*, Cm$^r$ | This study |
| pBBR1MCS | Expression vector, lacZ promoter, Cm$^r$ | This study |
| pBBR*cyaA* | pBBR1MCS harboring full-length *cyaA* (VP2987), Cm$^r$ | This study |
| pBBR*cyaB* | pBBR1MCS harboring full-length *cyaB*(VP1760), Cm$^r$ | This study |

**Figure Legends**

**Fig. 1. Growth pattern analysis in LBS and M9S D-Glucose.** Growth pattern analysis

of *V. parahaemolyticus* RIMD2210633 (closed circle), Δ*cyaA* (open circle), Δ*cyaB* (open

square), Δ*cyaB*/Δ*cyaA* (open triangle) and Δ*crp* (closed downward facing triangle) in **(A)**

LBS **(B)** M9S + 10mM D-glucose at 37°C with aeration for 24 hrs. For growth in minimal

media (M9), the strains were grown overnight in M9 3%NaCl (M9S) + 10mM glucose

(M9GS) at 37°C with aeration for 24 hrs.

**Fig. 2. Growth pattern analysis in alternative carbon sources.** Growth pattern analysis

of *V. parahaemolyticus* (closed circle), Δ*cyaA* (open circle), Δ*cyaB* (open square),

Δ*cyaB*/Δ*cyaA* (open triangle) and Δ*crp* (closed downward facing triangle) in **(A)** M9S +

10mM L-arabinose, **(B)** M9S + L-ribose **(C)** M9S + 10mM D-gluconate at 37°C with

aeration for 24 hrs. Inoculums were grown overnight in M9S + 10mM glucose at 37°C

with aeration.

**Fig. 3: Complementation analysis of *E. coli* Δ*cyaA*.** The *E. coli* Δ*cyaA* strain was

complemented separately with a functional copy of *cyaA* and *cyaB* from *V.*

*parahaemolyticus* and growth curves were conducted in **(A)** M9 + 10 mM glucose **(B)** M9

+ 10mM gluconate for 48 hrs at 37°C with aeration.

**Fig. 4. Growth pattern analysis in alternative carbon sources.** Growth pattern analysis

of *V. parahaemolyticus* (closed circle), Δ*cyaA* (open circle), Δ*cyaB* (open square),

Δ*cyaB*/Δ*cyaA* (open triangle) and Δ*crp* (closed downward facing triangle) in **(A)** M9S +

10mM L-arabinose, **(B)** M9S + L-ribose **(C)** M9S + 10mM D-gluconate at 37°C with

aeration for 24 hrs. Inoculums were grown overnight in LBS at 37°C with aeration.

**Fig. 5. Characterization of motility in *V. parahaemolyticus* mutants. (A)** Swimming

assays were conducted using wild type, Δ*crp*, Δ*cyaA*, Δ*cyaB*, and Δ*cyaB*/Δ*cyaA* mutants

grown at 37°C for 24 hrs. The diameter of the region covered by the bacterial strains

were measured and represented as bar graph **(B**) Swarming assays. Colonies of wild

type and mutant strains Δ*crp*, Δ*cyaA*, Δ*cyaB*, and Δ*cyaB*/Δ*cyaA* were assessed for

swarming motility in semi solid media over 48 hrs.

**Fig. 6. Capsule polysaccharide production**. Colonies of *V. parahaemolyticus*

RIMD2210633 and mutant strains were inoculated on HI plate containing Congo red

dye and grown at 37°C for 24 hrs.

**Fig. 7. Biofilm formation**. *V. parahaemolyticus* strains were grown for 24 hrs in HI broth.

The biofilms were visualized using crystal violet and then dissolved using DMSO, and

the optical density of the biofilms were measured at OD (600) and represented as bar

graphs. Error bars indicate standard error of mean (SEM). Unpaired Student *t* test was

conducted to obtain the *P* values. The significant difference in biofilm production is

denoted by asterisks. *, $P<0.05$, **, $P<0.01$, ***, $P<0.001$. NS, not significant.

**Fig. 8. Evolutionary analysis by Maximum Likelihood method of *cyaB***. Evolutionary

analyses were conducted in MEGA X. The evolutionary history was inferred by using

the Maximum Likelihood method and Kimura 2-parameter model. A discrete Gamma

distribution was used to model evolutionary rate differences among sites (5 categories

(+*G*, parameter = 0.5599)). The tree is drawn to scale, with branch lengths measured in

the number of substitutions per site. A discrete Gamma distribution was used to model

evolutionary rate differences among sites (3 categories (+*G*, parameter = 0.5711)). The

tree is drawn to scale, with branch lengths measured in the number of substitutions per

site. This analysis involved 61 nucleotide sequences. Codon positions included were

1st+2nd+3rd. All positions with less than 95% site coverage were eliminated (partial

deletion option). There was a total of 543 positions in the final dataset.

**Supplementary Data**

**Table S1.**

| Primer name and use | Sequence (5′-3′)ᵃ | Melting temp (ᵒC) | Size (bp) |
|---|---|---|---|
| **SOE PCR** | | | |
| VP1760 SOEA | GGA<u>TCTAGA</u>CAGCTCAACTGGGTTTGGTT | 62 | 408 |
| VP1760 SOEB | TTCAAACTGCCCTTGGAAATG | 54 | |
| VP1760 SOEC | CATTTCCAAGGGCAGTTTGAAGAAGCGGATAGAGAGCATCG | 66 | 513 |
| VP1760 SOED | CAC<u>GAGCTC</u>GCTGCTTATCTTTGCGCTTT | 64 | |
| VP1760 SOEFF | ACGGTGAAAGCCATCAAAAC | 54 | 1,708 |
| VP1760 SOEFR | CCTTCGGTCTTACAATAACCAATC | 54 | |
| VP2793 SOEA | <u>TCTAGA</u>TTGTGGTGATTCGTCTGCGT | 60 | 458 |
| VP2793 SOEB | TGTTGGGTCGGTTTGAGGTT | 57 | |
| VP2793 SOEC | AACCTCAAACCGACCCAACACTAATCTCTGCGCACGGTAA | 68 | 451 |
| VP2793 SOED | <u>GCATGC</u>CCATCTTCGTCAAGCAGGGT | 65 | |
| VP2793 SOEFF | CTGCACACCTTTGACGAAGC | 57 | 2,408 |
| VP2793 SOEFR | GAGTCGTTATGACGCCCGAT | 57 | |
| VP2987 SOEA | <u>TCTAGA</u>CAGGATGAGCGGCTTGAAGA | 61 | 513 |
| VP2987 SOEB | GCGATCGATACGCTGCTGGTT | 60 | |
| VP2987 SOEC | **AACCAGCAGCGTATCGATCGC**CACCCTGAAGAGGGTAACGC | 71 | 503 |
| VP2987 SOED | <u>GAGCTC</u>GCACCACACTCAGGAAAAGC | 64 | |
| VP2987 SOEFF | CGCACCTTGAGGGAGTTCAT | 57 | 4,042 |
| VP2987 SOEFR | GAGGATGTTGACCCACAGCA | 57 | |
| **Complementation** | | | |
| *cyaA* com F | CAA<u>GGATCC</u>TGTTCAACCAAGGAATTACCC | 61 | 2,563 |
| *cyaA* com R | GCA<u>TCTAGA</u>GGATATGTTTACGTTACGCG | 59 | |
| *cyaB* com F | CTT<u>GGATCC</u>AGTTTATGTAGTAGAGGCGA | 59 | 576 |
| *cyaB* com R | CAC<u>TCTAGA</u>GAAATAGCGACTATGCTGAT | 57 | |

ᵃUnderlined base pairs indicate restriction sites

**Fig. S1. Genomic context comparison of *crp, cyaA* and *cyaB* loci.** The genomic region

upstream and downstream of **(A)** *crp* and **(B)** *cyaA* **(C)** *cyaB* from *V. parahaemolyticus* was

compared with that of *crp* and *cyaA* from *V. cholerae, V. vulnificus* and *E. coli* and *cyaB*

from *V. harveyi* using tBLASTx, and figures were constructed using Easyfig.

**Fig. S2. Colony Morphology of *Vibrio parahaemolyticus* strains. A.** Single colony of *V.*

*parahaemolyticus* RIMD2210633 and its isogenic mutant strains Δ*cyaA*, Δ*cyaB*,

Δ*cyaB*/Δ*cyaA,* and Δ*crp* were plated on LBS agar plate and grown overnight at 37$^{O}$C

before the picture was taken. **B.** An inoculum from the glycerol stock of Δ*cyaB*/Δ*cyaA*

was plated on LBS agar and grown overnight at 37 $^{O}$C showing small and large

colonies. **C.** Comparison of Wild type (WT) and double mutant *cyaB*/*cyaA*. **D**. Addition

of 1mM cAMP restores large colony morphology to *cyaB*/*cyaA*.

**Fig. S3: CyaB amino acid sequence and crystal structure. A.** Alignment of CyaB among

*Aeromonas hydrophila, V. parahaemolyticus* and *Yersinia pestis.*

**Fig. S4**. **Phylogenetic comparison of CyaA and CyaB trees**. The CyaA tree ws

constructed using proteins sequences from the same set of species that contained CyaB.

Lines between the two trees of similar color represent species that cluster together on

the CyaA tree.

**Fig. 1**

## A. M9S + 10mM L-arabinose

## B. M9S + 10mM L-ribose

## C. M9S + 10mM D-gluconate

**Fig. 2**

**A. M9 + 10mM D-glucose**

**B. M9 + 10mM D-gluconate**

**Fig. 3**

## M9 + 10mM L-arabinose

## M9 + 10mM L-ribose

## M9 + 10mM D-gluconate

**Fig. 4**

## A. Swimming motility

## B. Swarming motility

WT        ΔcyaA     ΔcyaB  ΔcyaB/  Δcrp  ΔrpoN
                                    ΔcyaA

**Fig. 5**

# Capsule polysaccharide (CPS) production

| Rugose OP | Rugose OP | Smooth TR | Smooth TR | Smooth TR | Smooth TR |
|:---:|:---:|:---:|:---:|:---:|:---:|
| WT | ΔcyaA | ΔcyaB | ΔcyaB/ΔcyaA | Δcrp | ΔrpoN |

Fig. 6

## A. Biofilm formation assay

WT     Δ*cyaA*     Δ*cyaB*     Δ*cyaB/*    Δ*crp*     Δ*rpoN*
                                    Δ*cyaA*

## B. Biofilm quantification

Fig. 7

**Fig. 8**