Quantitative trait locus mapping of osmotic stress response in the wheat pathogen

Zymoseptoria tritici

Authors:

Jessica Stapley and Bruce A. McDonald

Plant Pathology Group, Institute of Integrative Biology, ETH Zurich, 8092 Zürich, Switzerland

Corresponding author: Jessica Stapley, Jessica.stapley@usys.ethz.ch

9
10
11
12

Abstract

Osmotic stress is a ubiquitous and potent stress for all living organisms. Fungal molecular

responses to salt stress in well studied model yeast species highlighted the importance of

the conserved HOG MAP kinase pathway. However, fewer studies have investigated the

genetic basis of salt tolerance in filamentous fungi. The main aim of this study was to

identify regions of the genome associated with tolerance to potassium chloride in the wheat

pathogen *Zymoseptoria tritici.* A secondary aim was to identify candidate genes affecting

salt tolerance within the most promising chromosomal regions. We achieved these aims

with a quantitative trait locus (QTL) mapping study using offspring from two crosses grown

in vitro in the presence or absence of osmotic stress imposed by 0.75M KCl. We identified

significant QTL for most of the traits in both crosses. Several QTL overlapped with QTL

identified in earlier studies for other traits and some QTL explained trait variation in both

the control and salt stress environments. A significant QTL on chromosome 3 explained

variation in colony radius at 8 days post inoculation (dpi) in the KCl environment as well as

colony radius KCl tolerance at 8 dpi. The QTL peak had a high LOD and encompassed an

interval containing only 36 genes. Six of these genes present promising candidates for

functional analyses.

Keywords:
Salt stress; plant pathogen; fungi; *Mycosphaerella graminicola*, QTL

1. Introduction

Osmotic stress is a common and potent environmental stress that can limit growth and

reproduction in many organisms. At the cellular level osmotic stress results in changes to

the movement of water into or out of the cell through osmosis. To counteract osmotic

stress, organisms need to accumulate compatible solutes (e.g. glycerol, erythritol, arabitol,

mannitol and trehalose) – that can balance osmosis and provide turgor pressure (Araújo et

al., 2020; Kogej et al., 2007). In fungal pathogens of plants, the pathogen may be exposed to

a relatively high salt concentration (0.5M) within the plant (see Araújo et al., 2020) as well

as experience changes in salt and osmolyte concentrations as plant tissues degrade (Jacob

et al., 2015). The genes and pathways involved in responding to the resulting osmotic

stresses may also affect plant-pathogen interactions and virulence. For example, turgor

pressure which is mediated by changes in vacuole size and number is important in both

plant tissue penetration and response to osmotic stress (Richards et al., 2010). Voltage-

gated chloride channels, which are involved in the regulation of cell volume, also influence

virulence in fungal pathogens of both plants and humans (Cañero and Roncero, 2008; Zhu

and Williamson, 2003). Knowledge of the genetic basis of osmotic sensitivity in plant

pathogens can thus provide important insights into the genetic basis of environmental

stress responses as well as disease development.

The immediate effects of osmotic stress in fungi include stalled growth, changes in the

recruitment of water, collapse of the cytoskeleton, altered plasma membranes and a

reduction in glycerol transport (Duran et al., 2010; Hohmann, 2002). Longer-term

adaptation to osmotic stress involves modifications of metabolism, plasma membranes, cell

wall thickness and cell wall melanisation, as well as an accumulation of mycosporines

(Hohmann, 2002; Kogej et al., 2007). Osmotic stress is also likely to activate many genes and

pathways involved in the 'general stress response'. For example, up to 10% of the genes

that were differentially expressed under different individual stressors were shared across

different osmotic stress treatments (Hohmann, 2002).

Although the key salt stress response pathways and genes have been identified and well

characterized in model yeasts, studies in filamentous fungi could provide novel insights into

the genes involved in responding to osmotic stress. The HOG pathway is an osmoresponsive

system and one of the most well-understood mitogen-activated protein (MAP) kinase

pathways (Hohmann, 2002). In yeast, two genes are central components of this pathway –

Pbs2 and Hog1p kinase. In plant pathogens, HOG signalling appears more complex than in

yeast, although it is still poorly understood (Jacob et al., 2015). Several filamentous

pathogenic fungi (including our model *Zymoseptoria tritici*) have multiple (up to 17) histidine

kinases, whereas *S. cerevisiae* has only one (Sln1p) (Jacob et al., 2015). Several studies have

identified transcription factors (TFs) involved in regulating the fungal response to salt stress,

as well as other stressors - including temperature and fungicide stress. For example, in

Aspergillus oryzae, GATA TFs were shown to be important in responding to salt and

temperature stress (Jiang et al., 2021). In the causal agent of Fusarium head blight

(*Fusarium graminearum*), the nutrient and stress factor 1 (NSf1) C_2H_2 zinc fingers protein

plays an important role in salt stress response, sexual-asexual reproduction, vegetative

growth, fungicide sensitivity and pathogenicity (Shi et al., 2021), and a Zn(2)-C6 fungal-type

DNA-binding TF (Fss1) is required for sodium and lithium tolerance (Son et al., 2015). Cell

wall melanisation also appears to confer benefits for osmotic stress tolerance. For example,

cell wall melanisation in halotolerant fungi helps to maintain high concentrations of glycerol

and contain the resulting turgor pressure by strengthening and preventing leakage through

the cell wall (Kogej et al., 2007). These examples coming from a relatively small number of

fungal species demonstrate how the molecular mechanisms of salt response can vary across

taxa and illustrate that additional work is needed to understand how fungi adapt to salt

stress.

To improve our understanding of osmotic stress response in fungi we performed a

quantitative trait loci (QTL) mapping study using the wheat pathogen *Z. tritici* growing under

salt stress. QTL studies have been instrumental in finding candidate genes for multiple traits

in this widespread and damaging wheat pathogen, including melanisation (Lendenmann et

al., 2014), fungicide sensitivity (Lendenmann et al., 2015), morphological switching

(Francisco et al., 2022) and oxidative stress (Zhong et al., 2021), and enabled the cloning and

functional validation of genes affecting virulence (Zhong et al., 2017; Meile et al. 2018),

melanisation (Krishnan et al. 2018), and morphological switching (Francisco et al., 2022).

QTL mapping identifies genomic regions that explain variation in a trait and that are likely to

harbour genes responsible for creating variation in the trait. The aim of this study was to

identify large effect loci associated with tolerance to salt stress in *Z. tritici* and identify

plausible candidate genes within the most promising QTL intervals.

2. Methods

2.1. QTL mapping crosses

Parent and progeny isolates used in this study are from two crosses that were described

previously (Lendenmann et al., 2014). The original choice of strains for making crosses was

based upon a comprehensive phenotyping campaign of an international collection of 150

strains, including 30 strains from Switzerland (Zhan et al., 2005). In this original phenotyping

campaign, we measured seven traits for each strain, including in planta virulence and

pycnidia size and density, and growth rates in vitro under different temperatures and in the

presence of fungicides. Based on data from these seven traits, four Swiss strains were

selected to make the QTL mapping crosses that yielded 700 offspring over the two crosses.

The four parents have since been tested under other environmental stressors (e.g. reactive

oxygen (Zhong et al., 2021)) including KCl (0.25-1.0M). In the presence of 0.75M KCl the

parents differed in both growth and melanisation traits (Table S1). The crosses were made

by coinfecting wheat leaves with the parent strains, with the resulting ascospores collected

from infected leaves and grown in vitro. One cross was between ST99CH1A5 and

ST99CH1E4 (herein referred to as 1A5 and 1E4 respectively) and the other was between

ST99CH3D7 and ST99CH3D1 (herein referred to as 3D7 and 3D1 respectively).

2.2. Genotyping

SNP genotype data for the progeny were obtained from a RAD sequence dataset that was

previously produced in our lab (first described in (Lendenmann et al., 2014). In brief, the

genome was cut using the restriction enzyme *Pst*I and the libraries were sequenced on an

Illumina HiSeq2000 with paired-end sequencing. Complete genome sequences of the

parental strains (Croll et al., 2013)(NCBI Biosample: SRS383146 (ST99CH3D1), SRS383147

(ST99CH3D7), SRS383142 (ST99CH1A5), and SRS383143 (ST99CH1E4)) were used to SNP

genotype the parents and offspring.

RADseq processing and variant discovery is described in detail at

https://github.com/jessstapley/QTL-mapping-Z.-tritici. In brief, RADseq reads were trimmed

of adaptors and low-quality sequence using trimmomatic (v0.35). The RADseq reads were

mapped to a reference genome using bwa mem (v0.7.17). Reads from progeny of the

3D7x3D1 cross were mapped to the 3D7 reference genome and progeny from the 1A5x1E4

cross were mapped to the 1A5 reference genome. Variant calling was done using the GATK

Germline Short Variant Discovery pipeline, following their Best Practice recommendations

(https://gatk.broadinstitute.org/hc/en-us/sections/360007226651-Best-Practices-

Workflows). After this we applied the following filters: only biallelic SNPs, parents had

alternative alleles, <50% missing genotypes per marker, <50% missing genotypes per

individual, mean read depth >3 and <30, depth quality (QD)>5, mapping quality >40, minor

allele frequency >0.02 and <0.80, and SNPs in regions where the SNP density is >3 in 10bp

were removed.

2.3. Generating linkage maps

Linkage maps were made using R (R Core Team, 2020), R Studio (RStudio Team, 2020) and

the R package qtl (v1.48-1 (Broman et al., 2003)). As the order and base pair positions of the

markers were known from the reference genomes, we only needed to calculate the

recombination distance between markers to create the linkage map. Before map

construction, we removed putative clones. To do this we calculated genetic similarity using

the 'comparegeno' function and removed individuals with >99% shared genotypes (83

progeny were removed from the 1A5x1E4 cross, 74 progeny were removed from the

3D7x3D1 cross). The linkage map was constructed for each chromosome separately using

the 'estimatemap' function with method 'morgan'. Then we performed multiple rounds of

cleaning to remove error-prone markers that caused map inflation. First, we removed

markers that had an estimated recombination distance above the 99.9% quantile. Then

recombination rate was estimated again and we inspected the maps. In regions of the map

where there was evidence of map inflation and spurious double recombination events, we

visualised these regions in an Integrated Genome Viewer (IGV, https://igv.org) to look at the

mapping alignments. Markers were removed from regions with poor alignments and

evidence of mismapping. A summary of the two linkage maps is provided in Supplementary

Material S1 and the complete maps are available at (https://github.com/jessstapley/QTL-

mapping-Z.-tritici). After applying all of these filters, the average marker spacing was 1217

bp (0.19 cM) and 529 bp (0.14 cM) for the 1A5x1E4 and 3D7x3D1 crosses, respectively.

2.4. Phenotyping

We compared colony growth and melanisation in a control environment and in the presence of potassium chloride (KCl) to simulate a salt stress environment. Potassium chloride (KCl) has been used extensively in the study of stress responses in fungi because it is environmentally ubiquitous and lacks the toxic effects of Na^+ (Araújo et al., 2020; Hohmann, 2002). Only non-clonal offspring strains were phenotyped (for 1A5x1E4 n= 259 strains; for 3D7x3D1 n=265 strains). The basic protocols for isolate recovery from -80^{0}C storage, growth in vitro and the measurements of colony size and colony melanisation were described previously (Zhong et al., 2021). In brief, Petri dishes containing Difco potato dextrose agar (PDA) were inoculated with 200μl of a spore solution (concentration 200 spores/ml) and grown at 18^{0}C for 8 and 12 days. In the salt stress environment, a sterile potassium chloride solution was added to the cooled and sterilized PDA to make a final concentration of 0.75M. Three replicates were grown in both the control and salt (KCl) environments.

Two traits were measured; colony area and colony grey value, using automated image analysis as described previously (Lendenmann et al., 2014; Zhong et al., 2021). Grey value was measured on a scale of 0-255, where darker, more melanised colonies have lower values (0 = black, 255 = white). Mean colony area (mm^2) and grey value were calculated from measurements of multiple colonies on each plate and then the mean was calculated across the three replicate plates to obtain a final mean colony area and mean grey value for each strain in the two environments. Colony area was converted to colony radius by dividing by π and taking the square root of this value. Measurements were taken at 8- and 12-days post inoculation (dpi). For the analysis we used three different types of data; (1) time point/age: the colony radius and grey value at 8 and 12 dpi; (2) daily rate measurements: the change in colony radius or grey value between 8 and 12 dpi (i.e. the difference in colony radius between 8 and 12 dpi/difference in days); and (3) KCl tolerance measurements: the ratio of a measurement (colony radius, grey value, growth rate or melanisation rate) between the KCl environment and the control environment (i.e. colony radius KCl tolerance at 8dpi = colony radius at 8 dpi in the KCl environment/colony radius at 8 dpi in the control environment, or growth rate KCl tolerance = growth rate in the KCl environment/growth

rate in the control environment). For the KCl tolerance measurements, values >1 indicate

that for that trait an isolate is more tolerant to KCl (higher value under salt stress), and

values <1 indicate that for that trait measured an isolate is more sensitive to KCl. In total

there were 18 traits (Table 1).

2.5. QTL mapping

QTL mapping was performed using the R (v 3.6.0) package 'qtl2' (v2_0.24) as described in

detail on github (https://github.com/jessstapley/QTL-mapping-Z.-tritici). We scanned the

genome for a single QTL per chromosome with the '*scan1*' function using a linear mixed

effect model, and we controlled for the relatedness of individuals (i.e. we included a

random polygenic effect) by fitting the kinship matrix as a random effect. Models that take

into account the genetic covariance between individuals can reduce the false discovery rate

in QTL scans and outperform models that do not include this information in the model

(Malosetti et al., 2011). The significance threshold for a QTL peak was determined by

permutation tests (n=1000) and we calculated a Bayes Credible Interval (95%) to identify the

interval size around the QTL peak. We used custom R scripts to extract a list of the genes

within the interval from the annotation (gff) files of each reference genome. The putative

encoded function of each gene was determined in previous analyses (for 1A5 see

(Plissonneau et al., 2018), for 3D7 see (Plissonneau et al., 2016)). The annotations were

obtained using Interproscan (https://www.ebi.ac.uk/interpro/) against multiple protein

databases and then screened using multiple methods (e.g. 8ignal

https://services.healthtech.dtu.dk/service.php?SignalP-5.0, Phobius

https://phobius.sbc.su.se/) to determine if the encoded proteins contained likely signal

peptides or secreted domains. The putative effect of a SNP in the coding sequence was

determined using SNPeff (http://pcingola.github.io/SnpEff/).

2.6. Identifying putative candidate genes in the 3D7x3D1 cross

Genes within a single QTL interval on chromosome 3 were further investigated to identify

the most likely candidate genes responsible for the phenotype. We identified the

orthologous genes in the 3D1 parent using an analysis performed across the genomes of 19

reference strains (Badet et al., 2020). We inspected the alignments in the region of interest

in IGV. We also blasted the 3D1 gene sequences (including ±200bp flanking regions) against

the 3D7 genome to calculate sequence similarity and identify indels.

2.7. RNAseq data analysis

We analysed RNAseq data that was previously created for the four parental strains during

infection of the susceptible wheat cultivar Drifter (Palma-Guerrero et al., 2017) as well as

cultures grown in vitro on two types of liquid media (yeast sucrose broth (YSB): yeast extract

10 g/L and sucrose 10 g/L, pH 6.8; and carbon-depleted minimal medium (MM), pH 5.8

(Francisco et al., 2019)). The raw sequence reads were downloaded from the Short Read

Archive (SRA, https://www.ncbi.nlm.nih.gov/sra, Bioproject: *invitro* SRP152081, *invivo*

SRP077418). RNAseq data processing and analysis is described in detail at github

(https://github.com/jessstapley/QTL-mapping-Z.-tritici). In brief, reads were trimmed and

then mapped to one of two reference genomes (reads from 3D1 and 3D7 were mapped to

the 3D7 reference genome while reads from 1E4 and 1A5 were mapped to the 1A5

reference genome). Then the number of reads mapping to each gene was counted using

R::Rsubread (Liao et al., 2019) and we tested for differential gene expression between

strains, 3D7 versus 3D1 and 1A5 versus 1E4, using R::EdgeR (Robinson et al., 2010).

3. Results

3.1. Effect of KCl on growth and melanisation

A total of 158,480 and 120,059 colonies from the 1A5x1E4 and 3D7x3D1 crosses,

respectively, were phenotyped (see Table S2 for details). Differences in growth and

melanisation between salt stress and control environments, and growth rate KCl tolerance

of and melanisation rate KCl tolerance can be seen in Figure 1. We used paired Wilcoxon

tests to compare mean trait values between the control and KCl treatments (Table S2). In

the presence of KCl, colony radius was smaller at 8 and 12 dpi in both crosses, and the

growth rate was reduced in the 3D7x3D1 cross, but not in the 1A5x1E4 cross, suggesting

that isolates from the 3D7x3D1 cross were more sensitive to KCl (Figure 1, Table S2). This is

consistent with the greater growth rate tolerance in the 1A5x1E4 cross compared to the

3D7x3D1 cross ($F_{(1,487)}$ = 9.19, *p*=0.002, growth rate KCl tolerance: 1A5x1E4 = 1.04, 3D7x3D1

= 0.96). In the KCl environment we observed less melanisation at both time points and a

similar reduced accumulation of melanin over time (melanisation rate) in both crosses

(Figure 1, Table S2, $F_{(1,486)}$ = 2.85, p=0.09, melanisation rate KCl tolerance: 1A5x1E4 = 0.196,

3D7x3D1 = 0.149).

3.2. Correlations between traits

All pairwise correlations (Pearson, non-parametric) between independent measurements

are provided in Tables S3 and S4. Excluding traits that are statistically or functionally

coupled, which includes: correlations between two time points of the same trait (e.g. colony

radius at 8 dpi and 12 dpi in the control), correlations between a rate and the corresponding

time point measurements (e.g. grey value at 12 dpi and melanisation rate in the control), or

correlations between rates and KCl tolerance measurements, we identified many

correlations between independent measurements (Table S3 and S4). Notably, growth rate in

the control environment was correlated with growth rate in the KCl environment in both

crosses (r(1A5x1E4) = 0.63 and r(3D7x3D1) = 0.62, Figure S2) – thus isolates that grow faster

in a benign environment also grow faster in the presence of salt stress, i.e. they have a

higher intrinsic growth rate. Melanisation rates showed a similar trend in 1A5x1E4 – isolates

that had higher melanisation rates in the benign environment also had higher melanisation

rates under KCl stress (Table S4, r = 0.19), but this was not observed in the 3D7x3D1 cross

(Table S3, r = 0.07, Figure S2).

We were particularly interested in the relationship between KCl tolerance and melanisation,

as melanin can help protect cells from osmotic stress. In both crosses, the mean grey value

in the control environment at 12 dpi was negatively correlated with growth rate KCl

tolerance (Tables S2 and S4, 1A5x1E4 r = -0.26 and 3D7x3D1 r = -0.43, Figure S2). The

relationship at 8 dpi was also negative in both crosses, but not significant in the 3D7x3D1

cross (Tables S3 and S4, 1A5x1E4 r = -0.44 and 3D7x31 r = -0.12, Figure S2). As lower mean

grey values correspond to higher amounts of melanin, these correlations indicate that

isolates that produce more melanin in a benign environment have a growth rate that is

more tolerant to KCl stress, although this relationship was not significant at 8 dpi in the

3D7x3D1 cross.

3.3. QTL results

We identified at least one significant QTL for all but three traits (melanisation rate and mean

grey value at 12 dpi in the KCl environment, and melanisation rate KCl tolerance at 12dpi) in

the 1A5x1E4 cross and all but one trait (melanisation rate KCl tolerance) in the 3D7x3D1

cross (Table 1). In general, the traits lacking significant QTL had relatively low heritability

(Table 1) and thus the power to detect a QTL for those traits was relatively low. All QTL

positions, interval sizes and numbers of genes within the interval are presented in Tables 2

and 3. Overlapping QTL were grouped and given a unique QTL identifier. Examples of how

traits measured in different environments can have overlapping QTL intervals are shown in

Figure 2. The QTL No. 4, 9 and 11 are each regions of the genome that explain variation in

growth rate in both the KCl and control environments. As expected, traits that were highly

correlated and/or statistically coupled often had a QTL peak at the same, or overlapping

positions (e.g. the 1A5x1E4 grey values in the control environment at 8 dpi and 12 dpi

mapped to the same position on Chr 2: 1438934bp; the 3D7x3D1 QTL for colony radius at 12

dpi and growth rate in the control environment mapped to Chr 8: 1100862bp). LOD plots for

all traits listed in Tables 2 and 3 are provided in the Supplementary Information S3 and S4.

In Figures 2 and 3 we provide example plots for only four traits (growth rate in KCl, growth

rate in the control, melanisation rate in KCl and melanisation rate in the control) for the two

crosses.

3.4. QTL shared across environments, traits, and time points

Several QTL were shared amongst benign and salt-stress environments, across time points

and traits. Only one of these shared QTL is novel to this study, the other shared QTL were

found in earlier studies investigating different stressors. The single novel QTL shared

between benign and salt-stress environments was found in the 3D7x3D1 cross on Chr 8

(1100.86-1221.33 Kb, QTL No. 9, Figure 2) and explained variation in seven traits including

growth in the control and KCl environments, as well as mean grey value in the control

environment and grey value KCl tolerance at 8 dpi (Figure 2, Table 2).

Of the QTL shared across benign and KCl environments that were identified here and found

in earlier studies using different stresses (e.g. cold, fungicide), the most striking – in terms of

the number of traits and the magnitude of the LODs, was the QTL on Chr 11 between

507.36-658.11 Kb (QTL No. 12) in the 3D7x3D1 cross. This QTL explained variation in 11 of

the 18 traits, and in both environments (Table 2). This QTL was identified previously and was

found to explain variation in growth, fungicide sensitivity and melanisation in this cross

(Lendenmann et al., 2015, 2014). The next largest, in terms of number of traits, was a QTL in

the 1A5x1E4 cross on Chr 8 between 89.573- 346.224 Kb (QTL No. 13) that explained

variation in growth rate, colony radius and grey value in both control and KCl environments

(Figure 3, Table 3). This QTL on Chr 8 was also found to explain variation in response to

oxidative stress (Zhong et al., 2021). Another notable QTL in terms of high LOD (>10) was on

Chr 10 (749.521– 1302.645 Kb, QTL No. 10) in the 3D7x3D1 cross, which explained variation

in melanisation rate and colony radius in the control environment (Table 2). This QTL was

previously found to explain variation in growth-related traits at 10^0C (Lendenmann et al.,

2014) and under oxidative stress (Zhong et al., 2021).

3.5. A unique QTL associated with the KCl environment

Several novel QTL that were unique to the KCl environment were also found (Table 2, 3). In

the 3D7x3D1 cross, two QTL were found for traits related to colony radius: the first on Chr 3

(QTL No. 3) explained variation in colony radius at 8 dpi and colony radius KCl tolerance at 8

and 12 dpi; and the second on Chr 4 (QTL No. 5) explained variation in colony radius at 8 dpi

and colony radius KCl tolerance at 8 dpi. For melanin-related traits, four QTL were found in

the 3D7x3D1 cross: the first on Chr 1 (QTL No. 1) explained variation in three traits

(melanisation rate, melanisation rate KCl tolerance and grey value at 12 dpi); the second on

Chr 2 (QTL No. 2) explained variation in grey value at 12 dpi; the third on Chr 6 (QTL No. 6)

explained variation in melanisation rate; and the fourth on Chr 8 (QTL No. 8) explained

variation in grey value at 8 dpi. In the 1A5x1E4 cross a total of five QTL related to colony

radius in KCl were found. The first three were on Chr 1: QTL No. 2 explained variation in

colony radius KCl tolerance at 12 dpi; QTL No. 3 explained variation in growth rate KCl

tolerance; and QTL No. 4 explained variation in colony radius at 12 dpi. The fourth unique

KCl environment QTL was on Chr 5 (QTL No. 10) and explained variation in growth rate KCl

tolerance; the fifth on Chr 6 (QTL No. 11) explained variation in radius at 8 dpi

To identify candidate genes associated with KCl stress, we focused attention on novel QTL

with a high LOD (>10), relatively narrow intervals, and that explained a relatively high

proportion of the variation for a KCl-associated trait. Only QTL No. 3 in the 3D7x3D1 cross

met these requirements (Table 2). This QTL had a high LOD (>10) for colony radius at 8 dpi in

the KCl environment and colony radius KCl tolerance at 8 dpi. The LOD peaked at 22.8 at

chromosome position 737773 bp. This QTL explained 33% of the phenotypic variance in

colony radius at 8 dpi and 23% of the variance in colony radius KCl tolerance at 8 dpi.

Isolates with the non-reference allele at all markers within the interval had smaller colonies

(F=113.1, df=1,233, p=<0.001, Figure 4c) and less colony radius KCl tolerance at 8 dpi

(F=113.1, df=1,233, p=<0.001). The Bayes confidence interval for the QTL for colony radius

in KCl at 8 dpi spanned 149.29 Kb and contained 94 SNPs and 36 genes (Table 2, Figure 4b).

For this interval we present data on the genes in the reference genome (3D7), the

orthologous genes in 3D1, the gene sequence similarity (including ±200bp of flanking

regions) between the parents, the gene's putative encoded function (PFAM database) and if

there is any evidence of differential gene expression between the parents across all

available datasets (Table 4 and see Table S5 for full details).

3.6. Candidate genes within QTL No. 3 in 3D7x3D1

Several genes had SNP variation predicted to have a high impact based on SNPEff analysis

(Table S6, *ZT3D7_G3622*, *ZT3D7_G3623*, *ZT3D7_G3624, ZT3D7_G3632, ZT3D7_G3633,*

ZT3D7_G3635, ZT3D7_G3649). A transposable element (6927bp) was found within 200bp of

ZT3D7_G3623, and one gene, *ZT3D7_G3649* is a singleton – with no orthologue present in

any of the other 18 *Z. tritici* reference genome sequences (Table S5). Based on predicted

protein function, and sequence and gene expression variation between the parents, we

consider six genes to be the most promising candidates for further analyses: *ZT3D7_G3622*,

ZT3D7_G3635, *ZT3D7_G3637*, *ZT3D7_G3639*, *ZT3D7_G3655* and *ZT3D7_G3658* (this order

reflects their relative positions on chromosome 3).

ZT3D7_G3622 has a loss of function SNP (Chr 3:598092) in 3D1 that is predicted to cause a

loss of a predicted start codon. The gene contains a single exon, encodes a protein length of

550 amino acids (aa) and is predicted to contain a fungal-specific transcription factor

domain and Zn(2)-C6 fungal-type DNA-binding domain (Table 4). The orthologue in the 3D1

genome is *ZT3D1_G3567*. It encodes a longer protein (722 aa) and has two exons, the first

exon's product encodes a Zn(2)-C6 fungal-type DNA-binding domain and the second exon's

product is the same as that of *ZT3D7_G3622*. The predicted protein in 3D1 is the same as

the orthologue In IPO323 (*Mycgr3T108446*). Although we do have reads mapping to this

region, and when grown in vitro on MM this gene is downregulated in 3D1 compared to 3D7

(Table S5), our RNA sequence data does not clearly support gene models in either 3D1 or

3D7, there are no clear exon/intron boundaries. This suggests that the annotation of this

gene may require manual curation.

ZT3D7_G3635 has a high impact SNP variant (Chr 3:681906) that is predicted to cause a

frameshift. The gene has 6 exons in 3D7 and encodes a protein length of 1078 aa. Based on

INTERPROSCAN and Ensemble Fungi, the protein has multiple predicted domains; the

largest is a protein kinase domain. The protein also contains a Serine/threonine-protein

kinase active site and the molecular function of the first GO term (GO:0004674) is protein

serine/threonine kinase activity – suggesting that this gene is likely to be involved in osmotic

stress response (Hohmann, 2002). The RNA sequence data supports the gene model. The

orthologue in 3D1 is *ZT3D1_G3580*, which has 3 exons and multiple predicted domains that

do not differ from those predicted for *ZT3D7_G3635*. The gene is up-regulated in vitro on

YSB and down-regulated in planta at 28d in 3D1 compared to 3D7 (Table S5).

ZT3D7_G3637 has one exon, encodes a protein length of 1285 aa, and includes multiple

predicted domains including a DEAD/DEAH box helicase, which is involved in plant

responses to osmotic stress (Nguyen et al., 2018). The orthologue in 3D1 (*ZT3D1_G3583*)

encodes a similar protein as *ZT3D7_G3637* – has the same length and has the same

predicted domains. There are no high impact SNPs between the parents, but several

moderate and low impact variants were identified (Table S6). The RNA sequence data

supports the annotation, and the gene is down regulated in 3D1 in both YSB and MM (Table

S5).

ZT3D7_G3639 has 4 exons in 3D7. The gene sequence similarity between 3D7 and 3D1 is

98.66% and there is an 8bp intronic deletion in 3D1. This gene belongs to the Major

Facilitator Superfamily (MFS) and encodes a protein length of 512 aa. According to the

Phobius analysis, the gene contains 12 transmembrane segments (Table S5). Proteins with

transmembrane activity and transmembrane segments are likely to play an important role

in sensing and regulating osmotic stress (Hohmann, 2002). The orthologue in 3D1 is

ZT3D1_G3580, which has the same number of exons, encodes the same protein length and

has the same predicted function. The RNA sequence data supports the gene model, and the

differential expression analysis found the gene is downregulated in planta at 12 and 14 days

in 3D1 compared to 3D7 (Table S5).

ZT3D7_G3655 has one exon and encodes a protein length of 848 aa. The gene contains 10

transmembrane segments (Table S5) and encodes a voltage-gated chloride channel domain

– thus it is likely to be involved in osmotic stress responses. The gene sequence between the

parents is almost identical, but there is one moderate effect SNP within the gene (Table S6).

When grown in vitro on MM this gene is upregulated in 3D1 compared to 3D7. The RNA

sequence data supports the gene model annotation.

ZT3D7_G3658 has four exons and encodes a protein length of 614 aa. Its orthologue in 3D1

(*ZT3D1_G3604*) has three exons and encodes 572 aa. The gene encodes a vacuolar protein

sorting-associated protein 17, Vps17. Vacuoles play an important role in osmotic stress

responses: their fission and fusion enable cells to maintain turgor pressure and regulate cell

volume (Richards et al., 2010). Comparison of the 3D1 gene sequence to the 3D1 genome

using BLAST revealed that the 3D7 gene sequence was identical, but the 3D1 orthologue

annotation is shorter and has only three exons. Our RNA sequence data supports the 3D1

annotation with only 3 exons, as there were no RNA sequence reads mapping to exon 1 of

the *ZT3D7_G36580*. This gene was downregulated in vitro in both MM and YSB in 3D1

compared to 3D7.

4. Discussion

We were able to map major effect loci affecting colony growth and melanisation in the

presence of potassium chloride in both crosses. We focussed on QTL that were unique to

the KCl environment to identify candidate genes that are specific to osmotic stress – as

opposed to finding genes that may be more generally related to intrinsic growth or melanin

production. A QTL on Chr 3 in the 3D7x3D1 cross was the most promising for identifying

genes specifically associated with KCl stress. We highlighted the six most promising

candidate genes in this QTL based on their predicted encoded functions, sequence similarity

and transcriptional profiles. We also identified several QTL shared across environments and

confirmed that several previously identified QTL, which explained variation in growth and

melanisation across different environmental stressors (Table 2 and 3, e.g. 3D7x3D1 Chr 11

and 1A5x1E4 Chr 8), also explained trait variation in the presence of osmotic stress.

4.1. Effect of KCl on colonies

In the presence of potassium chloride at 0.75M, colonies were typically smaller and less

melanised compared to the control environment (Figure 1, Table S2). A reduction in colony

radius in the KCl environment supports our hypothesis that this environment was stressful

to *Z. tritici*. Our results suggest that growth rate in the 1A5x1E4 cross was less affected by

KCl-induced osmotic stress, as mean growth rate was not reduced in the KCl environment,

compared to the 3D7x3D1 cross that exhibited a significantly reduced growth rate (Table

S2). However, we note that the overall difference in growth rate in the 3D7x3D1 progeny

between the control and KCl environments was relatively small (~6% reduction in growth,

Table S2).

When experiencing osmotic stress, an increase in melanin is thought to protect cells by

strengthening the fungal cell wall, but a reduced metabolic capacity under stress may limit

the ability of a cell to produce melanin. The mean colony grey values were higher, indicative

of lower levels of melanin, for strains grown in the KCl environment (Table S2). This suggests

that the *Z. tritici* response to osmotic stress did not involve an increased production of

melanin to protect the cells as was observed in halotolerant species (Kogej et al., 2007). We

did observe however, that isolates that were more melanised in the benign control

environment (lower Grey Value) were more tolerant to salt stress (Figure S2) – their growth

rates were higher and their reductions in melanisation were lower compared to isolates that

were less melanised in the control environment (Table S3, S4). This suggests that strains

having a greater intrinsic capacity to become melanised are more tolerant to salt stress.

4.2. Shared QTL

Some QTL that explained variation in growth and melanisation in the stressful environment

overlapped with QTL for trait variation in the control environment. Prominent examples

include the Chr 11 QTL No. 11 in the 3D7x3D1 cross and the Chr 8 QTL No. 13 in the

1A5x1E4 cross. These genomic regions may harbour genes that affect intrinsic growth of

colonies under a wide range of environments and/or genes with pleiotropic effects. Many of

the QTL shared across environments were identified in earlier experiments and discussed

previously (Lendenmann et al., 2015; Zhong et al., 2021). The QTL No. 11 was shown to be

due to differential expression of the *Zmr1* gene encoding a transcription factor that

regulates synthesis of DHN melanin in *Z. tritici* (Krishnan et al. 2018). *Zmr1* was shown to

have pleiotropic effects on melanin production, colony growth rates and fungicide

resistance (Krishnan et al. 2018).

4.3. QTL uniquely associated with the KCl environment

This QTL analysis enabled us to identity large effect loci that explain variation in traits under

KCl stress. Among the QTL uniquely associated with KCl stress, most had moderate LODs

with each QTL explaining only a small proportion of the phenotypic variance (3D7x3D1

maximum = 0.10, 1A5x1E4 maximum = 0.11). The exception was QTL No. 3 in the 3D7x3D1

cross, which explained up to 33% of the phenotypic variance. This QTL on chromosome 3

was associated with differences in colony radius at 8 dpi and colony radius KCl tolerance at 8

and 12 dpi. We targeted this QTL for further investigation, seeking to identify genes

specifically associated with KCl tolerance. It is interesting to note that no LOD peak was

detected for colony radius at 12 dpi, even though colony radius at 12 dpi was correlated

with colony radius at 8 dpi (Table S3, correlation = 0.58, p<0.001). As growth rate is based

on the change in colony radius between 8 and 12 dpi, we might expect growth rate in KCl to

share the QTL for colony radius at 8 dpi. To better understand why growth rate does not

share this QTL, we examined the trait correlations. Growth rate is highly correlated with

colony radius at 12 dpi (0.93) and less strongly correlated with colony radius at 8 dpi (0.26)

(Table S3). There is less variation in colony radius at 8 dpi, thus colony radius at 12 dpi has a

greater impact on the growth rate. The QTL No. 3 also explained variation in colony radius

KCl tolerance at 8 and 12 dpi, although the association with this trait at 12 dpi explained

relatively little phenotypic variation (0.08). Thus, the QTL peak is more strongly associated

with colony radius and colony radius KCl tolerance at an earlier stage of colony growth. A

similar pattern was observed in the chromosome 11 QTL associated with melanisation

(Krishnan et al., 2018; Lendenmann et al., 2014), with the QTL LOD score very high in the

younger colonies and decreasing as colonies aged. After cloning and characterizing the *Zmr1*

gene and its functions, this pattern could be explained based on the finding that the

difference in *Zmr1* expression (and the resulting difference in melanisation) was significantly

higher in younger colonies than in older colonies (Krishnan et al., 2018). We hypothesize

that QTL no. 3 associated with KCl tolerance is underpinned by a similar mechanism that is

specific to the earlier stage of colony growth.

4.4. Candidate genes in the QTL for colony radius and colony radius KCl tolerance at 8 dpi

in the 3D7x3D1 cross

Among the 36 genes found in the interval spanning this QTL No. 3, we consider six genes to

be the most promising candidates to be involved in sensing and responding to salt stress

and responsible for this QTL. Below we consider each in turn in the order of their relative

positions on chromosome 3.

The first candidate gene, *ZT3D7_G3622*, is predicted to encode a fungal-specific

transcription factor. The orthologue in the IPO323 reference genome (assembly MYCGR

v2.0) is *Mycgr3T108446* and it has 21 paralogs. Transcription factors (TFs) regulate gene

expression and are important regulators of molecular responses to environmental stimuli

(e.g. Hohmann, 2002; Shi et al., 2021), making this gene a good candidate. The *Zmr1* gene

identified in this same cross and shown to be responsible for differences in melanisation,

with greater effects in younger colonies, is a transcription factor (Krishnan et al. 2018). The

SNP variant in 3D1 (Chr 3:598092) is predicted to cause a loss of the start codon that is likely

to cause a loss of function. In *F. graminearum,* mutants lacking a similar TF (called *Fss1*)

were sensitive to high sodium and lithium, but they were not sensitive to elevated

potassium (Son et al., 2015).

The second candidate, *ZT3D7_G3635* is predicted to encode a serine/threonine-protein

kinase active site, which is known to regulate MAP kinases (Hohmann, 2002; Li et al., 2019).

A SNP variant found in the 3D1 allele is predicted to cause a frameshift that is likely to cause

a loss of function. The ortholog in the IPO323 genome is *Mycgr3G56270* and it has one

paralog in the reference genome. A serine/threonine kinase from *Aspergillus flavus*,

AflSte20, was recently functionally validated using a knockout mutant. This study showed

that the knockout had higher sensitivity to osmotic stress compared to the wild type (Li et

al., 2019), confirming the role of this serine/threonine-protein kinase in osmotic stress

response in a different fungal plant pathogen.

The third candidate, *ZT3D7_G3637* is predicted to encode a DEAD/DEAH box helicase

domain. The reference genome orthologue is *Mycgr3G99415* and it has 6 paralogs. The

DEAD – box helicases are the largest RNA helicase (RH) subfamily. In plants they play

multiple functional roles and are important in stress responses (Pandey et al., 2020). For

example, in *Arabidopsis thaliana AtRH17* is involved in salt-stress tolerance (Nguyen et al.,

2018), in rice *OsABP* is upregulated in response to salt (Macovei et al., 2012) and in barley

HvD1 accumulated in salt- and cold-stressed plants (Nakamura et al., 2004). In wheat, the

DEAD-box helicase *TaRH1* that was first identified as a component of the defence response

to stripe rust (*Puccinia striiformis f*. sp. *tritici*) was also found to be involved in plant

response to multiple stressors, including salt stress (Zhang et al., 2014). SNP variation found

between the parents was predicted to have only moderate effects, no high impact SNPs

were detected. We chose this gene as a promising candidate based mainly on its known

associations with salt stress in plants and the significant differences in expression found

between 3D1 and 3D7 while growing in vitro.

A gene encoding a Major Facilitator Superfamily (MFS) domain, *ZT3D7_G3639* is the fourth

candidate we highlight. The reference genome orthologue is *Mycgr3G85019* and it has 58

paralogs. MFS transporters facilitate the movement of small solutes across cell membranes

in response to chemiosmotic gradients – making them likely to play an important role in

sensing and regulating osmotic stress (Hohmann, 2002). We chose this gene as a promising

candidate based mainly on its known functions and associations with multiple stressors

including fungicides (Dos Santos et al., 2014; Omrane et al., 2015).

The fifth highlighted gene is *ZT3D7_G3655* that encodes a voltage gated chloride channel

domain and membrane transport domains. The reference genome orthologue is

Mycgr3T20761 and there are three paralogues in the IPO323 reference genome. Chloride

channels are important in animal cell volume regulation (Jentsch et al., 2002), fungal

oxidative stress response (Oddon et al., 2007), fungal pathogen virulence (Cañero and

Roncero, 2008; Zhu and Williamson, 2003) and plant immunity (Guo et al., 2014). The role of

chloride channels in response to salt stress is well established in animals (Jentsch et al.,

2002), and to a lesser degree in plants (Barbier-Brygoo et al., 2011), but less is known from

fungi. For one of the better-known families of chloride channels, the CLC family, a single

gene (*GEF1*) has been identified in yeast. In two pathogenic fungi (*Cryptococcus neoformans*

and *Fusarium oxysporum*) – the GEF1 homologue has been shown to influence virulence

(Cañero and Roncero, 2008; Zhu and Williamson, 2003). The gene sequence between the

parents is highly conserved, however gene expression differences were observed in minimal

media, so differences may exist in the regulatory sequence.

The final candidate we highlight is *ZT3D7_G3658*, which encodes a Vacuolar associated

protein 17, *Vps17*. The orthologous gene in the reference genome is *Mycgr3T108464* and

there are six paralogues. Comparing the DNA sequence of this gene against the 3D1 genome

revealed an identical sequence, but the annotation of the 3D7 and 3D1 differs – the 3D1

orthologue protein length is shorter and the gene has one fewer exons. Our RNA sequence

data suggests that the *ZT3D7_G3658* annotation may be incorrect - there were no reads

mapping to the first, small exon of this gene. This suggests that the annotation of this gene

may need manual curation before conducting further analyses. The gene was

downregulated in YSB and MM in 3D1 compared to 3D7, demonstrating that transcription

of this gene varies between the parents. Vacuoles play important roles in both osmotic

stress responses (Richards et al., 2010) and virulence (Li et al., 2021). Under salt stress the

vacuoles can undergo fusion and fission – changing in size and number (Richards et al.,

2010). *Vsp17* knockouts in *S. cerevisiae* had many small vacuoles but were resistant to

osmotic stress and could sporulate (Köhrer and Emr, 1993). We are unaware of any work

that has investigated the role of *Vsp17* in salt-stress responses of filamentous fungi.

4.5. QTL reproducibility

This dataset allowed us to explore the repeatability of previously identified QTL from other

published datasets. One earlier dataset used the same mapping populations but different

experimental conditions and treatments, with a different reference genome and different

genetic markers and map information (Lendenmann et al., 2015, 2014). A second earlier

dataset used the same isolates and experimental conditions, but different treatments and

different marker and map information (Zhong et al., 2021). Comparisons across these three

data sets yielded several consistent QTL. Our analyses confirmed that a region on

chromosome 11 in the 3D7x3D1 cross (Table 2, QTL No. 11) is associated with variation in

melanisation and growth in multiple environments (e.g. under fungicide stress in

Lendenmann et al., 2015; under oxidative stress in Zhong et al., 2021). This QTL contains the

Zmr1 transcription factor that was shown to regulate melanin production (Krishnan et al.,

2018). We hypothesize that *Zmr1* also explains the variation we observed under salt-stress.

A QTL associated with multiple growth and melanin traits in the 1A5x1E4 cross on

chromosome 8 (Table 3, QTL No. 13) overlaps with a previously identified QTL on

chromosome 8 that was associated with colony growth and tolerance to oxidative stress

(Zhong et al., 2021). These findings illustrate that many of our QTL are reproducible across

experiments and increases our confidence that we have identified biologically meaningful

genomic regions using our automated image analyses.

CRediT authorship contribution statement

Jessica Stapley: Formal analysis, Data curation, Writing - Original Draft, Writing - Review &

Editing, Resources, Data curation. Bruce A. McDonald: Conceptualization, Methodology,

Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper

Data availability

Phenotypic and SNP data are available at ETH Data repository (DOI: 10.3929/ethz-b-

000550424). All code and details of the analysis are available at

(https://github.com/jessstapley/QTL-mapping-Z.-tritici).

Acknowledgments

We thank Ziming Zhong, Tiziana Valeria Vonlanthen, Bethan Turnbull, Susanne Dora, Sarah

Furler, Alexandra Waltenspühl, and Jasmin Wiedmer for collecting the phenotypic data and

Javier Palma-Guerrero for supervision. We thank Marcello Zala for assistance in the

laboratory. This research was supported by the Swiss National Science Foundation

(31003A_155955 granted to BAM).

References

Araújo, C.A.S., Ferreira, P.C., Pupin, B., Dias, L.P., Avalos, J., Edwards, J., Hallsworth, J.E., Rangel, D.E.N., 2020. Osmotolerance as a determinant of microbial ecology: A study of phylogenetically diverse fungi. Fungal Biol. 124, 273–288. https://doi.org/10.1016/j.funbio.2019.09.001

Badet, T., Oggenfuss, U., Abraham, L., McDonald, B.A., Croll, D., 2020. A 19-isolate reference-quality global pangenome for the fungal wheat pathogen *Zymoseptoria tritici*. BMC Biol. 18, 12. https://doi.org/10.1186/s12915-020-0744-3

Barbier-Brygoo, H., De Angeli, A., Filleur, S., Frachisse, J.-M., Gambale, F., Thomine, S., Wege, S., 2011. Anion channels/transporters in plants: from molecular bases to regulatory networks. Annu. Rev. Plant Biol. 62, 25–51. https://doi.org/10.1146/annurev-arplant-042110-103741

Broman, K.W., Wu, H., Sen, S., Churchill, G.A., 2003. R/qtl: QTL mapping in experimental crosses. Bioinformatics 19, 889–890. https://doi.org/10.1093/bioinformatics/btg112

Cañero, D.C., Roncero, M.I.G., 2008. Influence of the chloride channel of *Fusarium oxysporum* on extracellular laccase activity and virulence on tomato plants. Microbiology 154, 1474–1481. https://doi.org/10.1099/mic.0.2007/015388-0

Croll, D., Zala, M., McDonald, B.A., 2013. Breakage-fusion-bridge Cycles and Large Insertions Contribute to the Rapid Evolution of Accessory Chromosomes in a Fungal Pathogen. PLOS Genet. 9, e1003567. https://doi.org/10.1371/journal.pgen.1003567

Dos Santos, S.C., Teixeira, M.C., Dias, P.J., Sá-Correia, I., 2014. MFS transporters required for multidrug/multixenobiotic (MD/MX) resistance in the model yeast: understanding their physiological function through post-genomic approaches. Front. Physiol. 5, 180–180. https://doi.org/10.3389/fphys.2014.00180

Duran, R., Cary, J.W., Calvo, A.M., 2010. Role of the Osmotic Stress Regulatory Pathway in Morphogenesis and Secondary Metabolism in Filamentous Fungi. Toxins 2, 367–381. https://doi.org/10.3390/toxins2040367

Francisco, C.S., Ma, X., Zwyssig, M.M., McDonald, B.A., Palma-Guerrero, J., 2019. Morphological changes in response to environmental stresses in the fungal plant pathogen *Zymoseptoria tritici*. Sci. Rep. 9, 9642. https://doi.org/10.1038/s41598-019-45994-3

Francisco, C.S., McDonald, B.A., Palma-Guerrero, J., 2022. A transcription factor and a phosphatase regulate temperature-dependent morphogenesis in a fungal plant pathogen (preprint). Molecular Biology. https://doi.org/10.1101/2022.02.07.479299

Guo, W., Zuo, Z., Cheng, X., Sun, J., Li, H., Li, L., Qiu, J.-L., 2014. The chloride channel family gene CLCd negatively regulates pathogen-associated molecular pattern (PAMP)-triggered immunity in Arabidopsis. J. Exp. Bot. 65, 1205–1215. https://doi.org/10.1093/jxb/ert484

Hohmann, S., 2002. Osmotic Stress Signaling and Osmoadaptation in Yeasts. Microbiol. Mol. Biol. Rev. 66, 300–372. https://doi.org/10.1128/MMBR.66.2.300-372.2002

Jacob, S., Foster, A.J., Yemelin, A., Thines, E., 2015. High osmolarity glycerol (HOG) signalling in *Magnaporthe oryzae*: Identification of MoYPD1 and its role in osmoregulation, fungicide action, and pathogenicity. Fungal Biol. 119, 580–594. https://doi.org/10.1016/j.funbio.2015.03.003

Jentsch, T.J., Stein, V., Weinreich, F., Zdebik, A.A., 2002. Molecular Structure and Physiological Function of Chloride Channels. Physiol. Rev. 82, 503–568. https://doi.org/10.1152/physrev.00029.2001

Jiang, C., Lv, G., Ge, J., He, B., Zhang, Z., Hu, Z., Zeng, B., 2021. Genome-wide identification
of the GATA transcription factor family and their expression patterns under
temperature and salt stress in *Aspergillus oryzae*. AMB Express 11, 56.
https://doi.org/10.1186/s13568-021-01212-w
Kogej, T., Stein, M., Volkmann, M., Gorbushina, A.A., Galinski, E.A., Gunde-Cimerman, N.,
2007. Osmotic adaptation of the halophilic fungus *Hortaea werneckii*: role of
osmolytes and melanization. Microbiology 153, 4261–4273.
https://doi.org/10.1099/mic.0.2007/010751-0
Köhrer, K., Emr, S., 1993. The yeast VPS17 gene encodes a membrane-associated protein
required for the sorting of soluble vacuolar hydrolases. J. Biol. Chem. 268, 559—569.
https://doi.org/10.1016/s0021-9258(18)54188-0
Krishnan, P., Meile, L., Plissonneau, C., Ma, X., Hartmann, F.E., Croll, D., McDonald, B.A.,
Sánchez-Vallet, A., 2018. Transposable element insertions shape gene regulation and
melanin production in a fungal pathogen of wheat. BMC Biol. 16, 78.
https://doi.org/10.1186/s12915-018-0543-2
Lendenmann, M.H., Croll, D., McDonald, B.A., 2015. QTL mapping of fungicide sensitivity
reveals novel genes and pleiotropy with melanization in the pathogen *Zymoseptoria*
tritici. Fungal Genet. Biol. 80, 53–67. https://doi.org/10.1016/j.fgb.2015.05.001
Lendenmann, M.H., Croll, D., Stewart, E.L., McDonald, B.A., 2014. Quantitative Trait Locus
Mapping of Melanization in the Plant Pathogenic Fungus *Zymoseptoria tritici*. G3
GenesGenomesGenetics 4, 2519–2533. https://doi.org/10.1534/g3.114.015289
Li, D., Qin, L., Wang, Y., Xie, Q., Li, N., Wang, S., Yuan, J., 2019. AflSte20 Regulates
Morphogenesis, Stress Response, and Aflatoxin Biosynthesis of *Aspergillus flavus*.
Toxins 11, 730. https://doi.org/10.3390/toxins11120730
Li, S., Zhang, S., Li, B., Li, H., 2021. The SNARE Protein CfVam7 Is Required for Growth,
Endoplasmic Reticulum Stress Response, and Pathogenicity of Colletotrichum
fructicola. Front. Microbiol. 12, 736066. https://doi.org/10.3389/fmicb.2021.736066
Liao, Y., Smyth, G.K., Shi, W., 2019. The R package Rsubread is easier, faster, cheaper and
better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res.
47, e47–e47. https://doi.org/10.1093/nar/gkz114
Macovei, A., Vaid, N., Tula, S., Tuteja, N., 2012. A new DEAD-box helicase ATP-binding
protein (OsABP) from rice is responsive to abiotic stress. Plant Signal. Behav. 7,
1138–1143. https://doi.org/10.4161/psb.21343
Malosetti, M., van Eeuwijk, F.A., Boer, M.P., Casas, A.M., Elía, M., Moralejo, M., Bhat, P.R.,
Ramsay, L., Molina-Cano, J.-L., 2011. Gene and QTL detection in a three-way barley
cross under selection by a mixed model with kinship information using SNPs. Theor.
Appl. Genet. 122, 1605–1616. https://doi.org/10.1007/s00122-011-1558-z
Nakamura, T., Muramoto, Y., Yokota, S., Ueda, A., Takabe, T., 2004. Structural and
transcriptional characterization of a salt-responsive gene encoding putative ATP-
dependent RNA helicase in barley. Plant Sci. 167, 63–70.
https://doi.org/10.1016/j.plantsci.2004.03.001
Nguyen, L., Seok, H.-Y., Woo, D.-H., Lee, S.-Y., Moon, Y.-H., 2018. Overexpression of the
DEAD-Box RNA Helicase Gene AtRH17 Confers Tolerance to Salt Stress in
Arabidopsis. Int. J. Mol. Sci. 19, 3777. https://doi.org/10.3390/ijms19123777
Oddon, D.M., Diatloff, E., Roberts, S.K., 2007. A CLC chloride channel plays an essential role
in copper homeostasis in *Aspergillus nidulans* at increased extracellular copper

concentrations. Biochim. Biophys. Acta BBA - Biomembr. 1768, 2466–2477.
https://doi.org/10.1016/j.bbamem.2007.05.022

Omrane, S., Sghyer, H., Audéon, C., Lanen, C., Duplaix, C., Walker, A.-S., Fillinger, S., 2015.
Fungicide efflux and the MgMFS1 transporter contribute to the multidrug resistance
phenotype in *Z ymoseptoria tritici* field isolates: Fungicide efflux & *MgMFS 1*
contribute to MDR in *Z. tritici*. Environ. Microbiol. 17, 2805–2823.
https://doi.org/10.1111/1462-2920.12781

Palma-Guerrero, J., Ma, X., Torriani, S.F.F., Zala, M., Francisco, C.S., Hartmann, F.E., Croll, D.,
McDonald, B.A., 2017. Comparative Transcriptome Analyses in *Zymoseptoria tritici*
Reveal Significant Differences in Gene Expression Among Strains During Plant
Infection. Mol. Plant-Microbe Interactions® 30, 231–244.
https://doi.org/10.1094/MPMI-07-16-0146-R

Pandey, S., Prasad, A., Sharma, N., Prasad, M., 2020. Linking the plant stress responses with
RNA helicases. Plant Sci. 299, 110607.
https://doi.org/10.1016/j.plantsci.2020.110607

Plissonneau, C., Hartmann, F.E., Croll, D., 2018. Pangenome analyses of the wheat pathogen
Zymoseptoria tritici reveal the structural basis of a highly plastic eukaryotic genome.
BMC Biol. 16, 5. https://doi.org/10.1186/s12915-017-0457-4

Plissonneau, C., Stürchler, A., Croll, D., 2016. The Evolution of Orphan Regions in Genomes
of a Fungal Pathogen of Wheat. mBio 7. https://doi.org/10.1128/mBio.01231-16

R Core Team, 2020. R: A language and environment for statistical computing.

Richards, A., Veses, V., Gow, N.A.R., 2010. Vacuole dynamics in fungi. Fungal Biol. Rev. 24,
93–105. https://doi.org/10.1016/j.fbr.2010.04.002

Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26,
139–140. https://doi.org/10.1093/bioinformatics/btp616

RStudio Team, 2020. RStudio: Integrated Development Environment for R.

Shi, D., Ren, W., Wang, J., Zhang, J., Mbadianya, J.I., Mao, X., Chen, C., 2021. The
transcription factor FgNsf1 regulates fungal development, virulence and stress
responses in Fusarium graminearum. J. Integr. Agric. 20, 2156–2169.
https://doi.org/10.1016/S2095-3119(20)63339-1

Son, H., Park, A.R., Lim, J.Y., Lee, Y.-W., 2015. Fss1 is involved in the regulation of an *ENA5*
homologue for sodium and lithium tolerance in *F usarium graminearum*: Roles of
Fss1 in *Fusarium graminearum*. Environ. Microbiol. 17, 2048–2063.
https://doi.org/10.1111/1462-2920.12757

Zhan, J., Linde, C.C., Jurgens, T., Merz, U., Steinebrunner, F., McDONALD, B.A., 2005.
Variation for neutral markers is correlated with variation for quantitative traits in the
plant pathogenic fungus Mycosphaerella graminicola. Mol. Ecol. 14, 2683–2693.
https://doi.org/10.1111/j.1365-294X.2005.02638.x

Zhang, X., Zhao, X., Feng, C., Liu, N., Feng, H., Wang, X., Mu, X., Huang, L., Kang, Z., 2014. The
cloning and characterization of a DEAD-Box RNA helicase from stress-responsive
wheat. Physiol. Mol. Plant Pathol. 88, 36–42.
https://doi.org/10.1016/j.pmpp.2014.07.004

Zhong, Z., Marcel, T.C., Hartmann, F.E., Ma, X., Plissonneau, C., Zala, M., Ducasse, A.,
Confais, J., Compain, J., Lapalu, N., Amselem, J., McDonald, B.A., Croll, D., Palma-
Guerrero, J., 2017. A small secreted protein in *Zymoseptoria tritici* is responsible for

avirulence on wheat cultivars carrying the Stb6 resistance gene. New Phytol. 214,
619–631. https://doi.org/10.1111/nph.14434
Zhong, Z., McDonald, B.A., Palma-Guerrero, J., 2021. Tolerance to oxidative stress is
associated with both oxidative stress response and inherent growth in a fungal
wheat pathogen. Genetics 217, iyaa022. https://doi.org/10.1093/genetics/iyaa022
Zhu, X., Williamson, P.R., 2003. A CLC-type chloride channel gene is required for laccase
activity and virulence in *Cryptococcus neoformans*: CLC-A of *Cryptococcus*
neoformans. Mol. Microbiol. 50, 1271–1281. https://doi.org/10.1046/j.1365-
2958.2003.03752.x
791
792

Figure 1. Trait variation in the control and salt stress environments across offspring in each

cross. Top panel: Points are mean colony radius in the control and KCl environment at 8 and

12 dpi, smoothed lines demonstrate linear change over time. Middle panel: Points are mean

grey value in the control and KCL environments at 8 and 12 dpi, smoothed lines

demonstrate linear change over time. Bottom panel: Growth rate KCl tolerance and

melanisation rate KCl tolerance (KCl tolerance = rate in KCl/rate in control conditions, see

section 2.4) between the two crosses.

Figure 2. LOD plots from interval mapping for representative traits a) growth rate and b)

melanisation rate in the control environment (grey) and in the KCl environment (orange) in

the 3D7x3D1 cross. Each sub-panel is a chromosome with the chromosome number

indicated at the top. Some numbers are missing because *Zymoseptoria tritici* carries eight

accessory chromosomes (Chr 14-21) that show presence/absence polymorphisms among

isolates. The dashed horizontal lines indicate the LOD significance thresholds for each trait

based on permutation. The numbers above the QTL peaks correspond to the QTL numbers

from Table 2. For LOD plots of all traits see Supplementary Information (S2).

Figure 3. LOD plots from interval mapping for representative traits a) growth rate and b)

melanisation rate in the control environment (grey) and in the KCl environment (orange) in

the 1A5x1E4 cross. Some numbers are missing because *Zymoseptoria tritici* carries eight

accessory chromosomes (Chr 14-21) that show presence/absence polymorphisms among

isolates. The dashed horizontal lines indicate the LOD significance thresholds for each trait

based on permutation. The numbers above the QTL peaks correspond to the QTL numbers

from Table 3. For LOD plots of all traits see Supplementary Information (S3) .

Figure 4. QTL on Chromosome 3 for three traits related to growth under osmotic stress in

the 3D7x3D1 cross. The three traits were colony radius at 8 dpi, colony radius KCl tolerance

at 8 and 12 dpi (KCl tolerance = colony radius in KCl/colony radius in control conditions , see

section 2.4). a) LOD peak and Bayes credible interval (shaded) from interval mapping, with

SNP positions indicated on the x-axis as black vertical lines, b) Alignment of the 36 genes

within the QTL interval between the 3D7 and 3D1 parental strains – orange-coloured genes

are the six most promising candidate genes described in the main text and c) allelic effects

825 of 94 SNPs within the QTL interval on mean colony radius in KCl at 8 dpi. Isolates carrying

826 the reference allele (ref=3D7) shown in grey have smaller colonies compared to the

827 alternative allele (alt=3D1) shown in orange. SNP position in base pairs is shown on the x-

828 axis. The marker with the highest LOD is at position 737773 bp.

830 Supplementary Material

831 S1. Summary of genetic linkage maps.

832 S2. Plots of phenotypic correlations.

833 S3. LOD plots from interval mapping for all traits in the 3D7x3D1 cross.

834 S4. LOD plots from interval mapping for all traits in the 1A5x1E4 cross.

Table 1. Summary of the QTL mapping results for all traits. Grey value and colony radius were measured at two time points: 8 and 12 dpi. Growth rate was measured as the change in colony radius/number of days(4). Melanisation rate was measured as the change in grey value/number of days(4). Colony radius/grey value KCl tolerance were calculated by dividing the value (grey value or colony radius) in the control environment by the value in the KCl environment. Growth rate/Melanisation KCl tolerance were calculated by dividing the rate (melanisation or growth) in the control environment by the rate in the KCl environment. The QTL peak columns show the chromosomal position of significant QTL peak for each trait in the 3D7x3D1 and 1A5x1E4 crosses respectively, and the final column contained estimated heritability (h^2) for each trait.

Data Type	Environment	DPI	Measurement	Measurement	QTL peak 3D7x3D1	h^2	QTL peak 1A5x1E4	h^2
Time point	Control	8	Colony radius	Colony radius in Control at 8dpi	3,8,10,11	0.53	2,3,8	0.56
Time point	Control	12	Colony radius	Colony radius in Control at 12dpi	3,7,8,10,11	0.62	7,8	0.48
Time point	KCl	8	Colony radius	Colony radius in KCl at 8dpi	3,4	0.64	6,	0.5
Time point	KCl	12	Colony radius	Colony radius in KCl at 12dpi	3,8	0.6	1,8	0.52
Time point	Control	8	Grey value	Grey value in Control at 8dpi	8,10,11	0.42	1,2,3,8	0.63
Time point	Control	12	Grey value	Grey value in Control at 12dpi	11	0.22	2,3,8	0.44
Time point	KCl	8	Grey value	Grey value in KCl at 8dpi	8,11	0.54	8	0.34
Time point	KCl	12	Grey value	Grey value in KCl at 12dpi	1,2,11	0.34	No Sig. QTL	0.29
Rate	Control	8-12	Growth rate	Growth rate in Control	3,8,11	0.61	7,8	0.33
Rate	KCl	8-12	Growth rate	Growth rate in KCl	3,8,11	0.45	8,	0.49
Rate	Control	8-12	Melanisation rate	Melanisation rate in Control	10	0.28	2,3,12	0.54
Rate	KCl	8-12	Melanisation rate	Melanisation rate in KCl	1,6	0.32	No Sig. QTL	0.27
Tolerance	Both	8	Grey value	Grey value KCl tolerance at 8 dpi	8,10	0.37	1,2,3,8	0.58
Tolerance	Both	12	Grey value	Grey value KCl tolerance at 12 dpi	11	0.18	No Sig. QTL	0.34
Tolerance	Both	8	Colony radius	Colony radius KCl tolerance at 8dpi	3,4,10	0.59	8	0.42
Tolerance	Both	12	Colony radius	Colony radius KCl tolerance at 12dpi	3,10,11	0.47	1	0.37
Tolerance	Both	8-12	Growth rate	Growth rate KCl tolerance	11	0.4	1,5	0.45
Tolerance	Both	8-12	Melanisation rate	Melanisation rate KCl tolerance	No Sig. QTL	0.14	1	0.32

Table 2. Description of all QTL identified from the 3D7x3D1 cross, including map and physical position of the LOD peak marker, the bayes credible interval length and the number of genes within the interval. PVE is the proportion of variance explained by the QTL. QTL number is a number assigned to a particular QTL peak which groups overlapping QTLs.

Type	Chr	Pos (cM)	LOD	Pos (bp)	No. markers	Interval Start (bp)	Interval end (bp)	Interval size (bp)	No. Genes	PVE	QTL number
Melanisation rate in KCl	1	457.43	6.18	1590235	2915	1035571	2769729	1734.158	598	0.10	1
Melanisation rate KCl tolerance	1	540.00	3.40	2295356	9000	788430	6389989	5601.559	1977	0.06	1
Grey value in KCl at 12dpi	1	576.00	5.99	2717234	519	2295128	2769729	474.601	174	0.10	1
Grey value in KCl at 12dpi	2	261.00	5.07	691536	141	679366	711981	32.615	15	0.09	2
Colony radius KCl tolerance at 12dpi	3	142.92	4.49	719549	553	396401	944712	548.311	125	0.08	3
Colony radius KCl tolerance at 8dpi	3	144.00	14.62	737773	105	588483	751983	163.5	36	0.23	3
Colony radius in KCl at 8dpi	3	144.15	22.83	737773	94	588483	737773	149.29	31	0.33	3
Growth rate in Control	3	415.00	5.31	2537296	2204	1982744	3347765	1365.021	462	0.09	4
Colony radius in Control at 12dpi	3	456.63	8.11	2993469	427	2971359	3196103	224.744	79	0.13	4
Colony radius in Control at 8dpi	3	482.67	7.09	3056536	428	2982086	3199430	217.344	76	0.12	4
Growth rate in KCl	3	485.65	4.93	3110053	845	2971359	3347765	376.406	132	0.08	4
Colony radius in KCl at 12dpi	3	485.65	4.99	3110053	3084	516017	3203947	2687.93	813	0.08	4
Colony radius KCl tolerance at 8dpi	4	400.00	4.28	1936846	1000	1845763	2448166	602.403	200	0.07	5
Colony radius in KCl at 8dpi	4	403.76	4.37	2013908	986	1867509	2448166	580.657	194	0.07	5
Melanisation rate in KCl	6	220.14	6.24	1767543	479	1568163	1981193	413.03	124	0.10	6
Colony radius in Control at 12dpi	7	186.78	4.47	1109088	1366	511101	1377451	866.35	265	0.08	7
Grey value in KCl at 8dpi	8	680.87	5.69	943669	732	502305	974947	472.642	123	0.10	8
Growth rate in Control	8	746.72	7.31	1100862	542	1071962	1221280	149.318	65	0.12	9
Colony radius in Control at 12dpi	8	746.72	10.77	1100862	274	1091579	1164224	72.645	29	0.17	9
Growth rate in KCl	8	747.72	9.66	1100862	667	1091692	1284370	192.678	77	0.16	9
Colony radius in KCl at 12dpi	8	751.61	9.79	1112633	670	1100862	1292397	191.535	76	0.16	9
Colony radius in Control at 8dpi	8	765.72	8.04	1221337	751	1091475	1354349	262.874	108	0.13	9
Grey value KCl tolerance at 8dpi	8	776.11	4.22	1350673	2608	962119	2165894	1203.775	457	0.07	9

Grey value in Control at 8dpi	8	776.11	4.35	1350706	2549	981398	2175005	1193.607	450	0.07	9
Grey value in Control at 8dpi	10	199.98	12.37	775475	160	769669	861699	92.03	32	0.20	10
Grey value KCl tolerance at 8dpi	10	199.98	10.68	775475	281	769669	919978	150.309	56	0.17	10
Melanisation rate in KCl	10	200.35	8.66	792011	299	769669	924425	154.756	59	0.14	10
Colony radius in Control at 8dpi	10	200.72	12.32	792272	299	769669	924425	154.756	59	0.19	10
Colony radius KCl tolerance at 8dpi	10	211.51	10.58	924412	283	792019	924425	132.406	53	0.17	10
Colony radius KCl tolerance at 12dpi	10	211.51	6.68	924414	1081	749521	1302645	553.124	211	0.11	10
Colony radius in Control at 12dpi	10	212.61	4.92	924425	540	749521	1054575	305.054	116	0.08	10
Colony radius in Control at 8dpi	11	183.07	4.20	507361	528	311631	698718	387.087	128	0.07	11
Colony radius KCl tolerance at 12dpi	11	188.28	5.60	581439	1340	74200	890544	816.344	268	0.09	11
Grey value in KCl at 12dpi	11	192.28	6.87	605810	408	463714	776764	313.05	99	0.11	11
Growth rate in Control	11	193.00	19.63	616039	5	616039	658119	42.08	13	0.29	11
Growth rate in KCl	11	193.00	5.47	616039	219	468760	698718	229.958	72	0.09	11
Colony radius in Control at 12dpi	11	193.00	16.79	616039	25	601600	658119	56.519	16	0.26	11
Grey value in Control at 12dpi	11	194.11	6.64	657530	498	514049	870059	356.01	121	0.11	11
Grey value KCl tolerance at 12dpi	11	194.11	5.55	657530	700	367519	870059	502.54	168	0.09	11
Grey value in Control at 8dpi	11	194.11	5.30	658070	244	463714	704331	240.617	77	0.09	11
Grey value in KCl at 8dpi	11	194.11	17.09	658119	5	616039	658119	42.08	13	0.26	11
Growth rate KCl tolerance	11	195.28	7.45	658119	370	500942	802650	301.708	98	0.12	11

Table 3. Description of all QTLs identified from the 1A5x1E4 cross, including map and physical position of the LOD peak marker, the bayes credible interval length and the number of genes within the interval. PVE is the proportion of variance explained by the QTL. QTL number is a number assigned to a particular QTL peak which identifies overlapping QTLs..

Type	Chr	Pos (cM)	LOD	Pos (bp)	No. markers	Interval Start (bp)	Interval end (bp)	Interval size (bp)	No. Genes	PVE	QTL number
Grey value in Control at 8dpi	1	328.91	5.69	572065	402	442697	727696	284.999	92	0.10	1
Grey value KCl tolerance at 8dpi	1	328.91	5.01	572065	405	442697	732023	289.326	93	0.08	1
Melanisation rate KCl tolerance	1	328.91	4.46	572065	463	461577	779936	318.359	102	0.08	1
Colony radius KCl tolerance at 12dpi	1	559.12	4.07	1849200	4654	399579	5725682	5326.103	1964	0.07	2
Growth rate KCl tolerance	1	797.00	4.80	3405315	3068	399544	3488556	3089.012	1151	0.08	3
Colony radius in KCl at 12dpi	1	1104.00	4.03	5792644	4774	372886	5956797	5583.911	2056	0.07	4
Melanisation rate in Control	2	70.00	6.56	574965	1211	405115	1914465	1509.35	509	0.11	5
Grey value in Control at 12dpi	2	204.66	4.55	1438934	618	1284536	2152463	867.927	281	0.08	6
Grey value in Control at 8dpi	2	204.66	18.08	1438934	185	1438863	1748215	309.352	95	0.27	6
Grey value KCl tolerance at 8dpi	2	204.66	15.97	1438934	184	1438934	1748215	309.281	95	0.24	6
Colony radius in Control at 8dpi	2	211.42	8.76	1631840	217	1438934	1776132	337.198	105	0.14	6
Grey value KCl tolerance at 8dpi	3	147.00	6.15	1325440	210	1243932	1684531	440.599	179	0.10	7
Grey value in Control at 8dpi	3	148.00	6.66	1325440	133	1325422	1592801	267.379	115	0.11	7
Colony radius in Control at 8dpi	3	163.74	5.83	1521535	725	355042	1592801	1237.759	374	0.10	8
Melanisation rate in Control	3	193.38	11.99	1711037	290	1489966	2105601	615.635	180	0.19	9
Grey value in Control at 12dpi	3	198.73	4.04	1928783	860	1743111	2927356	1184.245	361	0.07	9
Growth rate KCl tolerance	5	399.00	4.34	2638913	155	2511016	2658480	147.464	48	0.07	10
Colony radius in KCl at 8dpi	6	179.00	4.48	1512508	594	778583	1557528	778.945	256	0.08	11
Growth rate in Control	7	158.82	4.43	539824	255	437699	651318	213.619	80	0.07	12
Colony radius in Control at 12dpi	7	158.82	4.67	539837	254	437718	651318	213.6	80	0.08	12
Growth rate in KCl	8	76.80	8.79	274792	82	256529	346224	89.695	25	0.14	13
Colony radius in KCl at 12dpi	8	76.80	8.94	274792	67	256529	313758	57.229	17	0.15	13
Grey value in Control at 12dpi	8	78.58	4.92	302772	882	89684	921404	831.72	308	0.08	13

Grey value in Control at 8dpi	8	79.00	12.12	302772	66	270834	346224	75.39	22	0.19	13
Colony radius in Control at 12dpi	8	79.00	9.72	302772	80	260355	346224	85.869	24	0.16	13
Colony radius in Control at 8dpi	8	79.00	11.06	302772	415	260101	732444	472.343	163	0.18	13
Grey value KCl tolerance at 8dpi	8	79.00	8.26	302772	339	89573	346224	256.651	90	0.14	13
Colony radius KCl tolerance at 8dpi	8	137.00	5.05	832180	1177	589084	1648406	1059.322	396	0.08	13
Grey value in KCl at 8dpi	8	164.00	4.33	1377926	1406	305075	1623437	1318.362	485	0.07	13
Melanisation rate in Control	12	126.74	4.24	473688	275	441706	821603	379.897	101	0.07	14

Table 4 Genes within the QTL interval for colony radius at 8 dpi in KCl and colony radius KCl tolerance at 8 dpi in the 3D7x3D1 cross. Bold indicates genes that are the most promising candidate genes, as described in the main text. Gene sequence similarity between the two parental strains was obtained using BLAST. DGE: Differential gene expression column indicates if the gene was differentially expressed between parents in any one of the data sets used (in planta or in vitro, see Methods in main text). The annotations were obtained using Interproscan using multiple protein databases, here we show only the PFAM annotation.

Chr	Gene id 3D7	Gene start	Gene end	Gene id 3D1	Sequence similarity (%)	DGE	Annotation
3	**ZT3D7_G3622**	**598091**	**599743**	**ZT3D1_G3567**	**97.68**	**yes**	**Fungal-specific transcription factor domain**
3	ZT3D7_G3623	637954	638542	ZT3D1_G7387	96.23	NA	
3	ZT3D7_G3624	653468	654397	ZT3D1_G3569	90.78	yes	Protein of unknown function (DUF1349)
3	ZT3D7_G3625	655237	656628	ZT3D1_G3570	97.04	yes	
3	ZT3D7_G3626	667030	667567	NA		yes	Pterin 4 alpha carbinolamine dehydratase
3	ZT3D7_G3627	667812	668321	ZT3D1_G3571	98.46	yes	
3	ZT3D7_G3628	670445	671787	ZT3D1_G3572	100.00	yes	Transketolase, pyrimidine binding domain, Transketolase, C-terminal domain
3	ZT3D7_G3629	672336	673658	ZT3D1_G3573	99.83	NA	
3	ZT3D7_G3630	674777	676658	ZT3D1_G3574	99.96	yes	Dehydratase family
3	ZT3D7_G3631	677782	680135	ZT3D1_G3576	97.72	yes	ABC transporter, ABC transporter
3	ZT3D7_G3632	680734	681942	ZT3D1_G3577	97.73	yes	Epoxide hydrolase N terminus
3	ZT3D7_G3633	683647	685131	ZT3D1_G3578	81.12	NA	Phosphotransferase enzyme family
3	ZT3D7_G3634	685575	685937	ZT3D1_G3579	98.96	yes	
3	**ZT3D7_G3635**	**686721**	**692363**	**ZT3D1_G3580**	**99.07**	**yes**	**Protein kinase domain, Protein kinase C terminal domain, N-terminal half of MaoC dehydratase, MaoC like domain, C2 domain**
3	ZT3D7_G3636	693114	696535	ZT3D1_G3582	99.50	yes	Importin-beta N-terminal domain, Cse1
3	**ZT3D7_G3637**	**697014**	**700871**	**ZT3D1_G3583**	**97.44**	**yes**	**DEAD/DEAH box helicase, rRNA-processing arch domain, DSHCT (NUC185) domain, Ski2 N-terminal region, Helicase conserved C-terminal domain**
3	ZT3D7_G3638	701182	702047	ZT3D1_G3584	98.74	yes	short chain dehydrogenase
3	**ZT3D7_G3639**	**702431**	**704143**	**ZT3D1_G3585**	**97.16**	**yes**	**Major Facilitator Superfamily**
3	ZT3D7_G3640	705425	706303	ZT3D1_G3587	99.91	yes	
3	ZT3D7_G3641	706740	707975	ZT3D1_G3588	98.53	yes	
3	ZT3D7_G3642	708779	709336	ZT3D1_G3589	98.33	NS	
3	ZT3D7_G3643	709874	711886	ZT3D1_G3590	94.46	yes	Ubiquitin-conjugating enzyme
3	ZT3D7_G3644	712327	715212	ZT3D1_G3591	98.93	yes	Utp14 protein
3	ZT3D7_G3645	715430	715798	ZT3D1_G3592	99.74	no	Mitochondrial ribosomal protein L27

3	ZT3D7_G3646	716302	720497	ZT3D1_G3593	98.96	NA	Staphylococcal nuclease homologue, short chain dehydrogenase, Enoyl-(Acyl carrier protein) reductase, Tudor domain
3	ZT3D7_G3647	720900	723737	ZT3D1_G3594	99.38	yes	PQQ-like domain, ER membrane protein complex subunit 1, C-terminal
3	ZT3D7_G3648	725571	726437	ZT3D1_G3595	99.02	no	Ras family
3	ZT3D7_G3649	726651	727599	NA		yes	NmrA-like family
3	ZT3D7_G3650	727935	729979	ZT3D1_G3596	98.53	no	Heterokaryon incompatibility protein (HET)
3	ZT3D7_G3651	730698	731819	ZT3D1_G3597	99.93	yes	Transaldolase/Fructose-6-phosphate aldolase
3	ZT3D7_G3652	732329	733083	ZT3D1_G3598	96.53	yes	Copper/zinc superoxide dismutase (SODC)
3	ZT3D7_G3653	734230	736078	ZT3D1_G3599	99.73	yes	FGGY family of carbohydrate kinases, C-terminal domain, FGGY family of carbohydrate kinases, N-terminal domain
3	ZT3D7_G3654	737260	737987	ZT3D1_G3600	99.87	yes	
3	**ZT3D7_G3655**	**739211**	**741757**	**ZT3D1_G3601**	**99.97**	**yes**	**CBS domain, Voltage gated chloride channel**
3	ZT3D7_G3656	743352	744442	ZT3D1_G3602	99.73	yes	Protein of unknown function (DUF1295)
3	ZT3D7_G3657	744744	746896	ZT3D1_G3603	99.92	yes	Pro-kumamolisin, activation domain
3	**ZT3D7_G3658**	**748505**	**750639**	**ZT3D1_G3604**	**100**	**yes**	**PX domain, Vps5 C terminal like**