Parvovirus infection alters the nucleolar structure

Salla Mattola[1*], Simon Leclerc[1*], Satu Hakanen, [1] Vesa Aho[1], Colin R Parrish[2], Vihinen-

Ranta M[1].

[1] Department of Biological and Environmental Science and Nanoscience Center,
 University of Jyvaskyla, Jyvaskyla, Finland
[2] Baker Institute for Animal Health, Department of Microbiology and Immunology, College
of Veterinary Medicine, University of Cornell, Ithaca, USA

(*These authors contributed equally to this work).

#Address correspondence to Maija Vihinen-Ranta, maija.vihinen-ranta@jyu.fi

University of Jyvaskyla, Department of Biological and Environmental Science and

Nanoscience Center, P.O. Box 35, FI-40014 University of Jyvaskyla, Finland.

Running head: Parvovirus infection-induced changes in nucleoli

Keywords: Canine parvovirus, structural changes of nucleoli, viral NS2 protein

ABSTRACT

The nucleolus is a dynamic nuclear structure which plays important roles in ribosome biogenesis and cellular stress response to stressors such as viral infection. The nucleolus and nucleolar proteins are essential for the progression of infection by several viruses. Consequently, viral infection often induces alterations in nucleolar structure and composition. Here, we applied a deep learning algorithm segmentation and nucleolin labeling to analyze the nucleolar changes induced by autonomous parvovirus infection. Our results show that the size of nucleoli decreases and nucleolin is released into the nucleoplasm in late infection. Analyses of ki-67, one of the NS2-associated nucleolar proteins and a key factor in nucleolar organization, showed that the interaction between ki-67 with DNA increases in infection. The infection initiated by a viral clone lacking an intact NS2 failed to decrease the nucleolar size, however, the orientation of the nucleoli was changed. Our results suggest that parvoviruses modify and exploit nucleoli and nucleolar proteins during infection, and NS2 protein might play a role in the regulation of these processes.

(Words 169)

Authors summary

Autonomous parvoviruses with small genomes and only a small number of encoded proteins depend on the cellular nuclear machinery for their efficient replication. The nucleolus is an active nuclear structure involved in the essential functions of cells including synthesis of ribosomal RNA and regulation of cell growth and cellular stress. Many viruses target the nucleoli to manipulate cellular antiviral defense and to facilitate their replication by recruiting nucleolar proteins. Capsids of dependoparvoviruses, such as AAV, are assembled in the nucleolus, however, much less is known about how autonomous parvoviruses interact with the

nucleolus. Our results suggest that the progression of autonomous parvovirus infection results in structural alteration of nucleolar structure and redistribution of nucleolar proteins. Analyses of the nonstructural protein NS2 mutants in comparison to wild-type infection indicated that NS2 might play an important role in viral interactions with the nucleolus.

INTRODUCTION

Parvoviruses are small DNA viruses that undergo replication inside the host cell nucleus. Due to their relatively small genome size parvoviruses depend heavily on cellular factors and pathways for successful replication. Interaction with a multifunctional subnuclear compartment, nucleolus, is crucial for the replication and pathogenesis of several DNA viruses including herpesviruses (Boyne & Whitehouse, 2009; Callé et al., 2008; Pearson et al., 2011) and adenoviruses (Hindley et al., 2007; Lam et al., 2010; Lee et al., 2003; Okuwaki et al., 2001). Moreover, the capsids of adeno-associated viruses (AAV), a helper-dependent parvovirus, are assembled in the nucleolus (Johnson & Samulski, 2009; Wistuba et al., 1997). It is not clear, however, what is the role of nucleoli in autonomous parvovirus life cycle.

The nucleolus is the largest nuclear subcompartment which performs essential cellular functions involved in ribosome biogenesis such as transcription of ribosomal RNA genes, processing of precursor rRNAs, and ribosome subunit assembly (Dubois & Boisvert, 2016; Raška et al., 2006). Nucleoli are formed from specific chromosomal regions of acrocentric chromosomes (HSA13, 14, 15, 21, 22) termed as nucleolar-organizing regions (Henderson et al 1972). The mammalian nucleolus comprises three major morphological components, the fibrillar centre (FC), the dense fibrillar component (DFC) and the granular component (GC) (Hernandez-Verdun, 2011; Pederson, 2011; Raška et al., 2006). The FC contains several hundred copies of rRNA genes, arranged in tandem arrays of ribosomal gene repeats (McStay,

2016; Pederson, 2011). FCs and DFCs are involved in rDNA transcription and production of precursor ribosomal RNA (pre-rRNA), whereas further processing of pre-rRNA and assembly of ribosome subunits occur in GC (Hernandez-Verdun, 2011; Pederson, 2011; Raška et al., 2006). The nucleoli are dynamic structures and assemble through liquid-liquid phase separation when their mobile proteins constantly move between the nucleolus and the nucleoplasm (Falahati & Wieschaus, 2017; Feric et al., 2016; Lafontaine et al., 2021). The proteome of the human nucleoli contains approximately 4500 mobile proteins (Ahmad et al., 2009; F. M. Boisvert et al., 2010). The nucleoli have many additional functions besides ribosomal biogenesis including the regulation of mitosis and the cell-cycle, and acting as sensors for stressors such as oxidative stress, heat shock, nutrient deprivation (F.-M. Boisvert et al., 2007; Boulon et al., 2010; Iarovaia et al., 2019; Scherl et al., 2002) and viral infection (Salvetti & Greco, 2014).

Autonomous parvoviruses have non-enveloped icosahedral particles of ~25 nm in diameter containing a linear single-stranded DNA genome of ~5.0 kb. The viral DNA encodes two open reading frames for structural and nonstructural proteins (Cotmore et al., 1983). The canine parvovirus (CPV) capsid is composed of viral proteins VP1 and VP2, and capsids with DNA contain VP3, which is formed through the proteolytic cleavage of the VP2 N-terminus (Weichert et al., 1998). Autonomous parvoviruses encode two nonstructural proteins, NS1 and NS2 (Cotmore et al 1983). The CPV NS1 protein is a multifunctional protein that exhibits site-specific DNA binding, ATPase, helicase, and nickase activities (Niskanen et al., 2010, 2013) and is involved in the induction of apoptosis in host cells (Gupta et al., 2016; Saxena et al., 2013). The NS2 protein of an autonomously replicating parvovirus, minute virus of mice (MVM), is involved in capsid assembly, viral DNA replication and translation of viral mRNA (Cotmore et al., 1997; Naeger et al., 1990, 1993). Although the function of CPV NS2 is

currently mostly undefined our recent studies showed that it plays a role in chromatin remodeling and DNA damage response during replication (Mattola et al., 2022).

In this study the structural changes of nucleoli were monitored as CPV infection proceeded, particularly, the role of the NS2 protein in the process was clarified. We discovered that upon infection the nucleolar size was significantly altered and cellular DNA and nucleolar proteins were redistributed, and many of these changes were not detected if the virus lacked an intact NS2. The potential role of NS2 in the interaction between virus and nucleolus is also supported by the recent proximity-dependent biotin identification (BioID) findings showing that NS2 is associated with several nucleolar proteins.

RESULTS

Nucleolar size decreases in late infection

The nucleolus is a highly dynamic structure that responds by profound morphological and compositional alterations to cellular stresses such as viral infections (Lam et al., 2010; D. A. Matthews, 2001; Olausson et al., 2012; Salvetti & Greco, 2014). To characterize the nucleolar response to CPV infection-induced cellular stress, we examined the morphological alterations of nucleoli as CPV infection proceeded.

We studied the size and number of nucleoli in NLFK cells at 8, 12 and 24 hours post infection (hpi) by segmenting the labeled nucleolin signal using a deep learning algorithm (Fig. S1). The analyses showed that at 8 and 12 hpi the nucleolar volume was not significantly different from that in noninfected cells. However, a significant decrease was detected at 24 hpi (Fig. 1A and 1B). Simultaneously, the average number of nucleoli per nucleus remained similar to

noninfected cells at 8 and 12 hpi, however, a slight but statistically insignificant increase was detected at 24 hpi (Fig. 1C).

Figure 1. Progression of infection leads to changes in size and structure of nucleoli.
(A) Representative confocal images show the localization of the nucleoli (magenta) in noninfected and infected NLFK cell nuclei at 8, 12, and 24 hpi. Only nuclei are shown. The cells were immunolabeled with antibodies against nucleolin (magenta) and intact capsids (green). Blue corresponds to DAPI staining. Scale bars, 5 µm. The box-swarm plots represent (B) size defined by nucleoli volume and (C) number of nucleoli in noninfected and infected cells (n = 34, 25, 27, and 75 for noninfected, 8, 12, and 24 hpi, respectively). The 3D nucleoli were segmented using a deep learning algorithm. The error bars show the standard error of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as *** (p<0.001), * (p<0.05), or ns (not significant).

Altogether, our findings demonstrated a reduced nucleolar size and a slightly increased number of nucleoli in response to progression of infection, suggesting that infection induces important nucleolar alterations typical to nucleolar stress (Yang et al., 2018).

Nucleolin is released into the nucleoplasm from the nucleolus in late infection

The multifunctional phosphoprotein nucleolin is one of the most abundant nucleolar proteins (Biggiogera et al., 1990). Nucleolin has important roles in rDNA transcription, rRNA maturation, in the assembly and transport of ribosomes (Ginisty et al., 1999; Srivastava & Pollard, 1999). To participate in these processes nucleolin shuttles between the nucleolus and nucleoplasm (Chen & Huang, 2001). Relocalization of nucleolin out of the nucleoli is also triggered by cellular stress (Daniely & Borowiec, 2000; Jia et al., 2017) and by viral infection,

e.g. with herpes simplex virus type 1 (HSV-1) (Callé et al., 2008; Greco et al., 2012; Lymberopoulos & Pearson, 2007).

To gain insight into CPV infection-induced changes in the distribution of nucleolin, we analyzed noninfected and infected NLFK cells at 8, 12, and 24 hpi. Compared to noninfected cells, the total amount of intranuclear nucleolin was higher at 8and 12 hpi, but not at 24 hpi (Fig 2A). The nuclear distribution of nucleolin in infected cells was relatively similar to noninfected cells, however, nucleolin was slightly more concentrated close to the nuclear envelope (NE) at 12 hpi (Fig 2B). The amount of nucleolin in the segmented nucleoli was similarly significantly higher at 8 hpi and 12 hpi, but not at 24 hpi, in comparison to noninfected cells (Fig. 2C). Notably, the comparison of the ratios of nucleolar nucleolin to nuclear nucleolin in revealed a significant decrease the relative amount of nucleolar nucleolin at 24 hpi. This indicated a shift in the nucleolin distribution from the nucleolus to the nucleoplasm (Fig 2D).

Figure 2. Distribution of nucleolin changes as infection proceeds.
Analyses of temporal changes in infected cells at 8, 12 and 24 hpi (see Fig. 1A). (A) Total nuclear intensities of nucleolin, (B) nuclear distributions of nucleolin as a function of distance from the NE, (C) total nucleolar intensities of nucleolin in the segmented nucleoli, and (D) the nucleolar to the nuclear ratios of nucleolin are shown (n=19, 15, 17, and 28 for noninfected, 8, 12, and 24 hpi, respectively). The shaded areas in distribution graphs represent the standard error of the mean. The error bars show the standard error of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as **** (p<0.0001), ** (p<0.01), * (p<0.05) or ns (not significant).

To characterize potential structural alterations in nucleolar structure during CPV infection, we examined the changes in the chromatin organization inside the nucleolus. As expected, DNA was accumulated toward the NE at late stages of infection (Fig. 3A). Next, the analysis of segmented nucleolar region revealed that the amount of nucleolar DNA was significantly increased in comparison to noninfected cells at 8 and 12 hpi and decreased from 12 hpi to 24 hpi (Fig. 3B). To examine the organization of chromatin around and inside the nucleoli, we analyzed the distribution of DNA on both sides of the nucleolar border. In noninfected cells the DNA was distributed both outside and inside the nucleolus. In infected cells, nucleolin was distributed more to the nucleolar interior at 8 and 12 hpi, and to the exterior at 24 hpi (Fig. 3C).

Figure 3. Nucleolar chromatin organization is changed in infected cells.
Analyses of time-dependent changes in DNA intensity and distribution of infected cells (see Fig. 1A). A) Nuclear distribution of DNA in noninfected and infected NLFK cell nuclei at 8, 12, and 24 hpi (n =19, 15, 17, and 28, respectively). (B) The total nucleolar intensities of DNA in segmented nucleoli and (C) distribution of DNA close to the segmented nucleolar edge as a function of increasing distance from the nucleolar border are shown. The error bars and the shaded areas in the distribution graphs represent the standard error of the mean. The error bars show the standard error of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as *** (p<0.001), ** (p<0.01) or ns (not significant).

Taken together, these findings suggest that the shuttling of nucleolin from nucleoli to nucleoplasm is increased upon progression of infection, possibly due to the infection-induced cellular stress. Moreover, the infection-induced redistribution of DNA in close proximity and inside the nucleolus suggest that nucleolar structure changes during infection.

NS2 interactomes are localized to multiple nucleolar regions

Our previous BioID analyses revealed 122 high-confidence interactor proteins of NS2 (Bayesian false discovery rate, FDR ≤0.05) in infected and noninfected cells (Mattola et al., 2022). Comparison of the NS2 interacting proteins with the identified nucleolar proteins from the Human Protein Atlas (HPA) database (www.proteinatlas.org; Uhlen et al. 2010; Thul et al. 2017) revealed 26 nucleolar NS2 interactors in infected cells. The database is based on a study which identified 1,318 nucleolar proteins mostly located in FC, DFC or nucleolar rim in the interphase cells (Fig. 4A) (Stenström et al., 2020). Our analysis of the distribution of NS2-associated proteins demonstrated that 13 proteins were localized to the whole nucleolus, five proteins to the fibrillar center, and two proteins to the nucleolar rim. One protein showed dual localization to both the fibrillar center and nucleoplasm, and five proteins to the nucleolar rim and nucleolus (Fig. 4B). Further identification of NS2-associated proteins indicated that they have a role in multiple biological processes including transcription, RNA processing, DNA damage response and cell cycle (Fig. 4C, Table S1). The proteins involved in RNA processing were localized to the whole nucleolus, whereas proteins of DNA damage response were found from the fibrillar centers. Many of the proteins had multiple functions within several nucleolar regions. The number of nucleolar NS2-associated proteins was smaller in infected cells compared to that in NS2 transfected noninfected cells which suggest that actual number of interactors during infection is smaller than the possible number of interactors. Consequently, it is likely that some of these associated proteins are temporally linked to NS2 at a specific time

of the infection (Fig. S2A). In noninfected cells, NS2-associated proteins were localized to the whole nucleolus (21), to nucleolar rim (3), and five proteins had dual localization to both the fibrillar center and nucleoplasm, and five proteins to the nucleoli rim and nucleolus. The interactors associated to fibrillar centers were specific to that localization only (Fig. S2B). The top putative interactor of NS2 identified by BioID in infected cells was the proliferation marker protein Ki-67 (MKI67) (Mattola et al., 2022).

Figure 4. NS2-associated proteins have multiple localization sites in the nucleoli.
(A) A schematic image of the organization of the nucleolus and its subcompartments: the fibrillar center (FC), the dense fibrillar component (DFC), and the granular component (GC). (B) A Venn diagram showing the nucleolar localization of NS2-associated proteins identified by BioID in infected cells. The numbers of proteins localized into the whole nucleolus (blue), fibrillar center including FC and DFC (green), nucleolar rim (orange), or dual localization (light brown, light green) are shown. (C) The cumulative plot shows the enriched Gene Ontology (GO) terms for biological processes and nucleolar localization of NS2-associated proteins.

Altogether, proteins associated with NS2 are localized to multiple regions of the nucleolus most likely reflecting their multiple structural and functional features during infection.

Interaction between chromatin and ki-67 increases in late infection.

The ki-67 protein is a nucleolar heterochromatin anchoring protein which is specifically localized to the nucleolar rim of interphase cells (Stenström et al., 2020). Ki-67 is involved in the structural organization of chromatin and the nucleolus (Booth et al., 2014; Cuylen et al., 2016; MacCallum & Hall, 2000; Sobecki et al., 2016). As described previously, ki-67 is a prominent interactor for CPV NS2 (Mattola et al., 2022).

To examine virus-induced changes in nucleolar organization, we analyzed the distribution of ki-67 in noninfected and infected NLFK cells at 24 hpi. In noninfected cells ki-67 was accumulated in nucleoli, whereas in infected cells ki-67 was located at the perinuclear region, around the nucleoli and in the nucleoplasm (Fig. 5A). Both DNA and ki-67 showed similar nuclear distribution in infected cells including accumulation close to the NE (Figs 5B and 5C). Total intensities of DNA and ki-67 were significantly increased in comparison to noninfected cells (Fig 5D and 5E). Notably, the correlation between DNA and ki-67 increased significantly suggesting that infection facilitated tethering of ki-67 to the DNA (Fig 5F).

Taken together, we demonstrated that infection leads to an increase in the amount of ki-67, enhancement of ki-67 association with cellular DNA, and redistribution of ki-67 to the perichromosomal layer around the nucleoli and close to the NE. This supports a model where ki-67 is involved in the virus-induced change of DNA organization in nuclear and nucleosomal periphery.

Figure 5. Association of ki-67 with DNA increases in infection
(A) Representative confocal images showing the localization of DNA (blue), NS1 (green), and ki-67 (magenta) in noninfected and infected NLFK cells at 24 hpi. The cells were immunolabeled with antibodies against NS1 and ki-67. DNA was labeled with DAPI. Scale bars, 5 µm. The nuclear distributions of DNA (B) and ki-67 (C) as a function of increasing distance from the NE in noninfected and infected cells (n = 30). The shaded areas represent the standard error of the mean. Total intensities of DNA (D) and ki-67 (E) are shown together with the Pearson correlation coefficients between ki-67 and DNA (F). The error bars show the standard error of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as *** ($p < 0.001$) and * ($p < 0.05$).

NS2 plays a role in the infection-induced changes of nucleoli

Our previous BioID analysis suggested an association between CPV NS2 and several nucleolar proteins (Mattola et al., 2022). To further assess the role of NS2 in the nucleolar changes, we compared the nucleolar size and number in cells transfected with wild type (wt) infectious clone or with NS2 mutant clones. These mutants were designed to disrupt the expression of NS2 but not to affect sequence of NS1 (Wang et al., 1998). The NS2 mutants contained termination codons in the NS2 coding sequence of the infectious clone either upstream of the common NS1/2 splice donor or downstream of the NS2-specific splice acceptor. The original publication indicated that the NS2 splice donor mutant led to significantly deficient production

of viral capsids, capsid proteins and viral DNA. However, analyses of the splice acceptor indicated that the C-terminal domain of NS2 could be largely deleted with only modest effects on the viral replication in tissue culture cells and animals (Wang et al., 1998).

The previously shown nucleolar localization of NS2 in infected cells and NS2 association with nucleolar proteins (Mattola et al., 2022) suggest that NS2 might have a potential role in the infection-induced modification of the nucleoli described above. Here, we assessed how NS2 mutations affected the virus-induced changes of nucleoli in NLFK cells. Similarly, to infected cells at 24 hpi, the average size of segmented nucleoli was decreased and the number of nucleoli was increased in wt infectious clone transfected cells in comparison to nontransfected cells at 24 hours post transfection (hpt) (Fig. 6A). The quantitative analysis confirmed these changes in wt transfected cells and showed a similar decrease in nucleolar size in the NS2 splice acceptor mutant transfected cells. Notably, the size of nucleoli and their number did not change in the NS2 splice donor mutant transfected cells in comparison to nontransfected cells. (Figs 6B and C).

Figure 6. Intact NS2 is required for nucleolar changes
(A) Representative confocal images show nucleoli in nontransfected NLFK cells, and cells transfected with wt, NS2 splice donor and acceptor mutants at 24 hpt. The cells were immunolabeled with antibody against nucleolin (magenta) and intact viral capsids (green), and stained with DAPI (blue). Scale bars, 5 µm. The box-swarm plots represents (B) size and (C) number of nucleoli in nontransfected and transfected cells (n = 26, 47, 31 and 26 nucleoli for nontrasnfected, wt, and splice donor mutant and splice acceptor mutant, respectively). The 3D nucleoli were segmented using deep learning algorithm. The error bars show the standard error

of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as **** (p<0.0001), * (p<0.05) or ns (not significant).

Taken together, our results suggest that NS2 may play a role in the infection-induced change of nucleoli. However, the mechanisms by which NS2 is associated with the nucleolar alterations remains to be determined.

Nucleolar distribution of chromatin is affected by NS2 mutation

As shown above, infection induces changes in chromatin distribution in close proximity and within the nucleolus (Fig. 3). To further assess the possible role of NS2 in nucleolar organization of chromatin, we analyzed DNA distribution in cells transfected with wt or NS2 mutants.

Consistent with our infection data, the total nuclear intensity of DNA was increased in wt infectious clone transfected cells and also in wild-type-like splice acceptor mutant transfected NLFK cells at 24 hpt in comparison to nontransfected cells. The most significant increase in nuclear DNA was detected in the splice donor mutant transfected cells (Figs 7A). DNA showed a similar nuclear distribution in all transfected cells including accumulation close to the NE (Fig. 7B). The total amount of nucleolar DNA was most significantly increased in the splice donor transfected cells, and more moderately in wt and the splice acceptor transfected cells compared to nontransfected cells (Fig. 7C). Distribution analyses showed that consistently with 24 hpi (Fig. 3D), DNA was localized both to the inner and outer side of the nucleolar border in wt and the splice acceptor mutant transfected cells. In the splice donor mutant transfected cells, the DNA distribution was slightly shifted towards the nucleolar interior (Fig. 7D).

Figure 7. Chromatin organization close to and within the nucleolus is affected by NS2 mutation

(A) Total nuclear intensities of DNA and (B) nuclear distribution of DNA as a function of distance from the NE in nontransfected, wt, and NS2 splice donor and acceptor mutant transfected NLFK cells at 24 hpt (n =20, 19, 18, and 16, respectively). (C) Total nucleolar DNA intensities in segmented nucleoli and (D) distribution of DNA close to the segmented nucleolar edge are shown. The error bars and the shaded areas in the distribution graphs represent the standard error of the mean. The error bars show the standard error of the mean. Statistical significances were determined using Student´s t-test. The significance values shown are denoted as **** (p<0.0001), ** (p<0.01), * (p<0.05) or ns (not significant).

We showed previously that CPV NS2 has a potential role in controlling chromatin remodeling and DDR response (Mattola et al., 2022). Our current observations suggest that NS2 might be needed to organize chromatin packaging on the nucleolar interior and exterior.

Orientation of the nucleoli is altered in NS2 mutant infection

The nucleolus-nucleoplasm interface is known to be maintained by processes dependent on ATP consumption, and impacted by changes in the chromatin packaging and active transcription. In ATP depletion, the curvature and shape of the nucleoli are affected but not the average nucleolar area, eccentricity, or orientation. Orientation is unaffected by a wide range of biochemical perturbations (Caragine et al., 2019). Viral replication of autonomous parvoviruses is a highly ATP-dependent process, overseen by the NS1 protein. The ATP binding pocket of NS1 residing in the middle of the protein, a region not shared with the NS2 (Niskanen et al., 2010).

To further examine the participation of NS2 in the nucleolar organization we determined the orientation of the nucleoli in NS2 mutants transfected cells compared to wt transfected and nontransfected cells. Orientations of the nucleoli were calculated between the major nucleolar and nuclear axes and were given the values from 0 to 90 degrees (Fig 8A). Interestingly, we noticed that the orientation of nucleoli was often perpendicular to the orientation of the nucleus in the replication deficient splice donor mutant transfected NLFK cells (Fig. 8B). This observation was supported by our findings of the orientation of segmented nucleoli that the average nucleolar angle in the splice donor mutant transfected cells was significantly changed in comparison to nontransfected and wt transfected cells (Fig. 8C).

Figure 8. The orientation of the nucleoli is affected in NS2 mutant transfected cells.
(A) Schematic image describing the angle measurement between the nucleus and the nucleolus. (B) Representative segmented images with the represented nucleus direction (blue), nucleolus direction (green) and the angle (white). (C) The angle between the major nuclear and nucleolar axes in nontransfected, wt and NS2 splice donor and acceptor mutants transfected cells (n = 26, 47, 31 and 26 nucleoli respectively). Statistical significances were determined using Student´s t-test. The significance values shown are denoted as * (p<0.05) or ns (not significant).

The change in the orientation of nucleoli in the NS2 splice donor transfected cells is an unexpected finding. However, as shown previously the nucleoli are affected by the state of the surrounding nuclear DNA (Caragine et al., 2019). Altogether, consistent with our previous notion that NS2 affects the nuclear chromatin organization (Mattola et al., 2022), our observations support a model where the lack of intact NS2 leads to previously unreported changes in the nucleolar orientation upon CPV.

DISCUSSION

Many DNA viruses, such as HSV-1, adenovirus and AAV, target the nucleolus to interfere with host cell function and to recruit its proteins to facilitate viral infection (Callé et al., 2008; Lee et al., 2003; D. Matthews et al., 2011; Salvetti & Greco, 2014; Wistuba et al., 1997). The nucleolus plays currently mostly undefined roles during the autonomous parvovirus infection. Our findings demonstrated that CPV infection is able to induce temporal changes in the structure and protein content of the nucleolus by a process which is likely mediated by the viral NS2 protein.

The nucleolus is a dynamic structure, and the number and size of nucleoli change during the cell cycle (Caragine et al., 2019; Hernandez-Verdun, 2011). The size of nucleoli is also linked to nucleolar activity under stress (Boulon et al., 2010). Viral infection may also lead to reorganization of nucleolar architecture and change in its size (Greco, 2009; Hiscox, 2007).

Our findings demonstrated that, in addition to changing nucleolar size, CPV infection induced alterations in nucleolar chromatin organization and resulted in nucleolin dispersal from the nucleoli. The nucleolus is widely affected by viral infection and many viral proteins, including CPV NS2, localize into the nucleolus (Mattola et al., 2022), while various nucleolar proteins of infected host cell undergo redistribution (Salvetti & Greco, 2014). Nucleolin is a prominent marker for the nucleoli with as much as 90% of the protein localizing to the nucleoli and 5% to the nucleoplasm (Scherl et al., 2002). The nucleolar-nucleoplasmic shuttling of nucleolin is induced in cellular functions related to stress (Daniely et al., 2002; Daniely & Borowiec, 2000; Jia et al., 2017; Kim et al., 2005). The nuclear distribution of nucleolin is affected by virus infection. For example, nucleolin relocates to the cytoplasm in poliovirus infection (Waggoner & Sarnow, 1998) and adenovirus infection leads to viral protein V-mediated transport of nucleolin to the cytoplasm (Matthews, 2001). In HSV-1 infection, the viral protein UL24 is suggested to be involved in nucleolin dispersal (Lymberopoulos & Pearson, 2007). Nucleolin is also required for proper localization of human cytomegalovirus replication protein UL84 (Bender et al., 2014). Nucleolin promotes binding of HSV-1 and influenza A virus to the cell membrane (Chan et al., 2016; Greco et al., 2012; Kumar et al., 2016). The release of nucleolin from the nucleolus might be a result of a virus-induced heat shock response. This is supported by findings that oncolytic parvovirus H1 infection causes a release of inducible heat-shock protein HSP72 in susceptible human tumor cells (Moehler et al., 2003). The nucleolus is embedded in the chromatin network, which is suggested to support the formation and stabilization of its structure by maintaining the nucleolus-nucleoplasm interface (Caragine et al., 2019; Padeken & Heun, 2014; Qi & Zhang, 2021). It is highly likely that chromosomal organization in the vicinity of nucleolus is regulated by nucleolar DNA sequences of nucleolar-organizing regions (Németh & Längst, 2011). In HSV-1- and CPV -infected cells the formation of viral replication compartments is accompanied by extensive spatial reorganization of

chromatin (Ihalainen et al., 2009; Myllys et al., 2016) which might also affect the structure of nucleoli. Altogether, our data demonstrate that autonomous parvoviruses induce changes in the structure and protein content of nucleolus. We also speculate that the infection-induced changes of chromatin distribution inside and outside of the nucleoli might be due to infection-induced altering of the state of DNA in the surroundings of nucleolus or due to changes in nucleolar interactions with the chromatin. However, further experiments are required to explore these processes.

Our earlier BioID data linked CPV NS2 to several nucleolar proteins with multiple biological functions (Mattola et al., 2022). Spatial mapping of associated proteins has revealed their localization to nucleolar subcompartments such as the whole nucleolus, fibrillar center, and nucleoli rim. One of the most prominent interaction partners of NS2, ki-67 (Mattola et al., 2022), is located in the nuclear rim (Stenström et al., 2020). ki-67 is involved in the structural organization of chromatin and the nucleolus (Booth et al., 2014; Cuylen et al., 2016; MacCallum & Hall, 2000; Sobecki et al., 2016). The rim localization of ki-67 allows its interaction with the heterochromosomal layer around the nucleoli in interphase cells (Stenström et al., 2020). The expression and the functions of ki-67 are cell cycle dependent, with the protein level peaking in mitosis as ki-67 relocates to condensed chromosomes to facilitate the formation of the perichromosomal layer (Booth et al., 2014; Cuylen et al., 2016; Gerdes et al., 1984; Sobecki et al., 2017; Stamatiou & Vagnarelli, 2021). Our studies showed that interaction between ki-67 and chromatin is increased, and ki-67 was relocated to the heterochromosomal layer around the nucleoli and close to the NE at late stages of CPV infection. We speculate that the redistribution and increased association of ki-67 with chromatin might reflect the virus-induced regulation of cell cycle progression towards mitosis. This model is supported by our recent findings that infected cells escape from the virus-induced G2/M arrest and move to the

early stages of mitosis (Mattola et al., unpublished). Our studies of NS2-deficient mutant demonstrated that intact NS2 is required for the regulation of nucleolar size, organization of nucleolar chromatin and orientation of the nucleoli. Altogether these findings suggest that CPV NS2 could have a role in viral interactions affecting nucleolar structure and association with cellular chromatin.

Given that nucleolar proteins have many functions in RNA metabolism and chromatin organization, these proteins are usable for facilitation of the viral replication pathway. Our results suggest that autonomous parvoviruses such as CPV recruit nucleolar proteins in a process which is mediated by the NS2 protein and leads to alteration of the nucleolar structure. Our studies provide novel insights into interactions between autonomous parvoviruses and nucleoli which are potentially essential for the progression of infection.

MATERIALS AND METHODS

Cell lines, plasmids and viruses

Norden laboratory feline kidney (NLFK) cells were propagated as previously described (Mattola et al., 2022). CPV type 2 was generated from an infectious plasmid clone p265 (Parrish, 1991) and virus was produced as described earlier (Suikkanen et al., 2002). NS2 splice donor and acceptor mutants have been previously described (Wang et al., 1998). Transfection of NLFK cells was performed using JetOptimus reagent (Polyplus transfection, Illkirch, France).

Antibodies

The following primary antibodies against viral antigens and cellular proteins were used for confocal microscopy analyses: MAb anti-NS1 was a generous gift from Dr. Caroline Astell (University of British Columbia, Vancouver, Canada) (Yeung et al., 1991). MAb Anti-A3B10 antibody was used to detect intact capsids (Weichert et al., 1998). Anti-nucleolin rAb was used to label the nucleoli (Abcam, ab22858) (Cambrigde, UK). Ki-67 was detected with rAb anti-MKI-67 antibody (Abcam, ab15580). The primary antibodies were followed by goat anti-mouse or anti-rabbit Alexa448, Alexa546, and Alexa647 conjugated secondary antibodies (Thermo Fisher Scientific, Waltham, MA).

Immunolabelling and confocal microscopy

For microscopy sample preparation NLFK cells were cultured on coverslips at 37°C with 5 % $CO2$. Cells were either infected with CPV or transfected with CPV infectious clone or NS2 splice mutants and fixed at 8, 12 or 24 hpi and 24 hpt in 4 % paraformaldehyde for 12 minutes. Following fixation, cells were permeabilized with 0.1 % Triton X-100 in phosphate buffered saline supplemented with 0.1 % bovine serum albumin. After primary antibody immunolabeling cells were labeled with mouse or anti-rabbit secondary fluorescent antibodies. DNA was stained with Pro-Long Diamond anti-fade media with DAPI (Thermo Fisher Scientific).

Microscopy of immunolabeled samples was performed with Leica TCS SP8 FALCON (Leica microsystems, Mannheim, Germany) and Nikon A1R (Nikon, Tokyo, Japan) laser scanning confocal microscopes. The parameters for images acquired with Leica TCS SP8 FALCON were as follows: DAPI was excited with a 405-nm diode laser and the fluorescence was collected between 415-495 nm. Alexa488 and Alexa546 were excited with 499 nm and 557 nm wavelengths, respectively, of pulsed white light laser (80 MHz). The emission detection range

was adjusted to 505-561 nm for Alexa488 and 568-648 nm for Alexa546. Microscopy images with Nikon were acquired as follows: DAPI was excited with a 405 nm diode laser and collected with a 450/50 nm band-pass filter. Alexa488 was excited with a 488 nm argon laser, and fluorescence was collected with a 515/30 nm band-pass filter. Alexa 647 was excited with a 642 nm diode laser and detected with a 660 long-pass filter. Resolution for all images was 512 x 512 nm, with a pixel size between 60 nm in the x- and y-directions and 200 nm in the z-direction. For image analysis purposes a fixed optical configuration was used for all images of the same sample set.

Machine learning based segmentation

Human in the loop (HITL) based deep learning method was utilized for the nucleoli segmentations based on the nucleolin labeling (Bobes-Bascarán et al., 2021; Patel et al., 2019). Here we used a U-shape algorithm for the deep neural network (DNN). First, few cell nuclei were manually segmented to train the DNN. The DNN was used to predict the segmentations, based on images from the same dataset but including all planes of the microscopy stack into the segmented area. The mistakes of the algorithm were manually corrected, and the DNN was retrained. By each iteration, more images were added to the analysis, until the output was satisfying (DNN loss function). At this stage, the DNN was used to segment the nucleoli.

Image analyses

The nuclei were segmented from DAPI stain using Otsu's method (Otsu, 1979). Total fluorescence intensities in the nucleus and nucleoli were calculated by summing the pixel intensity values within the segmented regions. Label intensity distributions with respect to the nuclear or nucleolar border were determined by calculating the shortest distance of each pixel to the border and calculating mean intensity as a function of the distance using 2-pixel wide

distance bins for averaging. Due to the flat shape of fixed cells, the confocal section with the largest nuclear area was selected for the analyses on intensity with respect to the nuclear border and 2D Euclidean distance was used as a metric. Since there can be many nucleoli in one cell and the nucleoli can lie in various confocal planes, the whole 3D stack was used in the analyses with respect to the nucleolar border and 3D Euclidean distance was used as a metric. Correlations between DNA staining and antibody labels were determined by using Pearson correlation coefficients. For the nucleolar orientation analyses first the diameters of the nuclei in all the directions were determined, and the longest direction of the nuclei was designated as the zero degrees angle. The angles of the nucleoli were determined by determining the longest sections of the nucleoli and the orientations were compared to the orientation of the zero degrees diameter of the nuclei. The difference between the nucleoli and the zero degrees angle calibration point were represented as the orientation angles of the nucleoli. The number and size analyses of the nucleoli were based on machine learning based segmentations of the nucleoli. For all the analyses, the given statistics were calculated over cells.

BioID and analyzes

BioID of CPV NS2 protein was performed as previously descripted (Mattola et al., 2022).

FUNDING INFORMATION

REFERENCES

Ahmad, Y., Boisvert, F. M., Gregor, P., Cobley, A., & Lamond, A. I. (2009). NOPdb: Nucleolar Proteome Database--2008 update. *Nucleic Acids Research*, *37*(Database issue). https://doi.org/10.1093/NAR/GKN804

Bender, B. J., Coen, D. M., & Strang, B. L. (2014). Dynamic and nucleolin-dependent localization of human cytomegalovirus UL84 to the periphery of viral replication compartments and nucleoli. *Journal of Virology*, *88*(20), 11738–11747. https://doi.org/10.1128/JVI.01889-14

Biggiogera, M., Burki, K., Kaufmann, S. H., Shaper, J. H., Gas, N., Amalric, F., & Fakan, S. (1990). Nucleolar distribution of proteins B23 and nucolin in mouse preimplantation embryos as visualized by immunoelectron microscopy. *Development*, *110*(4), 1263–1270. https://doi.org/10.1242/dev.110.4.1263

Bobes-Bascarán, J., Mosqueira-Rey, E., & Alonso-Ríos, D. (2021). *Improving Medical Data Annotation Including Humans in the Machine Learning Loop*. 39. https://doi.org/10.3390/engproc2021007039

Boisvert, F. M., Lam, Y. W., Lamont, D., & Lamond, A. I. (2010). A quantitative proteomics analysis of subcellular proteome localization and changes induced by DNA damage. *Molecular & Cellular Proteomics : MCP*, *9*(3), 457–470. https://doi.org/10.1074/MCP.M900429-MCP200

Boisvert, F.-M., van Koningsbruggen, S., Navascués, J., & Lamond, A. I. (2007). *Acrocentric chromosome The multifunctional nucleolus*. https://doi.org/10.1038/nrm2184

Booth, D. G., Takagi, M., Sanchez-Pulido, L., Petfalski, E., Vargiu, G., Samejima, K., Imamoto, N., Ponting, C. P., Tollervey, D., Earnshaw, W. C., & Vagnarelli, P. (2014). Ki-67 is a PP1-interacting protein that organises the mitotic chromosome periphery. *ELife*, *2014*(3). https://doi.org/10.7554/ELIFE.01641.001

Boulon, S., Westman, B. J., Hutten, S., Boisvert, F. M., & Lamond, A. I. (2010). The nucleolus under stress. *Molecular Cell*, *40*(2), 216–227. https://doi.org/10.1016/J.MOLCEL.2010.09.024

Boyne, J. R., & Whitehouse, A. (2009). Nucleolar disruption impairs Kaposi's sarcoma-associated herpesvirus ORF57-mediated nuclear export of intronless viral mRNAs. *FEBS Letters*, *583*(22), 3549–3556. https://doi.org/10.1016/J.FEBSLET.2009.10.040

Callé, A., Ugrinova, I., Epstein, A. L., Bouvet, P., Diaz, J.-J., & Greco, A. (2008). Nucleolin is required for an efficient herpes simplex virus type 1 infection. *Journal of Virology*, *82*(10), 4762–4773. https://doi.org/10.1128/JVI.00077-08

Caragine, C. M., Haley, S. C., & Zidovska, A. (2019). Nucleolar dynamics and interactions with nucleoplasm in living cells. *ELife*, *8*. https://doi.org/10.7554/ELIFE.47533

Chan, C. M., Chu, H., Zhang, A. J., Leung, L. H., Sze, K. H., Kao, R. Y. T., Chik, K. K. H., To, K. K. W., Chan, J. F. W., Chen, H., Jin, D. Y., Liu, L., & Yuen, K. Y. (2016). Hemagglutinin of influenza A virus binds specifically to cell surface nucleolin and plays a role in virus internalization. *Virology*, *494*, 78–88. https://doi.org/10.1016/J.VIROL.2016.04.008

Chen, D., & Huang, S. (2001). Nucleolar components involved in ribosome biogenesis cycle between the nucleolus and nucleoplasm in interphase cells. *The Journal of Cell Biology*, *153*(1), 169–176. https://doi.org/10.1083/JCB.153.1.169

Cotmore, S. F., D'Abramo, A. M., Carbonell, L. F., Bratton, J., & Tattersall, P. (1997). The NS2 polypeptide of parvovirus MVM is required for capsid assembly in murine cells. *Virology*, *231*(2), 267–280. https://doi.org/10.1006/VIRO.1997.8545

Cotmore, S. F., Sturzenbecker, L. J., & Tattersall, P. (1983). The autonomous parvovirus MVM encodes two nonstructural proteins in addition to its capsid polypeptides. *Virology*, *129*(2), 333–343. https://doi.org/10.1016/0042-6822(83)90172-1

Cuylen, S., Blaukopf, C., Politi, A. Z., Müller-Reichert, T., Neumann, B., Poser, I., Ellenberg, J., Hyman, A. A., & Gerlich, D. W. (2016). Ki-67 acts as a biological surfactant to disperse mitotic chromosomes. *Nature*. https://doi.org/10.1038/nature18610

Daniely, Y., & Borowiec, J. A. (2000). Formation of a complex between nucleolin and replication protein a after cell stress prevents initiation of DNA replication. *Journal of Cell Biology*, *149*(4), 799–809. https://doi.org/10.1083/jcb.149.4.799

Daniely, Y., Dimitrova, D. D., & Borowiec, J. A. (2002). Stress-Dependent Nucleolin Mobilization Mediated by p53-Nucleolin Complex Formation. *Molecular and Cellular Biology*, *22*(16), 6014. https://doi.org/10.1128/MCB.22.16.6014-6022.2002

Dubois, M. L., & Boisvert, F. M. (2016). The nucleolus: Structure and function. *The Functional Nucleus*, 29–49. https://doi.org/10.1007/978-3-319-38882-3_2/FIGURES/2

Falahati, H., & Wieschaus, E. (2017). Independent active and thermodynamic processes govern the nucleolus assembly in vivo. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(6), 1335–1340. https://doi.org/10.1073/PNAS.1615395114

Feric, M., Vaidya, N., Harmon, T. S., Mitrea, D. M., Zhu, L., Richardson, T. M., Kriwacki, R. W., Pappu, R. v., & Brangwynne, C. P. (2016). Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell*, *165*(7), 1686–1697. https://doi.org/10.1016/J.CELL.2016.04.047

Gerdes, J., Lemke, H., Baisch, H., Wacker, H. H., Schwab, U., & Stein, H. (1984). Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. *Journal of Immunology (Baltimore, Md. : 1950)*, *133*(4), 1710–1715. https://pubmed.ncbi.nlm.nih.gov/6206131/

Ginisty, H., Sicard, H., Roger, B., & Bouvet, P. (1999). Structure and functions of nucleolin. *Journal of Cell Science*, *112*(6), 761–772. https://doi.org/10.1242/JCS.112.6.761

Greco, A. (2009). Involvement of the nucleolus in replication of human viruses. *Reviews in Medical Virology*, *19*(4), 201–214. https://doi.org/10.1002/RMV.614

Greco, A., Arata, L., Soler, E., Gaume, X., Couté, Y., Hacot, S., Callé, A., Monier, K., Epstein, A. L., Sanchez, J.-C., Bouvet, P., & Diaz, J.-J. (2012). Nucleolin interacts with US11 protein of herpes simplex virus 1 and is involved in its trafficking. *Journal of Virology*, *86*(3), 1449–1457. https://doi.org/10.1128/JVI.06194-11

Gupta, S. K., Sahoo, A. P., Rosh, N., Gandham, R. K., Saxena, L., Singh, A. K., Harish, D. R., & Tiwari, A. K. (2016). Canine parvovirus NS1 induced apoptosis involves mitochondria, accumulation of reactive oxygen species and activation of caspases. *Virus Research*, *213*, 46–61. https://doi.org/10.1016/j.virusres.2015.10.019

Hernandez-Verdun, D. (2011). Assembly and disassembly of the nucleolus during the cell cycle. *Nucleus (Austin, Tex.)*, *2*(3), 189–194. https://doi.org/10.4161/NUCL.2.3.16246

Hindley, C. E., Davidson, A. D., & Matthews, D. A. (2007). Relationship between adenovirus DNA replication proteins and nucleolar proteins B23.1 and B23.2. *The Journal of General Virology*, *88*(Pt 12), 3244–3248. https://doi.org/10.1099/VIR.0.83196-0

Hiscox, J. A. (2007). RNA viruses: hijacking the dynamic nucleolus. *Nature Reviews. Microbiology*, *5*(2), 119–127. https://doi.org/10.1038/NRMICRO1597

Iarovaia, O. v., Minina, E. P., Sheval, E. v., Onichtchouk, D., Dokudovskaya, S., Razin, S. v., & Vassetzky, Y. S. (2019). Nucleolus: A Central Hub for Nuclear Functions. *Trends in Cell Biology*, *29*(8), 647–659. https://doi.org/10.1016/J.TCB.2019.04.003

Ihalainen, T. O., Niskanen, E. A., Jylhävä, J., Paloheimo, O., Dross, N., Smolander, H., Langowski, J., Timonen, J., & Vihinen-Ranta, M. (2009). Parvovirus induced alterations in nuclear architecture and dynamics. *PLoS ONE*, *4*(6). https://doi.org/10.1371/journal.pone.0005948

Jia, W., Yao, Z., Zhao, J., Guan, Q., & Gao, L. (2017). New perspectives of physiological and pathological functions of nucleolin (NCL). *Life Sciences*, *186*(May), 1–10. https://doi.org/10.1016/j.lfs.2017.07.025

Johnson, J. S., & Samulski, R. J. (2009). Enhancement of adeno-associated virus infection by mobilizing capsids into and out of the nucleolus. *Journal of Virology*, *83*(6), 2632–2644. https://doi.org/10.1128/JVI.02309-08

Kim, K., Dimitrova, D. D., Carta, K. M., Saxena, A., Daras, M., & Borowiec, J. A. (2005). Novel checkpoint response to genotoxic stress mediated by nucleolin-replication protein a complex formation. *Molecular and Cellular Biology*, *25*(6), 2463–2474. https://doi.org/10.1128/MCB.25.6.2463-2474.2005

Kumar, D., Broor, S., & Rajala, M. S. (2016). Interaction of Host Nucleolin with Influenza A Virus Nucleoprotein in the Early Phase of Infection Limits the Late Viral Gene Expression. *PloS One*, *11*(10). https://doi.org/10.1371/JOURNAL.PONE.0164146

Lafontaine, D. L. J., Riback, J. A., Bascetin, R., & Brangwynne, C. P. (2021). The nucleolus as a multiphase liquid condensate. *Nature Reviews. Molecular Cell Biology*, *22*(3), 165–182. https://doi.org/10.1038/S41580-020-0272-6

Lam, Y. W., Evans, V. C., Heesom, K. J., Lamond, A. I., & Matthews, D. A. (2010). Proteomics Analysis of the Nucleolus in Adenovirus-infected Cells. *Molecular & Cellular Proteomics : MCP*, *9*(1), 117. https://doi.org/10.1074/MCP.M900338-MCP200

Lee, T. W. R., Blair, G. E., & Matthews, D. A. (2003). Adenovirus core protein VII contains distinct sequences that mediate targeting to the nucleus and nucleolus, and colocalization with human chromosomes. *Journal of General Virology*, *84*(12), 3423–3428. https://doi.org/10.1099/VIR.0.19546-0/CITE/REFWORKS

Lymberopoulos, M. H., & Pearson, A. (2007). Involvement of UL24 in herpes-simplex-virus-1-induced dispersal of nucleolin. *Virology*, *363*(2), 397–409. https://doi.org/10.1016/J.VIROL.2007.01.028

MacCallum, D. E., & Hall, P. A. (2000). The biochemical characterization of the DNA binding activity of pKi67. *The Journal of Pathology*, *191*(3), 286–298. https://doi.org/10.1002/1096-9896(2000)9999:9999<::aid-path628>3.0.co;2-j

Matthews, D. A. (2001). Adenovirus Protein V Induces Redistribution of Nucleolin and B23 from Nucleolus to Cytoplasm. *Journal of Virology*, *75*(2), 1031–1038. https://doi.org/10.1128/JVI.75.2.1031-1038.2001

Matthews, D., Emmott, E., & Hiscox, J. (2011). Viruses and the nucleolus. *Protein Reviews*, *15*, 321–345. https://doi.org/10.1007/978-1-4614-0514-6_14/FIGURES/1_14

Mattola, S., Mäntylä, E., Aho, V., Salminen, S., Oittinen, M., Salokas, K., Järvensivu, J., Hakanen, S., Ihalainen, T., Viiri, K., & Vihinen-Ranta, Maija. (n.d.). G2/M transition and apoptosis lead to facilitated nuclear egress of parvoviral capsids. *Manuscript*.

Mattola, S., Salokas, K., Aho, V., Mäntylä, E., Salminen, S., Hakanen, S., Niskanen, E. A., Svirskaite, J., Ihalainen, T. O., Airenne, K. J., Kaikkonen-Määttä, M., Parrish, C. R., Varjosalo, M., & Vihinen-Ranta, M. (2022). Parvovirus nonstructural protein 2 interacts with chromatin-regulating cellular proteins. *PLoS Pathogens*, *18*(4), e1010353. https://doi.org/10.1371/JOURNAL.PPAT.1010353

McStay, B. (2016). Nucleolar organizer regions: genomic "dark matter" requiring illumination. *Genes & Development*, *30*(14), 1598–1610. https://doi.org/10.1101/GAD.283838.116

Moehler, M., Zeidler, M., Schede, J., Rommelaere, J., Galle, P. R., Cornelis, J. J., & Heike, M. (2003). Oncolytic parvovirus H1 induces release of heat-shock protein HSP72 in susceptible human tumor cells but may not affect primary immune cells. *Cancer Gene Therapy*, *10*(6), 477–480. https://doi.org/10.1038/sj.cgt.7700591

Myllys, M., Ruokolainen, V., Aho, V., Smith, E. A., Hakanen, S., Peri, P., Salvetti, A., Timonen, J., Hukkanen, V., Larabell, C. A., & Vihinen-Ranta, M. (2016). Herpes simplex virus 1 induces egress channels through marginalized host chromatin. *Scientific Reports*, *6*(May), 1–9. https://doi.org/10.1038/srep28844

Naeger, L. K., Cater, J., & Pintel, D. J. (1990). The small nonstructural protein (NS2) of the parvovirus minute virus of mice is required for efficient DNA replication and infectious virus production in a cell-type-specific manner. *Journal of Virology*, *64*(12), 6166–6175. https://doi.org/10.1128/JVI.64.12.6166-6175.1990

Naeger, L. K., Salomé, N., & Pintel, D. J. (1993). NS2 is required for efficient translation of viral mRNA in minute virus of mice-infected murine cells. *Journal of Virology*, *67*(2), 1034–1043. https://doi.org/10.1128/JVI.67.2.1034-1043.1993

Németh, A., & Längst, G. (2011). Genome organization in and around the nucleolus. *Trends in Genetics : TIG*, *27*(4), 149–156. https://doi.org/10.1016/J.TIG.2011.01.002

Niskanen, E. A., Ihalainen, T. O., Kalliolinna, O., Häkkinen, M. M., & Vihinen-Ranta, M. (2010). Effect of ATP binding and hydrolysis on dynamics of canine parvovirus NS1. *Journal of Virology*, *84*(10), 5391–5403. https://doi.org/10.1128/JVI.02221-09

Niskanen, E. A., Kalliolinna, O., Ihalainen, T. O., Häkkinen, M., & Vihinen-Ranta, M. (2013). Mutations in DNA Binding and Transactivation Domains Affect the Dynamics of Parvovirus NS1 Protein. *Journal of Virology*, *87*(21), 11762–11774. https://doi.org/10.1128/jvi.01678-13

Okuwaki, M., Iwamatsu, A., Tsujimoto, M., & Nagata, K. (2001). Identification of nucleophosmin/B23, an acidic nucleolar protein, as a stimulatory factor for in vitro replication of adenovirus DNA complexed with viral basic core proteins. *Journal of Molecular Biology*, *311*(1), 41–55. https://doi.org/10.1006/JMBI.2001.4812

Olausson, K. H., Nistér, M., & Lindström, M. S. (2012). p53 -Dependent and -Independent Nucleolar Stress Responses. *Cells*, *1*(4), 774–798. https://doi.org/10.3390/CELLS1040774

Otsu, N. (1979). A Threshold Selection Method from Gray-Level Histograms. *IEEE Transactions on Systems, Man, and Cybernetics*, *9*(1), 62–66. https://doi.org/10.1109/TSMC.1979.4310076

Padeken, J., & Heun, P. (2014). Nucleolus and nuclear periphery: Velcro for heterochromatin. *Current Opinion in Cell Biology*, *28*(1), 54–60. https://doi.org/10.1016/J.CEB.2014.03.001

Parrish, C. R. (1991). Mapping specific functions in the capsid structure of canine parvovirus and feline panleukopenia virus using infectious plasmid clones. *Virology*, *183*(1), 195–205. https://doi.org/10.1016/0042-6822(91)90132-u

Patel, B. N., Rosenberg, L., Willcox, G., Baltaxe, D., Lyons, M., Irvin, J., Rajpurkar, P., Amrhein, T., Gupta, R., Halabi, S., Langlotz, C., Lo, E., Mammarappallil, J., Mariano, A. J., Riley, G., Seekins, J., Shen, L., Zucker, E., & Lungren, M. (2019). Human-machine partnership with artificial intelligence for chest radiograph diagnosis. *NPJ Digital Medicine*, *2*(1). https://doi.org/10.1038/S41746-019-0189-7

Pearson, A., Bourget, A., ben Abdeljelil, N., & Lymberopoulos, M. H. (2011). Role of the viral protein UL24 in nucleolar modifications induced by herpes simplex virus 1. *BMC Proceedings 2011 5:1*, *5*(1), 1–1. https://doi.org/10.1186/1753-6561-5-S1-P102

Pederson, T. (2011). The nucleolus. *Cold Spring Harbor Perspectives in Biology*, *3*(3), 1–15. https://doi.org/10.1101/CSHPERSPECT.A000638

Qi, Y., & Zhang, B. (2021). Chromatin network retards nucleoli coalescence. *Nature Communications*, *12*(1). https://doi.org/10.1038/S41467-021-27123-9

Raška, I., Shaw, P. J., & Cmarko, D. (2006). Structure and function of the nucleolus in the spotlight. *Current Opinion in Cell Biology*, *18*(3), 325–334. https://doi.org/10.1016/J.CEB.2006.04.008

Salvetti, A., & Greco, A. (2014). Viruses and the nucleolus: The fatal attraction. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, *1842*(6), 840–847. https://doi.org/10.1016/J.BBADIS.2013.12.010

Saxena, L., Kumar, G. R., Saxena, S., Chaturvedi, U., Sahoo, A. P., Singh, L. V., Santra, L., Palia, S. K., Desai, G. S., & Tiwari, A. K. (2013). Apoptosis induced by NS1 gene of Canine Parvovirus-2 is caspase dependent and p53 independent. *Virus Research*, *173*(2), 426–430. https://doi.org/10.1016/J.VIRUSRES.2013.01.020

Scherl, A., Couté, Y., Déon, C., Callé, A., Kindbeiter, K., Sanchez, J.-C., Greco, A., Hochstrasser, D., & Diaz, J.-J. (2002). Functional Proteomic Analysis of Human Nucleolus □ D. *Molecular Biology of the Cell*, *13*, 4100–4109. https://doi.org/10.1091/mbc.E02-05-0271

Sobecki, M., Mrouj, K., Camasses, A., Parisis, N., Nicolas, E., Llères, D., Gerbe, F., Prieto, S., Krasinska, L., David, A., Eguren, M., Birling, M. C., Urbach, S., Hem, S., Déjardin, J., Malumbres, M., Jay, P., Dulic, V., Lafontaine, D. L. J., … Fisher, D. (2016). The cell proliferation antigen Ki-67 organises heterochromatin. *ELife*, *5*(MARCH2016). https://doi.org/10.7554/ELIFE.13722

Sobecki, M., Mrouj, K., Colinge, J., Gerbe, F., Jay, P., Krasinska, L., Dulic, V., & Fisher, D. (2017). Cell-Cycle Regulation Accounts for Variability in Ki-67 Expression Levels. *Cancer Research*, *77*(10), 2722–2734. https://doi.org/10.1158/0008-5472.CAN-16-0707

Srivastava, M., & Pollard, H. B. (1999). Molecular dissection of nucleolin's role in growth and cell proliferation: new insights. *The FASEB Journal*, *13*(14), 1911–1922. https://doi.org/10.1096/FASEBJ.13.14.1911

Stamatiou, K., & Vagnarelli, P. (2021). Chromosome clustering in mitosis by the nuclear protein Ki-67. *Biochemical Society Transactions*, *49*(6), 2767–2776. https://doi.org/10.1042/BST20210717

Stenström, L., Mahdessian, D., Gnann, C., Cesnik, A. J., Ouyang, W., Leonetti, M. D., Uhlén, M., Cuylen-Haering, S., Thul, P. J., & Lundberg, E. (2020). Mapping the nucleolar proteome reveals a spatiotemporal organization related to intrinsic protein disorder. *Mol Syst Biol*, *16*, 9469. https://doi.org/10.15252/msb.20209469

Suikkanen, S., Sääjärvi, K., Hirsimäki, J., Välilehto, O., Reunanen, H., Vihinen-Ranta, M., & Vuento, M. (2002). Role of Recycling Endosomes and Lysosomes in Dynein-Dependent Entry of Canine Parvovirus. *Journal of Virology*, *76*(9), 4401–4411. https://doi.org/10.1128/JVI.76.9.4401-4411.2002

Waggoner, S., & Sarnow, P. (1998). Viral Ribonucleoprotein Complex Formation and Nucleolar-Cytoplasmic Relocalization of Nucleolin in Poliovirus-Infected Cells. *Journal of Virology*, *72*(8), 6699–6709. https://doi.org/10.1128/JVI.72.8.6699-6709.1998

Wang, D., Yuan, W., Davis, I., & Parrish, C. R. (1998). Nonstructural protein-2 and the replication of canine parvovirus. *Virology*, *240*(2), 273–281. https://doi.org/10.1006/VIRO.1997.8946

Weichert, W. S., Parker, J. S. L., Wahid, A. T. M., Chang, S. F., Meier, E., & Parrish, C. R. (1998). Assaying for structural variation in the parvovirus capsid and its role in infection. *Virology*, *250*(1), 106–117. https://doi.org/10.1006/VIRO.1998.9352

Wistuba, A., Kern, A., Weger, S., Grimm, D., & Kleinschmidt, J. A. (1997). Subcellular compartmentalization of adeno-associated virus type 2 assembly. *Journal of Virology*, *71*(2), 1341–1352. https://doi.org/10.1128/JVI.71.2.1341-1352.1997

Yang, K., Yang, J., & Yi, J. (2018). Nucleolar Stress: hallmarks, sensing mechanism and diseases. *Cell Stress*, *2*(6), 125–140. https://doi.org/10.15698/CST2018.06.139

Yeung, D. E., Brown, G. W., Tam, P., Russnak, R. H., Wilson, G., Clark-Lewis, I., & Astell, C. R. (1991). Monoclonal antibodies to the major nonstructural nuclear protein of minute virus of mice. *Virology*, *181*(1), 35–45. https://doi.org/10.1016/0042-6822(91)90467-P