Characterization of two conserved cell death elicitor families from the Dothideomycete fungal pathogens *Dothistroma septosporum* and *Fulvia fulva* (syn. *Cladosporium fulvum*)

Mariana Tarallo[1,*], Rebecca L. McDougal[2], Zhiyuan Chen[3], Yan Wang[3], Rosie E. Bradshaw[1,*],
Carl H. Mesarich[4,*]

[1]Laboratory of Molecular Plant Pathology/Bioprotection Aotearoa, School of Natural Sciences,
Massey University, Private Bag 11222, Palmerston North 4442, New Zealand

[2]Scion, Titokorangi Drive, Private Bag 3020, Rotorua 3046, New Zealand

[3]Department of Plant Pathology, Nanjing Agricultural University 210095, Nanjing, China

[4]Laboratory of Molecular Plant Pathology/Bioprotection Aotearoa, School of Agriculture and
Environment, Massey University, Private Bag 11222, Palmerston North 4442, New Zealand

*Correspondence:
Corresponding authors: M. Tarallo (m.tarallo@massey.ac.nz), R. E. Bradshaw
(r.e.bradshaw@massey.ac.nz) and C. H. Mesarich (c.mesarich@massey.ac.nz)
12

Keywords: Dothideomycetes, cell death-eliciting effectors, protein families, immune receptors,
plant-pathogen interactions, protein tertiary structure.

Abstract

Dothistroma septosporum (Ds) and *Fulvia fulva* (Ff; previously called *Cladosporium fulvum*) are two
closely related Dothideomycete fungal species that cause Dothistroma needle blight in pine and leaf
mold in tomato, respectively. During host colonization, these pathogens secrete virulence factors
termed effectors to promote infection. In the presence of corresponding host immune receptors,
however, these effectors activate plant defenses, including a localized cell death response that halts
pathogen growth. We identified two effector protein families, Ecp20 and Ecp32, which are conserved
between the two pathogens. The Ecp20 family has four paralogues in both species, while the Ecp32
family has four paralogues in *D. septosporum* and five in *F. fulva*. Both families have members that
are highly expressed during host infection. Members of the Ecp20 family have predicted structural
similarity to proteins with a β-barrel fold, including the Alt a 1 allergen from *Alternaria alternata*,
while members of the Ecp32 family have predicted structural similarity to proteins with a β-trefoil
fold, such as trypsin inhibitors and lectins. Using *Agrobacterium tumefaciens*-mediated transient
transformation assays, each family member was assessed for its ability to trigger cell death in leaves
of the non-host species *Nicotiana benthamiana* and *N. tabacum*. Using this approach, FfEcp20-2,
DsEcp20-3 and FfEcp20-3 from the Ecp20 family, and all members from the Ecp32 family, except
for the Ds/FfEcp32-4 pair, triggered cell death in both species. This cell death was dependent on
secretion of the effectors to the apoplast. In line with recognition by an extracellular immune
receptor, cell death triggered by Ds/FfEcp20-3 and FfEcp32-3 was compromised in *N. benthamiana*
silenced for *BAK1* or *SOBIR1*, which encode extracellular co-receptors involved in transducing
defense response signals following apoplastic effector recognition. We then investigated whether
DsEcp20-3 and DsEcp20-4 triggered cell death in the host species *Pinus radiata* by directly
infiltrating purified protein into pine needles. Strikingly, as in the non-host species, DsEcp20-3

triggered cell death, while DsEcp20-4 did not. Collectively, our study describes two new candidate
effector families with cell death-eliciting activity from *D. septosporum* and *F. fulva* and provides
evidence that members of these families are recognized by plant immune receptors.

1   Introduction

Understanding how plants and fungal pathogens interact at the molecular level can lead to
improvements in plant health. During infection, fungal pathogens deliver a collection of small,
proteinaceous virulence factors, termed effectors, into the apoplast and/or cells of their host plants to
promote the infection process (Rocafort et al., 2020; Wang et al., 2020). However, as fungal infection
has a negative impact on plant fitness, there is a strong selection pressure on the host to recognize
these effectors as invasion patterns and, in doing so, activate defense responses that provide
protection against the pathogen (Cook et al., 2015) In line with this, in resistant hosts, particular
effectors may be recognized by corresponding specific immune receptors, called resistance (R)
proteins, to activate plant defense responses. Often, one of these defense responses is the
hypersensitive response (HR), a localized cell death reaction that limits or halts pathogen growth
(Cook et al., 2015; Win et al., 2012). As these defense responses often lead to the pathogen being
unable to cause disease, such recognized effectors are typically referred to as avirulence (Avr)
determinants. To circumvent recognition, and therefore the ensuing plant defense responses, the
fungal invader must then modify its effector arsenal by secreting mutated or new effectors, shifting
the pressure to adapt back onto the plant host, resulting in a constant "arms race" in plant-pathogen
interactions (Coll et al., 2011; Jones & Dangl, 2006).

Progress in the generation of genomic, transcriptomic, and proteomic resources has enabled the rapid
identification of pathogen effectors, making it possible to screen them for elicitation or suppression
of plant defense responses, a process known as effectoromics. This, in turn, makes it possible to
discover host defense or susceptibility genes in different plant species. A search for conserved
effectors, sometimes called 'core' effectors, is essential as it can lead to broad spectrum resistance.
The tertiary structures of some effector proteins are conserved across microbial classes because they
have fundamental and important roles in infection, and for this reason are less likely to be mutated or
lost during microbial evolution (Nürnberger & Brunner, 2002). Therefore, plant resistance based on
the recognition of conserved effectors is more likely to be durable, as it is less likely to be overcome.

Dothistroma septosporum (Ds) is a hemibiotrophic pathogen that causes Dothistroma needle blight
(DNB), one of the most important foliar diseases of pine trees (Drenkhan et al., 2016). Based on
genomic information, *D. septosporum* is closely related to *Fulvia fulva* (Ff; syn. *Cladosporium*
fulvum) (de Wit et al., 2012). *F. fulva* is a biotrophic pathogen responsible for leaf mold of tomato
(*Solanum lycopersicum*), a disease that is mainly a problem in greenhouse or high tunnel
environments (Thomma et al., 2005). In both the *D. septosporum*–pine and *F. fulva*–tomato
pathosystems, infection begins with the germination of conidia on the leaf surface and penetration of
hyphae into the host through the stomata. Once inside, both pathogens colonize the apoplastic space
between mesophyll cells, where they secrete effector proteins (Kabir et al., 2015; Mesarich et al.,
2018).

F. fulva is one of the best studied filamentous fungal pathogens. The *F. fulva*–tomato interaction is a
model pathosystem in which early pioneering studies helped to establish the molecular basis of the
gene-for-gene concept of plant resistance to fungal pathogens (de Wit, 2016). In *F. fulva*, small,
secreted cysteine-rich proteins delivered to the host apoplast, termed extracellular proteins (Ecps)
based on their identification in apoplastic wash fluid samples, have been well-studied (Mesarich et

al., 2018). These include core effectors such as Ecp2-1, Ecp6 and Avr4 (de Jonge et al., 2010; Sánchez-Vallet et al., 2013; Stergiopoulos et al., 2010; van den Burg et al., 2006), homologs of which have been found in other plant-pathogenic fungi, including *D. septosporum* (Bolton et al., 2008; de Wit et al., 2012; Stergiopoulos et al., 2012). In contrast, few studies have identified and characterized effectors from *D. septosporum*. In one of these studies, DsEcp2-1, the ortholog of Ecp2-1 from *F. fulva*, was suggested to be an Avr protein responsible for eliciting defense responses in pine (Guo et al., 2020). In another of these studies, DsAvr4, the ortholog of Avr4 from *F. fulva*, was found to bind chitin, a function that is required to protect the fungal cell wall against hydrolysis by host chitinases (Mesarich et al., 2016; van den Burg et al., 2006).

In general, studies highlighting the role of effectors in molecular interactions between pathogens and gymnosperm hosts are few and far between. The close phylogenetic relationship between *D. septosporum* and *F. fulva*, however, provides a unique opportunity for comparative analysis of core effector functions across angiosperm and gymnosperm pathosystems. Recently, a set of candidate effector proteins from *D. septosporum* was transiently expressed in the non-host plants, *Nicotiana benthamiana* and *N. tabacum*, to identify those that are involved in triggering plant defense responses (Hunziker et al., 2021). Several of these proteins, all with orthologs in *F. fulva*, elicited a plant cell death response, suggesting that they are recognized by extracellular immune receptors in these non-host plants, triggering non-host resistance. Amongst these we found two proteins, Ds70057 and Ds70694, that were of particular interest, as they are orthologs of Ecps from *F. fulva* (Hunziker et al., 2021; Mesarich et al., 2018). Both proteins appear to be part of protein families – the Ecp20 family and the Ecp32 family (Hunziker et al., 2021). The Ecp20 family is made up of four members in both species, while the Ecp32 family is made up of four members in *D. septosporum* and five members in *F. fulva*.

In this study, we characterized and compared Ecp20 and Ecp32 family members from *D. septosporum* and *F. fulva* through prediction of their protein tertiary structures and by assessing their cell death-eliciting capacity in non-host angiosperm plants. We also used a novel assay involving a pine tissue culture system to determine whether two of the Ecp20 family members from *D. septosporum* trigger cell death in the gymnosperm host, *Pinus radiata*. This work provides insights into conservation of core effector function and recognition across a broad range of plant species.

2   Methods

2.1   Microorganisms and Plants

D. septosporum NZE10 (de Wit et al., 2012), GUA2 and COLN (Bradshaw et al., 2019), as well as *D. pini* CBS 116487 (BioSample ID SAMN02254964) and *F. fulva* 0WU (de Wit et al., 2012), were used to identify candidate effectors for characterization in this study. *Escherichia coli* DH5α (Taylor et al., 1993) and *Agrobacterium tumefaciens* GV3101 (Holsters et al., 1980) were used for gene cloning and *A. tumefaciens*-mediated transient transformation assays (ATTAs), respectively. *N. tabacum* Wisconsin 38 and *N. benthamiana* were used as model non-host plants for ATTAs, while *N. benthamiana* was used for tobacco rattle virus (TRV)-mediated gene silencing experiments. *P. radiata* clonal shoots, derived from family seed lots that are, compared to each other, relatively susceptible (S6 and S11) or tolerant (R4) to DNB, and grown from embryogenic tissue under sterile conditions on LPch agar (Hargreaves et al., 2004), were provided by Scion (New Zealand Forest Research Institute Ltd., Rotorua, New Zealand) and used in protein infiltration assays.

2.2   Bioinformatics

Members of the Ecp20 and Ecp32 protein families were identified from the predicted proteomes
and/or genomes of *D. septosporum* NZE10 and *F. fulva* 0WU (de Wit et al., 2012). Chromosome
locations of the *FfEcp20* and *FfEcp32* genes were identified using the *F. fulva* Race 5 genome
assembly (Zaccaron et al., 2022). Gene family members were also identified from genome sequences
of 18 other *D. septosporum* isolates (Bradshaw et al., 2019) and *D. pini* CBS 116487 (GenBank
assembly accession GCA_002116355.1), using reciprocal BLASTp and tBLASTn searches in
conjunction with an E-value cut-off of $<10^{-5}$. BLASTp was also employed to determine the level of
conservation for each family member across fungal species in the Joint Genome Institute (JGI)
MycoCosm database, using an E-value threshold of 10^{-5}. EffectorP v3.0 (Sperschneider & Dodds,
2021) was used to predict whether members of the Ecp20 and Ecp32 families are effectors. All *D.
septosporum* NZE10, *D. pini* CBS 116487 and *F. fulva* 0WU nucleotide and amino acid sequences
for Ecp20 and Ecp32 family members are shown in Supplementary Table S1.

Protein tertiary structure predictions were performed using AlphaFold2 (Jumper et al., 2021; Mirdita
et al., 2022), and the Dali server (Holm, 2020) was used to identify proteins with structural similarity
in the Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB). Here, a
Dali Z-score of 2 or greater was used to infer structural similarity. Protein tertiary structures were
visualized and rendered in PyMol v2.5, with alignments carried out using the CEalign tool (DeLano,
2002).

Protein sequence alignments were performed using Clustal Ω (Sievers et al., 2011) in Jalview
(www.jalview.org). Phylogenetic trees were constructed with the neighbor-joining method using
Geneious Software v9.1.8 (Kearse et al., 2012).

2.3   ATTA Expression Vectors

Candidate effector genes of *D. septosporum* NZE10, GUA2 and COLN, *D. pini* CBS 116487 and *F.
fulva* 0WU were cloned in the pICH86988 ATTA expression vector, which contains the nucleotide
sequence encoding the *N. tabacum* PR1α signal peptide for secretion into the plant apoplast, followed
by a 3×FLAG for detection via Western blotting (Weber et al. (2011), according to the method
described by Guo et al. (2020) in cases where the genes had no predicted introns, or synthesized
directly into pICH86988 by Twist Bioscience (San Francisco, CA, USA) in cases where the genes
had predicted introns. For the genes that were cloned into pICH86988, the mature coding sequence
was first amplified by polymerase chain reaction (PCR) from genomic DNA extracted from the
fungus, with genomic DNA extraction performed as described by Doyle and Doyle (1987). PCRs
were performed with Phusion High-Fidelity DNA polymerase (New England Biolabs, Beverly, MA,
USA), and PCR products were extracted from agarose gels using an E.Z.N.A.® Gel Extraction Kit
(Omega Bio-tek, Norcross, GA, USA). PCR primers were designed with Primer3 (Untergasser et al.,
2012) in Geneious v9.1.8 (Kearse et al., 2012). These primers, which were synthesized by Integrated
DNA Technologies (IDT; Supplementary Table S2), had 5' extensions containing a BsaI recognition
site and 4-base overhang specific for Golden Gate modular cloning (Engler et al., 2009) in between
the CaMV 35S promoter and octopine synthase terminator of pICH86988. Versions of the candidate
effector genes were also ligated into pICH86988 without the nucleotide sequence encoding the PR1α
signal peptide to prevent secretion of the proteins they encode into the apoplast. Constructs were then
transformed into *E. coli*, plasmids extracted using an E.Z.N.A.® Plasmid DNA Mini Kit I (Omega
Bio-tek), and correct assemblies confirmed by sequencing. ATTA expression vectors were
transformed into *A. tumefaciens* by electroporation as described previously (Guo et al., 2020).

2.4   ATTAs

To heterologously express candidate effector genes in *N. benthamiana* and *N. tabacum* using ATTAs,
single colonies of *A. tumefaciens* carrying pICH86988 constructs of interest were first inoculated into
selective lysogeny broth (LB) containing 50 µg/mL kanamycin, 10 µg/mL rifampicin and 30 µg/mL
gentamycin, then incubated overnight at 28ºC. Cells were subsequently collected by centrifugation at
2,500 *g* for 5 min and resuspended in 1 mL of infiltration buffer (10 mM $MgCl_2:6H_2O$, 10 mM MES-
KOH (Sigma-Aldrich, St. Louis, MO, USA), 100 µM acetosyringone (Sigma-Aldrich)) to an OD_{600}
of 0.5. Following resuspension, the abaxial surface of *N. benthamiana* and *N. tabacum* leaves was
infiltrated with the appropriate *A. tumefaciens* suspension using 1 mL needleless syringes. Here, the
extracellular elicitin INF1 from *Phytophthora infestans*, which is recognized by the LRK1 immune
receptor in *Nicotiana* species (Kamoun et al., 1997; Kanzaki et al., 2008), was used as a positive cell
death control, while the empty pICH86988 vector was used as a negative no cell death control. At
least 12–24 infiltration zones were tested for each candidate effector gene or control (consisting of
two leaves from each of two plants ×3–6 repeat experiments). Plants were monitored for up to 7 days,
and photographs taken using a Nikon D7000 camera. Infiltration zones were assessed on whether the
protein tested elicited a strong or weak cell death response, or no cell death. A strong cell death
response was scored when it was indistinguishable from the response elicited by INF1 and around the
entire infiltration zone, while a weak response was considered when the protein elicited a weaker
response around approximately 50% of the infiltration zone when compared with INF1. A no-cell
death response was indicated when the infiltration zone had the same response as the negative empty
pICH86988 vector control.

2.5 Western Blotting

To confirm that candidate effector proteins had been produced in *N. benthamiana* and *N. tabacum*
leaves using ATTAs, and to determine whether these proteins were of the predicted size, Western
blotting was conducted with total protein as described previously by Guo et al. (2020). For protein
separation by SDS-PAGE, 12% separating and 5% stacking gels were used, with proteins transferred
to PVDF membranes. Anti-FLAG® M2 primary antibody produced in mouse (Sigma-Aldrich),
together with anti-mouse secondary antibody produced in chicken (Santa Cruz Biotechnology,
Dallas, TX, USA) and SuperSignal® West Dura Extended Duration substrate (Thermo Fisher
Scientific, Waltham, MA, USA), were used for protein detection. Proteins were visualized using an
Azure Biosystems c600 Bioanalytical Imaging system (Azure Biosystems, Dublin, CA, USA).

2.6 Protein Expression and Purification

Genes encoding candidate effector proteins were amplified by PCR as above from the previously
generated ATTA expression vectors, without their native signal peptide sequences. Here, the 5' PCR
primers incorporated a FLAG-tag for detection by Western blotting. PCR primers are shown in
Supplementary Table S2. PCR amplicons were gel-extracted as above and ligated, using DNA ligase
(New England Biolabs), into the SmaI/EcoRI restriction sites of the expression vector pPic9-His_6
(Invitrogen, Carlsberg, CA, USA), which encodes an α-factor signal peptide for protein secretion and
a hexa-histidine (6×His) tag for protein purification. The ligated products were then transformed into
E. coli. The expression constructs were sequenced and contained the predicted mature sequence of
the candidate effectors fused to a 6×His tag at their N-terminus. The expression cassette was
linearized with SacI or SalI (New England Biolabs) restriction enzymes and transformed into *P.*
pastoris GS115, according to Kombrink (2012).

Protein expression in *P. pastoris* was performed according to Weidner et al. (2010). Briefly, protein
expression was induced in 200 mL of BMMY (Buffered Methanol-complex Medium), through the

addition of methanol to a concentration of 0.5% (v/v) every 24 h, for 72 h. The cells were then removed by centrifugation at 4,500 *g* for 30 min and the supernatant filter-sterilized by passage through a 0.22 µm membrane (ReliaPrep, Ahlstrom, Helsinki, Finland). Finally, the pH of the sterilized culture filtrate was adjusted to 8 with NaOH.

The secreted candidate effector protein (fused with an N-terminal 6×His tag) was purified from culture filtrate using immobilized metal ion affinity chromatography (IMAC), in conjunction with a Ni Sepharose™ 6 Fast Flow (GE Healthcare, Chicago, IL, USA) column, according to the manufacturer's protocol. Before loading, the column (Glass Econo-Column® - BioRad, Hercules, CA, USA) was packed with 5 mL of resin and equilibrated by washing with binding buffer (20 mM sodium phosphate, 0.5 M NaCl, pH 7.4). The culture filtrate was added to the column at a flow rate of 1 mL/min, and protein eluted with an elution buffer (20 mM sodium phosphate, 0.5 M NaCl and 500 mM imidazole, pH 7.4). The elution fractions were mixed, and elution buffer was added to a final volume of 50 mL to obtain sufficient volume for vacuum-infiltration of pine shoots. Western blotting was performed as above to determine the presence of each candidate effector protein in *P. pastoris* culture filtrate and after purification. Protein concentrations in culture filtrates and purified solutions were determined by using a Pierce™ Coomassie (Bradford) Protein Assay Kit (Thermo Fisher Scientific), as per the manufacturer's protocol.

2.7   Pine Infiltration with Purified Proteins

Proteins were infiltrated into tissue-cultured pine shoots at a concentration of approx. 20 µg/mL using the method developed by Hunziker et al. (2021). Briefly, clonal rootless pine shoots were maintained in LPch agar (Hargreaves et al., 2004) in glass jars, with each jar containing 6–8 shoots. For this experiment, clonal shoots originating from two relatively *Dothistroma*-susceptible families (S6 and S11) and one relatively *Dothistroma*-tolerant family (R4) (based on comparative field data) were used. The shoots were completely submerged in approx. 50 mL of each purified candidate protein or control solution (positive control: DsEcp32-3; negative control: elution buffer). Samples were exposed to vacuum in a glass chamber for 5 min to facilitate protein uptake. Shoots were then briefly rinsed in sterile MilliQ (MQ) water and subsequently placed back into the LPch agar in the glass jar. The experiment was conducted at 22°C in a room with natural light. Photos were taken 7 days after infiltration (dai) using a Nikon D7000 camera.

2.8   Tobacco Rattle Virus (TRV)-mediated Virus-induced Gene Silencing (VIGS)

Fragments used to generate virus-induced gene silencing (VIGS) constructs were amplified by PCR from cDNA or genomic DNA of *N. benthamiana* using PrimeSTAR GXL DNA Polymerase (Takara Bio Inc.) with the primers listed in Supplementary Table S2. The TRV VIGS system is composed of pTRV-RNA1, which encodes replication related proteins, and pTRV-RNA2, which harbors the viral coat protein, and the candidate fragment sequences that target silencing to the corresponding genes (Liu et al., 2002; MacFarlane, 1999). The purified fragments were cloned into the pTRV-RNA2 vector (Dong et al., 2007) using the ClonExpress II One Step Cloning Kit (Vazyme Biotech Co., Ltd, Nanjing, China). The resulting vectors were verified by sequencing and individually transformed into *A. tumefaciens* strain. DsEcp20-3 was screened against around 100 TRV VIGS constructs which targeted different *N. benthamiana* LRR-RLP and LRR-RLK-encoding genes. Information on this library of silencing constructs can be found in Supplementary Data 1 from Wang et al. (2018). Primers for *T26*, *T26-2*, TRV:*BAK1* (Heese et al., 2007) and TRV:*SOBIR1* (Liebrand et al., 2013) are shown in Supplementary Table S2.

A. tumefaciens strains carrying *N. benthamiana* gene silencing vectors (Wang et al., 2018) were
grown overnight in LB medium with the appropriate antibiotics. *A. tumefaciens* cells were pelleted as
mentioned above, resuspended, and incubated in infiltration medium (10 mM 2-[N-morpholino]
ethanesulfonic acid pH 5.6, 10 mM MgCl$_2$, 200 µM acetosyringone) for 3–4 h before
agroinfiltration. *A. tumefaciens* cultures expressing pTRV-RNA2 constructs were mixed in a 1:1 ratio
with *A. tumefaciens* culture expressing pTRV1 to a final OD$_{600}$ of 1.0 before infiltration into
cotyledons of four-leaf-stage *N. benthamiana* seedlings. *P. sojae* NPP1 and XEG1 (Fellbrich et al.,
2002; Ma et al., 2015), and *P. infestans* INF1 were used as cell death-eliciting controls for VIGS.
XEG1- and INF1-triggered cell death was known to be compromised in *N. benthamiana* leaves
treated with TRV:*BAK1* or TRV:*SOBIR1*, but not with TRV:*GFP*, which was used as a negative
control. (Wang et al., 2018).

2.9   RNA Isolation and Quantitative Reverse Transcription PCR

To determine the expression levels of *BAK1*, *SOBIR1* and *T26* in the TRV-mediated VIGS
experiment, *N. benthamiana* leaves of similar size were selected and ground to a powder in liquid
nitrogen. Total RNA was isolated from 100 mg of ground material using the E.Z.N.A.® Total RNA
Kit I (Omega Bio-tek) and used as a template for first strand cDNA synthesis with PrimeScript
Reverse Transcriptase (Takara Bio Inc., Kusatsu, Shiga, Japan). Quantitative reverse transcription
PCR (qRT-PCR) was performed on an ABI 7500 Fast Real-Time PCR system (Applied Biosystems
Inc., Foster City, CA, USA) using PrimeScript RT Master Mix (Takara Bio Inc.) with translation
elongation factor 1 alpha (*EF-1α*) as an endogenous control and the primers listed in Supplementary
Table S2. Data were analyzed using the $2^{-\Delta\Delta C_T}$ method (Livak & Schmittgen, 2001).

3   Results

3.1   Two *D. septosporum* Elicitors are Effector Candidates that Form Part of Protein Families with Orthologs in *F. fulva*

In our previous work, two small, secreted candidate effector proteins from *D. septosporum* NZE10,
Ds70694 and Ds70057, triggered cell death in the non-host species *N. tabacum* and *N. benthamiana*
(Hunziker et al., 2021). Both proteins are homologs of Ecps from *F. fulva* and are part of two
different protein families, Ecp20 and Ecp32, respectively (Mesarich et al., 2018). More specifically,
Ds70694 is an ortholog of FfEcp20-3 from *F. fulva* and was therefore named DsEcp20-3, whilst
Ds70057 is an ortholog of FfEcp32-3 and was therefore named DsEcp32-3.

The Ecp20 family has four paralogues in both species. Based on sequence similarities between them,
DsEcp20-3 and FfEcp20-3, together with the other orthologous pairs Ds/FfEcp20-1, Ds/FfEcp20-2
and Ds/FfEcp20-4, were grouped together (Figure 1A, Supplementary Figure S1A). Interspecies
pairwise amino acid identities between all family members from *D. septosporum* and *F. fulva* ranged
from 10.6% to 85.5%, while intraspecies pairwise identities between all family members ranged from
14.2% to 40.9% in *D. septosporum* and 11.8% to 44.2% in *F. fulva* (Supplementary Table S3). The
Ds/FfEcp20-1 genes contain one exon, with the intron located at the same position, while genes
Ds/FfEcp20-2, *-3* and *-4* do not have any introns (Supplementary Figure 2A). All genes encode a
protein with four cysteine residues, although the location of these varies slightly from protein pair to
protein pair (Supplementary Figures S1A and S2A).

Based on data from a previously published transcriptome study (Bradshaw et al., 2016), all genes of
the *D. septosporum* Ecp20 family are expressed during pine infection, with *DsEcp20-3* and *DsEcp20-*
4 being the two most highly expressed (Supplementary Table S4). *DsEcp20-3* showed strong up-

regulation in the Mid and Late *in planta* stages that mark the start and end of necrotrophic infection,
when compared with the Early (biotrophic) stage and *in vitro*, while *DsEcp20-4* is strongly up-
regulated in the Late *in planta* stage. For the corresponding genes in *F. fulva* during tomato infection,
as based on data from a previously published transcriptome study (Mesarich et al., 2018), *FfEcp20-2*
and *FfEcp20-3* were highly expressed throughout the timepoints analyzed, while *FfEcp20-4* was
highly expressed at the beginning of infection and downregulated at later time points, when
compared with *in vitro* (Supplementary Table S4). It should be noted that *FfEcp20-1* had no gene
model and therefore had no FPKM value assigned to it in the previous study (Mesarich et al., 2018).
Thus, no expression information was available for this gene.

The Ds/FfEcp32 family was analyzed in the same way as the Ds/FfEcp20 family above. The Ecp32
family has five paralogues in *F. fulva*, named FfEcp32-1 to -5, and four in *D. septosporum*, named
DsEcp32-1 and DsEcp32-3 to -5 (there is no ortholog of FfEcp32-2 in *D. septosporum*) (Figure 1B,
Supplementary Figure S1B). Interspecies pairwise amino acid identity between all family members
from *D. septosporum* and *F. fulva* ranged from 10.6% to 82%, while intraspecies pairwise identity
between members ranged from 22.1% to 42.1% in *D. septosporum* and 21.6% to 49% in *C. fulvum*
(Supplementary Table S3). Regarding the exon-intron structure, *Ds/FfEcp32-1*, *FfEcp32-2* and
Ds/FfEcp32-3 do not have any introns. In contrast, the *Ds/FfEcp32-4* gene pair has one intron at the
same position, while the *Ds/FfEcp32-5* gene pair has two introns at the same positions
(Supplementary Figure S2B). In terms of cysteine residues, the Ds/FfEcp32 proteins have anywhere
from none (Ds/FfEcp32-5) to five (Ds/FfEcp32-4), although in the case of DsEcp32-3, an additional
cysteine is present compared to the Ff ortholog (Supplementary Figures S1B and S2B).

All *DsEcp32* genes are also expressed during pine infection. However, only *DsEcp32-3* and
DsEcp32-5 are highly expressed *in planta*, especially at the Mid and Late necrotrophic stages when
compared with *in vitro*. *DsEcp32-1* and *DsEcp32-4* are only expressed at a low level *in planta*.
During *F. fulva* infection of tomato, *FfEcp32-1*, *FfEcp32-2* and *FfEcp32-3* are highly expressed at all
timepoints. *FfEcp32-4* was not expressed under any condition, while *FfEcp32-5* was only expressed
at 12 dpi, and was downregulated compared with *in vitro* (Supplementary Table S4).

Genomes from the two species are mesosyntenic (de Wit et al., 2012; Zaccaron et al., 2022) and, in
keeping with this, we found conservation of chromosome location between the two species for each
of the *Ds/FfEcp20* and *Ds/FfEcp32* gene pairs (Supplementary Figure S3). Taken together, sequence
similarity, intron position and chromosome localization suggest a common evolutionary origin for the
Ecp20 and *Ecp32* genes from both species.

Figure 1. Protein phylogenies of the *Dothistroma septosporum* and *Fulvia fulva* Ecp20 (A) and Ecp32 (B) families. The trees were constructed with the neighbor-joining method using Geneious Software v9.1.8. Bootstrap values are shown at the nodes as percentages. The scale bar represents 0.4 (A) and 0.2 (B) substitutions per site. Ds: *D. septosporum*; Ff: *F. fulva*.

3.2 Predicted Tertiary Structures of Ecp20 and Ecp32 Family Members Suggest Possible Roles in Virulence

We predicted the tertiary structures of all Ecp20 family members from *D. septosporum* NZE10 and *F. fulva* 0WU using AlphaFold2 (Figure 2; Supplementary Figure S4). All proteins from the Ecp20 family are predicted to be structurally similar to the characterized β-barrel fold of Alt a 1 from *Alternaria alternata* (RCSB protein data bank [PBD] ID: 3V0R) (Chruszcz et al., 2012) (Supplementary Table S4 and S5). Figure 2 shows the predicted structures for DsEcp20-3 and DsEcp20-4, as well as the alignments between DsEcp20-3 and the characterized structure of Alt a 1 from *A. alternata* and between the predicted structures of DsEcp20-3 and DsEcp20-4. The predicted structures for all the DsEcp20 and FfEcp20 proteins are shown in Supplementary Figure S4 along with characterized crystal structures of similar proteins from other fungal species.

The tertiary structure of Ecp32 proteins was also predicted using AlphaFold2 (Figure 3, Supplementary Figure S5). All Ecp32 family members had significant hits with trypsin inhibitors and lectins; these are proteins that share a β-trefoil fold and are known virulence factors in plant-pathogenic fungi (Renko et al., 2012; Sabotič et al., 2019; Varrot et al., 2013) (Supplementary Table S5). Figure 3 shows the predicted structures for DsEcp32-3 and DsEcp32-4, as well as the alignments between DsEcp32-3 and the characterized structure of a trypsin inhibitor from the plant species *Enterolobium contortisiliquum* (RCSB PBD ID: 4J2K) (Zhou et al., 2013), which is the top hit in the PDB for most Ds/FfEcp32 proteins. The alignment between the predicted structures of DsEcp32-3 and DsEcp32-4 is also shown in Figure 3, while all the predicted Ds/FfEcp32 structures can be found in Supplementary Figure S5 along with characterized structures of similar proteins.

(A) DsEcp20-3

(B) DsEcp20-3 x Alt a 1

(C) DsEcp20-4

(D) DsEcp20-3 x DsEcp20-4

Figure 2. Predicted protein tertiary structures of Ecp20 family members from *Dothistroma septosporum.* (A) Predicted structure of DsEcp20-3. (B) Alignment of the predicted DsEcp20-3 structure (pink) with the characterized structure of Alt a 1 (black) from *Alternaria alternata* (RCSB protein data bank [PBD] ID: 3V0R) (Chruszcz et al., 2012). (C) Predicted structure of DsEcp20-4. (D) Alignment of predicted DsEcp20-3 (pink) and DsEcp20-4 (blue) structures. Disulphide bonds are shown as yellow sticks. Protein structures were predicted with AlphaFold2, rendered in PyMol v2.5 and aligned using the CEalign tool (DeLano, 2002; Jumper et al., 2021; Mirdita et al., 2022). Structures on the left were colored in PyMol according to their AlphaFold2 pLDDT score: dark blue

366 for regions predicted with high confidence, light blue and green for regions of low confidence, and
367 red for very low confidence regions.

Figure 3. Predicted protein tertiary structure of Ecp32 family members from *Dothistroma*
septosporum. (A) Predicted structure of DsEcp32-3. (B) Alignment of the predicted DsEcp32-3
(pink) structure with the characterized structure of trypsin inhibitor (black) from *Enterolobium*

contortisiliquum (RCSB protein data bank [PBD] ID: 4J2K) (Zhou et al., 2013). (C) Predicted structure of DsEcp32-4. (D) Alignment of predicted DsEcp32-3 (pink) and DsEcp32-4 (blue) structures. Disulphide bonds are shown as yellow sticks. Protein structures were predicted with AlphaFold2, rendered in PyMol v2.5 and aligned using the CEalign tool (DeLano, 2002; Jumper et al., 2021; Mirdita et al., 2022). Structures on the left were colored in PyMol according to their AlphaFold2 pLDDT score: dark blue for regions predicted with high confidence, light blue and green for regions of low confidence, and red for very low confidence regions.

3.3 Ecp20 and Ecp32 Family Members are Conserved Across Other Fungal Species and *D. septosporum* isolates

The Ecp20 family is present in fungal species from other classes. DsEcp20-1 had 27 significant BLASTp hits in the JGI MycoCosm protein database. Two of these were from other Dothideomycete species (including the ortholog in *F. fulva*), while the rest were with from species in the Eurotiomycetes class (Supplementary Table S6). DsEcp20-2 and DsEcp20-3, the most closely related to each other among the DsEcp20 family members (Figure 1A), both had the same 28 hits in JGI. All of these matches were with other Dothideomycete species and, of these, 24 were hits to proteins from other plant pathogens (Supplementary Table S6). DsEcp20-4 had 1,006 hits in the JGI MycoCosm protein database, of which 241 were proteins from plant pathogens (Supplementary Table S6).

In general, the Ecp32 family is more widely conserved among other fungal species compared to the Ecp20 family. Individual searches in the JGI MycoCosm protein database for each of the DsEcp32 family members yielded approximately 1,000 hits for each family member (Supplementary Table S6). Those hits included proteins from fungal species with different lifestyles, ranging from saprophytes, to biotrophs, hemibiotrophs and necrotrophs. For example, DsEcp32-3 had 158 hits to proteins from fungal plant pathogens in JGI. Of these, 42 were with other members from the Mycosphaerellaceae family. These included *Cercospora zeae-maydis* with four paralogs, *Pseudocercospora* (*Mycosphaerella*) *eumusae* with six, *Pseudocercospora fijiensis* with five and *Zymoseptoria tritici* with three paralogs (Supplementary Table S6).

Dothistroma pini is also a causal agent of DNB (Barnes et al., 2004). Thus, we searched for orthologs of DsEcp20/32 encoded in the *D. pini* genome. All proteins from the *D. septosporum* Ecp20 and Ecp32 families had orthologs in the genome of *D. pini*, except Ecp32-5, for which the gene encoding this protein appears to be a pseudogene. The protein sequences were very similar between family members from the two *Dothistroma* species, with full-length pairwise amino acid identity varying from 81.1% to 93.8% (Supplementary Tables S1 and S3).

We next addressed the question of whether the *D. septosporum Ecp20* and *Ecp32* family genes are conserved in other isolates of this species and whether they are identical in sequence to NZE10. Genome sequences were available from 18 *D. septosporum* isolates collected from different geographic locations (Bradshaw et al., 2019). All members of the *D. septosporum Ecp20* and *Ecp32* families were present in the 18 other genomes. In each of these genomes, there was at least one single nucleotide polymorphism (SNP) leading to an amino acid difference in at least one family member for both Ecp20 and Ecp32 (Supplementary Table S7).

A total of 59 different SNPs were identified among the *Ecp20* family members across the 18 *D. septosporum* isolates, compared to NZE10 (Figure 4). *DsEcp20-1*, *DsEcp20-2* and *DsEcp20-4* had approximately the same number of SNPs: 11, 13 and 15, respectively, while *DsEcp20-3* had a total of 20. For *DsEcp20-1*, *DsEcp20-3* and *DsEcp20-4*, however, most SNPs (61%, 60% and 60%, respectively) were synonymous. In the case of *DsEcp20-2*, however, only 30% were synonymous

and a higher number of SNPs leading to an amino acid difference (9 out of 13; 70%) were observed
(Figure 4A). The *Ecp32* gene family had a higher number of DNA modifications across the 18 *D.*
septosporum genomes analyzed, with a total of 117 SNPs and 3 insertions. The two most
polymorphic genes were *DsEcp32-3* and *DsEcp32-5*. Interestingly, while *DsEcp32-3* had 31
sequence modifications, only five of these led to an amino acid difference (16%). All other family
members had 9 or 10 SNPs leading to an amino acid difference (Figure 4B).

The Guatemalan (GUA1 and GUA2) and Colombian (COLN and COLS) isolates showed the greatest
amino acid sequence diversity, when compared to strain NZE10, for both the Ecp20 and Ecp32
family members. For DsEcp20-1, the most diverse isolates were COLN and COLS, with identical
sequences, while COLN had the most sequence variation in DsEcp32-1. For DsEcp20-2 and -4, and
Ecp32-3, -4, and -5, both Guatemalan isolates were the most diverse and, for DsEcp20-3, GUA2 had
greater amino acid diversity (Supplementary Table S7).

Figure 4. Schematic diagram showing sequence variation in *Dothistroma septosporum* NZE10 *Ecp20* (A) and *Ecp32* (B) family members across 18 other *D. septosporum* isolates. Dark grey boxes indicate signal peptide positions and light grey boxes indicate intron positions. Black flags show the position of nucleotide variations that led to non-synonymous changes, with the specific amino acid differences indicated above the flags. White flags show the position of synonymous nucleotide variations. Red flags show the position of nucleotide insertions. Vertical black lines show the positions of cysteine residues. Nucleotide positions are shown under each gene. SP: signal peptide.

3.4 Other Members of the Ecp20 and Ecp32 Families from *D. septosporum* and *F. fulva* Trigger Cell Death Responses in Non-host Plants

Following on from a previous study in which DsEcp20-3 and DsEcp32-3 were shown to elicit cell death in the non-host plants *N. benthamiana* and *N. tabacum* using ATTAs (Hunziker et al., 2021),

we set out to determine whether other family members of both *D. septosporum* and *F. fulva* could
elicit the same response. To achieve this, all Ecp20 and Ecp32 family members were transiently
expressed in *N. benthamiana* and *N. tabacum* using ATTAs, and the percentages of plant responses
in three different categories recorded: strong cell death, weak cell death and no cell death (Figure 5;
Supplementary Figure S6). Here, orthologous pairs from both pathogens were tested on the same
leaf, so that differences between them could be better assessed.

As expected, the positive control INF1 triggered strong cell death responses and the negative control
empty vector did not elicit a response. Among Ecp20 family members, FfEcp20-2 triggered strong or
weak cell death responses in *N. benthamiana* in 91.7% of infiltration zones, whilst its ortholog
DsEcp20-2 did not elicit any cell death responses. Both FfEcp20-3 and DsEcp20-3 caused consistent
plant cell death in *N. benthamiana*, with DsEcp20-3 having 58% of infiltration zones showing strong
cell death across the entire infiltration zone (Figure 5A, 5B). In *N. tabacum*, DsEcp20-3 was the only
family member that triggered a cell death response (Supplementary Figure S6). Interestingly,
DsEcp20-4, one of the most highly expressed secreted proteins during infection of pine by *D.*
septosporum (Supplementary Table S4; Bradshaw et al. (2016)), and its ortholog from *F. fulva*,
FfEcp20-4, did not induce cell death responses in either *N. benthamiana* or *N. tabacum* (Figure 5;
Supplementary Figure S6).

The ability of Ecp32 family members to trigger cell death was also assessed in both *Nicotiana*
species. In this case, all proteins from the Ecp32 family, apart from the Ds/FfEcp32-4 protein pair,
induced some degree of cell death in both plant species. Interestingly, DsEcp32-1 consistently
triggered cell death in *N. benthamiana*, while its ortholog from *F. fulva*, FfEcp32-1, only caused a
strong cell death response in 3.5% of infiltration zones across all replicates (Figure 5C, 5D). In
contrast, in *N. benthamiana*, FfEcp32-3 was capable of triggering consistent strong cell death
responses, while DsEcp32-3 only trigged a strong response in approximately 20% of infiltration
zones. The Ds/FfEcp32-5 pair triggered similar degrees of cell death responses in *N. benthamiana*
(Figure 5D).

To determine whether apoplastic localization is required for members of the Ecp20 and Ecp32
families to trigger cell death, proteins that previously elicited a plant response, but lacked the PR1α
signal peptide for secretion to the apoplast, were transiently expressed in *N. benthamiana*. Without
the signal peptide, none of proteins tested were able to cause cell death (Supplementary Figure S7
and Supplementary Figure S8).

In cases where a cell death response was not observed at all, or where only weak cell death was
observed, a Western blot analysis was performed to confirm the presence of the proteins in the tissue
of the infiltrated plants (Supplementary Figure S8). All proteins that did not trigger cell death or only
triggered a weak cell death response were detected via Western blot, indicating that the lack of cell
death response was not due to absence of those proteins.

Taken together, these results indicate that Ecp20 and Ecp32 family members from two closely related
fungal pathogens, *D. septosporum* and *F. fulva*, differ in their ability to induce a cell death response
in non-host species. Furthermore, the finding that proteins without a signal peptide failed to trigger a
plant cell death response, suggests that secretion into the apoplastic environment of *N. benthamiana*
is essential for their cell death-eliciting activity.

Figure 5. Ecp20 and Ecp32 proteins from *Dothistroma septosporum* NZE10 and *Fulvia fulva* 0WU trigger cell death responses in the non-host plant *Nicotiana benthamiana*. Ecp20 (A) and Ecp32 (C) family members from *D. septosporum* and *F. fulva* were expressed in *N. benthamiana* using an *Agrobacterium tumefaciens*-mediated transient expression assay (ATTA) to assess their ability to elicit cell death. Representative images are shown (n = 12–24 infiltration zones), from at least three independent experiments. INF1, *Phytophthora infestans* elicitin positive cell death control; EV, empty vector negative no cell death control; Ds: *D. septosporum*; Ff: *F. fulva*. Graphs display the percentages of infiltration zones that showed cell death in response to Ecp20 (B) and Ecp32 (D) family members from *F. fulva* and *D. septosporum*, divided into three categories: strong cell death, weak cell death and no cell death. Photos were taken 7 days after infiltration.

3.5 Proteins from other *D. septosporum* Isolates and *D. pini* also Trigger Cell Death in Non-host Plants

Alleles of some *Ecp20* and *Ecp32* family genes identified from *D. pini*, and from isolates of *D. septosporum* other than NZE10, were tested for the ability of their encoded proteins to trigger cell death in *N. benthamiana*. For the proteins from *D. septosporum* NZE10 that consistently triggered the highest percentages of cell death (DsEcp20-3, DsEcp32-1 and DsEcp32-3), we tested orthologs of other *D. septosporum* alleles and found they triggered the same responses as those of *D. septosporum* NZE10 (Supplementary Figure S9). The same was true for the respective orthologs of DpEcp20-3, DpEcp32-1 and DpEcp32-3 from *D. pini*, which also elicited cell death responses in *N. benthamiana*. These results suggest conservation of cell death-eliciting activity between isolates of *D. septosporum* and between the two *Dothistroma* species for members of the Ecp20 and Ecp32 families.

3.6 Infiltration of Purified Ecp20 and Ecp32 Proteins from *D. septosporum* into *P. radiata* Triggers Cell Death

The pine infiltration method described in Hunziker et al. (2021) proved to be a suitable approach for testing the cell death-eliciting activity of candidate apoplastic effector proteins from pine pathogens in *P. radiata*. We previously showed that DsEcp32-3 triggered cell death in three *P. radiata* genotypes (S6, S11 and R4). Using the same method, we tested two additional proteins from the DsEcp20 family, DsEcp20-3 and DsEcp20-4, and repeated the infiltration of DsEcp32-3 as a positive cell death control (Figure 6, Supplementary Figure S10) (Hunziker et al., 2021).

The DsEcp32-3 positive control triggered cell death in all three pine genotypes, as expected (Figure 6). Infiltration of elution buffer (negative control) did not cause any damage or visible stress responses. Interestingly, both tested proteins gave the same response as they had shown in the non-host *Nicotiana* plants: DsEcp20-3 consistently elicited cell death in the same three pine genotypes used in the previous study, while DsEcp20-4 failed to elicit any response in the pine shoots.

Figure 6. *Dothistroma septosporum* candidate effectors DsEcp20-3 and DsEcp20-4 trigger the same responses in *Pinus radiata* shoot tissues following vacuum infiltration as in non-host plants. Whole shoots of *P. radiata* derived from family seedlots that are relatively susceptible (S6 and S11) or tolerant (R4) to *D. septosporum* infection were infiltrated with purified DsEcp20-3 and DsEcp20-4 proteins produced by heterologous expression in *Pichia pastoris*. Elution buffer was used as negative control and purified DsEcp32-3 protein as a positive control. Representative photos (from 18 to 24 pine shoots for each treatment) were taken 7 days after infiltration.

3.7 DsEcp20-3-triggered Cell Death in *N. benthamiana* requires Membrane-localized Receptor-like Proteins

To determine how cell death elicitors from the Ds/FfEcp20 and 32 family were perceived by plants, we silenced genes encoding the extracellular receptor-like kinase co-receptor genes *BAK1* and *SOBIR1*, as well as leucine-rich repeat (LRR) receptor-like genes, in *N. benthamiana* (Figure 7), in conjunction with agro-infiltration, as previously described by Wang et al. (2018). As expected, agro-infiltration of positive controls NPP1, INF1 and XEG triggered consistent strong cell death responses in *N. benthamiana* leaves treated with TRV:*GFP*, while INF1- and XEG1-triggered cell death was compromised in leaves treated with TRV:*BAK1* or TRV:*SOBIR1*. We also consistently found that DsEcp20-3-triggered cell death was compromised in *N. benthamiana* leaves in which the receptor-like kinase genes *BAK1* or *SOBIR1* were silenced (Figure 7B). In addition, we found that DsEcp20-3-triggered cell death was compromised in *N. benthamiana* leaves treated with TRV VIGS constructs TRV:*T26* and TRV:*T26-2* (Wang et al., 2018) (Figure 7D). TRV:*T26* and TRV:*T26-2* both target the same two *N. benthamiana* genes but in different regions of those genes. One of these target genes encodes an LRR-type extracellular receptor-like protein (Sol Genomics Network accession number - Niben101Scf03138g01049.1) and one an extracellular LRR (Niben101Scf01834g02009.1). We further screened other members of the Ecp20 and Ecp32 families (FfEcp20-3, DsEcp32-3 and FfEcp32-3), and found that cell death triggered by FfEcp20-3 and FfEcp32-3 was also significantly compromised in *N. benthamiana* leaves treated with TRV:*BAK1*, TRV:*SOBIR1* or either of the T26 constructs (Figure 7B and D). For DsEcp32-3-triggered cell death, we did not detect any reduction on TRV:*BAK1*- or TRV:*SOBIR1*-silenced plants (data not shown). Taken together, these results suggest that the cell death responses triggered by members of the Ecp20 and Ecp32 families from *D. septosporum* and *F. fulva* in *N. benthamiana* are the result of recognition by extracellular immune receptors, and that the transduction of defense response signals following this recognition requires the extracellular co-receptors BAK1 and SOBIR1.

Figure 7. Cell death triggered by DsEcp20-3 in *Nicotiana benthamiana* is compromised in leaves silenced for different receptor-like genes. Expression levels of (A) *BAK1* and *SOBIR1* and (C) *T26* constructs in plants silenced for these genes, relative to their expression in plants silenced for the

Green Fluorescent Protein (*GFP*) gene (absent in *N. benthamiana*; normalized to 1). Genes were
silenced using a Virus-Induced Gene Silencing (VIGS) method based on Tobacco Rattle Virus
(TRV). (C, D) Representative leaves showing cell death induced by expression of XEG1, INF1,
NPP1, DsEcp20-3, FfEcp20-3 and FfEcp32-3 in *N. benthamiana* treated with specific TRV-mediated
VIGS constructs as shown. Leaves (n = 8) were photographed three days after agro-infiltration (dai).
Experiments were repeated twice with similar results.

4   Discussion

Plant-pathogenic fungi secrete a collection of virulence factors, termed effector proteins, into their
hosts to promote the infection process (Lo Presti et al., 2015; Rocafort et al., 2020). While many of
these effectors appear to be singletons with no obvious sequence similarity to proteins present in the
effector repertoires to which they belong, others form part of families. Indeed, in *Venturia inaequalis*,
the biotrophic Dothideomycete pathogen responsible for scab disease of apple, more than half of the
predicted effector repertoire is made up of protein families based on sequence similarity alone (Deng
et al., 2017; Rocafort et al., 2022). Likewise, in *Ciborinia camelliae*, the necrotrophic Leotiomycete
pathogen responsible for flower blight disease of camellias, approximately one tenth of the predicted
secretome belongs to a single massively expanded candidate effector family (Denton-Giles et al.,
2020). Notably, effector families can also be associated with avirulence. For example, in the
hemibiotrophic Sordariomycete pathogen *Magnaporthe oryzae* (rice blast disease), the biotrophic
Leotiomycete pathogen *Blumeria graminis* (cereal powdery mildew disease), as well as the
biotrophic Pucciniomycete pathogen *Melampsora lini* (flax rust disease) and the hemibiotrophic
Dothideomycete pathogen *Leptosphaeria maculans* (stem canker disease of brassica), certain
avirulence effectors have been found to form part of protein families (Anderson et al., 2016;
Catanzariti et al., 2006; Dodds et al., 2004; Kang et al., 1995; Khang et al., 2008; Praz et al., 2017;
Spanu, 2017; Talbi et al., 2022).

The emergence of sequence-related effector families in fungi is often the result of duplication events
from an ancestral gene or from other family members and is typically followed by sequence
diversification through an accumulation of indels and synonymous/non-synonymous substitutions
and/or recombination events between different family members (Denton-Giles et al., 2020;
Franceschetti et al., 2017; Pendleton et al., 2014). Such sequence diversification is anticipated to have
roles in, for example, the diversification of effector functions, as well as the evasion of host
immunity. In some cases, this diversification can result in an almost complete loss of sequence
similarity between family members, with similarity only retained at the protein tertiary structure
level, as has recently been shown for several effector families of plant-pathogenic fungi using the
groundbreaking *ab initio* protein structure prediction tool, AlphaFold2 (Rocafort et al., 2022; Seong
& Krasileva, 2022). However, in many cases, sequence identity between family members is
maintained, even across different fungal species. These cross-species effector families are of high
interest, as their sequence conservation may indicate an important and/or conserved (i.e., core) role in
fungal virulence (Stergiopoulos et al., 2010).

In this study, we identified two candidate effector protein families, Ecp20 and Ecp32, which are
highly conserved at the primary sequence and tertiary structure levels between the two closely related
Dothideomycete fungal pathogens, *D. septosporum* and *F. fulva*. Despite their close phylogenetic
relationship, these two pathogens have very different hosts, with *D. septosporum* causing disease on
pine (a gymnosperm), and *F. fulva* causing disease on tomato (an angiosperm) (Drenkhan et al.,
2016; Thomma et al., 2005). An investigation of pre-existing gene expression data revealed that both
protein families have members encoded by genes that are highly expressed *in planta*, while others are

only moderately or lowly expressed. Such a finding is consistent with what was recently observed in
V. inaequalis, where gene family members are expressed at different levels during apple infection
(Rocafort et al., 2022).

Both the Ecp20 and Ecp32 families were found to have members that differed in their ability to
trigger cell death in the non-host species *N. benthamiana* and *N. tabacum*. The differential ability of
these family members to trigger a cell death response does not appear to be restricted to proteins from
D. septosporum and *F. fulva*. Indeed, it was recently determined that an ortholog of Ds/FfEcp20-4
from the hemibiotrophic Dothideomycete pathogen *Zymoseptoria tritici* (septoria leaf blotch disease
of wheat), named Zt4 (JGI protein ID 104697), triggers cell death in *N. benthamiana* (Kettles et al.,
2017). This is in contrast to Ds/FfEcp20-4, where, in our study, no cell death response was observed.
Similarly, a homolog of Ecp32-1 from *Z. tritici* (JGI protein ID 105896; no reciprocal ortholog in *D.
septosporum* or *F. fulva*) was not able to induce a cell death response in *N. benthamiana* (Kettles et
al., 2017) whereas in our study, DsEcp32-1 and FfEcp32-1 consistently and infrequently triggered
cell death, respectively. Of course, it remains possible that some of the family members triggered
other plant responses, such as changes in host gene expression or the accumulation of reactive
oxygen species (ROS), which were not measured in our assays. In any case, variation in the ability of
family members to elicit plant cell death is not restricted to the Ecp20 and Ecp32 families, with
variation also observed in, for example, the cell death-inducing capacity of CcSSP effector candidate
family members from *C. camelliae* in camellia petals (Denton-Giles et al., 2020).

BLASTp searches demonstrated that, like those shown above for *Z. tritici*, homologs of the Ecp20
and Ecp32 family members are present in other fungal species, with many of these species being
plant pathogens. Interestingly, as observed in *D. septosporum* and *F. fulva*, these homologs were also
frequently found to exist as protein families. Taken together, it is therefore likely that the Ecp20 and
Ecp32 proteins are core effectors, possibly contributing to conserved virulence functions in their
respective plant hosts. In addition to searching for homologs in other plant pathogens, we also
screened the genomes of other *D. septosporum* isolates and of *D. pini*, the other causal species of
DNB, and found that whilst Ecp20 and Ecp32 family members are polymorphic across other *D.
septosporum* isolates, all family members are present in these isolates and also in *D. pini*, with the
exception of Ecp32-5. There was also conservation of cell death-eliciting function between isolates
of *D. septosporum* and between the two *Dothistroma* species, since the respective orthologs showed
the same responses as those from the NZE10 strain, also triggering cell death in *N. benthamiana*.
Whether the identified sequence variation contributes to virulence function of these proteins outside
of cell death elicitation remains to be determined.

As the primary amino acid sequence of the Ecp20 and Ecp32 family members provided no insights
into their potential virulence function, we predicted the tertiary structures of these proteins using the
AlphaFold2 tool and compared them to proteins of characterized structure and/or function. Based on
this analysis, all eight *D. septosporum* and *F. fulva* Ecp20 family members were predicted to possess
structural similarity to PevD1, an effector protein with an Alt a 1 (β-barrel) fold from the
hemibiotrophic Sordariomycete pathogen *Verticillium dahliae* (Chruszcz et al., 2012; Han et al.,
2012). Interestingly, the Alt a 1 fold has also been found in MoHrip1, a plasma membrane-binding
virulence factor from *M. oryzae* that is highly expressed during rice infection (Zhang et al., 2017).
Just like FfEcp20-2, FfEcp20-3 and DsEcp20-3, the PevD1 and MoHrip1 proteins induce cell death
responses in *Nicotiana* species (Han et al., 2012; Zhang et al., 2017), suggesting that this cell death-
eliciting ability could be a conserved function of these proteins. It should be noted, however, that
proteins with an Alt a 1 fold, including PevD1, have also been shown to interact with and inhibit the
enzymatic activity of pathogenesis-related 5 (PR5) proteins from different host plants (Gómez-

Casado et al., 2014; Kumar & Mukherjee, 2020; Zhang et al., 2019). Whether the Ecp20 proteins
from *D. septosporum* and *F. fulva* perform a similar role in pathogen virulence remains to be
explored.

In contrast to members of the Ecp20 family, the Ecp32 proteins were predicted to be structurally
similar to proteins from diverse organisms with a β-trefoil fold. Proteins with a β-trefoil fold have the
ability to recognize different ligands, such as DNA, proteins and carbohydrates, and so can be
involved in different processes and interactions between organisms (Žurga et al., 2015). Among the
β-trefoil fold proteins, the most abundant members are lectins, carbohydrate-binding proteins and
hydrolase inhibitors, such as protease inhibitors (Renko et al., 2012). Lectins are proteins that
specifically bind carbohydrates and, in fungi, are associated with plant-pathogen interactions, fungal
growth, fruiting body development and, due to their toxicity, defense against parasitic or predatory
organisms (Sharon, 2007; Varrot et al., 2013). Lectins from different mushroom species, for example,
are known to be toxic to organisms such as *Caenorhabditis elegans* and *Drosophila melanogaster*
(Juillot et al., 2016; Schubert et al., 2012; Wohlschlager et al., 2011). These proteins bind to the
plasma membrane and, in doing so, induce cell death. Cyt toxins from *Bacillus thuringiensis*, for
instance, are insecticidal proteins with a β-trefoil fold that are known to interact with host membrane
lipids (Soberón et al., 2013). Even though the mechanism of lectin toxicity is not known, it is
hypothesized that this toxicity may be the direct result of host membrane damage. Taken together,
this could indicate a possible role Ecp32 proteins in triggering plant cell death, at least in part, by
interacting with and disrupting the host membrane.

As mentioned above, protease inhibitors can also adopt a β-trefoil fold, and in our study, it was found
that most Ds/FfEcp32 proteins had a predicted tertiary structure similar to that of a plant trypsin
inhibitor. Of interest, fungal pathogens frequently target host proteases involved in plant defense by
secreting effectors that serve as protease inhibitors, with well characterized examples including Pit2
from the biotrophic Ustilaginomycete *Ustilago maydis* (corn smut disease) and Avr2 from *F. fulva*
(Mueller et al., 2013; Tian et al., 2007; van Esse et al., 2008). While it is not yet known whether Pit2
or Avr2 adopt a β-trefoil fold, another possible function for the Ecp32 proteins could therefore
involve the inhibition of plant proteases during host colonization.

The fact that Ecp20 and Ecp32 family members of *D. septosporum*, *F. fulva* and *Z. tritici* are likely
apoplastic (Hunziker et al., 2021; Kettles et al., 2017; Mesarich et al., 2018), and that members of
both families across multiple fungal species trigger cell death in *Nicotiana* species (with this activity
dependent on their secretion to the apoplast), may suggest that each family presents a conserved
epitope that is recognized by an extracellular immune receptor in these non-host plants. In support of
recognition by an extracellular immune receptor, silencing of the extracellular co-receptors *NbBAK1*
or *NbSOBIR1* in *N. benthamiana*, which are required to transduce defense response signals following
apoplastic effector recognition (Liebrand et al., 2013; Liebrand et al., 2014), compromised
Ds/FfEcp20-3-dependent cell death. Notably, a compromised cell death response was also observed
for FfEcp32-3, but not DsEcp32-3. One possibility here is that the expression of *NbBAK1* and/or
NbSOBIR1 was not reduced to a level sufficient for the cell death response triggered by DsEcp32-3
to be compromised. Alternatively, given its predicted structural similarity to lectins, other processes
outside of immune receptor recognition, such as cell membrane perturbation, could be involved in
cell death elicitation by DsEcp32-3.

In line with the cell death response observed in the non-host species, we previously determined that
DsEcp32-3 can trigger cell death in pine and suggested that this protein might be a virulence factor in
the necrotrophic stage of DNB that helps to destroy needle tissue (Hunziker et al., 2021). With this in

mind, we decided to investigate whether two (purified) *D. septosporum* proteins from the Ecp20 family could also induce a cell death response in pine. Both proteins, DsEcp20-3 and DsEcp20-4, were of interest due to their high expression in the Mid and Late (necrotrophic) stages of *D. septosporum* infection in *P. radiata* (Bradshaw et al., 2016). The finding that DsEcp20-3 and DsEcp20-4 triggered different cell death responses in non-host plants, *N. benthamiana* and *N. tabacum*, also made them good candidates for this assay. Interestingly, DsEcp20-3, like DsEcp32-3, was able to trigger a cell death response in all pine genotypes tested. Taken together, these results suggest that the DsEcp20-3 and DsEcp32-3 proteins could also by recognized by an immune receptor (possibly conserved with *N. benthamiana*) or, as described above, participate in membrane perturbation in *P. radiata*. Consistent with what was observed in the non-host plants, DsEcp20-4 was unable to elicit a cell death response in the pine shoots. This may suggest that it does not have the conserved epitope required to elicit cell death following recognition by an immune receptor. To identify the regions or amino acid residues involved in cell death induction, and thus potential epitopes for recognition by cognate immune receptors, region swaps between cell death-eliciting and non-eliciting members of the Ecp20 and Ecp32 families, based on information gained from the prediction of surface-exposed residues by AlphaFold2, can now be performed in *Nicotiana* species or pine, similar to that described for the Avr4 effector protein of *D. septosporum* and *F. fulva* in *N. benthamiana* (Mesarich et al., 2016). This, in turn, might help to explain the differences in cell death-inducing capacity between, for example, CfEcp32-1 and DsEcp32-1, as well as DsEcp20-3 and DsEcp20-4.

As it is well known that plant cell death is a strategy exploited by necrotrophic and hemibiotrophic pathogens to acquire nutrients from their hosts during their necrotrophic stage or to facilitate a biotrophic–necrotrophic switch (Shao et al., 2021), one possibility is that the DsEcp20-3 and DsEcp32-3 proteins from *D. septosporum* are involved in these processes. In addition to their cell death-eliciting activities, support for this hypothesis is provided by the fact that the expression of both *DsEcp20-3* and *DsEcp32-3* is highly induced during the Mid and Late stages of pine infection by *D. septosporum*. Another possibility is that DsEcp20-3 may be acting as an Avr factor of *D. septosporum* recognized by the host, similar to that already proposed for the core effector DsEcp2-1 (Guo et al., 2020), while other family members, such as DsEcp20-4, which is also highly expressed during pine infection, might be virulence factors that interact with pine PR5 proteins to suppress host defenses. One way to approach these research questions is to generate mutants of *D. septosporum* deleted for a *DsEcp20* or *DsEcp32* gene (e.g., using CRISPR/Cas9 technology) to determine what impact their deletion or disruption has on necrotrophic infection by this fungus. The problem with this method is that because these genes form part of multigene families, functional redundancy may mask the role of the individual effectors (i.e. the deletion mutants may not show a reduction in virulence). However, it may be possible to create strains in which genes corresponding to several or all family members are deleted to assess whether fungal virulence, particularly in the necrotrophic infection stage, is compromised. Recently, we reported the first successful application of CRISPR/Cas9 genome editing to *D. septosporum* by disrupting a gene encoding a secreted protein that elicited cell death in *N. benthamiana* and pine (McCarthy et al., 2022), which will ultimately help with creating such strains.

In conclusion, our study describes two new candidate effector families with cell death-eliciting activity from *D. septosporum* and *F. fulva* and provides evidence to suggest that members of these families are recognized as invasion patterns by immune receptors of phylogenetically distinct plant species (angiosperms versus a gymnosperm). By characterizing the virulence (and potential avirulence) functions of such 'core' effectors, the knowledge gained could ultimately be used in durable disease resistance programs active against these two, and other, important plant pathogens.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6   Author Contributions

R.B., R.M., Y.W. and C.M. conceived and guided the study. M.T. and Z.C. designed and performed experiments and analyzed data. M.T., C.M. and R.B. led manuscript writing. All authors contributed to the manuscript.

7   Funding

This research was funded by Scion (New Zealand Forest Research Institute, Ltd., Rotorua, New Zealand) and through the Resilient Forests Research Program via Strategic Science Investment Funding from the New Zealand Ministry of Business Innovation and Employment (MBIE), grant number CO4X1703, and New Zealand Forest Grower Levy Trust funding, grant number QT-9108.

8   Acknowledgments

We thank Melissa Guo (Massey University, Palmerston North, New Zealand) and Lukas Hunziker (Curtin University, Perth, Australia) for input on experiments. Trevor S. Loo (Massey University, Palmerston North, New Zealand) is acknowledged for his advice on performing the IMAC, and Keiko Gough (Scion, Rotorua, New Zealand) for preparing the pine material.

9   References

Anderson, C., Khan, M. A., Catanzariti, A.-M., Jack, C. A., Nemri, A., Lawrence, G. J., Upadhyaya, N. M., Hardham, A. R., Ellis, J. G., Dodds, P. N. & Jones, D. A. (2016). Genome analysis and avirulence gene cloning using a high-density RADseq linkage map of the flax rust fungus, *Melampsora lini*. *BMC Genom., 17*(1), 1-20. doi: 10.1186/s12864-016-3011-9

Barnes, I., Crous, P. W., Wingfield, B. D. & Wingfield, M. J. (2004). Multigene phylogenies reveal that red band needle blight of *Pinus* is caused by two distinct species of *Dothistroma*, *D. septosporum* and *D. pini*. *Stud. Mycol., 50*(2), 551-565.

Bolton, M. D., Van Esse, H. P., Vossen, J. H., De Jonge, R., Stergiopoulos, I., Stulemeijer, I. J. E., Van Den Berg, G. C. M., Borrás-Hidalgo, O., Dekker, H. L., De Koster, C. G., De Wit, P. J. G. M., Joosten, M. H. A. J. & Thomma, B. P. H. J. (2008). The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Mol. Microbiol., 69*(1), 119-136. doi: 10.1111/j.1365-2958.2008.06270.x

Bradshaw, R. E., Guo, Y., Sim, A. D., Kabir, M. S., Chettri, P., Ozturk, I. K., Hunziker, L., Ganley, R. J. & Cox, M. P. (2016). Genome-wide gene expression dynamics of the fungal pathogen *Dothistroma septosporum* throughout its infection cycle of the gymnosperm host *Pinus radiata*. *Mol. Plant Pathol., 17*(2), 210-224. doi: 10.1111/mpp.12273

Bradshaw, R. E., Sim, A. D., Chettri, P., Dupont, P.-Y., Guo, Y., Hunziker, L., McDougal, R. L., Van der Nest, A., Fourie, A., Wheeler, D., Cox, M. P. & Barnes, I. (2019). Global population genomics of the forest pathogen *Dothistroma septosporum* reveal chromosome duplications in high dothistromin-producing strains. *Mol. Plant Pathol., 20*(6), 784-799. doi: 10.1111/mpp.12791

Catanzariti, A.-M., Dodds, P. N., Lawrence, G. J., Ayliffe, M. A. & Ellis, J. G. (2006). Haustorially
expressed secreted proteins from flax rust are highly enriched for avirulence elicitors. *Plant*
Cell, 18(1), 243-256. doi: 10.1105/tpc.105.035980

Chruszcz, M., Chapman, M. D., Osinski, T., Solberg, R., Demas, M., Porebski, P. J., Majorek, K. A.,
Pomés, A. & Minor, W. (2012). *Alternaria alternata* allergen Alt a 1: a unique β-barrel
protein dimer found exclusively in fungi. *J. Allergy Clin. Immunol., 130*(1), 241-247. doi:
10.1016/j.jaci.2012.03.047

Coll, N. S., Epple, P. & Dangl, J. L. (2011). Programmed cell death in the plant immune system. *Cell*
Death Differ., 18(8), 1247-1256. doi: 10.1038/cdd.2011.37

Cook, D. E., Mesarich, C. H. & Thomma, B. P. H. J. (2015). Understanding plant immunity as a
surveillance system to detect invasion. *Annu. Rev. Phytopathol., 53*(1), 541-563. doi:
10.1146/annurev-phyto-080614-120114

de Jonge, R., Peter van Esse, H., Kombrink, A., Shinya, T., Desaki, Y., Bours, R., van der Krol, S.,
Shibuya, N., Joosten Matthieu, H. A. J. & Thomma Bart, P. H. J. (2010). Conserved fungal
LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science, 329*(5994), 953-
955. doi: 10.1126/science.1190859

de Wit, P. J. (2016). *Cladosporium fulvum* effectors: weapons in the arms race with tomato. *Annu.*
Rev. Phytopathol., 54(1), 1-23. doi: 10.1146/annurev-phyto-011516-040249

de Wit, P. J., van der Burgt, A., Okmen, B., Stergiopoulos, I., Abd-Elsalam, K. A., Aerts, A. L.,
Bahkali, A. H., Beenen, H. G., Chettri, P., Cox, M. P., Datema, E., de Vries, R. P., Dhillon,
B., Ganley, A. R., Griffiths, S. A., Guo, Y., Hamelin, R. C., Henrissat, B., Kabir, M. S.,
Jashni, M. K., Kema, G., Klaubauf, S., Lapidus, A., Levasseur, A., Lindquist, E., Mehrabi, R.,
Ohm, R. A., Owen, T. J., Salamov, A., Schwelm, A., Schijlen, E., Sun, H., van den Burg, H.
A., van Ham, R. C., Zhang, S., Goodwin, S. B., Grigoriev, I. V., Collemare, J. & Bradshaw,
R. E. (2012). The genomes of the fungal plant pathogens *Cladosporium fulvum* and
Dothistroma septosporum reveal adaptation to different hosts and lifestyles but also
signatures of common ancestry. *PLoS Genet., 8*(11), e1003088. doi:
10.1371/journal.pgen.1003088

DeLano, W. L. (2002). Pymol: an open-source molecular graphics tool. *CCP4 Newsl. Protein*
Crystallogr., 40(1), 82-92.

Deng, C. H., Plummer, K. M., Jones, D. A. B., Mesarich, C. H., Shiller, J., Taranto, A. P., Robinson,
A. J., Kastner, P., Hall, N. E., Templeton, M. D. & Bowen, J. K. (2017). Comparative
analysis of the predicted secretomes of Rosaceae scab pathogens *Venturia inaequalis* and *V.*
pirina reveals expanded effector families and putative determinants of host range. *BMC*
Genom., 18(1), 339. doi: 10.1186/s12864-017-3699-1

Denton-Giles, M., McCarthy, H., Sehrish, T., Dijkwel, Y., Mesarich, C. H., Bradshaw, R. E., Cox,
M. P. & Dijkwel, P. P. (2020). Conservation and expansion of a necrosis-inducing small
secreted protein family from host-variable phytopathogens of the Sclerotiniaceae. *Mol. Plant*
Pathol., 21(4), 512-526. doi: 10.1111/mpp.12913

Dodds, P. N., Lawrence, G. J., Catanzariti, A.-M., Ayliffe, M. A. & Ellis, J. G. (2004). The
Melampsora lini AvrL567 avirulence genes are expressed in haustoria and their products are
recognized inside plant cells. *Plant Cell, 16*(3), 755-768. doi: 10.1105/tpc.020040

Dong, Y., Burch-Smith, T. M., Liu, Y., Mamillapalli, P. & Dinesh-Kumar, S. P. (2007). A ligation-
independent cloning tobacco rattle virus vector for high-throughput virus-induced gene

silencing identifies roles for NbMADS4-1 and -2 in Floral Development. *Plant Physiol., 145*(4), 1161-1170. doi: 10.1104/pp.107.107391

Doyle, J. J. & Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull., 19*, 11-15.

Drenkhan, R., Tomešová-Haataja, V., Fraser, S., Bradshaw, R. E., Vahalík, P., Mullett, M. S., Martín-García, J., Bulman, L. S., Wingfield, M. J., Kirisits, T., Cech, T. L., Schmitz, S., Baden, R., Tubby, K., Brown, A., Georgieva, M., Woods, A., Ahumada, R., Jankovský, L., Thomsen, I. M., Adamson, K., Marçais, B., Vuorinen, M., Tsopelas, P., Koltay, A., Halasz, A., La Porta, N., Anselmi, N., Kiesnere, R., Markovskaja, S., Kačergius, A., Papazova-Anakieva, I., Risteski, M., Sotirovski, K., Lazarević, J., Solheim, H., Boroń, P., Bragança, H., Chira, D., Musolin, D. L., Selikhovkin, A. V., Bulgakov, T. S., Keča, N., Karadžić, D., Galovic, V., Pap, P., Markovic, M., Poljakovic Pajnik, L., Vasic, V., Ondrušková, E., Piškur, B., Sadiković, D., Diez, J. J., Solla, A., Millberg, H., Stenlid, J., Angst, A., Queloz, V., Lehtijärvi, A., Doğmuş-Lehtijärvi, H. T., Oskay, F., Davydenko, K., Meshkova, V., Craig, D., Woodward, S. & Barnes, I. (2016). Global geographic distribution and host range of *Dothistroma* species: a comprehensive review. *For. Pathol., 46*(5), 408-442. doi: 10.1111/efp.12290

Engler, C., Gruetzner, R., Kandzia, R. & Marillonnet, S. (2009). Golden Gate Shuffling: A one-pot DNA shuffling method based on type IIs restriction enzymes. *PLoS One, 4*(5), e5553. doi: 10.1371/journal.pone.0005553

Fellbrich, G., Romanski, A., Varet, A., Blume, B., Brunner, F., Engelhardt, S., Felix, G., Kemmerling, B., Krzymowska, M. & Nürnberger, T. (2002). NPP1, a *Phytophthora*-associated trigger of plant defense in parsley and *Arabidopsis*. *Plant J., 32*(3), 375-390. doi: 10.1046/j.1365-313X.2002.01454.x

Franceschetti, M., Maqbool, A., Jiménez-Dalmaroni, M. J., Pennington, H. G., Kamoun, S. & Banfield, M. J. (2017). Effectors of filamentous plant pathogens: commonalities amid diversity. *Microbiol. Mol. Biol. Rev., 81*(2), e00066. doi: 10.1128/MMBR.00066-16

Gómez-Casado, C., Murua-García, A., Garrido-Arandia, M., González-Melendi, P., Sánchez-Monge, R., Barber, D., Pacios, L. F. & Díaz-Perales, A. (2014). Alt a 1 from *Alternaria* interacts with PR5 thaumatin-like proteins. *FEBS Lett., 588*(9), 1501-1508. doi: 10.1016/j.febslet.2014.02.044

Guo, Y., Hunziker, L., Mesarich, C. H., Chettri, P., Dupont, P.-Y., Ganley, R. J., McDougal, R. L., Barnes, I. & Bradshaw, R. E. (2020). DsEcp2-1 is a polymorphic effector that restricts growth of *Dothistroma septosporum* in pine. *Fungal Genet. and Biol., 135*, 103300. doi: 10.1016/j.fgb.2019.103300

Han, L., Liu, Z., Liu, X. & Qiu, D. (2012). Purification, crystallization and preliminary X-ray diffraction analysis of the effector protein PevD1 from *Verticillium dahliae*. *Acta Crystallogr. F: Struct. Biol. Commun., 68*(7), 802-805. doi: 10.1107/S1744309112020556

Hargreaves, C., Grace, L., Maas, S. v. d., Reeves, C., Holden, G., Menzies, M., Kumar, S. & Foggo, M. (2004). Cryopreservation of *Pinus radiata* zygotic embryo cotyledons: effect of storage duration on adventitious shoot formation and plant growth after 2 years in the field. *Can. J. For. Res., 34*(3), 600-608. doi: 10.1139/x03-226

Heese, A., Hann Dagmar, R., Gimenez-Ibanez, S., Jones Alexandra, M. E., He, K., Li, J., Schroeder Julian, I., Peck Scott, C. & Rathjen John, P. (2007). The receptor-like kinase SERK3/BAK1 is

a central regulator of innate immunity in plants. *PNAS, 104*(29), 12217-12222. doi:
10.1073/pnas.0705306104

Holm, L. (2020). "Using Dali for Protein Structure Comparison," in *Structural Bioinformatics:*
Methods and Protocols, ed. Z. Gáspári (New York, NY: Springer US), 29-42. doi:
10.1007/978-1-0716-0270-6-3

Holsters, M., Silva, B., Van Vliet, F., Genetello, C., De Block, M., Dhaese, P., Depicker, A., Inzé, D.,
Engler, G., Villarroel, R., Van Montagu, M. & Schell, J. (1980). The functional organization
of the nopaline *A. tumefaciens* plasmid pTiC58. *Plasmid, 3*(2), 212-230. doi: 10.1016/0147-
619X(80)90110-9

Hunziker, L., Tarallo, M., Gough, K., Guo, M., Hargreaves, C., Loo, T. S., McDougal, R. L.,
Mesarich, C. H. & Bradshaw, R. E. (2021). Apoplastic effector candidates of a foliar forest
pathogen trigger cell death in host and non-host plants. *Sci. Rep., 11*(1), 19958. doi:
10.1038/s41598-021-99415-5

Jones, J. D. G. & Dangl, J. L. (2006). The plant immune system. *Nature, 444*(7117), 323-329. doi:
10.1038/nature05286

Juillot, S., Cott, C., Madl, J., Claudinon, J., van der Velden, N. S. J., Künzler, M., Thuenauer, R. &
Römer, W. (2016). Uptake of *Marasmius oreades* agglutinin disrupts integrin-dependent cell
adhesion. *Biochim. Biophys. Acta, 1860*(2), 392-401. doi: 10.1016/j.bbagen.2015.11.002

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J.,
Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S.,
Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T.,
Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P. &
Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature,*
596(7873), 583-589. doi: 10.1038/s41586-021-03819-2

Kabir, M. S., Ganley, R. J. & Bradshaw, R. E. (2015). The hemibiotrophic lifestyle of the fungal pine
pathogen *Dothistroma septosporum*. *For. Pathol., 45*(3), 190-202. doi: 10.1111/efp.12153

Kamoun, S., van West, P., de Jong, A. J., de Groot, K. E., Vleeshouwers, V. G. A. A. & Govers, F.
(1997). A gene encoding a protein elicitor of *Phytophthora infestans* is down-regulated during
infection of potato. *Mol. Plant Microbe Interact., 10*(1), 13-20. doi:
10.1094/mpmi.1997.10.1.13

Kang, S., Sweigard, J. A. & Valent, B. (1995). The PWL host specificity gene family in the blast
fungus *Magnaporthe grisea*. *Mol. Plant Microbe Interact., 8*(6), 939-948. doi: 10.1094/mpmi-
8-0939

Kanzaki, H., Saitoh, H., Takahashi, Y., Berberich, T., Ito, A., Kamoun, S. & Terauchi, R. (2008).
NbLRK1, a lectin-like receptor kinase protein of *Nicotiana benthamiana*, interacts with
Phytophthora infestans INF1 elicitin and mediates INF1-induced cell death. *Planta, 228*(6),
977-987. doi: 10.1007/s00425-008-0797-y

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper,
A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Drummond, A. (2012).
Geneious basic: an integrated and extendable desktop software platform for the organization
and analysis of sequence data. *Bioinformatics, 28*(12), 1647-1649. doi:
10.1093/bioinformatics/bts199

Kettles, G. J., Bayon, C., Canning, G., Rudd, J. J. & Kanyuka, K. (2017). Apoplastic recognition of
multiple candidate effectors from the wheat pathogen *Zymoseptoria tritici* in the nonhost
plant Nicotiana benthamiana. *New Phytol., 213*(1), 338-350. doi: 10.1111/nph.14215

Khang, C. H., Park, S.-Y., Lee, Y.-H., Valent, B. & Kang, S. (2008). Genome organization and
evolution of the AVR-Pita avirulence gene family in the *Magnaporthe grisea* species
complex. *Mol. Plant Microbe Interact., 21*(5), 658-670. doi: 10.1094/mpmi-21-5-0658

Kombrink, A. (2012). "Heterologous production of fungal effectors in *Pichia pastoris*," in Plant
Fungal Pathogens: Methods and Protocols, ed. M. D. Bolton & B. P. H. J. Thomma (Totowa,
NJ: Humana Press), 209-217. doi: 10.1007/978-1-61779-501-5-13

Kumar, R. & Mukherjee, P. K. (2020). *Trichoderma virens* Alt a 1 protein may target maize
PR5/thaumatin-like protein to suppress plant defence: an in silico analysis. *Physiol. Mol.*
Plant Pathol., 112, 101551. doi: 10.1016/j.pmpp.2020.101551

Liebrand, T. W. H., van den Berg Grardy, C. M., Zhang, Z., Smit, P., Cordewener Jan, H. G.,
America Antoine, H. P., Sklenar, J., Jones Alexandra, M. E., Tameling Wladimir, I. L.,
Robatzek, S., Thomma Bart, P. H. J. & Joosten Matthieu, H. A. J. (2013). Receptor-like
kinase SOBIR1/EVR interacts with receptor-like proteins in plant immunity against fungal
infection. *PNAS, 110*(24), 10010-10015. doi: 10.1073/pnas.1220015110

Liebrand, T. W. H., van den Burg, H. A. & Joosten, M. H. A. J. (2014). Two for all: receptor-
associated kinases SOBIR1 and BAK1. *Trends Plant Sci., 19*(2), 123-132. doi:
10.1016/j.tplants.2013.10.003

Liu, Y., Schiff, M., Marathe, R. & Dinesh-Kumar, S. P. (2002). Tobacco *Rar1*, *EDS1* and
NPR1/NIM1 like genes are required for N-mediated resistance to tobacco mosaic virus. *Plant*
J., 30(4), 415-429. doi: 10.1046/j.1365-313X.2002.01297.x

Livak, K. J. & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the 2−ΔΔCT method. *Methods, 25*(4), 402-408. doi:
10.1006/meth.2001.1262

Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A., Reissmann,
S. & Kahmann, R. (2015). Fungal effectors and plant susceptibility. *Annu. Rev. Plant. Biol.,*
66, 513-545. doi: 10.1146/annurev-arplant-043014-114623

Ma, Z., Song, T., Zhu, L., Ye, W., Wang, Y., Shao, Y., Dong, S., Zhang, Z., Dou, D., Zheng, X.,
Tyler, B. M. & Wang, Y. (2015). A *Phytophthora sojae* glycoside hydrolase 12 protein is a
major virulence factor during soybean infection and is recognized as a PAMP. *Plant Cell,*
27(7), 2057. doi: 10.1105/tpc.15.00390

MacFarlane, S. A. (1999). Molecular biology of the tobraviruses. *J. Gen. Virol., 80*(11), 2799-2807.
doi: 10.1099/0022-1317-80-11-2799

McCarthy, H. M., Tarallo, M., Mesarich, C. H., McDougal, R. L. & Bradshaw, R. E. (2022).
Targeted gene mutations in the forest pathogen *Dothistroma septosporum* using
CRISPR/Cas9. *Plants, 11*(8) doi: 10.3390/plants11081016

Mesarich, C. H., Stergiopoulos, I., Beenen, H. G., Cordovez, V., Guo, Y., Karimi Jashni, M.,
Bradshaw, R. E. & de Wit, P. J. G. M. (2016). A conserved proline residue in
Dothideomycete Avr4 effector proteins is required to trigger a Cf-4-dependent hypersensitive
response. *Mol. Plant Pathol., 17*(1), 84-95. doi: 10.1111/mpp.12265

Mesarich, C. H., ￼kmen, B., Rovenich, H., Griffiths, S. A., Wang, C., Karimi Jashni, M.,
Mihajlovski, A., Collemare, J., Hunziker, L. & Deng, C. H. (2018). Specific hypersensitive
response–associated recognition of new apoplastic effectors from *Cladosporium fulvum* in
wild tomato. *Mol. Plant Microbe Interact., 31*(1), 145-162. doi: 10.1094/mpmi-05-17-0114-
FI

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. & Steinegger, M. (2022).
ColabFold: making protein folding accessible to all. *Nat. Methods*. doi: 10.1038/s41592-022-
01488-1

Mueller, A. N., Ziemann, S., Treitschke, S., Aßmann, D. & Doehlemann, G. (2013). Compatibility in
the *Ustilago maydis*-maize interaction requires inhibition of host cysteine proteases by the
fungal effector Pit2. *PLoS Pathog., 9*(2), e1003177. doi: 10.1371/journal.ppat.1003177

Nürnberger, T. & Brunner, F. (2002). Innate immunity in plants and animals: emerging parallels
between the recognition of general elicitors and pathogen-associated molecular patterns.
Curr. Opin. Plant Biol., 5(4), 318-324. doi: 10.1016/S1369-5266(02)00265-0

Pendleton, A. L., Smith, K. E., Feau, N., Martin, F. M., Grigoriev, I. V., Hamelin, R., Nelson, C. D.,
Burleigh, J. G. & Davis, J. M. (2014). Duplications and losses in gene families of rust
pathogens highlight putative effectors. *Front. Plant Sci., 5*(299). doi:
10.3389/fpls.2014.00299

Praz, C. R., Bourras, S., Zeng, F., Sánchez-Martín, J., Menardo, F., Xue, M., Yang, L., Roffler, S.,
Böni, R., Herren, G., McNally, K. E., Ben-David, R., Parlange, F., Oberhaensli, S., Flückiger,
S., Schäfer, L. K., Wicker, T., Yu, D. & Keller, B. (2017). AvrPm2 encodes an RNase-like
avirulence effector which is conserved in the two different specialized forms of wheat and rye
powdery mildew fungus. *New Phytol., 213*(3), 1301-1314. doi: 10.1111/nph.14372

Renko, M., Sabotič, J. & Turk, D. (2012). β-Trefoil inhibitors – from the work of Kunitz onward.
Biol. Chem., 393(10), 1043-1054. doi:10.1515/hsz-2012-0159

Rocafort, M., Bowen, J. K., Hassing, B., Cox, M. P., McGreal, B., de la Rosa, S., Plummer, K. M.,
Bradshaw, R. E. & Mesarich, C. H. (2022). The *Venturia inaequalis* effector repertoire is
expressed in waves, and is dominated by expanded families with predicted structural
similarity to avirulence proteins from other fungi. *bioRxiv*, 2022.2003.2022.482717. doi:
10.1101/2022.03.22.482717

Rocafort, M., Fudal, I. & Mesarich, C. H. (2020). Apoplastic effector proteins of plant-associated
fungi and oomycetes. *Curr. Opin. Plant Biol., 56*, 9-19. doi: 10.1016/j.pbi.2020.02.004

Sabotič, J., Renko, M. & Kos, J. (2019). β-Trefoil protease inhibitors unique to higher fungi. *Acta
Chim. Slov., 66*(1). doi: 10.17344/acsi.2018.4465

Sánchez-Vallet, A., Saleem-Batcha, R., Kombrink, A., Hansen, G., Valkenburg, D.-J., Thomma, B.
P. H. J. & Mesters, J. R. (2013). Fungal effector Ecp6 outcompetes host immune receptor for
chitin binding through intrachain LysM dimerization. *eLife, 2*, e00790. doi:
10.7554/eLife.00790

Schubert, M., Bleuler-Martinez, S., Butschi, A., Wälti, M. A., Egloff, P., Stutz, K., Yan, S., Collot,
M., Mallet, J.-M., Wilson, I. B. H., Hengartner, M. O., Aebi, M., Allain, F. H. T. & Künzler,
M. (2012). Plasticity of the β-trefoil protein fold in the recognition and control of invertebrate
predators and parasites by a fungal defence system. *PLoS Pathog., 8*(5), e1002706. doi:
10.1371/journal.ppat.1002706

Seong, K. & Krasileva, K. V. (2022). Comparative computational structural genomics highlights divergent evolution of fungal effectors. *bioRxiv*, 2022.2005.2002.490317. doi: 10.1101/2022.05.02.490317

Shao, D., Smith, D. L., Kabbage, M. & Roth, M. G. (2021). Effectors of plant necrotrophic fungi. *Front. Plant Sci., 12*. doi: 10.3389/fpls.2021.687713

Sharon, N. (2007). Lectins: carbohydrate-specific reagents and biological recognition molecules. *The J. Biol. Chem., 282*(5), 2753-2764. doi: 10.1074/jbc.X600004200

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J. D. & Higgins, D. G. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol., 7*, 539-539. doi: 10.1038/msb.2011.75

Soberón, M., López-Díaz, J. A. & Bravo, A. (2013). Cyt toxins produced by *Bacillus thuringiensis*: a protein fold conserved in several pathogenic microorganisms. *Peptides, 41*, 87-93. doi: 10.1016/j.peptides.2012.05.023

Spanu, P. D. (2017). Cereal immunity against powdery mildews targets RNase-Like proteins associated with haustoria (RALPH) effectors evolved from a common ancestral gene. *New Phytol., 213*(3), 969-971. doi: 10.1111/nph.14386

Sperschneider, J. & Dodds, P. N. (2021). EffectorP 3.0: prediction of apoplastic and cytoplasmic effectors in fungi and oomycetes. *bioRxiv*, 2021.2007.2028.454080. doi: 10.1101/2021.07.28.454080

Stergiopoulos, I., Kourmpetis, Y. A. I., Slot, J. C., Bakker, F. T., De Wit, P. J. G. M. & Rokas, A. (2012). In silico characterization and molecular evolutionary analysis of a novel superfamily of fungal effector proteins. *Mol. Biol. Evol., 29*(11), 3371-3384. doi: 10.1093/molbev/mss143

Stergiopoulos, I., van den Burg, H. A., Ökmen, B., Beenen, H. G., van Liere, S., Kema, G. H. J. & de Wit, P. J. G. M. (2010). Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *PNAS, 107*(16), 7610. doi: 10.1073/pnas.1002910107

Talbi, N., Fokkens, L., Audran, C., Petit-Houdenot, Y., Pouzet, C., Blaise, F., Gay, E., Rouxel, T., Balesdent, M.-H., Rep, M. & Fudal, I. (2022). The neighboring genes *AvrLm10A* and *AvrLm10B* are part of a large multigene family of cooperating effector genes conserved in Dothideomycetes and Sordariomycetes. *bioRxiv*, 2022.2005.2010.491286. doi: 10.1101/2022.05.10.491286

Taylor, R. G., Walker, D. C. & McInnes, R. R. (1993). *E. coli* host strains significantly affect the quality of small scale plasmid DNA preparations used for sequencing. *Nucleic Acids Res., 21*(7), 1677-1678. doi: 10.1093/nar/21.7.1677

Thomma, B. P. H. J., Van Esse, H. P., Crous, P. W. & De Wit, P. J. G. M. (2005). *Cladosporium fulvum* (syn. *Passalora fulva*), a highly specialized plant pathogen as a model for functional studies on plant pathogenic Mycosphaerellaceae. *Mol. Plant Pathol., 6*(4), 379-393. doi: 10.1111/j.1364-3703.2005.00292.x

Tian, M., Win, J., Song, J., van der Hoorn, R., van der Knaap, E. & Kamoun, S. (2007). A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-like apoplastic protease. *Plant Physiol., 143*(1), 364-377. doi: 10.1104/pp.106.090050

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M. & Rozen, S. G.
(2012). Primer3 - new capabilities and interfaces. *Nucleic Acids Res., 40*(15), e115. doi:
10.1093/nar/gks596

van den Burg, H., Harrison, S., Joosten, M., Vervoort, J. & de Wit, P. (2006). *Cladosporium fulvum*
Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during
infection. *Mol. Plant Microbe Interact., 19*(12), 1420-1430. doi: 10.1094/mpmi-19-1420

van Esse, H. P., Van't Klooster, J. W., Bolton, M. D., Yadeta, K. A., van Baarlen, P., Boeren, S.,
Vervoort, J., de Wit, P. J. G. M. & Thomma, B. P. H. J. (2008). The *Cladosporium fulvum*
virulence protein Avr2 inhibits host proteases required for basal defense. *Plant Cell, 20*(7),
1948-1963. doi: 10.1105/tpc.108.059394

Varrot, A., Basheer, S. M. & Imberty, A. (2013). Fungal lectins: structure, function and potential
applications. *Curr. Opin. Struct. Biol, 23*(5), 678-685. doi: 10.1016/j.sbi.2013.07.007

Wang, Y., Wang, Y. & Wang, Y. (2020). Apoplastic proteases: powerful weapons against pathogen
infection in plants. *Plant Commun., 1*(4), 100085. doi: 10.1016/j.xplc.2020.100085

Wang, Y., Xu, Y., Sun, Y., Wang, H., Qi, J., Wan, B., Ye, W., Lin, Y., Shao, Y., Dong, S., Tyler, B.
M. & Wang, Y. (2018). Leucine-rich repeat receptor-like gene screen reveals that *Nicotiana*
RXEG1 regulates glycoside hydrolase 12 MAMP detection. *Nature Commun., 9*(1), 594. doi:
10.1038/s41467-018-03010-8

Weber, E., Engler, C., Gruetzner, R., Werner, S. & Marillonnet, S. (2011). A modular cloning system
for standardized assembly of multigene constructs. *PLoS One, 6*(2), e16765. doi:
10.1371/journal.pone.0016765

Weidner, M., Taupp, M. & Hallam, S. J. (2010). Expression of recombinant proteins in the
methylotrophic yeast Pichia pastoris. *J. Vis. Exp., JoVE*(36), 1862. doi: 10.3791/1862

Win, J., Chaparro-Garcia, A., Belhaj, K., Saunders, D. G., Yoshida, K., Dong, S., Schornack, S.,
Zipfel, C., Robatzek, S., Hogenhout, S. A., & Kamoun, S. (2012). Effector biology of plant-
associated organisms: concepts and perspectives. *Cold Spring Harb. Symp. Quant. Biol.*, *77*,
235–247. doi: 10.1101/sqb.2012.77.015933

Wohlschlager, T., Butschi, A., Zurfluh, K., Vonesch, S. C., Auf dem Keller, U., Gehrig, P., Bleuler-
Martinez, S., Hengartner, M. O., Aebi, M. & Künzler, M. (2011). Nematotoxicity of
Marasmius oreades agglutinin (MOA) depends on glycolipid binding and cysteine protease
activity. *J. Biol. Chem., 286*(35), 30337-30343.

Zaccaron, A. Z., Chen, L.-H., Samaras, A. & Stergiopoulos, I. (2022). A chromosome-scale genome
assembly of the tomato pathogen *Cladosporium fulvum* reveals a compartmentalized genome
architecture and the presence of a dispensable chromosome. *Microb. Genom., 8*(4) doi:
10.1099/mgen.0.000819

Zhang, Y., Gao, Y., Liang, Y., Dong, Y., Yang, X. & Qiu, D. (2019). *Verticillium dahliae* PevD1, an
Alt a 1-like protein, targets cotton PR5-like protein and promotes fungal infection. *J. Exp.*
Bot., 70(2), 613-626. doi: 10.1093/jxb/ery351

Zhang, Y., Liang, Y., Dong, Y., Gao, Y., Yang, X., Yuan, J., Qiu, D. (2017). The *Magnaporthe*
oryzae Alt A 1-like protein MoHrip1 binds to the plant plasma membrane. *Biochem. Biophys.*
Res. Commun., 492(1), 55-60. doi: 10.1016/j.bbrc.2017.08.039

Zhou, D., Lobo, Y. A., Batista, I. F. C., Marques-Porto, R., Gustchina, A., Oliva, M. L. V. &
Wlodawer, A. (2013). Crystal structures of a plant trypsin inhibitor from *Enterolobium*

contortisiliquum (EcTI) and of its complex with bovine trypsin. *PLoS One, 8*(4), e62252. doi:
10.1371/journal.pone.0062252

Žurga, S., Pohleven, J., Kos, J. & Sabotič, J. (2015). β-Trefoil structure enables interactions between
lectins and protease inhibitors that regulate their biological functions. *J. Biochem., 158*(1), 83-
90. doi: 10.1093/jb/mvv025