# GWAS identifies candidate regulators of in planta regeneration in Populus trichocarpa

Michael F. Nagle[1], Jialin Yuan[2], Damanpreet Kaur[2], Cathleen Ma[1], Ekaterina Peremyslova[1], Yuan Jiang[3], Alexa Niño de Rivera[1], Sara Jawdy[4,5], Jin-Gui Chen[4,5,6], Kai Feng[4,5], Timothy B. Yates[4,5,6], Gerald A. Tuskan[4,5], Wellington Muchero[4,5,6], Li Fuxin[2], Steven H. Strauss[1]

[1] Department of Forest Ecosystems and Society, Oregon State University, Corvallis, OR, USA

[2] Department of Electrical Engineering and Computer Science, Oregon State University, Corvallis, OR, USA

[3] Statistics Department, Oregon State University, Corvallis, OR, USA

[4] Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

[5] Center for Bioenergy Innovation, Oak Ridge National Laboratory, Oak Ridge, TN, USA

[6] Bredesen Center for Interdisciplinary Research, University of Tennessee, Knoxville, TN, USA

**Abstract**

Plant regeneration is an important dimension of plant propagation, and a key step in the

production of transgenic plants. However, regeneration capacity varies widely among

genotypes and species, the molecular basis of which is largely unknown. While

association mapping methods such as genome-wide association studies (GWAS) have

long demonstrated abilities to help uncover the genetic basis of trait variation in plants,

the power of these methods relies on the accuracy and scale of phenotypic data used. To

enable a largescale GWAS of *in planta* regeneration in model tree *Populus*, we

implemented a workflow involving semantic segmentation to quantify regenerating plant

tissues (callus and shoot) over time. We found the resulting statistics are of highly non-

normal distributions, which necessitated transformations or permutations to avoid

violating assumptions of linear models used in GWAS. While transformations can lead to

a loss of statistical power, we demonstrate that this can be mitigated by the application of

the Augmented Rank Truncation method, or avoided altogether using the Multi-Threaded

Monte Carlo SNP-set (Sequence) Kernel Association Test to compute empirical *p*-values

in GWAS. We report over 200 statistically supported candidate genes, with top

candidates including regulators of cell adhesion, stress signaling, and hormone signaling

pathways, as well as other diverse functions. We demonstrate that sensitive genetic

discovery for complex developmental traits can be enabled by a workflow based on

computer vision and adaptation of several statistical approaches necessitated by to the

complexity of regeneration trait expression and distribution.

**Introduction**

Plant genetic engineering and gene editing have produced new varieties of crops with a

variety of valuable traits (NAS, 2016; Jaganathan *et al.*, 2018). However, the ability to

impart new traits by these methods is limited to crop species with genotypes that can

reliably undergo transformation and regeneration (TR). TR requires developmental

responses to a series of hormone treatments and amenability to gene insertion, and the

capacity for both varies greatly between and within species (Altpeter *et al.*, 2016). The

causes of this great variation in recalcitrance are poorly known; however, GWAS – with

its potential to identify genes whose variation plays a key role in capacity for TR – can

greatly enhance understanding of the TR process. In addition, the identified genes may

serve as "reagents" for overcoming recalcitrance, similar to how overexpression of

morphogenic regulator (MR) genes can enhance *in vitro* regeneration of transgenic shoot

in a variety of species (Gordon-Kamm *et al.*, 2019). *In planta* transformation methods

can also be enhanced by MR genes, including those in *Populus tomentosa* (Deng *et al.*,

2009), *Nicotiana benthamiana*, tomato, potato and grape (Maher *et al.*, 2020). However,

given the complexity and genotypic variation in TR capacity, it is likely that only a

fraction of the potentially useful MR genes have been identified.

To help identify the genes responsible for variation in TR, we conducted GWAS in a

population of 1,219 wild cottonwoods that had been re-sequenced by the US Department

of Energy, up to 917 of which were previously studied for a variety of traits (Zhang *et al.*,

2018a; Tuskan *et al.*, 2018; Muchero *et al.*, 2018; Bdeir *et al.*, 2019; Weighill *et al.*,

2019; Chhetri *et al.*, 2020). We focused on regeneration from cut stems, while

considering these may be a direct substrate for accelerated *in planta* transformation

systems, and because of the expectation that regeneration processes are likely to share

many elements whether induced *in planta* or *in vitro*. GWAS and related association

mapping methods have previously been applied to study variation in the rates of *in vitro*

regeneration in Arabidopsis, rice, wheat, rose, sorghum and poplar, among other plants

(reviewed by Lardon & Geelen, 2020).

Regeneration phenotypes are notoriously difficult to quantify, whether *in planta* or *in*

*vitro*. Calli and emerging shoots are often highly variable and complex in shape, color,

and size, and sequential measurements are hard to take without damaging or

contaminating regenerating tissues. This appears to have limited sample sizes in prior

GWAS studies of regeneration. For example, Tuskan et al. (2018) selected only 280

genotypes to phenotype callus growth from a re-sequenced GWAS population of 1,084 *P.*

*trichocarpa* genotypes. A similar GWAS of callus differentiation into shoots in *P.*

*euphratica* was limited to 297 genotypes (Zhang *et al.*, 2020). Nguyen et al. (2020) noted

the "extremely laborious" nature of phenotyping *in vitro* traits as a consideration in their

GWAS of callus formation across 96 rose genotypes (Nguyen *et al.*, 2020).

Because of the importance of a large and precise sample for statistical power in GWAS

(López-Cortegano & Caballero, 2019), we developed a computer vision method to

measure regeneration from sequential images of decapitated, regenerating stems. Over 40

published studies have made use of diverse computer vision methods in GWAS of plants,

including Arabidopsis, maize, wheat, rice, sorghum, soybean and barley, among others

(Guo *et al.*, 2022; Affortit *et al.*, 2022; reviewed by Xiao *et al.*, 2021). Developmental

traits related to biomass and growth of whole plants or specific tissues have commonly

been studied using various methods for segmentation of RGB images, including those

involving color thresholds (Yang *et al.*, 2015; Das *et al.*, 2015; Al-Tamimi *et al.*, 2016;

Guo *et al.*, 2018; Gage *et al.*, 2018; Pham *et al.*, 2019; Seethepalli *et al.*, 2020; Campbell

*et al.*, 2020; Ogawa *et al.*, 2021; Affortit *et al.*, 2022), machine learning approaches such

as support vector machines and random forests (Zou *et al.*, 2019; She *et al.*, 2019; Carlier

*et al.*, 2022; Xu *et al.*, 2022), and deep convolutional neural networks (DCNNs; Ma *et al.*,

2019a; Yasrab *et al.*, 2019; Xu *et al.*, 2022; Kamath *et al.*, 2022). In recent years, DCNNs

similar to that employed in the present study have demonstrated abilities to outperform

earlier methods (Adams *et al.*, 2020) and have become the dominant approach used for

diverse computer vision tasks. In the context of plant phenotyping, this was demonstrated

by the unparalleled performance of neural networks for the Leaf Segmentation Challenge

benchmark dataset (Aich & Stavness, 2017; Dobrescu *et al.*, 2017).

We report a GWAS of *in planta* regeneration in poplar, using a computer vision

workflow based on deep segmentation to quantify specific regenerating tissues, and four

association mapping methods to identify genetic markers associated with regeneration

traits. The non-normal and skewed characteristics of segmentation-based traits presented

a challenge for GWAS, which commonly relies on an assumption of normally-distributed

residuals. Due to this, we tested multiple GWAS methods with adaptations to avoid

violating this assumption while also preserving statistical power. We found over 200

statistically-supported gene candidates with diverse physiological roles that include

hormone signaling, plant stress response, control of cell division, and cell wall structure –

as well as many genes of unknown function.

**Materials and Methods**

<u>Experimental design</u>

The primary objective of our work here was to develop, test, and refine workflows using

computer vision to measure developing tissues during regeneration. The system also

needed to be sensitive enough to enable detailed genetic characterization of regeneration

traits. Toward this end, we first adapted a DCNN of the PSPNet architecture to compute

segmentation masks for regenerating plant tissues (callus, shoot and unregenerated stem)

on a high-throughput scale. Second, we evaluated the statistical considerations of using

these traits in GWAS, whether by models often relying on trait transformation to avoid

violating assumptions of normally-distributed residuals, or by models that avoid this

assumption altogether by binarization or permutation and computation of empirical $p$-

values. Finally, we applied these methods in a population with over 1,200 wild poplar

genotypes ideal for GWAS due to a remarkable number and density of single-nucleotide

polymorphisms (SNPs; up 34 million depending on GWAS method) and extremely low

linkage disequilibrium (LD; Zhang *et al.*, 2018a; Tuskan *et al.*, 2018; Muchero *et al.*,

2018; Bdeir *et al.*, 2019; Weighill *et al.*, 2019; Chhetri *et al.*, 2020). An overview of the

experimental population and analysis pipeline is shown in Fig. 1.

**Figure 1. Overview of *in planta* callus and shoot regeneration GWAS workflow.** The first row describes resources that were provided through other work prior to undertaking the GWAS. Images on the right of the "Phenotyping" panel are the RGB image, ground truth semantic label and overlay for one sample.

Plant materials

We utilized an expanded version of the previously reported re-sequenced *P. trichocarpa*

GWAS population (Zhang *et al.*, 2018a; Tuskan *et al.*, 2018; Muchero *et al.*, 2018; Bdeir

*et al.*, 2019; Weighill *et al.*, 2019; Chhetri *et al.*, 2020). The population was recently

expanded to include an additional 441 genotypes, filling a geographical gap that existed

in first release with 882 genotypes (Fig. 2). While this clone bank is kept at multiple

locations, phenotyping in this study only made use of the replicate in Corvallis, OR,

featuring a total of 1,307 clones in the population (out of 1,323) and for 1,219 of which

regeneration phenotyping was performed. Clones were grown at two field locations in

Corvallis, OR: one location planted in 2009 featuring the original GWAS population, and

another planted in 2015 featuring the newly added clones. Dormant cuttings were taken

in the winters of 2018, 2019, and 2020, frozen, and then rooted up to one year later.

Plants were then regularly pruned and fertilized to ensure there were healthy green leaves

suitable for sterilization and introduction into tissue culture. A second greenhouse

population was established and allowed to go dormant in winter; plants from this source

were occasionally used to replace plants in the main greenhouse population that provided

plant materials throughout the year.

**Figure 2. Origins of P. trichocarpa clones used to generate SNP set.** A total of 1,323 native genotypes were collected over a geographical range across the pacific northwest region of the USA and the southwest of Canada. Tree location is shown for 1,301 genotypes for which precise location data is available.

Sequencing and SNP set preparation

We performed SNP calling and analyzed the distribution of SNPs after resequencing of

406 additional genotypes by the DOE Joint Genome Institute, for a total of 1,323

genotypes in the complete SNP set (increased from 917 genotypes in the most recent SNP

set). DNA short-read sequencing data for all 1,323 genotypes was filtered and trimmed

with BBDuk (https://sourceforge.net/projects/bbmap/). Parameters used for BBDuk

included 'kmer=25', 'ktrim=r', and 'trimq=6'. Clean reads were then aligned to the P.

trichocarpa v3.0 reference genome with bwa-mem v0.7.17 (Li & Durbin, 2009).

Duplicate reads were then marked with Sambamba markdup v0.7.0 (Tarasov *et al.*,

2015). Next, BQSR was performed with GATK v4.1.3.0 BaseRecalibrator (Poplin *et al.*,

2018). Variants were then called by sample with GATK v4.1.3.0 HaplotypeCaller and

GVCFs were generated (Poplin *et al.*, 2018). The GVCFs were then consolidated with

GATK v4.1.3.0 GenomicsDBImport and joint calling was performed with GATK

v4.1.3.0 GenotypeGVCFs. Next, SNPs and insertion-deletion variants (indels) were

separated with GATK v4.1.3.0 SelectVariants. Indels were filtered with GATK v4.1.3.0

VariantFiltration with the following thresholds and flags: '-filter "QD < 2.0" --filter-name

"QD2" -filter "FS > 200.0" --filter-name "FS200" -filter "ReadPosRankSum < -20.0" --

filter-name "ReadPosRankSum-20"'. SNPs were filtered with GATK v4.1.3.0

VariantFiltration with the following thresholds and flags, '-filter "QD < 2.0" --filter-name

"QD2" -filter "SOR > 4.0" --filter-name "SOR4"  -filter "FS > 60.0" --filter-name "FS60"

-filter "MQ < 40.0" --filter-name "MQ40" -filter "MQRankSum < -12.5" --filter-name

"MQRankSum-12.5" -filter "ReadPosRankSum < -8.0" --filter-name "ReadPosRankSum-

8"'.  There was a total of 40.4M SNPs prior to filtering for minor allele frequency (MAF)

and additional quality criteria. The density and consistency of SNP data on each

chromosome were assessed using the R package Cmplot (Fig. S1) and by producing

histograms of gap sizes for each chromosome.

Assay of regeneration

Frozen stem cuttings were incubated at 4 °C for 2-4 weeks, then placed in 50mL Falcon

tubes with water for five weeks. Based on preliminary experiments (data not shown), we

found that treatment of the decapitated surface of each stem with 100μL of 0.5mg/mL

thidiazuron (TDZ) in water improved callus regeneration considerably (37% of genotypes

produced shoots, compared to 24% without TDZ). After application of TDZ to a given

stem tip, a 1.5mL microcentrifuge tube was inverted over the stem tip to prevent

desiccation during regeneration (as shown in Fig. 1). On a weekly basis beginning the

second week, decapitated stem surfaces were imaged from overhead using a Canon Rebel

Xsi DSLR camera attached to a rack mount.

Due to practical limitations on the numbers of clones that could be assayed for

regeneration simultaneously, subsets of the study genotypes (termed "phases") were

assayed at one time, with no more than 400 cuttings per phase. Images were taken on a

weekly basis from the second week through the fifth week, with the exceptions of weeks

four and five in the first phase and week four in the third phase. There were two replicate

plants measured for all but the first three phases, where only a single replicate was used.

Image segmentation

To perform annotation of images for computer vision, 249 images were randomly

sampled from the first seven phases and manually annotated using the Intelligent Deep

Annotator for Segmentation (IDEAS) graphical user interface that we previously

developed (Yuan *et al.*, 2022). As described in our prior work, these samples were used

to train a convolutional neural network (PSPNet) via TensorFlow v1.14 to segment

images of regenerating stem tips with each pixel labeled as one of four classes: callus,

shoot, unregenerated stem and background. The trained model was deployed to produce

inferred semantic masks for 11,791 images in the complete GWAS dataset, including

1,219 clones with up to four replicates and a median of two replicates, across four weekly

timepoints (Fig. 3). At each timepoint, two traits were computed: the proportion of total

plant area which consists of callus (henceforth, "callus area"), and of shoot ("shoot

area").

**Figure 3. False-color examples of segmentation from regenerating stems shown in cross-section.** For each section, shown are RGB images (left panels), inferred semantic mask labels (SEG, middle panels), and overlays of images and masks (OV, right panels) are shown for two replicates of genotype BESC-1013 across four timepoints of development, from week 2 (top) to week 5 (bottom). Semantic masks show unregenerated stem tissue (red), callus (blue) and shoot (green).

Computer vision trait preparation for GWAS

For replicated samples, the mean value of each trait across the two replicates was

computed and used in downstream analysis. For genotypes lacking replication due to

infection in one replicate, the single unreplicated trait value was used.

Additional traits were computed by performing principal component analysis (PCA)

using `stats::princomp` in R over three groups of traits: 1) callus area traits at all

timepoints; 2) shoot area traits at all timepoints; and 3) both callus area and shoot area at

all timepoints. Genotypes missing data for a given trait at any timepoint were excluded

from a given PCA. Scree plots were evaluated to estimate the number of PCs

representing significant proportions of trait variation.

The normality of traits was assessed using Q-Q plots, histograms, Shapiro-Wilks tests

and Pearson correlation coefficients computed against theoretical normal distributions

with the same mean and standard deviation as the given trait. To avoid severe violations

of normality that may lead to inflated error rates, all traits were transformed prior to

GWAS methods that depend on normality and linear model assumptions. The most basic

transformation applied was a removal of zero values followed by Box-Cox

transformation. For certain PC traits, a spike was observed at particular values, which

corresponded to genotypes with zero values for all traits used in the given PCA; these

genotypes were consequently removed. In cases where we determined that thresholding

or extreme outlier removal was necessary, these treatments were performed prior to the

Box-Cox transformation. In addition, as an alternative to Box-Cox transformations, rank-

based inverse normal (RB-INV) transformations were performed for difficult

distributions (Fig. S2, Table S1, S2).

Association Mapping

Because of the distinct assumptions and data types for which various GWAS methods are

suited, along with the non-normality of computer-vision-based traits and the risk of

sacrificing statistical power with transformations, we employed an analysis pipeline that

made use of four GWAS methods. These include Genome-wide Efficient Mixed Model

Association (GEMMA; Zhou & Stephens, 2012), which was parallelized using the GNU

Parallel (Tange, 2020) framework to simultaneously run each given trait on a CPU core.

Due to an assumption of normally distributed model predictions for GEMMA, this

method was only used to study versions of traits that had been transformed toward

normality, including the exclusion of all samples with zero values. The Generalized

Mixed Model Association Test (GMMAT; Chen *et al.*, 2016b) was used for single-

marker tests with the same kinship covariate as GEMMA; however, rather than using

continuous trait variables, GMMAT applies logistic regression and works with binarized

traits. Finally, as a means of performing tests that avoid any transformation or

binarization of traits, we applied our Multi-Threaded Monte Carlo Sequence Kernel

Association Test (MTMC-SKAT; github.com/naglemi/mtmcskat), an R extension of the

established SKAT method (Wu *et al.*, 2011; Ionita-Laza *et al.*, 2013) extending support

for efficient resampling on high-performance clusters. Using MTMC-SKAT, we tested

for the combined effect of adjacent SNPs within 3kb windows and computed empirical *p*-

values for SNPs indicated by an initial parametric test as being strongly associated with a

trait. With MTMC-SKAT, we tested and compared the control for population structure

via two types of covariates: a matrix of PCs from PCA of SNP data (P model) and a

matrix of theoretical subpopulation ancestry estimates from fastSTRUCTURE (Q model;

Raj *et al.*, 2014). MTMC-SKAT was deployed on the COMET high-performance cluster

made available through the NSF XSEDE program (Towns *et al.*, 2014). Additional

details on GWAS methods can be found in Supplementary Methods.

Statistically significant associations from the various pipelines were first determined by

computing Benjamini-Hochberg false discovery rate (FDR; $\alpha = 0.10$) and Bonferroni

thresholds ($\alpha = 0.05$). SNPs not found on contiguous assembled chromosomes were

excluded from computation from this point forward (approximately 3.6% of total SNPs).

The Bonferroni thresholds were computed given the number of tests equal to the number

of filtered SNPs (for single-marker tests GEMMA and GMMAT) and the number of 3kb

SNP windows (in the case of SKAT). We then extracted lists of SNPs with *p*-values

below these thresholds for interrogation.

We next evaluated the extent to which multiple SNPs supported the association of a

nearby gene, whether individual SNPs met the FDR or Bonferroni statistical thresholds or

not. We implemented the Augmented Rank Truncation (ART) method (Vsevolozhskaya

*et al.*, 2019) to scan Wald *p*-values from GEMMA and GMMAT and identify cases

where a SNP produces a *p*-value below $1*10^{-5}$ and is within 500bp of at least 5 additional

SNPs with *p*-values below $1*10^{-4}$ when considering the upper half of top-ranking SNPs

(according to *p*-values). While we initially computed Wald *p*-values with GMMAT for

only the top 100 or 1,000 SNPs according to a score test, additional Wald *p*-values were

computed for use in ART. For each of these 1kb windows, a combined *p*-value was

computed for the extracted SNPs. A Bonferroni threshold for ART *p*-values was

computed ($\alpha = 0.05$) from the approximate number of independent tests (contiguous

assembled genome size / ART window size). The Bonferroni threshold of $\sim1.27*10^{-7}$ was

computed using the number of independent tests of ($\sim3.94*10^{6}$ 1kb windows spanning

the ~394 Mb of contiguous assembled chromosomes) and is notably less conservative

than the Bonferroni threshold used for raw *p*-values from GEMMA/GMMAT

(henceforth, "conservative Bonferroni"), as computed from the total number of tests (as

low as $\sim4*10^{-9}$, given up to ~13 million SNPs on contiguous chromosomes remaining

after filtering; Supplementary Methods). To be emphasized, it is well-known that

Bonferroni thresholds for individual SNPs consider each SNP as an independent test,

leading to overcorrection since large numbers of SNPs are in LD.

To determine on a high-throughput scale which genes are likely to be responsible for

statistically significant quantitative trait loci (QTLs; either SNPs or SNP windows), we

used R scripts to reference genome and genome annotation data available through

Phytozome (phytozome.jgi.doe.gov; Tuskan *et al.*, 2006). In this workflow, the positions

of loci were evaluated for candidate genes only when these loci represent the "peak" of a

signal, determined by checking for any other loci within 30kb with a more significant *p*-

value. The candidate gene responsible for the significance of a given locus was assumed

by the workflow to be the gene that encompasses or is closest to the locus. The R package

InterMineR (Kyritsis *et al.*, 2019) was used to collect Phytozome data on gene function,

Arabidopsis homologs, and gene ontology terms and organized these by locus. The

GreeNC database was used to identify possible noncoding regulatory RNAs among gene

candidates (Di Marsico *et al.*, 2022). For the top gene candidates, particularly those

passing the conservative Bonferroni or FDR ($\alpha = 0.10$) thresholds, or those passing the

less conservative Bonferroni thresholds used for ART and among the five most-

significant GEMMA-ART or GMMAT-ART associations for a given trait, Integrative

Genomics Viewer (IGV; Robinson *et al.*, 2011; Thorvaldsdóttir *et al.*, 2013) was used to

manually investigate gene position relative to significant SNPs, including consideration

of other nearby genes, distance to the putative transcription start site, and direction of

transcription.

**Results**

<u>Trait transformations</u>

Prior to transformations, most traits displayed marked non-normal characteristics as

indicated by Q-Q plots, histograms, Shapiro-wilk tests, and Pearson correlation

coefficients of each distribution with a normal distribution with the same mean and

standard deviation. The improvement in normality after transformation was marked in

most cases (Table S1, S2; Fig. S2; data not shown). Non-normal characteristics of most

traits were reduced substantially by excluding genotypes with zero values and applying a

Box-Cox transformation (e.g., Fig. S2). For the traits of callus or shoot area at each

timepoint, based on visual inspection of histograms and Q-Q plots, the improvement in

metrics of normality was deemed adequate for linear models. All PCA-derived traits

necessitated additional treatments to avoid severe violations of the normality assumption

of linear models, including removal of outliers and in some cases removal of values

below an elbow in the frequency distribution (estimated as the position where the second

derivative of the probability frequency distribution is maximum; Table S2).

<u>Principal components as proxies for complex patterns of regeneration</u>

Scree plots and heat maps of loadings revealed common trends in regeneration across

timepoints and regenerating tissue types (callus and shoot). These results were obtained

for three different sets of PCA with different groups of traits: first, for both callus and

shoot area at all timepoints (Fig. 4), and then with callus and shoot data analyzed

independently over all timepoints (Fig. S3). In all three cases, the PC explaining the most

variation (PC1) represented a tendency of the tissue(s) included in PCA to regenerate

across all timepoints. Latter PCs provided proxies for more complex patterns of

regeneration. PC2 over callus traits appears to represent high levels of callus regeneration

at early, but not later timepoints. PC2 over all callus and shoot traits appears to represent

a tendency for callus to regenerate robustly, but to fail to develop into shoots. Subsequent

PCs, for each batch of traits, represented a relatively small proportion of variance

explained and were thus not analyzed for gene candidates.

**(A)**　　　　　　　　　　　　　　　　　　　　　　**(B)**

**Figure 4. PCA of *in planta* callus and shoot regeneration traits.** Results are shown from PCA over all callus and shoot traits across four weekly timepoints. **(A)** Scree plot; **(B)** Heat map of loadings from PCA.

**Figure 5. Linkage disequilibrium decay.** Each point represents the mean LD between SNPs of a given distance (x-axis). The blue line represents LD decay as computed with a spline function.

SNP set for *P. trichocarpa* provides comprehensive view of natural variation

The SNP set produced for this population displays polymorphism across all regions of

contiguous chromosomes represented in the reference genome (Fig. S1). Poplar clones

collected for the GWAS clone bank represent a wide range of geographic diversity,

nearly spanning the natural range of *P. trichocarpa* from British Columbia to the Pacific

Northwest of the United States (Fig. 2). There is clearly very strong natural

intrachromosomal recombination, as LD decay occurs rapidly and reaches $R^2 = 0.2$

within 2kb whether computed for the whole population or either of the two most

prominent subpopulations (Fig. 5, Fig. S4). Population structure analysis with

fastSTRUCTURE, cross-referenced with geography and phylogenetics, supported the

existence of 6 or 7 ancestral subpopulations, which were most distinct in southern

regions, while a greater degree of admixture was prevalent farther north and may be

responsible for the significantly lower LD in the "Oregon" subpopulation ($P < 0.001$; Fig.

S4). Our linear models supported a positive relationship between latitude and callus traits,

but were unable to detect such a relationship for shoot traits (Fig. S5-S10; Table S3;

Supplemental Results and Discussion).

<u>Genes implicated by significant quantitative trait loci (QTLs)</u>

We interrogated traits with $h^2_{SNP}$ above 0.10 for candidate genes (Table S4), except for

shoot area at week 2 (first timepoint) due to a sparse distribution and high level of noise

for this trait. We utilized three multiple-testing correction methods (from most to least

conservative: conservative Bonferroni, Benjamini-Hochberg FDR and ART-Bonferroni)

to provide three levels of confidence in associations to support prioritization of gene

candidates. Across the three GWAS methods GEMMA, GMMAT, and (MTMC-)SKAT,

we report a total of 13 unique QTL peaks with $p$-values passing the conservative

Bonferroni significance threshold, as well as 44 passing the FDR ($\alpha = 0.10$) threshold.

Among Bonferroni-significant associations, 6 are inside or within 5kb of a gene found in

the genome annotation, as well as 32 associations (~72.7%) meeting the FDR threshold

(Fig. 6-8, Table S5, S6). We found 137 unique QTL peaks from applying our

implementation of ART to GEMMA results (Table S5, S7), as well as 24 from applying

ART to GMMAT results (Table S5, S8). Several of the most promising candidates, based

on the biology of their homologs in Arabidopsis, are shown in Table 1.

**(A)**                                                     **(B)**

**Figure 6. Tallies of *in planta* callus and shoot associations across GWAS methods.** Shown are barplots summarizing the numbers of associations from each GWAS method, with two types of significance thresholds, as well as within a 5kb distance threshold of the nearest gene. QTL peaks were taken as the point with the lowest *p*-value at any given peak, where multiple points within the same peak may otherwise pass a given significance threshold. **(A)** QTL peaks passing the Benjamini-Hochberg threshold (FDR; α = 0.10) and/or conservative Bonferroni threshold; **(B)** QTL peaks passing ART-Bonferroni threshold (α = 0.05, *N* of 1kb windows in genome).

**Figure 7. Manhattan plot of GEMMA-ART results for week four callus area.** Shown is a Manhattan plot for GEMMA results for the trait of callus area at week four. Black and orange lines show Bonferroni significance thresholds for GEMMA results with independent SNPs, and for ART applied to GEMMA over 1kb windows of SNPs, respectively. Black circles represent tests of individual SNPs by GEMMA, while orange triangles represent 1kb windows tested by ART applied to GEMMA results.

**(A)**

**(B)**

**Figure 8. Close-up view of Manhattan plots for example *in planta* regeneration traits aligned with genome annotation.** Shown are show zoomed-in portions of Manhattan plots aligned to the genome annotation for P. trichocarpa (v3.1). Introns, untranslated regions and exons are respectively visualized with increasing thickness of bars. Labels in gray boxes were manually added to show gene IDs and the strand on which genes are found. **(A)** Results on chromosome 6 for GEMMA of Box-Cox-transformed trait Shoot PC2; **(B)** Results on chromosome 2 for GEMMA of Box-Cox transformed trait Callus/Shoot PC1, showing an association found significant via ART. These plots were made with Integrative Genomics Viewer (IGV) and boxes with Accession ID, gene name and strand were manually added underneath. Examples of plots for additional loci can be found in Fig. S14.

**Table 1. Fourteen *in planta* regeneration GWAS candidates highlighted for discussion, and Arabidopsis homologs relevant to regeneration.** Relevant literature is discussed for each of these candidates (Discussion). Distance (Dist.) of individual SNPs or SKAT/ART window centers from the transcription start site is shown for intergenic associations. Score and similarity percentage is shown for Smith-Waterman alignment of poplar gene candidates with Arabidopsis homologs, extracted from Phytozome (phytozome.jgi.doe.gov). Remaining gene candidates are summarized in Table S6-8.

[next page]

| | Gene candidates | | | | | | Arabidopsis homologs | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| Threshold | Trait | Method | Transf. | Dist. (bp) | QTL Pos. | Accession ID | Accession ID | Description | Score | Similarity |
| Bonf. | Callus Area wk. 2 | GMMAT | Binarized trait | 3 | 5' | Potri. 006G276200 | AT 3G12660 | FASCICLIN-like arabinogalactan protein 14 precursor (FLA14) | 156 | 60.40% |
| FDR (α=0.1) | Shoot Area wk. 4 | MTMC-SKAT | Untransformed | 0 | Exonic | Potri. 019G101900 | AT 4G28250 | expansin B3 | 409 | 86.90% |
| Bonf. | Shoot PC2 | GEMMA | Outliers removed, Box-Cox | 0 | Exonic | Potri. 006G254100 | AT 2G25490 | EIN3-BINDING F BOX PROTEIN 1 (EBF1) | 812 | 80.70% |
| FDR (α=0.1) | Shoot PC1 | MTMC-SKAT | Untransformed | 0 | Intragenic, non-exonic | Potri. 003G194600 | AT 3G12250 | TGACG MOTIF-BINDING FACTOR 6 (TGA6) | 520 | 91.00% |
| FDR (α=0.1) | Shoot PC1 | MTMC-SKAT | Untransformed | 4,877 | 3' | Potri. 015G041800 | AT 3G18165 | modifier of snc1,4 (MOS4) | 363 | 84.70% |
| FDR (α=0.1) | Shoot PC1 | MTMC-SKAT | Untransformed | 0 | Exonic | Potri. 002G070600 | AT 1G21326 | VQ motif-containing protein 3 (VQ3) | 105 | 65.00% |

**Table 1. Fourteen *in planta* regeneration GWAS candidates highlighted for discussion, and Arabidopsis homologs relevant to regeneration (continued)**

| | Gene candidates | | | | | | Arabidopsis homologs | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| Threshold | Trait | Method | Transf. | Dist. (bp) | QTL Pos. | Accession ID | Accession ID | Description | Score | Similarity |
| FDR (α=0.1) | Shoot PC2 | GEMMA | Outliers removed, thresholding, Box-Cox | 0 | Exonic | Potri. 002G173300 | AT 2G46560 | transducin family protein / WD-40 repeat family protein | 2357 | 66.80% |
| Bonf. | Callus Area wk. 5 | GEMMA | Box-Cox | 3,947 | 3' | Potri. 018G049600 | AT 5G35550 | TRANSPARENT TESTA 2 (TT2) | 196 | 69.90% |
| FDR (α=0.1) | Shoot PC1 | MTMC-SKAT | Untransformed | 0 | Exonic | Potri. 004G155400 | AT 1G75250 | RADIALIS-LIKE SANT/MYB 3 (RSM3) | 129 | 82.40% |
| FDR (α=0.1) | Callus 5w | GEMMA | Box-Cox | 1,192 | 5' | Potri. 010G105600 | AT 4G16110 | ARABIDOPSIS RESPONSE REGULATOR 2 (ARR2) | 382 | 75.70% |
| FDR (α=0.1) | Shoot PC2 | GEMMA | Outliers removed, thresholding, Box-Cox | 137 | 5' | Potri. 011G031100 | AT 1G11530 | C-TERMINAL CYSTEINE RESIDUE IS CHANGED TO A SERINE 1; thioredoxin | 96 | 69.30% |
| ART-Bonf. | Callus, Shoot PC1 | GEMMA | Box-Cox | 5,222 | 5' | Potri. 002G053400 | AT 1G03170 | FANTASTIC FOUR 2 (FAF2) | 103 | 54.60% |
| ART-Bonf. | Callus, Shoot PC1 | GEMMA | RB-INV | 5,222 | 5' | Potri. 002G053400 | AT 1G03170 | FANTASTIC FOUR 2 (FAF2) | 103 | 54.60% |

**Table 1. Fourteen *in planta* regeneration GWAS candidates highlighted for discussion, and Arabidopsis homologs relevant to regeneration (continued)**

| | *Gene candidates* | | | | | | *Arabidopsis homologs* | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| **Threshold** | **Trait** | **Method** | **Transf.** | **Dist. (bp)** | **QTL Pos.** | **Accession ID** | **Accession ID** | **Description** | **Score** | **Similarity** |
| ART-Bonf. | Callus Area wk. 4 | GEMMA | Box-Cox | 5,749 | 5' | Potri. 012G032900 | AT 4G27950 | CYTOKININ RESPONSE FACTOR 4 (CRF4) | 227 | 62.60% |
| ART-Bonf. | Shoot PC1 | GEMMA | RB-INV | 2,346 | 5' | Potri. 012G070400 | AT 3G52960 | PEROXIREDOXIN-II-E (PRXIIE) | 105 | 78.90% |

**Table 1. Fourteen *in planta* regeneration GWAS candidates highlighted for discussion, and Arabidopsis homologs relevant to regeneration (continued)**

**Discussion**

High-throughput phenomics support scale and precision of GWAS

Our mapping of complex developmental traits relied on computer vision methods to

precisely quantify traits that would be impractical to measure manually. Although the use

of an ordinal scoring system instead of segmentation is an alternative approach, this

would have risked the introduction of subjective biases and violation of linear model

assumptions—while sacrificing much precision and detail. The high-throughput

phenomics workflow used for this work was described, in part, by Yuan et al. (2022).

The IDEAS graphical interface for image annotation enabled the production of a large set

of training and validation examples (249 images in total) with pixelwise labels for callus,

shoot and unregenerated tissues. Here, we applied this method on a high-throughput scale

for GWAS and evaluated statistical consequences and necessary downstream adaptations,

such as appropriate trait transformations and GWAS methods. This training set enabled a

deep segmentation model that was used to automatically segment a total of 11,791

images. Although generation of the training samples was time-consuming, performing

manual segmentation for all images would have been time-prohibitive. Image annotation

using the GUI required approximately five minutes per image using IDEAS, for a total of

approximately 21 hours. Without a trained model for image segmentation, annotation of

the whole dataset would have required nearly 1,000 hours. This system or others that are

functionally comparable (Russell *et al.*, 2008; Dutta & Zisserman, 2019) can be made

more accessible and practical with innovations to reduce the number of clicks needed for

image annotation by further semi-automation of annotation. While there have been many

successes in the use of color thresholding methods that segment plants while requiring

relatively little or no training data (Yang *et al.*, 2015; Das *et al.*, 2015; Al-Tamimi *et al.*,

2016; Guo *et al.*, 2018; Gage *et al.*, 2018; Pham *et al.*, 2019; Seethepalli *et al.*, 2020;

Campbell *et al.*, 2020; Ogawa *et al.*, 2021; Affortit *et al.*, 2022), the highly variable

colors observed in callus and shoot rendered these methods inappropriate for our study.

Furthermore, in our previous work we reported a comparison of Random Forests (RF) to

our trained PSPNet model and found RF to underperform in segmentation accuracy

(Yuan *et al.*, 2022).

As reported in our prior work (Yuan *et al.*, 2022), overall accuracy in segmentation of the

"validation" set of images was 79.21% as measured by IoU, while relatively homogenous

stem tissues had IoU of 88.14%, and highly heterogenous callus tissues had 67.40%.

Imperfections in defining segmentation boundaries, whether manually or using a trained

model, leads to the introduction of statistical noise that will tend to reduce the computed

heritability of traits. However, the use of PCA over batches of traits reduced the impact

of this noise to produce traits of greater heritability. Indeed, while we studied six PC

traits and eight non-PC traits, four of the six traits with the highest heritability with were

PCs (Table S4).

Clearly, there is room for further advancement to make these phenomic workflows more

efficient and user-friendly. Advances in the architectures of deep segmentation neural

networks can contribute to improved accuracy in segmenting complex and heterogenous

tissues of interest to biologists. Further innovation in the interfaces used to generate

training sets can reduce the time cost needed for training these models. Once a training

set is obtained, the training of neural networks can require a labor-intensive process of

hyperparameter tuning. Recent developments such as Automated Machine Learning

(AutoML) can be used to automate the hyperparameter tuning process and simplify

training (Waring *et al.*, 2020; Karmaker *et al.*, 2021). Finally, the continued development

of robust and efficient GWAS methods that avoid linear model assumptions can help to

maximize the genetic insights gained from non-normal traits based on image

segmentation.

<u>Complementary GWAS approaches provide variety of insights</u>

Transformation of traits to approximate normality is commonly employed for biological

data during GWAS to avoid violating linear model assumption of normality of residuals.

In our study, because traits were computed as the proportion of plant tissue labeled by

segmentation masks as callus or shoot, and many genotypes did not produce either tissue

at early timepoints or any timepoints, the resulting distributions featured a mix of a zero

and nonzero values. Among traits in our study, the proportion of genotypes with zero

values ranged from 89 (for callus area at week five) to 1,106 (for shoot area at week two).

To avoid violations of the normality assumption arising from noncontinuous

distributions, genotypes featuring zero values were excluded from GEMMA tests for

each trait, but presence/absence tests were performed using GMMAT. GMMAT and

SKAT offer two complementary approaches to avoid problematic assumptions without

trait transformation, thus obviating the need to exclude totally recalcitrant genotypes and

thus suffer reduced statistical power.

Single-SNP methods including GEMMA and GMMAT share the advantage of providing

insights into the specific SNPs most likely to be causative with respect to the effect of a

gene on a trait. In most cases in our results, these appear to be regulatory SNPs in

promoters, suggesting that variation in gene expression, rather than sequence, is the

primary cause of trait variation. However, single-SNP methods suffer from relatively low

statistical power since by their nature they treat each SNP-trait relationship as an

independent test and do not consider combined effects of nearby SNPs. In contrast,

SKAT provides improved statistical power by allowing tests for the combined effects of

adjacent SNPs grouped into SNP windows, but only provides a single *p*-value for a whole

SNP window. Thus, our SKAT results do not make clear which SNPs in a given window

are responsible for trait variation, and as windows often overlap coding and regulatory

regions, we lack insight into whether SKAT-implicated candidates are responsible for

trait differences due to variation in their regulation or protein structure. Moreover, even

when a given window is entirely intergenic, we lack an ability for straightforward

investigation of specific promoter motifs that may be implicated by SKAT due to the lack

of single-SNP resolution. Finally, SKAT involves the upweighting of rare SNPs and

results are therefore less likely to feature top gene candidates regulated by common

variation (Wu *et al.*, 2011; Ionita-Laza *et al.*, 2013).

We therefore sought to employ a "best of both worlds" approach to improve the statistical

power of GEMMA and GMMAT by considering combined effects of adjacent common

SNPs without losing clarity into the specific SNPs most likely to be causative. To this

end, we employed ART as a post-hoc analysis of GEMMA and GMMAT results. As

ART involves the computation of combined *p*-values over SNP windows and does not

assume independence of SNPs, we obtained an increase in statistical power both via both

reduced *p*-values for SNP windows compared to individual SNPs (Vsevolozhskaya *et al.*,

2019), and by the ability to use a less-stringent Bonferroni threshold due to the number of

tests being equal to the number of 1kb SNP windows rather than the number of individual

SNPs. Our usage of ART enabled the detection of candidate genes including *FAF2*,

*CRF4* and *PRXIIE* (Table 1) that otherwise would have been missed in our study.

Although we are unaware of applied GWAS studies utilizing ART, our results

demonstrate the potential for this method to increase effective statistical power in

GWAS, which may be of particular benefit to studies using segmentation-based traits that

follow non-normal distributions and undergo transformations with a risk of reducing

power. As previously mentioned (Results), the ART-Bonferroni threshold represents the

least conservative of our three multiple-testing correction thresholds; associations

meeting this threshold alone thus represent our lowest-priority associations for

investigation and discussion.

<u>Candidate genes have diverse roles in signaling and development</u>

Our results indicate that natural variation in capabilities for *in planta* regeneration in

poplar is controlled by numerous genes with functionally diverse roles, including in cell

wall and membrane structure, hormone signaling, anthocyanin production and reactive

oxygen species (ROS) regulation. Very few gene of the gene candidates we found were

previously reported in two previous GWAS of related poplar traits, including several in a

GWAS of adventitious shoot and root in hydroponic culture (Sun *et al.*, 2019) and none

in a GWAS of *in vitro* callus regeneration (Tuskan *et al.*, 2018), likely due in part to the

different phenotyping methods, traits studied, and dependence of GWAS associations on

statistical approaches and experimental settings (Table S6-S9; Supplemental Results and

Discussion). Several of the most promising gene candidates we found, organized by

biological function of orthologs in Arabidopsis, are briefly discussed below.

*Regulation of cell wall adhesion*

Potri.006G276200 encodes a member of the FASCICLIN-LIKE ARABINOGALACTAN

(FLA) PROTEIN family and is implicated by a QTL three bases upstream of the

transcription start site. We report this association from GMMAT of callus area at week

two, the trait with greatest trait with greatest $h^2_{SNP}$ as estimated by GEMMA. The

significance of this QTL passes the most stringent multiple testing correction method

applied – the Bonferroni threshold ($\alpha = 0.05$) with each individual SNP considered an

independent test. No other QTLs associated with this trait meet the same threshold, nor

do any other QTLs from GMMAT with any trait in our study.

Many genes in the *FLA* family are differentially expressed during embryogenesis (Costa

*et al.*, 2019), but their regulation in the context of *in vitro* regeneration has received little

study. In Arabidopsis, *FLA1* was found to be upregulated during incubation on callus

induction media, while *FLA2* upregulation occurred upon transfer of explants to shoot

induction media (Johnson *et al.*, 2003). Loss-of-function of *FLA1* was reported to confer

an ability for efficient *in vitro* shoot regeneration to the otherwise recalcitrant Col-0

ecotype, while contrarily leading to loss of efficient regeneration in the regenerable

ecotype WS (Johnson *et al.*, 2011). Thus, effects of differential expression, as is likely to

be a consequence of the polymorphism from the SNP location, may be genotype-

dependent in poplar as well.

We found an association of shoot development (week four area and PC1) with a window

of SNPs including a portion of the promoter and first exon of Potri.019G101900 that is

related to Arabidopsis *EXPANSIN B3*. Expansins facilitate the process of pH-dependent

cell wall loosening, with various expansins expressed during different stages of

development. Perturbations of this gene superfamily have been studied in several plant

species, including Arabidopsis, tomatoes, rice, soybean, and tobacco. Overexpression

typically produces phenotypes of enhanced growth, such as increased size of plant cells

and tissues, as well as reduced fruit firmness. Knockout or knockdown, in contrast, leads

to reduced growth and increased firmness (Marowa *et al.*, 2016). Expansins are believed

to be key regulators of cell wall expansion downstream of auxin, a key hormone for

control of regeneration (Majda & Robert, 2018).

*Regulators of wound-responsive hormone signaling*

Potri.006G254100 is a putative homolog of *EIN3-binding F box protein 1 (EBF1)*.

Molecular evidence from Arabidopsis suggests that EBF1 facilitates ubiquitin-mediated

degradation of ETHYLENE-INSENSITIVE 3 (EIN3) and EIN3-LIKE 1 (EIL1;

Potuschak *et al.*, 2003; Shen *et al.*, 2016) and that this degradation is prevented when

EIN3 and EIL1 are stabilized by ETHYLENE-INSENSITIVE 2 (EIN2; An *et al.*, 2010).

Arabidopsis loss-of-function mutants of *EIN2* were used to supply cotyledon explant

material for an *in vitro* regeneration assay, which revealed an approximate fourfold

reduction in shoot regeneration in the mutants. The same assay revealed a roughly

threefold increase in shoot regeneration with loss-of-function of *HOOKLESS1 (HLS1)*, a

gene encoding a putative N-acetyltransferase with a role downstream of *EIN3* in

regulating a range of ethylene-regulated traits including apical hook development and *in*

*vitro* regeneration (Chatfield & Raizada, 2008). Also downstream of *EIN3* is positive and

negative regulation of numerous genes across at least eight hormone pathways including

ethylene signaling and the related jasmonate (JA) and salicylic acid (SA) signaling

pathways among others, suggesting that *EIN3* represents a key modulator of hormone

crosstalk (Chang *et al.*, 2013). Furthermore, *WOUND-INDUCED*

*DEDIFFERENTIATON (WIND)* transcription factors have been shown to regulate

ethylene, JA and SA pathways (Iwase *et al.*, 2021) and have been identified as regulators

of callus and shoot regeneration (Iwase *et al.*, 2011, 2016). In support of a key role of

*EIN3* in these pathways and in regeneration, we present at least eight gene candidates

implicated as interacting directly or indirectly with *EIN3* and upstream regulators of

EIN3 (Fig. 9). Our results, considered together with mutant studies in Arabidopsis,

suggest that these candidates regulate regeneration by mediating crosstalk between

ethylene, JA and SA signaling pathways.

**Figure 9. Proposed model integrating roles of *in planta* regeneration gene candidates.** Interactions involving Arabidopsis homologs of seven gene candidates (blue) and associated regulators were identified by literature review, providing an understanding of the broader context of hormone crosstalk between ethylene, JA, and SA pathways as they relate to regeneration. Node placement was assisted by the Force Atlas 2 algorithm as implemented in Gephi. Chemical structures for hormones were retrieved from ChemSpider (chemspider.com). Jasmonates are represented by (+)-7-Iso-Jasmonyl-L-isoleucine. This figure was produced using BioRender (biorender.com). Details for each of these gene candidates (blue) can be found in Table 1 and Discussion. Standard acronyms and abbreviations can be found on The Arabidopsis Information Resource (Berardini *et al.*, 2015) and are listed in Table S9. Evidence for interactions is summarized in Table S10.

Our GWAS results suggest a central role for SA and genes involved in SA signaling.

*NPR1* is a regulator of SA signaling via a mechanism that depends on at least three genes

with homologs implicated by QTLs in our GWAS (Fig. 9). Gene candidate

Potri.003G194600 encodes a homolog of TGACGT motif transcription factor *TGA6*.

*TGA6* and other members of the TGA family have been reported to interact with *NPR1*

(Zhang *et al.*, 1999; Subramaniam *et al.*, 2001; Fan & Dong, 2002; Rochon *et al.*, 2006;

Boyle *et al.*, 2009; Hussain *et al.*, 2018) to form a histone acetyltransferase complex

responsible for SA-associated epigenetic reprogramming (Jin *et al.*, 2018). Simultaneous

knockout of functionally redundant *TGA* genes (Zhang *et al.*, 2003b) or of *NPR1* (Cao *et*

*al.*, 1997) confers a loss of SA signaling, including SA-mediated pathogen resistance.

Moreover, a dominant mutation of the putative upstream regulator *SUPPRESSOR OF*

*NPR1, CONSTITUTIVE 1 (SNC1)* confers constitutive SA signaling, dwarf morphology,

and enhanced pathogen resistance (Zhang *et al.*, 2003a). This phenotype is reversed by

loss-of-function of *MODIFIER OF SNC1,4 (MOS4)*, a homolog of our gene candidate

Potri.015G041800 (Palma *et al.*, 2007).

Additional regulation of *EIN3* is believed to exist via phosphorylation of EIN3 protein by

the SA-regulated MAP KINASE 3 (MPK3) and MAP KINASE 6 (MPK6; Yoo *et al.*,

2008). MPK3 and MPK6 are also responsible for phosphorylation of many substrates in a

VQ MOTIF PROTEIN family related to our gene candidate Potri.002G070600. Although

the VQ proteins have not been studied in the context of *in vitro* regeneration to our

knowledge, many members are believed to function downstream of pathogen-associated

molecular patterns (PAMPs) and upstream of pathogen defense genes (Pecher *et al.*,

2014).

Our GWAS results also suggest a central role for anthocyanin and related genes. The SA

and JA pathways are linked with anthocyanin signaling by the activity of JAZ proteins in

negatively regulating MYB/bHLH/WD40 (MBW) protein complexes responsible for

transcriptional regulation of anthocyanin biosynthesis genes (Baudry *et al.*, 2004; Qi *et*

*al.*, 2011). We report two gene candidates homologous to MBW components,

Potri.002G173300 (encoding a WD-40 repeat family protein) and Potri.018G049600

(homolog of *TRANSPARENT TESTA 2*). Although we are unaware of these genes being

studied in the context of *in vitro* regeneration, MBWs regulate several steps of

anthocyanin biosynthesis downstream of naringenin chalcone, which is produced by

CHALCONE SYNTHASE (CHS; Yan *et al.*, 2021). CHS knockout in Arabidopsis

confers deficient *in vitro* shoot regeneration, with a light-dependent effect. The effects of

anthocyanins on shoot regeneration may be mediated by their effects of ROS scavenging

(Nameth *et al.*, 2013) and/or auxin accumulation (Brown *et al.*, 2001).

A functional relationship between *HLS1* (previously described; downstream of *EIN3*) and

*RSM1* (homolog of gene candidate Potri.004G155400) has been proposed due to

phenotypic similarities between *hls1* and *RSM1*-overexpressing Arabidopsis. Light-

deprived seedlings of both mutant lines presented various degrees of reduced hypocotyl

length, defective hook formation and defective gravitropism (Hamaguchi *et al.*, 2008).

However, whereas *HLS1* knockout is known to confer enhanced shoot regeneration in

Arabidopsis (Chatfield & Raizada, 2008), the effects of *RSM1* or *RSM* family

overexpression or knockout on shoot regeneration have not yet been reported to our

knowledge.

Several gene candidates from GWAS appear to affect cytokinin signaling.

Potri.010G105600 is a homolog of *ARABIDOPSIS RESPONSE REGULATOR 2 (ARR2)*,

which functions shortly downstream of cytokinin. B-type *ARR* genes such as *ARR2* share

some degree of functional redundancy and may each positively regulate *in vitro*

regeneration via transcriptional upregulation of key developmental genes such as

*WUSCHEL* (*WUS;* Xie *et al.*, 2018; reviewed by Nagle *et al.*, 2018). An additional level

of regulation over *WUS* expression exists via the *FANTASTIC FOUR (FAF)* gene family.

Overexpression of any of the four *FAF* genes (including *FAF2*, homolog of gene

candidate Potri.002G053400) leads to arrest of SAM development, possibly by inhibiting

*WUS* expression via an interaction with the feedback loop of regulation between *WUS*

and the *WUS* inhibitor *CLUVATA3* (Wahl *et al.*, 2010). Shoot meristem development is

also regulated by the *CYTOKININ RESPONSE FACTOR (CRF)* gene family (featuring

*CRF4*, a homolog of candidate Potri.012G032900), as shown by increased or reduced

rosette growth when other members of the *CRF* family are knocked out or overexpressed,

respectively. However, these experiments did not feature mutant analysis of the closely

related *CRF4* (Raines *et al.*, 2016).

*Reactive oxygen species (ROS) signaling*

At least two genes among our candidates appear to have roles in ROS regulation, which

may affect regeneration and other developmental processes by mediating post-

translational modifications of proteins involved in hormone signaling and/or by affecting

levels of oxidative damage to developing tissues. Potri.011G031100 and

Potr.012G070400 encode a putative thioredoxin-like protein and a peroxiredoxin,

respectively. Although we did not find reports of mutant phenotypes for closely related

genes in Arabidopsis in the context of regeneration or related processes, the thioredoxin

*DCC1* has been reported to affect *in vitro* shoot regeneration capacity in mutant lines as

well as across natural ecotypes of Arabidopsis (Zhang *et al.*, 2018b).

**Conclusions**

We report a GWAS of *in planta* regeneration in *P. trichocarpa* using a novel system for

phenotyping regeneration with computer vision, along with four complementary

statistical methods for association mapping. These analyses revealed over 200 candidate

genes, strongly implicating regulators of cell adhesion and stress signaling. Further

research using gene-editing or other transgenic technologies can validate these

associations and investigate the potential of these gene candidates to enhance

regeneration in genotypes of poplar and other plants that are recalcitrant to efficient

regeneration. While canonical regulators of *in vitro* regeneration tend to be involved in

auxin and cytokinin signaling pathways, our results suggest that stress pathways

downstream of ethylene, salicylic acid, and jasmonates may be of greatest importance to

the mode of *in planta* regeneration that we studied in *P. trichocarpa*. These pathways

have received little attention in studies where developmental regulator genes are used to

promote regeneration, and would appear to be promising avenues to pursue. Furthermore,

at least seven top candidates are likely closely-linked members of a genetic regulatory

network, separated from one another by no more than four degrees of direct interactions.

This, considered along with the complex nature of *in vitro* regeneration traits, suggests

that emerging multi-locus methods and epistasis tests may provide significantly greater

insights into the polygenic control of these traits. The scale and precision of this study

would not have been possible without computer vision, and our success in identifying

established regulators of regeneration demonstrates that semantic segmentation using

deep convolutional neural networks shows promise in supporting characterization of

regeneration and other complex developmental plant traits.

**Acknowledgements**

We thank the National Science Foundation Plant Genome Research Program for support

(IOS #1546900, Analysis of genes affecting plant regeneration and transformation in

poplar), and members of GREAT TREES Research Cooperative at OSU for its support of

the Strauss laboratory.

Support for the Poplar GWAS dataset is provided by the U.S. Department of Energy,

Office of Science Biological and Environmental Research (BER) via the Center for

Bioenergy Innovation (CBI) under Contract No. DE-PS02-06ER64304. The Poplar

GWAS Project used resources of the Oak Ridge Leadership Computing Facility and the

Compute and Data Environment for Science at Oak Ridge National Laboratory, which is

supported by the Office of Science of the U.S. Department of Energy under Contract No.

DE-AC05-00OR22725.

This work used the COMET high-performance cluster at the San Diego Supercomputing

Center (University of California, San Diego) made available through the Extreme Science

and Engineering Discovery Environment (XSEDE), which is supported by National

Science Foundation grant number ACI-1548562.

## Contributions of authors

Strauss, Fuxin, Jiang, and Muchero designed and directed the overall study, and obtained funding for its execution; Ma and Peremyslova designed and/or executed the phenotypic analyses; Nagle, Jiang, Yuan, and Kaur created, adapted, and executed the machine vision, computation, and data analysis pipelines; Nagle investigated gene candidates. Niño de Rivera assisted with inspecting results in IGV; Jawdy, Chen, Feng, Yates, Tuskan and Muchero supported acquisition and preparation of the GWAS population and SNP set; Nagle wrote the manuscript with editing from Strauss, and all others contributed further edits and revisions.

## References

**Adams J, Qiu Y, Xu Y, Schnable JC**. **2020**. Plant segmentation by supervised machine learning methods. *The Plant Phenome Journal* **3**: e20001.

**Affortit P, Effa-Effa B, Ndoye MS, Moukouanga D, Luchaire N, Cabrera-Bosquet L, Perálvarez M, Pilloni R, Welcker C, Champion A,** *et al.* **2022**. Physiological and genetic control of transpiration efficiency in African rice, Oryza glaberrima Steud. *Journal of Experimental Botany* **73**: 5279–5293.

**Aich S, Stavness I**. **2017**. Leaf Counting With Deep Convolutional and Deconvolutional Networks. *Proceedings of the IEEE International Conference on Computer Vision Workshops*: 2080–2089.

**Al-Tamimi N, Brien C, Oakey H, Berger B, Saade S, Ho YS, Schmöckel SM, Tester M, Negrão S**. **2016**. Salinity tolerance loci revealed in rice using high-throughput non-invasive phenotyping. *Nature Communications* **7**: 13342.

**Altpeter F, Springer NM, Bartley LE, Blechl AE, Brutnell TP, Citovsky V, Conrad LJ, Gelvin SB, Jackson DP, Kausch AP,** *et al.* **2016**. Advancing Crop Transformation in the Era of Genome Editing. *The Plant Cell* **28**: 1510–1520.

**An F, Zhao Q, Ji Y, Li W, Jiang Z, Yu X, Zhang C, Han Y, He W, Liu Y,** *et al.* **2010**. Ethylene-induced stabilization of ETHYLENE INSENSITIVE3 and EIN3-LIKE1 is mediated by proteasomal degradation of EIN3 binding F-box 1 and 2 that requires EIN2 in Arabidopsis. *The Plant Cell* **22**: 2384–2401.

**Armstrong JE, Crandell DR, Easterbrook DJ, Noble JB**. **1965**. Late Pleistocene
Stratigraphy and Chronology in Southwestern British Columbia and Northwestern
Washington. *GSA Bulletin* **76**: 321–330.

**Baudry A, Heim MA, Dubreucq B, Caboche M, Weisshaar B, Lepiniec L**. **2004**.
TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and
proanthocyanidin biosynthesis in Arabidopsis thaliana. *The Plant Journal* **39**: 366–380.

**Bdeir R, Muchero W, Yordanov Y, Tuskan GA, Busov V, Gailing O**. **2019**. Genome-
wide association studies of bark texture in Populus trichocarpa. *Tree Genetics &*
*Genomes* **15**: 14.

**Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E**. **2015**. The
arabidopsis information resource: Making and mining the "gold standard" annotated
reference plant genome. *genesis* **53**: 474–485.

**Boyle P, Le Su E, Rochon A, Shearer HL, Murmu J, Chu JY, Fobert PR, Després C**.
**2009**. The BTB/POZ Domain of the Arabidopsis Disease Resistance Protein NPR1
Interacts with the Repression Domain of TGA2 to Negate Its Function. *The Plant Cell*
**21**: 3700–3713.

**Brown DE, Rashotte AM, Murphy AS, Normanly J, Tague BW, Peer WA, Taiz L,**
**Muday GK**. **2001**. Flavonoids Act as Negative Regulators of Auxin Transport in Vivo in
Arabidopsis. *Plant Physiology* **126**: 524–535.

**Campbell MT, Grondin A, Walia H, Morota G**. **2020**. Leveraging genome-enabled
growth models to study shoot growth responses to water deficit in rice. *Journal of*
*Experimental Botany* **71**: 5669–5679.

**Cao H, Glazebrook J, Clarke JD, Volko S, Dong X**. **1997**. The Arabidopsis NPR1 gene
that controls systemic acquired resistance encodes a novel protein containing ankyrin
repeats. *Cell* **88**: 57–63.

**Carlier A, Dandrifosse S, Dumont B, Mercatoris B**. **2022**. Wheat Ear Segmentation
Based on a Multisensor System and Superpixel Classification. *Plant Phenomics* **2022**.

**Chang KN, Zhong S, Weirauch MT, Hon G, Pelizzola M, Li H, Huang SC, Schmitz**
**RJ, Urich MA, Kuo D, *et al.* 2013**. Temporal transcriptional response to ethylene gas
drives growth hormone cross-regulation in Arabidopsis (D Weigel, Ed.). *eLife* **2**: e00675.

**Chatfield SP, Raizada MN**. **2008**. Ethylene and shoot regeneration: hookless1
modulates de novo shoot organogenesis in Arabidopsis thaliana. *Plant Cell Reports* **27**:
655–666.

**Chen H, Wang C, Conomos MP, Stilp AM, Li Z, Sofer T, Szpiro AA, Chen W, Brehm JM, Celedón JC,** *et al.* **2016**. Control for Population Structure and Relatedness for Binary Traits in Genetic Association Studies via Logistic Mixed Models. *The American Journal of Human Genetics* **98**: 653–666.

**Chhetri HB, Furches A, Macaya-Sanz D, Walker AR, Kainer D, Jones P, Harman-Ware AE, Tschaplinski TJ, Jacobson D, Tuskan GA,** *et al.* **2020**. Genome-Wide Association Study of Wood Anatomical and Morphological Traits in Populus trichocarpa. *Frontiers in Plant Science* **11**: 1391.

**Costa M, Pereira AM, Pinto SC, Silva J, Pereira LG, Coimbra S**. **2019**. In silico and expression analyses of fasciclin-like arabinogalactan proteins reveal functional conservation during embryo and seed development. *Plant Reproduction* **32**: 353–370.

**Das A, Schneider H, Burridge J, Ascanio AKM, Wojciechowski T, Topp CN, Lynch JP, Weitz JS, Bucksch A**. **2015**. Digital imaging of root traits (DIRT): a high-throughput computing and collaboration platform for field-based root phenomics. *Plant Methods* **11**: 51.

**Deng W, Luo K, Li Z, Yang Y**. **2009**. A novel method for induction of plant regeneration via somatic embryogenesis. *Plant Science* **177**: 43–48.

**Di Marsico M, Paytuvi Gallart A, Sanseverino W, Aiese Cigliano R**. **2022**. GreeNC 2.0: a comprehensive database of plant long non-coding RNAs. *Nucleic Acids Research* **50**: D1442–D1447.

**Dobrescu A, Valerio Giuffrida M, Tsaftaris SA**. **2017**. Leveraging Multiple Datasets for Deep Leaf Counting. *2017 IEEE International Conference on Computer Vision Workshop (ICCVW)*: 2072–2079.

**Dutta A, Zisserman A**. **2019**. The VIA Annotation Software for Images, Audio and Video. In: MM '19. Proceedings of the 27th ACM International Conference on Multimedia. New York, NY, USA: Association for Computing Machinery, 2276–2279.

**Fan W, Dong X**. **2002**. In vivo interaction between NPR1 and transcription factor TGA2 leads to salicylic acid-mediated gene activation in Arabidopsis. *The Plant Cell* **14**: 1377–1389.

**Gage JL, White MR, Edwards JW, Kaeppler S, de Leon N**. **2018**. Selection Signatures Underlying Dramatic Male Inflorescence Transformation During Modern Hybrid Maize Breeding. *Genetics* **210**: 1125–1138.

**Gordon-Kamm B, Sardesai N, Arling M, Lowe K, Hoerster G, Betts S, Jones T**. **2019**. Using Morphogenic Genes to Improve Recovery and Regeneration of Transgenic Plants. *Plants* **8**: 38.

**Guo Z, Yang W, Chang Y, Ma X, Tu H, Xiong F, Jiang N, Feng H, Huang C, Yang**
**P,** *et al.* **2018**. Genome-Wide Association Studies of Image Traits Reveal Genetic
Architecture of Drought Resistance in Rice. *Molecular Plant* **11**: 789–805.

**Guo S, Zhou G, Wang J, Lu X, Zhao H, Zhang M, Guo X, Zhang Y**. **2022**. High-
Throughput Phenotyping Accelerates the Dissection of the Phenotypic Variation and
Genetic Architecture of Shank Vascular Bundles in Maize (Zea mays L.). *Plants* **11**:
1339.

**Hamaguchi A, Yamashino T, Koizumi N, Kiba T, Kojima M, Sakakibara H,**
**Mizuno T**. **2008**. A small subfamily of Arabidopsis RADIALIS-LIKE SANT/MYB
genes: a link to HOOKLESS1-mediated signal transduction during early morphogenesis.
*Bioscience, Biotechnology, and Biochemistry* **72**: 2687–2696.

**Hussain RMF, Sheikh AH, Haider I, Quareshy M, Linthorst HJM**. **2018**.
Arabidopsis WRKY50 and TGA Transcription Factors Synergistically Activate
Expression of PR1. *Frontiers in Plant Science* **9**: 930.

**Ionita-Laza I, Lee S, Makarov V, Buxbaum JD, Lin X**. **2013**. Sequence kernel
association tests for the combined effect of rare and common variants. *American Journal*
*of Human Genetics* **92**: 841–853.

**Iwase A, Harashima H, Ikeuchi M, Rymen B, Ohnuma M, Komaki S, Morohashi K,**
**Kurata T, Nakata M, Ohme-Takagi M,** *et al.* **2016**. WIND1 promotes shoot
regeneration through transcriptional activation of ENHANCER OF SHOOT
REGENERATION1 in Arabidopsis. *The Plant Cell* **29**: tpc.00623.2016.

**Iwase A, Kondo Y, Laohavisit A, Takebayashi A, Ikeuchi M, Matsuoka K, Asahina**
**M, Mitsuda N, Shirasu K, Fukuda H,** *et al.* **2021**. WIND transcription factors
orchestrate wound-induced callus formation, vascular reconnection and defense response
in Arabidopsis. *The New Phytologist* **232**: 734–752.

**Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M,**
**Sakakibara H, Sugimoto K,** *et al.* **2011**. The AP2/ERF transcription factor WIND1
controls cell dedifferentiation in arabidopsis. *Current Biology* **21**: 508–514.

**Jaganathan D, Ramasamy K, Sellamuthu G, Jayabalan S, Venkataraman G**. **2018**.
CRISPR for Crop Improvement: An Update Review. *Frontiers in Plant Science* **9**: 985.

**Jin H, Choi S-M, Kang M-J, Yun S-H, Kwon D-J, Noh Y-S, Noh B**. **2018**. Salicylic
acid-induced transcriptional reprogramming by the HAC–NPR1–TGA histone
acetyltransferase complex in Arabidopsis. *Nucleic Acids Research* **46**: 11712–11725.

**Johnson KL, Jones BJ, Bacic A, Schultz CJ**. **2003**. The Fasciclin-Like Arabinogalactan Proteins of Arabidopsis. A Multigene Family of Putative Cell Adhesion Molecules. *Plant Physiology* **133**: 1911–1925.

**Johnson KL, Kibble NAJ, Bacic A, Schultz CJ**. **2011**. A fasciclin-like arabinogalactan-protein (FLA) mutant of Arabidopsis thaliana, fla1, shows defects in shoot regeneration. *PloS One* **6**: e25154.

**Kamath R, Balachandra M, Vardhan A, Maheshwari U**. **2022**. Classification of paddy crop and weeds using semantic segmentation (JC and Stefania Tomasiello, Ed.). *Cogent Engineering* **9**: 2018791.

**Karmaker ("Santu") SK, Hassan MdM, Smith MJ, Xu L, Zhai C, Veeramachaneni K**. **2021**. AutoML to Date and Beyond: Challenges and Opportunities. *ACM Computing Surveys* **54**: 175:1-175:36.

**Kyritsis KA, Wang B, Sullivan J, Lyne R, Micklem G**. **2019**. InterMineR: an R package for InterMine databases. *Bioinformatics* **35**: 3206–3207.

**Lardon R, Geelen D**. **2020**. Natural Variation in Plant Pluripotency and Regeneration. *Plants* **9**: 1261.

**Lee T-H, Guo H, Wang X, Kim C, Paterson AH**. **2014**. SNPhylo: a pipeline to construct a phylogenetic tree from huge SNP data. *BMC Genomics* **15**: 162.

**Li H, Durbin R**. **2009**. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**: 1754–1760.

**Liu X, Huang M, Fan B, Buckler ES, Zhang Z**. **2016**. Iterative Usage of Fixed and Random Effect Models for Powerful and Efficient Genome-Wide Association Studies. *PLOS Genetics* **12**: e1005767.

**López-Cortegano E, Caballero A**. **2019**. Inferring the Nature of Missing Heritability in Human Traits Using Data from the GWAS Catalog. *Genetics* **212**: 891–904.

**Ma X, Deng X, Qi L, Jiang Y, Li H, Wang Y, Xing X**. **2019**. Fully convolutional network for rice seedling and weed image segmentation at the seedling stage in paddy fields. *PLOS ONE* **14**: e0215676.

**Maher MF, Nasti RA, Vollbrecht M, Starker CG, Clark MD, Voytas DF**. **2020**. Plant gene editing through de novo induction of meristems. *Nature Biotechnology* **38**: 84–89.

**Majda M, Robert S**. **2018**. The Role of Auxin in Cell Wall Expansion. *International Journal of Molecular Sciences* **19**: 951.

**Marowa P, Ding A, Kong Y**. **2016**. Expansins: roles in plant growth and potential
applications in crop improvement. *Plant Cell Reports* **35**: 949–965.

**Muchero W, Sondreli KL, Chen J-G, Urbanowicz BR, Zhang J, Singan V, Yang Y,**
**Brueggeman RS, Franco-Coronado J, Abraham N, *et al.* 2018**. Association mapping,
transcriptomics, and transient expression identify candidate genes mediating plant–
pathogen interactions in a tree. *Proceedings of the National Academy of Sciences* **115**:
11573–11578.

**Nagle M, Déjardin A, Pilate G, Strauss SH**. **2018**. Opportunities for Innovation in
Genetic Transformation of Forest Trees. *Frontiers in Plant Science* **9**: 1443.

**Nameth B, Dinka SJ, Chatfield SP, Morris A, English J, Lewis D, Oro R, Raizada**
**MN**. **2013**. The shoot regeneration capacity of excised Arabidopsis cotyledons is
established during the initial hours after injury and is modulated by a complex genetic
network of light signalling. *Plant, Cell & Environment* **36**: 68–86.

**National Academies of Sciences, Engineering, and Medicine, Division on Earth and**
**Life Studies, Board on Agriculture and Natural Resources, Committee on**
**Genetically Engineered Crops: Past Experience and Future Prospects**. **2016**.
*Genetically Engineered Crops: Experiences and Prospects*. 406-443: National
Academies Press.

**Nguyen THN, Winkelmann T, Debener T**. **2020**. Genetic analysis of callus formation
in a diversity panel of 96 rose genotypes. *Plant Cell, Tissue and Organ Culture* **142**:
505–517.

**Ogawa D, Sakamoto T, Tsunematsu H, Kanno N, Nonoue Y, Yonemaru J**. **2021**.
Haplotype analysis from unmanned aerial vehicle imagery of rice MAGIC population for
the trait dissection of biomass and plant architecture. *Journal of Experimental Botany* **72**:
2371–2382.

**Palma K, Zhao Q, Cheng YT, Bi D, Monaghan J, Cheng W, Zhang Y, Li X**. **2007**.
Regulation of plant innate immunity by three proteins in a complex conserved across the
plant and animal kingdoms. *Genes & Development* **21**: 1484–1493.

**Pecher P, Eschen-Lippold L, Herklotz S, Kuhle K, Naumann K, Bethke G, Uhrig J,**
**Weyhe M, Scheel D, Lee J**. **2014**. The Arabidopsis thaliana mitogen-activated protein
kinases MPK3 and MPK6 target a subclass of ’VQ-motif’-containing proteins to regulate
immune responses. *The New Phytologist* **203**: 592–606.

**Pham A-T, Maurer A, Pillen K, Brien C, Dowling K, Berger B, Eglinton JK, March**
**TJ**. **2019**. Genome-wide association of barley plant growth under drought stress using a
nested association mapping population. *BMC Plant Biology* **19**: 134.

**Poosapati S, Poretsky E, Dressano K, Ruiz M, Vazquez A, Sandoval E, Estrada-**
**Cardenas A, Duggal S, Lim J-H, Morris G, *et al.* 2022**. A sorghum genome-wide
association study (GWAS) identifies a WRKY transcription factor as a candidate gene
underlying sugarcane aphid (Melanaphis sacchari) resistance. *Planta* **255**: 37.

**Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV**
**der, Kling DE, Gauthier LD, Levy-Moonshine A, Roazen D, *et al.* 2018**. Scaling
accurate genetic variant discovery to tens of thousands of samples. *BioRxiv*: 201178.

**Potuschak T, Lechner E, Parmentier Y, Yanagisawa S, Grava S, Koncz C, Genschik**
**P**. **2003**. EIN3-dependent regulation of plant ethylene hormone signaling by two
arabidopsis F box proteins: EBF1 and EBF2. *Cell* **115**: 679–689.

**Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J,**
**Sklar P, de Bakker PIW, Daly MJ, *et al.* 2007**. PLINK: A Tool Set for Whole-Genome
Association and Population-Based Linkage Analyses. *The American Journal of Human*
*Genetics* **81**: 559–575.

**Qi T, Song S, Ren Q, Wu D, Huang H, Chen Y, Fan M, Peng W, Ren C, Xie D**. **2011**.
The Jasmonate-ZIM-domain proteins interact with the WD-Repeat/bHLH/MYB
complexes to regulate Jasmonate-mediated anthocyanin accumulation and trichome
initiation in Arabidopsis thaliana. *The Plant Cell* **23**: 1795–1814.

**Raines T, Shanks C, Cheng C-Y, McPherson D, Argueso CT, Kim HJ, Franco-**
**Zorrilla JM, López-Vidriero I, Solano R, Vaňková R, *et al.* 2016**. The cytokinin
response factors modulate root and shoot growth and promote leaf senescence in
Arabidopsis. *The Plant Journal* **85**: 134–147.

**Raj A, Stephens M, Pritchard JK**. **2014**. fastSTRUCTURE: variational inference of
population structure in large SNP data sets. *Genetics* **197**: 573–589.

**Revell LJ**. **2012**. phytools: an R package for phylogenetic comparative biology (and
other things). *Methods in Ecology and Evolution* **3**: 217–223.

**Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G,**
**Mesirov JP**. **2011**. Integrative genomics viewer. *Nature Biotechnology* **29**: 24–26.

**Rochon A, Boyle P, Wignes T, Fobert PR, Després C**. **2006**. The Coactivator Function
of Arabidopsis NPR1 Requires the Core of Its BTB/POZ Domain and the Oxidation of C-
Terminal Cysteines. *The Plant Cell* **18**: 3670–3685.

**Russell BC, Torralba A, Murphy KP, Freeman WT**. **2008**. LabelMe: A Database and
Web-Based Tool for Image Annotation. *International Journal of Computer Vision* **77**:
157–173.

**Seethepalli A, Guo H, Liu X, Griffiths M, Almtarfi H, Li Z, Liu S, Zare A, Fritschi**
**FB, Blancaflor EB,** *et al.* **2020**. RhizoVision Crown: An Integrated Hardware and
Software Platform for Root Crown Phenotyping. *Plant Phenomics* **2020**.

**She Y, Wang X, Gui T, Cai J**. **2019**. Lawn plant identification and segmentation based
on least squares support vector machine and multifeature fusion. *Journal of Electronic*
*Imaging* **28**: 023034.

**Shen X, Li Y, Pan Y, Zhong S**. **2016**. Activation of HLS1 by Mechanical Stress via
Ethylene-Stabilized EIN3 Is Crucial for Seedling Soil Emergence. *Frontiers in Plant*
*Science* **7**: 1571.

**Subramaniam R, Desveaux D, Spickler C, Michnick SW, Brisson N**. **2001**. Direct
visualization of protein interactions in plant cells. *Nature Biotechnology* **19**: 769–772.

**Sun P, Jia H, Zhang Y, Li J, Lu M, Hu J**. **2019**. Deciphering Genetic Architecture of
Adventitious Root and Related Shoot Traits in Populus Using QTL Mapping and RNA-
Seq Data. *International Journal of Molecular Sciences* **20**: 6114.

**Tange O**. **2020**. *GNU Parallel 20201122 ('Biden'); GNU Parallel is a general*
*parallelizer to run multiple serial command line programs in paralle lwithout changing*
*them.* Zenodo.

**Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P**. **2015**. Sambamba: fast
processing of NGS alignment formats. *Bioinformatics* **31**: 2032–2034.

**Thorvaldsdóttir H, Robinson JT, Mesirov JP**. **2013**. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Briefings in*
*Bioinformatics* **14**: 178–192.

**Towns J, Cockerill T, Dahan M, Foster I, Gaither K, Grimshaw A, Hazlewood V,**
**Lathrop S, Lifka D, Peterson GD,** *et al.* **2014**. XSEDE: Accelerating Scientific
Discovery. *Computing in Science & Engineering* **16**: 62–74.

**Tuskan GA, DiFazio S, Jansson S, Bohlmann J, Grigoriev I, Hellsten U, Putnam N,**
**Ralph S, Rombauts S, Salamov A,** *et al.* **2006**. The Genome of Black Cottonwood,
Populus trichocarpa (Torr. & Gray). *Science* **313**: 1596–1604.

**Tuskan GA, Mewalal R, Gunter LE, Palla KJ, Carter K, Jacobson DA, Jones PC,**
**Garcia BJ, Weighill DA, Hyatt PD,** *et al.* **2018**. Defining the genetic components of
callus formation: A GWAS approach. *PLoS ONE* **13**.

**Vsevolozhskaya OA, Hu F, Zaykin DV**. **2019**. Detecting Weak Signals by Combining
Small P-Values in Genetic Association Studies. *Frontiers in Genetics* **10**: 1051.

**Wahl V, Brand LH, Guo Y-L, Schmid M**. **2010**. The FANTASTIC FOUR proteins
influence shoot meristem size in Arabidopsis thaliana. *BMC Plant Biology* **10**: 285.

**Waring J, Lindvall C, Umeton R**. **2020**. Automated machine learning: Review of the
state-of-the-art and opportunities for healthcare. *Artificial Intelligence in Medicine* **104**:
101822.

**Weber JC, Stettler RF**. **1981**. Isoenzyme variation among ten populations of Populus
trichocarpa Torr. et Gray in the Pacific Northwest. *Silvae Genetica* **30**: 82–87.

**Weighill D, Tschaplinski TJ, Tuskan GA, Jacobson D**. **2019**. Data Integration in
Poplar: 'Omics Layers and Integration Strategies. *Frontiers in Genetics* **10**: 874.

**Wu MC, Lee S, Cai T, Li Y, Boehnke M, Lin X**. **2011**. Rare-Variant Association
Testing for Sequencing Data with the Sequence Kernel Association Test. *American*
*Journal of Human Genetics* **89**: 82–93.

**Xiao Q, Bai X, Zhang C, He Y**. **2021**. Advanced high-throughput plant phenotyping
techniques for genome-wide association studies: A review. *Journal of Advanced*
*Research* **35**: 215–230.

**Xie M, Chen H, Huang L, O'Neil RC, Shokhirev MN, Ecker JR**. **2018**. A B-ARR-
mediated cytokinin transcriptional network directs hormone cross-regulation and shoot
development. *Nature Communications* **9**: 1604.

**Xu Z, York LM, Seethepalli A, Bucciarelli B, Cheng H, Samac DA**. **2022**. Objective
Phenotyping of Root System Architecture Using Image Augmentation and Machine
Learning in Alfalfa (Medicago sativa L.). *Plant Phenomics* **2022**.

**Yan H, Pei X, Zhang H, Li X, Zhang X, Zhao M, Chiang VL, Sederoff RR, Zhao X**.
**2021**. MYB-Mediated Regulation of Anthocyanin Biosynthesis. *International Journal of*
*Molecular Sciences* **22**: 3103.

**Yang W, Guo Z, Huang C, Wang K, Jiang N, Feng H, Chen G, Liu Q, Xiong L**.
**2015**. Genome-wide association study of rice (Oryza sativa L.) leaf traits with a high-
throughput leaf scorer. *Journal of Experimental Botany* **66**: 5605–5615.

**Yasrab R, Atkinson JA, Wells DM, French AP, Pridmore TP, Pound MP**. **2019**.
RootNav 2.0: Deep learning for automatic navigation of complex plant root architectures.
*GigaScience* **8**: giz123.

**Yoo S-D, Cho Y-H, Tena G, Xiong Y, Sheen J**. **2008**. Dual control of nuclear EIN3 by
bifurcate MAPK cascades in C2H4 signalling. *Nature* **451**: 789–795.

**Yuan J, Kaur D, Zhou Z, Nagle M, Kiddle NG, Doshi NA, Behnoudfar A,**
**Peremyslova E, Ma C, Strauss SH,** *et al.* **2022**. Robust High-Throughput Phenotyping
with Deep Segmentation Enabled by a Web-Based Annotator. *Plant Phenomics* **2022**.

**Zhang Y, Fan W, Kinkema M, Li X, Dong X**. **1999**. Interaction of NPR1 with basic
leucine zipper protein transcription factors that bind sequences required for salicylic acid
induction of the PR-1 gene. *Proceedings of the National Academy of Sciences* **96**: 6523–
6528.

**Zhang Y, Goritschnig S, Dong X, Li X**. **2003a**. A gain-of-function mutation in a plant
disease resistance gene leads to constitutive activation of downstream signal transduction
pathways in suppressor of npr1-1, constitutive 1. *The Plant Cell* **15**: 2636–2646.

**Zhang Q, Su Z, Guo Y, Zhang S, Jiang L, Wu R**. **2020**. Genome-wide association
studies of callus differentiation for the desert tree, Populus euphratica. *Tree Physiology*
**40**: 1762–1777.

**Zhang Y, Tessaro MJ, Lassner M, Li X**. **2003b**. Knockout Analysis of Arabidopsis
Transcription Factors TGA2, TGA5, and TGA6 Reveals Their Redundant and Essential
Roles in Systemic Acquired Resistance. *The Plant Cell* **15**: 2647–2653.

**Zhang J, Yang Y, Zheng K, Xie M, Feng K, Jawdy SS, Gunter LE, Ranjan P,**
**Singan VR, Engle N,** *et al.* **2018a**. Genome-wide association studies and expression-
based quantitative trait loci analyses reveal roles of HCT2 in caffeoylquinic acid
biosynthesis and its regulation by defense-responsive transcription factors in Populus.
*New Phytologist* **220**: 502–516.

**Zhang H, Zhang TT, Liu H, Shi DY, Wang M, Bie XM, Li XG, Zhang XS**. **2018b**.
Thioredoxin-Mediated ROS Homeostasis Explains Natural Variation in Plant
Regeneration. *Plant Physiology* **176**: 2231–2250.

**Zhou X, Stephens M**. **2012**. Genome-wide efficient mixed-model analysis for
association studies. *Nature Genetics* **44**: 821–824.

**Zou K, Ge L, Zhang C, Yuan T, Li W**. **2019**. Broccoli Seedling Segmentation Based on
Support Vector Machine Combined With Color Texture Features. *IEEE Access* **7**:
168565–168574.

955