Title: Evolution of dispersal and the maintenance of fragmented metapopulations

Authors: Basile Finand, Thibaud Monnin, Nicolas Loeuille

Sorbonne Université, Université Paris Cité, Université Paris Est Créteil, CNRS, INRAE, IRD,

Institut d'Ecologie et des Sciences de l'Environnement de Paris (UMR7618), 75005 Paris,

France

ORCID numbers: BF 0000-0002-6834-1464; TM: 0000-0003-3762-942X; NL: 0000-0001-

9588-6542

corresponding author: finand.basile@gmail.com

ABSTRACT:

Because it affects dispersal risk and modifies competition levels, habitat fragmentation directly

constrains dispersal evolution. When dispersal is traded-off against competitive ability,

increased fragmentation is often expected to select higher dispersal. Such evolutionary effects

could favor the maintenance of the metapopulation by fostering spatial rescue effects. Using an

evolutionary model, we first investigate how dispersal evolves in a metapopulation when

fragmentation and aggregation of this fragmentation are fixed. Our results suggest that high

fragmentation indeed selects for dispersal increase, but this effect is largely reduced in

aggregated landscapes, to the point of being nonexistent at the highest aggregation levels.

Contrasted dispersal strategies coexist at high fragmentation levels and with no or low

aggregation. We then simulate time-varying fragmentation scenarios to investigate the

conditions under which evolutionary rescue of the metapopulation happens. Faster evolution of

dispersal favors the persistence of the metapopulation, but this effect is very reduced in

aggregated landscapes. Overall, our results highlight how the speed of evolution of dispersal

and the structuration of the fragmentation will largely constrain metapopulation survival in

changing environments.

Keywords: Dispersal evolution, fragmentation, metapopulation, evolutionary rescue, spatial

autocorrelation

 INTRODUCTION

 Dispersal, defined as the movement of individuals associated with gene flows across

 space (Ronce, 2007), is a key process in ecology and evolution. It has important consequences

 for population dynamics, changes in species distribution, maintenance of genetic diversity and

 for local adaptation (Travis *et al.*, 2013). Habitat loss and fragmentation result in decreased

 population sizes and gene flows, which undermines population viability and ultimately species

 survival. In landscapes that include suitable and unfavorable patches of varied size and

 distribution, dispersal allows individuals to move between suitable patches thereby favoring the

 survival of the metapopulation through spatial rescue effects (Levins, 1969). In a source-sink

 context, dispersal increases spatial occupancy as source populations allow the persistence of

 peripheral sink populations through dispersal (Pulliam, 1988). The maintenance of sink

 populations is especially important in the context of current changes as source-sink hierarchies

 could change in time. Given such environmental changes, dispersal helps the survival of species

 by allowing them to follow suitable niche conditions, thereby playing a key role in range

 expansions (Phillips *et al.*, 2006).

 Fragmentation creates spatial heterogeneities in several ways. It decreases the quantity

 of suitable habitat by decreasing the size and increasing the isolation of suitable patches, even

 though it increases their number (Fahrig, 2003). In our study, fragmentation is defined by the

 proportion of hostile vs. suitable locations (patches) and we systematically vary its degree of

 spatial aggregation. Dispersal evolution is affected by fragmentation, due to variations of

 various selective pressures. By definition, fragmentation increases spatial heterogeneity so that

 dispersing propagules encounter non-suitable patches more frequently. Theoretical and

 empirical studies suggest that such increases in dispersal costs and in spatial heterogeneity

 select decreased dispersal (Hastings, 1983; Travis and Dytham, 1999; Schtickzelle *et al.*, 2006;

 Bonte *et al.*, 2006; Cheptou *et al.*, 2008; Duputié and Massol, 2013). While such a

counterselection of dispersal was originally highlighted in theoretical models (Hastings, 1983;

Travis and Dytham, 1999), empirical evidence for such effects has accumulated in recent years,

for a large variety of species, from the weed *Crepis sancta* (Cheptou *et al.*, 2008)*,* to the

butterfly *Proclossiana eunomia* (Schtickzelle *et al.*, 2006) and the wolf spider *Pardosa*

monticola (Bonte *et al.*, 2006). Fragmentation however also increases inbreeding, kin

competition or temporal variation of the environment and all of these components usually select

for higher dispersal abilities (Hamilton and May, 1977; Charlesworth and Charlesworth, 1987;

Matthysen *et al.*, 1995; Gandon, 1999; Duputié and Massol, 2013; Cote *et al.*, 2017; Tung *et*

al., 2018; Oldfather *et al.*, 2021). In addition to the modulation of overall dispersal levels,

fragmentation can also, under certain conditions, maintain contrasted dispersal strategies

simultaneously. Previous investigations suggest that such a dispersal polymorphism evolves

under high fragmentation and high aggregation, with dispersing and non-dispersing individuals

coexisting within the same population (Bonte *et al.*, 2010). It principally appears because

aggregation produces a coexistence of many small patches and few large patches (Parvinen,

2002; Massol *et al.*, 2011; Parvinen *et al.*, 2020), or due to edge effects that select low dispersers

at the edge and high dispersers in central places (Travis and Dytham, 1999).

While these previous studies consider dispersal as an isolated trait, it is now widely

recognized that evolutionary changes in dispersal most often imply variations in phenotypic

traits that constrain other ecological interactions (Raffard *et al.*, 2022). It has been highlighted

that when colonization abilities are traded against competitive abilities, coexistence of a large

number of strategies is possible along this hierarchy (Tilman, 1994). Such a trade-off could for

instance occur because given a fixed quantity of energy, allocation could produce a large

number of small propagules (colonizer) or few large propagules (competitor) (eg, Smith and

Fretwell, 1974; Geritz *et al.*, 1999). For example, the weed *Crepis sancta* produces small and/or

large seeds. Small seeds have high wind dispersal due to their lightweight but low

competitiveness due to low resource storage. In contrast, large seeds have restricted dispersal

due to their weight but contain more resources (Cheptou *et al.*, 2008). In social insects, dispersal

and reproduction could follow from the production of many isolated queens that fly large

distances and have high mortality or through the split of the colony in a few propagules that

usually disperse on short distances but may be more efficient at gathering resources when

founding the new colonies (Cronin *et al.*, 2013, 2016). Fragmentation affects strategies along

the competition-colonization trade-off in different ways. First, it directly lowers the average

density at the metapopulation level, thereby changing competitive pressures. Second, it creates

isolated patches that act as a positive filter for the best dispersers. To our knowledge, only one

study considers how this competition/colonization trade-off affects the dispersal strategies

selected by fragmentation (Tilman *et al.*, 1994). This study shows that in a spatially variable

environment with an increase of fragmentation, the more competitive (and thus the less

dispersive) strategies disappear first, so that high dispersal strategies are selected.

Such results are obtained without considering explicit spatial structures as the position

of patches is not accounted for in Tilman *et al.*, 1994 (mean field approximation).

Fragmentation of the environment can however be an aggregated process, as human activities

such as urban development or agricultural exploitation are often concentrated in specific

locations. A previous work on metapopulations shows that the structuration of habitat

heterogeneities is crucial to study metapopulation responses to fragmentation (Hiebeler, 2000).

When environmental heterogeneities are spatially correlated (aggregation), predictions based

on mean-field approximation are often qualitatively incorrect when compared to spatially

explicit approaches (Hiebeler, 2000). In contrast, mean-field approximations yield correct

results in the case of randomly distributed fragmentation. Leaving out the

competition/colonization trade-off, the importance of aggregation in the evolution of dispersal

is highlighted by various studies. For example, in the context of correlated extinctions,

107 empirical work on the spider mite *Tetranychus urticae* and an associated theoretical model

108 show a selection for long-distance dispersal and a decrease of local dispersal compared to

109 spatially random extinctions (Fronhofer *et al.*, 2014). Travis and Dytham (1999) found a

110 decrease in dispersal with increased fragmentation, but an increase in dispersal with higher

111 aggregation. The risk to disperse outside of a large aggregate of suitable patches and into a

112 hostile environment is indeed lowered, so that aggregation modulates dispersal costs. Similarly,

113 (Bonte *et al.*, 2010) found a decrease of local and global dispersals with the increase of

114 fragmentation, and demonstrates that decreasing aggregation has the contrasted effect of

115 decreasing local dispersal and increasing global dispersal. To summarize, the study that

116 considers variations of dispersal strategies along a competition/colonization trade-off in

117 fragmented habitats use a spatially implicit (mean field) approach, while others use spatially

118 explicit landscapes but ignore possible competition/colonization trade-offs. The goal of our

119 study is therefore to integrate both aspects, that is to study the evolution of dispersal along the

120 competition/colonization trade-off given a spatially explicit structuration of the habitat.

121  Understanding this dispersal evolution has immediate consequences to better predict the

122 maintenance of metapopulations. For instance, a selected increase in dispersal favors the

123 exchange of individuals between patches and the colonization of empty patches (spatial rescue).

124 Extinction may also be prevented when natural selection favors adapted traits, i.e. evolutionary

125 rescue (Gomulkiewicz and Holt, 1995; Carlson *et al.*, 2014; Bell, 2017). Here, an evolutionary

126 increase of dispersal distances could avoid a population extinction in a climate change context

127 (Boeye *et al.*, 2012) or in a context of high mortality (Heino and Hanski, 2001). Given a

128 temporally increasing fragmentation, natural selection may favor high dispersal, as the

129 availability of empty and isolated patches constantly increases. Because only highly dispersive

130 strategies can reach them, such isolated patches act as filters that favor high dispersal (Heino

131 and Hanski, 2001). Consistent with this theoretical prediction, a temporal increase of

fragmentation led to higher dispersal in *Drosophila melanogaster* experiments (Tung *et al.*,

2018). Conversely, if evolution were to lead to less dispersal, it would potentially decrease

metapopulation persistence (Gyllenberg *et al.*, 2002). The implication of the evolution of

dispersal for metapopulation persistence in a world that becomes increasingly fragmented is

therefore an important, unresolved issue.

Using metapopulation simulations, we studied how the spatio-temporal structuration of

fragmented environments acts on dispersal evolution given a competition/colonization trade-

off. First, we fixed fragmentation and aggregation levels and investigated how dispersal

evolved. Second, we varied fragmentation over time to test whether dispersal evolution can

prevent extinction (evolutionary rescue), under various rates of evolution of dispersal. We

hypothesize that, in a fixed environment, higher fragmentation selects for an increase in

dispersal because more empty patches will become available to colonizers and inaccessible to

competitors. In addition, competition could be relaxed in fragmented landscapes as the average

occupancy is lowered. However, if the fragmentation is aggregated, large groups of suitable

patches could persist in the landscape. Such a situation is favorable to competitors and should

decrease the selection toward lower dispersal or lead to dispersal polymorphism with

competitors dominating aggregated patches while colonizers remain favored in isolated

patches. When fragmentation increases over time, we hypothesize that the occurrence of

evolutionary rescue depends on the speed of evolution of dispersal, which needs to be faster

than the speed of fragmentation to counterbalance its effects.

MATERIAL AND METHODS

Our simulations consider a spatially explicit environment consisting of a grid of 50x50

patches wrapped into a torus to avoid edge effects (Fig. 1). Each patch can be in one of three

possible states: unsuitable, suitable and empty or suitable and occupied. Only suitable and

empty patches are available to dispersing individuals. Importantly, we define fragmentation as

the percentage of unsuitable patches. This definition of fragmentation is classically considered

in the literature and is directly linked to other components often used to describe fragmentation

such as the number of independent patches, their size or their isolation (Fahrig, 2003). For a

given level of fragmentation, we independently vary the degree of aggregation of unsuitable

patches. Examples of landscapes can be found in the upper left part of Fig. 1. Unsuitable patches

are distributed randomly or with a set percentage of aggregation (created with a fractal

Brownian motion) using the *NLMR* and *landscapetools* package (Sciaini *et al.*, 2018). A higher

aggregation means that a suitable patch is more likely to be close to another one compared to

the random expectation.

Individuals are characterized by two traits: colonization and competition capacities

(both integer values) directly linked through the colonization/competition trade-off. Each time

step is divided in three parts: colonization, competition and extinction (see Fig. 1).

(1) Colonization. The colonization capacity defines the radius (number of patches) of the

area around the individual where its offsprings are dispersed. We assume that offspring

will potentially colonize all empty but suitable patches within this range. This implicitly

assumes that individuals with larger colonization capacity are not limited by the number

of offspring they produce. This assumption is not biologically unrealistic. For a given

amount of energy devoted to reproduction, colonizer strategies could be based on small

but numerous propagules while competitive strategy would rely on a few bigger

propagules (Smith and Fretwell, 1974). Plant species can for instance produce a lot of

small seeds that, being light, could disperse far or few big seeds that disperse near

(Cheptou *et al.*, 2008). Such a trade-off is not restricted to plants and could also apply

to various animal species. Consider, for example, ant species. Some species produce

181 new queens that disperse alone by flying, whereas others produce new queens that

182 disperse with a group of workers by walking. In the first case, the dispersing propagule

183 is made of a single queen and has a long-distance dispersal while in the second case, the

184 dispersive propagule encompasses not only the queen, but also all accompanying

185 workers, so that its energy cost is high. In the first case, the mother colony produces a

186 lot of new queens which is not the case in the second (Cronin *et al.*, 2013).

187 (2) Competition. We assume that only empty suitable patches can be colonized by

188 propagules. Given that individuals fill all suitable surrounding patches with their

189 offspring, suitable empty patches are often reached by several offspring simultaneously.

190 We then assume that the competitive hierarchy favors the strategy that has the smallest

191 dispersal distance (competition-colonization trade-off, lower part of Fig. 1). The

192 surviving individual inherits the dispersal strategy of its parents, except in the case of

193 rare mutation events. When mutation occurs, the dispersal range of the mutant

194 individual is enhanced or reduced by one cell, with equal probability. We assume that

195 established individuals (ie, occupied patches) cannot be displaced by incoming

196 propagules, regardless of their traits.

197 (3) Extinction. We assume that extinction probability does not depend on the dispersal trait.

198 Occupied suitable patches therefore become empty (but still suitable) with a fixed

199 probability e at each time step (e=0.05).

200 Each landscape is populated, at the beginning of the simulation, with ten populations

201 that are randomly distributed on suitable patches. These populations are assumed to be highly

202 dispersive (colonization capacity of twelve). We verified that initial conditions (number of

203 populations and initial colonization capacity) do not affect the equilibrium outcome

204 (Supplementary information Figure 1).

Scenario 1: Evolution of dispersal in fixed fragmented landscapes

In the first scenario, we fix the landscape and study how dispersal evolution depends on

the levels of fragmentation and aggregation. Fragmentation corresponds to a specified

percentage of unsuitable patches (i.e. 0, 20, 40, 60, 80, 90, 95 or 99% of patches are assumed

unsuitable). These unsuitable patches are aggregated at varying degrees (0, 20, 40, 60, or 80%).

Twenty different landscapes are generated for each combination of fragmentation and

aggregation. The mutation rate is set at 0.1. Each simulation lasts 50 000-time steps. Because

simulations are stochastic, they never reach a completely stable equilibrium, but we checked

that 50 000 time steps allowed the system to reach a stable regime that can be characterized

(see examples of simulations on Fig. 2b,c,d). We record the mean dispersal capacities of

individuals during the simulation and the occupancy of each dispersal strategy.

Scenario 2: Evolutionary rescue under progressively increasing fragmentation

In the second scenario, we progressively increase the level of fragmentation over time.

We systematically manipulate the rates of fragmentation and of mutation to investigate

conditions under which dispersal evolution can delay extinction. The grid is supposed to be

fully suitable at the onset of the simulation and for the first 200-time steps. We then

progressively increase fragmentation until the metapopulation becomes extinct. As in the first

scenario, the increase in fragmentation occurs with random or aggregated distributions of

unsuitable patches (levels of aggregation: 0%, 10%, 20%, 40%, 60%, 80%). Rates of

fragmentation correspond to the probability that a suitable patch becomes unsuitable within a

given time step. We tested three rates of fragmentation (0.0001, 0.001, 0.01). As evolutionary

rescue is construed as a race between the speed of the disturbance and the speed of adaptation

(Gomulkiewicz and Holt, 1995), we also systematically manipulate the speed of evolution by

considering different rates of mutation (0.001, 0.01, 0.1). We replicate each combination of

aggregation, fragmentation rate and mutation rate forty times. We record the percentage of

fragmentation at population extinction as an index of the resistance of the metapopulation to

the disturbance. Higher values of this index show that evolution of dispersal allowed the

metapopulation to survive higher levels of the disturbance. Evolutionary rescue occurs if

metapopulations with evolution of dispersal resist higher disturbance levels than

metapopulations without dispersal evolution. For each set of simulations, we also record the

variations of dispersal strategies (occupancy of the various dispersal phenotypes) over time to

identify the path that evolutionary rescue takes.

RESULTS

Evolution of dispersal in fixed fragmented landscapes

Higher fragmentation selects for increasing mean dispersal distances. In non-

fragmented landscapes, competitive strategies eventually dominate so that dispersal distance

quickly evolves close to one (Fig. 2a and b). Such a strategy remains dominant for all low levels

of fragmentation (0 to 60%). High dispersal is selected under higher fragmentation, especially

strongly when fragmentation is random (7.27±1.08 patches at 99%, red line in Fig. 2a, see also

Fig. 2c), which is congruent with Tilman *et al.* (1994). However, adding aggregation strongly

decreases this selection effect. For instance, a little bit of aggregation (20%, orange line in Fig.

2a, see also Fig. 2d) suffices to lower the selected dispersal distance in very fragmented

landscapes to 2.09±1.02 patches. Higher aggregation (40 to 80%) further decreases the selected

dispersal distance, so that fragmentation hardly has any effect on selected dispersal when

aggregation is high (blue lines, Fig. 2a). Aggregation therefore qualitatively changes the results

of mean field models (such as Tilman *et al.*, 1994).

Evolution of polymorphism in fragmented landscapes.

Beyond the observed variations in mean dispersal distances, the long-term variability of

dispersal strategies also depends on fragmentation and aggregation. Particularly, when

fragmentation is sufficient, we observe the coexistence of several dispersal strategies

(polymorphism, Fig. 3). In all cases of polymorphisms, we observe similar patterns. Suitable

patches that are close to one another sustain the less dispersive strategies, while isolated patches

act as filters that favor the more dispersive strategies. The set of polymorphic strategies however

vary depending on the relative positions of patches. For instance, given a very high

fragmentation (99%) with no or little aggregation (20% or less), random few suitable patches

are close to one another by chance (purple patches, Fig. 3b). Because the distances among these

patches is still quite important, the dispersal strategy they sustain is still quite high (around 5).

Other patches are more isolated (blue patches, Fig. 3c) and act as a selective pressure favoring

very high dispersal distances (around 9). When fragmentation is slightly lower (80 to 95%) and

aggregation slightly higher (20 to 40%), large aggregates of suitable patches occur in the

landscape (red patches, Fig. 3e) and favor competitive strategies (dispersal distance around 1).

The remaining suitable patches are isolated and favor a continuum of more dispersive strategies.

Evolutionary rescue under progressively increasing fragmentation

When fragmentation increases over time, fast dispersal evolution allows a longer

persistence of the metapopulation, i.e. an evolutionary rescue. Fig. 4 shows this evolutionary

rescue as the difference (orange arrows) between the scenario with no evolution (mutation rate

equal zero) and the three scenarios with more or less rapid evolution. Intuitively, evolutionary

rescue occurs and is strongest when there is no aggregation, fragmentation rate is low, and

mutation rate is high (mean difference of 3.04% between scenarios without and with evolution,

Fig. 4a). Evolutionary rescue is largely decreased when fragmentation rate is higher (a

difference of 0.63%, Fig. 4c). Variations in the potential of evolutionary rescue are not

continuous. Rather, a threshold mutation rate can be identified relative to the fragmentation

rate. Under our set of parameter values, evolutionary rescue occurs when the mutation rate is

ten times higher than the fragmentation rate (Fig. 4a,b,c, blue arrows). Finally, we stress that

aggregation largely constrains evolutionary rescue. No potential for evolutionary rescue can be

identified in aggregated landscapes (Fig. 4d,e,f).

DISCUSSION

Our study shows an increase of dispersal capacities in fragmented landscapes in the

context of competition/colonization trade-offs. Aggregation acts as an opposite force, as

decreased dispersal is selected in more aggregated landscapes. At high fragmentation and low

aggregation, different strategies can be selected and can coexist, with better competitors in

aggregated patches and better colonizers in isolated patches. Such an evolution of

polymorphism allows a good global coverage of available space. When fragmentation increases

with time, the rapid evolution of dispersal facilitates the survival of the metapopulation but this

evolutionary rescue effect can only be observed in non-aggregated landscapes and when

fragmentation rate is not too high.

The selection of more dispersive strategies in fragmented landscapes in a context of

competition/colonization trade-offs is congruent with Tilman *et al.* (1994) which also relies on

this trade-off. Other studies on the evolution of dispersal in spatially heterogeneous systems,

but in the absence of a competition/colonization trade-off, show a reverse pattern, as dispersal

is then counter selected because dispersal costs are enhanced by spatial heterogeneities

(Hastings, 1983; Travis and Dytham, 1999). This highlights that patterns of selection strongly

depend on the trade-off structure associated with dispersal traits. Recent works highlight the

importance of dispersal syndromes (Stevens *et al.*, 2014; Raffard *et al.*, 2022), i.e. the fact that

dispersal traits may be directly coupled to traits defining ecological interactions. The

competition/colonization trade-off falls within this category, as variations of dispersal are

directly coupled to competition hierarchies. Our results therefore highlight how such a

syndrome could lead, for some landscapes, to the selection of higher dispersal, while works that

consider evolution of dispersal alone (eg, Hastings, 1983; Travis and Dytham, 1999) would

produce the reverse pattern. Application of either framework of course depends on the types of

organisms that are considered and whether dispersal traits are competitively costly.

We explain the selection of higher dispersal in fragmented landscapes by two

mechanisms. First, fragmentation decreases overall occupancy, thus reducing the average

competition level. The advantage of competitive strategies is therefore reduced. Second,

fragmentation intensifies the strength of the competition for space which favors colonizers.

Isolated patches in fragmented landscapes can only be exploited by highly dispersive strategies.

Effects of fragmentation on mean levels of dispersal have led to contrasted results in empirical

studies (Cheptou *et al.*, 2017). Our results of a selection of higher dispersal is for instance

coherent with empirical studies of nuthatches (*Sitta europaea*) in Belgium (Matthysen *et al.*,

1995). Similarly, metapopulation study of the Glanville fritillary in Finland shows that isolated

patches of the metapopulation act as positive filters for the more dispersive strategies (Hanski

et al., 2004) and that this variation can be linked to allelic variations constraining flight

metabolism (Haag *et al.*, 2005). Conversely, a decrease of dispersal at higher fragmentation

levels has been observed in various animal or plant species (Schtickzelle *et al.*, 2006; Bonte *et*

al., 2006; Cheptou *et al.*, 2008). Spatial heterogeneity and competition decrease are two forces

that act with opposite effects, the former decreasing dispersal (higher dispersal cost) while the

latter increases it (decreased abundancies in fragmented landscapes, relax competition). The

importance of each force varies among species and needs to be systematically considered to

better predict changes in dispersal.

We found that aggregation largely reduces the selection of dispersal strategies, to the

extent that such a selection cannot even be detected when aggregation is larger than 40%. This

points out the importance of spatially explicit models. In a previous work, Hiebeler (2000)

showed how mean field approximations provide accurate occupancy predictions for random

fragmented landscapes, but not when aggregation exists. Similarly, we show here that while

our results on dispersal evolution in random landscapes are coherent with mean field

approximations (Tilman *et al.*, 1994), such approximations do not provide qualitatively

adequate variations when aggregation takes place. We explain the reduction of dispersal due to

aggregation by the fact that it favors the replacement of colonizers by competitors because of a

high probability to find a favorable patch next to another favorable patch. The landscape is there

continuous, so that competition is selected in such localities. Such a result is in line with Bonte

et al. (2010) who found an increase of local dispersal (our competitive strategy) and a decrease

of long-distance dispersal (our colonizer strategy) in aggregated fragmentation scenarios. We

therefore completely agree with the necessity of spatially explicit approaches to better

understand the dynamics of fragmented metapopulations (Hiebeler, 2000). Here, a simple

mean-field approach would yield an overestimation of dispersal evolution and of associated

evolutionary rescue effects.

We observe dispersal polymorphism when fragmentation is high and aggregation low

to intermediate. Such landscapes contain a mix of large aggregates of patches and of isolated

patches. The strategy favored in the aggregates of patches is more competitive, and the dispersal

distance selected there depends on how loose the aggregates are. When aggregation produces

continuous aggregates, the most competitive strategies are favored (dispersal distance close to

1, Fig. 3f), while when aggregates are looser, selected dispersal distances could be higher (Fig.

3f). In all cases, isolated patches favor more dispersive strategies. These results are coherent

with previous theoretical studies that show how fragmentation can favor dispersal

polymorphism. In particular, some of them showed that polymorphism is prevalent when few

large patches (our patch aggregates) co-occur with small patches (our isolated patches)

(Parvinen, 2002; Bonte *et al.*, 2010; Massol *et al.*, 2011; Parvinen *et al.*, 2020).

Evolutionary rescue can be construed as a race between speed of adaptation and of

environmental deterioration. Hence, the faster the evolution and the slower the perturbation, the

more likely the rescue. We observe that such expectations are met when fragmentation is

random (no aggregation). Dispersal evolution delays the extinction of the population when

fragmentation rate is low and mutation rate high. Fast selection of good dispersers then occurs.

As these are adapted to occupy isolated patches, such strategies foster spatial rescue in the

highly fragmented landscapes. Slow evolution would not allow that. At the onset of

fragmentation, the grid is continuous, and mostly occupied by competitors. If fragmentation is

too fast, there is no time for dispersers to appear and become selected and to fill isolated patches.

Interestingly, our study shows that rescue happens as a threshold phenomenon, being only

possible when mutation rates are higher (10 times higher in our model) than perturbation rates.

No evolutionary rescue occurs when fragmentation is aggregated, for two reasons. First,

aggregation delays extinction by itself even without evolution. Under high fragmentation and

aggregation levels, suitable patches make small continuous groups that facilitate the local

persistence of competitors. Second, in an aggregated context, dispersal evolution is absent or

strongly constrained (blue curves on Fig. 2a) so that little evolutionary potential exists. In the

absence of such an adaptive potential, evolutionary rescue cannot act. Our results on the

possibility of rescue through evolutionary changes of dispersal agree with former theoretical

works where fragmentation either stemmed from climatic changes (Boeye *et al.*, 2012) or from

heterogeneities in mortality (Heino and Hanski, 2001). While in the actual context of fast

environmental changes, it may seem complicated for species to evolve quickly enough (10

times faster than the perturbation), several examples of fast evolution of dispersal in fragmented

systems have been reported (reviewed in Cheptou *et al.*, 2017). Whether such evolutions are

sufficient to affect long term metapopulation persistence is however unknown. The fact that

evolutionary rescue does not happen here in aggregated landscapes also has important

implications. The current fragmentation of habitats is a complex non-random process that may

be frequently auto-correlated in space, therefore producing aggregated structures. For instance,

the construction of additional urban areas next to existing urban areas creates aggregated

landscapes. Studies from the Tabriz Metropolitan Area (Iran) show that the destruction of

suitable habitats surrounding the cities result in the creation of aggregated non-suitable patches

(Dadashpoor *et al.*, 2019a, 2019b). Aggregation of fragmentation can also be linked to the

displacement and development of agricultural activities. In Beijing City, China, landscape

patterns show important and complicated changes in the distribution of urban and agricultural

lands. Economic development expands cultivated land and construction into forests and

grasslands resulting in aggregated and less diverse landscapes (Li *et al.*, 2017). We propose that

when fragmentation happens in such aggregated ways, evolution will likely play a minor role

in the maintenance of the metapopulation.

Our study highlights the importance of considering dispersal syndromes (here through

the competition/colonization trade-off) and the structuration of habitat fragmentation to better

understand how dispersal evolves in disturbed landscapes. We acknowledge that our model is

quite simple and can only be used to provide baseline scenarios. For instance, fragmentation

can create changes not only in competition intensity, but also in other community aspects (eg,

presence of mutualists and enemies, see Cheptou *et al.*, 2017). While we simply focus on the

colonization-competition trade-off, evolutionary changes can also involve other phenotypic

traits. Colonization of empty patches, usually free of conspecifics, could for instance lead to

the fast evolution of intrinsic growth rates (Williams *et al.*, 2019). We hope that the results we

present here will motivate efforts to better understand the multidimensionality of dispersal

evolution and its implications for the future of biodiversity.

409 **Figure 1**: Illustration of a simulation in the context of the competition/colonization tradeoff in
410 a fixed environment scenario. The box represents the competition/colonization process when
411 two individuals arrive on the same patch.

Figure 2: Dispersal (mean +/- SD) at the end of simulations (at equilibrium) versus
environment fragmentation and aggregation (a), and over time for the 20 replicates for
conditions of no fragmentation and no aggregation (b), 99% of fragmentation and 20% of
aggregation (c) or 99% of fragmentation and no aggregation (d).

Figure 3: 2 examples of simulations showing dispersal polymorphism. (a) and (d) show the presence of dispersal strategies over time in two separate simulations. (b) and (e) represent the grid at the end of the corresponding simulations. (c) and (f) show the relative abundance of each strategy (mean percentage +/- SD) over the last 5 000-time steps. For (a) and (d) the intensity of black represents the proportion of each strategy for the given time. It is log-transformed for (d). For (b) and (e) grey patches are unsuitable, white patches are suitable and empty, and coloured patches are suitable and occupied by populations differing in dispersal strategies (ranging from low dispersal in red to high dispersal in blue). Dispersal strategies are similarly color coded in panels (c) and (f). The first row of panels (a-c) shows an example with two equally abundant dispersal distance strategies (dispersing at 5 and 9 patches). Conditions are fragmentation of 99% and no aggregation. The second row of panels (d-f) shows an example where one dispersal distance strategy (at 1.1 patches) dominates (fragmentation of 95% and aggregation of 20%).

Figure 4: Percentage of fragmentation at population extinction versus mutation rate,
fragmentation rate and aggregation. Orange arrows represent the difference of fragmentation at
population extinction between no evolution and fast evolution. Blue arrows represent the steps
between the two mutation rates where there is a difference in fragmentations at population
extinction, i.e., where the mutation rate is ten times higher than the fragmentation rate.

REFERENCES

Bell G (2017). Evolutionary Rescue. *Annu Rev Ecol Evol Syst* **48**: 605–627.

Boeye J, Travis JMJ, Stoks R, Bonte D (2012). More rapid climate change promotes
evolutionary rescue through selection for increased dispersal distance. *Evol Appl* **6**: 353–
364.

Bonte D, Borre J Vanden, Lens L, Jean-Pierre Maelfait (2006). Geographical variation in
wolf spider dispersal behaviour is related to landscape structure. *Anim Behav* **72**: 655–
662.

Bonte D, Hovestadt T, Poethke HJ (2010). Evolution of dispersal polymorphism and local
adaptation of dispersal distance in spatially structured landscapes. *Oikos* **119**: 560–566.

Carlson SM, Cunningham CJ, Westley PAH (2014). Evolutionary rescue in a changing world.
Trends Ecol Evol **29**: 521–530.

Charlesworth D, Charlesworth B (1987). Inbreeding depression and its evolutionary
consequences. *Annu Rev Ecol Syst* **18**: 237–268.

Cheptou P-O, Carrue O, Rouifed S, Cantarel A (2008). Rapid evolution of seed dispersal in
an urban environment in the weed Crepis sancta. *Proc Natl Acad Sci* **105**: 3796–3799.

Cheptou P-O, Hargreaves AL, Bonte D, Jacquemyn H (2017). Adaptation to fragmentation:
evolutionary dynamics driven by human influences. *Philos Trans R Soc B Biol Sci* **372**:
20160037.

Cote J, Bestion E, Jacob S, Travis J, Legrand D, Baguette M (2017). Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. *Ecography (Cop)* **40**: 56–
73.

Cronin AL, Loeuille N, Monnin T (2016). Strategies of offspring investment and dispersal in
a spatially structured environment: a theoretical study using ants. *BMC Ecol* **16**: 4.

Cronin AL, Molet M, Doums C, Monnin T, Peeters C (2013). Recurrent Evolution of
Dependent Colony Foundation Across Eusocial Insects. *Annu Rev Entomol* **58**: 37–55.

Dadashpoor H, Azizi P, Moghadasi M (2019a). Land use change, urbanization, and change in
landscape pattern in a metropolitan area. *Sci Total Environ* **655**: 707–719.

Dadashpoor H, Azizi P, Moghadasi M (2019b). Analyzing spatial patterns, driving forces and
predicting future growth scenarios for supporting sustainable urban growth: Evidence
from Tabriz metropolitan area, Iran. *Sustain Cities Soc* **47**: 101502.

Duputié A, Massol F (2013). An empiricist's guide to theoretical predictions on the evolution
of dispersal. *Interface Focus* **3**: 20130028.

Fahrig L (2003). Effects of Habitat Fragmentation on Biodiversity. *Annu Rev Ecol Evol Syst*
34: 487–515.

Fronhofer EA, Stelz JM, Lutz E, Poethke HJ, Bonte D (2014). Spatially correlated extinctions
select for less emigration but larger dispersal distances in the spider mite Tetranychus
urticae. *Evolution (N Y)* **68**: 1838–1844.

Gandon S (1999). Kin Competition, the Cost of Inbreeding and the Evolution of Dispersal. *J*

Theor Biol **200**: 345–364.

Geritz SAH, van der Meijden E, Metz JAJ (1999). Evolutionary Dynamics of Seed Size and
Seedling Competitive Ability. *Theor Popul Biol* **55**: 324–343.

Gomulkiewicz R, Holt RD (1995). When does Evolution by Natural Selection Prevent
Extinction ? *Evolution (N Y)* **49**: 201–207.

Gyllenberg M, Parvinen K, Dieckmann U (2002). Evolutionary suicide and evolution of
dispersal in structured metapopulations. *J Math Biol* **45**: 79–105.

Haag CR, Saastamoinen M, Marden JH, Hanski I (2005). A candidate locus for variation in
dispersal rate in a butterfly metapopulation. *Proc R Soc B Biol Sci* **272**: 2449–2456.

Hamilton WD, May RM (1977). Dispersal in stable habitats. *Nature* **269**: 578–581.

Hanski I, Erälahti C, Kankare M, Ovaskainen O, Sirén H (2004). Variation in migration
propensity among individuals maintained by landscape structure. *Ecol Lett* **7**: 958–966.

Hastings A (1983). Can spatial variation alone lead to selection for dispersal? *Theor Popul*
Biol **24**: 244–251.

Heino M, Hanski I (2001). Evolution of Migration Rate in a Spatially Realistic
Metapopulation Model. *Am Nat* **157**: 495–511.

Hiebeler D (2000). Populations on fragmented landscapes with spatially structured
heterogeneities: landscape generation and local dispersal. *Ecology* **81**: 1629–1641.

Levins R (1969). Some Demographic and Genetic Consequences of Environmental
Heterogeneity for Biological Control. *Bull Entomol Soc Am* **15**: 237–240.

Li H, Peng J, Yanxu L, Yi'na H (2017). Urbanization impact on landscape patterns in Beijing
City, China: A spatial heterogeneity perspective. *Ecol Indic* **82**: 50–60.

Massol F, Duputié A, David P, Jarne P (2011). Asymmetric patch size distribution leads to
disruptive selection on dispersal. *Evolution (N Y)* **65**: 490–500.

Matthysen E, Adriaensen F, Dhondt AA (1995). Dispersal Distances of Nuthatches , Sitta
europaea , in a Highly Fragmented Forest Habitat. **72**: 375–381.

Oldfather MF, Van Den Elzen CL, Heffernan PM, Emery NC (2021). Dispersal evolution in
temporally variable environments: implications for plant range dynamics. *Am J Bot* **108**:
1584–1594.

Parvinen K (2002). Evolutionary branching of dispersal strategies in structured
metapopulations. *J Math Biol* **45**: 106–24.

Parvinen K, Ohtsuki H, Wakano JY (2020). Evolution of dispersal in a spatially
heterogeneous population with finite patch sizes. *Proc Natl Acad Sci* **117**: 7290–7295.

Phillips BL, Brown GP, Webb JK, Shine R (2006). Invasion and the evolution of speed in
toads. *Nature* **439**: 803–803.

Pulliam HR (1988). Sources, Sinks, and Population Regulation. *Am Nat* **132**: 652–661.

Raffard A, Bestion E, Cote J, Haegeman B, Schtickzelle N, Jacob S (2022). Dispersal
syndromes can link intraspecific trait variability and meta-ecosystem functioning. *Trends*
Ecol Evol **37**: 322–331.

Ronce O (2007). How Does It Feel to Be Like a Rolling Stone? Ten Questions About
Dispersal Evolution. *Annu Rev Ecol Evol Syst* **38**: 231–253.

Schtickzelle N, Mennechez G, Baguette M (2006). Dispersal depression with habitat
fragmentation in the bog fritillary butterfly. *Ecology* **87**: 1057–1065.

Sciaini M, Fritsch M, Scherer C, Simpkins CE (2018). NLMR and landscapetools: An
integrated environment for simulating and modifying neutral landscape models in R (N
Golding, Ed.). *Methods Ecol Evol* **9**: 2240–2248.

Smith CC, Fretwell SD (1974). The Optimal Balance between Size and Number of Offspring.
Am Nat **108**: 499–506.

Stevens VM, Whitmee S, Le Galliard J-F, Clobert J, Böhning-Gaese K, Bonte D, *et al.*
(2014). A comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial
animals (J Chase, Ed.). *Ecol Lett* **17**: 1039–1052.

Tilman D (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology* **75**:
2–16.

Tilman D, May RM, Lehman CL, Nowak MA (1994). Habitat destruction and the extinction
debt. *Nature* **371**: 65–66.

Travis JMJ, Delgado M, Bocedi G, Baguette M, Bartoń K, Bonte D, *et al.* (2013). Dispersal
and species' responses to climate change. *Oikos* **122**: 1532–1540.

Travis JMJ, Dytham C (1999). Habitat persistence, habitat availability and the evolution of
dispersal. *Proc R Soc B Biol Sci* **266**: 723–728.

Tung S, Mishra A, Shreenidhi PM, Sadiq MA, Joshi S, Sruti VRS, *et al.* (2018). Simultaneous
evolution of multiple dispersal components and kernel. *Oikos* **127**: 34–44.

Williams JL, Hufbauer RA, Miller TEX (2019). How Evolution Modifies the Variability of
Range Expansion. *Trends Ecol Evol* **34**: 903–913.

548