Hyaluronic acid biosynthesis promotes an invasive, stem-like cancer cell phenotype by

broadly altering metabolism.

Adrian A. Shimpi[1*], Matthew L. Tan[1*], Michael Vilkhovoy[2], David Dai[2], L. Monet Roberts[1, 2],

Joe Kuo[2], Lingting Huang[2], Jeffrey D. Varner[2], Matthew Paszek[2,3], Claudia Fischbach[1,3,†]

[1]Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca,

New York, 14853, USA

[2]Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University,

Ithaca, New York, 14853, USA

[3]Kavli Institute at Cornell for Nanoscale Science, Cornell University, Ithaca, New York, 14853,

USA

*Contributed equally to this work

[†]Corresponding author:

Claudia Fischbach-Teschl

cf99@cornell.edu

The authors declare no potential conflicts of interest.

Word Count: 4034 without Abstract and Materials and Methods

Figures and Tables: 6 main figures, 7 supplementary figures, 1 supplementary table, 3

supplementary files

Abstract

Metastasis is the leading cause of breast cancer-related deaths and driven by invasion, but how

cancer stem-like cells (CSCs) and their metabolic phenotype regulate breast cancer invasion

remains unclear. Using microfluidic invasion models, metabolomics, and computational flux

balance analysis (FBA), we investigated the functional links between the stem-like, metabolic,

and invasive phenotype of breast cancer cells. Our results suggest that CSCs are more invasive

than non-CSCs and that broad metabolic changes associated with overproduction of hyaluronic

acid (HA) rather than transcriptional changes caused by hypoxia-inducible factor 1α (HIF1α)-

related signaling play a role in this process. Accordingly, overexpression of hyaluronic acid

synthases (HAS) 2 or 3 induced a breast cancer metabolic phenotype that promoted stemness and

invasion. These changes were clinically significant as a gene signature defined by HAS2/3

overexpression correlated with increased invasion and was predictive of worse patient survival.

Collectively, this study suggests that HA overproduction enabled by metabolic reprogramming

negatively influences prognosis in breast cancer patients by enhancing stemness and invasion

and expands the current paradigm implicating HIF1α signaling as the major driver of these

processes.

Introduction

Despite advancements in treatment options, breast cancer remains the second leading cause of cancer-related deaths in women[1]. Mortality in breast cancer is driven by metastasis and relapse, during which cancer cells in the primary tumor invade into surrounding tissues and disseminate into distant sites to form secondary tumors. Recent studies suggests that this process is mediated by a subpopulation of cancer cells known as cancer stem-like cells (CSCs)[2]. CSCs are characterized by the expression of stem-cell associated markers (e.g. NANOG, SOX2, OCT4, aldehyde dehydrogenase [ALDH]) self-renewal properties, and increased invasiveness that collectively promote tumor metastasis and subsequent relapse[3,4]. However, mechanisms that contribute to the emergence and maintenance of the CSC phenotype with consequences on tumor invasion are poorly understood.

Aberrant cellular metabolism is one of the hallmarks of cancer and has historically been described as the Warburg effect where cancer cells undergo glycolytic metabolism despite the presence of oxygen[5]. However, it is now clear that tumor cells are metabolically plastic allowing them to dynamically adapt their metabolism to accommodate spatiotemporal changes in oxygen and nutrient availability[6]. Furthermore, CSCs exhibit metabolic phenotypes distinct from non-CSCs, which influences their ability to maintain redox homeostasis providing attractive therapeutic targets[7–9]. How metabolic reprogramming promotes or maintains stem-like properties in breast cancer cells and which consequences these changes have on tumor invasion, however, remains unclear.

Increased hyaluronic acid (HA) production can induce a CSC phenotype, and previous studies

suggest that metabolic flux into the hexosamine biosynthesis pathway (HBP) and corresponding

changes in hypoxia-inducible factor 1α (HIF1α) signaling play a role in this process[10]. Whether

or not HA biosynthesis affects metabolism more broadly, however, remains undefined due to

limited methods that allow studies of metabolism at the scale of genome-wide networks.

Accumulation of HA and increased expression of HA synthase 2 (HAS2) and

glutamine:fructose-6-phosphate amidotransferase 1 (GFAT1), the rate-limiting enzyme of HBP,

have been correlated with worse breast cancer patient survival[11–13]. These results suggest a

prominent role for HAS2, but HA is produced by several HAS isoforms that vary in enzyme

kinetics, nutrient sensitivity, and the formation of microvilli-like structures on the cell surface

suggestive of differences in cell-surface retention[14–17]. Moreover, clinical studies suggest that

increased HA production irrespective of the HAS isoform is associated with worse tumor

differentiation[11,12]. Which role HAS isotype-independent metabolic reprogramming plays in this

process, how HA overproduction affects metabolism besides altering HBP and HIF1α, and

which effect these changes have on the CSC phenotype and invasion have been understudied.

Here, we use a combination of engineered cell lines, microfabricated culture models,

computational approaches, and patient survival analysis to investigate the interconnectedness of

the CSC phenotype, HA production, and metabolism and its influence on cancer cell invasion.

We demonstrate that increased HA production promotes stemness due to broad metabolic

changes. These changes stimulate cancer cell invasive potential irrespective of HAS isoform and

were associated with worse patient survival. Collectively, our results suggest that HA

overproduction due to metabolic reprogramming negatively influences prognosis in breast cancer

patients by altering invasion.

Results

Cancer Stem-Like Cells Exhibit Increased Invasive Potential

To investigate differences in invasion between CSCs and their non-differentiated counterparts,

we utilized the CSC reporter cell line GFP-NANOG MDA-MB-231 in which green fluorescent

protein (GFP) expression is controlled by the NANOG promoter[18] (Fig. 1a). Using fluorescence-

activated cell sorting (FACS) GFP-NANOG MDA-MB-231 cells were sorted into GFP^{Null},

GFP^{Low} (bottom 5% GFP expressing), and GFP^{High} (top 5% GFP expressing cells) populations to

enrich for different stem-like states (Fig. 1a) whose phenotype was maintained over 5 days of

culture (Supplementary Fig. 1). Moreover, the GFP^{High} cell population proliferated more slowly

than the non-CSC GFP^{Null} population, consistent with a more quiescent phenotype (Fig. 1b). To

study potential differences in the invasive phenotype of these different cell populations, we

monitored tumor cell invasion in response to a morphogen gradient using a microfluidic collagen

type I hydrogel model (Fig. 1c). In this setup, GFP^{High} NANOG MDA-MB-231 cells invaded

into the 3D collagen hydrogel region of the device more readily than their less stem-like

counterparts, validating that CSCs exhibit increased invasive potential (Fig. 1c,d). Cell migration

and invasion are energetically intensive processes requiring increased metabolic consumption[19].

Accordingly, inhibiting energy production with the glycolysis inhibitor 2-deoxyglucose (2-DG)

reduced both collagen hydrogel invasion and random 2D migration of unsorted GFP-NANOG

MDA-MB-231 cells (Fig. 1e,f). These results indicate that CSCs exhibit increased invasive and

migratory potential in our experimental setup that was dependent on glycolytic metabolism.

Cancer Stem-Like Cells Have Altered Metabolism

To characterize the metabolic phenotype of cancer cells as a function of their stem-like

phenotype, extracellular metabolite production and consumption rates were measured in sorted

GFP-NANOG cells (Fig. 2a,b). The more stem-like cells consumed more glucose and produced

more lactate relative to their non-stem-like counterparts indicative of increased glycolysis (Fig.

2a). Real-time metabolic analysis with the Agilent Seahorse (Seahorse) Analyzer of extracellular

acidification rate (ECAR) and oxygen consumption rate (OCR) further suggested that CSCs not

only exhibited increased glycolysis, but also oxidative phosphorylation (Fig. 2b, Supplementary

Fig. 2). Acccordingly, culturing in glucose-free media decreased the percentage of GFP[High] cells

relative to media containing physiological levels of glucose (Fig. 2c). This approach also

decreased the ALDH bright (ALDH[Br]) fraction (a marker for breast CSCs[20,21]) of wildtype

MDA-MB-231 cells confirming that our results were not an artefact of the GFP-NANOG

reporter cell line. Moreover, inhibition of glycolysis by 2-DG reduced the GFP[High] population

further corroborating that the CSC phenotype is intimately linked to glucose metabolism (Fig.

2d).

As increased HA production has been associated with both metabolic reprogramming and the

CSC phenotype[10,13], we next assessed how HA production correlates with the CSC phenotype.

Immunofluorescence (IF) image analysis identified that GFP expression in GFP-NANOG MDA-

MB-231 cells positively correlated with the amount of cell surface-associated HA (Fig. 2e).

These differences were likely due to changes in HA biosynthesis and subsequent cell surface

retention as analysis of secreted HA by ELISA indicated low levels across all experimental

conditions that were not significantly different from each other although more stem-like cells

seemed to secrete slightly less HA (Supplementary Fig. 3a). Together, these results suggest that

more stem-like cells exhibit increased glucose metabolism relative to their less stem-like

counterparts and that these changes correlate with elevated levels of cell surface-associated HA.

Hyaluronic Acid Production Correlates with Stemness

To better quantify the relationship between glucose metabolism, cell surface-associated HA, and

stemness, GFP-NANOG MDA-MB-231 cells were stained for HA and then subjected to flow

cytometry and targeted metabolomics. Consistent with the IF results, the amount of cell surface-

associated HA directly correlated with GFP expression levels (Fig. 3a). Importantly, sorting the

GFP-NANOG MDA-MB-231 cells into high and low HA producing cells for targeted

metabolomics revealed broad changes of intracellular metabolites with hierarchical clustering

separating the two HA production phenotypes (Fig. 3b). In particular, both upper glycolytic and

tricarbocylic acid (TCA) cycle metabolites such as glucose-6-phosphate (G6P), fructose-6-

phosphate (F6P), oxaloacetate (OAA), and citrate (CIT) were increased in the highly HA

producing cells (Fig. 3c). Combining these results with the broad metabolic alterations of CSCs

described above (Fig. 2), these data suggest a direct link between glycolysis-dependent HA

synthesis and the CSC phenotype (Fig. 3d). To more directly test how HA synthesis affects

stemness, GFP-NANOG MDA-MB-231 cells were cultured with the HA synthesis inhibitor 4-

methylumbelliferone (4-MU, 0.5 mM). This approach resulted in a decrease of stem-like GFP[High]

cells consistent with previous studies using primary murine breast cancer cells cells[22]

(Supplementary Fig. 3b). However, this effect was not significant in our hands possibly due to

differences in the utilization of UDP-glucuronic acid (UDP-GlcUA, a subunit of HA) by

different cell lines[23], which is depleted by 4-MU treatment[24].

To more directly determine if increasing production of HA promotes stemness, HAS2 and HAS3

were stably overexpressed in MDA-MB-231 and the non-malignant breast epithelial cell line

MCF10A. While HA is synthesized by three HAS isoforms (HAS1, 2, 3), HAS 1 varies in HA

synthesis and secretion rate from HAS 2 and 3. In contrast, HAS2 and HAS3 exhibit similar

sensitivities and responses to precursor availability and thus were used here to study their effects

on the metabolic reprogramming[14–16]. Overexpression of HAS2/3 increased the defining

characteristic of CSCs in both cell lines: self-renewal as measured through sphere formation in a

limited dilution assay (Fig. 3e, Supplementary Fig. 4a,b). Additionally, overexpression of both

HAS2/3 markedly increased the fraction of CD44$^+$/CD24$^-$ MCF10A cells, which characterizes an

invasive breast CSC population with more mesenchymal characteristics[25,26] (Fig. 3f,

Supplementary Fig. 4d). This population could not be assessed in HAS overexpressing MDA-

MB-231 as this cell line contains an intrinsically high fraction of CD44+/CD24- cells (>85%),

making changes difficult to quantify[27,28] (Supplementary Fig. 4c). Nevertheless, HAS3

overexpression increased the ALDH^{Br} fraction in MCF10A, albeit in a non-statistically

significant trend (Supplementary Fig. 4e). Together, these results suggest that HA

overproduction increases stem-like cell properties in breast cancer cells regardless of cell line

and HAS2/3 isoform.

Increased HA Production by CSCs is Associated with Increased Glucose Conversion and ATP

Production

Given our results that metabolic reprogramming of CSCs supports their energy demands during

invasion (Fig. 2) and that a more stem-like phenotype is associated with increased HA

production (Fig. 3), we speculated that increased HA production promotes a more energetic CSC

phenotype. Probing the individual contribution of HA biosynthetic pathways to the metabolic

state of CSCs solely by measuring different metabolite levels, however, is challenging given the

interconnectedness of most metabolic pathways. To circumvent these limitations and delineate

the contribution of HA biosynthesis to CSC metabolism more directly, a flux balance analysis

(FBA) model was constructed. FBA is a widely utilized mathematical approach to model the

flow of metabolites (flux) through a genome-scale network of metabolic pathways, including

glycolysis, oxidative phosphorylation, hexosamine biosynthesis, and amino acid consumption[29],

and has been successfully used to investigate cancer metabolism[30,31]. In contrast to traditional

metabolomics, FBA also enables the simulation of a desired metabolic phenotype by adjusting

model parameters such as the objective function and flux constraints. To develop the model for

our study, extracellular metabolomics of GFP[High] and GFP[Null] cells were performed over 72

hours to define metabolite consumption/production profiles used to constrain flux values for

more and less stem-like MDA-MB-231 breast cancer cells, respectively (Fig. 4a). Additionally,

the objective function of the FBA model was set to maximize HA synthesis to study both the

capacity of CSCs and non-CSCs to produce HA and the associated changes in metabolic flux.

Results from the FBA model indicated that the more stem-like cells (GFP[High]) increased flux

through the upper stages of glycolysis (i.e., conversion of glucose to fructose-6-phosphate), ATP

generation, lactate production/secretion, and glutamine uptake relative to the non-stem-like cells

(GFP[Null]) (Fig. 4b, Supplementary File 1). Additionally, the FBA model indicated increased flux

through all HA synthesis intermediate steps for the GFP[high] cells relative to the GFP[null] cells (Fig.

4c). Collectively, these data suggest that the increased capacity of CSCs to synthesize HA is

related to increased glucose conversion but simultaneously allows these cells to produce ATP

more efficiently. To confirm the predictive value of the FBA model, a Seahorse real-time ATP

rate assay was performed. This analysis verified that the mitochondrial and overall ATP

production rate was increased in GFP^{High} versus GFP^{Null} cells (Fig. 4d,e). Together, these

findings suggest that the metabolic phenotype of more stem-like cancer cells leading to increased

HA biosynthesis promotes ATP production by these cells. FBA models are suitable for

predicting these changes.

Metabolic Changes resulting from Increased HA Production Stimulate a Stem-like Breast

Cancer Cell Phenotype to Promote Invasion.

Our aforementioned results suggest functional links between the stem-like and invasive

phenotype of tumor cells (Fig. 1), CSCs and metabolism (Fig. 2), CSCs and HA (Fig. 3), and HA

and metabolism (Fig. 4). Furthermore, HA has been previously associated with increased

invasion[32,33]. However, whether or not these single observations are mechanistically connected

remained to be determined. Therefore, we next hypothesized that the metabolic phenotype

induced by HA overproduction increases the stem-like phenotype of breast cancer cells to

promote invasion. Indeed, Seahorse analysis revealed that MCF10A and MDA-MB-231 cells

overexpressing HAS2 and HAS3 had increased ECAR compared to their parental controls

suggesting an increase in glycolytic energy production (Fig. 5a). Interestingly, OCR was

unchanged in HAS2/3-overexpressing MCF10A but increased in HAS2/3-overexpressing MDA-

MB-231 cells (Supplementary Fig. 5a), consistent with an increase in OCR in GFP^{High} vs.

GFP^{Low} and GFP^{Null} MDA-MB-231 (Fig. 2b). Treatment with 2-DG decreased both cell surface-

associated and secreted HA in MCF10A cells (Supplementary Fig. 5b, c). Notably, 2-DG

treatment of MCF10A cells decreased the CD44$^+$/CD24$^-$ fraction in the HA overproducing cells

but had no effect on the parental control cells suggesting that the enhanced glycolysis associated

with HA overproduction is critical to maintaining stemness (Fig. 5b). Furthermore, HA

overproduction increased both total and glycolytic ATP production (Fig. 5c, Supplementary Fig.

5d) that 2-DG reversed to similar levels as in MCF10A control cells (Fig. 5c). A similar trend

was noted for MDA-MB-231 cells, although 2-DG decreased glycolytic ATP production only in

the HAS3 overexpressing cells. Overexpression of HAS2/3 also increased invasion of both

MCF10A and MDA-MB-231 (Fig. 5d, e), consistent with their increased stem-like phenotype

(Fig. 3d, e). This effect was inhibited by 2-DG treatment, implicating a functional consequence

of HA-mediated stemness and metabolism in tumor cell invasion (Fig. 5d, e). As 2-DG treatment

did not affect the ATP production rate in parental cells (Fig. 5c), these changes in invasion can

be attributed to differences in metabolism rather than compromised cell viability. Together this

data suggests that HA overproduction induces a glycolytic phenotype that is crucial for CSC-

mediated invasion.

HA Overproduction Transcriptome Changes Predict Worse Patient Survival

Our data suggest that both HAS2 and 3 induce phenotypes associated with increased invasion

and, thus, malignancy. RNA sequencing was conducted on the MCF10A cells overexpressing

HAS2 and HAS3 and their parental control to identify transcriptional changes that would allow

us to interrogate the contribution of HA overproduction to stemness, invasion, and patient

outcomes. Principle component analysis (PCA) and hierarchical clustering indicated that the

transcriptome of HA overproducing cells differed significantly from their parental control (Fig.

6a, b, Supplementary Fig. 6a, Supplementary File 2). Gene set enrichment analysis (GSEA) with

the Hallmarks gene sets from the Molecular Signature Database revealed enrichment of a wide

range of pathways for both HAS2 and HAS3-overexpressing cells, including those previously

associated with stemness and invasion such as NF-κB[34], hypoxia[35], PI3K signaling[36], IL6-

STAT3 signaling, and reactive oxygen species[7,8] (Fig 6c). Since HIF1α signaling has been

previously implicated in an HA-dependent increase in the CSC phenotype[10], we performed

GSEA for HIF1α target genes[37–41]. Interestingly, neither HAS2 nor HAS3 transcriptomes were

enriched for HIF1α target genes (Fig. 6d), which have previously been suggested as drivers of

HA-dependent stemness[10]. These results further support our findings that the cellular phenotypes

investigated in this study are due to broader metabolic changes and cannot be attributed to

glycolysis-driven changes in hypoxia-related signaling.

To determine the clinical relevance of our findings, a 72-gene signature was generated from the

genes upregulated in both HAS2 and HAS3-overexpressing cells (log_2-fold change > 1, p-value

< 0.05) compared to their parental control (Fig. 6e, Supplementary Table 1). Overrepresentation

analysis of this gene signature surprisingly revealed no significant enrichment of genes

associated with most metabolic pathways from the KEGG database (Supplementary Fig. 6b)

possibly due to the fact that energy production can be regulated beyond transcriptional

changes[42,43]. Subsequently, we calculated enrichment scores for this HA overproduction gene

signature and a curated list of gene signatures involved in migration, cytoskeleton dynamics, and

metabolism (Supplementary File 3, Supplementary Fig. 7a) using single sample GSEA

(ssGSEA)[44] for chemotherapy-naïve patients in the METABRIC cohort. Tumorous tissue from

patients enriched for our HA overproduction signature were simultaneously enriched for several

migration and actin cytoskeleton gene sets suggesting increased tumor invasion in these patients,

while metabolic gene sets unrelated to glycosaminoglycan synthesis were not enriched after

correcting for random associations (Fig. 6f, Supplementary Fig. 7b,c). Importantly, our HA

overproduction signature also predicted worse patient survival consistent with their increased

expression of invasion-related gene sets (Fig. 6f,g). Gene signatures specific to upregulation of

either HAS2 or HAS3 also predicted worse patient survival that was not seen in the parental

control (Supplementary Fig. 6c). Collectively, our results suggest that genes associated with

increased HA biosynthesis correlate with an enrichment of migratory genes and predict worse

patient survival regardless of HAS2/3 isoform.

Discussion

Although CSCs are known to exhibit a distinct metabolic phenotype from non-CSCs, the exact

nature of their metabolic alterations and consequences on tumor invasion, a hallmark of

metastasis and worse patient prognosis, remain unclear. Indeed, past studies have reported that

CSCs have either a primarily glycolytic phenotype[9,45] or increased mitochondrial oxidative

metabolism[46,47]. Furthermore, the functional link between CSC metabolism and invasion is

understudied although invasion is known to be a highly energetic process and is often driven by

more stem-like cancer cells. Here, we demonstrate that CSCs can increase glycolytic and

oxidative metabolism simultaneously and that these changes in metabolism promote stemness

and invasion due to increased HA production. Moreover, we identified that these cellular

changes correlated with a gene expression signature that was independent of differential

expression of HIF1α target genes, but predictive of patient survival.

The energetic demands associated with HA-mediated invasion can induce broad metabolic

changes. For example, increased HA production rapidly depletes UDP-sugar substrates, which

cancer cells may compensate for by increasing glycolysis to maintain flux into the HBP and thus,

the pool of UDP-sugars[10]. Our FBA model corroborates this phenomenon and provides further

insight into metabolic states associated with CSCs such as an increased HA production capacity.

The resulting metabolic phenotypes associated with overexpressing HAS2 or HAS3 in cells

suggest that the glycolytic demand dominates over possible changes in oxidative metabolism

previously associated with stemness[8,48,49]. Furthermore, our finding that HA overproduction

induces phenotypic and transcriptomic changes that correlate with invasion may help explain

why glycolytic, mesenchymal breast CSCs localize to the leading edge of tumors and worsen

patient prognosis[8,26]. Together, these results suggest that HA production enables a more invasive,

malignant CSC metabolic phenotype.

To interrogate alterations in other metabolic pathways associated with glycolysis and oxidative

phosphorylation, we utilized FBA to predict changes in metabolic fluxes induced by stemness.

FBA is especially proficient in enabling the study of cancer metabolism as it simulates metabolic

phenotypes based on real-world constraints such as cell growth rate and glucose uptake[29,50]. The

FBA model developed here indicated that stem-like cells exhibited an increased capacity for HA

production, and that this increase in HA production was associated with broad metabolic

alterations. Interestingly, our model suggested that although the conversion of glucose to

pyruvate is decreased in stem-like cells when maximizing HA production, the production of both

lactate and acetyl-CoA from pyruvate is increased. An alternative source of pyruvate are the

malic enzymes, which convert malate to pyruvate while producing the reducing agent NADPH[51].

Indeed, our FBA model indicated an increase in the flux through malic enzyme 2 (ME2), which

operates within the mitochondria (Supplementary File 1). Increased ME2 activity has been

previously associated with the loss of cellular senescence and increased tumorigenesis[52,53].

Additionally, ME2 plays a crucial role in the production of glutamine which provides a source of

n-acetylglucosamine, one of the building blocks of HA[54]. The FBA model developed here

supports our results that CSCs are associated with increased HA production and energy

production. However, any connection with malic enzyme activity will need to be experimentally

verified in future experiments, as is typical in interpreting FBA predictions.

Throughout the metastatic cascade, HA has been implicated in promoting invasive phenotypes

and to support survival of circulating cells in the vasculature[32,33,55]. While HA-dependent

changes of cancer malignancy and stemness have been primarily attributed to upregulation of

HAS2[13,22], our results indicate that HAS3 similarly promotes a stem-like state (Fig. 3). Indeed,

the fact that 2-DG inhibited the tumor cell stem-like phenotype in both HAS2/3-overexpressing

cells suggests that the metabolic reprogramming enacted by HA overproduction may be the

central regulator of the tumor stem-like phenotype (Fig. 5b). HA production is regulated by

cytoplasmic pools of UDP sugars, which are elevated in cancerous breast tissue[56,57]. The

conversion of UDP-glucose to UDP-glucuronic acid can regulate metastasis through *SNAI1*

mRNA stabilization[58], which we identified as upregulated in HA overproducing cells (Fig. 6,

Supplementary File 2). Interestingly, degradation of HA produced by cancer cells promotes

glucose uptake that can promote migration and further fuel metabolism[59]. Combined with our

results, this suggests HA intermediates generated by HA-induced metabolic reprogramming may

help regulate metastatic phenotypes, but further experiments delineating the contributions of HA-

associated metabolic intermediates on the CSC phenotype are required.

Transcriptomic analysis of HAS2 and HAS3 overexpressing MCF10As indicated an enrichment

of multiple stemness-associated gene sets (e.g. IL6-JAK-STAT3 signaling and hypoxia), but

metabolic genes were not differentially expressed. This discrepancy with our experimental

observations may be explained by the fact that metabolic changes are not only regulated

transcriptionally but also by enzyme activity levels and localization[42,43]. For example, the activity

of phosphofructokinase (PFK), a key enzyme controlling glycolytic flux[60], is regulated by

glycosylation with O-linked β-N-acetylglucosamine (O-GlcNAc)[61]. Our FBA model indicates

that CSCs have an increased HBP capacity to produce UDP-GlcNAc, the sugar donor for O-

GlcNAcylation, which would be expected to decrease PFK activity and thus glycolysis[61].

Instead, our results suggest an increase rather than decrease in glycolysis, possibly because flux

rates are not regulated soley by cytoplasmic enzymes levels, but also through their relative

activities and substrates availability[42,60,62]. Furthermore, cytoskeletal rearrangement, which is

critical for invasion, can independently control glycolytic flux by mediating enzyme

degradation[63] or sequestration[43]. Our results suggest that the glycolytic CSC phenotype

associated with HA overproduction is not regulated transcriptionally, but whether this is enacted

by cytoskeletal dynamics or changes in the relative activity of different metabolic enzymes

requires further investigation.

Our data suggests that cell states associated with increased glycolysis enrich for highly invasive

CSCs that are characterized by increased HA production. Interestingly, HA produced by CSCs is

primarily retained on the cell surface rather than excreted into the surrounding environment (Fig.

3, Supplementary Fig. 3). Consequentially, the retention of HA on the cell surface contributes to

glycocalyx thickness whose biophysical properties impact the interactions between cells and

their surrounding environment including cell-extracellular matrix interactions necessary for

migration[17,64]. In particular, HA-dependent changes of the glycocalyx can promote adhesion-

independent or ameboid migration by altering the friction required for force generation against

extracellular structures[65]. Alternatively, changes in glycocalyx thickness impact surface receptor

diffusion patterns and generate kinetic traps that lead to receptor clustering to strengthen

adhesion assembly and activate mechanosignaling[66,67]. Further studies to delineate the

contributing biophysical properties are needed to determine the influence of HA on predominant

migration mode. Moreover, migration is an energy-intensive process and the necessary ATP

utilization can determine migration decisions. Therefore, the increased ATP production observed

in the stem-like and HA-overproducing cells highlights the need to decouple glycocalyx-

mediated, biophysical determinants from metabolic reprogramming. Overall our results provide a

possible explanation for why CSCs are associated with increased metastasis. Nevertheless,

further characterization of the biophysical and metabolic phenotype of CSCs and their

contributions to invasion and metastasis are necessary to define additional therapeutic targets.

Acknowledgements

Thank you to all members of the Fischbach lab for valuable feedback and discussion of the

research. The authors would also thank Dr. Andrea Di Micheli, David McKellar, and Dr. Ben

Cosgrove for assistance with RNA-sequencing data processing. The work described was

supported by National Cancer Institute through the Center on the Physics of Cancer Metabolism

(1U54CA210184), the National Science Foundation (NSF) through Graduate Research

Fellowship Program awarded to A.A.S and L.M.R. (DGE-1650441), the U.S. Department of

Education through the Graduate Assistance in Areas of National Need (GAANN) Fellowship

awarded to M.L.T., the Breast Cancer Coalition of Rochester, and the Cornell NanoScale

Science & Technology Facility (CNF), a member of the National Nanotechnology Coordinated

Infrastructure (NNCI) supported by the National Science Foundation (Grant NNCI-2025233).

This work utilized facilities at the Cornell Institute of Biotechnology including the Cornell

University Biotechnology Resource Center (BRC) Imaging Facility (RRID:SCR_021741) for

image acquisition using the Zeiss LSM 710 Confocal Microscope (NIH 1S10RR025502), the

BRC Genomics Facility (RRID:SCR_021727) for sequencing, and the BRC Flow Cytometry

Facility (RRID:SCR_021740) for cell sorting with BD Biosciences FACSAria Fusion. Mass

spectrometric analysis of targeted metabolomics was performed at the Weill Cornell Medicine

Proteomics and Metabolomics Core Facility.

Author Contributions

A.A.S., M.L.T., and C.F. designed the study. A.A.S. and M.L.T. conducted most of the

experiments. M.V. and D.D. conducted extracellular metabolomics for FBA analysis. L.M.R.

and M.P. performed FACS for HA production. J.K., L.H., and M.P. generated plasmids for HAS

overexpression. L.M.R. and A.A.S. generated cell lines. J.V. and M.L.T. performed FBA

analysis. A.A.S., M.L.T., and C.F. analyzed the data and wrote the manuscript. All authors

discussed the results and commented on the manuscript.

Materials and Methods

Cell Culture

MCF10A (ATCC) cells were cultured in DMEM/F12 supplemented with 5% Horse Serum, 1%

penicillin/streptomycin, 10 µg/mL Insulin, 0.5 µg/mL Hydrocortisone, 100 ng/mL Cholera toxin,

and 20 ng/mL human epidermal growth factor (EGF). MDA-MB-231 (ATCC) and the GFP-

NANOG MDA-MB-231 (a kind gift from Dr. Ofer Reizes[18]) cells were cultured in DMEM

supplemented with 10% FBS and 1% penicillin/streptomycin. HAS overexpressing cells were

cultured in their respective medias containing 1 µg/mL doxycycline (Santa Cruz Biotechnology).

For glycolytic inhibition studies, cells were treated with media containing 2-deoxyglucose

(MilliporeSigma) matching the glucose concentration in the media (25mM or 20mM for DMEM,

DMEM/F12 respectively) for 24 hours unless otherwise noted.

Generated Cell Lines

cDNAs for human HAS2 and HAS3 were generated and cloned into the lentiviral vector pLV

HygroR tetOn to create stably transduced, tetracycline-inducible MCF10A cell lines as

previously described[17]. cDNAs for human HAS2 and HAS3 were also fabricated and inserted

into a tetracycline-inducible PiggyBac expression vector to generate pPB huHAS2-IRES2-

mScarlet-IRES2-NeoR and pPB huHAS3-IRES2-mScarlet-IRES2-NeoR through custom gene

synthesis (Twist Biosience). Generation of the MDA-MB-231 HAS overexpressing cells was

conducted using either the pPB huHAS2-IRES2-mScarlet-IRES2-NeoR or pPB huHAS3-IRES2-

mScarlet-IRES2-NeoR or with the Piggybac transponase using the Nucleofector Cell Line Kit V

(Lonza). Stably transfected cells were selected using 1 µg/mL puromycin (MilliporeSigma) or

800 µg/mL G418 (ThermoFisher). MDA-MB-231 HAS overexpressing cells were then sorted

post-selection on mScarlet expression levels.

FACS and Flow Cytometry

GFP-NANOG MDA-MB-231 cells were trypsinized and resuspended at 10×10^6 cells/mL in

FACS buffer (2.5% FBS/PBS, 2mM EDTA) and processed on the BD FACSAria Fusion Cell

Sorter. The top 5% and bottom 5% of the GFP+ fraction were designated as GFP[High] and GFP[Low]

respectively, while the non-GFP expressing population were designated as GFP[Null]. Cells were

sorted into cell culture media, recounted, seeded, and allowed to rest for 24 hours before use in

experiments.

Cell sorting for targeted metabolomics was conducted on GFP-NANOG MDA-MB-231 cells

stained for HA using Alexfluor-568 (ThermoFisher) conjugated HA binding protein (HABP, 40

µg/mL, MilliporeSigma). The high HA (top 5%) or low HA (bottom 5%) fractions were sorted

using a BD FACSAria Fusion Cell Sorter.

ALDH activity was determined for MDA-MB-231 and MCF10A cells using the Aldefluor™

Assay (STEMCELL Technologies) according to manufacturer instructions with incubation

conducted for 30min at 37°C. Analysis was conducted on a BD Accuri C6 Plus analyzer.

The fraction of CD44+/CD24- cells was determined by trypsinizing and resuspending cells in

FACS buffer at 10×10^6 cells/mL followed by incubation with antibodies against human CD44

(APC-conjugated, Clone G44-26, 1:5, BD Biosciences) and CD24 (PE-Cy7-conjugated, Clone

ML5, 1:20, BD Biosciences). Gates were determined using the isotype controls mouse anti-

IgG2b κ (APC-conjugated, Clone 27-35, 1:5, BD Biosciences) and mouse anti-IgG2a κ (PE-

Cy7-conjugated, Clone G155-178, 1:20, BD Biosciences). Cells were analyzed on a BD Accuri

C6 Plus Analyzer.

Invasion and Migration Assays

To prepare microfluidic invasion assays, rat tail Type I collagen (Corning) was neutralized with

1 N NaOH and diluted with 1X DMEM to a final concentration of 2.5 mg/mL. Before

neutralization and dilution, 10x DMEM was added to collagen as a pH indicator to a final

concentration of 10% v/v. Microfluidic 3D cell culture devices (AIM Biotech) were then injected

with the 2.5 mg/mL rat tail Type I collagen solution into the center channel of the chip. The

chips were then incubated at 4°C for 30 minutes, followed by incubation at 37°C for 30 minutes

to complete collagen polymerization. For MDA-MB-231 invasion, DMEM containing 1% FBS

and 1% penicillin/streptomycin was injected into the left flanking media channel, and DMEM

containing 10% FBS and 1% penicillin/streptomycin was injected into the right flanking media

channel. For MCF10A invasion, MCF10A media devoid of Horse Serum and EGF was injected

into the left flanking media channel, and fully supplemented MCF10A media was injected into

the right flanking media channel. 10 μL of a 2.5×10^5 cell suspension of GFP-NANOG MDA-

MB-231 or MCF10A cells were then injected into both ports of the left channel (20 μL total).

Cells were allowed to invade through the hydrogel channel for 3 - 5 days before fixation in a 4%

paraformaldehyde solution. Media was exchanged every 24 hours. After fixation, chips were

stained, imaged, and individual cell invasion distance was measured using ImageJ by segmenting

nuclei.

To assess random migration ability, GFP-NANOG MDA-MB-231 cells were plated on

fibronectin-coated (30 μg/mL) glass 96 well plates and placed in an Incucyte S3 (Sartorius) live

cell imaging system. Images were obtained in 20-minute intervals over 24 hours. Individual cell

tracking was performed using ImageJ to determine migration velocity (motility) and random

migration paths.

Immunofluorescence

GFP-NANOG MDA-MB-231 cells were plated on fibronectin-coated (30 µg/mL) glass

coverslips at 2500 cells/cm^2. Cells were then fixed in 4% paraformaldehyde (PFA)/PBS (w/v) for

20 min at room temperature, blocked with 1% bovine serum albumin (BSA)/PBS (w/v) for 1 hr

at room temperature, and incubated overnight with AlexaFluor-568 conjugated hyaluronic acid

binding protein (HABP, 13.3 µg/mL, MilliporeSigma) at 4°C. Afterwards cells were incubated

with DAPI (2.5 µg/mL, ThermoFisher) for 30 min at room temperature.

For invasion experiments, devices were fixed in 4% paraformaldehyde (PFA)/PBS for 30 min at

room temperature, permeabilized with 0.1% Triton X-100/PBS (v/v) for 15 min at room

temperature, blocked with 1% bovine serum albumin (BSA)/PBS (w/v) for 1 hr at room

temperature, and incubated with DAPI (2.5 µg/mL) and either AlexaFluor-568 (ThermoFisher)

or AlexaFluor-647 (ThermoFisher) phalloidin to visualize F-actin.

Confocal Microscopy and Image Analysis

Images were acquired on a Zeiss LSM 710 confocal microscope with either a LD LCI Plan-

Apochromat 25×/0.8 Imm Korr DIC M27 or C-Apochromat W M27 10x/0.45 objective. Images

were analyzed using ImageJ with custom scripts. Briefly, for single cell intensity measurements,

z-stacks were sum projected and cells were segmented based on HA intensity. Cell clusters were

manually corrected to individual cells, while overlapping or edge-located cells were excluded

from analysis.

For invasion experiments, the C-Apochromat W M27 10x/0.45 objective was used at 0.6x zoom,

and a z-stack was obtained along a 2 mm length in the center of the device. Z-stacks were

maximum intensity projected, and invasion was assessed based on nuclei displacement along the

x-axis (across the hydrogel region). Invasion distance was normalized to the appropriate control

in each set of replicates per experiment, and data was pooled together across device replicates to

obtain averages per condition.

Metabolic Analysis

To obtain an initial measure of glycolytic ability, sorted GFP-NANOG MDA-MB-231 cells were

seeded in 24 well plates at 10,000 cells/cm^2 in 1 mL of media. Glucose concentration was

measured using a GlucCell Glucose Monitoring System (CESCO Bioengineering), and lactate

concentration was obtained using a colormetric lactate assay kit (Sigma).

Targeted metabolomics was conducted on GFP-NANOG MDA-MB-231 cells 48 hours post

sorting for HA production. Media was collected and non-adherent cells were pelleted at 500 x g

for 4 minutes. Meanwhile, 0.5 mL of 80% methanol (MetOH) was added onto adherent cells and

incubated at -80°C, After aspirating supernatant, the cell pellet was resuspended in 0.5 mL of

80% MetOH and added to the respective well of adherent cells in plate and incubated at -80°C

for 15 minutes. Cells were then scraped using a cell scraper and collected into an Eppendorf tube

and pelleted at 20,000 x g for 10 minutes at 4°C. Supernatant was then transferred to 2 mL

screw-cap vial and dried overnight under vaccum at room temperature. Following overnight

drying, samples were then dried for 2.5 hours in SpeedVac SPD 1030 at room temperature and

then stored at -80°C until analyzed by Weill Cornell Medicine Proteomics and Metabolomics

Core Facility.

Real-time changes in metabolism were tested using a Seahorse XFe96 Analyzer in conjunction

with the Seahorse XF Glycolysis Stress Test Assay Kit and the Seahorse XF Real-time ATP

Assay Rate Kit (Agilent). Manufacturer instructions were followed to perform each of the

assays. Wildtype and modified MDA-MB-231 cells were seeded on a Seahorse XFe96 Cell

Culture microplate at 20,000 cells per well in standard media and allowed to attach overnight.

For experiments with MCF10A, cells were seeded at 20,000 cells per well. HAS overexpressing

cells had media changed 24 hours after seeding to include 1 µg/mL doxycycline. Media was then

changed to the specific Seahorse assay media, and plates were prepared according to

manufacturer instructions for each assay kit. Relevant metabolic values from each assay were

calculated using template worksheets provided by Agilent. After the assays were complete, DNA

was extracted from each well using Caron's Buffer (25 mM Tris-HCl, 0.4 M NaCl, 0.5% (w/v)

sodium dodecylsulfate), and total DNA content measured using the fluorometric Quantifluor

dsDNA Assay (VWR) and converted to cell number for normalization.

Flux Balance Analysis (FBA)

To generate a computational model of metabolic fluxes in sorted GFP-NANOG MDA-MB-231

cells, cells were sorted as described above and seeded into 24 well plates at 10,000 cells/cm^2.

Cells were allowed to attach overnight before a fresh media change. Media was collected 72

hours after the initial media change and stored at -80°C before performing extracellular

metabolomics to measure levels of glucose, lactate, and the 20 amino acids. Glucose was

measured using Contour next EZ Blood Glucose Monitoring System (Ascensia Diabetes Care)

using 5 µL as the sample volume. Lactate and amino acid concentrations were assayed using an

Acquity UPLC H-Class System equipped with QDa and tunable UV (TUV) detectors controlled

by Empower 3 software (Waters Corporation). Specifically, extracellular amino acids were

analyzed using a Waters AccQ-Tag Ultra Derivatization Kit (Waters) according to the

manufacturer's recommendations. Derivatized samples were injected onto an AccQ-Tag Ultra

C18 column (1.7 μm, 2.1 mm x 100 mm, Waters) and detected by an Acquity TUV detector

(Waters) at 260 nm. Amino acids were identified by known retention times of standards and

concentrations were determined by comparison with calibration standard curves. For the lactate

measurements, samples were first deproteinized by treatment with an equal volume of

trichloroacetic acid followed by centrifugation at 12,000 x g for 10 minutes. 200 μL of the

supernatant was then combined with 600 μL of acetonitrile (ACN) before injecting 2 μL into the

LC-MS system. Lactate was quantified using a standard curve ranging from 0.05mM to 1mM.

Separations were performed on an Acquity UPLC BEH Amide Column (1.7 μm, 2.1 mm x 100

mm, Waters). Solvent A consisted of 50:50 ACN:Water and solvent B consisted of 95:5

ACN:Water. The solvent gradient started at 0.01% solvent A and 99.9% solvent B, raised to 40%

A in 0.5 minutes, further raised to 70% A in 1.5 minutes, and returned to initial conditions over

0.1 minute and held for 3 minutes to re-equilibrate the column. The flow rate was set to 0.6

mL/min, the autosampler was set to 5°C, and the column was set to 50°C. The mass-to-charge

ratio (m/z) of lactate was 88.9. Analysis was performed in negative ion mode with a cone voltage

of 15V and probe temperature of 600°C.

Extracellular metabolomics data of all 20 amino acids, lactate, and glucose was then used to

constrain a flux balance analysis by imposing bounds on allowable fluxes. Growth rate and O2

fluxes were constrained from cell counts and Seahorse data respectively. The model used in this

study was implemented in the Julia programming language[68], where the linear programming

problem was solved using the GNU Linear Programming Kit (GLPK) package

(https://www.gnu.org/software/glpk/). The stoichiometric matrix and metabolic growth

requirements were derived from a previously developed Core Cancer model[31]. The objective

function was set to maximize HA production subject to experimentally estimated rates of uptake,

secretion, and cell growth to define the phenotype of interest; 1000 simulations were performed

for each condition. A subset of the FBA results representing the energetic pathways of interest

was incorporated into a graphical representation (Fig. 4). The full FBA results can be found in

Supplementary File 1.

Limited Dilution Assay

Cells were serially diluted and seeded into ultra-low attachment 96-well plates (Corning) in 200

μL of serum-free DMEM/F-12 containing 2% B27, 10 ng/mL basic fibroblast growth factor, 20

ng/mL epidermal growth factor, 10 μg/mL insulin, and 1 μg/mL doxycycline. Sphere formation

was assessed after two weeks of culture, and the number of spheres was counted using a phase-

contrast microscope. The stem cell frequency was determined using the extreme limited dilution

algorithm[69].

RNA-Sequencing

Parental (NeoR-rtTA) and HA overproducing MCF10A cells were seeded on 10-cm dishes at

5000 cells/cm^2 and allowed to adhere overnight. Media was refreshed to include 1 μg/mL of

doxycycline to induce HAS2 and HAS3 expression and cultured for an additional 48 hours. RNA

was isolated using the Qiagen RNeasy kit according to manufacturer instructions. RNA libraries

were prepared using the Illumina TruSeq RNA Kit, and single-ended 75bp read lengths were

sequenced on the Illumina NextSeq 500 system.

Sequence Alignment and Gene Set Enrichment Analysis

Reads were trimmed using TrimGalore version 0.4.4

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and aligned to the human

reference genome GRCh38 (ENSEMBL) using STAR version 2.6.0a[70]. Reads of genomic

features were counted using featureCounts[71], and differential gene expression was determined

using DESeq2[72]. Differentially expressed genes in HA overproducing cells were defined as a

log-2 fold change greater than 1 and an adjusted p-value less than 0.0001 compared to the

parental cells. Genes common to both HAS2 and HAS3 were combined to generate the HA

overproducing gene signature for survival analysis. Individual HAS2 and HAS3 gene signatures

were defined as the differentially expressed genes with log 2-fold change greater than 2 or 5,

respectively, and a p-value less than 0.0001.

Gene set enrichment analysis (GSEA) was conducted with a ranked list generated by taking the

sign of the fold change multiplied by the log-10 of the adjusted p-value. The list was inputted to

the GSEA Java applet (http://software.broadinstitute.org/gsea/index.jsp) using the

GSEAPreRanked tool and the Hallmarks gene sets from MSigDB v.7.0. Gene sets were

considered significantly enriched with a p-value and FDR value ≤ 0.05.

Patient Survival and Enrichment Analysis

Patient data from the METABRIC cohort was extracted from the Cancer Genomics Data Server.

Patients were limited to those having not received chemotherapy. Gene signature scores were

calculated using the single-sample GSEA (ssGSEA)[44] with the GSVA package[73]. The top and

bottom quartiles of the ssGSEA scores for the 72-gene HA overproduction signature were

designated as high and low scores, respectively, for survival and enrichment analysis. Kaplan-

Meier survival analysis was conducted with the *survival* package in R using a Cox proportional

hazard model with statistical significance determined using a log-rank test. Stratified patients

were further analyzed for enrichment of curated lists of migration, cytoskeletal, and metabolic

gene signatures obtained from the MSigDB v7.0. To correct for random associations, 300

randomly selected gene signatures with the same number of genes (15-500 genes) as gene sets in

the curated list had ssGSEA enrichment scores additionally calculated. An emprical cumulative

distribution function was established using the *ecdf* function in R and a p-value cutoff was

determined where 95% of values fell below.

Statistical analysis

All experiments were performed with at least three independent biological replicates unless

otherwise noted. Pairwise comparisons were conducted using a Mann-Whitney U test unless

otherwise noted. Multiple comparisons were evaluated with either a Kruskal-Wallis Test or two-

way ANOVA with Dunn's post hoc analysis. Results were considered statistically significant

with a p-value less than 0.05. Unless otherwise noted, all data points are plotted mean +/- the

standard deviation. All statistically analysis was performed using GraphPad Prism v9.3 or R.

628
References

1. Siegel, R. L., Miller, K. D. & Jemal, A. Cancer statistics, 2020. *CA. Cancer J. Clin.* **70**, 7–
30 (2020).

2. Batlle, E. & Clevers, H. Cancer stem cells revisited. *Nat. Med.* **23**, 1124–1134 (2017).

3. Pattabiraman, D. R. & Weinberg, R. A. Tackling the cancer stem cells — what challenges
do they pose? *Nat. Rev. Drug Discov.* **13**, 497–512 (2014).

4. Wu, C.-W. (Ken), Wang, M.-L. & Chiou, S.-H. Targeting cancer stem cells: emerging
role of Nanog transcription factor. *Onco. Targets. Ther.* **6**, 1207 (2013).

5. DeBerardinis, R. J. & Chandel, N. S. Fundamentals of cancer metabolism. *Sci. Adv.* **2**,

e1600200 (2016).

6.   Fendt, S.-M., Frezza, C. & Erez, A. Targeting Metabolic Plasticity and Flexibility
Dynamics for Cancer Therapy. *Cancer Discov.* **10**, 1797–1807 (2020).

7.   Diehn, M. *et al.* Association of reactive oxygen species levels and radioresistance in
cancer stem cells. *Nature* **458**, 780–783 (2009).

8.   Luo, M. *et al.* Targeting Breast Cancer Stem Cell State Equilibrium through Modulation
of Redox Signaling. *Cell Metab.* **28**, 69-86.e6 (2018).

9.   Peiris-Pagès, M., Martinez-Outschoorn, U. E., Pestell, R. G., Sotgia, F. & Lisanti, M. P.
Cancer stem cell metabolism. *Breast Cancer Res.* **18**, 55 (2016).

10.   Chanmee, T. *et al.* Hyaluronan Production Regulates Metabolic and Cancer Stem-like
Properties of Breast Cancer Cells via Hexosamine Biosynthetic Pathway-coupled HIF-1
Signaling. *J. Biol. Chem.* **291**, 24105–24120 (2016).

11.   Auvinen, P. *et al.* Hyaluronan in Peritumoral Stroma and Malignant Cells Associates with
Breast Cancer Spreading and Predicts Survival. *Am. J. Pathol.* **156**, 529–536 (2000).

12.   Auvinen, P. *et al.* Hyaluronan synthases (HAS1–3) in stromal and malignant cells
correlate with breast cancer grade and predict patient survival. *Breast Cancer Res. Treat.*
143, 277–286 (2014).

13.   Chokchaitaweesuk, C., Kobayashi, T., Izumikawa, T. & Itano, N. Enhanced hexosamine
metabolism drives metabolic and signaling networks involving hyaluronan production and
O-GlcNAcylation to exacerbate breast cancer. *Cell Death Dis.* **10**, 803 (2019).

14.   Itano, N. *et al.* Three Isoforms of Mammalian Hyaluronan Synthases Have Distinct
Enzymatic Properties. *J. Biol. Chem.* **274**, 25085–25092 (1999).

15.   Rilla, K. *et al.* Hyaluronan Synthase 1 (HAS1) Requires Higher Cellular UDP-GlcNAc
Concentration than HAS2 and HAS3*. *J. Biol. Chem.* **288**, 5973–5983 (2013).

16.   Rilla, K. *et al.* Hyaluronan production enhances shedding of plasma membrane-derived
microvesicles. *Exp. Cell Res.* **319**, 2006–2018 (2013).

17.   Shurer, C. R. *et al.* Physical Principles of Membrane Shape Regulation by the Glycocalyx.
Cell **177**, 1757-1770.e21 (2019).

18.   Thiagarajan, P. S. *et al.* Development of a Fluorescent Reporter System to Delineate
Cancer Stem Cells in Triple-Negative Breast Cancer. *Stem Cells* **33**, 2114–2125 (2015).

19.   Romani, P., Valcarcel-Jimenez, L., Frezza, C. & Dupont, S. Crosstalk between
mechanotransduction and metabolism. *Nat. Rev. Mol. Cell Biol.* **22**, 22–38 (2021).

20.   Ginestier, C. *et al.* ALDH1 Is a Marker of Normal and Malignant Human Mammary Stem

Cells and a Predictor of Poor Clinical Outcome. *Cell Stem Cell* **1**, 555–567 (2007).

21.   Vincent, K. M., Findlay, S. D. & Postovit, L. M. Assessing breast cancer cell lines as
tumour models by comparison of mRNA expression profiles. *Breast Cancer Res.* **17**, 114
(2015).

22.   Chanmee, T. *et al.* Excessive Hyaluronan Production Promotes Acquisition of Cancer
Stem Cell Signatures through the Coordinated Regulation of Twist and the Transforming
Growth Factor β (TGF-β)-Snail Signaling Axis. *J. Biol. Chem.* **289**, 26038–26056 (2014).

23.   Kultti, A. *et al.* 4-Methylumbelliferone inhibits hyaluronan synthesis by depletion of
cellular UDP-glucuronic acid and downregulation of hyaluronan synthase 2 and 3. *Exp.*
Cell Res. **315**, 1914–1923 (2009).

24.   Kakizaki, I. *et al.* A Novel Mechanism for the Inhibition of Hyaluronan Biosynthesis by
4-Methylumbelliferone. *J. Biol. Chem.* **279**, 33281–33289 (2004).

25.   Al-Hajj, M., Wicha, M. S., Benito-Hernandez, A., Morrison, S. J. & Clarke, M. F.
Prospective identification of tumorigenic breast cancer cells. *Proc. Natl. Acad. Sci.* **100**,
3983–3988 (2003).

26.   Liu, S. *et al.* Breast Cancer Stem Cells Transition between Epithelial and Mesenchymal
States Reflective of their Normal Counterparts. *Stem Cell Reports* **2**, 78–91 (2014).

27.   Li, W. *et al.* Unraveling the roles of CD44/CD24 and ALDH1 as cancer stem cell markers
in tumorigenesis and metastasis. *Sci. Rep.* **7**, 13856 (2017).

28.   Sheridan, C. *et al.* CD44+/CD24-breast cancer cells exhibit enhanced invasive properties:
an early step necessary for metastasis. *Breast Cancer Res.* **8**, R59 (2006).

29.   Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.* **28**,
245–248 (2010).

30.   Asgari, Y., Zabihinpour, Z., Salehzadeh-Yazdi, A., Schreiber, F. & Masoudi-Nejad, A.
Alterations in cancer cell metabolism: The Warburg effect and metabolic adaptation.
Genomics **105**, 275–281 (2015).

31.   Zielinski, D. C. *et al.* Systems biology analysis of drivers underlying hallmarks of cancer
cell metabolism. *Sci. Rep.* **7**, 1–14 (2017).

32.   Brett, M.-E. *et al.* In vitro elucidation of the role of pericellular matrix in metastatic
extravasation and invasion of breast carcinoma cells. *Integr. Biol.* **10**, 242–252 (2018).

33.   Offeddu, G. S. *et al.* The cancer glycocalyx mediates intravascular adhesion and
extravasation during metastatic dissemination. *Commun. Biol.* **4**, 255 (2021).

34.   Korkaya, H., Liu, S. & Wicha, M. S. Breast cancer stem cells, cytokine networks, and the
tumor microenvironment. *J. Clin. Invest.* **121**, 3804–3809 (2011).

35.   Pang, M.-F. *et al.* Tissue Stiffness and Hypoxia Modulate the Integrin-Linked Kinase ILK
706   to Control Breast Cancer Stem-like Cells. *Cancer Res.* **76**, 5277–5287 (2016).

707   36.   Yoon, C. *et al.* PI3K/Akt pathway and Nanog maintain cancer stem cells in sarcomas.
708   *Oncogenesis* **10**, 12 (2021).

709   37.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection.
710   *Cell Syst.* **1**, 417–425 (2015).

711   38.   Miroshnikova, Y. A. *et al.* Tissue mechanics promote IDH1-dependent HIF1α–tenascin C
712   feedback to regulate glioblastoma aggression. *Nat. Cell Biol.* **18**, 1336–1345 (2016).

713   39.   Schaefer, C. F. *et al.* PID: the Pathway Interaction Database. *Nucleic Acids Res.* **37**,
714   D674–D679 (2009).

715   40.   Semenza, G. L. Hypoxia-inducible factor 1: oxygen homeostasis and disease
716   pathophysiology. *Trends Mol. Med.* **7**, 345–350 (2001).

717   41.   Slemc, L. & Kunej, T. Transcription factor HIF1A: downstream targets, associated
718   pathways, polymorphic hypoxia response element (HRE) sites, and initiative for
719   standardization of reporting in scientific literature. *Tumor Biol.* **37**, 14851–14861 (2016).

720   42.   Hackett, S. R. *et al.* Systems-level analysis of mechanisms regulating yeast metabolic
721   flux. *Science (80-.).* **354**, aaf2786–aaf2786 (2016).

722   43.   Hu, H. *et al.* Phosphoinositide 3-Kinase Regulates Glycolysis through Mobilization of
723   Aldolase from the Actin Cytoskeleton. *Cell* **164**, 433–446 (2016).

724   44.   Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven
725   cancers require TBK1. *Nature* **462**, 108–112 (2009).

726   45.   Chen, C.-L. *et al.* NANOG Metabolically Reprograms Tumor-Initiating Stem-like Cells
727   through Tumorigenic Changes in Oxidative Phosphorylation and Fatty Acid Metabolism.
728   *Cell Metab.* **23**, 206–219 (2016).

729   46.   Pastò, A. *et al.* Cancer stem cells from epithelial ovarian cancer patients privilege
730   oxidative phosphorylation, and resist glucose deprivation. *Oncotarget* **5**, 4305–4319
731   (2014).

732   47.   Vlashi, E. *et al.* Metabolic differences in breast cancer stem cells and differentiated
733   progeny. *Breast Cancer Res. Treat.* **146**, 525–534 (2014).

734   48.   Dong, C. *et al.* Loss of FBP1 by Snail-Mediated Repression Provides Metabolic
735   Advantages in Basal-like Breast Cancer. *Cancer Cell* **23**, 316–331 (2013).

736   49.   Ciavardelli, D. *et al.* Breast cancer stem cells rely on fermentative glycolysis and are
737   sensitive to 2-deoxyglucose treatment. *Cell Death Dis.* **5**, e1336–e1336 (2014).

50. Lewis, N. E. & Abdel-Haleem, A. M. The evolution of genome-scale models of cancer
metabolism. *Front. Physiol.* **4**, 1–7 (2013).

51. Simmen, F. A., Alhallak, I. & Simmen, R. C. M. Malic enzyme 1 (ME1) in the biology of
cancer: it is not just intermediary metabolism. *J. Mol. Endocrinol.* **65**, R77–R90 (2020).

52. Jiang, P., Du, W., Mancuso, A., Wellen, K. E. & Yang, X. Reciprocal regulation of p53
and malic enzymes modulates metabolism and senescence. *Nature* **493**, 689–693 (2013).

53. Cheng, C.-P. *et al.* The mechanisms of malic enzyme 2 in the tumorigenesis of human
gliomas. *Oncotarget* **7**, 41460–41472 (2016).

54. Sarfraz, I. *et al.* Malic enzyme 2 as a potential therapeutic drug target for cancer. *IUBMB*
Life **70**, 1076–1083 (2018).

55. Turley, E. A., Wood, D. K. & McCarthy, J. B. Carcinoma Cell Hyaluronan as a "Portable"
Cancerized Prometastatic Microenvironment. *Cancer Res.* **76**, 2507–2512 (2016).

56. Oikari, S. *et al.* UDP-sugar accumulation drives hyaluronan synthesis in breast cancer.
Matrix Biol. **67**, 63–74 (2018).

57. Caon, I. *et al.* Sirtuin 1 reduces hyaluronan synthase 2 expression by inhibiting nuclear
translocation of NF-κB and expression of the long-noncoding RNA HAS2–AS1. *J. Biol.*
Chem. **295**, 3485–3496 (2020).

58. Wang, X. *et al.* UDP-glucose accelerates SNAI1 mRNA decay and impairs lung cancer
metastasis. *Nature* **571**, 127–131 (2019).

59. Sullivan, W. J. *et al.* Extracellular Matrix Remodeling Regulates Glucose Metabolism
through TXNIP Destabilization. *Cell* **175**, 117-132.e21 (2018).

60. Tanner, L. B. *et al.* Four Key Steps Control Glycolytic Flux in Mammalian Cells. *Cell*
Syst. **7**, 49-62.e8 (2018).

61. Yi, W. *et al.* Phosphofructokinase 1 Glycosylation Regulates Cell Growth and
Metabolism. *Science (80-.).* **337**, 975–980 (2012).

62. Liberti, M. V. *et al.* A Predictive Model for Selective Targeting of the Warburg Effect
through GAPDH Inhibition with a Natural Product. *Cell Metab.* **26**, 648-659.e8 (2017).

63. Park, J. S. *et al.* Mechanical regulation of glycolysis via cytoskeleton architecture. *Nature*
578, 621–626 (2020).

64. Kuo, J. C.-H., Gandhi, J. G., Zia, R. N. & Paszek, M. J. Physical biology of the cancer cell
glycocalyx. *Nat. Phys.* **14**, 658–669 (2018).

65. Bergert, M. *et al.* Force transmission during adhesion-independent migration. *Nat. Cell*
Biol. **17**, 524–529 (2015).

66.   Paszek, M. J. *et al.* The cancer glycocalyx mechanically primes integrin-mediated growth
and survival. *Nature* **511**, 319–325 (2014).

67.   Barnes, J. M. *et al.* A tension-mediated glycocalyx–integrin feedback loop promotes
mesenchymal-like glioblastoma. *Nat. Cell Biol.* **20**, 1203–1214 (2018).

68.   Bezanson, J., Edelman, A., Karpinski, S. & Shah, V. B. Julia: A Fresh Approach to
Numerical Computing. *SIAM Rev.* **59**, 65–98 (2017).

69.   Hu, Y. & Smyth, G. K. ELDA: Extreme limiting dilution analysis for comparing depleted
and enriched populations in stem cell and other assays. *J. Immunol. Methods* **347**, 70–78
(2009).

70.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

71.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

72.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

73.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
microarray and RNA-Seq data. *BMC Bioinformatics* **14**, 7 (2013).

788
789

790
791
Figure 1: CSCs have increased invasive potential that is sensitive to metabolic challenge. a)
Schematic of the cancer stem cell reporter line GFP-NANOG MDA-MB-231 and sorting
strategy. **b)** Growth curve of sorted cells as measured by DNA amount, normalized to the first
measurement at t = 12 hrs (n = 3 samples). **c)** Schematic of the microfluidic device to analyze
cell invasion in response to a morphogen gradient generated by applying serum-containing
medium to the right channel only. **d)** Invasion distance of sorted GFP-NANOG MDA-MB-231
into collagen type I over 5 days (n = 4 fields of views, 1 device per condition). **e)** Invasion of
GFP-NANOG MDA-MB-231 cells into collagen type I treated with or without 2-DG for 5 days
(n = 3 devices per condition). **f)** Random migration of GFP-NANOG MDA-MB-231 treated with
or without 2-DG for 24 hours (n = 5 representative cells for migration plots, n = 20 cells per
condition for motility). Scale bar = 250 μm. * p< 0.05, ** p<0.01, *** p<0.001, **** p<0.0001
803

804
805
806
Figure 2: Cancer stem-like cells exhibit altered metabolism compared to non-stem-like
cancer cells. a) Glucose and lactate concentrations in media conditioned by sorted GFP-
NANOG MDA-MB-231 cells 48 hrs post sort as measured by a GlucCell glucose meter and a
lactate colorimetric assay (n = 3 samples). **b)** ECAR and OCR measurements of sorted GFP-
NANOG cells obtained during Seahorse-based glycolysis stress test (n > 6 samples). **c)** Fraction
of GFP positive (GFP+) GFP-NANOG MDA-MB-231 (n = 6) (i) and aldehyde dehydrogenase
high (ALDH+) (n = 3) (ii) parental MDA-MB-231 when cultured with 100 mg/dL glucose
(+glucose) or glucose-free media (-glucose) for 72 hours. GFP+ and ALDH+ cells were
determined by image analysis and Aldefluor assay, respectively. **d)** Fraction of GFP+ GFP-
NANOG MDA-MB-231 cells treated with or without 2-deoxyglucose (2-DG) (n = 3). **e)**
Immunofluorescence analysis of HA in the MDA-MB-231 NANOG reporter line. (n = 248 cells)
Scale bar = 50 μm. * p< 0.05, ** p<0.01, *** p<0.001, **** p<0.0001.
819
820

821
822
823
Figure 3: Increased HA production correlates with an increase in CSCs. a) Flow cytometry
analysis of cellular GFP levels categorized by their levels of cell surface-associated HA. **b)**
Heatmap representing changes of intracellular metabolites as measured by metabolomics of
sorted HA^{High} and HA^{Low} GFP-NANOG MDA-MB-231. **c)** Graphical representation of selected
metabolites in the central carbon metabolic pathway ratios between HA^{High} and HA^{Low} cells. **d)**
Schematic representing the theorized relationship between glycolysis, HA synthesis, and
stemness. **e)** Representative images of spheres formed through a limiting dilution assay of HAS2
and HAS3 overexpressing MDA-MB-231 and the corresponding sphere number. (n = 11) **f)**
Flow cyometry analysis of the percentage of the CD44^{+}/CD24^{-} population of parental or HA
overproducing MCF10A cells (n = 5). * p<0.05, ** p< 0.01, **** p<0.001. Scale bar = 200 μm.
834

835
836
837
Figure 4: Flux balance analysis predicts increased HA production in cancer stem-like cells
and ATP production. a) Workflow schematic. Sorted GFP-NANOG cells were cultured for 72
hrs before performing extracellular metabolomics using LC-MS. This data was used to constrain
a computational flux balance analysis model. **b)** Flux balance analysis model of the bioenergetic
pathway of sorted GFP-NANOG MDA-MB-231 derived from extracellular metabolomics
obtained over 72 hours of culture. Fluxes shown are GFP[High] relative to GFP[Null]. Grey
metabolites indicate relevant metabolite fluxes constrained based on extracellular metabolomic
profiles. Oxygen was constrained using values obtained from the Agilent Seahorse assay. For the
total FBA model constraints and results, see Supplementary File 1. **c)** Expanded HA synthesis
pathway flux balance analysis of GFP[High] relative to GFP[Null] (dashed box in B). Same legend as
in B. **d)** ATP production rate and **e)** rate index derived from measurements using the Agilent
Seahorse ATP real-time production rate assay kit (n $\geq$ 5). * p<0.05, ** p< 0.01.
850

Figure 5: Increased glycolytic metabolism necessary for HA production stimulates stemness and invasion of breast cancer cells. a) Extracellular acidification rate measurements of parental MCF10A or HA overproducing cells using the Agilent Seahorse XF Analyzer. **b)** Percentage of CD44$^+$/CD24$^-$ cells of parental MCF10A or HA overproducing cells with or without 20mM 2-DG as measured by flow cytometry. **c)** ATP production rate of either MCF10A or MDA-MB-231 cells overexpressing HAS2 or HAS3 pre-treated with or without 2-DG (20mM for MCF10A, 25mM for MDA-MB-231) as measured using the Real-Time ATP Rate Assay in the Agilent Seahorse XF Analyzer. (n ≥ 8) **d)** Invasion of the MCF10A and **e)** MDA-MB-231 HA overproducing cells. Representative immunofluorescence images and corresponding quantification of invasion after 5 days are shown (n = 3 devices per condition). Scale bar: 200 μm, * p< 0.05, ** p<0.01, *** p<0.001, **** p<0.0001

863
864
Figure 6: Gene signatures of HA overproducing cells correlate with worse patient prognosis regardless of HAS isoform. a) Principle Component Analysis plot of RNA sequencing results of the MCF10A or HA overproducing cells. **b**) Heatmap of z-score normalized gene expression of differentially expressed genes with hierarchical clustering of both samples and genes. **c**) Gene Set Enrichment Analysis of the Hallmarks gene signature database for genes upregulated in either HAS2 or HAS3 overexpressing cells contrasted against Parental MCF10A cells. **d**) Gene Set Enrichment Analysis of published HIF1α target genes for HAS2 or HAS3 overexpressing cells. All results are not significant. (FDR q-values ≥ 0.25) **e**) Venn diagram displaying the total number of significantly increased genes (FDR ≤ 0.05, log2(Fold Change) ≥ 1) in either HAS2 or HAS3 overexpressing cells. The top 4 most significant genes excluding HAS2/3 are shown in the table. **f**) Ranked enrichment analysis of a curated list of

migration and metabolism gene sets for tumors from the METABRIC cohort. The degree of
enrichment is determined as the $-\log_{10}$(p-value) multiplied by the sign of the difference in
enrichment scores (ES) of patients stratified into either low or high enrichment of the HA
overproduction gene signature. The dashed red line denotes the empirically determined cutoff for
non-random enrichment. **g)** METABRIC patient survival probability predicted by the
overlapping 72-gene signature of HA overexpressing cells. Patient signature scores were
stratified into quartiles, and a log-rank test was used to determine statistical significance.