Convergent approaches to delineate the metabolic regulation of tumor invasion by

hyaluronic acid biosynthesis

Adrian A. Shimpi[1*], Matthew L. Tan[1*], Michael Vilkhovoy[2], David Dai[2], L. Monet Roberts[1, 2],

Joe Kuo[2], Lingting Huang[2], Jeffrey D. Varner[2], Matthew Paszek[2,3], Claudia Fischbach[1,3,†]

[1]Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca,

New York, 14853, USA

[2]Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University,

Ithaca, New York, 14853, USA

[3]Kavli Institute at Cornell for Nanoscale Science, Cornell University, Ithaca, New York, 14853,

USA

*Contributed equally to this work

[†]Corresponding author:

Claudia Fischbach-Teschl

cf99@cornell.edu

The authors declare no potential conflicts of interest.

Abstract

Metastasis is the leading cause of breast cancer-related deaths and often driven by invasion and cancer-stem like cells (CSCs). Both the CSC phenotype and invasion have been associated with increased hyaluronic acid (HA) production. How these independent observations are connected, and which role metabolism plays in this process remains unclear due in part to the lack of convergent approaches that integrate engineered model systems, computational tools, and cancer biology. Using microfluidic invasion models, metabolomics, computational flux balance analysis (FBA), and bioinformatic analysis of patient data we investigated the functional links between the stem-like, invasive, and metabolic phenotype of breast cancer cells as a function of HA biosynthesis. Our results suggest that CSCs are more invasive than non-CSCs and that broad metabolic changes caused by overproduction of HA play a role in this process. Accordingly, overexpression of hyaluronic acid synthases (HAS) 2 or 3 induced a metabolic phenotype that promoted breast cancer cell stemness and invasion *in vitro* and upregulated a transcriptomic signature that was predictive of increased invasion and worse survival in patients. Collectively, this study suggests that HA overproduction leads to metabolic adaptations that help satisfy the energy demands necessary for 3D invasion of breast cancer stem cells further highlighting the importance of engineered model systems and multidisciplinary approaches in cancer research.

Introduction

Despite advancements in treatment options, breast cancer remains the second leading cause of cancer-related deaths in women[1]. Mortality in breast cancer is driven by metastasis and relapse, during which cancer cells in the primary tumor invade into surrounding tissues and disseminate into distant sites to form secondary tumors. The pathogenesis of metastasis can be attributed to intratumoral heterogeneity, where phenotypic diversity enables a subset of cells to become invasive and resistant to traditional therapies[2,3]. In particular, the emergence of cancer cells with stem-like properties (CSCs) contributes to metastasis because of their self-renewal and invasive properties[4]. CSCs are identified by their expression of stem cell markers (e.g. NANOG, SOX2, OCT4, aldehyde dehydrogenase [ALDH]) and influenced by features of the tumor microenvironment including extracellular matrix (ECM)[5,6]. For example, CSC invasion into the surrounding stroma is controlled by ECM microarchitecture and stiffness[7,8] including collagen fiber alignment at the tumor periphery[7,9,10]. However, the emergence and maintenance of the CSC phenotype and their consequences on 3D collagen invasion are poorly understood as studies often isolate tumor cells from the complex microenvironmental conditions that influence their behavior *in vivo*.

One key component of the tumor microenvironment influencing tumor cell phenotype and invasion is hyaluronic acid (HA). While much prior work has focused on how HA secreted by stromal cells regulates tumorigenesis[11–13], cancer cells themselves also produce HA. In fact, overproduction of HA by tumor cells enriches for a CSC phenotype[14,15] and correlates with worse patient prognosis[11,16]. Although several studies have delineated the specific signaling pathways by which HA modulates cell behavior[15,17], excess production of HA also regulates tumor cells via

biophysical mechanisms[18,19]. For example, biosynthesis and pericellular retention of HA as part of

the glycocalyx allows tumor cells to invade and extravasate more effectively[20,21]. However, which

metabolic adaptations tumor cells may use to increase HA-mediated tumor cell invasion and the

role cancer cell stemness plays in this process remains unclear.

Aberrant cellular metabolism is a hallmark of cancer that has been independently tied to increased

tumor cell invasion, stemness, and HA biosynthesis. Therefore, we speculated that metabolism

may serve as an overarching regulator of HA-mediated tumor cell invasion. Specifically, we

hypothesized that metabolic adaptations increase cancer cell stemness, which phenotypically can

more readily satisfy the energetic demands of 3D invasion. Prior findings suggest that CSCs

exhibit metabolic phenotypes that are distinct from non-CSCs[22–24] and that effective tumor cell

invasion through dense ECM requires metabolic adaptations including increased glucose uptake

and ATP generation[25,26]. While increased glucose uptake due to aerobic glycolysis is a common

feature of tumor cell metabolism, this typically results in the fermentation of glucose to lactate

(known as the Warburg Effect) which produces ATP less efficiently than oxidative

phosphorylation per glucose molecule. This energetically disadvantageous state must then

encourage tumor cells to develop compensatory mechanisms to generate the necessary energy for

3D invasion.

Because increased HA production of CSCs is known to direct intermediate products of glycolysis

into the hexosamine biosynthetic pathway (HBP)[17] it is possible that the resulting metabolic

rewiring enables their 3D invasion by providing alternative strategies for ATP production.

However, identifying broad metabolic changes beyond conventional biochemical methods

requires integrating computational approaches that can model the flow of metabolites through

relevant large-scale metabolic networks, while simulating a desired metabolic phenotype for

subsequent experimental validation. Here, we integrate engineered cell lines, microfabricated

culture models, and computational approaches including flux-balance-analysis (FBA) to

investigate the interconnectedness of the tumor cell phenotype, HA production, and metabolism

and its influence on cancer cell invasion. We demonstrate that metabolic adaptations associated

with increased HA production promote a stem-like phenotype in cancer cells, which can more

readily satisfy the energy demands necessary for 3D invasion. Bioinformatic analysis of clinical

data from publicly available datasets further indicated that these changes correlated with increased

invasive potential and worse survival in patients. Collectively, our results suggest that HA

overproduction due to metabolic reprogramming negatively influences prognosis in breast cancer

patients by altering invasion, and further motivate the need for utilizing a multidisciplinary toolset

to study intratumoral heterogeneity and its role in invasion.

Results

Cancer Stem-Like Cells Exhibit Increased Invasive Potential

To investigate differences in invasion between CSCs and their non-differentiated counterparts, we

utilized the CSC reporter cell line GFP-NANOG MDA-MB-231 in which green fluorescent

protein (GFP) expression is controlled by the NANOG promoter[27] (Fig. 1a). Using fluorescence-

activated cell sorting (FACS) GFP-NANOG MDA-MB-231 cells were sorted into GFP^{Null}, GFP^{Low}

(bottom 5% GFP expressing), and GFP^{High} (top 5% GFP expressing cells) populations to enrich

for different stem-like states (Fig. 1a) whose phenotype was maintained over 5 days of culture

(Supplementary Fig. 1). Moreover, the GFP^{High} cell population proliferated more slowly than the

non-CSC GFP[Null] population, consistent with a more quiescent phenotype (Fig. 1b). To study

potential differences in the invasive phenotype of these different cell populations, we monitored

tumor cell invasion in response to a morphogen gradient using a microfluidic collagen type I

hydrogel model (Fig. 1c). In this setup, GFP[High] NANOG MDA-MB-231 cells invaded into the 3D

fibrillar collagen hydrogel region (Fig. 1d) of the device more readily than their less stem-like

counterparts, validating that CSCs exhibit increased 3D invasive potential (Fig. 1e). Cell migration

and invasion are energetically intensive processes requiring increased metabolic consumption[28].

Accordingly, inhibiting energy production with the glycolysis inhibitor 2-deoxyglucose (2-DG)

reduced both collagen hydrogel invasion and random 2D migration of unsorted GFP-NANOG

MDA-MB-231 cells (Fig. 1f,g). These results indicate that CSCs exhibit increased invasive and

migratory potential in our experimental setup that was dependent on glycolytic metabolism.

Cancer Stem-Like Cells Have Altered Metabolism

To characterize the metabolic phenotype of cancer cells as a function of their stem-like phenotype,

extracellular metabolite production and consumption rates were measured in sorted GFP-NANOG

cells (Fig. 2a,b). The more stem-like cells consumed more glucose and produced more lactate

relative to their non-stem-like counterparts indicative of increased glycolysis (Fig. 2a). Real-time

metabolic analysis with the Agilent Seahorse (Seahorse) Analyzer of extracellular acidification

rate (ECAR) and oxygen consumption rate (OCR) further suggested that CSCs not only exhibited

increased glycolysis, but also oxidative phosphorylation (Fig. 2b, Supplementary Fig. 2).

Acccordingly, culturing in glucose-free media decreased the percentage of GFP[High] cells relative

to media containing physiological levels of glucose (Fig. 2c). This approach also decreased the

ALDH bright (ALDH[Br]) fraction (a marker for breast CSCs[29,30]) of wildtype MDA-MB-231 cells

confirming that our results were not an artefact of the GFP-NANOG reporter cell line. Moreover,

inhibition of glycolysis by 2-DG reduced the GFP^{High} population further corroborating that the

CSC phenotype is intimately linked to glucose metabolism (Fig. 2d).

As increased HA production has been associated with both metabolic reprogramming and the CSC

phenotype[17,31], we next assessed how HA production correlates with the CSC phenotype.

Immunofluorescence (IF) image analysis identified that GFP expression in GFP-NANOG MDA-

MB-231 cells positively correlated with the amount of cell surface-associated HA (Fig. 2e). These

differences were likely due to changes in HA biosynthesis and subsequent cell surface retention as

analysis of secreted HA by ELISA indicated low levels across all experimental conditions that

were not significantly different from each other although more stem-like cells seemed to secrete

slightly less HA (Supplementary Fig. 3a). Together, these results suggest that more stem-like cells

exhibit increased glucose metabolism relative to their less stem-like counterparts and that these

changes correlate with elevated levels of cell surface-associated HA.

Hyaluronic Acid Production Correlates with Stemness

To better quantify the relationship between glucose metabolism, cell surface-associated HA, and

stemness, GFP-NANOG MDA-MB-231 cells were stained for HA and then subjected to flow

cytometry and targeted metabolomics. Consistent with the IF results, the amount of cell surface-

associated HA directly correlated with GFP expression levels (Fig. 3a). Importantly, sorting the

GFP-NANOG MDA-MB-231 cells into high and low HA producing cells for targeted

metabolomics revealed broad changes of intracellular metabolites with hierarchical clustering

separating the two HA production phenotypes (Fig. 3b). In particular, both upper glycolytic and

tricarbocylic acid (TCA) cycle metabolites such as glucose-6-phosphate (G6P), fructose-6-

phosphate (F6P), oxaloacetate (OAA), and citrate (CIT) were increased in the highly HA

producing cells (Fig. 3c). Combining these results with the broad metabolic alterations of CSCs

described above (Fig. 2), these data suggest a direct link between glycolysis-dependent HA

synthesis and the CSC phenotype (Fig. 3d). To more directly test how HA synthesis affects

stemness, GFP-NANOG MDA-MB-231 cells were cultured with the HA synthesis inhibitor 4-

methylumbelliferone (4-MU, 0.5 mM). These results trended towards a decreased fraction of stem-

like GFP[High] cells consistent with previous studies on stemness and invasion[15,20] (Supplementary

Fig. 3b).

To more directly determine if increasing production of HA promotes stemness, HAS2 and HAS3

were stably overexpressed in MDA-MB-231 and the non-malignant breast epithelial cell line

MCF10A. While HA is synthesized by three HAS isoforms (HAS1, 2, 3), HAS 1 varies in HA

synthesis and secretion rate from HAS 2 and 3. In contrast, HAS2 and HAS3 exhibit similar

sensitivities and responses to precursor availability and thus were used here[32–34]. Overexpression

of HAS2/3 increased the defining characteristic of CSCs in both cell lines. More specifically,

overexpression of HAS2/3 increased MDA-MB231 self-renewal as measured through sphere

formation in a limited dilution assay (Fig. 3e, Supplementary Fig. 4a,b) and increased the fraction

of CD44[+]/CD24[-] MCF10A cells, which characterizes an invasive breast CSC population with more

mesenchymal characteristics[35,36] (Fig. 3f, Supplementary Fig. 4d). This population could not be

assessed in HAS overexpressing MDA-MB-231 as this cell line contains an intrinsically high

fraction of CD44+/CD24- cells (>85%), making changes difficult to quantify[37,38] (Supplementary

Fig. 4c). Finally, HAS3 overexpression also trended towards an increase in the ALDH[Br] fraction

of MCF10A (Supplementary Fig. 4e). Together, these results suggest that HA overproduction

increases stem-like cell properties in breast cancer cells regardless of cell line and HAS2/3 isoform.

Increased HA Production by CSCs is Associated with Increased Glucose Conversion and ATP

Production

Given our results that metabolic reprogramming of CSCs supports their energy demands during

invasion (Fig. 2) and that a more stem-like phenotype is associated with increased HA production

(Fig. 3), we speculated that increased HA production promotes a more energetic CSC phenotype.

Probing the individual contribution of HA biosynthetic pathways to the metabolic state of CSCs

solely by measuring different metabolite levels, however, is challenging given the

interconnectedness of most metabolic pathways. To circumvent these limitations and delineate the

contribution of HA biosynthesis to CSC metabolism more directly, a flux balance analysis (FBA)

model was constructed. FBA is a widely utilized mathematical approach to model the flow of

metabolites (flux) through a genome-scale network of metabolic pathways, including glycolysis,

oxidative phosphorylation, hexosamine biosynthesis, and amino acid consumption[39], and has been

successfully used to investigate cancer metabolism[40,41]. In contrast to traditional metabolomics,

FBA also enables the simulation of a desired metabolic phenotype by adjusting model parameters

such as the objective function and flux constraints. To develop the model for our study,

extracellular metabolomics of GFP[High] and GFP[Null] cells were performed over 72 hours to define

metabolite consumption/production profiles used to constrain flux values for more and less stem-

like MDA-MB-231 breast cancer cells, respectively (Fig. 4a). Additionally, the objective function

of the FBA model was set to maximize HA synthesis to study both the capacity of CSCs and non-

CSCs to produce HA and the associated changes in metabolic flux. Results from the FBA model

indicated that the more stem-like cells (GFP^{High}) increased flux through the upper stages of

glycolysis (i.e., conversion of glucose to fructose-6-phosphate), ATP generation, lactate

production/secretion, and glutamine uptake relative to the non-stem-like cells (GFP^{Null}) (Fig. 4b,

Supplementary File 1). Additionally, the FBA model indicated increased flux through all HA

synthesis intermediate steps for the GFP^{high} cells relative to the GFP^{null} cells (Fig. 4c). Collectively,

these data suggest that the increased capacity of CSCs to synthesize HA is related to increased

glucose conversion but simultaneously allows these cells to produce ATP more efficiently. To

confirm the predictive value of the FBA model, a Seahorse real-time ATP rate assay was

performed. This analysis verified that the mitochondrial and overall ATP production rate was

increased in GFP^{High} versus GFP^{Null} cells (Fig. 4 d,e). Together, these findings suggest that the

metabolic phenotype of more stem-like cancer cells leading to increased HA biosynthesis

promotes ATP production by these cells. FBA models are suitable for predicting these changes.

Metabolic Changes resulting from Increased HA Production Stimulate a Stem-like Breast Cancer

Cell Phenotype to Promote Invasion.

Our aforementioned results suggest functional links between the stem-like and invasive phenotype

of tumor cells (Fig. 1), CSCs and metabolism (Fig. 2), CSCs and HA (Fig. 3), and HA and

metabolism (Fig. 4). However, whether these single observations are mechanistically connected

remained to be determined. Therefore, we next hypothesized that the metabolic phenotype induced

by HA overproduction leads to changes in energy production that increase the stem-like phenotype

of breast cancer cells to promote invasion. Indeed, Seahorse analysis revealed that MCF10A and

MDA-MB-231 cells overexpressing HAS2 and HAS3 had increased ECAR compared to their

parental controls suggesting an increase in glycolytic energy production (Fig. 5a). Interestingly,

OCR was unchanged in HAS2/3-overexpressing MCF10A but increased in HAS2/3-

overexpressing MDA-MB-231 cells (Supplementary Fig. 5a), consistent with an increase in OCR

in GFP^{High} vs. GFP^{Low} and GFP^{Null} MDA-MB-231 (Fig. 2b). Treatment with 2-DG decreased both

cell surface-associated and secreted HA in MCF10A cells (Supplementary Fig. 5b, c). Notably, 2-

DG treatment of MCF10A cells decreased the CD44$^+$/CD24$^-$ fraction in the HA overproducing

cells but had no effect on the parental control cells suggesting that the enhanced glycolysis

associated with HA overproduction is critical to maintaining stemness (Fig. 5b). Furthermore, HA

overproduction increased both total and glycolytic ATP production (Fig. 5c, Supplementary Fig.

5d) that 2-DG reversed to similar levels as in MCF10A control cells (Fig. 5c). A similar trend was

noted for MDA-MB-231 cells, although 2-DG decreased glycolytic ATP production only in the

HAS3 overexpressing cells. Overexpression of HAS2/3 also increased invasion of both MCF10A

and MDA-MB-231 (Fig. 5d, e), consistent with their increased stem-like phenotype (Fig. 3d, e).

This effect was inhibited by 2-DG treatment, implicating a functional consequence of HA-

mediated stemness and metabolism in tumor cell invasion (Fig. 5d, e). As 2-DG treatment did not

affect the ATP production rate in parental cells (Fig. 5c), these changes in invasion can be

attributed to differences in metabolism rather than compromised cell viability. Together this data

suggests that HA overproduction induces a glycolytic phenotype that is crucial for CSC-mediated

invasion.

HA Overproduction Transcriptome Changes Predict Worse Patient Survival

As our data implied that HAS2 and 3 overexpression induced more invasive tumor cell phenotypes

we next tested if and how these findings may correlate with differences in clinical prognosis. To

this end, RNA sequencing was conducted on the MCF10A cells overexpressing HAS2 and HAS3

and their parental control to identify transcriptional changes that would allow us to interrogate the

contribution of HA overproduction to stemness, invasion, and patient outcomes. Principle

component analysis (PCA) and hierarchical clustering indicated that the transcriptome of HA

overproducing cells differed significantly from their parental control (Fig. 6a, b, Supplementary

Fig. 6a, Supplementary File 2). Gene set enrichment analysis (GSEA) with the Hallmarks gene

sets from the Molecular Signature Database revealed enrichment of a wide range of pathways for

both HAS2 and HAS3-overexpressing cells, including those previously associated with stemness

and invasion such as NF-κB[42], hypoxia[8], PI3K signaling[43], IL6-STAT3 signaling, and reactive

oxygen species[22,23] (Fig 6c). Since HIF1α signaling has been previously implicated in an HA-

dependent increase in the CSC phenotype[17], we performed GSEA for HIF1α target genes[44–48].

Interestingly, neither HAS2 nor HAS3 transcriptomes were enriched for HIF1α target genes (Fig.

6d), which have previously been suggested as drivers of HA-dependent stemness[17]. These results

further support our findings that the cellular phenotypes investigated in this study are due to

broader metabolic and energetic changes and cannot be attributed to glycolysis-driven changes in

hypoxia-related signaling.

To determine the clinical relevance of our findings, a 72-gene signature was generated from the

genes upregulated in both HAS2 and HAS3-overexpressing cells (log_2-fold change > 1, p-value <

0.05) compared to their parental control (Fig. 6e, Supplementary Table 1). Overrepresentation

analysis of this gene signature revealed no significant enrichment of genes associated with most

metabolic pathways from the KEGG database (Supplementary Fig. 6b). While surprising these

results can be explained by the fact that metabolism is significantly regulated by posttranscriptional

changes[49,50]. Subsequently, we calculated enrichment scores for this HA overproduction gene

signature and a curated list of gene signatures involved in migration, cytoskeleton dynamics, and

metabolism (Supplementary File 3, Supplementary Fig. 7a) using single sample GSEA

(ssGSEA)[51] for chemotherapy-naïve patients in the METABRIC cohort. Tumorous tissue from

patients enriched for our HA overproduction signature were simultaneously enriched for several

migration and actin cytoskeleton gene sets after accounting for random associations suggesting

increased tumor invasion in these patients. Metabolic gene sets except for glycosaminoglycan

synthesis were not enriched in patients consistent with our findings that metabolic pathways were

not overrepresented in our HA production signature nor enriched by both HAS2/3 overexpression

in MCF10A cells (Fig. 6f, Supplementary Fig. 6b, Supplementary Fig. 7b,c). Importantly, our HA

overproduction signature also predicted worse patient survival consistent with their increased

expression of invasion-related gene sets (Fig. 6f,g). Gene signatures specific to upregulation of

either HAS2 or HAS3 also predicted worse patient survival that was not seen in the parental control

(Supplementary Fig. 6c). Collectively, our results suggest that genes associated with increased HA

biosynthesis correlate with an enrichment of migratory genes and predict worse patient survival

regardless of HAS2/3 isoform and transcriptional regulation of metabolic gene sets.

Discussion

Although CSCs have been associated with HA production and altered metabolism, the exact nature

of these connections to tumor heterogeneity and consequences on tumor cell invasion remain

unclear. Because of the breadth of expertise required to probe each of these aspects, it is infeasible

to rely on a single model system or analytical technique to perform a comprehensive investigation

of these connections. Furthermore, the systems used must be compatible to enable a holistic

evaluation of these aspects. Here, we have used a suite of multidisciplinary approaches that include

engineered cell lines, *in vitro* 3D cell culture models, computational metabolic modeling, and

genomic tools to uncover the relationship between tumor cell phenotype, HA production,

metabolic reprogramming, and 3D invasion. We demonstrate that CSCs can increase glycolytic

and oxidative metabolism simultaneously and that the resulting changes in HA production support

3D invasion by supporting more efficient ATP production. Moreover, we identified that these

cellular changes correlated with a gene expression signature that was predictive of patient survival.

Our data suggests that cell states associated with increased glycolysis enrich for highly invasive,

HA-producing CSCs. Interestingly, HA produced by CSCs is primarily retained on the cell surface

rather than excreted into the surrounding environment (Fig. 3, Supplementary Fig. 3).

Consequentially, the retention of HA on the cell surface contributes to glycocalyx thickness whose

biophysical properties impact the interactions between cells and their surrounding environment

including cell-extracellular matrix interactions necessary for migration[18,52]. In particular, HA-

dependent changes of the glycocalyx can promote adhesion-independent or ameboid migration by

altering the friction required for force generation against extracellular structures[53], which may

further decrease the energy needed for migration. Alternatively, changes in glycocalyx thickness

impact surface receptor diffusion patterns and accessibility to impact both adhesion-mediated and

receptor tyrosine kinase-mediated signaling[54–56]. Further studies to delineate the contributing

biophysical properties are needed to determine the influence of HA on predominant migration

mode.

Throughout the metastatic cascade, HA has been implicated in promoting invasive phenotypes and

to support survival of circulating cells in the vasculature[19–21]. While HA-dependent changes of

cancer malignancy and stemness have been primarily attributed to upregulation of HAS2[15,31], our

findings indicate that HAS3 similarly promotes a stem-like state (Fig. 3). Indeed, our results that

2-DG inhibited the stem-like phenotype in both HAS2/3-overexpressing mammary epithelial cells

suggests that the metabolic reprogramming enacted by HA overproduction may be a central

regulator of the tumor stem-like phenotype (Fig. 5b). Interestingly, exogenous degradation of HA

produced by cancer cells promotes glucose uptake that can further promote migration[56]. This

mechanism possibly provides a positive feedback mechanism by which increased HA production

regardless of HAS isoform stimulates 3D invasion. Further studies are necessary to decouple the

metabolic programming associated with HA production and degradation on the CSC phenotype.

To interrogate alterations in other metabolic pathways associated with glycolysis and oxidative

phosphorylation, we utilized FBA to predict changes in metabolic fluxes induced by stemness.

FBA is especially proficient in enabling the study of cancer metabolism as it simulates metabolic

phenotypes based on real-world constraints such as cell growth rate and glucose uptake[39,57]. The

FBA model developed here indicated that stem-like cells exhibited an increased capacity for HA

production, and that this increase in HA production was associated with broad metabolic

alterations. Interestingly, our model suggested that although the conversion of glucose to pyruvate

is decreased in stem-like cells when maximizing HA production, the production of both lactate

and acetyl-CoA from pyruvate is increased. An alternative source of pyruvate are the malic

enzymes, which convert malate to pyruvate while producing the reducing agent NADPH[58]. Indeed,

our FBA model indicated an increase in the flux through malic enzyme 2 (ME2) (Supplementary

File 1), which has been previously associated with the loss of cellular senescence and increased

tumorigenesis[59,60]. Although any connection with ME2 will need to be experimentally verified in

future experiments, the FBA model developed here was able to provide additional insights into the

metabolic alterations associated with stemness-related HA production.

The energetic demands associated with HA-mediated invasion can induce broad metabolic

changes. For example, increased HA production rapidly depletes UDP-sugar substrates, which

cancer cells may compensate for by increasing glycolysis to maintain flux into the HBP and thus,

the pool of UDP-sugars[17]. Our FBA model corroborates this phenomenon and provides further

insight into metabolic states associated with CSCs such as an increased HA production capacity.

The resulting metabolic phenotypes associated with overexpressing HAS2 or HAS3 in cells

suggest that the glycolytic demand dominates over possible changes in oxidative metabolism

previously associated with stemness[23,61,62]. Furthermore, our finding that HA overproduction

induces phenotypic and transcriptomic changes that correlate with invasion may help explain why

glycolytic, mesenchymal breast CSCs localize to the leading edge of tumors and worsen patient

prognosis[23,36]. Together, these results suggest that HA production enables a more invasive,

malignant CSC metabolic phenotype.

Transcriptomic analysis of HAS2 and HAS3 overexpressing MCF10As indicated an enrichment

of multiple stemness-associated gene sets (e.g. IL6-JAK-STAT3 signaling and hypoxia), but

metabolic genes were not differentially expressed. This discrepancy with our experimental

observations may be explained by how metabolic changes are not only regulated transcriptionally

but also by enzyme activity levels, localization, and substrate availability[49,50,63,64]. Furthermore,

cytoskeletal rearrangement, which is critical for invasion, can independently control glycolytic

flux by mediating enzyme degradation[65] or sequestration[50]. Our results suggest that the glycolytic

CSC phenotype associated with HA overproduction is not regulated transcriptionally, but whether

this is enacted by cytoskeletal dynamics or changes in the relative activity of different metabolic

enzymes requires further investigation.

Acknowledgements

Thank you to all members of the Fischbach lab for valuable feedback and discussion of the

research. The authors would also thank Dr. Andrea Di Micheli, David McKellar, and Dr. Ben

Cosgrove for assistance with RNA-sequencing data processing. The work described was supported

by National Cancer Institute through the Center on the Physics of Cancer Metabolism

(1U54CA210184), the National Science Foundation (NSF) through Graduate Research Fellowship

Program awarded to A.A.S and L.M.R. (DGE-1650441), the U.S. Department of Education

through the Graduate Assistance in Areas of National Need (GAANN) Fellowship awarded to

M.L.T., the Breast Cancer Coalition of Rochester, and the Cornell NanoScale Science &

Technology Facility (CNF), a member of the National Nanotechnology Coordinated Infrastructure

(NNCI) supported by the National Science Foundation (Grant NNCI-2025233). This work utilized

facilities at the Cornell Institute of Biotechnology including the Cornell University Biotechnology

Resource Center (BRC) Imaging Facility (RRID:SCR_021741) for image acquisition using the

Zeiss LSM 710 Confocal Microscope (NIH 1S10RR025502), the BRC Genomics Facility

(RRID:SCR_021727)   for   sequencing,   and   the   BRC   Flow   Cytometry   Facility

(RRID:SCR_021740) for cell sorting with BD Biosciences FACSAria Fusion. Mass spectrometric

analysis of targeted metabolomics was performed at the Weill Cornell Medicine Proteomics and

Metabolomics Core Facility.

Author Contributions

A.A.S., M.L.T., and C.F. designed the study. A.A.S. and M.L.T. conducted most of the

experiments. M.V. and D.D. conducted extracellular metabolomics for FBA analysis. L.M.R. and

M.P. performed FACS for HA production. J.K., L.H., and M.P. generated plasmids for HAS

overexpression. L.M.R. and A.A.S. generated cell lines. J.V. and M.L.T. performed FBA analysis.

A.A.S., M.L.T., and C.F. analyzed the data and wrote the manuscript. All authors discussed the

results and commented on the manuscript.

Materials and Methods

Cell Culture

MCF10A (ATCC) cells were cultured in DMEM/F12 supplemented with 5% Horse Serum, 1%

penicillin/streptomycin, 10 µg/mL Insulin, 0.5 µg/mL Hydrocortisone, 100 ng/mL Cholera toxin,

and 20 ng/mL human epidermal growth factor (EGF). MDA-MB-231 (ATCC) and the GFP-

NANOG MDA-MB-231 (a kind gift from Dr. Ofer Reizes[27]) cells were cultured in DMEM

supplemented with 10% FBS and 1% penicillin/streptomycin. HAS overexpressing cells were

cultured in their respective medias containing 1 µg/mL doxycycline (Santa Cruz Biotechnology).

For glycolytic inhibition studies, cells were treated with media containing 2-deoxyglucose

(MilliporeSigma) matching the glucose concentration in the media (25mM or 20mM for DMEM,

DMEM/F12 respectively) for 24 hours unless otherwise noted.

Generated Cell Lines

cDNAs for human HAS2 and HAS3 were generated and cloned into the lentiviral vector pLV

HygroR tetOn to create stably transduced, tetracycline-inducible MCF10A cell lines as previously

described[52]. cDNAs for human HAS2 and HAS3 were also fabricated and inserted into a

tetracycline-inducible PiggyBac expression vector to generate pPB huHAS2-IRES2-mScarlet-

IRES2-NeoR and pPB huHAS3-IRES2-mScarlet-IRES2-NeoR through custom gene synthesis

(Twist Biosience). Generation of the MDA-MB-231 HAS overexpressing cells was conducted

using either the pPB huHAS2-IRES2-mScarlet-IRES2-NeoR or pPB huHAS3-IRES2-mScarlet-

IRES2-NeoR or with the Piggybac transponase using the Nucleofector Cell Line Kit V (Lonza).

Stably transfected cells were selected using 1 µg/mL puromycin (MilliporeSigma) or 800 µg/mL

G418 (ThermoFisher). MDA-MB-231 HAS overexpressing cells were then sorted post-selection

on mScarlet expression levels.

FACS and Flow Cytometry

GFP-NANOG MDA-MB-231 cells were trypsinized and resuspended at 10×10^6 cells/mL in FACS

buffer (2.5% FBS/PBS, 2mM EDTA) and processed on the BD FACSAria Fusion Cell Sorter. The

top 5% and bottom 5% of the GFP+ fraction were designated as GFP^{High} and GFP^{Low} respectively,

while the non-GFP expressing population were designated as GFP^{Null}. Cells were sorted into cell

culture media, recounted, seeded, and allowed to rest for 24 hours before use in experiments.

Cell sorting for targeted metabolomics was conducted on GFP-NANOG MDA-MB-231 cells

stained for HA using Alexfluor-568 (ThermoFisher) conjugated HA binding protein (HABP, 40

µg/mL, MilliporeSigma). The high HA (top 5%) or low HA (bottom 5%) fractions were sorted

using a BD FACSAria Fusion Cell Sorter.

ALDH activity was determined for MDA-MB-231 and MCF10A cells using the Aldefluor™

Assay (STEMCELL Technologies) according to manufacturer instructions with incubation

conducted for 30min at 37°C. Analysis was conducted on a BD Accuri C6 Plus analyzer.

The fraction of CD44+/CD24- cells was determined by trypsinizing and resuspending cells in

FACS buffer at $10x10^6$ cells/mL followed by incubation with antibodies against human CD44

(APC-conjugated, Clone G44-26, 1:5, BD Biosciences) and CD24 (PE-Cy7-conjugated, Clone

ML5, 1:20, BD Biosciences). Gates were determined using the isotype controls mouse anti-IgG2b

κ (APC-conjugated, Clone 27-35, 1:5, BD Biosciences) and mouse anti-IgG2a κ (PE-Cy7-

conjugated, Clone G155-178, 1:20, BD Biosciences). Cells were analyzed on a BD Accuri C6 Plus

Analyzer.

Invasion and Migration Assays

To prepare microfluidic invasion assays, rat tail Type I collagen (Corning) was neutralized with 1

N NaOH and diluted with 1X DMEM to a final concentration of 2.5 mg/mL. Before neutralization

and dilution, 10x DMEM was added to collagen as a pH indicator to a final concentration of 10%

v/v. Microfluidic 3D cell culture devices (AIM Biotech) were then injected with the 2.5 mg/mL

rat tail Type I collagen solution into the center channel of the chip. The chips were then incubated

at 4°C for 30 minutes, followed by incubation at 37°C for 30 minutes to complete collagen

polymerization. For MDA-MB-231 invasion, DMEM containing 1% FBS and 1%

penicillin/streptomycin was injected into the left flanking media channel, and DMEM containing

10% FBS and 1% penicillin/streptomycin was injected into the right flanking media channel. For

MCF10A invasion, MCF10A media devoid of Horse Serum and EGF was injected into the left

flanking media channel, and fully supplemented MCF10A media was injected into the right

flanking media channel. 10 μL of a $2.5x10^5$ cell suspension of GFP-NANOG MDA-MB-231 or

MCF10A cells were then injected into both ports of the left channel (20 μL total). Cells were

allowed to invade through the hydrogel channel for 3 - 5 days before fixation in a 4%

paraformaldehyde solution. Media was exchanged every 24 hours. After fixation, chips were

stained, imaged, and individual cell invasion distance was measured using ImageJ by segmenting

nuclei.

To assess random migration ability, GFP-NANOG MDA-MB-231 cells were plated on

fibronectin-coated (30 µg/mL) glass 96 well plates and placed in an Incucyte S3 (Sartorius) live

cell imaging system. Images were obtained in 20-minute intervals over 24 hours. Individual cell

tracking was performed using ImageJ to determine migration velocity (motility) and random

migration paths.

Immunofluorescence

GFP-NANOG MDA-MB-231 cells were plated on fibronectin-coated (30 µg/mL) glass coverslips

at 2500 cells/cm^2. Cells were then fixed in 4% paraformaldehyde (PFA)/PBS (w/v) for 20 min at

room temperature, blocked with 1% bovine serum albumin (BSA)/PBS (w/v) for 1 hr at room

temperature, and incubated overnight with AlexaFluor-568 conjugated hyaluronic acid binding

protein (HABP, 13.3 µg/mL, MilliporeSigma) at 4°C. Afterwards cells were incubated with DAPI

(2.5 µg/mL, ThermoFisher) for 30 min at room temperature.

For invasion experiments, devices were fixed in 4% paraformaldehyde (PFA)/PBS for 30 min at

room temperature, permeabilized with 0.1% Triton X-100/PBS (v/v) for 15 min at room

temperature, blocked with 1% bovine serum albumin (BSA)/PBS (w/v) for 1 hr at room

temperature, and incubated with DAPI (2.5 µg/mL) and either AlexaFluor-568 (ThermoFisher) or

AlexaFluor-647 (ThermoFisher) phalloidin to visualize F-actin.

Confocal Microscopy and Image Analysis

Images were acquired on a Zeiss LSM 710 confocal microscope with either a LD LCI Plan-

Apochromat 25×/0.8 Imm Korr DIC M27 or C-Apochromat W M27 10x/0.45 objective. Images

were analyzed using ImageJ with custom scripts. Briefly, for single cell intensity measurements,

z-stacks were sum projected and cells were segmented based on HA intensity. Cell clusters were

manually corrected to individual cells, while overlapping or edge-located cells were excluded from

analysis.

For invasion experiments, the C-Apochromat W M27 10x/0.45 objective was used at 0.6x zoom,

and a z-stack was obtained along a 2 mm length in the center of the device. Z-stacks were

maximum intensity projected, and invasion was assessed based on nuclei displacement along the

x-axis (across the hydrogel region). Invasion distance was normalized to the appropriate control

in each set of replicates per experiment, and data was pooled together across device replicates to

obtain averages per condition.

Metabolic Analysis

To obtain an initial measure of glycolytic ability, sorted GFP-NANOG MDA-MB-231 cells were

seeded in 24 well plates at 10,000 cells/cm^2 in 1 mL of media. Glucose concentration was measured

using a GlucCell Glucose Monitoring System (CESCO Bioengineering), and lactate concentration

was obtained using a colormetric lactate assay kit (Sigma).

Targeted metabolomics was conducted on GFP-NANOG MDA-MB-231 cells 48 hours post

sorting for HA production. Media was collected and non-adherent cells were pelleted at 500 x g

for 4 minutes. Meanwhile, 0.5 mL of 80% methanol (MetOH) was added onto adherent cells and

incubated at -80°C, After aspirating supernatant, the cell pellet was resuspended in 0.5 mL of 80%

MetOH and added to the respective well of adherent cells in plate and incubated at -80°C for 15

minutes. Cells were then scraped using a cell scraper and collected into an Eppendorf tube and

pelleted at 20,000 x g for 10 minutes at 4°C. Supernatant was then transferred to 2 mL screw-cap

vial and dried overnight under vaccum at room temperature. Following overnight drying, samples

were then dried for 2.5 hours in SpeedVac SPD 1030 at room temperature and then stored at -80°C

until analyzed by Weill Cornell Medicine Proteomics and Metabolomics Core Facility.

Real-time changes in metabolism were tested using a Seahorse XFe96 Analyzer in conjunction

with the Seahorse XF Glycolysis Stress Test Assay Kit and the Seahorse XF Real-time ATP Assay

Rate Kit (Agilent). Manufacturer instructions were followed to perform each of the assays.

Wildtype and modified MDA-MB-231 cells were seeded on a Seahorse XFe96 Cell Culture

microplate at 20,000 cells per well in standard media and allowed to attach overnight. For

experiments with MCF10A, cells were seeded at 20,000 cells per well. HAS overexpressing cells

had media changed 24 hours after seeding to include 1 µg/mL doxycycline. Media was then

changed to the specific Seahorse assay media, and plates were prepared according to manufacturer

instructions for each assay kit. Relevant metabolic values from each assay were calculated using

template worksheets provided by Agilent. After the assays were complete, DNA was extracted

from each well using Caron's Buffer (25 mM Tris-HCl, 0.4 M NaCl, 0.5% (w/v) sodium

dodecylsulfate), and total DNA content measured using the fluorometric Quantifluor dsDNA

Assay (VWR) and converted to cell number for normalization.

Flux Balance Analysis (FBA)

To generate a computational model of metabolic fluxes in sorted GFP-NANOG MDA-MB-231

cells, cells were sorted as described above and seeded into 24 well plates at 10,000 cells/cm^2. Cells

were allowed to attach overnight before a fresh media change. Media was collected 72 hours after

the initial media change and stored at -80°C before performing extracellular metabolomics to

measure levels of glucose, lactate, and the 20 amino acids. Glucose was measured using Contour

next EZ Blood Glucose Monitoring System (Ascensia Diabetes Care) using 5 μL as the sample

volume. Lactate and amino acid concentrations were assayed using an Acquity UPLC H-Class

System equipped with QDa and tunable UV (TUV) detectors controlled by Empower 3 software

(Waters Corporation). Specifically, extracellular amino acids were analyzed using a Waters AccQ-

Tag Ultra Derivatization Kit (Waters) according to the manufacturer's recommendations.

Derivatized samples were injected onto an AccQ-Tag Ultra C18 column (1.7 μm, 2.1 mm x 100

mm, Waters) and detected by an Acquity TUV detector (Waters) at 260 nm. Amino acids were

identified by known retention times of standards and concentrations were determined by

comparison with calibration standard curves. For the lactate measurements, samples were first

deproteinized by treatment with an equal volume of trichloroacetic acid followed by centrifugation

at 12,000 x g for 10 minutes. 200 μL of the supernatant was then combined with 600 μL of

acetonitrile (ACN) before injecting 2 μL into the LC-MS system. Lactate was quantified using a

standard curve ranging from 0.05mM to 1mM. Separations were performed on an Acquity UPLC

BEH Amide Column (1.7 μm, 2.1 mm x 100 mm, Waters). Solvent A consisted of 50:50

ACN:Water and solvent B consisted of 95:5 ACN:Water. The solvent gradient started at 0.01%

solvent A and 99.9% solvent B, raised to 40% A in 0.5 minutes, further raised to 70% A in 1.5

minutes, and returned to initial conditions over 0.1 minute and held for 3 minutes to re-equilibrate

the column. The flow rate was set to 0.6 mL/min, the autosampler was set to 5°C, and the column

was set to 50°C. The mass-to-charge ratio (m/z) of lactate was 88.9. Analysis was performed in

negative ion mode with a cone voltage of 15V and probe temperature of 600°C.

Extracellular metabolomics data of all 20 amino acids, lactate, and glucose was then used to

constrain a flux balance analysis by imposing bounds on allowable fluxes. Growth rate and O2

fluxes were constrained from cell counts and Seahorse data respectively. The model used in this

study was implemented in the Julia programming language[66], where the linear programming

problem was solved using the GNU Linear Programming Kit (GLPK) package

(https://www.gnu.org/software/glpk/). The stoichiometric matrix and metabolic growth

requirements were derived from a previously developed Core Cancer model[41]. The objective

function was set to maximize HA production subject to experimentally estimated rates of uptake,

secretion, and cell growth to define the phenotype of interest; 1000 simulations were performed

for each condition. A subset of the FBA results representing the energetic pathways of interest was

incorporated into a graphical representation (Fig. 4). The full FBA results can be found in

Supplementary File 1.

Limited Dilution Assay

Cells were serially diluted and seeded into ultra-low attachment 96-well plates (Corning) in 200

μL of serum-free DMEM/F-12 containing 2% B27, 10 ng/mL basic fibroblast growth factor, 20

ng/mL epidermal growth factor, 10 μg/mL insulin, and 1 μg/mL doxycycline. Sphere formation

was assessed after two weeks of culture, and the number of spheres was counted using a phase-

contrast microscope. The stem cell frequency was determined using the extreme limited dilution

algorithm[67].

RNA-Sequencing

Parental (NeoR-rtTA) and HA overproducing MCF10A cells were seeded on 10-cm dishes at 5000

cells/cm^2 and allowed to adhere overnight. Media was refreshed to include 1 µg/mL of doxycycline

to induce HAS2 and HAS3 expression and cultured for an additional 48 hours. RNA was isolated

using the Qiagen RNeasy kit according to manufacturer instructions. RNA libraries were prepared

using the Illumina TruSeq RNA Kit, and single-ended 75bp read lengths were sequenced on the

Illumina NextSeq 500 system.

Sequence Alignment and Gene Set Enrichment Analysis

Reads   were   trimmed   using   TrimGalore   version   0.4.4

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and aligned to the human

reference genome GRCh38 (ENSEMBL) using STAR version 2.6.0a[68]. Reads of genomic features

were counted using featureCounts[69], and differential gene expression was determined using

DESeq2[70]. Differentially expressed genes in HA overproducing cells were defined as a log-2 fold

change greater than 1 and an adjusted p-value less than 0.0001 compared to the parental cells.

Genes common to both HAS2 and HAS3 were combined to generate the HA overproducing gene

signature for survival analysis. Individual HAS2 and HAS3 gene signatures were defined as the

differentially expressed genes with log 2-fold change greater than 2 or 5, respectively, and a p-

value less than 0.0001.

Gene set enrichment analysis (GSEA) was conducted with a ranked list generated by taking the

sign of the fold change multiplied by the log-10 of the adjusted p-value. The list was inputted to

the   GSEA   Java   applet   (http://software.broadinstitute.org/gsea/index.jsp)   using   the

GSEAPreRanked tool and the Hallmarks gene sets from MSigDB v.7.0. Gene sets were considered

significantly enriched with a p-value and FDR value ≤ 0.05.

Patient Survival and Enrichment Analysis

Patient data from the METABRIC cohort was extracted from the Cancer Genomics Data Server. Patients were limited to those having not received chemotherapy. Gene signature scores were calculated using the single-sample GSEA (ssGSEA)[71] with the GSVA package[72]. The top and bottom quartiles of the ssGSEA scores for the 72-gene HA overproduction signature were designated as high and low scores, respectively, for survival and enrichment analysis. Kaplan-Meier survival analysis was conducted with the *survival* package in R using a Cox proportional hazard model with statistical significance determined using a log-rank test. Stratified patients were further analyzed for enrichment of curated lists of migration, cytoskeletal, and metabolic gene signatures obtained from the MSigDB v7.0. To correct for random associations, 300 randomly selected gene signatures with the same number of genes (15-500 genes) as gene sets in the curated list had ssGSEA enrichment scores additionally calculated. An empricial cumulative distribution function was established using the *ecdf* function in R and a p-value cutoff was determined where 95% of values fell below.

Statistical analysis

All experiments were performed with at least three independent biological replicates unless otherwise noted. Pairwise comparisons were conducted using a Mann-Whitney U test unless otherwise noted. Multiple comparisons were evaluated with either a Kruskal-Wallis Test or two-way ANOVA with Dunn's post hoc analysis. Results were considered statistically significant with a p-value less than 0.05. Unless otherwise noted, all data points are plotted mean +/- the standard deviation. All statistically analysis was performed using GraphPad Prism v9.3 or R.

614
References

1.   Siegel, R. L., Miller, K. D. & Jemal, A. Cancer statistics, 2020. *CA. Cancer J. Clin.* **70**, 7–
30 (2020).

2.   Hinohara, K. & Polyak, K. Intratumoral Heterogeneity: More Than Just Mutations. *Trends*
Cell Biol. **29**, 569–579 (2019).

3.   Turajlic, S. & Swanton, C. Metastasis as an evolutionary process. *Science (80-.).* **352**,
169–175 (2016).

4.   Batlle, E. & Clevers, H. Cancer stem cells revisited. *Nat. Med.* **23**, 1124–1134 (2017).

5.   Pattabiraman, D. R. & Weinberg, R. A. Tackling the cancer stem cells — what challenges
do they pose? *Nat. Rev. Drug Discov. 2014 137* **13**, 497–512 (2014).

6.   Wang, M. L., Chiou, S. H. & Wu, C. W. Targeting cancer stem cells: emerging role of
Nanog transcription factor. *Onco. Targets. Ther.* **6**, 1207 (2013).

7.   Ray, A., Slama, Z. M., Morford, R. K., Madden, S. A. & Provenzano, P. P. Enhanced
Directional Migration of Cancer Stem Cells in 3D Aligned Collagen Matrices. *Biophys. J.*
112, 1023–1036 (2017).

8.   Pang, M.-F. *et al.* Tissue Stiffness and Hypoxia Modulate the Integrin-Linked Kinase ILK
to Control Breast Cancer Stem-like Cells. *Cancer Res.* **76**, 5277–5287 (2016).

9.   Conklin, M. W. *et al.* Aligned Collagen Is a Prognostic Signature for Survival in Human
Breast Carcinoma. *Am. J. Pathol.* **178**, 1221–1232 (2011).

10.   Provenzano, P. P. *et al.* Collagen reorganization at the tumor-stromal interface facilitates
local invasion. *BMC Med.* **4**, 38 (2006).

11.   Auvinen, P. *et al.* Hyaluronan in Peritumoral Stroma and Malignant Cells Associates with
Breast Cancer Spreading and Predicts Survival. *Am. J. Pathol.* **156**, 529–536 (2000).

12.   Marusyk, A. *et al.* Spatial Proximity to Fibroblasts Impacts Molecular Features and
Therapeutic Sensitivity of Breast Cancer Cells Influencing Clinical Outcomes. *Cancer*
Res. **76**, 6495–6506 (2016).

13.   Jacobetz, M. A. *et al.* Hyaluronan impairs vascular function and drug delivery in a mouse
model of pancreatic cancer. *Gut* **62**, 112–120 (2013).

14.   Hartheimer, J. S., Park, S., Rao, S. S. & Kim, Y. Targeting Hyaluronan Interactions for
Glioblastoma Stem Cell Therapy. *Cancer Microenviron.* **12**, 47–56 (2019).

15.   Chanmee, T. *et al.* Excessive Hyaluronan Production Promotes Acquisition of Cancer
646   Stem Cell Signatures through the Coordinated Regulation of Twist and the Transforming
647   Growth Factor β (TGF-β)-Snail Signaling Axis. *J. Biol. Chem.* **289**, 26038–26056 (2014).

648   16.   Auvinen, P. *et al.* Hyaluronan synthases (HAS1–3) in stromal and malignant cells
649   correlate with breast cancer grade and predict patient survival. *Breast Cancer Res. Treat.*
650   **143**, 277–286 (2014).

651   17.   Chanmee, T. *et al.* Hyaluronan Production Regulates Metabolic and Cancer Stem-like
652   Properties of Breast Cancer Cells via Hexosamine Biosynthetic Pathway-coupled HIF-1
653   Signaling. *J. Biol. Chem.* **291**, 24105–24120 (2016).

654   18.   Kuo, J. C.-H., Gandhi, J. G., Zia, R. N. & Paszek, M. J. Physical biology of the cancer cell
655   glycocalyx. *Nat. Phys.* **14**, 658–669 (2018).

656   19.   Turley, E. A., Wood, D. K. & McCarthy, J. B. Carcinoma Cell Hyaluronan as a "Portable"
657   Cancerized Prometastatic Microenvironment. *Cancer Res.* **76**, 2507–2512 (2016).

658   20.   Brett, M.-E. *et al.* In vitro elucidation of the role of pericellular matrix in metastatic
659   extravasation and invasion of breast carcinoma cells. *Integr. Biol.* **10**, 242–252 (2018).

660   21.   Offeddu, G. S. *et al.* The cancer glycocalyx mediates intravascular adhesion and
661   extravasation during metastatic dissemination. *Commun. Biol.* **4**, 255 (2021).

662   22.   Diehn, M. *et al.* Association of reactive oxygen species levels and radioresistance in
663   cancer stem cells. *Nature* **458**, 780–783 (2009).

664   23.   Luo, M. *et al.* Targeting Breast Cancer Stem Cell State Equilibrium through Modulation
665   of Redox Signaling. *Cell Metab.* **28**, 69-86.e6 (2018).

666   24.   Peiris-Pagès, M., Martinez-Outschoorn, U. E., Pestell, R. G., Sotgia, F. & Lisanti, M. P.
667   Cancer stem cell metabolism. *Breast Cancer Res.* **18**, 55 (2016).

668   25.   Zanotelli, M. R. *et al.* Regulation of ATP utilization during metastatic cell migration by
669   collagen architecture. *Mol. Biol. Cell* **29**, 1–9 (2018).

670   26.   Cunniff, B., McKenzie, A. J., Heintz, N. H. & Howe, A. K. AMPK activity regulates
671   trafficking of Mitochondria to the leading edge during cell migration and matrix invasion.
672   *Mol. Biol. Cell* **27**, 2662–2674 (2016).

673   27.   Thiagarajan, P. S. *et al.* Development of a Fluorescent Reporter System to Delineate
674   Cancer Stem Cells in Triple-Negative Breast Cancer. *Stem Cells* **33**, 2114–2125 (2015).

675   28.   Romani, P., Valcarcel-Jimenez, L., Frezza, C. & Dupont, S. Crosstalk between
676   mechanotransduction and metabolism. *Nat. Rev. Mol. Cell Biol.* **22**, 22–38 (2021).

677   29.   Ginestier, C. *et al.* ALDH1 Is a Marker of Normal and Malignant Human Mammary Stem
678   Cells and a Predictor of Poor Clinical Outcome. *Cell Stem Cell* **1**, 555–567 (2007).

30.   Vincent, K. M., Findlay, S. D. & Postovit, L. M. Assessing breast cancer cell lines as
tumour models by comparison of mRNA expression profiles. *Breast Cancer Res.* **17**, 114
(2015).

31.   Chokchaitaweesuk, C., Kobayashi, T., Izumikawa, T. & Itano, N. Enhanced hexosamine
metabolism drives metabolic and signaling networks involving hyaluronan production and
O-GlcNAcylation to exacerbate breast cancer. *Cell Death Dis.* **10**, 803 (2019).

32.   Rilla, K. *et al.* Hyaluronan Synthase 1 (HAS1) Requires Higher Cellular UDP-GlcNAc
Concentration than HAS2 and HAS3. *J. Biol. Chem.* **288**, 5973–5983 (2013).

33.   Rilla, K. *et al.* Hyaluronan production enhances shedding of plasma membrane-derived
microvesicles. *Exp. Cell Res.* **319**, 2006–2018 (2013).

34.   Itano, N. *et al.* Three Isoforms of Mammalian Hyaluronan Synthases Have Distinct
Enzymatic Properties. *J. Biol. Chem.* **274**, 25085–25092 (1999).

35.   Al-Hajj, M., Wicha, M. S., Benito-Hernandez, A., Morrison, S. J. & Clarke, M. F.
Prospective identification of tumorigenic breast cancer cells. *Proc. Natl. Acad. Sci.* **100**,
3983–3988 (2003).

36.   Liu, S. *et al.* Breast Cancer Stem Cells Transition between Epithelial and Mesenchymal
States Reflective of their Normal Counterparts. *Stem Cell Reports* **2**, 78–91 (2014).

37.   Li, W. *et al.* Unraveling the roles of CD44/CD24 and ALDH1 as cancer stem cell markers
in tumorigenesis and metastasis. *Sci. Rep.* **7**, 13856 (2017).

38.   Sheridan, C. *et al.* CD44+/CD24-breast cancer cells exhibit enhanced invasive properties:
an early step necessary for metastasis. *Breast Cancer Res.* **8**, R59 (2006).

39.   Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.* **28**,
245–248 (2010).

40.   Asgari, Y., Zabihinpour, Z., Salehzadeh-Yazdi, A., Schreiber, F. & Masoudi-Nejad, A.
Alterations in cancer cell metabolism: The Warburg effect and metabolic adaptation.
Genomics **105**, 275–281 (2015).

41.   Zielinski, D. C. *et al.* Systems biology analysis of drivers underlying hallmarks of cancer
cell metabolism. *Sci. Reports 2017 71* **7**, 1–14 (2017).

42.   Korkaya, H., Liu, S. & Wicha, M. S. Breast cancer stem cells, cytokine networks, and the
tumor microenvironment. *J. Clin. Invest.* **121**, 3804–3809 (2011).

43.   Yoon, C. *et al.* PI3K/Akt pathway and Nanog maintain cancer stem cells in sarcomas.
Oncogenesis **10**, 12 (2021).

44.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection.
Cell Syst. **1**, 417–425 (2015).

45.   Miroshnikova, Y. A. *et al.* Tissue mechanics promote IDH1-dependent HIF1α–tenascin C
feedback to regulate glioblastoma aggression. *Nat. Cell Biol.* **18**, 1336–1345 (2016).

46.   Schaefer, C. F. *et al.* PID: the Pathway Interaction Database. *Nucleic Acids Res.* **37**,
D674–D679 (2009).

47.   Semenza, G. L. Hypoxia-inducible factor 1: oxygen homeostasis and disease
pathophysiology. *Trends Mol. Med.* **7**, 345–350 (2001).

48.   Slemc, L. & Kunej, T. Transcription factor HIF1A: downstream targets, associated
pathways, polymorphic hypoxia response element (HRE) sites, and initiative for
standardization of reporting in scientific literature. *Tumor Biol.* **37**, 14851–14861 (2016).

49.   Hackett, S. R. *et al.* Systems-level analysis of mechanisms regulating yeast metabolic
flux. *Science (80-.).* **354**, aaf2786–aaf2786 (2016).

50.   Hu, H. *et al.* Phosphoinositide 3-Kinase Regulates Glycolysis through Mobilization of
Aldolase from the Actin Cytoskeleton. *Cell* **164**, 433–446 (2016).

51.   Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven
cancers require TBK1. *Nature* **462**, 108–112 (2009).

52.   Shurer, C. R. *et al.* Physical Principles of Membrane Shape Regulation by the Glycocalyx.
Cell **177**, 1757-1770.e21 (2019).

53.   Bergert, M. *et al.* Force transmission during adhesion-independent migration. *Nat. Cell
Biol.* **17**, 524–529 (2015).

54.   Paszek, M. J. *et al.* The cancer glycocalyx mechanically primes integrin-mediated growth
and survival. *Nature* **511**, 319–325 (2014).

55.   Barnes, J. M. *et al.* A tension-mediated glycocalyx–integrin feedback loop promotes
mesenchymal-like glioblastoma. *Nat. Cell Biol.* **20**, 1203–1214 (2018).

56.   Sullivan, W. J. *et al.* Extracellular Matrix Remodeling Regulates Glucose Metabolism
through TXNIP Destabilization. *Cell* **175**, 117-132.e21 (2018).

57.   Lewis, N. E. & Abdel-Haleem, A. M. The evolution of genome-scale models of cancer
metabolism. *Front. Physiol.* **4**, 1–7 (2013).

58.   Simmen, F. A., Alhallak, I. & Simmen, R. C. M. Malic enzyme 1 (ME1) in the biology of
cancer: it is not just intermediary metabolism. *J. Mol. Endocrinol.* **65**, R77–R90 (2020).

59.   Jiang, P., Du, W., Mancuso, A., Wellen, K. E. & Yang, X. Reciprocal regulation of p53
and malic enzymes modulates metabolism and senescence. *Nature* **493**, 689–693 (2013).

60.   Cheng, C.-P. *et al.* The mechanisms of malic enzyme 2 in the tumorigenesis of human
gliomas. *Oncotarget* **7**, 41460–41472 (2016).

61.   Dong, C. *et al.* Loss of FBP1 by Snail-Mediated Repression Provides Metabolic
Advantages in Basal-like Breast Cancer. *Cancer Cell* **23**, 316–331 (2013).

62.   Ciavardelli, D. *et al.* Breast cancer stem cells rely on fermentative glycolysis and are
sensitive to 2-deoxyglucose treatment. *Cell Death Dis. 2014 57* **5**, e1336–e1336 (2014).

63.   Liberti, M. V. *et al.* A Predictive Model for Selective Targeting of the Warburg Effect
through GAPDH Inhibition with a Natural Product. *Cell Metab.* **26**, 648-659.e8 (2017).

64.   Tanner, L. B. *et al.* Four Key Steps Control Glycolytic Flux in Mammalian Cells. *Cell
Syst.* **7**, 49-62.e8 (2018).

65.   Park, J. S. *et al.* Mechanical regulation of glycolysis via cytoskeleton architecture. *Nature*
578, 621–626 (2020).

66.   Bezanson, J., Edelman, A., Karpinski, S. & Shah, V. B. Julia: A Fresh Approach to
Numerical Computing. *SIAM Rev.* **59**, 65–98 (2017).

67.   Hu, Y. & Smyth, G. K. ELDA: Extreme limiting dilution analysis for comparing depleted
and enriched populations in stem cell and other assays. *J. Immunol. Methods* **347**, 70–78
(2009).

68.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

69.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

70.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

71.   Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven
cancers require TBK1. *Nature* **462**, 108–112 (2009).

72.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
microarray and RNA-Seq data. *BMC Bioinformatics* **14**, 7 (2013).

771
772

773
774
Figure 1: CSCs have increased invasive potential that is sensitive to metabolic challenge. a)
Schematic of the cancer stem cell reporter line GFP-NANOG MDA-MB-231 and sorting
strategy. **b)** Growth curve of sorted cells as measured by DNA amount, normalized to the first
measurement at t = 12 hrs (n = 3 samples). **c)** Schematic of the microfluidic device to analyze
cell invasion in response to a morphogen gradient generated by applying serum-containing
medium to the right channel only. **d)** Representative confocal reflectance microscopy image of a
fibrillar collagen hydrogel in microfluidic invasion device. **e)** Invasion distance of sorted GFP-
NANOG MDA-MB-231 into collagen type I over 5 days (n = 4 fields of views, 1 device per
condition). **f)** Invasion of GFP-NANOG MDA-MB-231 cells into collagen type I treated with or
without 2-DG for 5 days (n = 3 devices per condition). **g)** Random migration of GFP-NANOG
MDA-MB-231 treated with or without 2-DG for 24 hours (n = 5 representative cells for
migration plots, n = 20 cells per condition for motility). Scale bar = 250 μm. * p< 0.05, **
p<0.01, *** p<0.001, **** p<0.0001

789
790
791
Figure 2: Cancer stem-like cells exhibit altered metabolism compared to non-stem-like
cancer cells. a) Glucose and lactate concentrations in media conditioned by sorted GFP-
NANOG MDA-MB-231 cells 48 hrs post sort as measured by a GlucCell glucose meter and a
lactate colorimetric assay (n = 3 samples). **b)** ECAR and OCR measurements of sorted GFP-
NANOG cells obtained during Seahorse-based glycolysis stress test (n > 6 samples). **c)** Fraction
of GFP positive (GFP+) GFP-NANOG MDA-MB-231 (n = 6) (i) and aldehyde dehydrogenase
high (ALDH+) (n = 3) (ii) parental MDA-MB-231 when cultured with 100 mg/dL glucose
(+glucose) or glucose-free media (-glucose) for 72 hours. GFP+ and ALDH+ cells were
determined by image analysis and Aldefluor assay, respectively. **d)** Fraction of GFP+ GFP-
NANOG MDA-MB-231 cells treated with or without 2-deoxyglucose (2-DG) (n = 3). **e)**
Immunofluorescence analysis of HA in the MDA-MB-231 NANOG reporter line. (n = 248 cells)
Scale bar = 50 μm. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.
804
805

806
807
808
Figure 3: Increased HA production correlates with an increase in CSCs. a) Flow cytometry
analysis of cellular GFP levels categorized by their levels of cell surface-associated HA. **b)**
Heatmap representing changes of intracellular metabolites as measured by metabolomics of
sorted HA^{High} and HA^{Low} GFP-NANOG MDA-MB-231. **c)** Graphical representation of selected
metabolites in the central carbon metabolic pathway ratios between HA^{High} and HA^{Low} cells. **d)**
Schematic representing the theorized relationship between glycolysis, HA synthesis, and
stemness. **e)** Representative images of spheres formed through a limiting dilution assay of HAS2
and HAS3 overexpressing MDA-MB-231 and the corresponding sphere number. (n = 11) **f)**
Flow cytometry analysis of the percentage of the CD44^{+}/CD24^{-} population of parental or HA
overproducing MCF10A cells (n = 5). * p<0.05, ** p< 0.01, **** p<0.001. Scale bar = 200 μm.
819

820
821
822
Figure 4: Flux balance analysis predicts increased HA production in cancer stem-like cells
and ATP production. a) Workflow schematic. Sorted GFP-NANOG cells were cultured for 72
hrs before performing extracellular metabolomics using LC-MS. This data was used to constrain
a computational flux balance analysis model. **b)** Flux balance analysis model of the bioenergetic
pathway of sorted GFP-NANOG MDA-MB-231 derived from extracellular metabolomics
obtained over 72 hours of culture. Fluxes shown are GFP[High] relative to GFP[Null]. Grey
metabolites indicate relevant metabolite fluxes constrained based on extracellular metabolomic
profiles. Oxygen was constrained using values obtained from the Agilent Seahorse assay. For the
total FBA model constraints and results, see Supplementary File 1. **c)** Expanded HA synthesis
pathway flux balance analysis of GFP[High] relative to GFP[Null] (dashed box in B). Same legend as
in B. **d)** ATP production rate and **e)** rate index derived from measurements using the Agilent
Seahorse ATP real-time production rate assay kit ($n \geq 5$). * $p<0.05$, ** $p< 0.01$.
835

Figure 5: Increased glycolytic metabolism necessary for HA production stimulates stemness and invasion of breast cancer cells. a) Extracellular acidification rate measurements of parental MCF10A or HA overproducing cells using the Agilent Seahorse XF Analyzer. **b)** Percentage of CD44$^+$/CD24$^-$ cells of parental MCF10A or HA overproducing cells with or without 20mM 2-DG as measured by flow cytometry. **c)** ATP production rate of either MCF10A or MDA-MB-231 cells overexpressing HAS2 or HAS3 pre-treated with or without 2-DG (20mM for MCF10A, 25mM for MDA-MB-231) as measured using the Real-Time ATP Rate Assay in the Agilent Seahorse XF Analyzer. (n ≥ 8) **d)** Invasion of the MCF10A and **e)** MDA-MB-231 HA overproducing cells. Representative immunofluorescence images and corresponding quantification of invasion after 5 days are shown (n = 3 devices per condition). Scale bar: 200 μm, * p< 0.05, ** p<0.01, *** p<0.001, **** p<0.0001

848
849
Figure 6: Gene signatures of HA overproducing cells correlate with worse patient
prognosis regardless of HAS isoform. a) Principle Component Analysis plot of RNA
sequencing results of the MCF10A or HA overproducing cells. **b)** Heatmap of z-score
normalized gene expression of differentially expressed genes with hierarchical clustering of both
samples and genes. **c)** Gene Set Enrichment Analysis of the Hallmarks gene signature database
for genes upregulated in either HAS2 or HAS3 overexpressing cells contrasted against Parental
MCF10A cells. **d)** Gene Set Enrichment Analysis of published HIF1α target genes for HAS2 or
HAS3 overexpressing cells. All results are not significant. (FDR q-values ≥ 0.25) **e)** Venn
diagram displaying the total number of significantly increased genes (FDR ≤ 0.05, log2(Fold
Change) ≥ 1) in either HAS2 or HAS3 overexpressing cells. The top 4 most significant genes
excluding HAS2/3 are shown in the table. **f)** Ranked enrichment analysis of a curated list of

migration and metabolism gene sets for tumors from the METABRIC cohort. The degree of
enrichment is determined as the $-\log_{10}$(p-value) multiplied by the sign of the difference in
enrichment scores (ES) of patients stratified into either low or high enrichment of the HA
overproduction gene signature. The dashed red line denotes the empirically determined cutoff for
non-random enrichment. **g)** METABRIC patient survival probability predicted by the
overlapping 72-gene signature of HA overexpressing cells. Patient signature scores were
stratified into quartiles, and a log-rank test was used to determine statistical significance.