Proposal and extensive test of a calibration protocol for crop phenology models

Daniel Wallach[1], Taru Palosuo[2], Peter Thorburn[3], Henrike Mielenz[4], Samuel Buis[5], Zvi Hochman[3], Emmanuelle Gourdain[6], Fety Andrianasolo[6], Benjamin Dumont[7], Roberto Ferrise[8], Thomas Gaiser[9], Cecile Garcia[6], Sebastian Gayler[10], Matthew Harrison[11], Santosh Hiremath[12], Heidi Horan[3], Gerrit Hoogenboom[13,14], Per-Erik Jansson[15], Qi Jing[16], Eric Justes[17], Kurt-Christian Kersebaum[18,19], Marie Launay[20], Elisabet Lewan[20], Ke Liu[11], Fasil Mequanint[10], Marco Moriondo[22], Claas Nendel[18,19,23], Gloria Padovan[8], Budong Qian[16], Niels Schütze[24], Diana-Maria Seserman[18], Vakhtang Shelia[13,14], Amir Souissi[25], Xenia Specka[18], Amit Kumar Srivastava[9], Giacomo Trombi[8], Tobias K.D. Weber[10], Lutz Weihermüller[26], Thomas Wöhling[24,27], Sabine J. Seidel[9*]

[1]INRAE, UMR AGIR, Castanet Tolosan, France. ORCID 0000-0003-3500-8179

[2]Natural Resources Institute Finland (Luke), Helsinki, Finland

[3]CSIRO Agriculture and Food, Brisbane, Queensland, Australia

[4]Institute for Crop and Soil Science, Federal Research Centre for cultivated Plants, Julius Kühn-Institut (JKI), Braunschweig, Germany

[5]INRAE, UMR 1114 EMMAH, Avignon, France

[6]ARVALIS - Institut du végétal Paris, France

[7]Plant Sciences & TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium

[8]Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, Italy

[9]Institute of Crop Science and Resource Conservation, University of Bonn, Germany

[10]Institute of Soil Science and Land Evaluation, Biogeophysics, University of Hohenheim, Stuttgart, Germany

[11]Tasmanian Institute of Agriculture, University of Tasmania Launceston, Australia

[12]Aalto University School of Science, Espoo, Finland;

[13]Agricultural and Biological Engineering Department, University of Florida, Gainesville, Florida, USA

[14]Food Systems Institute, University of Florida, Gainesville, Florida, USA

[15]Royal Institute of Technology (KTH), Stockholm, Sweden

[16]Ottawa Research and Development Centre, Agriculture and Agri-Food Canada, Ottawa, Canada

[17]CIRAD, UMR SYSTEM, Montpellier, France

[18]Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

[19]Global Change Research Institute CAS, Brno, Czech Republic[19]INRAE, US 1116 AgroClim, Avignon,
France

[20]INRAE, US 1116 AgroClim, Avignon, France

[21]Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Uppsala,
Sweden

[22]CNR-IBE, Firenze, Italy

[23]Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[24]Institute of Hydrology and Meteorology, Chair of Hydrology, Technische Universität Dresden, Dresden,
Germany

[25]Swift Current Research and Development Centre, Agriculture and Agri-Food Canada, Swift Current,
Saskatchewan, Canada.

[26]Institute of Bio- and Geosciences - IBG-3, Agrosphere, Forschungszentrum Jülich GmbH, Jülich,
Germany

[27]Lincoln Agritech Ltd., Hamilton, New Zealand
50

51   [*]Correspondence to sabine.seidel@uni-bonn.de
52

Abstract

A major effect of environment on crops is through crop phenology, and therefore, the capacity to predict phenology for new environments is important. Mechanistic crop models are a major tool for such predictions, but calibration of crop phenology models is difficult and there is no consensus on the best approach. Here we propose an original, , detailed approach, a protocol, for calibration of such models. The protocol covers all the steps in the calibration work-flow, namely choice of default parameter values, choice of objective function, choice of parameters to estimate from the data, calculation of optimal parameter values and diagnostics. The major innovation is in the choice of which parameters to estimate from the data, which combines expert knowledge and data-based model selection. First, almost additive parameters are identified and estimated. This should make bias (average difference between observed and simulated values) nearly zero. These are "obligatory" parameters, that will definitely be estimated. Then candidate parameters are identified, which are parameters likely to explain the remaining discrepancies between simulated and observed values. A candidate is only added to the list of parameters to estimate if it leads to a reduction in BIC (Bayesian Information Criterion), which is a model selection criterion. A second original aspect of the protocol is the specification of documentation for each stage of the protocol. The protocol was applied by 19 modeling teams to three data sets for wheat phenology. All teams first calibrated their model using their "usual" calibration approach, so it was possible to compare usual and protocol calibration. Evaluation of prediction error was based on data from sites and years not represented in the training data. Compared to usual calibration, calibration following the new protocol reduced the variability between modeling teams by 22% and significantly reduced prediction error.

Key words

crop model, prediction error, protocol, model ensemble, variability

1. Introduction

Plant phenology is a major aspect of plant response to environment, and a major determinant of plant response to climate change. This includes phenology of natural vegetation, which is a dominant aspect of plant ecology (Cleland et al. 2007) and has been shown to be affected by warming (Piao et al. 2019; Menzel et al. 2020; Stuble et al. 2021) and phenology of cultivated crops. For the latter, phenology must be taken into account for crop management (Sisheber et al. 2022), choice of cultivar or cultivar characteristics adapted to a particular region (Zhang et al. 2022) and for evaluating the impact of climate change on crop production (Rezaei et al. 2018). It is thus important to be able to predict phenology as a function of environment, in particular as a function of climate.

A number of mechanistic crop models have been developed, which include simulation of phenology, and such models are regularly used to evaluate management options (McNunn et al. 2019) or the effect of climate change on crops, including wheat (Asseng et al. 2013), rice, (Li et al. 2015), maize (Bassu et al. 2014) and others. Such models are particularly important for taking into account an increasing diversity of combinations of weather events (Webber et al. 2020).

Mechanistic models in general, and models used to simulate crop phenology in particular (we will refer to such models as crop phenology models, though they are usually embedded within more general crop models), are based on our understanding of the processes and their inter-linkages that drive the evolution of the system. This conceptual understanding usually builds on detailed experiments that study specific aspects of the system (e.g. Brisson et al., 2003 for the crop model STICS). The set of model equations is referred to as "model structure" (Tao et al. 2018).

In addition to model structure, simulation requires values for all the model parameters. In essentially all uses of crop models, the model is first calibrated using observed data that is related to the target population for which predictions are required, for example observations for the specific variety of interest and/or for the particular set of growing environments of interest. Calibration normally only concerns a fairly small subset of the parameters in a crop model, but is essentially always necessary because mechanistic models are only approximations, without universally valid parameter values (Fath and Jorgensen 2011; Wallach 2011).

There are therefore two main tracks to improvement of crop phenology models. The first is through improvement of model structure through improved understanding of the underlying processes, and the second is through improvement of the model parameters. For a fixed data set, improvement of model parameters implies improvement of model calibration, and that is the topic here.

Calibration of crop models is usually patterned on methods developed for regression models in statistics. However, the application of statistical methods to system models is not straightforward. Among the difficulties are the fact that system models often have multiple output variables that can be compared to observed results (e.g., dates of heading and dates of flowering for crop phenology models), that errors for different variables in the same environment are often correlated and that there are usually many parameters, often more than the number of data points available. While the details differ, these problems apply to essentially all system models, not only crop phenology models but also full crop models, hydrological models, ecology models and models in other fields. No doubt as a result, there are no widely accepted standard methods for calibration of system models. It has been found, for example, that there is a wide diversity of calibration approaches between modeling teams furnished with identical data, even between modeling teams using the same model structure (Confalonieri et al. 2016; Wallach et al. 2021a, b), where a modeling team is a group of people working together on or with a crop model.

Because of the importance of calibration and the lack of standard approaches for calibration, there have been many studies published that make recommendations as to how to calibrate crop models or

system models in other fields. One type of study is model-specific, and identifies the most important
parameters to estimate for a particular model (Ahuja and Ma, 2011). Other studies have focused on the
methodology of identifying the most important parameters through sensitivity analysis (Khorashadi
Zadeh et al. 2022), on the choice between frequentist and Bayesian paradigms (Gao et al. 2021), on the
form of the objective function, or on the numerical algorithm for searching for the best parameter values
(Rafiei et al. 2022). A recent study has shown that different modeling teams make different choices for all
the steps of the calibration procedure, (Wallach et al., 2021c). That study showed that the modeling
community is far from having a consensus on how to calibrate phenology models, and emphasized the
need to treat calibration holistically, taking into account all aspects.

The purpose of this study was to define and test an original, detailed, comprehensive procedure
for crop phenology model calibration that could be applied to a wide range of crop phenology models. We
refer to this new procedure as a "protocol", to emphasize that it contains detailed instructions for
calibration. It builds on the recommendations in Wallach et al. (2021c) but goes beyond those more
general recommendations, most importantly in proposing an original approach for choosing the
parameters to estimate, which is arguably the most important calibration decision. A second major
innovation of the protocol is the definition of documentation tables for each step of the calibration
procedure, which can be used both for communication within a modeling team and to inform users of the
calibrated model. We tested the protocol in a large multi-model ensemble study where the objective was to
simulate wheat phenology in multiple environments (combinations of site and sowing date) in France and
in Australia. Nineteen modeling teams participated, and were all able to implement the protocol, showing
that the protocol, though detailed, is nonetheless sufficiently flexible to be applicable to a wide range of
models. The protocol was tested in comparison with the "usual" calibration procedure of each modeling
team, which was possible because each modeling team had previously calibrated their model using the
same data as here. To our knowledge this is the first example of such a stringent test for a new calibration
procedure. It provides a realistic test of whether the proposed protocol really improves calibration. The
current study showed that the protocol reduced the variability between modeling teams compared to usual
calibration approaches and, most importantly, that it significantly reduced prediction error compared to
usual calibration approaches.

2. Materials and Methods

Data sets

Three data sets for wheat phenology were used here, each with data from multiple environments,
i.e. combinations of site, year and sowing date. The first two data sets were from cultivar trials in France,
for the two winter wheat varieties Apache and Bermude (Wallach et al 2021a). For both cultivars there
were the same 14 calibration environments (6 different sites, sowing years 2010, 2011, 2014-2016 but
not every year was represented for every site) and 8 evaluation environments (5 different sites, sowing
years 2012 and 2013). The calibration and evaluation subsets had neither site nor year in common, so the
evaluation is a rigorous test of how well a modeling team can simulate phenology for out-of-sample
environments. The observed data were days from sowing to beginning of stem elongation (BBCH30 on
the BBCH scale, Meier 1997) and to middle of heading (BBCH 55).

The third data set had data for the spring wheat variety Janz from a multi-location multi-year
multiple sowing date trial in Australia (Lawes et al. 2016; Wallach et al. 2021b). The calibration data
were from four sites in 2010 and 2011, with three sowing dates per site (overall 24 environments). The
evaluation data were from six sites in 2012, with three sowing dates per site (overall 18 environmets).
Once again, the calibration and evaluation subsets had neither site nor year in common. In the original
trials, BBCH development stage was observed once weekly in each environment. Based on those data, a
graph of BBCH stage versus day was produced, and interpolation was used to obtain the day for each
integer BBCH stage from the earliest to the latest recorded in each environment. Those dates were
provided to the modeling teams.

Modeling teams

Nineteen modeling teams, using 16 different model structures (Table 1), participated in this study,
which was carried out within the Agricultural Model Intercomparison and Improvement Project (AgMIP;
www.agmip.org). The modeling teams are identified only by a code ("M1", "M2" etc.) without indicating
which model structure they used, since it would be misleading to give the impression that the results are
determined solely by model structure.

Table 1: List of model structures used by participating modeling teams

Model structure	Version(s)	References
AgroC	May 2018	(Herbst et al. 2008; Klosterhalfen et al. 2017)
APSIM	7.8, 7.9, 7.10	(Keating et al. 2003; Holzworth et al. 2014)
AquaCrop	4.0	(Vanuytrecht et al. 2014)
CERES-Wheat	DSSAT V4.7.	(Hoogenboom et al., 2019a, 2019b; Jones et al., 2003)
CoupModel	Version 5.4.4	(P.-E. Jansson 2012; Senapati et al. 2016; Coucheney et al. 2018)
CROPSIM-Wheat	DSSAT V4.7	(Hoogenboom et al., 2019a, 2019b; Jones et al., 2003)
Cropsyst	3.04.08	(Stockle et al. 2001)
HERMES	4.27	(Kersebaum, 2007; Kersebaum, 2011)
LINTUL	LINTUL5	(Wolf 2012)
MONICA	2.02	(Nendel et al. 2011; Specka et al. 2015, 2019)
PANORAMIX	R version	(Gate 1995; Chatelin et al. 2005)
SPASS	Expert-N 5.0	(Wang 1997)
CERES	Expert-N 5.0	(Godwin et al.)
SSM-Wheat		(Soltani et al. 2013)

STICS	8_5_0	(Brisson et al. 2009; Coucheney et al. 2015)
WOFOST	7.1.7	(Boogaard et al. 1998)

Goodness-of-fit and evaluation of predictions

Goodness-of-fit refers to how well a calibrated model fits the data used for calibration. Prediction accuracy refers to how well a calibrated model simulates for environments different than those in the calibration data set,. Since the evaluation environments here are for sites and years not represented in the calibration data, the test of simulated values against the evaluation data truly reflects how well a model can predict for new environments.

For both goodness-of-fit and out-of-sample prediction, our basic evaluation metric is the sum of squared errors (SSE) and the related quantities mean squared error (MSE) and root mean squared error (RMSE), where

$$SSE = \sum_i \sum_j \left(y_{ij} - \hat{y}_{ij} \right)^2$$
$$MSE = SSE / n \qquad (1)$$
$$RMSE = \sqrt{MSE}$$

The sum for SSE is over variables and environments i.e. over all measurements for which there are corresponding simulated values. Here y_{ij} is the observed value of variable i for environment j, $\hat{y}_{ij}$ is the corresponding simulated value and n is the number of terms in the sum. We also look at the decomposition of MSE as the sum of three terms, namely squared bias (bias²), a term related to the difference in standard deviations of the observed and simulated values (SDSD) and a term related to the correlation of observed and simulated vales (LCS) (Kobayashi 2004).

In addition, we compare the simulated results in this study with two simple benchmark models. The first (the "naive" model) is simply the average number of days to each stage in the calibration data of each data set. This is used as the prediction model for all environments of that data set. The often used Nash Sutcliffe modeling efficiency is one minus the ratio of MSE of a model to MSE of the naive model. The naive model ignores all variability between environments, so it is a very low bar as a benchmark. We therefore also use a more sophisticated benchmark, the "onlyT" model, introduced in Wallach et al. (2021a). This benchmark model assumes that the sum of degree days above a threshold of 0°C from sowing to each stage is fixed for spring wheat. For winter wheat, a simple vernalization model is used to determine the start of development. Vernalization is 0 if daily mean temperature is below -4°C, increases linearly to 1 at 3°C, remains at 1 to 10°C, decreases linearly to 0 at 17°C and is 0 above 17. °C. When the sum of daily vernalization reaches 50, vernalization is complete (van Bussel et al. 2015; Wallach et al. 2021a). Then the fixed number of degree days applies after vernalization is completed. Both benchmark models are quite easily parameterized based on calibration data, and then easily applied to new environments.

Simulation exercise

The participants received input data (daily weather at the field, soil characteristics, management details and, where possible, initial conditions) for all environments of every data set. Also, the observed

data from the calibration environments were provided to all participants. The participants were asked to use those data to calibrate their models using the calibration protocol described in detail below, and then to simulate and report days after sowing to stages BBCH10, BBCH30, and BBCH55 for the French calibration and evaluation environments, and to stages BBCH10, BBCH30, BBCH65, and BBCH90 for the Australian calibration and evaluation environments. Days to emergence (BBCH10) was included to have an example of a variable for which there were no calibration data. The BBCH stages 30 and 55 requested for the French environments represent stages that are used for fertilizer decisions in France. The BBCH stages 30, 65, and 90 requested for the Australian environments represent major transitions that are explicitly simulated by many models.

Seventeen of the 19 participating modeling teams participated in previous phases of this project, and in that framework had already calibrated their model with the same data as here using their usual calibration approach (Wallach et al. 2021a, b). The two remaining teams also calibrated their models using their usual approach before beginning to use the protocol proposed here. It is the results of the usual calibration method that are compared here to the results of using the proposed protocol. At no time were the evaluation data shown to participants, neither in previous studies nor in the present study.

The protocol does not impose a specific software solution. However, several participants used trial and error in their usual approach and requested help in finding and implementing an automated search algorithm, since that is required for the protocol. To answer this need, the CroptimizR R package (Buis et al. 2021) was modified to do the protocol calculations, and many of the participants used this software.

In addition to the individual models, we report on two ensemble models, created by taking the mean (the e-mean model) or the median (the e-median model) of the simulated values. These ensemble models were calculated both for the usual and protocol calibration results.

AICc and BIC

The protocol prescribes a model selection criterion to decide which parameters to estimate. The corrected Akaike Information Crition (AICc) and the Bayesian Information Criterion (BIC) are two different criteria that are often used for model selection (Chakrabarti and Ghosh 2011). Both are based on model error, with a penalization term that increases with the number of estimated parameters. Assuming that model errors are normally distributed, the criteria are

$$AICc = n\ln(MSE) + 2p + \frac{2p(p+1)}{n-p-1} \tag{2}$$

$$BIC = n\ln(MSE) + p\ln(n)$$

where n is the number of data points and p is the number of calibrated parameters. These criteria are only used for comparing models calibrated using the same data.

There have been comparisons between these criteria, but there does not seem to be one that systematically performs better than the other, for choosing the model that predicts best (Kuha 2004). In applying the protocol here, participants were asked to perform the calculations twice, once using the AICc criterion and once using the BIC criterion to choose the parameters to estimate. In almost all cases, the two criteria led to exactly the same choice of parameters. In the few cases where the criteria led to different choices, the final models had very similar RMSE for the evaluation data, with a very slight advantage to BIC (Supplementary tables S24-S25). All results shown here are based on the BIC criterion.

3. Results and discussion

Description of protocol

The protocol is based on the recommendations in Wallach et al. (2021c), and follows the same list of steps (Figure 1), but has important additions, in particular for the choice of parameters to estimate and the documentation to be produced,

Figure 1. Schematic diagram of steps in proposed calibration protocol.

Step 1. Choose default parameter values

Since most parameters retain their default values, the choice of default values for those parameters that affect phenology is important. For phenology, one would want to have reasonable approximations to the cycle length for the cultivar in question, to photoperiod dependence and to vernalization requirements. This information and more is usually available from the cultivar developer. The documentation for step 1 (see example in Table 2) specifies the characteristics of the cultivar being modeled and those of the cultivar used to provide default parameter values.

280 **Table 2: Example of protocol documentation for Step 1, "Choose default parameter**
281 **values". The first row shows cultivar characteristics of the observed cultivar. The second**
282 **row shows characteristics of the cultivar that provides the default parameter values. This**
283 **example is for the French data set and modeling team M21.**

Variety	Characteristics
Variety of target population: Apache	A soft winter wheat. Stem elongation – semi-early. Heading – early. Vernalization requires 40 days where full vernalization occurs if daily average temperature is between 3°C and 10°C. There is no vernalization below -4°C or above 17°C. Otherwise there is a proportional reduction in vernalization effectiveness.
Default variety : Soissons	Soissons seems to be close to Apache in terms of vernalization requirements and earliness.

Step 2 Identify correspondence between observed and simulated variables

In the simplest case, there is a simulated variable that corresponds directly to each observed
development stage. The documentation for step 2 is a table with one row for each observed variable,
showing the corresponding simulated variable if any (see example in Table 3) .

Table 3: Example of documentation for protocol Step 2, "Identify correspondence
between observed and simulated variables". The table has one row for each measured
variable, showing the corresponding simulated variable if any. This example is for the
French data sets and modeling team M21.

Measured variable	Corresponding simulated value
Days to BBCH stage 30	Days to end juvenile stage
Days to BBCH stage 55	Days at maximum LAI

297
298

Step 3. Define the objective function

The objective function of the protocol is squared error summed over development stages and
environments, which is the objective function of ordinary least squares (OLS) regression and is often used
in crop model calibration. A major choice of the protocol is to include in the objective function all the
observed development stages that have a simulated equivalent, including stages that are not of primary
interest. A first reason is that often the same calibrated model will be used for several different objectives,
so measured variables that are not of central interest in the current study may be important in future
studies. Furthermore, using more variables makes the model a better representation of multiple aspects of
the system dynamics, which is likely to improve all simulations. The choice in the protocol is to use OLS
and to avoid estimating additional parameters related to variance and covariance of errors. However, one
should check residual errors to evaluate the extent of heteroscedasticity or correlation of errors. Since the
objective function is the sum of squared errors over the variables from step two, no new decisions are
required here and no additional documentation is required.

Step 4. Choose which parameters to estimate

This is arguably the most difficult, and the most important, decision of the calibration approach.
Here we propose a novel approach which combines expert knowledge with a statistical model selection
criterion. This approach distinguishes two categories of parameters to estimate: the nearly additive,
obligatory parameters (those that will definitely be estimated) and the candidate parameters (those that
will be tested, and only changed from the default value if the improvement in the fit to the calibration data
is sufficiently large).

Step 4a Identify the obligatory parameters

The obligatory parameters are parameters that are nearly additive, i.e. such that changing the
parameter has a similar effect for all environments for some variable in the objective function. Usually, a
parameter that represents degree days to a measured stage is a good choice as an obligatory parameter for
time to that stage. Estimating a truly additive parameter (that adds the same constant amount to days to a
stage for all environments) will exactly eliminate bias for that stage (i.e. the mean of simulated values will
exactly equal the mean of observed values). Estimating an almost additive parameter will almost eliminate
bias. Once bias is nearly eliminated, one may already have a fairly reasonable fit to the data. Each almost
additive parameter must affect a different variable or combination of variables. There cannot be more
almost additive parameters than the number of variables in the objective function. Otherwise, the
parameters would be very poorly estimated, or non-estimable. The protocol does allow fewer almost
additive parameters than observed variables. In that case bias is only nearly eliminated on average over
several variables, and not for each variable. For each obligatory parameter one must provide the default
value and what one considers a reasonable range for that parameter (for an example of choice of bounds
see Tao et al. 2013). The documentation for step 4a is a table with one row for each obligatory parameter
(see example in Table 4).

Table 4*:* **Example of documentation for protocol step Step 4a, "Identify the**
obligatory parameters". These are parameters that are almost additive, i.e. that have nearly
the same effect for all environments. There is one row for each obligatory parameter. The
number of obligatory parameters cannot exceed the number of observed variables which
have a simulated equivalent, and each obligatory parameter must be nearly additive for a
different variable or combination of variables. This example is for the French data set,
variety Apache for modeling team M21.

Name of obligatory parameter	explanation	Default value (lower-upper limits)
stlevamf	Degree days sowing to end juvenile stage	233 (150-400)
stamflax	Degree days sowing to maximum LAI	354 (150-500)

Step 4b. Identify candidate parameters
The role of the candidate parameters is to reduce the variability between environments that
remains after estimation of the obligatory parameters. It is the role of the modeler to identify the candidate
parameters and to order them by amount of variability likely to be explained. In the calculation step (step
5), each candidate parameter is tested, and only those that lead to a reduction in the BIC criterion are
retained for estimation. Otherwise, the parameter is kept at its default value. The documentation for step
4b is a table with one row for each candidate parameter (see example in Table 5) .

Table 5*:* **Example of documentation for protocol Step 4b, "Identify candidate**
parameters". These are parameters that seem likely to explain a substantial part of the variability
between environments that remains after fitting the obligatory parameters. There is one row for
each candidate, which should be in the order of presumed importance. This example is for the
French data set, variety Apache for modeling team M21.

Candidate parameter (include units and a brief explanation)	Default value (Lower - upper bounds)
jvc (days): number of vernalizing days	38 (25 – 60)
sensrsec (no unit): index of root sensitivity to drought (1=insensitive)	0.5 (0 – 1)
belong (degree.day-1): parameter of the curve of coleoptile elongation	0.012 (0.005 – 0.03)
JVCmini (days): minimum vernalizing days required	7.0 (2 – 15)
stressdev (no unit): maximum phasic delay allowed due to stresses	0 (0 – 1)

Step 5. Calculation of the optimal parameter values

The protocol prescribes the use of a simplex algorithm for searching for the optimal parameter
values. The Nelder-Mead simplex method (Nelder and Mead 1965) is a robust, derivative-free method,
which is appropriate for crop models which may have multiple discontinuities. The results of the simplex
are sensitive to starting values (Wang and Shoup 2011), so the protocol calls for multiple starting points.
In the first calculation step, the obligatory parameters are estimated and the BIC value is calculated. This
is the initial list of parameter to estimate. Then each candidate parameter is tested in turn. If estimating the
new candidate together with the previous list of parameters to estimate leads to a reduction in BIC, the
candidate is added to the list of parameters to estimate. If not, the candidate returns to its default value and
will not be estimated. The documentation for step 5 is a table with one row for each step in the calculation
(see example in Table 6).

401   **Table *6:* Example of documentation for Step 5 "Calculation of the optimal parameter**
402 **values". The first line shows the optimization results for the obligatory parameters, and the**
403 **resulting sum of squared errors and BIC criterion. Each subsequent line corresponds to a**
404 **candidate parameter. If estimating the candidate together with the previously selected**

parameters leads to a decrease in BIC compared to the smallest value so far, the candidate

is added to the list of parameters to estimate. If not, the candidate returns to its default

value and is not considered further. In this example, the first candidate parameter (jvc) is

accepted, and all the subsequent candidate parameters increase BIC, and are therefore,

rejected. The model finally chosen (minimum BIC) has three estimated parameters. This

Estimated.parameters	Initial.parameter.values	Final.values	Sum.of. squared. errors	BIC
stlevamf, stamflax	multiple	227, 360	405	81.47
stlevamf, stamflax, jvc	**227, 360, multiple**	**212, 367, 55.91**	**349**	**80.64**
stlevamf, stamflax, jvc, sensrsec	212, 367, 55.91, multiple	209, 367, 58.40, 0.057	322	81.71
stlevamf, stamflax, jvc, belong	212, 367, 55.91, multiple	212, 367, 55.91, 0.012	349	83.97
stlevamf, stamflax, jvc, jvcmini	212, 367, 55.91, multiple	197, 362, 55.28, 20.88	319	81.45
stlevamf, stamflax, jvc, stressdev	212, 367, 55.91, multiple	212, 367, 55.91, 0.00	349	83.97

example is for the French data set, variety Apache, modeling team M21.

Step 6 Examine goodness-of-fit

Many diagnostics are possible and useful. We emphasize particularly a graph of simulated versus observed values (see example in fig 2), calculated MSE and its decomposition for each variable (see example in table 7) and comparison with the two benchmark models (see example in table 8).

Table 7: First example of documentation for protocol step Step 6, "Examine goodness-of-fit". In this table there is one row for each observed variable with a simulated equivalent, showing mean squared error (MSE) and its decomposition into three terms. Of particular interest is the bias² contribution, which should be small if there is an almost

428 **obligatory parameter corresponding to this variable. This example is for the French data**
429 **set, variety Apache, modeling team M21.**

	MSE (days²)	bias² (days²)	SDSD (days²)	LCS (days²)
BBCH30	19.64	0.25	5.93	13.46
BBCH55	5.29	0.02	0.03	5.24

Table 8: Second example of documentation for protocol Step 6, "Examine goodness-
of-fit". In this table there is one row for each observed variable with a simulated equivalent,
showing root mean squared error (RMSE) for the calibrated model and for two benchmark
models. The "naïve" benchmark assumes that all environments have the same number of
days to the given development stage, equal to the average of the observed days to that stage.
The "onlyT" benchmark assumes a constant number of degree days to the stage in
question, with a simple vernalization calculation in the case of winter wheat. This example
is for the French data set, variety Apache, modeling team M21.

variable	naive RMSE (days)	onlyT RMSE (days)	M21 RMSE (days)
BBCH30	12.5	8.4	3.1
BBCH55	8.3	9.5	3.8

Comparison of protocol and usual calibration

Using usual or protocol calibration led to important differences in the calibrated models. For example, the number of estimated parameters in the final model was different between protocol and usual calibration (Supplementary Figure S1, table S1). The differences between simulated values after usual and protocol calibration were small for BBCH10, for which there were no calibration data. For the other stages, the simulated values differed appreciably (Supplementary Figure S2, Table S2).

Goodness of fit and evaluation for usual and protocol calibration

Figures 2 and 3 show RMSE using usual and protocol calibration for the French and Australian data sets respectively (results by modeling team are in Supplementary Tables S3-S8). Table 9 shows RMSE values for each data set, averaged over modeling teams, for usual and protocol calibration for the calibration and evaluation data. The protocol reduces RMSE by 10-22% compared to the usual calibration method. The p values for a one-sided paired t-test of the hypothesis that RMSE is larger for usual calibration than for protocol calibration are also shown. On average over stages other than BBCH10, all three data sets have significantly larger RMSE values with usual calibration than with protocol calibration for the calibration data ($p<0.05$). For the evaluation data, $p<0.01$ for the two French data sets, but $p=0.15$ for the Australian data set. The table also shows the proportion of modeling teams where RMSE is larger for the usual calibration than for the protocol calibration. Looking at the averages over stages and then averaging over data sets, 75% of models have lower RMSE for protocol calibration than for usual calibration for the calibration data, and 66% for the evaluation data. Almost all modeling teams did better than the two benchmark models for all stages, both for usual calibration and protocol calibration, with slightly better results for protocol calibration (Supplementary Tables S3-S8). Since the protocol specifically aims to reduce bias, one would expect squared bias to be a smaller fraction of MSE for protocol calibration than for usual calibration, and this is the case, both for the calibration data and the evaluation data (Supplementary Tables S9-S23).

Figure 2. Root mean squared error (RMSE) of results following usual or protocol calibration, for each modeling team, separately for calibration and evaluation data of the French data sets.

Figure 3. Root mean squared error (RMSE) of results following usual or protocol calibration, for each modeling team, separately for calibration and evaluation data of the Australian data set.

Table 9*: **Comparison of errors for usual and protocol calibration. The table shows**
root mean squared error (RMSE) averaged over modeling teams for each stage and for the
average over stages, separately for the calibration and evaluation data. For each of
calibration and evaluation data, the first column is RMSE for simulations using the usual
calibration method, the second column is RMSE using protocol calibration, and the third
column is the p value of a one sided paired t test, that tests whether RMSE for usual
calibration is larger than for protocol calibration. Below the p value is the fraction of
modeling teams for which RMSE for usual calibration is larger than for protocol
calibration.

		Calibration data			Evaluation data		
		usual RMSE	protocol RMSE	p-value RMSE$_u$ > RMSE$_p$	usual RMSE	protocol RMSE	p-value RMSE$_u$ > RMSE$_p$
France Apache	BBCH30	.7	6	0.007 10/16	.7	.2 6	0.20 9/16
	BBCH55		4	0.02 11/17	.3	.8 5	0.005 13/17
	average	.4	5	0.004 14/17		6	0.006 14/17
France Bermude	BBCH30	.1	4 5.	0.05 11/16	.8	.5 7	0.07 9/16
	BBCH55	.8	7 5.	0.04 11/17	.9	.1 6	0.08 11/17
	average		6 5.	0.029 13/17	.8	.8 6	0.008 11/17
Australia	BBCH30	3.3	2.0 1	0.09 9/17	5.1	4.8 1	0.38 8/17
	BBCH65	1.4	0 9.	0.06 14/19	1.1	1.4 1	0.60 10/19
	BBCH90	1.9	9 9.	0.096 11/18	.0	.2 6	0.20 12/18
	average	2.2	0.2 1	0.049 13/19	1.7	0.7 1	0.15 10/19

Comparison of usual and protocol calibration for ensemble of models

The choice of usual or protocol calibration has little effect on the predictive accuracy of the ensemble models e-mean and e-median. Averaged over development stages and over data sets, for the evaluation data, RMSE for e-median is respectively 5.7 and 5.8 days for usual and protocol calibration. The values for RMSE of e-mean are 6.1 and 6.2 days for usual and protocol calibration, respectively (Supplementary Tables S4, S6, S8).

Recently, many crop model studies have been based on ensembles of models (Jägermeyr et al. 2021). Many studies have found that the ensemble mean and median are good predictors, sometimes better than even the best individual model (Martre et al. 2015; Wallach et al. 2018; Farina et al. 2021). It has thus, become quite common to base projections of climate change impact on crop production on the ensemble median (e.g. Asseng et al., 2019). The e-mean and e-median results here, compared to the individual modeling teams, are in keeping with previous results. The e-median model is among the better predictors though not the very best, and is somewhat better than e-mean.

The variability between simulated results for different modeling teams is shown in Table 10. The standard deviation is similar for usual and protocol calibration for BBCH10, for which there are no data for calibration, but is systematically smaller for protocol calibration for the other stages. Considering the average over stages other than BBCH10 and taking the average over data sets, protocol calibration reduced the standard deviation of simulated values by 31% for the calibration data and by 22% for the evaluation data.

Table 10*: Variability of simulated values. This table shows the standard deviations
of values simulated by the different modeling teams (days), for each simulated development
stage and for the average over stages other than BBCH10. BBCH is not considered, since
there were no observed values for BBCH10. Separate standard deviations are given for the
calibration and evaluation data, and simulation using usual calibration or protocol
calibration.

| | | | Calibration data | | Evaluation data | |
			usual	protocol	usual	protocol
France Apache	10	BBCH	4.2	4.1	4.8	5.2
	30	BBCH	6.4	4.3	6.2	5.5
	55	BBCH	4.5	3.0	6.3	3.7
		average	5.4	3.6	6.2	4.6
France Bermude	10	BBCH	4.3	4.4	4.8	6.6
	30	BBCH	6.9	4.3	6.7	6.2
	55	BBCH	4.8	3.5	5.8	4.3
		average	5.9	3.9	6.2	5.3
Australia	10	BBCH	8.6	7.5	9.6	8.0
	30	BBCH	11.3	7.3	10.2	8.1
	65	BBCH	10.3	7.4	8.2	7.1
	90	BBCH	11.2	9.3	11.1	6.9
		average	10.9	8.0	9.8	7.4

4. Conclusions

We propose an original, detailed protocol for calibration of crop phenology models and show that
it can be applied by a wide range of wheat models and for data sets with different structures. While the
application here is to wheat phenology models, the same protocol could undoubtedly be used more
generally for phenology models of other crops.

Comparison with usual calibration practices shows that, on average over modeling teams, the
protocol leads to a better fit to the calibration data and to a better fit to out-of-sample data. Application of
the protocol would be advantageous not just for individual modeling studies, but also for studies based on
ensembles of models, including projections of climate impact. In particular, we have shown that if all
modeling teams use the protocol, between-model variability can be substantially reduced, thus reducing
the uncertainty of projections.

Author Declarations

Funding

This study was implemented as a co-operative project under the umbrella of the Agricultural Model Intercomparison and Improvement Project (AgMIP). This work was supported by the Academy of Finland through projects AI-CropPro (316172 and 315896) and DivCSA (316215) and Natural Resources Institute Finland (Luke) through a strategic project EFFI, the German Federal Ministry of Education and Research (BMBF) in the framework of the funding measure "Soil as a Sustainable Resource for the Bioeconomy - BonaRes", project "BonaRes (Module B, Phase 3): BonaRes Centre for Soil Research, subproject B" (grant 031B1064B), the BonaRes project "I4S" (031B0513I) of the Federal Ministry of Education and Research (BMBF), Germany, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 2070 -390732324 EXC (PhenoRob), the Ministry of Education, Youth and Sports of Czech Republic through SustES - Adaption strategies for sustainable ecosystem services and food security under adverse environmental conditions (project no. CZ.02.1.01/0.0/0.0/16_019/000797), the Agriculture and Agri-Food Canada's Project J-002303 "Sustainable crop production in Canada under climate change" under the Interdepartmental Research Initiative in Agriculture, the JPI FACCE MACSUR2 project, funded by the Italian Ministry for Agricultural, Food, and Forestry Policies (D.M. 24064/7303/15 of 6/Nov/2015), and the INRAE CLIMAE meta-programme and AgroEcoSystem department. The order in which the donors are listed is arbitrary.

Conflicts of interests / Competing interests
The authors have no financial or proprietary interests in any material discussed in this article.

Code availability / Data availability
The datasets generated during and/or analyzed during the current study are not publicly available but are available from the authors on reasonable request.

Ethics approval/declarations
Not applicable

Consent to participate
Not applicable

Consent for publication
Not applicable

Authors' contributions

Daniel Wallach: Conceptualization, methodology, project administration, writing - original draft, validation. Taru Palosuo, Henrike Mielenz, Peter Thorburn, Samuel Buis, Sabine Seidel: Conceptualization, methodology, project administration, writing , review &editing. All other authors: Simulations, model expertise, writing, review & editing.

References

Ahuja LR, Ma L (eds) (2011) Methods of Introducing System Models into Agricultural Research.
American Society of Agronomy and Soil Science Society of America, Madison, WI, USA

Asseng S, Ewert F, Rosenzweig C, et al (2013) Uncertainty in simulating wheat yields under climate
change. Nat Clim Chang 3:827–832. doi: 10.1038/nclimate1916

Asseng S, Martre P, Maiorano A, et al (2019) Climate change impact and adaptation for wheat protein.
Glob Chang Biol 25:155–173. doi: 10.1111/gcb.14481

Bassu S, Brisson N, Durand J-L, et al (2014) How do various maize crop models vary in their responses to
climate change factors? Glob Chang Biol 20:2301–20. doi: 10.1111/gcb.12520

Brisson N, Gary C, Justes E, et al (2003) An overview of the crop model stics. Eur J Agron 18:309–332.
doi: 10.1016/S1161-0301(02)00110-7

Buis S, Lecharpentier P, Vezy R, et al (2021) SticsRPacks/CroptimizR: v0.4.0. doi:
10.5281/ZENODO.5121194

Chakrabarti A, Ghosh JK (2011) AIC, BIC and Recent Advances in Model Selection. Philos Stat 583–
605. doi: 10.1016/B978-0-444-51862-0.50018-6

Cleland EE, Chuine I, Menzel A, et al (2007) Shifting plant phenology in response to global change.
Trends Ecol. Evol. 22:357–365

Confalonieri R, Orlando F, Paleari L, et al (2016) Uncertainty in crop model predictions: What is the role
of users? Environ Model Softw 81:165–173. doi: 10.1016/j.envsoft.2016.04.009

Farina R, Sándor R, Abdalla M, et al (2021) Ensemble modelling, uncertainty and robust predictions of
organic carbon in long‐term bare‐fallow soils. Glob Chang Biol 27:904–928. doi:
10.1111/gcb.15441

Fath B, Jorgensen SE (2011) Fundamentals of ecological modelling: Applications in environmental
management and research. 4th edition. Elsevier, Amsterdam

Gao Y, Wallach D, Hasegawa T, et al (2021) Evaluation of crop model prediction and uncertainty using
Bayesian parameter estimation and Bayesian model averaging. Agric For Meteorol 311:108686. doi:
10.1016/J.AGRFORMET.2021.108686

Jägermeyr J, Müller C, Ruane AC, et al (2021) Climate impacts on global agriculture emerge earlier in
new generation of climate and crop models. Nat Food 2:873–885. doi: 10.1038/s43016-021-00400-y

Khorashadi Zadeh F, Nossent J, Woldegiorgis BT, et al (2022) A fast and effective parameterization of
water quality models. Environ Model Softw 149:105331. doi: 10.1016/J.ENVSOFT.2022.105331

Kobayashi K (2004) Comments on another way of partitioning mean squared deviation proposed by
Gauch et al. (2003). With reply. Agron J 96:1206–1207

Kuha J (2004) AIC and BIC: Comparisons of Assumptions and Performance. Sociol Methods Res
33:188–229. doi: 10.1177/0049124103262065

Lawes RA, Huth ND, Hochman Z (2016) Commercially available wheat cultivars are broadly adapted to
location and time of sowing in Australia's grain zone. Eur J Agron 77:38–46. doi:
10.1016/J.EJA.2016.03.009

Li T, Hasegawa T, Yin X, et al (2015) Uncertainties in predicting rice yield by current crop models under
a wide range of climatic conditions. Glob Chang Biol 21:1328–41. doi: 10.1111/gcb.12758

Martre P, Wallach D, Asseng S, et al (2015) Multimodel ensembles of wheat growth: many models are
better than one. Glob Chang Biol 21:911–25. doi: 10.1111/gcb.12768

McNunn G, Heaton E, Archontoulis S, et al (2019) Using a Crop Modeling Framework for Precision
Cost-Benefit Analysis of Variable Seeding and Nitrogen Application Rates. Front Sustain Food Syst
3:108. doi: 10.3389/fsufs.2019.00108

Menzel A, Yuan Y, Matiu M, et al (2020) Climate change fingerprints in recent European plant
phenology. Glob Chang Biol 26:2599–2612. doi: 10.1111/gcb.15000

Nelder JA, Mead R (1965) A Simplex Method for Function Minimization. Comput J 7:308–313. doi:
10.1093/comjnl/7.4.308

Piao S, Liu Q, Chen A, et al (2019) Plant phenology and global climate change: current progresses and
challenges. Glob Chang Biol gcb.14619. doi: 10.1111/gcb.14619

Rafiei V, Nejadhashemi AP, Mushtaq S, et al (2022) An improved calibration technique to address high
dimensionality and non-linearity in integrated groundwater and surface water models. Environ
Model Softw 149:105312. doi: 10.1016/J.ENVSOFT.2022.105312

Rezaei EE, Siebert S, Hüging H, Ewert F (2018) Climate change effect on wheat phenology depends on
cultivar change. Sci Rep 8:4891. doi: 10.1038/s41598-018-23101-2

Sisheber B, Marshall M, Mengistu D, Nelson A (2022) Tracking crop phenology in a highly dynamic
landscape with knowledge-based Landsat–MODIS data fusion. Int J Appl Earth Obs Geoinf
106:102670. doi: 10.1016/J.JAG.2021.102670

Stuble KL, Bennion LD, Kuebbing SE (2021) Plant phenological responses to experimental warming – A
synthesis. Glob Chang Biol gcb.15685. doi: 10.1111/gcb.15685

Tao F, Rötter RP, Palosuo T, et al (2018) Contribution of crop model structure, parameters and climate
projections to uncertainty in climate change impact assessments. Glob Chang Biol 24:1291–1307.
doi: 10.1111/gcb.14019

van Bussel LGJ, Stehfest E, Siebert S, et al (2015) Simulation of the phenological development of wheat
and maize at the global scale. Glob Ecol Biogeogr 24:1018–1029. doi: 10.1111/geb.12351

Wallach D (2011) Crop model calibration: A statistical perspective. Agron J 103:1144–1151

Wallach D, Martre P, Liu B, et al (2018) Multimodel ensembles improve predictions of crop-environment-
management interactions. Glob Chang Biol 24:5072–5083. doi: 10.1111/gcb.14411

Wallach D, Palosuo T, Thorburn P, et al (2021a) How well do crop modeling groups predict wheat
phenology, given calibration data from the target population? Eur J Agron 124:126195. doi:
https://doi.org/10.1016/j.eja.2020.126195

Wallach D, Palosuo T, Thorburn P, et al (2021b) Multi-model evaluation of phenology prediction for
wheat in Australia. Agric For Meteorol 298–299:108289. doi:
10.1016/J.AGRFORMET.2020.108289

Wallach D, Palosuo T, Thorburn P, et al (2021c) The chaos in calibrating crop models: Lessons learned
from a multi-model calibration exercise. Environ Model Softw 145:105206. doi:
10.1016/J.ENVSOFT.2021.105206

Wallach D, Palosuo T, Thorburn P, et al (2021d) Multi model evaluation of phenology prediction for
wheat in Australia. Agric For Meteorol in press: doi: 10.1101/2020.06.06.133504

Wang PC, Shoup TE (2011) Parameter sensitivity study of the Nelder–Mead Simplex Method. Adv Eng
Softw 42:529–533. doi: 10.1016/J.ADVENGSOFT.2011.04.004

Webber H, Lischeid G, Sommer M, et al (2020) No perfect storm for crop yield failure in Germany.
Environ Res Lett 15:. doi: 10.1088/1748-9326/aba2a4

Zhang L, Zhang Z, Tao F, et al (2022) Adapting to climate change precisely through cultivars renewal for
rice production across China: When, where, and what cultivars will be required? Agric For Meteorol
316:108856. doi: 10.1016/J.AGRFORMET.2022.108856

691