Enhancing associative learning in rats with a computationally designed training

protocol

Xu O. Zhang; Yili Zhang; Douglas S. Engelke; Claire E. Cho; Paul Smolen; John H

Byrne; Fabricio H. Do-Monte[#].

Department of Neurobiology and Anatomy, McGovern Medical School, The University of

Texas Health Science Center at Houston, Houston, Texas 77030

#Corresponding author:

Fabricio H. Do Monte, DVM, PhD.

Phone: 713-500-5613

Email: fabricio.h.domonte@uth.tmc.edu

Competing Interests: The authors declare no competing interests.

Author Contributions: X.O.Z, D.S.E., and C.E.C performed and analyzed the

behavioral experiments. Y.Z. implemented the computational model and ran all

simulations. P.S. helped design and implement the computational model. F.H.D-M and

J.H.B supervised and contributed to all aspects of this study. All the authors participated

in the design of the experiments. X.O.Z. and F.H.D-M interpreted the data and prepared

the manuscript with comments from all the co-authors.

Classification: Biological Sciences/Neuroscience

Keywords: Associative memory, fear conditioning, fear extinction, long-term

potentiation, learning model, spaced learning.

This PDF file includes:

Main Text

Figures 1 to 5, Supplementary Figures 1 to 3

Abstract

Associative learning requires the activation of protein kinases with distinct temporal

dynamics. Learning protocols with computationally designed intertrial intervals (ITIs)

that maximize the interaction between fast-activated protein kinase A (PKA) and slow-

activated extracellular signal-regulated kinase (ERK) enhance nonassociative learning

in *Aplysia*. Here, we tested whether an optimal learning protocol with irregular ITIs,

predicted computationally to increase the overlap between PKA and ERK signaling in

rat hippocampus, would enhance associative learning in mammals. We simulated

~1000 training protocols with irregular ITIs and identified an optimal protocol, predicted

to induce stronger associative learning than standard protocols with fixed ITIs. With

auditory fear conditioning, we showed that male adult rats exposed to the optimal

conditioning protocol exhibited stronger fear memory retrieval and impaired fear

memory extinction, compared to rats that received either massed or spaced

conditioning protocols with fixed ITIs. With fear extinction, we likewise observed that

fear conditioned rats exposed to the optimal extinction protocol showed improved

extinction of contextual fear memory, compared to rats that received standard extinction

protocols. Together, these findings demonstrate the capacity of a behavioral

intervention driven by a computational model of memory-related signaling pathways to

enhance associative learning in mammals and may provide greater insight into

strategies to improve cognition in humans.

Significance Statement

To test if associative learning in mammals can be enhanced by predetermined training

protocols, we used a computational model of the temporal dynamics of essential

biochemical cascades for learning and memory to predict an optimal training protocol in

rats. Using auditory fear conditioning and fear extinction tests, we demonstrated that the

optimal protocol enhanced the acquisition as well as the extinction of fear memories.

Our results suggest that a non-invasive behavioral intervention guided by computational

modeling can be used to enhance associative learning in mammals, with potential

clinical relevance for facilitating memory formation as well as enhancing extinction-

based therapies in humans.

MAIN TEXT

INTRODUCTION

Long-term memory (LTM) formation that lasts for days to years is believed to be

mediated by synaptic plasticity including long-term potentiation (LTP) or its invertebrate

analogue long-term facilitation (LTF), which require gene expression and protein

synthesis (1-5). Studies in the last decades have investigated LTP/LTF and their

underlying molecular processes as potential targets to enhance learning or restore

memory deficits in laboratory animals. However, traditional interventions using systemic

cognitive enhancers or intracerebral pharmacological manipulations (6-10) are either

based on trial-and-error approaches in specific model systems or are highly invasive,

making the translation of such interventions difficult for human applications. The recent

development of neural recording and optogenetic techniques has enabled precise

control of neurons based on their intrinsic firing rates in order to enhance learning or

modify memories (11-14), but the use of these techniques in humans remains

unforeseeable in the near future.

An alternative approach to enhance learning and memory is to develop computational

models to predict optimal training protocols based on the intracellular molecular

cascades that underlie LTM formation and LTP/LTF induction (15-17). Previous studies

have identified activation of PKA (18-20) and of the mitogen-activated protein kinase

(MAPK) isoform ERK (21-23) as essential cascades for LTF. These two pathways

converge to phosphorylate transcriptional factors such as cAMP responsive element

binding protein (CREB), which subsequently induce expression of multiple plasticity-

related genes (24-26). The observation that that these two pathways exhibit distinct

kinetics of activation (17, 20, 27), suggests that the temporal activity patterns, and

activation overlap, of these pathways may constitute an important target to enhance

associative learning. Accordingly, our previous study demonstrated that a

computationally designed protocol with irregular ITIs predicted to maximize the overlap

of PKA and ERK activities enhances LTF and nonassociative learning, specifically the

long-term sensitization of the tail elicited siphon-withdrawal reflex in *Aplysia* (28).

Substantial similarities between molecular processes of LTF in invertebrates and LTP in

mammals make it plausible that the same strategies used to enhance LTF and

nonassociative learning in invertebrates could enhance LTP and associative learning in

mammals. PKA activation in rodent hippocampus and amygdala is required for LTP and

LTM (29, 30). Similarly, activation of ERK/MAPK cascades and their cross-talk with PKA

kinase are required for the phosphorylation of CREB and LTP induction (31, 32). Albeit

with different kinetics than observed in invertebrates, PKA and ERK activation in

mammals also differ considerably in temporal dynamics (33-35), providing a similar

opportunity to maximize their overlap with irregular ITIs in an attempt to enhance

memory formation.

Here, we tested if the invertebrate LTF model (28) can be adapted to computationally

design an optimal associative learning protocol in mammals. Empirical data from the

literature were used to model PKA and ERK dynamics in rat hippocampus (33-35), a

critical brain region implicated in the formation of associative memories in mammals

(36-38). Then we simulated ~1000 different training protocols with irregular ITIs and

identified an optimal protocol, predicted to induce stronger memory than standard

training protocols with fixed ITIs. Using auditory fear conditioning and fear extinction

paradigms in rats, we confirmed that our computationally designed protocol enhanced

the acquisition as well as the extinction of fear memories in mammals. This strategy

may have potential clinical relevance for interventions aiming at facilitating memory

formation in psychiatric disorders associated with cognitive impairment in humans, as

well as to enhance extinction-based therapies in patients suffering from fear-related

disorders.

RESULTS

A computational model based on PKA and ERK dynamics in rats identified an

optimal partial fear conditioning protocol

Our model previously used in *Aplysia* (28) was adapted to simulate the dynamics of

PKA and ERK during the process of LTP induction in rat hippocampus (**Fig. 1A**). In this

model, the stimulus (*Stim*), which represents tetanic stimuli, rapidly activates PKA via

the cAMP pathway and slowly activates ERK via the Raf-MEK pathway. The synergistic

interaction between PKA and ERK pathways is quantified by a variable *inducer*, which

corresponds to the efficacy of the stimulus in inducing LTP. ERK kinetics were

described by differential equations (see Methods) with parameter values reproducing

empirical findings that ERK activity reaches peak levels 15~20 min after BDNF

treatment or tetanic stimuli in rat hippocampus acute slices (35, 39). Equations

describing PKA kinetics simulated data showing that PKA is transiently activated within

2 min after LTP induction in slices from rat hippocampus or within 5 min *in vivo* after

spatial discrimination task training (33, 34). We used this model to simulation fear

conditioning in rats, in which the pairing of a conditioned stimulus (CS) with an

unconditioned stimulus (US) is represented by *Stim*, and the conditioned responses are

proportional to the peak value of *inducer*. We observed that a single CS-US pairing

produced little overlap between PKA and ERK pathways (**Fig. 1B1**). We then simulated

~1000 partial fear conditioning protocols, with 4 trials and varying ITIs, to identify one

predicted to have optimal ITIs for triggering downstream gene activation. To compare,

we also simulated a Full Conditioning (FC) protocol with 8 trials and a fixed ITI of 270 s,

and a Partial Conditioning (PC) protocol with 4 trials and the same ITI of 270 s, which

resemble previous protocols used for full and partial fear conditioning in rats (40, 41).

Our simulation identified an Optimal Partial Conditioning (OPC) protocol with 4 trials and

irregular ITIs of 8, 8, and 16 min, which was able to produce higher peak *inducer* than

the standard PC protocol (**Fig. 1B2-1B4**). Comparing OPC and PC in Fig. 1B3-B4, both

protocols produced pERK curves with similar peak levels (orange). However, PKA

activity induced by the last trial of OPC had a much larger overlap with the pERK curve

than did PKA induced by PC. Therefore, OPC induced a much higher peak level of

inducer (purple). Based on our simulation, we predicted that the computationally

designed OPC protocol would produce stronger long-term memory in rats than the

standard PC protocol.

The optimal partial conditioning protocol induces stronger fear memory than a

standard partial conditioning protocol

To test if the computationally designed OPC protocol could enhance associative

learning in rats, we used a classical auditory fear conditioning paradigm. Because the

hippocampus is required for context encoding during fear conditioning (42-44), we used

two different chambers (context A and context B) to assess the contribution of the

context to the auditory fear memory. The two contexts differed in size, floor texture,

visual decoration of the walls, and odor cues (**Fig. 2A**). On Day 0, rats were pre-

exposed to context A for 20 min to familiarize with the chamber where fear conditioning

took place in the following day. After matching their baseline freezing and locomotor

activity during the familiarization session (**Fig. S1**), rats were assigned to one of the

three experimental groups described above in our simulation experiments: Full

Conditioning (FC), Partial Conditioning (PC), and Optimal Partial Conditioning (OPC).

On Day 1, rats were placed into context A and exposed to one nonreinforced

habituation tone (3 kHZ, 75 dB, 30 s) followed by distinct fear conditioning protocols (44

min duration, **Fig. 2B**). The FC group received eight presentations of CS (3 kHZ tone,

75 dB, 30 s) that co-terminated with an US (footshock, 0.7 mA, 0.5 s) with ITIs of 270 s.

The PC group received four similar CS-US pairings with the same ITIs of 270 s, and

remained in the chamber until the end of the session to equate the total context

exposure time. The OPC group received four similar CS-US pairings with ITIs of 8, 8,

and 16 min. All groups reached high levels of freezing (> 65%, **Fig. 2C**) and reduced

locomotion at the end of the fear acquisition session. On Day 2, rats were placed in

context B and given two CS presentations (ITI of 150 s, 7 min duration) in the absence

of US to test the retrieval of tone-associated fear memory in a novel context. Compared

to the FC group, both PC and OPC groups exhibited less freezing during the second CS

presentation (Two-way repeated measures (RM) ANOVA followed by Tukey's post-hoc

test. $F_{(2, 37)}$ = 3.751, P = 0.033; FC vs. OPC: p = 0.029; FC vs. PC: p = 0.002; OPC vs.

PC: p = 0.527). Nevertheless, both FC and OPC groups showed reduced average

speed during the retrieval test compared to the standard PC group, suggesting stronger

fear retrieval in a novel context (**Fig. 2D;** One-way ANOVA followed by Tukey's post

hoc test. $F_{(2, 37)}$ = 10.87, P < 0.001; FC vs. OPC: p = 0.131; FC vs. PC: p < 0.001; OPC

vs. PC: p = 0.024). On Day 3, rats were returned to context A for an extinction training

session where they received twelve CS presentations (ITIs of 150 s, 37 min duration).

Although no significant differences in freezing levels were observed among the groups

during each CS presentation (Two-way RM ANOVA. $F_{(22, 407)}$ = 1.169, P = 0.271), a

within session extinction analysis comparing the first four CSs to the last four CSs

revealed impaired extinction learning in the FC and OPC groups, when compared to the

PC group. This impairment in the acquisition of extinction in the FC and OPC groups

was characterized by sustained freezing levels across the session, which differed from

the PC group that showed reduced freezing levels from the first to the last CSs

presentation (**Fig. 2E left;** Paired Student's t test. FC: t_{13} = 2.153, p = 0.051; PC: t_{11} =

4.191, p = 0.001; OPC: t_{13} = 0.869, p = 0.400**)**. Interestingly, the OPC group exhibited

persistent freezing levels during the extinction training session as indicated by a

reduced relative change in freezing from the beginning to the end of the session, when

compared to the PC group (**Fig. 2E right;** One-way ANOVA followed by Tukey's post-

hoc test. $F_{(2, 37)}$ = 3.315, P = 0.047; FC vs. OPC: p = 0.487; FC vs. PC: p = 0.320; OPC

vs. PC: p = 0.037). Despite the persistent freezing levels across the extinction training

session observed in the OPC group, no significant differences were found between the

OPC and the PC groups during either the extinction retrieval tests performed on Days 4

and 5 (Two-way RM ANOVA. context A: $F_{(2, 37)}$ = 0.097, P = 0.907; context B: $F_{(2, 37)}$ =

2.984, P = 0.063) or the spontaneous recovery tests performed on Days 29 and 30 (**Fig.**

2C; Two-way RM ANOVA followed by Tukey's post hoc test. context A: $F_{(2, 37)}$ = 3.259,

P = 0.049; FC vs. OPC: p = 0.027; FC vs. PC: p = 0.038; OPC vs. PC: p = 0.999;

context B: $F_{(2, 37)}$ = 0.774, P = 0.469). Taken together, these data demonstrate that the

computationally designed training protocol (OPC) is able to enhance fear acquisition in

rats, thereby resulting in a stronger fear memory that is more resistant to extinction.

The optimal partial conditioning protocol induces stronger fear memory than a

spaced partial conditioning protocol

Previous studies have demonstrated that spaced learning protocols result in stronger

memories than standard (i.e., massed) learning protocols in both humans (45, 46) and

rodents (47-49, but also see 50). Thus, an alternative explanation for the augmented

fear memory observed in the OPC group in Fig. 2 could be simply a trial-spacing effect,

because our standard PC group used massed CS-US pairings. Both protocols produced

similar peak values of the pERK curves, but different overlap of PKA and ERK activities,

suggesting a way in which appropriately spaced protocols might advantageously

enhance overlap of kinase activities with distinct dynamics. Indeed, there was a broad

range of effective 4-trial protocols (**Fig. S2**), with one protocol with fixed 11 min 10 s ITIs

producing a nearly identical peak level of inducer as the OPC protocol, and a higher

peak level of inducer than other protocols with fixed ITIs from 1 min to 21 min (**Fig. 3A1,**

Fig. S2A). We termed this protocol the spaced partial conditioning (SPC) protocol. We

also examined the effects of different stimulus intensities on predicting optimal training

trials. The 1,000 different protocols were re-simulated with a reduced intensity of

stimulus. Interestingly, the protocol with 4 trials and irregular ITIs of 8, 8, and 16 min

still produced the greatest peak level of inducer (**Fig. 3A2, Fig. S2B**). Equally

interesting was that with weaker stimuli, the OPC protocol became relatively more

effective than the SPC protocol. Therefore, we predicted that empirically, the irregular

intervals of the OPC may produce more effective conditioning than the SPC protocol. To

test this hypothesis, we conducted an experiment to compare the OPC and SPC

protocols using a reduced shock intensity (0.5 mA instead of the 0.7 mA used in Fig. 2)

(**Fig. 3B**). Rats were assigned to OPC or SPC groups by matching their baseline

freezing and locomotor activity during the familiarization session (**Supplementary Fig.**

1B). During the fear acquisition, the SPC group received 4 CS-US pairings with ITIs of

11 min and 10 s, whereas the OPC group was exposed to the same protocol as in Fig.

2 (*i.e.*, 4 CS-US pairings with ITIs of 8, 8, and 16 min) (**Fig. 3C**). We found that the

difference in the ITIs between the OPC and SPC groups was sufficient to result in

distinct levels of fear memory acquisition. Rats exposed to OPC showed higher CS

freezing levels during the third CS-US pairing of the fear acquisition session (**Fig. 3D**

Day 1; Two-way RM ANOVA followed by Tukey's planned comparison. $F_{(4, 64)}$ = 1.550,

P = 0.339; OPC vs. SPC: p = 0.019). In addition, fear memory retrieval was increased

when rats in the OPC group were returned to context A for an extinction training session

on Day 3, as indicated by higher freezing during the first two CS presentations,

compared to the SPC group (**Fig. 3D Inset;** Welch's t test. $t_{12.43}$ = 2.227, p = 0.045).

Both OPC and SPC groups exhibited the same levels of freezing by the end of the

extinction training session (**Fig. 3D Day 3;** Two-way RM ANOVA followed by Tukey's

post hoc test. $F_{(11, 176)}$ = 2.997, P = 0.001; Tone 12, OPC vs. SPC: p > 0.999), as well as

during the extinction retrieval tests performed in context A (**Fig. 3D Day 4;** Two-way RM

ANOVA. $F_{(2, 37)}$ = 0.097, P = 0.907) or context B (**Fig. 3D Day 5;** Two-way RM ANOVA.

$F_{(2, 37)}$ = 2.984, P = 0.629). However, fear memory spontaneously recovered when rats

in the OPC group were retested in context A approximately 3 weeks later, as indicated

by higher freezing during the first CS presentation compared to SPC-trained rats (**Fig.**

3D Day 29; Two-way RM ANOVA followed by Tukey's planned comparison. $F_{(2, 32)} =$

1.809, P = 0.180; OPC vs. SPC: p = 0.030). In summary, these data suggest that the

enhancement in fear memory acquisition observed with our computationally designed

OPC protocol cannot simply be attributed to a trial-spacing effect or differences in the

delay to remove the animals from the chamber, but it is rather associated with the

maximized overlap between PKA and ERK signaling.

The computationally designed optimal partial conditioning protocol also

enhanced fear extinction

Extinction is a new learning that temporarily inhibits the initial associative memory (51,

52). Because previous studies suggest that the molecular mechanisms underlying fear

extinction learning are similar to those underlying fear acquisition (53-58), we

hypothesized that the computationally designed protocol would also enhance the

acquisition of extinction, suppressing the original fear memory. We therefore used the

OPC protocol with 4 trials and irregular ITIs of 8, 8, and 16 min as an Optimal Partial

Extinction (OPE) protocol (**Fig. 4B4**). We compared the OPE protocol with a Full

Extinction (FE, **Fig. 4B2**) protocol using 12 trials and ITIs of 150 s, and a Partial

Extinction (PE, **Fig. 4B3**) protocol using 4 trials and the same ITIs of 150 s, which

resemble previous protocols used for full and partial fear extinction in rats (59-61).

Similar to the simulation results for the conditioning protocols, the OPE protocol

triggered higher peak *inducer* than the PE protocol with equal ITIs of 150 s, and was

comparable to the FE protocol (**Fig. 4B**). We thus predicted the OPE protocol would

result in stronger extinction of fear memory than the standard PE protocol.

To test this prediction, we designed an experiment comparing the efficacy of different

extinction protocols (**Fig. 5A**). We used conditioned suppression of reward-seeking

behavior as an additional measure of fear memory because lever-press suppression is

more sensitive than freezing during fear extinction paradigms (62, 63). In addition, lever

presses help to maintain a constant level of activity so that freezing can be more reliably

measured along the session (64). Since lever presses for reward are trained in a

specific context, and extinction memory is context-dependent (65, 66), we used only

one context in this experiment so that lever press suppression can be consistently

assessed across sessions and the extinction memory can be retrieved after the

extinction learning. Rats were initially trained to press a lever to receive sucrose pellets

in a variable interval schedule of 60 s. After 7 days of training, rats reached the same

levels of lever pressing (~10 presses/min) and locomotor activity (**Supplementary Fig.**

1C). On Day 8, they underwent a fear conditioning session which included five

nonreinforced habituation tones (3 kHZ, 75 dB, 30 s duration) followed by seven CS-US

pairings (ITIs of 150 s, 37 min duration). In simulations, the OPE group showed peak

levels of *inducer* comparable to the FE group and higher than the PE group when strong

stimuli were used. We therefore increased the footshock intensity from 0.7 mA to 1.0

mA, which resulted in stronger fear acquisition and consequently higher freezing levels

during the extinction session. This stimulus intensity also helped to avoid a "floor effect"

in the following tests. On Day 9, rats were assigned to three experimental groups by

matching their freezing and lever presses during the fear conditioning session: Full

Extinction (FE), Partial Extinction (PE), and Optimal Partial Extinction (OPE) (One-way

ANOVA. $F_{(14, 231)}$ = 0.404, P = 0.973). On Day 10, rats were returned to the same

chamber for a fear extinction session (39 min duration). The FE group received twelve

CSs with ITIs of 150 s; the PE group received four CSs with ITIs of 150 s and remained

in the chamber until the end of the session; and the OPE group received four CSs with

ITIs of 8, 8, and 16 min (**Fig. 5B**). Our results revealed that at the end of the CS

presentations, freezing levels reduced from ~50% to ~20% in the FE groups, whereas

the PE and OPE groups maintained the same levels of freezing throughout the four CSs

(**Fig. 5C, Day 8**). Similarly, the FE group showed a significant increase in lever presses

(from 0.1 to 1.5 press/min) during the CS presentations from the beginning to the end of

the session, whereas CS lever presses remained unaltered in the PE and OPE groups

(**Supplementary Fig. 3A&B**; Paired Student's t test; FE: t_{11} = 2.837, p = 0.016; PE: t_{10}

= 1.000, p = 0.341; OPE: t_{12} = 1.237, p = 0.239). However, the OPE group showed a

significant increase in lever presses during the 30-s periods that preceded the CS

presentations (pre-CS) when comparing the first two pre-CS periods to the last pre-CS

period of the extinction training session (**Fig. 5D&E**; Paired Student's t test; FE: t_{11} =

0.994, p = 0.341; PE: t_{10} = 1.111, p = 0.293; OPE: t_{12} = 3.773, p = 0.003), suggesting

enhanced within-session extinction of contextual fear memory (67). On Days 10 and 35,

rats were returned to the same chamber and received two CSs with an ITI of 150 s to

test the strength of fear extinction memory during extinction retrieval and spontaneous

recovery tests, respectively. No significant difference in CS freezing or CS lever presses

were observed among the three groups during these tests (**Fig. 5C, Day 10&35;** Two-

way RM ANOVA. Day 10: Freezing, $F_{(2, 33)}$ = 0.207, P = 0.814; Lever Presses, $F_{(2, 33)}$ =

0.069, P = 0.933; Day 35: Freezing, $F_{(2, 33)}$ = 0.535, P = 0.591; Lever Presses, $F_{(2, 33)}$ =

1.085, P = 0.350). However, the OPE group showed increased pre-CS lever pressing

rate either in the first pre-CS period (**Fig. 5D;** Two-way RM ANOVA followed by Tukey's

planned comparison. $F_{(2, 33)}$ = 2.897, P = 0.069; Tone 1, a, FE vs. PE, p = 0.791; b, FE

vs. OPE, p = 0.025; c, OPE vs. PE, p = 0.005) or when averaging the two pre-CS

periods during the spontaneous recovery test (**Fig. 5F**; Kruskal-Wallis's test followed by

Dunn's post hoc test. $H(2)$ = 6.414, P = 0.041; a: FE vs. OPE, p = 0.531; b: FE vs. PE, p

= 0.709; c: OPE vs. PE, p = 0.035), when compared to the FE and PE groups on day

35. These results suggest that, although the tone-associated memory was similar

among the groups, our computationally designed OPE protocol was able to enhance the

extinction of contextual fear memory, as indicated by enhanced within-session

extinction of conditioned suppression and by reduced conditioned suppression

approximately 3 weeks after the extinction training session.

DISCUSSION

Studies on nonassociative learning in invertebrates have provided significant insights

about the molecular mechanisms of learning and memory due to their simplicity and

tractability (2, 5, 68). Long-term nonassociative learning in different regions of the

nervous system or across different species shares many intracellular molecular

cascades including cAMP-PKA and Raf-MEK-ERK signaling (21, 25). These two

pathways are also required for associative learning in invertebrates and mammals (29,

31, 69). The conservation of these mechanisms makes these molecular cascades

potential candidates for a universal intervention to enhance learning and memory. We

have previously demonstrated that a computationally designed protocol that maximizes

the interaction between PKA and ERK pathways enhances LTF and nonassociative

learning in *Aplysia* (28). Here, we extended the feasibility of this computational

approach to associative learning in mammals by adapting the simplified mathematical

model used in *Aplysia* to simulate the dynamics of PKA and ERK in rat hippocampus

based on empirical data available in the literature (33-35). After simulating ~1000

different training protocols of auditory fear conditioning, we identified an optimal protocol

with irregular ITIs which resulted in stronger fear acquisition and extinction-resistant

memory formation in rats, when compared to either massed or spaced training protocols

with equal ITIs. Using a separate fear extinction paradigm, we also showed that an

optimal training protocol was sufficient to induce enhanced extinction of contextual fear

memory in rats, when compared to standard or partial extinction protocols with equal

ITIs. Our results demonstrate the power of a simplified model of intracellular signaling

cascades in guiding associative learning across species, attesting the essential role of

the interaction between PKA and ERK pathways in both nonassociative and associative

learning. In addition, our findings highlight the potential clinical relevance of using

computational modeling to enhance associative learning in patients with memory

deficits, as well as to enhance extinction-based exposure therapies in patients with

anxiety disorders.

Pioneering studies have demonstrated that training protocols using spaced ITIs result in

stronger memory acquisition than those using standard massed ITIs, a well-established

phenomenon described as the "trial-spacing effect" (see review by 15). Data

demonstrating this effect in animals and humans illustrate the relationship between the

duration of the ITIs and the strength of memory formation as an inverted-U-shape curve

(70, 71), consistent with our simulated results (Fig. 1C). Previous findings have

attributed this spacing effect to the different efficacy of massed vs. spaced protocols in

inducing LTP/LTF through plasticity-related signaling. For instance, spaced training

protocols in rodents lead to increased activation of ERK (72), followed by CREB

phosphorylation and the expression of downstream genes, which have not been

reported following massed training protocols in the same species (73). A computational

model based on slice electrophysiology from rat hippocampal data has found that the

peak of ERK activation aligns with the optimal ITI for LTP induction (39). Although that

study provided a possible mechanism for the trial-spacing effect and a potential

approach to identify more effective protocols for associative learning, it relied on fixed

ITIs that are not necessarily the optimal intervals to induce synaptic plasticity and LTM.

In fact, our simulations showed that an optimal protocol with irregular ITIs was superior

to most spaced protocols with the same number of trials and equal ITIs. In the fear

conditioning experiment, the optimal protocol induced stronger and persistent fear

memory in rats, when compared to a spaced protocol predicted to be the most efficient

among protocols with equal ITIs. These results suggest that the trial-spacing effect in

mammals can be at least partially explained by enhanced overlap between PKA and

ERK pathways, which are critical for CREB activation (32). More importantly, our

findings demonstrate that protocols that include irregular ITIs can be a better approach

to enhance learning than previous protocols using equal ITIs.

The enhanced performance of our optimal protocol (OPC) compared to a spaced

protocol (SPC) during fear conditioning appears constrained by the intensity of stimuli.

The model predicts higher performance for OPC when weak stimuli are used, but

comparable performance when strong stimuli are used (**Fig. 3A**). Drawing a line to

distinguish weak and strong footshock intensities is not straightforward because the

relationship between fear memory and footshock intensity is neither monotonic nor

linear (74). Nevertheless, we observed clear differences in the efficacy of the optimal

fear conditioning protocol when using footshocks of different intensities. When we

compared the optimal partial conditioning protocol to a massed partial conditioning

protocol using a standard footshock intensity (0.7 mA, **Fig.2**) commonly used in

previous studies (30), we found only small differences in the conditioned responses (CS

freezing) between the two protocols. However, when we used a lower footshock

intensity (0.5 mA, **Fig.3**) to compare the optimal partial conditioning protocol to a

spaced partial conditioning protocol, we found a significant increase in CS freezing in

the OPC group during fear acquisition, retrieval, and spontaneous recovery tests.

Considering the robustness of the trial-spacing effect, it is possible that differences

between the OPC and the massed PC groups at the higher footshock intensity were

masked by a "ceiling effect". These data suggest that our approach could be more

beneficial for learning protocols relying on relatively weak stimuli. Additional studies

need to be conducted to understand the mechanism by which the intensity of stimuli

determines the enhanced performance of protocols with irregular ITIs compared to

protocols with fixed ITIs.

The association between the CS and US is primarily mediated by the lateral amygdala

where LTP induces enhanced CS responses (see review by 75, 76). Nevertheless,

another important and distinguishable component of fear memory formation is the

context in which the association has occurred. In the fear conditioning experiment,

enhanced CS freezing by our computationally designed protocol was only observed in

the same context where fear conditioning took place, suggesting that the memory

facilitation effect induced by our optimal protocol is context dependent. Similarly, in the

fear extinction experiment, enhancement by the optimal protocol was only observed

during the pre-CS lever pressing, a more sensitive index of contextual fear memory

during extinction and spontaneous recovery (62, 63, 67, 77, 78). These results are

consistent with the fact that the model is based on empirical data from rat hippocampus

currently available in the literature. The hippocampus plays major roles in context

encoding during fear conditioning, fear extinction, and the time-dependent

reappearance of fear following extinction training (*i.e.*, spontaneous recovery) (see

reviews by 44, 79). The observation that the temporal dynamics of ERK pathways differ

across brain regions involved in fear memory, with peaks occurring at 20 min after

stimulation in the hippocampus versus 60 min after fear conditioning in the lateral

amygdala (35, 80, 81) , suggests that the model may be able to predict optimal

protocols targeting specific components of fear memory based on the dynamics of

intracellular signaling cascades in distinct brain regions. Future studies will test this

possibility by modeling the molecular cascades in the lateral amygdala to preferentially

target the acquisition and extinction of CS-associated memories.

It is worth noting that the model is an extremely simplified one. We did not consider

many other important molecular cascades critical for LTP and memory formation. For

example, calcium/calmodulin-dependent protein kinase II (CaMKII) and protein kinase C

(PKC) are also required for LTP induction (16, 82, 83). In addition, we built the model

with empirical data of PKA and ERK dynamics from the literature, which are based on

ex vivo analyses and have limited temporal resolution. However, despite these

limitations, we demonstrated that the simplified model using essential biochemical

cascades for learning and memory was sufficient to enhance associative learning in

three different experiments in mammals. A more complex model that incorporates a

wider range of intracellular and extracellular processes based on *in vivo* real-time data

may strengthen the predictive ability of simulations. Further experiments should also

investigate whether the memory enhancing effects of our optimal protocol vary across

subjects of different sexes and ages, as well as its efficacy in animal models of cognitive

impairment. Together, our results suggest the possibility of using similar model-driven,

non-invasive behavioral approaches in preclinical and clinical studies aimed at

enhancing learning or restoring memory deficits in humans.

MATERIALS AND METHODS

Animals

All experimental procedures were approved by the Center for Laboratory Animal

Medicine and Care of The University of Texas Health Science Center at Houston.

National Institutes of Health guidelines for the care and use of laboratory animals were

strictly followed in order to minimize any potential discomfort and suffering. A total of

120 male Long-Evans hooded adult rats (Charles Rivers Laboratories) 3-5 months of

age and weighing 330-450 g were used. Rats were kept in a 12-hour light/12-hour dark

cycle with food and water *ad libitum*. Experiments were conducted during the light

phase. For the Optimal Extinction experiment, animals were maintained on a restricted

diet of 18 g per day of standard laboratory rat chow to increase their motivation during

the lever press training. Rats' weights were monitored weekly to make sure that all

animals maintained their weight under food restriction.

Model Development

The mathematical model for the activation of kinase cascades critical for long-term

memory (LTM) was modified from a previous model of the signaling cascades for the

induction of long-term synaptic facilitation (28). As mentioned above, the induction of LTM

for fear conditioning is known for requiring the activation of multiple kinase cascades with

different temporal dynamics. In the model, the cyclic AMP (cAMP)-PKA pathway is rapidly

activated after training, whereas the Raf- MEK-ERK pathway is slowly activated, and both

are required for the induction of LTM after fear conditioning and extinction (**Fig. 1A, Fig.**

4A).

PKA Pathway.   The dynamics of cAMP activation upstream of PKA following

training and the cAMP-dependent activation of PKA are described by Eqs. 1-4. Inactive

PKA is a holoenzyme (PKA$_{RC}$, Eq. 2), consisting of regulatory (PKA$_R$, Eq. 3) and catalytic

(PKA$_C$, Eq. 4) subunits.   In response to training, represented by the variable *Stim* in Eq.

1, cAMP is activated (Eq. 1). Active cAMP binds to the regulatory subunit of PKA, leading

to the release of free, active catalytic subunit (Fig. 1A) (Eqs. 2-4). '*Stim*' represents the

neurotransmitters released by training that function to activate kinase cascades.

490
$$\frac{d[cAMP]}{dt} = \lambda \frac{[Stim]}{[Stim] + K_{Stim}} - k_{b,cAMP}[cAMP] \qquad \text{(Eq. 1)}$$

491
$$\frac{d[PKA_{RC}]}{dt} = k_{b,PKA}[PKA_C][PKA_R] - k_{f,PKA}[PKA_{RC}][cAMP]^2 \qquad \text{(Eq. 2)}$$

492
$$\frac{d[PKA_R]}{dt} = k_{f,PKA}[PKA_{RC}][cAMP]^2 - k_{b,PKA}[PKA_C][PKA_R] \qquad \text{(Eq. 3)}$$

493
$$\frac{d[PKA_C]}{dt} = k_{f,PKA}[PKA_{RC}][cAMP]^2 - k_{b,PKA}[PKA_C][PKA_R] \qquad \text{(Eq. 4)}$$

Parameter values are modified from (28) to activate PKA immediately after training, but

with PKA activity quickly returning to the basal in ~5 min (Fig. 1B) (33, 34): λ = 14.6

μM/min, K_{STIM} = 85 μM, $k_{b,cAMP}$ = 4 min^{-1}, $k_{f,PKA}$ = 20 μM^{-2}min^{-1}, $k_{b,PKA}$ = 12

μM^{-1}min^{-1}

ERK Pathway. The activation of ERK by training '*Stim*' is via sequential activation

of the upstream kinases Raf and MEK (Fig. 1A). Raf activates the MAP kinase kinase

MEK, MEK in turn activates the MAP kinase ERK. The differential equations describing

the activation of Raf, MEK, and ERK (Eqs. 5-12) are similar to those in (28). However, a

discrete time delay in activation of Raf was removed from the phosphorylation of Raf (Eq.

5). Instead, based on empirical data (35, 39), parameters describing the activation of Raf

were adjusted from (28) so that the ERK activation curve reached the peak around 20

min post-training (Fig. 1B).

507
$$\frac{d[Raf^p]}{dt} = k_{f,Raf}[Raf][Stim] - k_{b,Raf}[Raf^p]$$
(Eq. 5)

508
$$[Raf] = [Raf]_{total} - [Raf^p]$$
(Eq. 6)

509
$$\frac{d[MEK]}{dt} = \frac{k_{b,MEK}[MEK^p]}{[MEK^p] + K_{MEK,2}} - \frac{k_{f,MEK}[Raf^p][MEK]}{[MEK] + K_{MEK,1}}$$
(Eq. 7)

510
$$\frac{d[MEK^{pp}]}{dt} = \frac{k_{f,MEK}[Raf^p][MEK^p]}{[MEK^p] + K_{MEK,1}} - \frac{k_{b,MEK}[MEK^{pp}]}{[MEK^{pp}] + K_{MEK,2}}$$
(Eq. 8)

511
$$[MEK^p] = [MEK]_{total} - [MEK] - [MEK^{pp}]$$
(Eq. 9)

512
$$\frac{d[ERK]}{dt} = \frac{k_{b,ERK}[ERK^p]}{[ERK^p] + K_{ERK,2}} - \frac{k_{f,ERK}[MEK^{pp}][ERK]}{[ERK] + K_{ERK,1}}$$
(Eq. 10)

513
$$\frac{d[ERK^{pp}]}{dt} = \frac{k_{f,ERK}[MEK^{pp}][ERK^p]}{[ERK^p] + K_{ERK,1}} - \frac{k_{b,ERK}[ERK^{pp}]}{[ERK^{pp}] + K_{ERK,2}}$$
(Eq. 11)

514
$$[ERK^p] = [ERK]_{total} - [ERK] - [ERK^{pp}]$$
(Eq. 12)

Parameter values: $k_{f,Raf}$ = 0.001 $\mu M^{-1}min^{-1}$, $k_{b,Raf}$ = 0.05 min^{-1}, $[Raf]_{total}$ = 0.5 μM,

$k_{f,MEK}$ = 0.41 min^{-1}, $k_{b,MEK}$ = 0.04 $\mu M/min$, $K_{MEK,1}$ = 0.20 μM,

$K_{MEK,2}$ = 0.19 μM, $[MEK]_{total}$ = 0.5 μM, $k_{f,ERK}$ = 0.41 min⁻¹,

$k_{b,ERK}$ = 0.12 μM/min, $K_{ERK,1}$ = 0.19 μM,

$K_{ERK,2}$ = 0.21 μM, $[ERK]_{total}$ = 0.5 μM.

As in (28), a variable '*inducer*' was used to quantify the overlap of activation

between PKA and ERK, which together regulate the gene expression necessary for the

induction of LTM.

523                    $inducer = k_{inducer}\ [PKA_C]\ [ERK^{pp}]$ (Eq. 13)

where $k_{inducer}$ = 1 μM⁻¹.

To determine which protocols could more effectively activate *inducer,* four-trial

protocols with three ITIs, each ranging from 2–20 min in steps of 2 min, were simulated.

Combining these permutations yielded 10^3 protocols. For each protocol, the maximal

(peak) overlap between PKA and ERK was quantified as the peak level of *inducer*

produced. Also, the value of *Stim* was varied from 100 to 300 μM to represent weak and

strong trainings and to test the robustness of protocols. Based on the maximal overlap,

a protocol with ITIs of 8, 8, and 16 min was selected and was denoted optimal partial

conditioning (OPC). We assumed that fear conditioning and extinction activate the same

signaling pathways, so we predicted this protocol would similarly enhance extinction.

<u>Numerical methods.</u>   Fourth-order Runge-Kutta integration was used for

integration of all differential equations with a time step of 3 s. Further time step reduction

did not lead to significant improvement in accuracy. The steady-state levels of variables

were determined after at least one simulated day, prior to any manipulations. The model

was programmed in XPPAUT (http://www.math.pitt.edu/~bard/xpp/xpp.html) (84) and

simulated on Dell Precision T1700 microcomputers.

Behavioral Tasks

Optimal Conditioning

Apparatuses

Two distinct chambers (context A and context B) positioned inside sound attenuating

boxes were used during the Optimal Conditioning experiments. Context A consisted of a

small operant chamber (34 cm high x 25 cm wide x 23 cm deep, 200 lux, Med

Associates, see schematic drawing in Fig. 2A top) with one of the two aluminum walls

covered by black adhesive paper and two transparent acrylic walls, and a metal grid

floor beneath which a microcentrifuge tube (Eppendorf) containing 50 µl of 10% amyl

acetate (Sigma-Aldrich) was positioned. Context B consisted of a larger acrylic operant

chamber (40 cm high x 50 cm wide x 26 cm deep, 20 lux, Med Associates, see

schematic drawing in Fig. 2A bottom) with one of its walls covered by a black and white

stripped paper, and a floor made of a white acrylic board beneath which a

microcentrifuge tube (Eppendorf) containing 50 µl of deionized water was positioned.

Procedures

On day 0, rats were placed in context A for a 20-min familiarization session. Next, rats

were assigned to three experimental groups by matching their baseline freezing and

locomotor activity during the familiarization session: Full Conditioning (FC), Partial

Conditioning (PC), and Optimal Partial Conditioning (OPC). On day 1, rats were placed

into context A and exposed to one nonreinforced habituation tone (3 kHz, 75 dB, 30 s)

followed by distinct fear conditioning protocols (44 min duration). The FC group received

eight presentations of a conditioned stimulus (CS, 3 kHz tone, 75 dB, 30 s) that co-

terminated with an unconditioned stimulus (US, footshock, 0.7 mA, 0.5 s), with fixed ITIs

of 270 s. The PC group received four similar CS-US pairings with the same ITIs of 270

s, and remained in the chamber until the end of the session. The OPC group received

four similar CS-US pairings with ITIs of 8, 8, and 16 min. On day 2, rats were placed in

context B and given two CS presentations (ITI of 150 s, 7 min duration) in the absence

of US to test the retrieval of tone-associated fear memory in a novel context. On day 3,

rats were returned to context A for an extinction training session where they received

twelve CS presentations (ITIs of 150 s, 37 min duration). The first four CSs were

compared to the last four CSs to assess extinction learning within the same session. On

day 4 and 5, rats were placed back in context A and B, respectively, and exposed to an

extinction retrieval session (similar to day 2). On day 29 and 30, rats were placed back

in context A and B, respectively, and exposed to a spontaneous recovery session

(similar to day 2). On day 1 and day 2, rats in the same group were trained

simultaneously in four chambers with the group order counterbalanced to avoid

interference from different protocols in neighboring chambers. On the following days,

rats were simultaneously tested in four chambers regardless of the group assignment.

Each rat was tested in the exact same chamber across the days. Footshocks, tones,

intertrial intervals, and session duration were controlled by an automated video tracking

system (ANY-maze, Stoelting), which also quantified the percentage of time freezing,

distance traveled, average speed and maximum speed. All rats passed the criteria of

20% of freezing during at least one CS of the fear conditioning session and the first two

CSs of the extinction session.

Optimal vs. Spaced Partial Conditioning

Apparatus and Procedures

The same apparatus and contexts described above were used. Following familiarization

on Day 0, rats were randomly assigned to two groups: Optimal Partial Conditioning

(OPC) and Spaced Partial Conditioning (SPC). The same procedures described above

were used here, except that: *i)* the footshock intensity was reduced from 0.7 mA to 0.5

mA to decrease the possibility of ceiling effects; *ii)* the SPC group received four CS-US

pairings with ITIs of 11 min and 10 s. The same automated video tracking system (ANY-

maze, Stoelting) described above was used for protocol control and behavioral

quantification. One rat that failed the criteria of 20% of freezing during at least one CS of

the fear conditioning session and the first two CSs of the extinction session was

excluded from the analyses.

Optimal Extinction

Lever-press training

Rats were placed in an acrylic/aluminum operant chamber (34 cm high x 25 cm wide x

23 cm deep, Med Associates, see schematic drawing in Fig. 4A) and trained to press a

lever for sucrose on a fixed ratio of one pellet for each press. Next, animals were trained

in a variable interval schedule of reinforcement that was gradually reduced across the

days (one pellet every 15 s, 30 s, or 60 s) until they reached a minimum criterion of 10

presses/min after 7 days of training. All sessions lasted 30 min and were performed on

consecutive days. Sucrose pellet delivery, variable intervals, and session duration were

controlled by an automated video tracking system (ANY-maze, Stoelting).

Apparatus and Procedure

On day 8, rats were placed into the same chamber where they had previously

undergone lever presses training. Animals were exposed to five nonreinforced

habituation tones (3 kHZ, 75 dB, 30 s duration) followed by seven CS-US pairings (ITIs

of 150 s, 37 min duration). The footshock intensity was increased from 0.7 mA to 1.0-

1.2 mA to result in stronger fear acquisition and consequently higher freezing levels

during the extinction training session in the next day. Rats were assigned to three

experimental groups by matching their freezing and lever presses during the fear

conditioning session: Full Extinction (FE), Partial Extinction (PE), and Optimized Partial

Extinction (OPE). On day 9, rats were returned to the same chamber for a fear

extinction session (39 min duration). The FE group received twelve CSs with ITIs of 150

s; the PE group received four CSs with ITIs of 150 s and remained in the chamber until

the end of the session; and the OPE group received four CSs with ITIs of 8, 8, and 16

min. On day 10 and 35, rats were placed into the same chamber and received two CSs

with an ITI of 150 s to test the strength of fear extinction memory during extinction

retrieval and spontaneous recovery tests, respectively. The same automated video

tracking system (ANY-maze, Stoelting) described above was used for protocol control

and behavioral quantification. Two rats that never reached the criterium of 20% of

freezing during at least one CS of the fear conditioning session and the first two CSs of

the extinction session were excluded from the analyses.

Quantification and statistical analysis

Rats were recorded with digital video cameras (Logitech C920) and behavioral indices were measured using automated video-tracking system (ANY-maze). Lever presses per minute were calculated by measuring the number of presses during the 30 s cue (or 30 s pre-cue) multiplied by two. All graphics and numerical values reported in the figures are presented as mean ± standard error of the mean (SEM). Grubbs' tests were used to identify outliers ($p < 0.05$) in each experiment and one rat in Fig. 2 was removed after identified by the test. Shapiro-Wilk test were performed to determine whether the data have a normal distribution. F tests before pair-wise comparison and Brown-Forsythe tests before multiple group comparison were performed to test if groups have the same variance. Statistical significance for parametric distributions was determined with paired Student's t test, Welch's t test, one-way analysis of variance (ANOVA) or two-way repeated-measures ANOVA followed by Tukey post-hoc comparisons, whereas Kruskal-Wallis test followed by Dunn's post-hoc test was used for non-parametric distributions (Prism 7), as indicated for each experiment. Sample size was based on estimations from previous literature and experience.

Data availability

All data that support the findings presented in this study are available from the corresponding author on reasonable request. Source codes will be submitted to the ModelDB database (85), and to GitHub (https://github.com/Owenxz/Zhang-XO-Enhanced-Learning-2022.git).

ACKNOWLEDGEMENT

We thank Nikita Watson and Sharon Gordon for their technical and administrative

assistance. We also thank current members of the Byrne and Do Monte and Labs for

their valuable comments on the manuscript. This work was supported by NIH grant

R01-NS102490 to J.H.B, and NIH grant R00-MH105549, NIH grant R01-MH120136, a

Brain & Behavior Research Foundation grant (NARSAD Young Investigator), and a

Rising STARs Award from UT System to F.H.D-M.

REFERENCES

1.   S. J. Martin, P. D. Grimwood, R. G. Morris, Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* **23**, 649-711 (2000).

2.   E. R. Kandel, The molecular biology of memory storage: a dialogue between genes and synapses. *Science* **294**, 1030-1038 (2001).

3.   M. A. Lynch, Long-term potentiation and memory. *Physiol Rev* **84**, 87-136 (2004).

4.   C. M. Alberini, Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev* **89**, 121-145 (2009).

5.   J. H. Byrne, R. D. Hawkins, Nonassociative learning in invertebrates. *Cold Spring Harb Perspect Biol* **7** (2015).

6.   S. Sharif, A. Guirguis, S. Fergus, F. Schifano, The use and impact of cognitive enhancers among university students: a systematic review. *Brain Sci* **11** (2021).

7.   J. C. Lauterborn *et al.*, Chronic ampakine treatments stimulate dendritic growth and promote learning in middle-aged rats. *J Neurosci* **36**, 1636-1646 (2016).

8.   J. L. McGaugh, L. F. Petrinovich, Effects of drugs on learning and memory. *Int Rev Neurobiol* **8**, 139-196 (1965).

9.   G. Lynch, C. D. Cox, C. M. Gall, Pharmacological enhancement of memory or cognition in normal subjects. *Front Syst Neurosci* **8**, 90 (2014).

10.   S. M. Fernandez *et al.*, Estradiol-induced enhancement of object memory consolidation involves hippocampal extracellular signal-regulated kinase activation and membrane-bound estrogen receptors. *J Neurosci* **28**, 8660-8667 (2008).

11.   M. T. Brown *et al.*, Ventral tegmental area GABA projections pause accumbal

cholinergic interneurons to enhance associative learning. *Nature* **492**, 452-456

(2012).

12.   S. Nabavi *et al.*, Engineering a memory with LTD and LTP. *Nature* **511**, 348-352

(2014).

13.   K. Lee *et al.*, Parvalbumin interneurons modulate striatal output and enhance

performance during associative learning. *Neuron* **93**, 1451-1463 e1454 (2017).

14.   X. Liu *et al.*, Optogenetic stimulation of a hippocampal engram activates fear

memory recall. *Nature* **484**, 381-385 (2012).

15.   P. Smolen, Y. Zhang, J. H. Byrne, The right time to learn: mechanisms and

optimization of spaced learning. *Nat Rev Neurosci* **17**, 77-88 (2016).

16.   P. Smolen, D. A. Baxter, J. H. Byrne, Comparing theories for the maintenance of

late LTP and long-term memory: computational analysis of the roles of kinase

feedback pathways and synaptic reactivation. *Front Comput Neurosci* **14**,

569349 (2020).

17.   Y. Zhang, P. D. Smolen, L. J. Cleary, J. H. Byrne, Quantitative description of the

interactions among kinase cascades underlying long-term plasticity of *Aplysia*

sensory neurons. *Sci Rep* **11**, 14931 (2021).

18.   S. Schacher, V. F. Castellucci, E. R. Kandel, cAMP evokes long-term facilitation

in *Aplysia* sensory neurons that requires new protein synthesis. *Science* **240**,

1667-1669 (1988).

19. B. A. Goldsmith, T. W. Abrams, Reversal of synaptic depression by serotonin at

Aplysia sensory neuron synapses involves activation of adenylyl cyclase. *Proc*

Natl Acad Sci U S A **88**, 9021-9025 (1991).

20. U. Muller, T. J. Carew, Serotonin induces temporally and mechanistically distinct

phases of persistent PKA activity in *Aplysia* sensory neurons. *Neuron* **21**, 1423-

1434 (1998).

21. K. C. Martin *et al.*, MAP kinase translocates into the nucleus of the presynaptic

cell and is required for long-term facilitation in *Aplysia*. *Neuron* **18**, 899-912

(1997).

22. S. K. Sharma, T. J. Carew, The roles of MAPK cascades in synaptic plasticity

and memory in *Aplysia*: facilitatory effects and inhibitory constraints. *Learn Mem*

11, 373-378 (2004).

23. S. K. Sharma *et al.*, Differential role of mitogen-activated protein kinase in three

distinct phases of memory for sensitization in *Aplysia*. *J Neurosci* **23**, 3899-3907

(2003).

24. P. K. Dash, B. Hochner, E. R. Kandel, Injection of the cAMP-responsive element

into the nucleus of *Aplysia* sensory neurons blocks long-term facilitation. *Nature*

345, 718-721 (1990).

25. D. Bartsch *et al.*, Aplysia CREB2 represses long-term facilitation: relief of

repression converts transient facilitation into long-term functional and structural

change. *Cell* **83**, 979-992 (1995).

26. R. Y. Liu, L. J. Cleary, J. H. Byrne, The requirement for enhanced CREB1

expression in consolidation of long-term synaptic facilitation and long-term

excitability in sensory neurons of *Aplysia*. *J Neurosci* **31**, 6871-6879 (2011).

27. G. T. Philips, E. I. Tzvetkova, T. J. Carew, Transient mitogen-activated protein

kinase activation is confined to a narrow temporal window required for the

induction of two-trial long-term memory in *Aplysia*. *J Neurosci* **27**, 13701-13705

(2007).

28. Y. Zhang *et al.*, Computational design of enhanced learning protocols. *Nat*

Neurosci **15**, 294-297 (2011).

29. G. E. Schafe, J. E. LeDoux, Memory consolidation of auditory pavlovian fear

conditioning requires protein synthesis and protein kinase A in the amygdala. *J*

Neurosci **20**, RC96 (2000).

30. T. Abel *et al.*, Genetic demonstration of a role for PKA in the late phase of LTP

and in hippocampus-based long-term memory. *Cell* **88**, 615-626 (1997).

31. J. P. Adams, J. D. Sweatt, Molecular psychology: roles for the ERK MAP kinase

cascade in memory. *Annu Rev Pharmacol Toxicol* **42**, 135-163 (2002).

32. S. Impey *et al.*, Cross talk between ERK and PKA is required for Ca^{2+} stimulation

of CREB-dependent transcription and ERK nuclear translocation. *Neuron* **21**,

869-883 (1998).

33. E. D. Roberson, J. D. Sweatt, Transient activation of cyclic AMP-dependent

protein kinase during hippocampal long-term potentiation. *J Biol Chem* **271**,

30436-30441 (1996).

34.   S. I. Vázquez, A. Vázquez, S. Peña de Ortiz, Different hippocampal activity

profiles for PKA and PKC in spatial discrimination learning. *Behavioral*

Neuroscience **114**, 1109-1118 (2000).

35.   Y. Wang *et al.*, A molecular brake controls the magnitude of long-term

potentiation. *Nat Commun* **5**, 3051 (2014).

36.   B. Milner, L. R. Squire, E. R. Kandel, Cognitive neuroscience and the study of

memory. *Neuron* **20**, 445-468 (1998).

37.   A. Mayes, D. Montaldi, E. Migo, Associative memory and the medial temporal

lobes. *Trends Cogn Sci* **11**, 126-135 (2007).

38.   K. Nakazawa *et al.*, Requirement for hippocampal CA3 NMDA receptors in

associative memory recall. *Science* **297**, 211-218 (2002).

39.   S. M. Ajay, U. S. Bhalla, A role for ERKII in synaptic pattern selectivity on the

time-scale of minutes. *Eur J Neurosci* **20**, 2671-2680 (2004).

40.   J. A. Detert, N. D. Kampa, J. R. Moyer, Jr., Differential effects of training intertrial

interval on acquisition of trace and long-delay fear conditioning in rats. *Behav*

Neurosci **122**, 1318-1327 (2008).

41.   T. B. Lonsdorf *et al.*, Don't fear 'fear conditioning': Methodological considerations

for the design and analysis of studies on human fear acquisition, extinction, and

return of fear. *Neurosci Biobehav Rev* **77**, 247-285 (2017).

42.   R. G. Phillips, J. E. LeDoux, Differential contribution of amygdala and

hippocampus to cued and contextual fear conditioning. *Behav Neurosci* **106**,

274-285 (1992).

43.   N. R. Selden, B. J. Everitt, L. E. Jarrard, T. W. Robbins, Complementary roles for

the amygdala and hippocampus in aversive conditioning to explicit and

contextual cues. *Neuroscience* **42**, 335-350 (1991).

44.   S. Maren, K. L. Phan, I. Liberzon, The contextual brain: implications for fear

conditioning, extinction and psychopathology. *Nat Rev Neurosci* **14**, 417-428

(2013).

45.   N. J. Cepeda, H. Pashler, E. Vul, J. T. Wixted, D. Rohrer, Distributed practice in

verbal recall tasks: A review and quantitative synthesis. *Psychol Bull* **132**, 354-

380 (2006).

46.   M. Raman *et al.*, Teaching in small portions dispersed over time enhances long-

term knowledge retention. *Med Teach* **32**, 250-255 (2010).

47.   M. J. Anderson, S. A. Jablonski, D. B. Klimas, Spaced initial stimulus

familiarization enhances novelty preference in Long-Evans rats. *Behav*

Processes **78**, 481-486 (2008).

48.   L. Jiang *et al.*, Spaced training enhances contextual fear memory via activating

hippocampal 5-HT2A receptors. *Front Mol Neurosci* **12**, 317 (2019).

49.   M. T. Scharf *et al.*, Protein synthesis is required for the enhancement of long-

term potentiation and long-term memory by spaced training. *J Neurophysiol* **87**,

2770-2777 (2002).

50.   C. K. Cain, A. M. Blouin, M. Barad, Temporally massed CS presentations

generate more fear extinction than spaced presentations. *J Exp Psychol Anim*

Behav Process **29**, 323-333 (2003).

51.   L. de Oliveira Alvares, F. H. Do-Monte, Understanding the dynamic and destiny

of memories. *Neurosci Biobehav Rev* **125**, 592-607 (2021).

52.   G. J. Quirk, D. Mueller, Neural mechanisms of extinction learning and retrieval.

Neuropsychopharmacology **33**, 56-72 (2008).

53.   D. E. Berman, Y. Dudai, Memory extinction, learning anew, and learning the new:

dissociations in the molecular machinery of learning in cortex. *Science* **291**,

2417-2419 (2001).

54.   J. F. Flood, M. E. Jarvik, E. L. Bennett, A. E. Orme, M. R. Rosenzweig, Protein

synthesis inhibition and memory for pole jump active avoidance and extinction.

Pharmacol Biochem Behav **7**, 71-77 (1977).

55.   J. Radulovic, N. C. Tronson, Molecular specificity of multiple hippocampal

processes governing fear extinction. *Rev Neurosci* **21**, 1-17 (2010).

56.   E. Santini, R. U. Muller, G. J. Quirk, Consolidation of extinction learning involves

transfer from NMDA-independent to NMDA-dependent memory. *J Neurosci* **21**,

9009-9017 (2001).

57.   G. Szapiro, M. R. Vianna, J. L. McGaugh, J. H. Medina, I. Izquierdo, The role of

NMDA glutamate receptors, PKA, MAPK, and CAMKII in the hippocampus in

extinction of conditioned fear. *Hippocampus* **13**, 53-58 (2003).

58.   N. C. Tronson *et al.*, Regulatory mechanisms of fear extinction and depression-

like behavior. *Neuropsychopharmacology* **33**, 1570-1583 (2008).

59.   E. Santini, H. Ge, K. Ren, S. Pena de Ortiz, G. J. Quirk, Consolidation of fear

extinction requires protein synthesis in the medial prefrontal cortex. *J Neurosci*

24, 5704-5710 (2004).

60.   A. Holmes, G. J. Quirk, Pharmacological facilitation of fear extinction and the

819   search for adjunct treatments for anxiety disorders--the case of yohimbine.

820   *Trends Pharmacol Sci* **31**, 2-7 (2010).

821   61.   F. H. Do-Monte, G. Manzano-Nieves, K. Quinones-Laracuente, L. Ramos-

822   Medina, G. J. Quirk, Revisiting the role of infralimbic cortex in fear extinction with

823   optogenetics. *J Neurosci* **35**, 3607-3615 (2015).

824   62.   M. Mast, R. J. Blanchard, D. C. Blanchard, The relationship of freezing and

825   response suppression in a CER situation. *The Psychological Record* **32**, 151

826   (1982).

827   63.   F. Sotres-Bayon, D. Sierra-Mercado, E. Pardilla-Delgado, G. J. Quirk, Gating of

828   fear in prelimbic cortex by hippocampal and amygdala inputs. *Neuron* **76**, 804-

829   812 (2012).

830   64.   G. J. Quirk, G. K. Russo, J. L. Barron, K. Lebron, The role of ventromedial

831   prefrontal cortex in the recovery of extinguished fear. *J Neurosci* **20**, 6225-6231

832   (2000).

833   65.   C. A. Podlesnik, M. E. Kelley, C. Jimenez-Gomez, M. E. Bouton, Renewed

834   behavior produced by context change and its implications for treatment

835   maintenance: A review. *J Appl Behav Anal* **50**, 675-697 (2017).

836   66.   T. D. Goode, S. Maren, Animal models of fear relapse. *ILAR J* **55**, 246-258

837   (2014).

838   67.   M. A. Morgan, J. E. LeDoux, Differential contribution of dorsal and ventral medial

839   prefrontal cortex to the acquisition and extinction of conditioned fear in rats.

840   *Behav Neurosci* **109**, 681-688 (1995).

68. T. J. Carew, C. L. Sahley, Invertebrate learning and memory: from behavior to

molecules. *Annu Rev Neurosci* **9**, 435-487 (1986).

69. R. D. Hawkins, J. H. Byrne, Associative learning in invertebrates. *Cold Spring*

Harb Perspect Biol **7** (2015).

70. P. P. Verkoeijen, R. M. Rikers, H. G. Schmidt, Limitations to the spacing effect:

demonstration of an inverted u-shaped relationship between interrepetition

spacing and free recall. *Exp Psychol* **52**, 257-263 (2005).

71. P. B. Barela, Theoretical mechanisms underlying the trial-spacing effect in

Pavlovian fear conditioning. *J Exp Psychol Anim Behav Process* **25**, 177 (1999).

72. S. M. Ajay, U. S. Bhalla, A propagating ERKII switch forms zones of elevated

dendritic activation correlated with plasticity. *HFSP J* **1**, 49-66 (2007).

73. D. Genoux *et al.*, Protein phosphatase 1 is a molecular constraint on learning

and memory. *Nature* **418**, 970-975 (2002).

74. M. Davis, D. I. Astrachan, Conditioned fear and startle magnitude: effects of

different footshock or backshock intensities used in training. *J Exp Psychol Anim*

Behav Process **4**, 95-103 (1978).

75. S. Maren, Synaptic mechanisms of associative memory in the amygdala. *Neuron*

47, 783-786 (2005).

76. J. P. Johansen, C. K. Cain, L. E. Ostroff, J. E. LeDoux, Molecular mechanisms of

fear learning and memory. *Cell* **147**, 509-524 (2011).

77. N. Padilla-Coreano, F. H. Do-Monte, G. J. Quirk, A time-dependent role of

midline thalamic nuclei in the retrieval of fear memory. *Neuropharmacology* **62**,

457-463 (2012).

78. A. M. Woods, M. E. Bouton, Immediate extinction causes a less durable loss of

performance than delayed extinction following either fear or appetitive

conditioning. *Learn Mem* **15**, 909-920 (2008).

79. M. E. Bouton, S. Maren, G. P. McNally, Behavioral and neurobiological

mechanisms of Pavlovian and instrumental extinction learning. *Physiol Rev* **101**,

611-681 (2021).

80. G. E. Schafe *et al.*, Activation of ERK/MAP kinase in the amygdala is required for

memory consolidation of pavlovian fear conditioning. *J Neurosci* **20**, 8177-8187

(2000).

81. B. Di Benedetto *et al.*, Activation of ERK/MAPK in the lateral amygdala of the

mouse is required for acquisition of a fear-potentiated startle response.

Neuropsychopharmacology **34**, 356-366 (2009).

82. R. C. Malenka *et al.*, An essential role for postsynaptic calmodulin and protein

kinase activity in long-term potentiation. *Nature* **340**, 554-557 (1989).

83. S. Wang, T. Sheng, S. Ren, T. Tian, W. Lu, Distinct roles of PKCι/λ and PKMζ in

the initiation and maintenance of hippocampal long-term potentiation and

memory. *Cell Rep* **16**, 1954-1961 (2016).

84. B. Ermentrout, *Simulating, analyzing, and animating dynamical systems : a guide

to XPPAUT for researchers and students*, Software, environments, tools (Society

for Industrial and Applied Mathematics, Philadelphia, 2002), pp. xiv, 290 p.

85. R. A. McDougal, T. M. Morse, M. L. Hines, G. M. Shepherd, ModelView for

ModelDB: online presentation of model structure. *Neuroinformatics* **13**, 459-470

(2015).

Figure 1. Computational simulations of PKA and ERK pathways predict an optimal protocol for

fear conditioning. A, Schematic of the model. Stimulus (*Stim*) activates PKA via cAMP and activates

ERK via Raf-MEK. The variable *inducer* quantifies the PKA/ERK interaction (activity overlap). **B1**,

Simulated time courses of activated ERK (pERK, orange traces, μM) and activated PKA (PKAc, green

traces) in response to one trial of *Stim* (μM). **B2**, Simulated time courses of pERK (orange traces, μM),

PKAc (green traces) and *inducer* (red traces, nM) in response to an 8-trial protocol with regular ITIs of 4.5

min (full conditioning, FC). **B3**, Simulated time courses of pERK (orange traces), PKAc (green traces),

and *inducer* (red traces) in response to a 4-trial protocol with regular ITIs of 4.5 min (partial conditioning,

PC). **B4**, Simulated time courses of pERK (orange traces), PKAc (green traces) and *inducer* (red traces)

in response to a 4-trial protocol with computationally designed intervals (optimal partial conditioning,

OPC).

Figure 2. Computationally designed protocol enhanced fear conditioning in rats. **A,** Schematics of

the enhanced fear conditioning experimental procedures. Top: tests that were conducted in Context A.

Bottom: tests that were conducted in Context B. **B,** Schematics of the fear conditioning protocols for full

conditioning (FC, n = 14), partial conditioning (PC, n = 12), and optimal partial conditioning groups (OPC,

n = 14). Following a habituation (Hab.) tone, rats received multiple trials of a conditioned stimulus (CS, 3

kHz tone, 30 s) that co-terminated with an unconditioned stimulus (US, footshock, 0.7 mA, 0.5 s). **C,**

Freezing levels during CS presentations of each group across the experiment. Two-way Repeated-

Measure ANOVA for each day followed by Tukey's post hoc test. Letters a, b, and c indicate pairwise

post hoc tests with $p < 0.05$: a, FC vs. OPC; b, FC vs. PC; c, OPC vs. PC. **D,** OPC group shows lower

average speed compared to PC group. One-way ANOVA followed by Tukey post hoc test. Letters b and c

indicate pairwise tests with $p < 0.05$: b, FC vs. PC ; c, OPC vs. PC. **E,** OPC group is resistant to extinction

while PC group shows significantly more extinction. Left, the freezing levels during the first 4 and last 4

CS presentations. Paired Student's t test. Letter e indicates test with $p < 0.05$: PC, Last 4 CS vs. First 4

CS. Right, normalized change of the freezing level during extinction, as indicated by the difference of the

freezing levels between last 4 and first 4 CS presentations as a percentage of the freezing level of the

first 4 CS presentations. One-way ANOVA followed by Tukey's post hoc test. * $p<0.05$. Letter c indicates

pairwise test with $p < 0.05$: OPC vs. PC. Data shown here and in subsequent illustrations as mean ±

SEM.

Figure 3. **Optimal partial conditioning protocol induced stronger fear memory than spaced partial**

conditioning protocol in rats. **A**, *Inducer* peak levels from the FC, PC, and OPC protocols, compared to

peak levels from partial conditioning protocols of 4 trials with regular ITIs varying from 1 to 21 min, using

standard intensity stimulus (**A1**) or reduced intensity stimulus (**A2**). The peak *inducer* values of FC, PC,

and OPC are labeled by dashed lines (black: FC; red: OPC; purple: PC). The blue curve gives peak

inducer values for the protocols with regular ITIs, and the curve peaks at the dark blue dot and arrow,

representing the SPC protocol with equal ITIs of 11 min and 10 s. **B,** Schematics of the fear conditioning

procedures. Top: tests that were conducted in Context A. Bottom: tests that were conducted in Context B.

C, Schematics of the fear conditioning protocols for optimal partial conditioning (OPC, n = 8), and spaced

partial conditioning groups (SPC, n = 10, ITI = 670 s). Following a habituation (Hab.) tone, rats received 4

trials of a conditioned stimulus (CS, 3 kHz tone, 30 s) that co-terminated with an unconditioned stimulus

(US, footshock, 0.5 mA, 0.5 s). **D,** Freezing levels during CS presentations of each group across the

experiment; Two-way repeated-measure ANOVA for each day followed by Tukey's pairwise post hoc test,

* $p < 0.05$. Inset: the average freezing level during the first two CS presentations; Welch's t test, * $p < 0.05$.

Figure 4. Prediction of enhanced protocol for fear extinction. A, Schematic of the model. **B1**, Simulated time courses of activated ERK (pERK, orange traces, μM) and activated PKA (PKAc, green traces) in response to one trial of *Stim* (μM). **B2**, Simulated time courses of pERK (orange traces, μM), PKAc (green traces) and *inducer* (red traces, nM) in response to 12-trial protocol with regular intervals of 2.5 min (full extinction, FE). **B3**, Simulated time courses of pERK (orange traces), PKAc (green traces) and *inducer* (red traces) in response to 4-trial protocol with regular intervals of 2.5 min (partial extinction, PE). **B4**, Simulated time courses of pERK (orange traces), PKAc (green traces) and *inducer* (red traces) in response to 4-trial protocol with computationally designed intervals (optimal partial extinction, OPE).

Figure 5. Computationally designed protocol enhanced fear extinction in rats. **A.** Schematic of the enhanced fear extinction experimental procedures. Rats were trained to constantly press a lever to retrieve sucrose pellets in a variable interval 60 (VI60) schedule, where the average interval between each sucrose delivery is 60 s. **B,** Schematics of the fear extinction protocols for full extinction (FE, n = 12), partial extinction (PE, n = 11), and optimal partial extinction groups (OPE, n = 13), during which rats received multiple trials of a conditioned stimulus (CS, 3 kHz tone, 30 s). **C,** Freezing levels during CS presentations of each group across the experiment. No significant difference between groups was found

by Two-way RM ANOVA. **D,** Lever presses rates during the 30s of each CS presentation (pre-CS period)

of each group across the experiment. Two-way repeated-measure ANOVA for each day followed by

Tukey's pairwise post hoc test. Letters a and c indicate pairwise tests with $p < 0.05$: a, FE vs. OPE ; c,

OPE vs. PE. **E,** OPE group shows significant increase of pre-CS lever presses comparing the last CS

presentation to the average lever press rate of the first two CS presentations. Paired Student's t test.

Letter f indicates test with $p < 0.05$: OPC, Last 4 CS vs. First 4 CS. **F,** OPE group shows higher lever

presses during pre-CS period compared to PE group in spontaneous recovery test. One-way ANOVA

followed by Tukey's post hoc test. Letter c indicates pairwise comparison with * $p<0.05$: OPE vs. PE.

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

     SUPPLEMENTARY FIGURES

1025

Supplementary Figure 1. No difference of baseline activity between groups. A, No differences in

total freezing level (left), average speed (middle), and maximum speed (right) between the three groups

were observed during familiarization of the enhanced conditioning experiment. **B,** No differences in total

freezing level (left), average speed (middle), and maximum speed (right) were seen between the three

groups during familiarization of the enhanced vs. spaced conditioning experiment. **C,** No differences in

total freezing level (left), average speed (middle), and rate of lever presses (right) were observed between

the three groups during the last day of lever presses training of the enhanced extinction experiment. One-

way ANOVA. Data shown as mean ± SEM.

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

Supplementary Figure 2. **Histogram of peak levels of *inducer* from 1,000 protocols. A**, Standard

intensity stimulus (*Stim* = 300 μM). The range of peak levels of *inducer* (0 - 220 nM) was subdivided into

14 bins, and the number of simulations that produced a peak concentration of *inducer* in each subdivision

was plotted. The arrows indicate which bins contained the peak concentrations produced by the partial

conditioning, spaced partial conditioning (SPC), and the optimal partial conditioning (OPC) protocols. **B,**

Reduced intensity stimulus (*Stim* = 200 μM). The range of peak levels of *inducer* (0 - 160 nM) was

subdivided into 14 bins, and the number of simulations that produced a peak concentration of *inducer* in

each subdivision was plotted. Insets below the main plots illustrate in detail the difference in peak *inducer*

for SPC *vs*. OPC, which is negligible in **A**.

1062

1063

1064

1065

1066

1067

1068

1069

Supplementary Figure 3. Lever press during CS presentations in enhanced extinction experiment.

A, Lever presses rates during CS presentations of each group across the experiment. Two-way repeated-

measure ANOVA for each day found no difference among groups. **B,** FE group shows significant

increase of CS lever presses comparing the last CS presentation to the average lever press rate of the

first two CS presentations. Paired Student's t test. Letter d indicates test with $p < 0.05$: d, FE, Last 4 CS

vs. first 4 CS.

1076