Enhancing associative learning in rats with a computationally designed training

protocol

Xu O. Zhang; Yili Zhang; Claire E. Cho; Douglas S. Engelke; Paul Smolen; John H

Byrne; Fabricio H. Do-Monte[#].

Department of Neurobiology and Anatomy, McGovern Medical School, The University of

Texas Health Science Center at Houston, Houston, Texas 77030

#Corresponding author:

Fabricio H. Do Monte, DVM, PhD.

Phone: 713-500-5613

Email: fabricio.h.domonte@uth.tmc.edu

Abbreviated title: Enhancing learning with a computational model

Conflict of Interests: The authors declare no competing financial interests.

22   **Keywords:** Associative   memory,   fear   conditioning,   fear   extinction,   long-term

23   potentiation, computational model, spaced learning.

Abstract

Background - Learning requires the activation of protein kinases with distinct temporal dynamics. In *Aplysia,* nonassociative learning can be enhanced by a computationally designed learning protocol with intertrial intervals (ITIs) that maximize the interaction between fast-activated protein kinase A (PKA) and slow-activated extracellular signal-regulated kinase (ERK). Whether a similar strategy can enhance associative learning in mammals is unknown.

Methods - We simulated 1,000 training protocols with varying ITIs to predict an optimal protocol based on empirical data for PKA and ERK dynamics in rat hippocampus. Adult male rats received the optimal protocol or control protocols in auditory fear conditioning and fear extinction experiments. Immunohistochemistry was performed to evaluate phosphorylated cAMP responsive element binding (pCREB) protein levels in brain regions implicated in fear acquisition.

Results - Rats exposed to the optimal conditioning protocol with irregular ITIs exhibited impaired extinction memory acquisition within the session using a standard footshock intensity, and stronger fear memory retrieval and spontaneous recovery with a weaker footshock intensity, compared to rats that received massed or spaced conditioning protocols with fixed ITIs. Rats exposed to the optimal extinction protocol displayed improved extinction of contextual fear memory and reduced spontaneous recovery compared to rats that received standard extinction protocols. Moreover, the optimal conditioning protocol increased pCREB levels in the dentate gyrus (DG) of the dorsal hippocampus, suggesting enhanced induction of long-term potentiation.

46 <u>Conclusion</u> - These findings demonstrate that a computational model-driven behavioral

47 intervention can enhance associative learning in mammals, and may provide insight into

48 strategies to improve cognition in humans.

INTRODUCTION

Long-term memory (LTM) formation is believed to be mediated by synaptic plasticity including long-term potentiation (LTP) or its invertebrate analogue long-term facilitation (LTF), which require gene expression and protein synthesis (1-5). Studies over decades have investigated LTP/LTF and their underlying molecular processes as potential targets to enhance learning or restore memory deficits in laboratory animals. However, interventions using systemic cognitive enhancers or intracerebral pharmacological manipulations (6-10) are either based on trial-and-error approaches in specific model systems or are highly invasive, preventing or hindering translation to humans.

An alternative approach to enhance learning and memory is to develop computational models to predict optimal learning protocols based on dynamics of intracellular molecular cascades that underlie LTM formation and LTP/LTF induction (11-13). Studies have identified activation of PKA (14-16) and of the mitogen-activated protein kinase (MAPK) isoform ERK (17-19) as essential cascades for LTF. These two pathways converge to phosphorylate transcriptional factors such as CREB, which subsequently induce plasticity-related genes during LTF induction (20-24). These two pathways exhibit distinct kinetics of activation (13, 16, 25), suggesting that the temporal activity patterns, and activation overlap, of these pathways may constitute an important target to enhance associative learning. We previously demonstrated that a computationally designed protocol with irregular ITIs, predicted to maximize overlap of PKA and ERK activities, enhances LTF and nonassociative learning, specifically long-term sensitization of the tail elicited siphon-withdrawal reflex, in *Aplysia* (26).

Substantial similarities between molecular processes of LTF in invertebrates and

LTP in mammals make it plausible that strategies used to enhance LTF and

nonassociative learning in invertebrates could enhance LTP and associative learning in

mammals. PKA activation in rodent hippocampus and amygdala is required for LTP and

LTM (27, 28). Activation of ERK/MAPK cascades and their cross-talk with PKA are

required for CREB phosphorylation, induction of plasticity-related genes, and LTP

induction (29-32). Although the dynamics of PKA and ERK activation differ between

invertebrates and mammals (33-35), and may vary across different brain subregions and

behavioral protocols in rodents, the chronological order of activation for these two

intracellular molecules is evolutionarily-conserved across species. In fact, a PKA peak

precedes the nuclear translocation of ERK and the subsequent activation of CREB in

Aplysia (16, 25),   mice (29), and rats (36-38) which provides strong support for our

model.

Therefore, we tested if our invertebrate LTF model (26) can be adapted to

computationally design an optimal associative learning protocol in mammals. Because

the majority of the above literature on LTP induction has focused on fear conditioning

experiments, we employed this well-established paradigm to investigate associative

learning (39) and used extant empirical data to model PKA and ERK dynamics in rat

hippocampus (33-35), a critical brain region implicated in the formation of associative

memories (40-42). We simulated 1,000 different training protocols with varying ITIs to

identify an optimal protocol with irregular ITIs that maximizes the overlap of PKA and

ERK activities, thereby predicting the formation of stronger memory than training

protocols with the same number of stimuli and fixed ITIs. We then tested this optimal

protocol in auditory fear conditioning and fear extinction paradigms in rats.

RESULTS

A computational model based on PKA and ERK dynamics identified an optimal

fear conditioning protocol

The model in (26) was adapted to simulate PKA and ERK dynamics in rat hippocampus

during LTP induction (**Fig. 1A**). To the best of our knowledge, hippocampus is the only

rat brain region in which ERK dynamics has been measured with a temporal resolution

of minutes following a stimulus. We simulated fear conditioning, in which the pairing of a

conditioned stimulus (CS) with an unconditioned stimulus (US) is represented by *Stim*.

The conditioned responses are assumed to be proportional to the peak value of *inducer*.

A single CS-US pairing produced little overlap between PKA and ERK pathways (**Fig.**

1B1). We then simulated 1,000 fear conditioning protocols, with 4 trials and both fixed

and irregular ITIs. These simulations included a Short Conditioning (SC) protocol with 4

trials and a fixed ITI of 270 s. As a reference control, we also simulated a Regular

Conditioning (RC) protocol with 8 trials and the same ITI of 270 s. These protocols

respectively resemble previous protocols used for short and regular fear conditioning in

rats (43, 44). Simulations identified an optimal conditioning protocol with 4 trials and

irregular ITIs of 8, 8, and 16 min (**Fig. 1B2-1B4**), termed Optimal Short Conditioning

(OSC). PKA activity induced by the last trial of OSC had a much larger overlap with the

pERK curve than did PKA induced by SC. Therefore, OSC induced a much higher peak

level of *inducer* (red). We therefore predicted OSC would produce stronger long-term

memory in rats than standard SC.

The optimal short conditioning protocol increases fear memory compared to the standard short conditioning protocol

Because the hippocampus is required for context encoding during auditory fear conditioning (45, 46), we used two different chambers (context A and context B) to assess the contribution of the context to the auditory fear memory(**Fig. 2A**). Our pilot experiments and a series of recent fear conditioning studies have shown that female rats exhibit higher active defensive responses (i.e., darting) and lower passive defensive responses (i.e., freezing) than male rats (47-50). We therefore performed our experiments in males which exhibit enhanced freezing behavior, the main index of fear memory adopted in our study. On Day 0, rats were familiarized with context A for 20 min. They were pre-assigned to Short Conditioning (SC), Optimal Short Conditioning (OSC), or Regular Conditioning (RC) groups so that baseline freezing and locomotion were similar among groups (**Table 1**). On Day 1, rats were placed into context A and exposed to one nonreinforced habituation tone followed by the SC, OSC, or RC protocols (**Fig. 2B**). All groups reached high levels of freezing (**Fig. 2C**) and reduced locomotion at the end of the fear acquisition session.

On Day 2, rats were placed in context B and given two CS presentations to test the retrieval of tone-associated fear memory in a novel context. Compared to the RC group, both SC and OSC groups exhibited less freezing during the second CS presentation (**Fig. 2C**). Yet, both RC and OSC groups showed reduced average speed during the retrieval test compared to the SC group, suggesting generalized fear responses in a novel context (**Table 1**).

140   On Day 3, in context A, rats underwent a retrieval and extinction training session 141   with twelve CS presentations. Only the RC group showed a significantly higher pre-tone 142   freezing level, suggesting a robust contextual fear memory (**Table 1**). However, the 143   three groups exhibited the same levels of freezing during the first two CS presentations, 144   suggesting tone-evoked fear retrieval was comparable among the groups (**Fig. 2C**). 145   Nevertheless, a within session extinction analysis comparing the first four CSs to the 146   last four CSs revealed impaired extinction learning in the OSC group as compared to 147   the SC group, with the OSC group displaying smaller reduction of freezing levels from 148   the first to the last CS presentation (**Fig. 2D)**. Despite this extinction impairment, no 149   significant differences were found between the OSC and the SC groups during 150   extinction retrieval tests performed on Days 4 and 5 or the spontaneous recovery tests 151   performed on Days 29 and 30 (**Fig. 2C**). The RC group showed higher averaged CS 152   freezing compared to SC group during both extinction retrieval tests (**Table 1**).

153   In summary, the OSC protocol demonstrated modest enhancement of fear 154   acquisition, indicated by reduced locomotion and stronger, more resistant CS-evoked 155   fear memory during within-session extinction. However, this enhancement didn't persist 156   during extinction retrieval and spontaneous recovery tests. These results suggest OSC 157   produces a more robust initial fear memory but without a lasting effect on extinction 158   memory.

The optimal short conditioning protocol induces stronger fear memory than a 161   **spaced short conditioning protocol**

Spaced learning protocols result in stronger memories than massed learning protocols

in both humans (51, 52) and rodents (53-55, but also see 56), raising the possibility that

the augmented fear memory observed in the OSC group could simply be attributed to a

trial-spacing effect. Our simulations predicted a broad range of effective simulated 4-trial

protocols (**Fig. 3**). One protocol with relatively long, fixed ITIs of 11 min 10 s produced a

nearly identical peak level of *inducer* as the OSC protocol, and a higher peak level of

inducer than other protocols with fixed ITIs from 1 min to 21 min (**Fig. 3A, Fig. 4A1**). We

termed this protocol the Spaced Short Conditioning (SSC) protocol. Further simulations

with reduced stimulus intensity showed that the OSC protocol still produced the greatest

peak level of *inducer* and was relatively more effective compared to SSC, with these

weaker stimuli (**Fig. 3B, Fig. 4A2**). Therefore, we predicted that the irregular intervals of

OSC may produce more effective conditioning than does regularly spaced SSC.

To test this prediction, we empirically compared the SSC and OSC protocols

using a reduced shock intensity (0.5 mA instead of the 0.7 mA used in Fig. 2) (**Fig. 4B-**

C). Rats exposed to OSC showed higher CS freezing during the second and third CS-

US pairing of the fear acquisition session (**Fig. 4D, Day 1)**. In addition, fear memory

retrieval was increased when rats in the OSC group were returned to context A for an

extinction training session on Day 3, as indicated by higher freezing during the first two

CS presentations, compared to the SSC group (**Fig. 4D, Inset**). Both OSC and SSC

groups exhibited the same levels of freezing by the end of the extinction training session

(**Fig. 4D, Day 3**), as well as during the extinction retrieval tests performed in context A

(**Fig. 4D, Day 4**) or context B (**Fig. 4D, Day 5**). However, the OSC group showed

greater spontaneous recovery of fear memory in context A approximately 3 weeks later,

as indicated by higher freezing during the first CS presentation compared to SSC-

trained rats (**Fig. 4D, Day 29**). In summary, these data suggest the enhancement in fear

memory acquisition observed with our OSC protocol cannot simply be attributed to a

trial-spacing effect or differences in the delay to remove the animals from the chamber,

but it is rather associated with the maximized overlap between PKA and ERK signaling.

The optimal short conditioning protocol induces greater levels of phosphorylated

CREB in DG than a spaced short conditioning protocol

The computational model predicts that the increased overlap of PKA and ERK activation

in OSC protocol leads to greater LTP induction (*inducer*) in rat hippocampus. This

increased overlap plausibly augments phosphorylation of CREB (29), a transcription

factor required for synaptic plasticity and LTP (22, 57, See review by 58). We therefore

used immunohistochemistry to quantify levels of pCREB following fear conditioning in

No-Shock control (NS), and in SSC and OSC groups (**Fig. 5A**) in brain regions

previously implicated in the acquisition of fear memory such as the amygdala, the

paraventricular nucleus of the thalamus, and the dorsal hippocampus (59, 60, see review

by 61). During the fear acquisition session, rats exposed to SSC and OSC protocols

exhibited higher freezing levels, indicative of successful fear acquisition compared to

the NS group, with the OSC group showing even higher freezing levels than the SSC

group (**Fig. 5B**), consistent with Fig. 4.

Immunohistochemistry revealed high levels of pCREB expression in the SSC and

OSC groups, as compared to the NS group, across different brain regions including the

lateral, basal, and central nuclei of the amygdala (**Fig. 5C**), and the paraventricular

nucleus of the thalamus (PVT, **Fig. 5D**), suggesting that increased pCREB levels in

SSC and OSC groups are mediated by the CS-US pairing. However, we did not

observe differences in pCREB levels between the OSC and SSC groups in the

amygdala or PVT (**Table 1**), and pCREB levels were similar among the three groups in

CA1 hippocampal subfields (**Fig. 5E**). Interestingly, rats exposed to the OSC protocol

exhibited increased pCREB expression in the DG, as compared to the SSC and NS

groups (**Fig. 5E**). Given that increased pCREB in the DG has been associated with

enhanced LTP in rodents (62), these data support the hypothesis that the augmentation

in fear memory acquisition observed with the OSC protocol is mediated by higher levels

of LTP induction in the rat hippocampus.

The optimal protocol also enhanced fear extinction

Extinction is a new learning that temporarily inhibits the initial associative memory (63,

64). Extinction also relies on PKA and ERK signaling to induce plasticity-related gene

transcription (65-67). We predicted that the computationally designed protocol would

also enhance the acquisition of extinction, thereby suppressing the original fear memory

(**Fig. 6A-B1**). We compared an Optimal Short Extinction (OSE) protocol, comprising

four trials with irregular ITIs of 8, 8, and 16 minutes (**Fig. 6B3**), against a Regular

Extinction (RE, **Fig. 6B4**) protocol using 12 trials and ITIs of 150 s, and a Short

Extinction (SE, **Fig. 6B2**) protocol using 4 trials and the same ITIs of 150 s, which

resemble previous protocols used for regular and short fear extinction in rats (68-70). In

simulations, the OSE protocol triggered higher peak *inducer* than the SE protocol, and

was comparable to the RE protocol (**Fig. 6B1-3**). We thus predicted OSE would result in

stronger extinction of fear memory than does the standard SE.

To test this prediction, we designed an experiment comparing extinction protocol

efficacy using conditioned suppression of reward-seeking behavior (**Fig. 7A**), which is

more sensitive than freezing in fear extinction paradigms (71, 72). Lever presses also

help to maintain constant activity for reliable freezing measurement along the session

(73). Since lever presses for reward are trained in a specific context, and extinction

memory is context-dependent (74, 75), we used only one context in this experiment.

Rats were initially trained to press a lever to receive sucrose pellets in a variable interval

schedule of 60 s until they reached the same levels of lever pressing and locomotor

activity (see Supp. Methods). On Day 8, rats underwent a fear conditioning session

which included five nonreinforced habituation tones followed by seven CS-US pairings.

On Day 9, rats were pre-assigned to three experimental groups for extinction (**Fig. 7B**)

so that freezing levels, average speeds, and lever-press rates were similar during the

fear conditioning session (**Table 1**). At the end of the extinction session, freezing levels

reduced from ~50% to ~25% in the RE group, whereas the SE and OSE groups

maintained the same levels of freezing throughout the four CSs (**Fig. 7C**, Day 9).

Similarly, the RE group showed a significant increase in lever presses during the CS

presentations at the end of the session, whereas CS lever presses remained unaltered

in the SE and OSE groups (**Fig. 7D** *left*). However, the OSE group showed a significant

increase in lever presses during the 30-s periods preceding the CS presentations (pre-

CS, **Fig. 7D** *right*), suggesting enhanced within-session extinction of contextual fear

memory (76).

On Days 10 and 35, rats were returned to the same chamber to test the strength

of fear extinction memory during extinction retrieval and spontaneous recovery tests,

respectively. No significant difference in CS freezing or CS lever presses were observed

among the three groups (**Fig. 7C, Day 10&35**). However, the OSE group showed

increased pre-CS lever pressing during the spontaneous recovery test (**Fig. 7E *right***),

when compared to the SE and RE groups. These results suggest that, although the

tone-associated memory was similar among the groups, the OSE protocol enhanced

contextual fear memory extinction, evidenced by enhanced within-session extinction of

conditioned suppression and reduced conditioned suppression three weeks post-

extinction training.

DISCUSSION

A computationally designed protocol that maximizes the interaction between PKA and

ERK pathways enhances LTF and nonassociative learning in *Aplysia* (26). Here, we

extended this strategy to associative learning in mammals by adapting the simplified

mathematical model used in *Aplysia* to simulate the dynamics of PKA and ERK in rat

hippocampus based on empirical data in the literature (33-35). Through a series of

computational simulations and experimental validations, we discovered that an

optimized training protocol, predicted to enhance the overlap of PKA and ERK

activation, can facilitate acquisition and extinction of conditioned fear.

Immunohistochemistry demonstrated that the augmented fear memory induced by the

optimal protocol was associated with increased expression levels of pCREB in the DG

subfield of the dorsal hippocampus. Our results demonstrate the power of a simplified

model of intracellular signaling cascades in describing associative learning across species, attesting the essential role of the interaction between PKA and ERK pathways in both nonassociative and associative learning.

Training protocols using spaced ITIs result in stronger memory acquisition than those using standard massed ITIs, a well-established phenomenon described as the "spacing effect" (see review by 11). The spacing effect has been linked to the varying efficacy of massed and spaced protocols in triggering LTP/LTF via biochemical signaling, and a previous computational model suggests the optimal ITI aligns with peak ERK activation (77-79). However, that model assumed fixed ITIs, which may not be ideal for synaptic plasticity and long-term memory. Our optimized training protocol, with irregular ITIs, induced stronger and persistent fear memory in rats, as compared to a spaced protocol that had been predicted to be the most effective among protocols with equal ITIs. These results suggest that the spacing effect in mammals can be at least partially explained by enhanced overlap between the PKA and ERK pathways, which are critical for CREB activation (29). Our study also indicates that learning protocols with irregular inter-trial intervals (ITIs) may be more effective than those with equal ITIs. Although spaced protocols are known for facilitating fear memory acquisition, there is a lot of controversy when it comes to fear extinction. Whereas some studies have demonstrated that spaced intervals between the CSs facilitate extinction memory and attenuate renewal and spontaneous recovery of fear (56, 80-82), others have shown the opposite, impaired extinction memory and increased recovery of fear (56, 81) . We therefore selected a control group with massed trials to better control for our optimized protocol.

The enhanced performance of the optimal (OSC) protocol appeared to be constrained by the intensity of the stimuli. The model predicted higher performance when weak stimuli were used but comparable performance with strong stimuli (**Fig. 4A**). Drawing a line to distinguish weak and strong footshock intensities is not straightforward because the relationship between fear memory and footshock intensity is neither monotonic nor linear (83). Nevertheless, we observed clear differences in the efficacy of the optimal fear conditioning protocol when using footshocks of different intensities. When we compared the OSC protocol to a massed short conditioning protocol using a standard footshock intensity (0.7 mA, **Fig. 2**) commonly used in previous studies (28), we found only small differences in the conditioned responses (CS freezing) between the two protocols. However, when we used a lower footshock intensity (0.5 mA, **Fig. 4**) to compare the optimal short conditioning protocol to a spaced short conditioning protocol, we found a significant increase in CS freezing in the OSC group during fear acquisition, retrieval, and spontaneous recovery tests. Considering the robustness of the spacing effect, it is possible that differences between the OSC and the massed SC groups at the higher footshock intensity were in part masked by a "ceiling effect". Alternatively, previous studies in rats have demonstrated that the hippocampus is required for tone-evoked fear memory when a weak (0.4 mA) but not a strong (0.9 mA) footshock is used (84), which could also explain the differences we observed at the two intensities. Either way, our data suggest that our approach could be more beneficial for learning protocols relying on relatively weak stimuli. Additional studies need to be conducted to understand the mechanism by which the intensity of stimuli governs the enhanced performance of protocols with irregular ITIs compared to protocols with fixed ITIs.

The association between the CS and US is primarily mediated by the lateral

amygdala where LTP induces enhanced CS responses (see review by 39, 85).

Nevertheless, another important and distinguishable component of fear memory

formation is the context in which the association has occurred. Interestingly, we only

observed enhanced conditioned responses in the context in which fear conditioning

occurred, suggesting the memory facilitation effect is context-dependent. Similarly, in

the fear extinction experiment, enhancement by the optimal protocol was only observed

during the pre-CS lever pressing, a more sensitive index of contextual fear memory

during extinction and spontaneous recovery (71, 72, 76, 86, 87). This is consistent with

the hippocampus' role in encoding context during fear conditioning and extinction, and

in the time-dependent reappearance of fear post-extinction (*i.e.*, spontaneous recovery),

on which our model is based (see reviews by 46, 88). Accordingly, immunohistochemical

results showed greater pCREB expression in the OSC group specifically in the DG of

the dorsal hippocampus, a region implicated in the acquisition and extinction of

contextual fear memory (89, 90). Increasing CREB expression in DG neurons has been

demonstrated to improve contextual fear acquisition (91), whereas inactivating DG

neurons that overexpress CREB disrupts contextual fear retrieval (92).

Another hippocampal subregion implicated in contextual fear is the CA1 (93).

Although the OSC, SSC and NC groups displayed similar levels of pCREB expression

in CA1, the single post-training time point (15 min) we used does not suffice to rule out

the possibility of distinct pCREB dynamics in CA1 compared to DG. CA1 pCREB peaks

at 30 min and disappears at 90 min following fear conditioning (94, 95). Further

experiments comparing distinct post-training intervals may better reveal the dynamics of

pCREB expression in different hippocampal subregions, as well as in other areas involved in fear acquisition (e.g., amygdala, PVT), which in the present study exhibited similar levels of pCREB expression in the OSC and SSC groups. The observation that the dynamics of ERK pathways differ across brain regions involved in fear memory, with peaks occurring at 20 min after a single stimulation in the hippocampus versus 60 min in the lateral amygdala (35, 96, 97), suggests that the model may be able to predict optimal protocols targeting specific components of fear memory based on the dynamics of intracellular signaling cascades in corresponding brain regions. Future studies will test this possibility by modeling the molecular cascades in the lateral amygdala and the medial prefrontal cortex to preferentially target the acquisition and extinction of CS-associated memories, respectively.

Our model is an obvious simplification of the mechanisms underlying fear conditioning and extinction. It does not include other molecular cascades critical for LTP and memory formation, such as calcium/calmodulin-dependent kinase II (CaMKII) or protein kinase C (PKC), due to the lack of empirical data for simulating their dynamics of activation (12, 98, 99). The model was constructed using empirical PKA and ERK dynamics from the literature, which are based on *ex vivo* analyses and have limited temporal resolution. Furthermore, the model assumes the molecular mechanisms for fear memory acquisition are similar to those for extinction. Despite these limitations, the simplified model was sufficient to predict enhanced activation of the LTP-related transcription factor CREB in the DG and to facilitate associative learning in three different experiments. A more complex model that incorporates a wider range of intracellular and extracellular processes based on *in vivo* data would likely have

enhanced predictive ability. Further experiments should also investigate the memory

enhancing effects of computationally designed training protocols in different types of

associative memories, including discrimination learning and backwards conditioning. In

addition, it will be important to test whether the effects observed with optimal protocols

vary across subjects of different sexes and ages, as well as protocol efficacy in animal

models of cognitive impairment. Together, our results suggest the possibility of using

similar model-driven, non-invasive behavioral approaches in studies aimed at

enhancing learning or restoring memory deficits in humans, or aimed at improving

extinction-based exposure therapies for anxiety disorders.

MATERIALS AND METHODS (see additional details in Supplementary Methods).

Animals

The study involved 120 male Long-Evans rats, aged 3-5 months, kept in a 12-hour

light/dark cycle. All procedures followed the National Institutes of Health guidelines for

animal care and were approved by the Center for Laboratory Animal Medicine and Care

of The University of Texas Health Science Center at Houston.

Model Development

The mathematical model, adapted from (26), describes the activation of the PKA and

ERK pathways. Parameters were adjusted based on empirical data (33-35, 78) for

hippocampus. Simulations used fourth-order Runge-Kutta integration for differential

equations, and were conducted in XPPAUT (100) on Dell Precision T1700

microcomputers.

Behavioral Tasks

Rats were exposed to fear conditioning in two distinct chambers. Three groups – RC

(eight tone-shock pairings, 270s intervals), SC (four pairings, 270s intervals), and OSC

(four pairings, 8, 8, and 16 min intervals) - were used. Optimal vs. Spaced Short

Conditioning compared OSC with SSC (four pairings, 11min 10s intervals) with

footshock intensity at 0.5 mA. Extinction involved lever-press training, followed by fear

conditioning. These rats were divided into three groups: RE (twelve tone presentations,

150s intervals), SE (four tone presentations, 150s intervals), and OSE (four tone

presentations, 8, 8, and 16 min intervals).

Immunohistochemistry

Rat brains were prepared post-conditioning for immunohistochemistry to analyze

pCREB-positive cells. Brains were sectioned and treated with rabbit anti-pCREB serum

(1:1000; EMD Millipore 06-519) and biotinylated goat anti-rabbit IgG antibody (1:200,

Vector Labs, BA-1000-1.5), and revealed by ABC kit (1:50, Vector Labs, PK-6100) and

DAB-Ni solution (Vector Labs, SK-4100). Images were captured using a Nikon

microscope and analyzed with QuPath software (101) for automatic cell detection and

quantification. pCREB-positive cells in specified brain regions were counted and

densities calculated.

Quantification and statistical analysis

Grubbs' tests (102) were used to identify outliers in each experiment. Shapiro-Wilk tests were performed to determine normal distribution. Equal variance was confirmed through F tests and Brown-Forsythe tests. Statistical significance was determined using t-tests, ANOVA, or Kruskal-Wallis tests with relevant post-hoc comparisons, as appropriate. Sample size was based on prior literature and experience.

Data availability

All data that support the findings presented in this study are available from the corresponding author on reasonable request. Source codes will be submitted to GitHub (https://github.com/Owenxz/Zhang-XO-Enhanced-Learning-2022.git).

Author Contributions: X.O.Z, D.S.E., and C.E.C performed and analyzed the behavioral experiments. X.O.Z and C.E.C performed and analyzed the immunohistochemistry experiments. Y.Z. implemented the computational model and ran all simulations. P.S. helped design and implement the computational model. F.H.D-M and J.H.B supervised and contributed to all aspects of this study. All the authors participated in the design of the experiments. X.O.Z. and F.H.D-M interpreted the data and prepared the manuscript with comments from all the co-authors.

Acknowledgments: This work was supported by the Russell and Diana Hawkins Family Foundation Discovery Fellowship and the Dr. John J. Kopchick Fellowship to X.O.Z, NIH grant R01-NS102490 to J.H.B, and NIH grant R00-MH105549, NIH grant

R01-MH120136, a Brain & Behavior Research Foundation grant (NARSAD Young

Investigator), and a Rising STARs Award from UT System to F.H.D-M. We thank Nikita

Watson and Sharon Gordon for their technical and administrative assistance. We also

thank current members of the Byrne and Do Monte labs for their valuable comments on

the manuscript. This manuscript has been posted on BioRxiv as a preprint (doi:

https://doi.org/10.1101/621540).

REFERENCES

1.   Martin SJ, Grimwood PD, Morris RG (2000): Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci.* 23:649-711.

2.   Kandel ER (2001): The molecular biology of memory storage: a dialogue between genes and synapses. *Science.* 294:1030-1038.

3.   Lynch MA (2004): Long-term potentiation and memory. *Physiol Rev.* 84:87-136.

4.   Alberini CM (2009): Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev.* 89:121-145.

5.   Byrne JH, Hawkins RD (2015): Nonassociative learning in invertebrates. *Cold Spring Harb Perspect Biol.* 7.

6.   McGaugh JL, Petrinovich LF (1965): Effects of drugs on learning and memory. *Int Rev Neurobiol.* 8:139-196.

7.   Fernandez SM, Lewis MC, Pechenino AS, Harburger LL, Orr PT, Gresack JE, et al. (2008): Estradiol-induced enhancement of object memory consolidation involves hippocampal extracellular signal-regulated kinase activation and membrane-bound estrogen receptors. *J Neurosci.* 28:8660-8667.

8.   Lynch G, Cox CD, Gall CM (2014): Pharmacological enhancement of memory or cognition in normal subjects. *Front Syst Neurosci.* 8:90.

9.   Lauterborn JC, Palmer LC, Jia Y, Pham DT, Hou B, Wang W, et al. (2016): Chronic ampakine treatments stimulate dendritic growth and promote learning in middle-aged rats. *J Neurosci.* 36:1636-1646.

10.   Sharif S, Guirguis A, Fergus S, Schifano F (2021): The use and impact of cognitive enhancers among university students: a systematic review. *Brain Sci.* 11.

11.   Smolen P, Zhang Y, Byrne JH (2016): The right time to learn: mechanisms and

optimization of spaced learning. *Nat Rev Neurosci*. 17:77-88.

12.   Smolen P, Baxter DA, Byrne JH (2020): Comparing theories for the maintenance

of late LTP and long-term memory: computational analysis of the roles of kinase

feedback pathways and synaptic reactivation. *Front Comput Neurosci*. 14:569349.

13.   Zhang Y, Smolen PD, Cleary LJ, Byrne JH (2021): Quantitative description of the

interactions among kinase cascades underlying long-term plasticity of *Aplysia* sensory

neurons. *Scientific reports*. 11:14931.

14.   Schacher S, Castellucci VF, Kandel ER (1988): cAMP evokes long-term

facilitation in *Aplysia* sensory neurons that requires new protein synthesis. *Science*.

240:1667-1669.

15.   Goldsmith BA, Abrams TW (1991): Reversal of synaptic depression by serotonin

at Aplysia sensory neuron synapses involves activation of adenylyl cyclase. *Proc Natl*

Acad Sci U S A. 88:9021-9025.

16.   Muller U, Carew TJ (1998): Serotonin induces temporally and mechanistically

distinct phases of persistent PKA activity in *Aplysia* sensory neurons. *Neuron*. 21:1423-

1434.

17.   Martin KC, Michael D, Rose JC, Barad M, Casadio A, Zhu H, et al. (1997): MAP

kinase translocates into the nucleus of the presynaptic cell and is required for long-term

facilitation in *Aplysia*. *Neuron*. 18:899-912.

18.   Sharma SK, Carew TJ (2004): The roles of MAPK cascades in synaptic plasticity

and memory in *Aplysia*: facilitatory effects and inhibitory constraints. *Learn Mem*.

11:373-378.

19.   Sharma SK, Sherff CM, Shobe J, Bagnall MW, Sutton MA, Carew TJ (2003):

Differential role of mitogen-activated protein kinase in three distinct phases of memory

for sensitization in *Aplysia*. *J Neurosci*. 23:3899-3907.

20.   Dash PK, Hochner B, Kandel ER (1990): Injection of the cAMP-responsive

element into the nucleus of *Aplysia* sensory neurons blocks long-term facilitation.

Nature. 345:718-721.

21.   Bartsch D, Ghirardi M, Skehel PA, Karl KA, Herder SP, Chen M, et al. (1995):

Aplysia CREB2 represses long-term facilitation: relief of repression converts transient

facilitation into long-term functional and structural change. *Cell*. 83:979-992.

22.   Liu RY, Cleary LJ, Byrne JH (2011): The requirement for enhanced CREB1

expression in consolidation of long-term synaptic facilitation and long-term excitability in

sensory neurons of *Aplysia*. *J Neurosci*. 31:6871-6879.

23.   Guan Z, Giustetto M, Lomvardas S, Kim JH, Miniaci MC, Schwartz JH, et al.

(2002): Integration of long-term-memory-related synaptic plasticity involves bidirectional

regulation of gene expression and chromatin structure. *Cell*. 111:483-493.

24.   Mohamed HA, Yao W, Fioravante D, Smolen PD, Byrne JH (2005): cAMP-

response elements in Aplysia creb1, creb2, and Ap-uch promoters: implications for

feedback loops modulating long term memory. *The Journal of biological chemistry*.

280:27035-27043.

25.   Philips GT, Tzvetkova EI, Carew TJ (2007): Transient mitogen-activated protein

kinase activation is confined to a narrow temporal window required for the induction of

two-trial long-term memory in *Aplysia*. *J Neurosci*. 27:13701-13705.

26.   Zhang Y, Liu RY, Heberton GA, Smolen P, Baxter DA, Cleary LJ, et al. (2011):

Computational design of enhanced learning protocols. *Nat Neurosci*. 15:294-297.

27.   Schafe GE, LeDoux JE (2000): Memory consolidation of auditory pavlovian fear

conditioning requires protein synthesis and protein kinase A in the amygdala. *J*

Neurosci. 20:RC96.

28.   Abel T, Nguyen PV, Barad M, Deuel TA, Kandel ER, Bourtchouladze R (1997):

Genetic demonstration of a role for PKA in the late phase of LTP and in hippocampus-

based long-term memory. *Cell*. 88:615-626.

29.   Impey S, Obrietan K, Wong ST, Poser S, Yano S, Wayman G, et al. (1998):

Cross talk between ERK and PKA is required for Ca^{2+} stimulation of CREB-dependent

transcription and ERK nuclear translocation. *Neuron*. 21:869-883.

30.   Tao X, Finkbeiner S, Arnold DB, Shaywitz AJ, Greenberg ME (1998): Ca2+ influx

regulates BDNF transcription by a CREB family transcription factor-dependent

mechanism. *Neuron*. 20:709-726.

31.   Adams JP, Sweatt JD (2002): Molecular psychology: roles for the ERK MAP

kinase cascade in memory. *Annu Rev Pharmacol Toxicol*. 42:135-163.

32.   Bambah-Mukku D, Travaglia A, Chen DY, Pollonini G, Alberini CM (2014): A

positive autoregulatory BDNF feedback loop via C/EBPbeta mediates hippocampal

memory consolidation. *J Neurosci*. 34:12547-12559.

33.   Roberson ED, Sweatt JD (1996): Transient activation of cyclic AMP-dependent

protein kinase during hippocampal long-term potentiation. *The Journal of biological*

chemistry. 271:30436-30441.

34.   Vázquez SI, Vázquez A, Peña de Ortiz S (2000): Different hippocampal activity

profiles for PKA and PKC in spatial discrimination learning. *Behavioral Neuroscience*.

114:1109-1118.

35.   Wang Y, Zhu G, Briz V, Hsu YT, Bi X, Baudry M (2014): A molecular brake

controls the magnitude of long-term potentiation. *Nature communications*. 5:3051.

36.   Vossler MR, Yao H, York RD, Pan MG, Rim CS, Stork PJ (1997): cAMP

activates MAP kinase and Elk-1 through a B-Raf- and Rap1-dependent pathway. *Cell*.

89:73-82.

37.   Roberson ED, English JD, Adams JP, Selcher JC, Kondratick C, Sweatt JD

(1999): The mitogen-activated protein kinase cascade couples PKA and PKC to cAMP

response element binding protein phosphorylation in area CA1 of hippocampus. *J

Neurosci*. 19:4337-4348.

38.   Ahmed T, Frey JU (2005): Plasticity-specific phosphorylation of CaMKII, MAP-

kinases and CREB during late-LTP in rat hippocampal slices in vitro.

Neuropharmacology. 49:477-492.

39.   Johansen JP, Cain CK, Ostroff LE, LeDoux JE (2011): Molecular mechanisms of

fear learning and memory. *Cell*. 147:509-524.

40.   Milner B, Squire LR, Kandel ER (1998): Cognitive neuroscience and the study of

memory. *Neuron*. 20:445-468.

41.   Nakazawa K, Quirk MC, Chitwood RA, Watanabe M, Yeckel MF, Sun LD, et al.

(2002): Requirement for hippocampal CA3 NMDA receptors in associative memory

recall. *Science*. 297:211-218.

42.   Mayes A, Montaldi D, Migo E (2007): Associative memory and the medial

temporal lobes. *Trends Cogn Sci.* 11:126-135.

43.   Detert JA, Kampa ND, Moyer JR, Jr. (2008): Differential effects of training

intertrial interval on acquisition of trace and long-delay fear conditioning in rats. *Behav*

Neurosci. 122:1318-1327.

44.   Do-Monte FH, Quinones-Laracuente K, Quirk GJ (2015): A temporal shift in the

circuits mediating retrieval of fear memory. *Nature.* 519:460-463.

45.   Selden NR, Everitt BJ, Jarrard LE, Robbins TW (1991): Complementary roles for

the amygdala and hippocampus in aversive conditioning to explicit and contextual cues.

Neuroscience. 42:335-350.

46.   Maren S, Phan KL, Liberzon I (2013): The contextual brain: implications for fear

conditioning, extinction and psychopathology. *Nat Rev Neurosci.* 14:417-428.

47.   Gruene TM, Flick K, Stefano A, Shea SD, Shansky RM (2015): Sexually

divergent expression of active and passive conditioned fear responses in rats. *eLife.* 4.

48.   Pellman BA, Schuessler BP, Tellakat M, Kim JJ (2017): Sexually Dimorphic Risk

Mitigation Strategies in Rats. *eNeuro.* 4.

49.   Colom-Lapetina J, Li AJ, Pelegrina-Perez TC, Shansky RM (2019): Behavioral

Diversity Across Classic Rodent Models Is Sex-Dependent. *Front Behav Neurosci.*

13:45.

50.   Mitchell JR, Trettel SG, Li AJ, Wasielewski S, Huckleberry KA, Fanikos M, et al.

(2022): Darting across space and time: parametric modulators of sex-biased

conditioned fear responses. *Learn Mem.* 29:171-180.

51.   Cepeda NJ, Pashler H, Vul E, Wixted JT, Rohrer D (2006): Distributed practice in

verbal recall tasks: A review and quantitative synthesis. *Psychol Bull*. 132:354-380.

52.   Raman M, McLaughlin K, Violato C, Rostom A, Allard JP, Coderre S (2010):

Teaching in small portions dispersed over time enhances long-term knowledge

retention. *Med Teach*. 32:250-255.

53.   Scharf MT, Woo NH, Lattal KM, Young JZ, Nguyen PV, Abel T (2002): Protein

synthesis is required for the enhancement of long-term potentiation and long-term

memory by spaced training. *J Neurophysiol*. 87:2770-2777.

54.   Anderson MJ, Jablonski SA, Klimas DB (2008): Spaced initial stimulus

familiarization enhances novelty preference in Long-Evans rats. *Behav Processes*.

78:481-486.

55.   Jiang L, Wang L, Yin Y, Huo M, Liu C, Zhou Q, et al. (2019): Spaced training

enhances contextual fear memory via activating hippocampal 5-HT2A receptors. *Front

Mol Neurosci*. 12:317.

56.   Cain CK, Blouin AM, Barad M (2003): Temporally massed CS presentations

generate more fear extinction than spaced presentations. *J Exp Psychol Anim Behav

Process*. 29:323-333.

57.   Bourtchuladze R, Frenguelli B, Blendy J, Cioffi D, Schutz G, Silva AJ (1994):

Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive

element-binding protein. *Cell*. 79:59-68.

58.   Kida S (2012): A Functional Role for CREB as a Positive Regulator of Memory

Formation and LTP. *Exp Neurobiol*. 21:136-140.

59.   Fanselow MS, LeDoux JE (1999): Why we think plasticity underlying Pavlovian

fear conditioning occurs in the basolateral amygdala. *Neuron*. 23:229-232.

60.   Penzo MA, Robert V, Tucciarone J, De Bundel D, Wang M, Van Aelst L, et al.

(2015): The paraventricular thalamus controls a central amygdala fear circuit. *Nature*.

519:455-459.

61.   Do Monte FH, Quirk GJ, Li B, Penzo MA (2016): Retrieving fear memories, as

time goes by. *Molecular psychiatry*. 21:1027-1036.

62.   Davis S, Vanhoutte P, Pages C, Caboche J, Laroche S (2000): The MAPK/ERK

cascade targets both Elk-1 and cAMP response element-binding protein to control long-

term potentiation-dependent gene expression in the dentate gyrus in vivo. *J Neurosci*.

20:4563-4572.

63.   Quirk GJ, Mueller D (2008): Neural mechanisms of extinction learning and

retrieval. *Neuropsychopharmacology*. 33:56-72.

64.   de Oliveira Alvares L, Do-Monte FH (2021): Understanding the dynamic and

destiny of memories. *Neurosci Biobehav Rev*. 125:592-607.

65.   Flood JF, Jarvik ME, Bennett EL, Orme AE, Rosenzweig MR (1977): Protein

synthesis inhibition and memory for pole jump active avoidance and extinction.

Pharmacol Biochem Behav. 7:71-77.

66.   Szapiro G, Vianna MR, McGaugh JL, Medina JH, Izquierdo I (2003): The role of

NMDA glutamate receptors, PKA, MAPK, and CAMKII in the hippocampus in extinction

of conditioned fear. *Hippocampus*. 13:53-58.

67.   Tronson NC, Schrick C, Fischer A, Sananbenesi F, Pages G, Pouyssegur J, et

al. (2008): Regulatory mechanisms of fear extinction and depression-like behavior.

Neuropsychopharmacology. 33:1570-1583.

68.   Santini E, Ge H, Ren K, Pena de Ortiz S, Quirk GJ (2004): Consolidation of fear

extinction requires protein synthesis in the medial prefrontal cortex. *J Neurosci*.

24:5704-5710.

69.   Holmes A, Quirk GJ (2010): Pharmacological facilitation of fear extinction and the

search for adjunct treatments for anxiety disorders--the case of yohimbine. *Trends*

Pharmacol Sci. 31:2-7.

70.   Do-Monte FH, Manzano-Nieves G, Quinones-Laracuente K, Ramos-Medina L,

Quirk GJ (2015): Revisiting the role of infralimbic cortex in fear extinction with

optogenetics. *J Neurosci*. 35:3607-3615.

71.   Mast M, Blanchard RJ, Blanchard DC (1982): The relationship of freezing and

response suppression in a CER situation. *The Psychological Record*. 32:151.

72.   Sotres-Bayon F, Sierra-Mercado D, Pardilla-Delgado E, Quirk GJ (2012): Gating

of fear in prelimbic cortex by hippocampal and amygdala inputs. *Neuron*. 76:804-812.

73.   Quirk GJ, Russo GK, Barron JL, Lebron K (2000): The role of ventromedial

prefrontal cortex in the recovery of extinguished fear. *J Neurosci*. 20:6225-6231.

74.   Goode TD, Maren S (2014): Animal models of fear relapse. *ILAR J*. 55:246-258.

75.   Podlesnik CA, Kelley ME, Jimenez-Gomez C, Bouton ME (2017): Renewed

behavior produced by context change and its implications for treatment maintenance: A

review. *J Appl Behav Anal*. 50:675-697.

76.   Morgan MA, LeDoux JE (1995): Differential contribution of dorsal and ventral

medial prefrontal cortex to the acquisition and extinction of conditioned fear in rats.

Behav Neurosci. 109:681-688.

77.   Ajay SM, Bhalla US (2007): A propagating ERKII switch forms zones of elevated

dendritic activation correlated with plasticity. *HFSP J*. 1:49-66.

78.   Ajay SM, Bhalla US (2004): A role for ERKII in synaptic pattern selectivity on the

time-scale of minutes. *Eur J Neurosci*. 20:2671-2680.

79.   Genoux D, Haditsch U, Knobloch M, Michalon A, Storm D, Mansuy IM (2002):

Protein phosphatase 1 is a molecular constraint on learning and memory. *Nature*.

418:970-975.

80.   Urcelay GP, Wheeler DS, Miller RR (2009): Spacing extinction trials alleviates

renewal and spontaneous recovery. *Learn Behav*. 37:60-73.

81.   Lang AJ, Craske MG (2000): Manipulations of exposure-based therapy to reduce

return of fear: a replication. *Behav Res Ther*. 38:1-12.

82.   Gerhard DM, Meyer HC (2021): Extinction trial spacing across days differentially

impacts fear regulation in adult and adolescent male mice. *Neurobiol Learn Mem*.

186:107543.

83.   Davis M, Astrachan DI (1978): Conditioned fear and startle magnitude: effects of

different footshock or backshock intensities used in training. *J Exp Psychol Anim Behav*

Process. 4:95-103.

84.   Quinn JJ, Wied HM, Ma QD, Tinsley MR, Fanselow MS (2008): Dorsal

hippocampus involvement in delay fear conditioning depends upon the strength of the

tone-footshock association. *Hippocampus*. 18:640-654.

85. Maren S (2005): Synaptic mechanisms of associative memory in the amygdala. *Neuron.* 47:783-786.

86. Woods AM, Bouton ME (2008): Immediate extinction causes a less durable loss of performance than delayed extinction following either fear or appetitive conditioning. *Learn Mem.* 15:909-920.

87. Padilla-Coreano N, Do-Monte FH, Quirk GJ (2012): A time-dependent role of midline thalamic nuclei in the retrieval of fear memory. *Neuropharmacology.* 62:457-463.

88. Bouton ME, Maren S, McNally GP (2021): Behavioral and Neurobiological Mechanisms of Pavlovian and Instrumental Extinction Learning. *Physiol Rev.* 101:611-681.

89. Kheirbek MA, Drew LJ, Burghardt NS, Costantini DO, Tannenholz L, Ahmari SE, et al. (2013): Differential control of learning and anxiety along the dorsoventral axis of the dentate gyrus. *Neuron.* 77:955-968.

90. Bernier BE, Lacagnina AF, Ayoub A, Shue F, Zemelman BV, Krasne FB, et al. (2017): Dentate Gyrus Contributes to Retrieval as well as Encoding: Evidence from Context Fear Conditioning, Recall, and Extinction. *J Neurosci.* 37:6359-6371.

91. Restivo L, Tafi E, Ammassari-Teule M, Marie H (2009): Viral-mediated expression of a constitutively active form of CREB in hippocampal neurons increases memory. *Hippocampus.* 19:228-234.

92. Park S, Kramer EE, Mercaldo V, Rashid AJ, Insel N, Frankland PW, et al. (2016): Neuronal Allocation to a Hippocampal Engram. *Neuropsychopharmacology.* 41:2987-2993.

93.   Hunsaker MR, Kesner RP (2008): Dissociations across the dorsal-ventral axis of

CA3 and CA1 for encoding and retrieval of contextual and auditory-cued fear. *Neurobiol*

Learn Mem. 89:61-69.

94.   Sindreu CB, Scheiner ZS, Storm DR (2007): Ca2+ -stimulated adenylyl cyclases

regulate ERK-dependent activation of MSK1 during fear conditioning. *Neuron*. 53:79-89.

95.   Chaaya N, Jacques A, Belmer A, Beecher K, Ali SA, Chehrehasa F, et al. (2019):

Contextual Fear Conditioning Alter Microglia Number and Morphology in the Rat Dorsal

Hippocampus. *Front Cell Neurosci*. 13:214.

96.   Schafe GE, Atkins CM, Swank MW, Bauer EP, Sweatt JD, LeDoux JE (2000):

Activation of ERK/MAP kinase in the amygdala is required for memory consolidation of

pavlovian fear conditioning. *J Neurosci*. 20:8177-8187.

97.   Di Benedetto B, Kallnik M, Weisenhorn DM, Falls WA, Wurst W, Holter SM

(2009): Activation of ERK/MAPK in the lateral amygdala of the mouse is required for

acquisition of a fear-potentiated startle response. *Neuropsychopharmacology*. 34:356-

366.

98.   Malenka RC, Kauer JA, Perkel DJ, Mauk MD, Kelly PT, Nicoll RA, et al. (1989):

An essential role for postsynaptic calmodulin and protein kinase activity in long-term

potentiation. *Nature*. 340:554-557.

99.   Wang S, Sheng T, Ren S, Tian T, Lu W (2016): Distinct roles of PKCι/λ and

PKMζ in the initiation and maintenance of hippocampal long-term potentiation and

memory. *Cell Rep*. 16:1954-1961.

100.   Ermentrout B (2002): *Simulating, analyzing, and animating dynamical systems :*

a guide to XPPAUT for researchers and students. Philadelphia: Society for Industrial

and Applied Mathematics.

101.   Bankhead P, Loughrey MB, Fernandez JA, Dombrowski Y, McArt DG, Dunne

PD, et al. (2017): QuPath: Open source software for digital pathology image analysis.

Scientific reports. 7:16878.

102.   Grubbs FE (1969): Procedures for Detecting Outlying Observations in Samples.

Technometrics. 11:1-21.

Figure 1. Computational simulations of PKA and ERK pathways predict an optimal protocol for fear conditioning. A, Schematic of the model. Stimulus (*Stim*, µM) rapidly activates PKA via cAMP pathway, and activates ERK more slowly via Raf-MEK pthway. The synergistic interaction between PKA and ERK pathways is quantified by a variable *inducer*, which corresponds to the efficacy of the stimulus in inducing LTP. ERK kinetics were described by differential equations (see Supplementary Methods) with parameter values reproducing empirical findings that ERK activity reaches peak levels 15~20 min after BDNF treatment or tetanic stimuli in rat hippocampus acute slices (35, 78). Equations describing PKA kinetics simulated data showing that PKA is transiently activated within 2 min after LTP induction in

slices from rat hippocampus or within 5 min *in vivo* after spatial discrimination task training (33,

34). **B1**, Simulated time courses of activated ERK (pERK, µM) and activated PKA (PKAc, µM) in

response to one trial of *Stim*. **B2**, Simulated time courses of pERK, PKAc, and *inducer* in

response to a 4-trial protocol with regular ITIs of 4.5 min (short conditioning, SC). **B3**, Simulated

time courses of pERK, PKAc, and *inducer* (nM) in response to a 4-trial protocol with

computationally designed intervals (optimal short conditioning, OSC). **B4**, Simulated time

courses of pERK, PKAc, and *inducer* (nM) in response to an 8-trial protocol with regular ITIs of

4.5 min (regular conditioning, RC).

Figure 2. Computationally designed protocol partially enhanced fear conditioning in rats.

A, Schematics of the enhanced fear conditioning experimental procedures. Top: tests that were

conducted in Context A. Bottom: tests that were conducted in Context B. **B,** Schematics of the

fear conditioning protocols for SC (n = 12), OSC (n = 14), and RC (n = 14) groups. Following a

habituation (Hab.) tone, rats received multiple trials of a CS (3 kHz tone, 30 s) that co-

terminated with a US (footshock, 0.7 mA, 0.5 s). **C,** Freezing levels during CS presentations of

each group across the experiment. Two-way Repeated-Measure ANOVA for each day followed

by Holm-Sidak's post hoc test. Letters *a, b*, and *c* indicate pairwise post hoc tests with p < 0.05:

a, OSC vs. SC; *b*, OSC vs. RC; *c*, SC vs. RC. '×' denotes the pre-tone freezing levels. **D,** OSC

group is resistant to extinction while SC group shows significantly more extinction. *Left*, freezing

levels during the first 4 and last 4 CS presentations. Paired Student's t-test. Letter *e* indicates

test with p < 0.05: SC, Last 4 CS vs. First 4 CS. *Center*, normalized change of the freezing level

during extinction, as indicated by the difference of the freezing levels between last 4 and first 4

CS presentations as a percentage of the freezing level of the first 4 CS presentations ((Last 4

CS – First 4 CS)/First 4 CS). One-way ANOVA followed by Tukey's post hoc test. Letter *a*

indicates pairwise test with p < 0.05: OSC vs. SC. *Right*, the number of cues required to reach

to a 50% reduction of the original freezing level (average of freezing level during the first two

cues). Kruskal-Wallis's test followed by Dunn's multiple comparisons test. Letter *a* indicates

pairwise test with p < 0.05: OSC vs. SC. Data shown here and in the next illustrations as mean

± SEM.

Figure 3. **Histogram of peak levels of *inducer* from 1,000 protocols.** **A**, Standard intensity stimulus (*Stim* = 300 µM). The range of peak levels of *inducer* (0 – 220 nM) was subdivided into 14 bins, and the number of simulations that produced a peak concentration of *inducer* in each subdivision was plotted. The arrows indicate which bins contained the peak concentrations produced by the short conditioning (SC), spaced short conditioning (SSC), and optimal short conditioning (OSC) protocols. **B,** Reduced intensity stimulus (*Stim* = 200 µM). The range of peak levels of *inducer* (0 – 160 nM) was subdivided into 14 bins, and the number of simulations that produced a peak concentration of *inducer* in each subdivision was plotted. Insets below the main plots illustrate in detail the difference in peak *inducer* for SSC *vs*. OSC, which is negligible in **A**.

Figure 4. **Optimal short conditioning protocol induced stronger fear memory than spaced**

short conditioning protocol in rats. **A**, *Inducer* peak levels from the RC, SC, and OSC

protocols, compared to peak levels from short conditioning protocols of 4 trials with regular ITIs

varying from 1 to 21 min, using standard intensity stimulus (**A1**) or reduced intensity stimulus

(**A2**). The peak *inducer* values of RC, SC, and OSC are labeled by dashed lines (black: RC; red:

OSC; purple: SC). The blue curve gives peak *inducer* values for the protocols with regular ITIs,

and the curve peaks at the dark blue dot and arrow, representing the SSC protocol with equal

ITIs of 11 min 10s. **B,** Schematics of the fear conditioning procedures. Top: tests that were

conducted in Context A. Bottom: tests that were conducted in Context B. **C,** Schematics of the

fear conditioning protocols for SSC (n = 10, ITI = 11 min 10 s) and OSC (n = 8) groups.

Following a habituation (Hab.) tone, rats received 4 trials of a CS (3 kHz tone, 30 s) that co-

terminated with a US (footshock, 0.5 mA, 0.5 s). **D,** Freezing levels during CS presentations of

each group across the experiment; '×' denotes the pre-tone freezing levels. Two-way repeated-

measure ANOVA for each day followed by Holm-Sidak's pairwise post hoc test, * $p<0.05$. Inset:

the average freezing level during the first two CS presentations; Welch's t-test, * $p<0.05$.

Figure 5. **Immunohistochemistry quantification of pCREB in rats exposed to optimal and spaced short conditioning protocols**. **A,** Schematic of the immunohistochemistry procedure. **B,** Freezing levels during CS presentation for No-Shock control (NS, n = 4), Spaced Short

Conditioning (SSC, n = 6), and Optimal Short Conditioning (OSC, n = 5) groups during fear

conditioning; Two-way RM ANOVA followed by Holm-Sidak's post-hoc test. Letters *a, b*, and *c*

indicate pairwise post hoc tests with $p < 0.05$: *a*, OSC vs. SSC; *b*, SSC vs. NS; *c*, OSC vs. NS.

C, Average pCREB density in the lateral (LA), basolateral (BLA), and central (CeA) subregions

of the amygdala for NS, SSC, and OSC groups; Two-way RM ANOVA followed by Holm-Sidak's

post-hoc test. Letters *b* and *c* indicate pairwise post hoc tests with $p < 0.05$: *b*, SSC vs. NS; *c*,

OSC vs. NS. **D,** Average pCREB density in the paraventricular nucleus of the thalamus (PVT)

for NS, SSC, and OSC groups; One-way ANOVA followed by Tukey's post-hoc test. Letters

b and *c* indicate pairwise post hoc tests with $p < 0.05$: *b*, SSC vs. NS; *c*, OSC vs. NS. **E, *Left*,**

Average pCREB density across hippocampal subfields for NS, SSC, and OSC groups; Two-way

RM ANOVA followed by Holm-Sidak's post-hoc test. Letters *a* and *c* indicate pairwise post hoc

tests with $p < 0.05$: *a*, OSC vs. SSC; *c*, OSC vs. NS**; *Right*,** Representative images of pCREB

immunostaining in distinct subfields of the dorsal hippocampus (*top left*) in the NS group (*top*

right), SSC group (*bottom left*), and OSC group (*bottom right*).

Figure 6. Prediction of enhanced protocol for fear extinction. A, Schematic of the model.

B1, Simulated time courses of activated ERK (pERK, μM) and activated PKA (PKAc, μM) in

response to one trial of *Stim* (μM). **B2**, Simulated time courses of pERK, PKAc, and *inducer* in

response to 4-trial protocol with regular intervals of 2.5 min (short extinction, SE). **B3**, Simulated

time courses of pERK, PKAc, and *inducer* in response to 4-trial protocol with computationally

designed intervals (optimal short extinction, OSE). **B4**, Simulated time courses of pERK, PKAc,

and *inducer* (nM) in response to 12-trial protocol with regular intervals of 2.5 min (regular

extinction, RE).

Figure 7. The computationally designed protocol partially enhanced fear extinction in rats. **A.** Schematic of the enhanced fear extinction experimental procedures. Rats were trained

to constantly press a lever to retrieve sucrose pellets in a variable interval 60 (VI60) schedule,

where the average interval between each sucrose delivery is 60 s. **B,** Schematics of the fear

extinction protocols for SE (n = 11), OSE (n = 13), and FE (n = 12) groups, during which rats

received multiple trials of a CS (3 kHz tone, 30 s). **C,** Freezing levels during CS presentations of

each group across the experiment. No significant difference between groups was found by Two-

way RM ANOVA. '×' denotes the pre-tone freezing levels. **D,** Lever press rates during the first

two and the last CS presentation (*left*) or the 30 s before the first two and the last CS

presentation (pre-CS period) of each group. OSE group shows significant increase of pre-CS

lever presses comparing the last CS presentation to the average lever press rate of the first two

CS presentations. Paired Student's t-test. Letters *f* and *d* indicate tests with p < 0.05: *f*, RE, Last

CS vs. First 2 CS; *d*, OSE, Last pre-CS vs. First 2 pre-CS. **E,** OSE group shows higher lever

presses during pre-CS period compared to SE group in spontaneous recovery test. No

difference was found for the CS lever presses among the three groups ($H(2)$ = 2.102, P =

0.349). Kruskal-Wallis's test followed by Dunn's post hoc test. Letter *a* indicates pairwise

comparison with *p<0.05: OSE vs. SE.

Table 1. Statistical Analyses Results

Figure	Experiment Day	Description	Statistical test	Omnibus test	Omnibus p value	Post-hoc test	Post-hoc comparisons	Post-hoc p
2	Day 0	Baseline freezing	One-way ANOVA	F(2,37) = 0.018	0.982	n/a	n/a	n/a
		Baseline average speed	One-way ANOVA	F(2,37) = 0.062	0.939	n/a	n/a	n/a
		Baseline maximum speed	One-way ANOVA	F(2,37) = 0.044	0.956	n/a	n/a	n/a
	Day 2	Pre-tone freezing	Kruskal-Wallis	H(2) = 0.887	0.143	n/a	n/a	n/a
		Second CS freezing	Two-way RM ANOVA	F(2,37) = 3.751	0.033	Holm-Sidak	OSC vs. SC / OSC vs. RC / SC vs. RC	0.282 / 0.022 / 0.002
		Average speed	One-way ANOVA	F(2,37) =10.87	< 0.001	Tukey	OSC vs. SC / OSC vs. RC / SC vs. RC	0.024 / 0.131 / <0.001
	Day 3	Pre-tone freezing	Kruskal-Wallis	H(2) = 13.93	< 0.001	Dunn	OSC vs. SC / OSC vs. RC / SC vs. RC	>0.999 / 0.020 / 0.001
		First 2 CS freezing	Two-way RM ANOVA	F(22,407) = 1.169	0.271	n/a	n/a	n/a
		First 4 vs. last 4 CS freezing	Paired student's t-test	OSC: t(13) = 0.869 / SC: t(11) = 4.191 / RC: t(13) = 2.153	0.400 / 0.001 / 0.051	n/a	n/a	n/a
		Relative change in freezing	One-way ANOVA	F(2,37) = 3.315	0.047	Tukey	OSC vs. SC / OSC vs. RC / SC vs. RC	0.037 / 0.487 / 0.320
		Cue at half-reduced freezing	Kruskal-Wallis	H(2) = 7.08	0.029	Dunn	OSC vs. SC / OSC vs. RC / SC vs. RC	0.039 / >0.099 / 0.095
	Day 4	Extinction retrieval CTX A	Two-way RM ANOVA	F(2,37) = 0.097	0.907	n/a	n/a	n/a
		Average CS freezing	One-way ANOVA	F(2,37) = 3.944	0.028	Tukey	OSC vs. SC / OSC vs. RC / SC vs. RC	0.508 / 0.508 / 0.023

	Day	Measure	Test	Statistic	=		Post-hoc	Comparison	
	Day 5	Extinction retrieval CTX B	Two-way RM ANOVA	F(2,37) = 2.984		0.063	n/a	n/a	n/a
		Average CS freezing	One-way ANOVA	F(2,37) = 4.067		0.025	Tukey	OSC vs. SC OSC vs. RC SC vs. RC	0.437 0.236 0.020
	Day 29	Spontaneous recovery	Two-way RM ANOVA	F(2,37) = 3.259		0.049	Holm-Sidak	OSC vs. SC OSC vs. RC SC vs. RC	0.976 0.030 0.030
	Day 30	Spontaneous recovery	Two-way RM ANOVA	F(2,37) = 0.774		0.469	n/a	n/a	n/a
4	Day 0	Baseline freezing	Unpaired student's t-test	t(16) = 0.988		0.337	n/a	n/a	n/a
		Baseline average speed	Unpaired student's t-test	t(16) = 0.419		0.680	n/a	n/a	n/a
		Baseline maximum speed	Unpaired student's t-test	t(16) = 0.225		0.825	n/a	n/a	n/a
	Day 1	Freezing 2nd CS-US pairing	Two-way RM ANOVA	F(4,64) = 1.550		0.339	Holm-Sidak planned comparison	OSC vs. SSC	0.040
		Freezing 3rd CS-US pairing	Two-way RM ANOVA	F(4,64) = 1.550		0.339	Holm-Sidak planned comparison	OSC vs. SSC	0.019
	Day 3	First 2 CS freezing	Welch's t-test	t(12.43) = 2.227		0.045	n/a	n/a	n/a
		Freezing end of session (tone 12)	Two-way RM ANOVA	F(11,176) = 2.997		0.001	Holm-Sidak	OSC vs. SSC	0.987
	Day 4	Extinction retrieval CTX A	Two-way RM ANOVA	F(1,16) = 0.349		0.563	n/a	n/a	n/a
	Day 5	Extinction retrieval CTX B	Two-way RM ANOVA	F(1,16) = 1.130		0.304	n/a	n/a	n/a
	Day 29	Freezing first CS presentation	Two-way RM ANOVA	F(1,16) = 1.809		0.197	Holm-Sidak planned comparison	OSC vs. SSC	0.037
5	Day 0	Baseline freezing	One-way ANOVA	F(2,12) = 0.970		0.407	n/a	n/a	n/a
		Baseline average speed	One-way ANOVA	F(2,12) = 3.706		0.056	n/a	n/a	n/a
		Baseline maximum speed	One-way ANOVA	F(2,12) = 3.839		0.051	n/a	n/a	n/a

	Day 1	Freezing 2nd CS-US pairing	Two-way RM ANOVA	$F(8,48) = 1.976$	0.070	Holm-Sidak planned comparison	OSC vs. SSC SSC vs. NS OSC vs. NS	0.144 0.029 0.001
		Freezing 3rd CS-US pairing	Two-way RM ANOVA	$F(8,48) = 1.976$	0.070	Holm-Sidak planned comparison	OSC vs. SSC SSC vs. NS OSC vs. NS	0.312 0.002 <0.001
		Freezing 4th CS-US pairing	Two-way RM ANOVA	$F(8,48) = 1.976$	0.070	Holm-Sidak planned comparison	OSC vs. SSC SSC vs. NS OSC vs. NS	0.020 0.037 <0.001
		Lateral amygdala immunohistochemistry	Two-way RM ANOVA	$F(4,24) = 2.844$	0.046	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.838 <0.001 <0.001
		Basal amygdala immunohistochemistry	Two-way RM ANOVA	$F(4,24) = 2.844$	0.046	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.420 0.085 0.028
		Central amygdala immunohistochemistry	Two-way RM ANOVA	$F(4,24) = 2.844$	0.046	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.402 0.010 0.059
		PVT immunohistochemistry	One-way ANOVA	$F(2,12) = 16$	< 0.001	Tukey	OSC vs. SSC SSC vs. NS OSC vs. NS	0.681 <0.001 0.002
		Hippocampus subfields immunohistochemistry	Two-way RM ANOVA	$F(3,36) = 55.99$	< 0.001	Holm-Sidak	CA1 vs. DG CA2 vs. DG CA3 vs. DG	<0.001 <0.001 <0.001
		CA1 immunohistochemistry	Two-way RM ANOVA	$F(6,36) = 2.487$	0.041	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.860 0.416 0.416
		CA2 immunohistochemistry	Two-way RM ANOVA	$F(6,36) = 2.487$	0.041	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.811 0.153 0.187
		CA3 immunohistochemistry	Two-way RM ANOVA	$F(6,36) = 2.487$	0.041	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.968 0.465 0.465
		DG immunohistochemistry	Two-way RM ANOVA	$F(6,36) = 2.487$	0.041	Holm-Sidak	OSC vs. SSC SSC vs. NS OSC vs. NS	0.049 0.059 <0.001

7	Day 8	Fear conditioning freezing and lever presses	One-way ANOVA	F(14,231) = 0.404	0.973	n/a	n/a	n/a
		Freezing levels	Kruskal-Wallis	H(2) = 0.188	0.910	n/a	n/a	n/a
		Average speed	One-way ANOVA	F(2,33) = 0.042	0.958	n/a	n/a	n/a
		Lever press rate	One-way ANOVA	F(2,33) = 0.480	0.622	n/a	n/a	n/a
	Day 9	CS lever press rate beginning to end	Paired student's t-test	SE: t(10) = 1.000 OSE: t(12) = 1.237 RE: t(11) = 2.837	0.341 0.239 0.016	n/a	n/a	n/a
		Pre-CS lever press rate beginning to end	Paired student's t-test	SE: t(10) = 1.111 OSE: t(12) = 3.773 RE: t(11) = 0.994	0.293 0.003 0.341	n/a	n/a	n/a
	Day 10	CS freezing	Two-way RM ANOVA	F(2,33) = 0.207	0.814	n/a	n/a	n/a
		CS lever press rate	Two-way RM ANOVA	F(2,33) = 0.069	0.933	n/a	n/a	n/a
	Day 35	CS freezing	Two-way RM ANOVA	F(2,33) = 0.535	0.591	n/a	n/a	n/a
		CS lever press rate	Two-way RM ANOVA	F(2,33) = 1.085	0.350	n/a	n/a	n/a
		pre-CS lever press rate (tone 1)	Two-way RM ANOVA	F(2,33) = 2.897	0.069	Holm-Sidak planned comparison	OSE vs. SE OSE vs. RE SE vs. RE	0.005 0.019 0.516
		Average pre-CS lever press rate	Kruskal-Wallis	H(2) = 6.414	0.041	Dunn	OSE vs. SE OSE vs. RE SE vs. RE	0.035 0.531 0.709