1 **A circular zone of attachment to the extracellular matrix provides directionality to the**
2 **motility of *Toxoplasma gondii* in 3D**
3
4
5

6 Rachel V. Stadler[1], Shane R. Nelson[2], David M. Warshaw[2], Gary E. Ward[1]*
7
8 [1] Department of Microbiology and Molecular Genetics, and
9 [2] Department of Molecular Physiology and Biophysics,
10 University of Vermont Larner College of Medicine, Burlington, Vermont, U.S.A. 05405
11
12 * Correspondence to: Gary.Ward@uvm.edu
13
14

<u>Abstract</u>

Toxoplasma gondii is a protozoan parasite that infects 30-40% of the world's population. Infections are
typically subclinical but can be severe and, in some cases, life threatening. Central to the virulence of *T.*
gondii is an unusual form of substrate-dependent motility that enables the parasite to invade cells of its
host and to disseminate throughout the body. A hetero-oligomeric complex of proteins that functions in
motility has been characterized, but how these proteins work together to drive forward motion of the
parasite remains controversial. A key piece of information needed to understand the underlying
mechanism(s) is the directionality of the forces that a moving parasite exerts on the external environment.
The linear motor model of motility, which has dominated the field for the past two decades, predicts
continuous anterior-to-posterior force generation along the length of the parasite. We show here using
three-dimensional traction force mapping that the predominant forces exerted by a moving parasite are
instead periodic and directed in towards the parasite at a fixed circular location within the extracellular
matrix. These highly localized forces, which are generated by the parasite pulling on the matrix, create a
visible constriction in the parasite's plasma membrane. We propose that the ring of inward-directed force
corresponds to a circumferential attachment zone between the parasite and the matrix, through which the
parasite propels itself to move forward. The combined data suggest a closer connection between the
mechanisms underlying parasite motility and host cell invasion than previously recognized. In parasites
lacking the major surface adhesin, TgMIC2, neither the inward-directed forces nor the constriction of the
parasite membrane are observed. The trajectories of the TgMIC2-deficient parasites are less straight than
those of wild-type parasites, suggesting that the annular zone of TgMIC2-mediated attachment to the
extracellular matrix normally constrains the directional options available to the parasite as it migrates
through its surrounding environment.

Introduction

Parasites of the phylum Apicomplexa cause pervasive human disease and are responsible for approximately one million deaths annually [1, 2]. *Toxoplasma gondii* is the most prevalent apicomplexan parasite, infecting 30-40% of the human population worldwide [3-5]. *T. gondii* infections are typically subclinical in immunocompetent individuals but may cause blindness [6, 7]. In immunocompromised individuals and the developing fetus, the disease can be life-threatening [5, 8, 9].

T. gondii is a single-cell protozoan, and the pathogenesis associated with acute infection is caused by the highly motile tachyzoite life-cycle stage. Like other apicomplexan parasites, the *T. gondii* tachyzoite uses a unique form of substrate-dependent motility to invade into and egress from cells of its host, to migrate across biological barriers, and to disseminate throughout the infected host. Motility is, therefore, central to the pathogenesis of toxoplasmosis [10-12]. The parasite has no cilia or flagella and can move at speeds of up to 3 μm/s without the leading edge protrusions that drive the substrate-dependent motility of other eukaryotic cells. Within a three-dimensional (3D) extracellular matrix, the parasite moves along a complex helical trajectory characterized by regular oscillations in velocity and changes in trajectory curvature and torsion [13].

Motility is driven, at least in part, by a complex of proteins known as the glideosome, which is anchored to the inner membrane complex at the parasite periphery (Figure 1A). In the "linear motor" model of motility, short actin filaments are bound through the bridging protein TgGAC to the cytosolic tails of transmembrane adhesins, such as TgMIC2, and translocated towards the posterior end of the parasite via a class XIV myosin, TgMyoA [14-17]. If the extracellular domains of the transmembrane adhesins are attached to ligands on the substrate, the rearward translocation of actin results in forward movement of the parasite. Other apicomplexan parasites, including those that cause malaria (*Plasmodium spp.*) and cryptosporidiosis (*Cryptosporidium spp.*), express similar glideosome proteins, suggesting that the mechanism of motility is conserved [18].

Although abundant evidence supports the importance of TgMyoA and other components of the glideosome in motility [12, 16, 19-23], several recent observations have raised questions about whether these proteins generate force as described by the linear motor model and, if so, whether this is the only motility mechanism available to the parasite [19, 24-28]. Because parasite motility is essential for virulence, elucidation of the mechanism(s) underlying motility is of both fundamental cell biological interest and potential clinical relevance. One key piece of missing information needed for a full mechanistic understanding of motility is the directionality of the forces the parasite exerts on the external environment as it moves along its 3D helical trajectory. The linear motor model predicts continuous anterior-to-posterior forces along the entire periphery of the moving parasite. We show here using 3D traction force mapping that the predominant forces exerted by the parasite are instead organized into a

71 highly localized circumferential ring of inward-directed force at a fixed location within the matrix. These
72 inward-directed forces create a visible constriction of the parasite's plasma membrane, reflecting a
73 circular zone of tight attachment of the parasite to the matrix through which the parasite passes through as
74 it progresses forward along its helical trajectory. In the absence of this circular attachment zone, the
75 parasite is still able to move but this movement loses much of its directionality.
76
77 **Results**
78
Development of a quantitative 3D traction force mapping assay
As a first step towards visualizing the forces exerted by parasites moving in 3D, we used a bead
displacement assay in the well-established Matrigel extracellular matrix model [13, 27, 29]. Fluorescent
microspheres embedded within the Matrigel were pulled towards a moving parasite from nearby anterior,
posterior, and lateral locations within the matrix, and these microspheres returned to their original
positions after the parasite had passed (Figure 1B, Videos 1&2). These results demonstrate that the
parasite does indeed exert a detectable force on the surrounding matrix. Unexpectedly, we also noted
constrictions in the body of the fluorescent parasite as it moved (Video 2). These constrictions were even
more apparent by brightfield microscopy (Figure 1C, Video 3). Each constriction formed at the apical end
of the parasite and remained stationary relative to the matrix as the parasite moved through, reminiscent
of the "moving junction" through which the parasite penetrates during host cell invasion (discussed
further below).
To develop a more quantitative force mapping assay and explore the relationship between the
constrictions and matrix deformation, we used Alexa-Fluor 647-conjugated fibrinogen [30] to generate a
fluorescent 3D fibrin matrix (Figure 2A and Video 4). The use of a fluorescent matrix enabled mapping of
matrix displacements at all spatial points in our imaged volume rather than at the limited number of
discrete positions offered by fluorescent microspheres. Parasite motility in matrices generated using 2.25
mg/ml fibrinogen was indistinguishable from that observed in Matrigel (Figure 2B and Suppl. Figure 1A-
B). Visual comparison of successive image volumes confirmed that parasites moving through fluorescent
fibrin also deform the matrix (Figure 2C and Video 5) and undergo constrictions (see below) similar to
those observed in Matrigel.
To calculate and map the deformations of the fluorescent fibrin matrix, we used the Fast Iterative
Digital Volume Correlation (FIDVC) algorithm developed by Christian Franck's group
(https://github.com/FranckLab/FIDVC) to iteratively determine the 3D fibrin displacement fields between
consecutive volumetric image stacks in our time series [31]. Each image volume is divided into 16,807
subvolumes (49 x 49 x 7 subvolumes in x, y, z). The algorithm compares the voxel intensity pattern

between corresponding subvolumes at two consecutive timepoints, calculating one 3D displacement

vector per subvolume. The data can be displayed as a map for the entire imaging volume or for any 2D

plane (*x-y*, *x-z*, or *y-z*; see Video 6). Most of the 2D images below show the displacement maps in the *x-y*

plane on one of the seven *z* subvolume levels.

To filter out background noise and identify displacements caused by the parasite, we calculated

displacement vectors during periods when no parasites were moving within our imaging volume and used

these to establish a displacement detection threshold for each dataset (see Methods and Figure 3 for

details). Using this approach, the limit of detection for 3D displacement magnitudes in the system ranges

from 42 - 46 nm; a similar 3D displacement detection threshold (43 - 46 nm) was observed in samples

containing no parasites (see Figure 6E below), validating the approach. When analyzing data in the *x-y*

plane only, the background cutoff was calculated using the *x-y* displacement magnitudes (rather than the

x-y-z displacement magnitudes); this lowered the displacement detection threshold to 28 - 31 nm.

In order to translate matrix displacements into the magnitude of forces generated by the parasite, we

determined the viscoelasticity of the fibrin matrix using laser trapping (see Methods). The fibrin gel

behaves as a predominantly elastic matrix (Suppl. Figure 2) with an elastic modulus of 15.6 pN/µm and a

viscous modulus of 0.09 pN*s/µm. The minimal detectable force magnitude in this system is therefore

0.66-0.72 pN in 3D and 0.45-0.49 pN in the *x-y* plane.

123   Parasite constrictions are tightly linked to periodic bursts of matrix deformation and motility

With this system for 3D traction force mapping in hand, we analyzed the pattern and directionality of

forces produced by wild-type parasites as they moved through the fibrin matrix. Plotting fibrin

displacement as *x-y* maps showed that the matrix deforms in towards the parasite from all *x-y* directions at

discrete, periodic time points along the trajectory, before relaxing back to its initial position (*e.g.*, Figures

4A-D and Videos 7, 8). Viewing the data as projections along each axis demonstrates that the matrix is

simultaneously pulled in towards the parasite in all three dimensions (Suppl. Figure 3 and Video 6). In

approximately 50% of cases, inward deformation occurred between two consecutive image volumes (1.6

seconds apart) and was followed in the next image volume by relaxation (*e.g.*, Figure 4A-C). In most

other cases, the inward deformation was held through multiple time points before relaxing (Suppl. Figure

4). During periods of motility when the matrix was not being pulled inward or relaxing outward, little

deformation of the matrix above background was detectable (*e.g.*, Figure 4C, leftmost and rightmost

panels). This observation suggests either that the parasite pulls on the matrix only during some portions of

its trajectory or that the parasite produces both strong and weak deformations, and the weak deformations

– which could be continuous – are below our level of detection.

When the displacement maps were overlaid onto the images of the moving parasites, it was
immediately apparent that the timing of each large matrix deformation coincided with the presence of a
parasite constriction and that the displacement vectors pointed primarily in towards the constriction
(Figure 4C, D; additional examples are shown in Suppl. Figure 5). The fact that the displacements were
directed inwards rather than away from the parasite indicates that the pinching of the parasite membrane
results from pulling forces generated by the parasite rather than the parasite squeezing through a narrow
pore within the matrix, which would push the matrix away. When a constriction persisted over multiple
time points, the fibrin continued to deform into the constriction and/or to hold the deformation over those
same time points as the parasite passed through (*e.g.*, Figure 4D).

The time it took a parasite to move completely through a constriction ranged from 1.6-56.2 seconds
(average 6.8 ± 5.2 seconds, measured in brightfield; n=99 parasites, 188 constrictions) with the longer
times due to a stall in forward progression of a constricting parasite. The time between one constriction
finishing and another beginning was also variable (Figure 5A). In 46% of the cases, the end of one
constriction was followed immediately by the beginning of another, while in 14% of the cases a second
constriction would start before the first constriction finished (Figure 5A, B and Video 9). Most
importantly, all parasites that move at least one body length undergo a constriction and the total trajectory
length is directly proportional to the number of constrictions observed (Figure 5C), strongly suggesting
that the parasite-generated forces that create the constriction play an important role in forward movement.

The much reduced width of a constricting parasite (2.6 ± 0.4 μm *vs.* 1.4 ± 0.2 μm at the parasite's
widest point before and during a constriction, respectively; Figure 5D, n=15 parasites, 19 constrictions)
led us to question whether the constriction presents a physical obstacle to the forward motion of the
nucleus (normal nuclear diameter measured by DNA staining = 2.0 ± 0.3 μm). In parasites labeled with
both a fluorescent DNA stain and a fluorescently-labeled antibody against the major surface protein,
TgSAG1 (to visualize the constriction), the normally round parasite nucleus did indeed become thinner
and more elongate as it passed through the constriction and regained its round shape once it was through
(Figure 5E). The ratio of the nuclear dimensions perpendicular and parallel to the long axis of the parasite
was 0.93 ± 0.08 in a non-constricting parasite and 0.50 ± 0.06 during a constriction (n=9 parasites, 15
constrictions).

Interestingly, the distribution of the fluorescent anti-TgSAG1 antibody on the parasite surface
changed as the parasite moved: much of the fluorescent signal appeared to be "swept" from anterior to
posterior at the constriction as the parasite moved through (Suppl. Figure 6 and Video 10). The
redistributed antibody was ultimately shed and deposited behind the moving parasite in the form of a
fluorescent helical trail within the Matrigel (Suppl. Figure 6 and Video 10).

Parasites that do not generate constrictions do not detectably deform the matrix

Since TgMyoA is thought to play a central role in force production during parasite motility (Figure 1A), we conducted force mapping experiments using a parasite line lacking TgMyoA [20]. As previously reported [27], these parasites are much less motile in 3D than wild-type parasites: only 18 of 769 TgMyoA knockout parasites examined (2%) moved one body length or more during a 302-second time course, compared to 99 of 278 wild-type parasites (36%). Of the 18 TgMyoA knockout parasites that moved at least one body length, none showed clear constrictions at a fixed point within the matrix like those seen in wild-type parasites (Figure 6A). The pattern of movement was also different from wild-type parasites; rather than moving along smooth helical trajectories, the TgMyoA knockout parasites moved in tightly twisting arcs or in a stair-like pattern consisting of right-angled turns associated with a sharp bend in the body of the parasite (Figure 6A and Video 11). In stark contrast to the wild-type parasites, we also saw no evidence by traction force mapping of a ring of inward-directed forces produced by the TgMyoA knockout parasites. In fact, we did not detect any displacement of the matrix above background noise by these parasites (Figure 6B and Suppl. Figure 7, n=16 parasites), suggesting that any forces produced by the moving knockout parasites are below the limit of detection in our traction force assay.

We also tested whether the surface adhesin, TgMIC2, plays a role in generating the constriction-associated inward forces on the matrix. The motility defect in parasites lacking TgMIC2 [29] is less severe than the defect in parasites lacking TgMyoA, with 36 of 297 TgMIC2 knockout parasites (12%) capable of moving more than one body length (see also [29, 32]). However, these parasites often moved in an erratic, stop and start pattern, bending sharply and moving in tightly twisting arcs (Figure 6A and Video 12). The TgMIC2 knockout parasites also failed to form constrictions and did not detectably deform the fibrin matrix (Figure 6C, n=36 parasites, Suppl. Figures 8A-C, n=18 parasites) despite being able to move at maximum speeds similar to wild-type parasites [29]. Taken together, these data confirm that motility is significantly altered, but not completely eliminated, in parasites lacking TgMyoA or TgMIC2. Furthermore, the data demonstrate that both proteins are required to produce the constriction and the constriction-associated inward forces on the matrix observed during the motility of wild-type parasites.

TgMIC2-deficient parasites move less directionally

The larger number of TgMIC2 knockout parasites capable of long runs of motility (compared to the MyoA knockouts) enabled us to analyze their phenotype in greater detail. The nuclei of parasites lacking TgMIC2 did not change shape as they moved in 3D (Figure 7A and Suppl. Figure 8D), providing further evidence that these parasites do not undergo the constrictions seen in wild-type parasites. Given their apparent propensity for tight turns, we also quantified the ability of the TgMIC2 knockout parasites to

move directionally, by comparing displacement distance (first to last point) to total trajectory length.

While parasites with and without TgMIC2 traveled along trajectories of similar mean length (38 ± 9 µm

vs. 42 ± 8 µm, respectively), parasites lacking TgMIC2 moved approximately half as far from their

starting point as wild-type parasites (Figure 7B and Suppl. Figure 8E). These data suggest that the

circumferential zone of attachment to the matrix, which requires TgMIC2, functions to convert

meandering motility into more linear, directed forward progression.

<u>Discussion</u>

The linear motor model of apicomplexan motility predicts that a parasite moves by pulling

continuously on the matrix in an anterior-to-posterior direction along the length of the parasite. It was,

therefore, unexpected that the predominant forces revealed by our new 3D traction force mapping assay

were instead periodic and highly localized, pointing inwards from all directions towards a circumferential

location on the parasite surface. This annular band of parasite-generated, inward-directed force forms at a

fixed position within the matrix and creates a visible constriction at the parasite periphery. The

constriction passes from the anterior to the posterior end of the parasite as the parasite moves forward.

Typically, when the constriction reaches the parasite's posterior end a new constriction forms at the apical

end and the cycle repeats. These results suggest that the moving parasite periodically assembles ring-

shaped zones of tight attachment to the matrix, which deform the matrix and serve as stationary platforms

that the parasite propels itself through, one body length at a time.

Attachment to the substrate, force generation, and motility in 2D *vs.* 3D. Both malaria

sporozoites and *T. gondii* tachyzoites are capable of moving in circles on 2D surfaces. The pioneering

work of Munter, Frischknecht and colleagues combined traction force mapping with reflection

interference contrast microscopy (RICM) during the 2D circular gliding of malaria sporozoites [33]. They

concluded that the parasite adheres to the substrate at its apical and posterior ends; each of these adhesion

sites produces stalling forces parallel to the long axis of the parasite, and disengagement of the adhesion

sites drives forward motion. Traction forces perpendicular to the direction of movement were also

detected at the center of the parasite but these were interpreted as nonspecific and nonproductive. The

apical and posterior adhesion sites were shown to move forward *with* the parasite, rather than remaining

stationary with respect to the substrate.

T. gondii tachyzoites demonstrate a second form of productive motility on 2D surfaces called helical

gliding: the banana-shaped parasite starts with its left side in contact with the substrate, moves forward

one body length while rotating along its longitudinal axis until it lies on its right side, and then flips back

to its left side to repeat the cycle [34]. This form of motility is the equivalent of the fully helical

trajectories seen in 3D but constrained in 2D by contact of the curved parasite with the rigid substrate. In

RICM studies of 2D helical gliding [21, 35] the tachyzoite was seen to first attach to the substrate at its

apical end. The zone of attachment then expanded along the length of the parasite and, as the parasite

moved forward and its apical end twisted up off the cover slip, the zone of attachment shrunk until it was

reduced to a spot at the posterior end. In contrast to *Plasmodium* sporozoites, the position of the

tachyzoite adhesion site remained fixed relative to the substrate and the parasite slid over this attachment

site as the parasite moved forward. 2D traction force mapping revealed forces parallel, but not

perpendicular, to the longitudinal axis of the tachyzoite; these traction forces, together with spring-like

forces generated by the parasite's microtubule cytoskeleton, were postulated to produce forward motion

[35].

When these previous studies of *T. gondii* 2D motility are combined with the 3D results presented

here, a consensus emerges that the basic unit of motility consists of apical attachment to the substrate,

followed by apical-to-posterior translocation of the adhesion site, which is fixed in its position relative to

the substrate. When the adhesion site reaches the posterior end of the parasite, a new apical attachment

site is formed, and the cycle begins again. The adhesion site in a 3D matrix is circular, and the pulling

forces exerted by the parasite on the matrix manifest as a visible constriction in the parasite's plasma

membrane. In 2D, no deformation of the parasite has been reported at the adhesion site, likely reflecting

differences in attachment to the substrate via a point/patch on the parasite surface *vs*. circumferential

attachment. We also saw no evidence in 3D for a persistent stall force at the rear of the parasite. The

biggest difference between the 2D and 3D force mapping data is that, in 3D, all force vectors point

towards one highly localized circumferential region on the surface of the parasite, and many of these

force vectors contained a strong component perpendicular to the long axis of the parasite.

Munera Lopez et al [36] also recently noted that tachyzoites show constrictions as they move through

Matrigel. These authors ascribed the constrictions to the parasite having to squeeze through small pores in

the matrix. Our force mapping results and the regularity with which the constrictions form at the apical

end of the parasite argue strongly that the constrictions do not result from the parasite pushing its way

through pores, but rather from parasite-generated pulling forces on the matrix. Annular constrictions that

remain stationary relative to the surrounding environment have also previously been reported along the

length of *Plasmodium* ookinetes as they move through peritrophic membrane and microvillar network of

the mosquito midgut [37-39], suggesting that parasite-generated circular zones of attachment to the

environment may be a conserved feature of apicomplexan motility in 3D.

3D motility and invasion: variations on a theme. During the invasion of host cells by *T. gondii* and

other apicomplexan parasites, proteins secreted from the parasite's apical organelles assemble a ring-

shaped zone of tight attachment between the membranes of the two cells. This "moving junction" does

not in fact move: it is anchored to the host cytoskeleton and is hypothesized to provide a fixed platform

against which the parasite exerts force [40-45]. Parasite surface adhesins, which are engaged with the

junction via their extracellular domains, are thought to be translocated by MyoA in an anterior-to-

posterior direction, pushing the parasite through the junction and into the host cell. The body of the

parasite narrows dramatically as it passes through the moving junction, bearing a striking resemblance to

the constrictions seen during 3D motility.

Two central players in moving junction formation in *T. gondii* are TgAMA1, an adhesin secreted onto

the parasite surface from the micronemes, and TgRON2, which is secreted by the parasites into the host

cell plasma membrane and serves the ligand to which TgAMA1 binds. Neither TgAMA1 [46, 47] nor

TgRON2 (RVS and GEW, unpublished data) plays any detectable role in parasite motility. Instead,

TgMIC2 – another microneme protein secreted onto the parasite surface – likely plays an analogous role

during motility to that of TgAMA1 during invasion, with TgMIC2 binding to ligands in the extracellular

matrix rather than to TgRON2 in the host cell membrane.

As the parasite moves forward during both invasion and 3D motility, it simultaneously rotates around

its long axis [13, 34, 48]. Our results provide a possible explanation for this rotational motion. In both

cases, parasite receptors bind to their cognate ligands in the form of a circular ring of attachment that is

fixed in position relative to the host cell (in the case of invasion) or the extracellular matrix (in the case of

3D motility). It has long been proposed that the TgMyoA motor driving invasion and motility is arranged

helically within the parasite, along the spiraling subcortical microtubules (*e.g.*, [35]). Alternatively, a

recent modeling study suggests that the unique geometry of the parasite periphery can generate a self-

organized helical pattern of anterior-to-posterior actin flow [49]. In either case, if the actin filaments flow

helically and are connected to the tails of transmembrane adhesins engaged circumferentially with the

host cell surface or extracellular matrix, the forward motion of the parasite either into the cell or through

the matrix will by definition be accompanied by rotational twist. Some portion of the perpendicular force

that we observed directed towards the annular constriction in our traction force assays may in fact reflect

torque on the matrix produced by the parasite as it twists through this circular zone of attachment.

As a parasite invades a host cell, antibodies are "shaved" off the parasite surface at the moving

junction by an as yet unknown mechanism [50]. Remarkably, a similar phenomenon is observed during

3D motility: fluorescent antibody against the surface protein TgSAG1 is lost from the parasite surface

anterior to the constriction and shed as a helical trail of fluorescence behind the moving parasite (Suppl.

Figure 6 and Video 10).

The numerous parallels between invasion and 3D motility noted here suggest that the two processes

may be more mechanistically similar than previously recognized [14, 21], with implications for our

understanding of each. For example, our data suggest that 3D motility can be best described as the

parasite undergoing sequential invasion-like events through circular zones of attachment to the matrix;

during invasion the parasite only needs to move a single body length to penetrate into the host cell, but

during motility the parasite can string multiple constriction events together to move multiple body lengths

(Figure 5C). The data also raise the possibility that the characteristic constriction of the parasite seen at

the moving junction during invasion does not reflect the parasite being squeezed as it *pushes* itself

through the small diameter opening into the host cell, as is commonly thought (*e.g.,* [51, 52]), but rather

the parasite attaching to and *pulling* on the host cell plasma membrane as it penetrates.

The linear motor model. The data presented here with wild-type parasites are at least partially

consistent with the linear motor model of motility, in which: (a) actin polymerization is nucleated at the

apical tip of the parasite by formin1 [21]; (b) TgMyoA translocates the actin filaments in an anterior-to-

posterior direction along the length of the parasite; and (c) binding of the rearward flowing actin filaments

to transmembrane adhesins [16], whose extracellular domains are attached to the substrate, drives the

forward motion of the parasite (Figure 1A). While anterior-to-posterior actin flux may occur along the

entire length of the parasite periphery [49, 53, 54], we propose a modification to the linear motor model

in which force is generated, in whole or in part, by the rearward translocation of the subset of actin

filaments that are coupled to adhesins at the circular ring of attachment. We were unable to detect any

anterior-to-posterior bias in the force vectors at the constriction, which would be expected if the parasite

pulls against this zone of attachment in order to move forward. These forces may be below our limit of

detection (~0.72 pN). Alternatively, compression of the fibrin immediately adjacent to the constriction

might increase the local elastic modulus of the matrix, making it more difficult to see proximal

displacements because of a local increase in the displacement detection threshold. Because the fibrin gel

behaves primarily as an elastic matrix (Suppl. Figure 2), the relationship between acceleration and force is

complex and we cannot calculate with any confidence how much total displacement / force would be

required to achieve a given parasite velocity and relate this to the displacement detection threshold of the

assay.

The regularity with which the parasite generates the circular zones of attachment on its surface,

yielding on average one constriction per body length of motion (Figure 5D), could be due to periodic

bursts of formin1-mediated actin polymerization at the parasite's apical tip [21] or to periodic release of

adhesins such as TgMIC2 from the micronemes. Alternatively, the observation that constrictions occur

back-to-back almost 50% of the time (Figure 5A), may suggest that a new constriction forms at the apical

tip before the prior constriction has ended, but this new constriction stalls as the TgMyoA (or possibly

TgMyoH [21]) to which it is connected pulls against the adhesive forces of the more posterior previous

constriction. Future real-time analysis of actin polymerization and microneme secretion in motile

parasites, together with higher sensitivity force mapping, will be necessary to further interrogate the

periodic nature of ring assembly.

The lack of a motility-associated constriction in TgMyoA-deficient parasites could reflect an
inability either to assemble ring of adhesins at the apical tip of these parasites or, more likely, to move
through the incipient ring once it has formed. Alternatively, it could be that the ring-shaped structures that
form during motility are contractile, and contractility requires TgMyoA. The inward deformation of the
fibrin matrix towards the constriction reported here is consistent with a contractile ring, as is the physical
constriction of the parasite plasma membrane. Intriguingly, the speed of the parasite's forward motion
increases once the constriction passes the halfway point on the parasite's longitudinal axis (Suppl. Figure
9, n=20 parasites, 24 constrictions). This observation suggests that physical squeezing of the tapering
posterior end of the parasite by the narrow (and potentially contractile) constriction may contribute to the
parasite's forward motion.
If the coupling of actin flow to the circular zone of attachment is important for forward motion, how
do the parasites lacking TgMIC2 that move do so at near normal speeds [29] in the absence of
constrictions? First, it is possible that both the wild-type and mutant parasites generate small forces
distributed over their entire surface (in addition to larger forces that may be generated at the constriction
in wildtype), which are collectively sufficient to drive motility but individually below our level of
detection. Alternatively, the loss of TgMyoA or TgMIC2 might lead to compensatory changes in the
expression of other genes with overlapping or redundant function [25, 55, 56]. The TgMIC2 KO parasites
do in fact show changes in the expression of several other microneme and motility-associated proteins
[29]. Finally, it is possible that parasites lacking TgMyoA and/or TgMIC2 use a motility mechanism that
is entirely different from that of wild-type parasites (*e.g.*, [20, 24]) and does not involve a constriction but
is nevertheless capable of supporting motility and sustaining the parasite's lytic cycle. By analogy with
observations from animal cells [57], parasites might use different mechanisms for motility in different
situations such as squeezing through tight junctions [10] *vs.* migrating through loose interstitial tissues. If
the motility of the mutants is driven by a different mechanism than that of wild-type parasites, higher
sensitivity 3D traction force mapping may reveal informative differences in their pattern of force
generation.
A guidance system for motility? The forward motion of the TgMIC2 knockout parasites, which
lack the circular attachment zone, is significantly more disorganized than the motility of wild-type
parasites. The mutant parasites can still generate relatively long trajectories, but the trajectories take them
less far from their starting point than wild-type parasites. As discussed above, attachment to the matrix via
a circular ring of adhesion likely contributes to the rotational twisting of the parasite as it moves forward
and thereby to the helicity of the parasite's trajectory. Extensive studies of the helical swimming behavior
of bacteria has shown that a curved cell shape and helical trajectory are a particularly efficient way for
small cells to move through viscous media [58-62]. Engagement of *T. gondii* with the extracellular matrix

through a circular band of adhesion may therefore not only create a fixed platform for the parasite to pull

against and move forward; it may also function as part of a guidance system to help the parasite move

efficiently through the various environments it encounters as it disseminates through the tissues of the

unfortunate organisms it infects.

<u>Methods</u>

Parasite and Cell Culture. Parasites were propagated by serial passage in human foreskin fibroblasts

(HFFs). HFFs were grown to confluence in Dulbecco's Modified Eagle's Medium (DMEM) (Life

Technologies, Carlsbad, CA) containing 10% (vol/vol) heat-inactivated fetal bovine serum (FBS) (Life

Technologies, Carlsbad, CA), 10 mM HEPES pH 7, and 100 units/ml penicillin and 100 ug/ml

streptomycin, as previously described [63]. Prior to infection with *T. gondii*, the medium was changed to

DMEM supplemented with 10mM HEPES pH 7, 100 units/ml penicillin and 100 ug/ml streptomycin, and

1% (vol/vol) FBS.

Pitta Chamber Assembly. 22×22 coverglasses were washed with Alconox detergent, rinsed with tap

water, deionized water, and ethanol (100%), and air dried. Two strips of double-sided tape (Scotch 3M,

St. Paul, MN) were placed 3 mm apart on a glass slide, and the coverglass was placed on top of the tape

and pressed firmly to ensure a complete seal. The flow cell volume was approximately 10 µl.

Matrigel Matrix. Pitta chambers containing parasites embedded in polymerized Matrigel were prepared

as previously described [13]. Briefly, parasites were harvested from infected HFFs via syringe release and

passed through a 3 µm Nucleopore filter (Whatman, Piscataway NJ). Parasites were pelleted and

resuspended at in Live Cell Imaging Solution (LCIS) buffer. For fluorescence imaging, the LCIS

contained 0.5 mg/ml Hoechst 33342 (Thermo Scientific, Waltham, MA) and/or a 1:20 dilution of

Alexa546-conjugated anti-TgSAG1 (see Table 1). Fluorescent anti-TgSAG1 (100µg) was prepared using

the Alexa Fluor 546 Antibody Labeling Kit (Thermo Scientific, Waltham, MA) as per the manufacturer's

instructions. Parasites were incubated with the Hoechst 33342 (10 minutes) and/or anti-TgSAG1(15

minutes) at room temperature, then mixed with Matrigel and LCIS on ice in a 1:3:3 (vol/vol/vol) ratio and

immediately added to a Pitta chamber. The Pitta chamber was incubated for three minutes at 35°C in a

custom-build heated microscope enclosure (UVM Instrumentation and Model Facility, Burlington, VT)

before imaging.

Fibrin Matrix. Unlabeled and Alexa-Fluor 647-labeled fibrinogen (both from Thermo Scientific,

Waltham, MA) were each dissolved in 0.1M sodium bicarbonate (pH 8.3) to 15 mg/ml. Aliquots of 50 µl

were flash frozen and stored at -80C until use. Fibrin gels [30] containing parasites were prepared in Pitta

chambers as follows. The parasite culture medium was replaced with LCIS buffer before harvesting via

syringe release and filtering. 500 µl of the parasite suspension were pelleted and resuspended in LCIS

buffer to achieve a higher parasite concentration. Parasites were mixed with fibrinogen (final fibrinogen

concentration of 2.25, 4.5, or 9 mg/ml). LCIS buffer containing thrombin (Sigma, Burlington, MA) and

FBS was added to the parasite-fibrinogen suspension for final concentrations of $1.5x10^8$ parasites per ml,

1 unit/ml of thrombin, and 10% FBS. The mixture was immediately pipetted into a Pitta chamber and

allowed to polymerize at room temperature for three minutes before mounting on the microscope and

imaging.

A Nikon A1R-ER point-scanning confocal microscope was used to visualize the porosity of the

fluorescent fibrin matrix (Suppl. Figure 1A), Using Galvano scanning, image (1024 × 1024 pixel) stacks

were captured with 60× Apo (0.1 µm/pixel, NA 1.49l) objective with 0.25 µm spacing over 25 µm. The

LUNV laser was used at wavelength 633 nm and the pinhole size was 28.10 µm.

Image Acquisition. All other imaging was done on a Nikon Eclipse TE300 widefield epifluorescence

microscope (Nikon Instruments, Melville, NY) equipped with a NanoScanZ piezo Z stage insert (Prior

Scientific, Rockland, MA). See Table 1 for details of the imaging setup. Time-lapse video stacks were

collected with an iXON Life 888 EMCCD camera (Andor Technology, Belfast, Ireland) using NIS

Elements software v.5.11 (Nikon Instruments, Melville, NY). Fluorescently labeled parasites were

imaged using a pE-4000 LED illuminator (CoolLED, Andover England) and a 89402 Quad filter

(Chroma, Bellows Falls, VT). Stacks consisting of individual images (1024 pixel× 384 pixel) captured

0.5-1 µm apart in z, covering a total of 10-40 µm, were collected using either a 20× PlanApo λ (0.65

pixel/µm, NA 0.75) or 60 × PlanApo λ (0.22 pixel/µm, NA 1.4) objective as described in Table 1. The

same volume was successively imaged 60-360 times over the course of 64-302 seconds. The camera was

set to trigger mode, no binning, readout speed of 35 MHz, conversion gain of 3.8x, and EM gain setting

of 300.

Table 1: Imaging parameters for the different experiments described

Experiment	Objective	Fluorochrome (Excitation/emission wavelengths)	Image spacing in z	Exposure time per image	Number of Image stacks	Total time	Volume (x, y, z)

Microspheres (Matrigel)	60×	DragonGreen (490/507-530nm) tdTomato parasites (550/579-608 nm)	41 slices, 1μm apart	16 ms	60	64s	225.3 μm × 84.5 μm × 40 μm
Fibrin vs Matrigel and TgMIC2 KO directionality	20×	Hoechst 33342 (385/420-449 nm)	41 x 1 μm	16 ms	120	80s	665.6 μm × 249.6 μm × 40 μm
Force Mapping, WT (Fibrin)	60×	tdTomato parasites (550/579-608 nm) Fibrin (635/666-723 nm)	50 x 0.5 μm	16 ms	60	96s	225.3 μm × 84.5 μm × 24.5 μm
WT, TgMyoA KO, TgMIC2 KO; Brightfield (Matrigel, fibrin)	60×	N/A: Brightfield	21 x 1 μm	40 ms	360	302s	225.3 μm × 225.3 μm × 20 μm
Nuclear size vs constriction (Matrigel)	20×	Hoechst 33342 (385/420-449 nm) Anti-SAG1 Alexa 548 (550/579-608 nm)	41 x 1 μm	16 ms	60	80s	665.6 μm × 249.6 μm × 40 μm
TgMyoA KO Force Map (Fibrin)	60×	Hoechst 33342 (385/420-449 nm) Fibrin (635/666-723 nm)	50 x 0.5 μm	16 ms	60	96s	225.3 μm × 84.5 μm × 24.5 μm
TgMIC2 KO Force Map (Fibrin)	60×	YFP cytosol (490/507-530 nm) Fibrin (635/666-723 nm)	50 x 0.5 μm	16 ms	60	96s	225.3 μm × 84.5 μm × 24.5 μm

Tracking Parasite Motility. Datasets were analyzed in Imaris ×64 v. 9.2.0 (Bitplane AG, Zurich,

Switzerland). Fluorescently labeled parasite nuclei were tracked using the ImarisTrack module within a

1018 pixel × 380 pixel region of interest to prevent artifacts from tracking objects near the border. Spot

detection used an estimated spot diameter of $3.0 \times 3.0 \times 6.0$ µm (*x, y, z)*. A maximum distance of 6.0 µm

and a maximum gap size of 2 frames were applied to the tracking algorithm. Tracks with durations under

10 seconds or displacements of less than 2 µm were discarded to avoid tracking artifacts and parasites

moving by Brownian motion, respectively [13]. Accurate tracking was confirmed by visual inspection of

parasite movements superimposed on their calculated trajectories.

For the comparison between Matrigel and fibrin, we analyzed three biological replicates, each consisting

of three technical replicates. Student's T-tests were used to determine statistical significance between

fibrin and Matrigel samples.

Fibrin Deformation with FIDVC. The Fast Iterative Digital Volume Correlation (FIDVC) algorithm

was used to calculate 3D fibrin displacements [31] by comparing two consecutive image volumes. The

image volumes were cropped to 384×384 pixels (*x,y)* $\times 48$ slices (*z*) before running FIDVC. The initial

interrogation window size was set to $32 \times 32 \times 32$ and the program was run incrementally. The

displacements arrows were plotted using a combination of previously described MATLAB code for 2D

quiver color-coded for directionality [30] and custom MATLAB code. To filter out background signal, we

calculated the mean magnitude of the 16,807 displacement vectors for each timepoint comparison in each

dataset. The displacement detection threshold for that dataset was set as three standard deviations above

the lowest of these mean values.

Rheology. A Lumicks C-Trap laser trap was used to determine the viscoelasticity of the fibrin. First, the

power density spectrum (PDS) was calculated with 0.91 µm styrene beads in a flow cell. The beads were

embedded in a fibrin gel, and a bead was captured in the laser trap. The flow cell was oscillated on the

stage in the *y* dimension at different frequencies (1, 5, 10, 20, 50, 100 Hz), each for 5 seconds, at a fixed

amplitude of 100 nm. The trap position and force in the *y* dimension were captured at 78 kHz. The elastic

(in-phase) and viscous (out-of-phase) moduli were calculated using the bead's force trace. The data were

analyzed in R studio. A sliding two-sided window filter, size 78, was applied to both traces. The position

signal was fitted with the function $position = A * \sin(t * f - phase)$ using nonlinear least-squares with

fit parameters for amplitude (A), frequency (f), and phase. The force readout was fit with the function

$force = B * \sin(t * f - phase) + C * \cos(t * f - phase)$ using nonlinear least-squares with fitting

parameters B and C, where B is the in-phase (elastic) component of the force. The C fit parameter divided

by the velocity of the trap equals the viscous modulus.

Figure 1: **Parasites moving in 3D deform the surrounding Matrigel and undergo periodic constrictions**. **(A)** In the linear motor model of motility, the TgMyoA motor (TgMyoA and its associated light chains, TgMLC1 and either TgELC1 or TgELC2) is anchored to the parasite's inner membrane complex (IMC) via TgGAP45 and the transmembrane proteins TgGAP40 and TgGAP50. The lumenal portion of GAP50 is thought to interact with GAPM, a protein that spans the inner IMC membrane and likely connects the entire glideosome to the underlying parasite cytoskeleton. Short actin filaments located between the parasite plasma membrane and the IMC are connected to ligands on the substrate through a linker protein, possibly GAC, which binds to the cytosolic tails of surface adhesins such as TgMIC2. The TgMyoA motor displaces the actin filaments rearward; because the motor is connected to the IMC and the actin is connected to the substrate, this causes the parasite to move forward relative to the substrate. The vertical blue and red squiggles on TgGAP45 and TgMLC1 denote lipid anchors that attach the proteins to the IMC [23, 28, 64]). Figure modified from [28] and re-used here under the terms of a Creative Commons Attribution 4.0 International license (doi/10.1128/mSphere.00823-20). **(B)** Sequential time series images in a single *z*-plane demonstrate that Dragon Green-labeled microspheres embedded within the Matrigel are displaced towards moving tdTomato-expressing parasites; note displacement of the green bead closest to the fixed crosshair in the merged images as the parasite moves from top left to bottom right. See Video 1 for the entire time series, and Video 2 for a second example of bead displacement by a moving parasite. **(C)** Brightfield images showing a constriction progressing from the anterior to the posterior end of a parasite as the parasite moves forward one body length; see Video 3 for entire time series. All wild-type parasites analyzed (n=99) formed at least one constriction per body length of motility. Scale bars = 5 µm, timestamps in seconds.

Figure 2: **(A)** Confocal imaging of a 2.25 mg/ml fluorescent fibrin gel. A maximum fluorescence intensity projection of 51 *z*-slices captured 0.25 µm apart is shown; scale bar = 10 µm. **(B)** Maximum fluorescence intensity projections showing the trajectories of tdTomato-expressing wildtype parasites moving in 3D in Matrigel (top) *vs.* fibrin (bottom). See Suppl. Figure 1 for quantitative comparison of the motility parameters in the two matrices. Scale bar = 40 µm. **(C)** A parasite moving in a 1.3 mg/ml fluorescent fibrin matrix (see Video 4) visibly deforms the matrix, as evident from the non-coincident fluorescence signals in the highlighted area at the time points indicated (merge). Scale bar = 10 µm, timestamps in seconds.

Figure 3: **Determining threshold of detection in 3D traction force mapping. (A)** A fluorescent fibrin matrix containing wild-type parasites was imaged over 96 seconds (60 successive image volumes). FIDVC was then used to calculate the 16,807 3D fibrin displacement vectors between successive pairs of

time points. The magnitudes of the 16,807 *x-y* displacement vectors are plotted for each of the 59 pairwise timepoint comparisons. This dataset contained a single moving parasite, which moved intermittently during the first part of the time course but not at the end, as indicated. For each time course, the two consecutive timepoints that gave the lowest mean *x-y* displacement vector magnitude were used to set the background threshold for that dataset: any displacements less than three standard deviations (3SD) above this mean were considered background noise. **(B)** Expanded view of the 16,807 datapoints from the two time point comparisons shown in purple and green in panel (A): no parasite movement was seen between timepoints 6 and 7, while a single constricting parasite was seen moving between timepoints 7 and 8. Arrow indicates the 3SD noise detection threshold; all points above this threshold are colored blue in panels (A) and (B). **(C)** Displacement vectors with magnitudes greater than the 3SD threshold for the two highlighted timepoint comparisons were mapped back onto the imaging volume; the cluster of displacement vectors evident in the timepoints 7-8 comparison corresponds to the position of the single moving parasite in the volume.

Figure 4: 3D traction force mapping in fluorescent fibrin reveals that the matrix is periodically pulled in towards the constriction during parasite motility. **(A)** Sequential time series images in a single *z*-plane of a tdTomato-expressing parasite moving in fibrin (boxed). Scale bar = 10 μm, timestamps in seconds. **(B)** Force maps from the corresponding *z* subvolume plane of the data shown in (A); each map depicts the motions of the fluorescent fibrin matrix between the two consecutive timepoints, as indicated by the brackets. Arrow length (multiplied 24-fold for display) indicates the magnitude of matrix displacement and arrow color the directionality (see color wheel). **(C)** Zoomed images showing the force maps from the boxed region of panel B, after background subtraction (see text), overlaid on the parasite images from panel A. Length of arrows indicating displacement magnitude are multiplied 15-fold for display. Note inward displacement of the matrix in the second panel, outward displacement (relaxation) of the matrix in the third panel, and no detectable displacement vectors in the fourth panel even though the parasite continues to move. See Video 7 for real-time playback of the displacement vectors overlaid on the moving parasite. **(D)** Zoomed overlays of the force maps (background subtracted) and images of a moving parasite from a second dataset. The inwardly displaced matrix in this example did not relax back to its original position within the time frame of the experiment. Arrow lengths indicating displacement magnitude are multiplied 15-fold for display. See Video 8 for real-time playback of the displacement vectors overlaid on the moving parasite. Scale bars in C and D = 5 μm. Empty red arrowheads indicate position of the constriction.

Figure 5: The constrictions are tightly linked with motility and are sufficiently narrow to deform the parasite nucleus as it passes through. (A) The time interval between constrictions during motility of

individual parasites (n=188 constrictions, 99 parasites). A negative time interval (black bars) corresponds

to the presence of two constrictions in the same parasite for the indicated amount of time. Green bars

denote back-to-back constrictions and grey bars denote a delay between constrictions for the indicated

amounts of time. For 27% of the constrictions (not plotted), no subsequent constriction occurred within

the time frame of the experiment. **(B)** Example of a parasite with two concurrent constrictions. Brightfield

images; scale bar = 5 µm; timestamps in seconds. **(C)** Number of constrictions observed *vs*. distance

traveled, expressed in terms of binned parasite body lengths (*i.e.*, multiples of 7 µm). **(D)** Width of non-

constricting parasites halfway along their longitudinal axis *vs*. width at the constriction halfway along the

parasite's longitudinal axis (n=19 constrictions, 15 parasites). **(E)** Dimensions of the nucleus (determined

by Hoechst 33342 staining) in non-constricting parasites *vs*. nuclei passing through a constriction (n=9

parasites, 21 constrictions). Nuclear length and width are defined as the diameters parallel and

perpendicular to the long axis of the parasite, respectively. Samples were compared by Student's two-

tailed t-test.

<u>**Figure 6**</u>: **3D motility and force mapping of parasites lacking TgMyoA or TgMIC2.** **(A)** Brightfield

images showing TgMyoA knockout and TgMIC2 knockout parasites moving within a Matrigel matrix

without a detectable constriction. A wild-type parasite undergoing a typical constriction is shown for

comparison. The TgMyoA knockout parasite makes right-angled "stairstep" turn, and the TgMIC2

knockout parasite moves erratically in a tightly twisting arc. Scale bar = 5 µm, timestamps in seconds.

See Videos 11 and 12 for the entire time series for the TgMyoA and TgMIC2 knockout parasites. **(B-E)**

As in Figure 3A, the plots show the magnitudes of the 16,807 fibrin *x-y* displacement vectors between

each pair of successive image volumes generated in experiments using: (B) TgMyoA knockout parasites;

(C) TgMIC2 knockout parasites; (D) wild-type parasites; and (E) no added parasites (fibrin only). Periods

during the 96-second time course when parasites were moving are indicated. For each time course, the

two consecutive timepoints that gave the lowest mean displacement magnitude were used to set the

background threshold: any displacements less than three standard deviations (3SD) above this mean were

considered noise for that dataset. Examples of the force maps surrounding motile TgMyoA and TgMIC2

knockout parasites are shown in Suppl. Figures 7 and 8, respectively.

Figure 7: Knockout of TgMIC2 results in the loss of the constriction and less directional motility.

(A) Representative plots of nuclear shape (ratio of the nuclear diameters perpendicular *vs*. parallel to the

long axis of the parasite) in one wildtype and one TgMIC2 knockout parasite (black circles and green

squares, respectively), as the parasites moved over time. The wild-type parasite underwent multiple

concurrent constrictions along its trajectory. For the combined data from multiple parasites, see Suppl.

Figure 8D. **(B)** Ratio of parasite displacement (D) to trajectory length (L) for wildtype (n= 1399) and

TgMIC2 knockout (n= 165) parasites (black circles and green squares, respectively). The calculated

means (± SD) are from three independent biological replicates; see Suppl. Figure 8E. In panels (A) and

(B), "WT" refers to the TgMIC2 LoxP parasites [29] before treatment with rapamycin to excise *MIC2*.

<u>Supplemental figure legends</u>

<u>Suppl. Figure 1</u>: Comparison of parasite motility in fluorescent fibrin *vs*. Matrigel. (A) Comparison

of the maximum and mean speeds of parasites in 2.25 mg/ml fibrin *vs.* Matrigel. Horizontal bars indicate

mean (± SD) from 3 independent biological replicates (Matrigel n=853, Fibrin n=913). $p > 0.05$ for both

maximum and mean speed, Student's two-tailed t-test. **(B)** Comparison of the proportion of parasites

moving in 2.25 mg/ml fibrin *vs.* Matrigel. Horizontal bars indicate mean (± SD) from 3 independent

biological replicates (Matrigel n=853, Fibrin n=913). $p > 0.05$, Student's two-tailed t-test.

<u>Suppl. Figure 2</u>: Rheological properties of the fibrin matrix. The viscoelastic properties of the fibrin

matrix were measured by laser trapping of 0.91 μm styrene beads. **(A)** The laser trap position at 1 Hz (top

panel) and the resulting force as measured by the laser trap (bottom) show a primarily in-phase force

response. **(B)** The laser trap position and output force at 1-20Hz were used to calculate the elastic and

viscous properties of the matrix; these frequencies were chosen to encompass the approximate speed of a

moving parasite. Bars indicate mean (± SD).

<u>Suppl. Figure 3</u>: All displacement vectors point in towards a moving parasite. Matrix displacement

vectors from all subvolumes (49 (*x)* x 49 (*y*) X 7 (*z*)) in the image volume were calculated by FIDVC and

projected onto the *x-y, x-z* and *y-z* planes, as indicated (axis labels are in μm). Length of arrows indicate

displacement magnitude and are multiplied 10-fold for display. The data correspond to the parasite shown

in the second panel of Figure 4C and frame 2 of video 7. The moving parasite is outlined on each z slice

with red circles, and the position of the constriction is indicated by either an empty arrowhead (*x-y*) or a

black dot (*y-z*).

<u>Suppl. Figure 4</u>: Pattern of pulling, holding and release of the matrix during individual constriction

events. The number of consecutive frames in which: the matrix displaced in towards the parasite (pull);

no further matrix displacement was observed (hold); and the matrix moved away from the parasite

towards its original position (release) are indicated. n=9 parasites, 25 constrictions.

<u>Suppl. Figure 5</u>: Additional examples showing that matrix displacement is directed primarily in

towards the constriction in moving parasites. Empty red arrowheads indicate position of the

constriction and white arrows the direction of parasite travel. Length of black arrows indicating

displacement magnitude are multiplied 15-fold for display. Scale bars = 5 µm.

Suppl. Figure 6: **Fluorescent antibody against TgSAG1 is depleted from the parasite surface**

anterior to the constriction and shed into the Matrigel matrix. The top panels show the anti-TgSAG1

fluorescent signal; red empty arrowhead marks the location of the constriction. Middle panels show the

Hoechst 33342-stained nucleus, which stays in focus throughout the time series. Bottom panels show the

merged images (red = anti-TgSAG1, blue = Hoechst 33342). Scale bar = 5 µm; timestamps in seconds.

See Video 10 for the entire time series. All moving parasites examined (n=9) deposited trails containing

anti-SAG1 in the Matrigel.

Suppl. Figure 7: **The small number of TgMyoA knockout parasites that move more than one body**

length produce no detectable force on the fibrin matrix. (A) Sequential fluorescence images in a single

z-plane of a moving TgMyoA knockout parasite stained with Hoechst 33342 (to label the parasite

nucleus), **(B)** the corresponding force maps, and **(C)** the zoomed images showing the force maps from

panel B (boxed region), with background subtraction, overlaid on the parasite images from panel A. A red

crosshair placed at a fixed position in the images in panel (C) illustrates movement of the parasite. The

highlighted parasite continued to move for another 24 sec; no signal above background was seen on the

force maps at any time along its trajectory. Length of arrows indicating displacement magnitude are

multiplied 24-fold in (B) and 15-fold in (C) for display. Scale bar = 5 µm, timestamps in seconds.

Suppl. Figure 8: **TgMIC2 knockout parasites show no evidence of constrictions, produce no**

detectable force on the fibrin matrix and move less directionally. (A) Sequential fluorescence images

in a single *z*-plane of a moving TgMIC2 knockout parasite expressing YFP, **(B)** the corresponding force

maps, and **(C)** the zoomed images showing the force maps from panel B (boxed region), with background

subtraction, overlaid on the parasite images from panel A. A red crosshair placed at a fixed position in the

images in panel (C) illustrates movement of the parasite. The highlighted parasite continued to move for

another 18 sec; no signal above background was seen on the force maps at any time along its trajectory.

Length of arrows indicating displacement magnitude are multiplied 24-fold in (C) and 15-fold in (D) for

display. Scale bars = 10 µm (panel A), 5 µm (panel C); timestamps in seconds. **(D)** Ratio of nuclear

diameter perpendicular (width) *vs.* parallel (length) to the long axis of the parasite in moving wild-type

(WT; n=9) and TgMIC2 knockout (n=10) parasites, over time. For the wild-type parasites, triangles

indicate time points when the nucleus was passing through a constriction and circles when the nucleus

was not passing through a constriction. **(E)** Ratio of parasite displacement to trajectory length for wild-

type and TgMIC2 knockout parasites (three biological replicates, each consisting of three technical

replicates). The horizontal bars indicate means (+/- SD), which were compared by Student's two-sided t-

test. Note that in panels D and E, "WT" refers to the TgMIC2 LoxP parasites [29] before treatment with
rapamycin to excise *MIC2*.

Suppl. Figure 9: Parasites move faster after the constriction has past the midway point on the
parasite's longitudinal axis. The number of time points when the constriction was present either anterior
or posterior to the midway point along the parasite's longitudinal axis were determined. Only parasites
that progressed through one complete anterior-to-posterior constriction and neither stopped partway nor
initiated a second constriction while the original constriction was present were analyzed (n=20 parasites,
24 constrictions). It takes a constriction on average longer (*i.e.*, more timepoints) to progress from the
anterior tip of the parasite to the midway point along the parasite surface than it does from the midway
point to the posterior end. Bars indicate mean ± standard deviation; the data were compared by Student's
two-tailed t-test.

653
654   **Videos**
655
Video 1: Microsphere (green) being displaced towards a parasite (red) moving in Matrigel within in a
single *z* plane. Playback is in real time; scale bar = 5 µm. Single frames from this video are shown in
Figure 1B.
Video 2: Maximum intensity projection in *z* of microspheres (green) being displaced due to parasite (red)
movement within the Matrigel. Playback is in real time; scale bar = 5 µm.
Video 3: Brightfield imaging of a parasite undergoing a single constriction. Playback is in real time; scale
bar = 5 µm. Single frames from this video are shown in Figure 1C.
Video 4: Confocal fluorescence imaging of a 2.25 mg/ml fluorescent fibrin gel. 51 *x-y* slices captured
0.25 µm apart in *z* were reassembled into the volumetric view shown. Dimensions of the imaged volume
are shown at bottom left. See Figure 2A for maximum intensity projection of these data.
Video 5: Deformation of fluorescent fibrin (green) by a parasite (red; bottom) as it moves up through the
image stack. Playback is in real time; scale bar = 10 µm. Single frames from this video are shown in
Figure 2C.
Video 6: 3D displacement map surrounding the moving parasite shown in the second panel of Figure 4C
and frame 2 of Video 7. Length of arrows indicating displacement magnitude are multiplied 10-fold for
display. The parasite is outlined on each z plane with red circles. Single frames from this video displaying
projections onto the *x-y, x-z* and *y-z* planes are shown in Suppl. Figure 3.

Video 7: *x-y* displacement map overlaid on the moving parasite shown in Figure 4A-C. Playback is in real

time. Arrow size corresponds to relative displacement magnitude and arrow color to displacement

direction as described in Figure 4.

Video 8: *x-y* displacement map overlaid on the moving parasite shown in Figure 4D. Playback is in real

time. Arrow size corresponds to relative displacement magnitude and arrow color to displacement

direction as described in Figure 4.

Video 9: Brightfield imaging of a parasite undergoing two constrictions at the same time. Playback is in

real time; scale bar = 5 μm. Single frames from this video are shown in Figure 5B.

Video 10: Parasite labeled with Alexa546-conjugated anti-TgSAG1 (red) and Hoechst 33342 (blue)

undergoing a constriction. Playback is in real time; scale bar = 5 μm. Note the helical trail of shed

fluorescent antibody behind the moving parasite. Single frames from this video are shown in Suppl.

Figure 6.

Video 11: Brightfield imaging of a moving TgMyoA knockout parasite. Playback is in real time; scale

bar = 5 μm. Single frames from this video are shown in Figure 6A.

Video 12: Brightfield imaging of a moving TgMIC2 knockout parasite. Playback is in real time; scale bar

= 5 μm. Single frames from this video are shown in Figure 6A.

Acknowledgements. This work was supported by U.S. Public Health Service grants: AI139201 and

AI137767 (G.E.W): GM141743 and S10OD026884 for the Lumicks C-Trap (D.M.W); and T32AI055402

and F31AI145214 (R.V.S.), as well as American Heart Association grant 19PRE34370071 (R.V.S.) We

thank Nicole Bouffard and Dr. Douglas Taatjes of the UVM Larner College of Medicine Microscopy

Imaging Center (RRID# SCR_018821) for helpful advice and assistance with the Nikon A1R-ER

confocal microscope, which is supported by U.S. Public Health Service grant 1S10OD025030-01 from

the National Center for Research Resources. We thank: Dr. Markus Meissner for sharing the TgMyoA

and TgMIC2 knockout parasites; Dr. David Sibley for anti-TgSAG1 antibody; Drs. Mark Rould, Alex

Dunn, Christina Hueschen, Li-av Segev Zarko, John Boothroyd, Leanna Owen, Markus Meissner, and

Ulrich Schwarz for helpful discussions; Anne Snyder, Drs. Robyn Kent, Christina Hueschen, and Chris

Huston for helpful feedback on the manuscript. Finally, we thank Dr. Christian Franck for making the

FIDVC code used in this work freely available through GitHub.

References

1.   World malaria report. Geneva: World Health Organization, 2021.

2.   Sow SO, Muhsen K, Nasrin D, Blackwelder WC, Wu Y, Farag TH, et al. The Burden of
Cryptosporidium Diarrheal Disease among Children < 24 Months of Age in Moderate/High Mortality

Regions of Sub-Saharan Africa and South Asia, Utilizing Data from the Global Enteric Multicenter
Study (GEMS). PLoS Negl Trop Dis. 2016;10(5):e0004729. Epub 2016/05/25. doi:
10.1371/journal.pntd.0004729. PubMed PMID: 27219054; PubMed Central PMCID:
PMCPMC4878811

3. Bigna JJ, Tochie JN, Tounouga DN, Bekolo AO, Ymele NS, Youda EL, et al. Global, regional, and
country seroprevalence of Toxoplasma gondii in pregnant women: a systematic review, modelling and
meta-analysis. Sci Rep. 2020;10(1):12102. Epub 2020/07/23. doi: 10.1038/s41598-020-69078-9.
PubMed PMID: 32694844; PubMed Central PMCID: PMCPMC7374101.

4. Flegr J, Prandota J, Sovičková M, Israili ZH. Toxoplasmosis--a global threat. Correlation of latent
toxoplasmosis with specific disease burden in a set of 88 countries. PloS one. 2014;9(3):e90203. Epub
2014/03/26. doi: 10.1371/journal.pone.0090203. PubMed PMID: 24662942; PubMed Central
PMCID: PMCPMC3963851.

5. Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of *Toxoplasma gondii*
seroprevalence and implications for pregnancy and congenital toxoplasmosis. International journal for
parasitology. 2009;39(12):1385-94. Epub 2009/05/13. doi: 10.1016/j.ijpara.2009.04.003. PubMed
PMID: 19433092.

6. Rothova A. Ocular involvement in toxoplasmosis. British Journal of Ophthalmology. 1993;77(6):371-
7. doi: doi: 10.1136/bjo.77.6.371.

7. Su E, Honda A, Latkany P. Ocular Disease due to *Toxoplasma gondii*. In: Weiss LM, Kim K, editors.
Toxoplasma gondii - The Model Apicomplexan: Perspectives and Methods. Second ed. London, UK:
Academic Press; 2014. p. 161-92.

8. Daher W, Plattner F, Carlier MF, Soldati-Favre D. Concerted action of two formins in gliding motility
and host cell invasion by *Toxoplasma gondii*. PLoS Pathog. 2010;6(10). PubMed PMID: 20949068.

9. Elsheikha HM, Marra CM, Zhu XQ. Epidemiology, Pathophysiology, Diagnosis, and Management of
Cerebral Toxoplasmosis. Clin Microbiol Rev. 2021;34(1). Epub 2020/11/27. doi: 10.1128/cmr.00115-
19. PubMed PMID: 33239310.

10. Barragan A, Brossier F, Sibley LD. Transepithelial migration of *Toxoplasma gondii* involves an
interaction of intercellular adhesion molecule 1 (ICAM-1) with the parasite adhesin MIC2. Cell
Microbiol. 2005;7(4):561-8. PubMed PMID: 15760456.

11. Harker KS, Ueno N, Lodoen MB. *Toxoplasma gondii* Dissemination: A Parasite's Journey through the
Infected Host. Parasite immunology. 2015;37:141-9. Epub 2014/11/20. doi: 10.1111/pim.12163.
PubMed PMID: 25408224.

12. Meissner M, Schluter D, Soldati D. Role of *Toxoplasma gondii* myosin A in powering parasite gliding
and host cell invasion. Science. 2002;298(5594):837-40. PubMed PMID: 12399593.

13. Leung JM, Rould MA, Konradt C, Hunter CA, Ward GE. Disruption of *TgPHIL1* alters specific
parameters of *Toxoplasma gondii* motility measured in a quantitative, three-dimensional live motility
assay. PloS one. 2014;9(1):e85763. Epub 2014/02/04. doi: 10.1371/journal.pone.0085763. PubMed
PMID: 24489670; PubMed Central PMCID: PMC3906025.

14. Frenal K, Dubremetz JF, Lebrun M, Soldati-Favre D. Gliding motility powers invasion and egress in
Apicomplexa. Nat Rev Microbiol. 2017;15(11):645-60. doi: 10.1038/nrmicro.2017.86. PubMed
PMID: 28867819.

15. Heintzelman MB. Gliding motility in apicomplexan parasites. Seminars in cell & developmental
biology. 2015. Epub 2015/10/03. doi: 10.1016/j.semcdb.2015.09.020. PubMed PMID: 26428297.

16. Jacot D, Tosetti N, Pires I, Stock J, Graindorge A, Hung YF, et al. An Apicomplexan Actin-Binding
Protein Serves as a Connector and Lipid Sensor to Coordinate Motility and Invasion. Cell Host
Microbe. 2016;20(6):731-43. doi: 10.1016/j.chom.2016.10.020. PubMed PMID: 27978434.

17. Sibley LD. Intracellular parasite invasion strategies. Science. 2004;304(5668):248-53. PubMed
PMID: 15073368.

18. Kappe S, Bruderer T, Gantt S, Fujioka H, Nussenzweig V, Ménard R. Conservation of a gliding
motility and cell invasion machinery in Apicomplexan parasites. J Cell Biol. 1999;147(5):937-44.
Epub 1999/12/01. doi: 10.1083/jcb.147.5.937. PubMed PMID: 10579715; PubMed Central PMCID:
PMCPMC2169348.

19. Andenmatten N, Egarter S, Jackson AJ, Jullien N, Herman JP, Meissner M. Conditional genome
engineering in *Toxoplasma gondii* uncovers alternative invasion mechanisms. Nature methods.
2012;10(2):125-7. Epub 2012/12/25. doi: 10.1038/nmeth.2301. PubMed PMID: 23263690; PubMed
Central PMCID: PMC3605914.

20. Egarter S, Andenmatten N, Jackson AJ, Whitelaw JA, Pall G, Black JA, et al. The *Toxoplasma* Acto-
MyoA motor complex is important but not essential for gliding motility and host cell invasion. PloS
one. 2014;9(3):e91819. Epub 2014/03/19. doi: 10.1371/journal.pone.0091819. PubMed PMID:
24632839; PubMed Central PMCID: PMC3954763.

21. Tosetti N, Dos Santos Pacheco N, Soldati-Favre D, Jacot D. Three F-actin assembly centers regulate
organelle inheritance, cell-cell communication and motility in Toxoplasma gondii. eLife. 2019;8.
Epub 2019/02/13. doi: 10.7554/eLife.42669. PubMed PMID: 30753127; PubMed Central PMCID:
PMCPMC6372287.

22. Dobrowolski JM, Sibley LD. *Toxoplasma* invasion of mammalian cells is powered by the actin
cytoskeleton of the parasite. Cell. 1996;84(6):933-9.

23. Frenal K, Polonais V, Marq JB, Stratmann R, Limenitakis J, Soldati-Favre D. Functional dissection
of the apicomplexan glideosome molecular architecture. Cell Host Microbe. 2010;8(4):343-57.
PubMed PMID: 20951968.

24. Gras S, Jimenez-Ruiz E, Klinger CM, Schneider K, Klingl A, Lemgruber L, et al. An endocytic-
secretory cycle participates in Toxoplasma gondii in motility. PLoS Biol. 2019;17(6):e3000060. Epub
2019/06/25. doi: 10.1371/journal.pbio.3000060. PubMed PMID: 31233488.

25. Meissner M, Ferguson DJ, Frischknecht F. Invasion factors of apicomplexan parasites: essential or
redundant? Current opinion in microbiology. 2013;16(4):438-44. Epub 2013/06/04. doi:
10.1016/j.mib.2013.05.002. PubMed PMID: 23727286.

26. Tardieux I, Baum J. Reassessing the mechanics of parasite motility and host-cell invasion. J Cell Biol.
2016;214(5):507-15. doi: 10.1083/jcb.201605100. PubMed PMID: 27573462; PubMed Central
PMCID: PMCPMC5004448.

27. Whitelaw JA, Latorre-Barragan F, Gras S, Pall GS, Leung JM, Heaslip A, et al. Surface attachment,
promoted by the actomyosin system of *Toxoplasma gondii* is important for efficient gliding motility
and invasion. BMC Biol. 2017;15(1):1. doi: 10.1186/s12915-016-0343-5. PubMed PMID: 28100223;
PubMed Central PMCID: PMCPMC5242020.

28. Rompikuntal PK, Kent RS, Foe IT, Deng B, Bogyo M, Ward GE. Blocking Palmitoylation of
Toxoplasma gondii Myosin Light Chain 1 Disrupts Glideosome Composition but Has Little Impact
on Parasite Motility. mSphere. 2021;6(3). Epub 2021/05/21. doi: 10.1128/mSphere.00823-20.
PubMed PMID: 34011689.

29. Gras S, Jackson A, Woods S, Pall G, Whitelaw J, Leung JM, et al. Parasites lacking the micronemal
protein MIC2 are deficient in surface attachment and host cell egress, but remain virulent *in vivo*
[version 1]. Wellcome Open Research. 2017;2:32. doi: 10.12688/wellcomeopenres.11594.1.

30. Owen LM, Adhikari AS, Patel M, Grimmer P, Leijnse N, Kim MC, et al. A cytoskeletal clutch
mediates cellular force transmission in a soft, three-dimensional extracellular matrix. Mol Biol Cell.
2017;28(14):1959-74. Epub 2017/06/09. doi: 10.1091/mbc.E17-02-0102. PubMed PMID: 28592635;
PubMed Central PMCID: PMCPMC5541846.

31. Bar-Kochba E, Toyjanova J, Andrews E, Kim K-S, Franck C. A Fast Iterative Digital Volume
Correlation Algorithm for Large Deformations. Experimental Mechanics2015. p. 261-74.

32. Huynh MH, Carruthers VB. *Toxoplasma* MIC2 is a major determinant of invasion and virulence. PLoS
Pathog. 2006;2(8):e84. PubMed PMID: 16933991.

33. Munter S, Sabass B, Selhuber-Unkel C, Kudryashev M, Hegge S, Engel U, et al. *Plasmodium*
sporozoite motility is modulated by the turnover of discrete adhesion sites. Cell Host Microbe.
2009;6(6):551-62. PubMed PMID: 20006843.

34. Hakansson S, Morisaki H, Heuser J, Sibley LD. Time-lapse video microscopy of gliding motility in
Toxoplasma gondii reveals a novel, biphasic mechanism of cell locomotion. Mol Biol Cell.
1999;10(11):3539-47.

35. Pavlou G, Touquet B, Vigetti L, Renesto P, Bougdour A, Debarre D, et al. Coupling Polar Adhesion
with Traction, Spring and Torque Forces Allows High Speed Helical Migration of the Protozoan
Parasite Toxoplasma. ACS Nano. 2020;14:7121-39. Epub 2020/05/21. doi:
10.1021/acsnano.0c01893. PubMed PMID: 32432851.

36. Munera Lopez J, Tengganu IF, Liu J, Murray JM, Arias Padilla LF, Zhang Y, et al. An apical protein,
Pcr2, is required for persistent movement by the human parasite Toxoplasma gondii. PLoS Pathog.
2022;18(8):e1010776. Epub 2022/08/23. doi: 10.1371/journal.ppat.1010776. PubMed PMID:
35994509; PubMed Central PMCID: PMCPMC9436145.

37. Freyvogel TA. Shape, movement in situ and locomotion of plasmodial ookinetes. Acta Trop.
1966;23(3):201-22. Epub 1966/01/01. PubMed PMID: 4380662.

38. Vlachou D, Zimmermann T, Cantera R, Janse CJ, Waters AP, Kafatos FC. Real-time, in vivo analysis
of malaria ookinete locomotion and mosquito midgut invasion. Cell Microbiol. 2004;6(7):671-85.
Epub 2004/06/10. doi: 10.1111/j.1462-5822.2004.00394.x. PubMed PMID: 15186403.

39. Zieler H, Dvorak JA. Invasion in vitro of mosquito midgut cells by the malaria parasite proceeds by a
conserved mechanism and results in death of the invaded midgut cells. Proceedings of the National
Academy of Sciences of the United States of America. 2000;97(21):11516-21. Epub 2000/10/12. doi:
10.1073/pnas.97.21.11516. PubMed PMID: 11027351; PubMed Central PMCID: PMCPMC17232.

40. Alexander DL, Mital J, Ward GE, Bradley P, Boothroyd JC. Identification of the moving junction
complex of *Toxoplasma gondii*: a collaboration between distinct secretory organelles. PLoS Pathog.
2005;1(2):e17. doi: 10.1371/journal.ppat.0010017. PubMed PMID: 16244709.

41. Besteiro S, Dubremetz JF, Lebrun M. The moving junction of apicomplexan parasites: a key structure
for invasion. Cell Microbiol. 2011;13(6):797-805. PubMed PMID: 21535344.

42. Dubremetz JF. Host cell invasion by *Toxoplasma gondii* [see comments]. Trends Microbiol.
1998;6(1):27-30.

43. Lebrun M, Michelin A, El Hajj H, Poncet J, Bradley PJ, Vial H, et al. The rhoptry neck protein RON4
re-localizes at the moving junction during *Toxoplasma gondii* invasion. Cell Microbiol.
2005;7(12):1823-33. PubMed PMID: 16309467.

44. Porchet-Hennere E, Torpier, G. [Relationships between *Toxoplasma* and its host cell]. Protistologica.
1983;19:357-70.

45. Aikawa M, Miller LH, Johnson J, Rabbege J. Erythrocyte entry by malarial parasites. A moving
junction between erythrocyte and parasite. J Cell Biol. 1978;77(1):72-82.

46. Bargieri DY, Andenmatten N, Lagal V, Thiberge S, Whitelaw JA, Tardieux I, et al. Apical membrane
antigen 1 mediates apicomplexan parasite attachment but is dispensable for host cell invasion. Nature
communications. 2013;4:2552. Epub 2013/10/11. doi: 10.1038/ncomms3552. PubMed PMID:
24108241.

47. Mital J, Meissner M, Soldati D, Ward GE. Conditional expression of *Toxoplasma gondii* apical
membrane antigen-1 (TgAMA1) demonstrates that TgAMA1 plays a critical role in host cell invasion.
Mol Biol Cell. 2005;16(9):4341-9. doi: 10.1091/mbc.E05-04-0281. PubMed PMID: 16000372.

48. Pavlou G, Biesaga M, Touquet B, Lagal V, Balland M, Dufour A, et al. Toxoplasma Parasite Twisting
Motion Mechanically Induces Host Cell Membrane Fission to Complete Invasion within a Protective
Vacuole. Cell Host Microbe. 2018;24(1):81-96.e5. Epub 2018/07/17. doi:
10.1016/j.chom.2018.06.003. PubMed PMID: 30008293.

49. Hueschen CL, Zarko L-aS, Chen J-H, LeGros MA, Larabell CA, Boothroyd JC, et al. Emergent Actin
Flows Explain Diverse Parasite Gliding Modes. bioRxiv. 2022:2022.06.08.495399. doi:
10.1101/2022.06.08.495399.

50. Dubremetz JF, Rodriguez C, Ferreira E. *Toxoplasma gondii*: redistribution of monoclonal antibodies
on tachyzoites during host cell invasion. Exp Parasitol. 1985;59(1):24-32.

51. Herm-Gotz A, Weiss S, Stratmann R, Fujita-Becker S, Ruff C, Meyhofer E, et al. *Toxoplasma gondii*
myosin A and its light chain: a fast, single-headed, plus-end-directed motor. EMBO J.
2002;21(9):2149-58.

52. Mordue DG, Desai N, Dustin M, Sibley LD. Invasion by *Toxoplasma gondii* Establishes a Moving
Junction That Selectively Excludes Host Cell Plasma Membrane Proteins on the Basis of Their
Membrane Anchoring. J Exp Med. 1999;190(12):1783-92.

53. Quadt KA, Streichfuss M, Moreau CA, Spatz JP, Frischknecht F. Coupling of Retrograde Flow to
Force Production During Malaria Parasite Migration. ACS Nano. 2016;10(2):2091-102. Epub
2016/01/23. doi: 10.1021/acsnano.5b06417. PubMed PMID: 26792112.

54. Stadler RV, Nelson SR, Warshaw DM, Ward GE. A circular zone of attachment to the extracellular
matrix acts as a guidance system for Toxoplasma gondii 3D motility. bioRxiv.
2022:2022.06.08.495400. doi: 10.1101/2022.06.08.495400.

55.   Frenal K, Soldati-Favre D. Plasticity and Redundancy in Proteins Important for *Toxoplasma* Invasion.
PLoS Pathog. 2015;11(8):e1005069. Epub 2015/08/14. doi: 10.1371/journal.ppat.1005069. PubMed
PMID: 26270966; PubMed Central PMCID: PMC4536160.

56.   Lamarque MH, Roques M, Kong-Hap M, Tonkin ML, Rugarabamu G, Marq JB, et al. Plasticity and
redundancy among AMA-RON pairs ensure host cell entry of *Toxoplasma* parasites. Nature
communications. 2014;5:4098. Epub 2014/06/18. doi: 10.1038/ncomms5098. PubMed PMID:
24934579.

57.   Lammermann T, Bader BL, Monkley SJ, Worbs T, Wedlich-Soldner R, Hirsch K, et al. Rapid
leukocyte migration by integrin-independent flowing and squeezing. Nature. 2008;453(7191):51-5.
doi: 10.1038/nature06887. PubMed PMID: 18451854.

58.   Berg HC, Turner L. Movement of microorganisms in viscous environments. Nature.
1979;278(5702):349-51. Epub 1979/03/22. doi: 10.1038/278349a0. PubMed PMID: 370610.

59.   Ferrero RL, Lee A. Motility of Campylobacter jejuni in a viscous environment: comparison with
conventional rod-shaped bacteria. J Gen Microbiol. 1988;134(1):53-9. Epub 1988/01/01. doi:
10.1099/00221287-134-1-53. PubMed PMID: 3053972.

60.   Hazell SL, Lee A, Brady L, Hennessy W. Campylobacter pyloridis and gastritis: association with
intercellular spaces and adaptation to an environment of mucus as important factors in colonization of
the gastric epithelium. J Infect Dis. 1986;153(4):658-63. Epub 1986/04/01. doi:
10.1093/infdis/153.4.658. PubMed PMID: 3950447.

61.   Kaiser GE, Doetsch RN. Letter: Enhanced translational motion of Leptospira in viscous environments.
Nature. 1975;255(5510):656-7. Epub 1975/06/19. doi: 10.1038/255656a0. PubMed PMID: 1134561.

62.   Kimsey RB, Spielman A. Motility of Lyme disease spirochetes in fluids as viscous as the extracellular
matrix. J Infect Dis. 1990;162(5):1205-8. Epub 1990/11/01. doi: 10.1093/infdis/162.5.1205. PubMed
PMID: 2230247.

63.   Roos DS, Donald RG, Morrissette NS, Moulton AL. Molecular tools for genetic dissection of the
protozoan parasite *Toxoplasma gondii*. Methods in Cell Biology. 451994. p. 27-63.

64.   Foe IT, Child MA, Majmudar JD, Krishnamurthy S, van der Linden WA, Ward GE, et al. Global
Analysis of Palmitoylated Proteins in *Toxoplasma gondii*. Cell Host Microbe. 2015;18(4):501-11.
Epub 2015/10/16. doi: 10.1016/j.chom.2015.09.006. PubMed PMID: 26468752.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7