Title: A conserved membrane protein negatively regulates Mce1 complexes in mycobacteria

Authors: Yushu Chen[a], Shu-Sin Chng[a,b,*]

Affiliations:

[a] Department of Chemistry, National University of Singapore, Singapore 117543.

[b] Singapore Center for Environmental Life Sciences Engineering, National University of Singapore

(SCELSE-NUS), Singapore 117456.

* To whom correspondence should be addressed. Email: chmchngs@nus.edu.sg

Author contributions: Y.C. and S.-S.C. designed research; Y.C. performed research; Y.C. and S.-S.C.

analyzed data; Y.C. and S.-S.C. wrote the paper.

Competing Interest Statement: The authors declare that no competing interests exist.

Classification: Biological Sciences (Biochemistry)

Keywords: mammalian cell entry, ABC transporter, fatty acid uptake, negative regulator

This PDF file includes

Main Text

Figures 1 to 5

Abstract

Tuberculosis (TB) remains one of the deadliest infectious diseases, posing a serious threat to global health. *Mycobacterium tuberculosis*, the causative agent of TB, is an intracellular pathogen that relies on various mechanisms to survive and persist within the host environment. Among the many virulence factors that contribute to pathogenesis, mycobacteria encode Mce systems, putative transporters that are important for lipid uptake. The molecular basis for Mce function(s) is poorly understood. To gain insights into the composition and architecture of mycobacterial Mce systems, we characterized the putative Mce1 complex involved in fatty acid transport. Using affinity purification, we show that the Mce1 system in *Mycobacterium smegmatis* is an ATP-binding cassette transporter complex comprising YrbE1A/B as heterodimeric transmembrane domains (TMDs) and MceG as nucleotide-binding domains (NBDs), associated with possible heterohexameric assemblies of MCE-domain substrate-binding proteins. We demonstrate that Msmeg_6540 and Mce1A are functionally redundant homologs, likely forming distinct Mce1 complexes that contribute to fatty acid uptake. Finally, we establish that the membrane protein Msmeg_0959, herein renamed to Mce1N, negatively regulates Mce1 function by blocking MceG access to YrbE1B. This regulatory mechanism is conserved in *M. tuberculosis* Mce1N (Rv0513). Our work offers molecular understanding of Mce complexes, shedding light on lipid metabolism and its regulation in mycobacteria.

Introduction

Tuberculosis (TB) has been one of the top killers among single infectious diseases in humans, alone claiming 1.3 million lives out of 10 million new cases in 2020 (1). Apart from the active cases, it is estimated that one quarter of the world population are carriers of latent TB, which may progress into active infections (2). TB is primarily caused by *Mycobacterium tuberculosis*, an organism known for its ability to adapt to the host environment and modulate the host immune system (3). Its success as a pathogen can be owed to diverse virulence factors encoded in the genome of *M. tuberculosis*. In particular, many of these virulence factors promote host adaptation by contributing to the acquisition and catabolism of important nutrients including metals, sugars and lipids (4).

Mycobacterial *mce* operons have been recognized as a group of virulence factors important for lipid uptake (5). These operons are widespread throughout the *Mycobacterium* genus, with different numbers present in different species (6). *Mycobacterium smegmatis,* an environmental microorganism, contains six *mce* operons, while *Mycobacterium leprae*, the causative agent of leprosy, has only one such operon (*mce1*) in its highly degenerate genome. *M. tuberculosis* contains four *mce* operons, individually important determinants of virulence and collectively critical for pathogenesis (7-10). For example, the *mce1* and *mce4* operons are implicated in fatty acid and cholesterol uptake, respectively (9, 11, 12). In general, an *mce* operon consists of eight genes, two *yrbE* (e.g., *yrbE1A/B*) and six *mce* genes (e.g., *mce1A–F),* with different numbers of accessory genes (6) that may be required for function (12, 13) (e.g., *mam1A–D* after the *mce1* operon). Furthermore, a gene encoding a transcriptional regulator (e.g., *mce1R*) can be found upstream of some *mce* operons (6, 14, 15) (*SI Appendix*, Fig. S1). Our understanding of how these operons facilitate lipid uptake is limited.

It is believed that *mce* operons encode ATP-binding cassette (ABC) transporters that are analogous to the MlaFEDB complex in Gram-negative bacteria (6, 16), which is important for retrograde transport of phospholipids (17, 18). In MlaFEDB, MlaF, MlaE and MlaD represent the nucleotide-binding domains (NBDs), transmembrane domains (TMDs), and substrate-binding proteins (SBPs), respectively (19, 20). In Mce systems, YrbEA/Bs and MceA–Fs would form the corresponding TMDs and SBPs based on homology; here, MceG, a protein not encoded within *mce* operons, is thought to constitute the NBDs that energize all these putative transporters (21, 22). MlaD and MceA–F all contain a single mammalian cell

70 entry (MCE) domain. Interestingly, MlaD and other MCE-domain proteins, such as PqiB and YebT/LetB, have been shown to form homohexamers that provide a hydrophobic tunnel each, presumably for lipid substrate transport (19, 20, 23, 24). The fact that *mce* operons encode six Mce proteins MceA–F hints at the possible formation of heterohexameric MCE assemblies within each putative transporter. Despite similarities to MlaFEDB, however, there has been a lack of biochemical studies on mycobacterial Mce systems, and their exact compositions and overall architectures remain unknown.

76 Here, we biochemically characterized the putative Mce1 complex in *M. smegmatis.* We demonstrate that Mce1 is an ABC transporter comprising a core complex of YrbE1A/B-MceG, associated with a likely heterohexamer of Mce1A–F. We reveal that Msmeg_6540 is a functional homolog that can possibly replace Mce1A in the complex, and both proteins contribute to the fatty acid uptake function of Mce1. Furthermore, we identify Msmeg_0959, herein renamed to Mce1N, as a negative regulator of the Mce1 system; it prevents MceG from associating with the complex via competitive binding to YrbE1B, thereby modulating transport activity. Our work provides critical insights into the architecture, function and regulation of Mce systems in mycobacteria.

85 **Results**

86 **Key proteins encoded in the *mce1* operon co-purify as a complex**

87 To probe the composition of the putative Mce1 complex, we expressed His-tagged YrbE1A or YrbE1B in *M. smegmatis* using an integrative plasmid and conducted affinity purification following cell lysis and detergent extraction from total membranes. This was done in a strain that presumably up-regulates the *mce1* operon due to the lack of Mce1R, a known negative regulator (14). We found that His-tagged YrbE1B, but not YrbE1A, pulled down two distinct protein bands at ~45 and ~19 kDa (Fig. 1). Tandem mass spectrometry (MS/MS) sequencing of the excised bands indicated that they were MceG (39.0 kDa) and a protein of unknown function, Msmeg_0959 (19.1 kDa), respectively. We did not observe additional bands corresponding to other expected members of the Mce1 complex, possibly due to low cellular levels, or instability of the complex *ex vivo*. To detect potential low abundance proteins specifically co-purified with His-tagged YrbE1B, we directly subjected entire gel lanes divided into two sections to MS/MS protein identification. Remarkably, this facilitated the detection of all other putative Mce1 components, including

YrbE1A and Mce1A–F, similar to a recent report identifying YrbE4A/B, Mce4A–F and MceG as components

of the Mce4 complex (13). Interestingly, a seventh MCE-domain protein Msmeg_6540 (6) was also co-

purified with His_6-YrbE1B, while none of the putative Mce-associated membrane proteins (Mam1A–D) was

identified. Our data suggest that the Mce1 complex likely comprises the core ABC transporter YrbE1A/B-

MceG associated with a presumed heterohexamer of MCE-domain proteins.

Msmeg_6540 and Mce1A are two redundant components of the Mce1 system

Msmeg_6540 shares 80% primary sequence identity with Mce1A, compared to less than 20% with

other MCE-domain proteins in the Mce1 complex and ~25% with MceA proteins in other putative *M.*

smegmatis Mce complexes (*SI Appendix*, Table S4). Another Mce1A paralog Msmeg_5818 (6), which is

only 60% identical, was not co-purified with His-tagged YrbE1B (Fig. 1). This suggests that Msmeg_6540

could be a functional homolog of Mce1A. To study the role of Msmeg_6540, we tested the effects of

deleting *msmeg_6540* on fatty acid uptake, a key function of the Mce1 complex (11, 12). Deletion of

yrbE1A/B or *mceG* resulted in reduced [^{14}C]-palmitic acid uptake (Fig. 2A), and retarded cell growth

specifically when palmitate was the sole carbon source (Fig. 2B). Interestingly, we found that [^{14}C]-palmitic

acid uptake was less efficient in cells lacking either *mce1A* or *msmeg_6540*, and in fact further reduced in

the Δ*mce1A*Δ*6540* double mutant, comparable to levels in Δ*yrbE1A/B* or Δ*mceG* cells (Fig. 2A). In addition,

while both single deletion mutants exhibited slower growth when using palmitate as the sole carbon source,

the Δ*mce1A*Δ*6540* strain grew at an even lower rate, again similar to Δ*yrbE1A/B* or Δ*mceG* strains (Fig.

2B). Our results therefore reveal that Mce1A and Msmeg_6540 are functionally redundant. Curiously,

removing Msmeg_6540 has a stronger impact on fatty acid uptake than removing Mce1A, suggesting

Mce1 complexes containing Msmeg_6540 in place of Mce1A could play a more important role in *M.*

smegmatis.

YrbE1A/B-MceG constitute the core ABC transporter

Despite extensive efforts, we were not successful in over-expressing the *mce1* operon along with

mceG in *E. coli* and purifying the full Mce1 complex. Instead, we were able to isolate the core ABC

transporter when we co-expressed YrbE1A, His-tagged YrbE1B and MceG. This complex eluted as a

single peak on size exclusion chromatography (Fig. 3A); we showed that the purified complex exhibits ATP

hydrolytic activity (Fig. 3B), implying proper assembly of YrbE1A/B-MceG. Interestingly, we found that His-

tagged YrbE1B alone, but not YrbE1A, could pull down MceG when co-expressed in *E. coli*, indicating that

the TMD-NBD interactions are mainly mediated through YrbE1B (Fig. 3C). Consistent with this idea, an

AlphaFold2 structural model of the YrbE1A/B-MceG complex (25-27), which resembles the homologous

MlaFE structure from Gram-negative bacteria (RMSD=2.934 Å; *SI Appendix*, Fig. S2) (28), presented a

smaller buried area between MceG and YrbE1A (699 Å^2) compared to YrbE1B (837 Å^2) (Fig. 3D and *SI*

Appendix, Table S5). We conclude that the TMDs and NBD proteins of the Mce1 system interact

asymmetrically to form a stable functional ABC transporter.

Msmeg_0959 is a negative regulator of Mce1

In addition to MceG, Msmeg_0959 was another protein abundantly co-purified with His-tagged

YrbE1B expressed in *M. smegmatis*. Msmeg_0959 is a putative conserved membrane protein with

unknown function; it contains two predicted transmembrane helices (TM1: W39-G59; TM2: L65-T90) and

a C-terminal cytoplasmic domain (S91-P178) (*SI Appendix*, Fig. S3). We initially hypothesized that it might

function to stabilize the Mce1 complex, analogous to the role of the cytoplasmic protein MlaB in the Gram-

negative MlaFEDB complex (19). To test this idea, we attempted to isolate a stable complex containing

YrbE1A/B, MceG and Msmeg_0959. To our surprise, when these four proteins were co-expressed and

purified from *E. coli*, Msmeg_0959, but noticeably not MceG, was co-purified with His-tagged YrbE1B (Fig.

4A). This observation suggests that Msmeg_0959 may destabilize the Mce1 complex instead. In support

of this notion, when Msmeg_0959 was overexpressed in *M. smegmatis*, much less MceG was co-purified

with His-tagged YrbE1B (Fig. 4B). We conclude that Msmeg_0959 may disrupt YrbE1A/B-MceG

interactions.

Since TMD-NBD interaction within the Mce1 complex is largely mediated through YrbE1B, we

presumed that Msmeg_0959 directly interacts with YrbE1B. We generated an AlphaFold2 structural model

comprising YrbE1B and Msmeg_0959 (25-27), predicting possible interactions between the

transmembrane domain of Msmeg_0959 and that of YrbE1B opposite its dimerization surface with YrbE1A

(Fig. 4C). The structural model revealed that the cytoplasmic domain of Msmeg_0959 may contact a short

helix of YrbE1B sitting at the membrane-water boundary. This helix may be a determinant of the interaction

between YrbE1B and Msmeg_0959 since the amino acid sequence of this region is very different from that

in YrbE1A, which did not pull down Msmeg_0959 in *M. smegmatis* (*SI Appendix*, Fig. S4). Mutating part

of this helix (RLVAEIGMGT) to a flexible loop (GGSSG) did not affect the stability of the YrbE1A/B-MceG

complex *in vitro*. However, this mutation prevented co-purification of Msmeg_0959 with His-tagged YrbE1B

from *E. coli*, at the same time alleviating the destabilizing effect of Msmeg_0959 on YrbE1A/B-MceG

interactions (Fig. 4A). Notably, overlay of the models of YrbE1B-Msmeg_0959 and YrbE1A/B-MceG

highlighted substantial steric clashes between MceG and the cytoplasmic domain of Msmeg_0959 (Fig.

4C), indicating that their respective interactions with YrbE1B are likely competitive and mutually exclusive.

Taken together, our results establish that Msmeg_0959 specifically binds YrbE1B and prevents association

of the NBD (MceG) to this ABC transporter.

Our biochemical experiments suggest that Msmeg_0959 may be a negative regulator of the Mce1

complex. Consistent with this idea, Msmeg_0959 overexpression inhibited [^{14}C]-palmitic acid uptake in WT

cells, but not residual uptake observed in Δ*yrbE1A/B* cells (Fig. 4D). Furthermore, we found that

overexpression of Rv0513, the *M. tuberculosis* homolog of Msmeg_0959, in *M. smegmatis* similarly

reduced the amount of MceG associated with YrbE1B (Fig. 4B), and slowed down [^{14}C]-palmitic acid

uptake (Fig. 4D). In contrast, overexpression of Ml2418c, the truncated *M. leprae* homolog lacking a large

portion of the C-terminal cytoplasmic domain (*SI Appendix*, Fig. S3), had no such effects (Fig. 4B and D).

We therefore conclude that Msmeg_0959/Rv0513, herein renamed to Mce1N (Mce1 negative regulator),

inhibits the fatty acid uptake function of the Mce1 transporter(s) via a conserved mechanism that involves

prevention of YrbE1B-MceG interaction.

Discussion

In this study, we set out to characterize the Mce1 complex in *M. smegmatis*. We have shown that the

complex likely comprises the core inner membrane ABC transporter YrbE1A/B-MceG, associated with a

putative heterohexamer composed of Mce1A–F. Interestingly, we have found that Msmeg_6540, a

homolog of Mce1A, can also be co-purified as part of the Mce1 complex, and have demonstrated that it is

functionally redundant with Mce1A in fatty acid uptake. Furthermore, we have discovered a negative

regulator of the Mce1 complex, Mce1N, which physically hinders the dominant TMD-NBD interaction (i.e.,

YrbE1B-MceG) in the ABC transporter. Our work provides novel insights into the architecture and

regulation of mycobacterial Mce complexes.

Based on analogy to other MCE-domain proteins, such as MlaD, PqiB (20) and YebT/LetB (23, 24)

involved in lipid transport in Gram-negative bacteria (*SI Appendix*, Fig. S5*A*), Mce1A–F likely form a

hexamer. Each of these Mce proteins contains a single MCE domain (like MlaD) and an extended helical

domain (like PqiB), followed by the presence of an additional C-terminal domain of varied sizes. According

to AlphaFold2 prediction (25, 26, 29), Mce1A–F can indeed form a heterohexamer (*SI Appendix*, Fig. S5*B*).

In contrast, a homohexameric model of Mce1A alone could not be generated. In the AlphaFold2 model,

the heterohexamer adopts an architecture that comprises a hexameric MCE ring connected to a tube

formed by a bundle of the helical domains; a continuous hydrophobic tunnel runs through the entire

assembly that might extend across the periplasm (Fig. 5 and *SI Appendix*, Fig. S5*B* and *C*), consistent

with its function in fatty acid uptake. In this predicted model, the C-terminal domains of the six Mce proteins

organize to form a possible substrate entry point at the tip (*SI Appendix*, Fig. S5*B*). Interestingly, *M.*

smegmatis possesses Msmeg_6540 as a functional homolog of Mce1A, hinting towards the presence of

two variants of Mce1 complexes (6), containing Mce1B–F and either Mce1A or Msmeg_6540 (Fig. 5).

Differences between the primary sequences of Mce1A and Msmeg_6540 lie mainly in the extreme C-

terminal domain, with differing residues mapped to both the exterior and interior of the 'substrate entry

point' in the AlphaFold2 models (*SI Appendix*, Fig. S5*D*). We propose that Mce1 complexes containing

either Mce1A or Msmeg_6540 may have distinct substrate specificities, perhaps advantageous for an

environmental microorganism like *M. smegmatis*.

Despite the fact that there are known ABC transporters with heterodimeric TMDs, it has not been

appreciated that such asymmetry can give rise to major differences in affinity between each TMD and NBD

pair—in the case of the Mce1 transporter, the NBD dimer may be predominantly recruited by YrbE1B. This

might be applicable to other ABC transporters with heterodimeric TMDs since TMD-NBD interfaces of

different areas were also observed within some of them (*SI Appendix*, Table S5). That TMD-NBD

association in the Mce1 transporter relies on the YrbE1B-MceG interaction offers a compelling reason why

the negative regulator Mce1N also selectively binds YrbE1B. Based on our modeling and biochemical data,

the cytoplasmic domain of Mce1N likely interacts with YrbE1B extensively at a short helix between the

latter's N-terminal amphipathic interface helix and TM1. This short helix is nominally a part of TM1 but is

presented at an angle due to a kink introduced by a flexible loop within TM1. Remarkably, this structural

feature may be exclusively conserved across YrbEBs, but not YrbEAs (*SI Appendix*, Fig. S6). This raises

an interesting possibility that beyond the most conserved Mce1 complex, Mce1N may also regulate

additional Mce systems by binding to the corresponding YrbEB, perhaps providing a way to redistribute

MceG among all Mce systems.

The discovery of a negative regulator that competes with the NBD for binding to the TMDs may be

unique to Mce complexes, as known regulatory mechanisms for ABC transporters typically involve

modulation of ATPase activity instead of removing the NBD from the complex (19, 30-34). How and when

the Mce1 complex is modulated would directly depend on the levels of Mce1N (Fig. 5), which is highly

conserved across the *Mycobacterium* genus (present in both fast- and slow-growing, pathogenic and non-

pathogenic species). We note that *mce1N* may be in an operon together with four upstream essential

genes involved in porphyrin biosynthesis (*hemACDB*). Porphyrin biosynthesis has been reported to

increase, suggesting possible up-regulation of Mce1N (Msmeg_0959), when *M. smegmatis* cells enter

dormancy (35). In this situation, negative regulation by Mce1N might be a strategy that cells use to inhibit

some Mce complexes to conserve ATP. Alternatively, Mce1N levels may correlate with the need for lipid

uptake. Outside the host, where fatty acid substrates may be scarce, it might be preferable for

mycobacterial cells to keep the Mce1 system in a routinely OFF state and avoid futile ATP hydrolysis.

During infection, however, cells may require a fully active Mce1 complex to utilize host fatty acids. We do

not yet know of any environment-dependent regulation of Mce1N. However, it is noteworthy that in *M.*

leprae, which possesses Mce1 as the only Mce system (6), Mce1N (Ml2418c) lacks the C-terminal

cytoplasmic domain and is likely non-functional. Consistently, we have shown that Ml2418c was unable to

inhibit the Mce1 complex when overexpressed in *M. smegmatis* (Fig. 4). Combined with the fact that

mce1R, the gene encoding the transcriptional repressor of the *mce1* operon (*SI Appendix*, Fig. S1), is

absent in *M. leprae*, the pathogen may have lost its ability to down-regulate the Mce1 system, which can

facilitate constitutive fatty acid uptake during its obligate parasitic life cycle.

We have uncovered a new regulatory mechanism for Mce1 (Fig. 5), and potentially other Mce systems,

which may have implications for how mycobacterial cells coordinate lipid uptake processes. Despite the

presence of a transcriptional regulator for some *mce* operons (*SI Appendix*, Fig. S1), all Mce transporters

apparently share the same ATPase MceG (22). This indicates that mycobacterial cells have the ability to

modulate the levels of each complex, yet adjust their activities simultaneously. In the same vein, we

speculate that Mce1N could also be a shared negative regulator to counter MceG binding/activation, and

primarily functions to deactivate all Mce complexes simultaneously. Whether Mce1N has preferences for

specific Mce systems remains to be examined. Nevertheless, it likely contributes significantly to the overall

regulation towards maintaining a balanced metabolic flux of different lipid substrates required for

mycobacterial survival (12, 36), which may involve other yet-to-be-identified uptake transporters beyond

Mce systems (37).

Materials and Methods

Affinity purification

Harvested *M. smegmatis* or *E. coli* cells were resuspended in cold TBS (Tris-buffered saline, 20 mM Tris-

HCl pH 8.0, 150 mM NaCl) containing lysozyme, phenylmethylsulfonyl fluoride, and DNase I, and lysed

using a French Press (GlenMills) at 20,000 psi. After removal of unbroken cells by centrifugation, the

supernatant was subjected to ultra-centrifugation at 95,000 x g for 45 min. Membrane proteins were

extracted from the pellet using TBS-A (TBS containing 5 mM $MgCl_2$ and 10% glycerol) with 1% n-dodecyl-

β-D-maltoside (DDM). The suspension was subjected to a second round of ultra-centrifugation at 89,000

x g for 45 min. The supernatant was loaded onto TALON metal affinity resin (Takara Bio) pre-equilibrated

with TBS. The mixture was allowed to drain by gravity. The resin was then washed ten times using high-

salt (300 mM NaCl) TBS-A with 8 mM imidazole and 0.05% DDM and subsequently eluted using TBS-A

with 200 mM imidazole and 0.05% DDM.

Palmitic acid uptake

The palmitic acid uptake assay was adapted from (12). Briefly, *M. smegmatis* cells were grown to stationary

phase in Sauton's medium with palmitate as the carbon source. The cells were pelleted and resuspended

in PBS (10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 2.7 mM KCl, 137 mM NaCl, pH 7.4) supplemented with 0.05%

Triton X-100 and 0.1% fatty acid-free bovine serum albumin (BSA) yielding a final OD_{600} of 0.8. After [^{14}C]-

palmitic acid (Perkin Elmer) was added, the cells were incubated at 37 ºC with shaking. Cells were harvested at different time points by centrifugation and washed once with ice-cold PBS containing Triton X-100 and BSA. The cells were finally resuspended in the same buffer and aliquoted into plastic scintillation vials for counting using a MicroBeta2 microplate counter (Perkin Elmer).

Acknowledgements

We thank Jia Fu Erh and Wei Shen Ho (National University of Singapore) for constructing plasmids and early affinity purification attempts. We are grateful to Beijing Genomics Institute (BGI) Research for providing DNA synthesis and assembly support. We thank Ross Tomaino (Taplin MS facility, Harvard Medical School) for help with protein identification. This work was supported sequentially by the Singapore Ministry of Education Academic Research Fund Tier 2 grants (MOE2019-T2-1-128 and MOE000116) (to S.-S.C.).

References

1. World Health Organization, Global Tuberculosis Report 2021, (2021).

2. R. M. Houben, P. J. Dodd, The global burden of latent tuberculosis infection: a re-estimation using mathematical modelling. *PLoS Med.* **13**, e1002152 (2016).

3. Q. Chai, Y. Zhang, C. H. Liu, *Mycobacterium tuberculosis*: an adaptable pathogen associated with multiple human diseases. *Front. Cell. Infect. Microbiol.* **8**, 158 (2018).

4. M. A. Forrellad *et al.*, Virulence factors of the *Mycobacterium tuberculosis* complex. *Virulence* **4**, 3-66 (2013).

5. L. I. Klepp, J. S. y Sabio, F. Bigi, Mycobacterial MCE proteins as transporters that control lipid homeostasis of the cell wall. *Tuberculosis (Edinb)* **132**, 102162 (2022).

6. N. Casali, L. W. Riley, A phylogenomic analysis of the *Actinomycetales mce* operons. *BMC Genomics* **8**, 60 (2007).

7. N. Shimono *et al.*, Hypervirulent mutant of *Mycobacterium tuberculosis* resulting from disruption of the *mce1* operon. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 15918-15923 (2003).

8. A. Gioffre *et al.*, Mutation in *mce* operons attenuates *Mycobacterium tuberculosis* virulence. *Microbes*

Infect. **7**, 325-334 (2005).

9. A. K. Pandey, C. M. Sassetti, Mycobacterial persistence requires the utilization of host cholesterol.

Proc. Natl. Acad. Sci. U. S. A. **105**, 4376-4380 (2008).

10. R. H. Senaratne *et al.*, *Mycobacterium tuberculosis* strains disrupted in *mce3* and *mce4* operons are

attenuated in mice. *J. Med. Microbiol.* **57**, 164-170 (2008).

11. M. A. Forrellad *et al.*, Role of the Mce1 transporter in the lipid homeostasis of *Mycobacterium*

tuberculosis. *Tuberculosis (Edinb)* **94**, 170-177 (2014).

12. E. V. Nazarova *et al.*, Rv3723/LucA coordinates fatty acid and cholesterol uptake in *Mycobacterium*

tuberculosis. *eLife* **6** (2017).

13. L. Rank, L. E. Herring, M. Braunstein, Evidence for the mycobacterial mce4 transporter being a

multiprotein complex. *J. Bacteriol.* **203**, e00685-00620 (2021).

14. N. Casali, A. M. White, L. W. Riley, Regulation of the *Mycobacterium tuberculosis mce1* operon. *J.*

Bacteriol. **188**, 441-449 (2006).

15. M. P. Santangelo *et al.*, Study of the role of Mce3R on the transcription of *mce* genes of *Mycobacterium*

tuberculosis. *BMC Microbiol.* **8**, 38 (2008).

16. J. C. Malinverni, T. J. Silhavy, An ABC transport system that maintains lipid asymmetry in the Gram-

negative outer membrane. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 8009-8014 (2009).

17. X. Tang *et al.*, Structural insights into outer membrane asymmetry maintenance in Gram-negative

bacteria by MlaFEDB. *Nat. Struct. Mol. Biol.* **28**, 81-91 (2021).

18. W.-Y. Low, S. Thong, S.-S. Chng, ATP disrupts lipid-binding equilibrium to drive retrograde transport

critical for bacterial outer membrane asymmetry. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

19. S. Thong *et al.*, Defining key roles for auxiliary proteins in an ABC transporter that maintains bacterial

outer membrane lipid asymmetry. *eLife* **5** (2016).

20. D. C. Ekiert *et al.*, Architectures of Lipid Transport Systems for the Bacterial Outer Membrane. *Cell*

169, 273-285 e217 (2017).

21. S. M. Joshi *et al.*, Characterization of mycobacterial virulence genes through genetic interaction

mapping. *Proc Natl Acad Sci U S A* **103**, 11760-11765 (2006).

22. J. Garcia-Fernández, K. Papavinasasundaram, B. Galán, C. M. Sassetti, J. L. García, Unravelling the

pleiotropic role of the MceG ATPase in *Mycobacterium smegmatis*. *Environ. Microbiol.* **19**, 2564-2576

(2017).

23. C. Liu, J. Ma, J. Wang, H. Wang, L. Zhang, Cryo-EM structure of a bacterial lipid transporter YebT. *J.*

Mol. Biol. **432**, 1008-1019 (2020).

24. G. L. Isom *et al.*, LetB structure reveals a tunnel for lipid transport across the bacterial envelope. *Cell*

181, 653-664 (2020).

25. R. Evans *et al*., Protein complex prediction with AlphaFold-Multimer. BioRxiv [Preprint] (2022).

https://www.biorxiv.org/content/10.1101/2021.10.04.463034v2 (accessed 6 June 2022).

26. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589

(2021).

27. M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nat. Methods* 10.1038/s41592-

022-01488-1 (2022).

28. C. Zhou *et al.*, Structural insight into phospholipid transport by the MlaFEBD complex from *P.*

aeruginosa. *J. Mol. Biol.* **433**, 166986 (2021).

29. M. A. Cianfrocco, M. Wong, C. Youn, R. Wagner, A. E. Leschziner, "COSMIC²: A science gateway for

cryo-electron microscopy structure determination" in *Proceedings of the Practice and Experience in*

Advanced Research Computing 2017 on Sustainability, Success and Impact (Association for

Computing Machinery, New York, NY, 2017), pp. 1-5.

30. R. R. Crawford, P. K. Potukuchi, E. G. Schuetz, J. D. Schuetz, Beyond competitive inhibition: regulation

of ABC transporters by kinases and protein-protein interactions as potential mechanisms of drug-drug

interactions. *Drug Metab. Dispos.* **46**, 567-580 (2018).

31. N. S. Kadaba, J. T. Kaiser, E. Johnson, A. Lee, D. C. Rees, The high-affinity *E. coli* methionine ABC

transporter: structure and allosteric regulation. *Science* **321**, 250-253 (2008).

32. S. Gerber, M. Comellas-Bigler, B. A. Goetz, K. P. Locher, Structural basis of trans-inhibition in a

molybdate/tungstate ABC transporter. *Science* **321**, 246-250.

33. S. Chen, M. L. Oldham, A. L. Davidson, J. Chen, Carbon catabolite repression of the maltose

transporter revealed by X-ray crystallography. *Nature* **499**, 364-368 (2013).

34. S. B. Ko *et al.*, Gating of CFTR by the STAS domain of SLC26 transporters. *Nat. Cell. Biol.* **6**, 343-350

(2004).

35. V. D. Nikitushkin *et al.*, The main pigment of the dormant *Mycobacterium smegmatis* is porphyrin.

FEMS Microbiol. Lett. **363** (2016).

36. W. Lee, B. C. VanderVen, R. J. Fahey, D. G. Russell, Intracellular Mycobacterium tuberculosis exploits

host-derived fatty acids to limit metabolic stress. *J. Biol. Chem.* **288**, 6788-6800 (2013).

37. A. Martin, J. Daniel, The ABC transporter Rv1272c of *Mycobacterium tuberculosis* enhances the import

of long-chain fatty acids in *Escherichia coli*. *Biochem. Biophys. Res. Commun.* **496**, 667-672 (2018).

Figure 1. Affinity purification with His₆-YrbE1B defines the composition of Mce1 complexes. SDS-PAGE and α-His immunoblot analyses of proteins affinity-purified from *M. smegmatis* Δ*mce1R* cells expressing His-tagged YrbE1A (p*his-yrbE1A*) or YrbE1B (p*his-yrbE1B)*. pMV306hsp was used as the empty vector control. For protein identification from gel bands, the top three proteins detected in the p*his-yrbE1B* lane are listed. For protein identification from gel sections (blue: 35–90 kDa, green: 18–35 kDa), proteins with more than ten total peptides detected and significantly enriched in the p*his-yrbE1B* lane are listed. CBB, Coomassie brilliant blue.

Figure 2. Both Msmeg_6540 and Mce1A are important for fatty acid uptake. (A) [^{14}C]-palmitic acid

uptake kinetics of indicated *M. smegmatis* strains. Each data point (mean ± standard deviation) represents

results from three technical replicates. The uptake profiles are representative of at least three independent

experiments. **(B)** Growth curves of the same strains in **(A)** in Sauton's medium containing palmitate as the

sole carbon source or (inset) 7H9 medium containing glycerol. Each data point (mean ± standard deviation)

represents results from three biological replicates.

Figure 3. YrbE1A/B-MceG form a stable asymmetric complex with ATPase activity. (A) Size exclusion chromatographic profile of the purified complex comprising YrbE1A, His$_6$-YrbE1B and MceG. SDS-PAGE analysis of the peak corresponding to the complex (black arrow head) is shown. **(B)** Enzyme-coupled ATPase assay of 0.1 µM YrbE1A/B-MceG and YrbE1A/B-MceG$_{K43A}$ in detergent micelles at 37 ºC. Average ATP hydrolysis rates from three technical triplicates were plotted against ATP concentrations, and fitted to an expanded Michaelis-Menten equation that includes a term for Hill coefficient (n). YrbE1A/B-MceG: K$_m$ = 284 ± 311µM, k_{cat} = 0.184 ± 0.125 µM ATP s^{-1}/µM complex, n = 1.28 ± 0.57. The data for the mutant failed to fit to the equation with significant confidence. Each data point is shown as mean ± standard deviation. **(C)** SDS-PAGE and α-His immunoblot analyses of proteins affinity-purified from *E. coli* cells over-expressing His$_6$-YrbE1A or His$_6$-YrbE1B together with MceG. CBB, Coomassie brilliant blue. **(D)** An AlphaFold2 structural model of the YrbE1A/B-MceG complex (*green*: YrbE1A, *magenta*: YrbE1B, *cyan* and *orange*: MceG), with respective calculated interface areas indicated.

Figure 4. Msmeg_0959 negatively regulates the Mce1 transporter. (A) SDS-PAGE and α-His

immunoblot analyses of proteins affinity-purified from *E. coli* cells over-expressing YrbE1A, MceG, and

either wild-type His$_6$-YrbE1B (WT) or the helix-to-loop mutant His$_6$-YrbE1B (GS), in the presence (+) or

absence (-) of Msmeg_0959. **(B)** SDS-PAGE and α-His immunoblot analyses of proteins affinity-purified

from *M. smegmatis* cells over-expressing Mce1N from different species (Rv0513 from *M. tuberculosis* and

Ml2418c from *M. leprae*) and His$_6$-YrbE1B as the bait protein. **(C)** An overlay of AlphaFold2 models of

YrbE1B (*magenta*)-MceG (*orange* surface) and YrbE1B-Msmeg_0959 (*blue*). The helix-to-loop mutation

located on the putative short helix at the YrbE1B-Msmeg_0959 interface is highlighted in *yellow*. The steric

clash between MceG and Msmeg_0959 is indicated by *red* arrows. **(D)** [^{14}C]-palmitic acid uptake kinetics

of indicated *M. smegmatis* strains overexpressing Mce1N from different mycobacterial species. Each data

point (mean ± SD) represents results from three technical replicates. The uptake profiles are representative

of three independent experiments.

Figure 5. A proposed model of Mce1 complexes and their regulation by Mce1N. Mce1 complexes are ABC transporters for fatty acids, with YrbE1A/B as TMDs, MceG as NBDs, and heterohexameric MCE-domain proteins as SBPs. There are two variants of Mce1 complexes in *M. smegmatis*, each containing Msmeg_6540 or Mce1A in addition to Mce1B–F. The relative order of protomers within heterohexameric SBP complexes is indicated based on the AlphaFold2 models (*SI Appendix*, Fig. S5*B*). Mce1N is a negative regulator of the Mce1 system; it specifically binds to YrbE1B and prevents its interaction with MceG.