Conformational changes induced by chaperone interactions facilitate lipid transport critical for bacterial outer membrane asymmetry

Jiang Yeow[1] and Shu-Sin Chng[1,2,*]

Affiliation:

[1]Department of Chemistry, National University of Singapore, Singapore 117543

[2]Singapore Center for Environmental Life Sciences Engineering, National University of Singapore (SCELSE-NUS), Singapore 117456

[*]To whom correspondence should be addressed: chmchngs@nus.edu.sg

Author Contributions: J.Y. and S.-S.C designed research; J.Y. performed wet lab experiments described in this work; J.Y. and S.-S.C. analyzed and discussed data; J.Y. and S.-S.C. wrote the paper.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences; Biochemistry

Keywords: bacteria; outer membrane; lipid transport; phospholipid; lipid asymmetry; membrane protein; protein complex; channel

This PDF file includes:

Main Text

Figures 1 to 6

Abstract

The outer membrane (OM) of Gram-negative bacteria is an asymmetric lipid bilayer with outer leaflet lipopolysaccharides (LPS) exposed to extracellular milieu and inner leaflet phospholipids (PLs) facing the periplasm. This unique lipid asymmetry is the key to its innate drug resistance, rendering the OM impermeable to external insults, including antibiotics and bile salts. To maintain this OM barrier, the OmpC-Mla system removes mislocalized PLs from the OM outer leaflet, and transports them back to the inner membrane (IM); in the first step, the OM OmpC-MlaA complex transfers PLs to the periplasmic chaperone MlaC. This process likely occurs via a hydrophilic channel in MlaA, yet mechanistic details have remained elusive. Here, we obtain a molecular view of the architecture of the MlaA-MlaC transient complex by mapping the interaction surfaces between MlaA and MlaC in *Escherichia coli*. We show that electrostatic interactions are important for MlaC recruitment, and that MlaC eventually binds MlaA at the periplasmic face in a manner that juxtaposes the MlaA channel and the MlaC lipid binding cavity. We further provide evidence for conformational changes in the MlaA channel that correlate with functional states of MlaA, as well as interactions with MlaC binding and OM porins. Our work offers novel insights into the molecular mechanism for lipid transfer by the OmpC-MlaA complex for overall retrograde transport of PLs to the IM to maintain OM lipid asymmetry.

Introduction

Gram-negative bacteria assemble two lipid bilayers in their cell envelopes - the inner membrane (IM) and the outer membrane (OM) that encapsulate the cytoplasm and periplasm, respectively. At the OM, lipopolysaccharides (LPS) pack tightly with divalent cations and occupy the outer leaflet, while phospholipids (PL) reside mostly in the inner leaflet, presenting a unique bilayer with extreme asymmetry (1-3). The outer leaflet of LPS forms a stable layer with drastically reduced permeability, offering an effective protective barrier against the entry of toxic compounds, such as detergents and antibiotics. Proper establishment and maintenance of lipid asymmetry is thus critical for OM barrier function, which allows Gram-negative bacteria to survive in the most hostile environments, including the mammalian intestinal tract.

How OM lipid asymmetry is achieved has been extensively studied. It is well known that LPS are transported and inserted into the outer leaflet of the OM via the Lpt protein bridge machinery (4). Other major OM components, i.e. β-barrel proteins and OM lipoproteins, are assembled into the membrane via the Bam (5) and Lol (6) pathways, respectively. It is less understood how exactly PLs are brought to the OM (7), but recent studies suggest the involvement of a redundant set of AsmA-like proteins in this process (8-10). Evidently, balancing the levels of all these OM components during active cell growth and division would be critical to ensure OM stability, and hence lipid asymmetry. In this regard, the Tol-Pal complex has been implicated in lipid homeostasis, possibly via removal of excess PLs from the OM (11). When coordination across these pathways is sub-optimal, PLs may become mislocalized to the outer leaflet of the OM, forming PL bilayer patches that compromise OM lipid asymmetry and barrier function (12, 13). To cope with these defects, additional cellular mechanisms are deployed to eliminate PLs that end up aberrantly in the outer leaflet of the OM. The OM phospholipase, PldA, hydrolyzes acyl chains from these PLs (14), while the OM acyltransferase, PagP, transfers an acyl chain from outer leaflet PLs to LPS and phosphatidylglycerol (PG) (15, 16). Finally, the OmpC-Mla system, an established PL trafficking pathway, extracts PLs from the outer leaflet of the OM and shuttles them back to the IM (17, 18).

The OmpC-Mla system, first identified in *Escherichia coli* based on homology to the TGD2 pathway in *Arabidopsis thaliana*, comprises the OM lipoprotein MlaA in complex with osmoporin OmpC, the periplasmic chaperone MlaC, and the IM ATP-binding cassette (ABC) transporter MlaFEDB (17-19).

Removing *ompC* or any *mla* component results in PL accumulation in the outer leaflet of the OM of *E. coli*,

indicating a role in maintaining OM lipid asymmetry (17, 18). To achieve that, the OmpC-MlaA complex

somehow extracts PLs from the outer leaflet of the OM, hands them over to MlaC, which in turn transfers

these PLs into the IM via the MlaFEDB complex (20-22). Many recent biochemical and structural studies

have provided detailed insights into ATP-dependent PL transfer steps at the IM (19, 20, 23-30). In particular,

we now know that when PL-bound MlaC arrives at the IM, it can spontaneously transfer the lipid molecule to

the binding cavity of MlaFEDB (21). Even though this initial process is reversible (20, 21, 25), ATP

binding/hydrolysis by the MlaFEDB complex ultimately catalyzes the transfer of PLs into the IM, driving

overall retrograde transport from the OM (20, 21). How ATP binding/hydrolysis may be coupled to PL

transport in the MlaFEDB complex can be partly inferred from recently solved structures (20, 26-30), though

detailed mechanistic understanding is still limited.

Lipid transfer at the OM is somewhat less understood, especially how retrograde PL movement from

the outer leaflet of the OM to MlaC can be facilitated in an energy-independent manner. The OM lipoprotein

MlaA interacts with trimeric porins OmpC and OmpF, where OmpC appears to play a more critical role in

Mla function (18). It has been established that, unlike canonical lipoproteins, MlaA resides entirely within the

OM bilayer and makes extensive contacts with porins at their dimeric interfaces. Moreover, MlaA itself forms

a hydrophilic channel in the OM, which may be gated by conformational movements of an adjacent β-hairpin

loop motif (31, 32). The MlaA channel may provide a plausible path for PL translocation across the OM;

however, we do not yet have clear ideas of how MlaA actually sits in the OM lipid bilayer, as well as the

expected conformational changes that should occur in the protein during PL transport. MlaC binds PLs with

presumed high affinity (24, 25, 33), and has been demonstrated to interact with MlaA in vivo (24) and in vitro

(23). Yet it remains elusive how exactly MlaA interacts with MlaC to allow spontaneous, productive transfer

of PLs.

Here, we elucidate a molecular picture of the transient MlaA-MlaC complex during PL transfer at the

OM. Using in vivo photo- and disulfide-crosslinking, we demonstrate that MlaC interacts with MlaA directly

at its periplasmic base, in a manner that juxtaposes the lipid binding cavity of MlaC with the hydrophilic

channel of MlaA. In addition, mutational analyses reveal that MlaA recruits MlaC to the OM via additional

electrostatic interactions between a C-terminal tail helix of MlaA and a surface charged patch on MlaC, an

observation corroborated by an AlphaFold2 predicted model of the interaction complex. Furthermore, we

establish solvent-accessibility changes in a key channel residue in MlaA that are influenced by interactions

with MlaC and porins (OmpC/F), as well as functionality of MlaA itself. Collectively, our work provides

mechanistic insights into how MlaC interacts with MlaA to induce conformational changes that may facilitate

retrograde PL transfer from the outer leaflet of the OM to MlaC to maintain lipid asymmetry.

Results

MlaC interacts with α-helices at the periplasmic opening of the hydrophilic channel of MlaA

To receive PLs from MlaA, MlaC would have to gain access to the hydrophilic MlaA channel from

the periplasmic side, albeit transiently. To map the region(s) on MlaA that contacts MlaC, we replaced ~27

residues on the periplasmic face of *E. coli* MlaA separately with *para*-L-benzoylphenylalanine (*p*Bpa) (34),

and examined which positions allow photoactivatable crosslinks with MlaC. We found five positions on MlaA

that demonstrated strong photoactivatable crosslinks with MlaC when substituted with *p*Bpa (**Fig. 1A and**

1B). Another five positions exhibited weaker crosslinks to MlaC (**Fig. S1**). Interestingly, these residues are

all located on the C-terminal helices (Q195-N226) of MlaA extending from the channel opening in the

periplasmic face (**Fig. 1C**), suggesting that MlaC can be juxtaposed to receive lipids from MlaA at the base

of the channel.

We have also previously established the residues on MlaC that contact MlaA via the same

photocrosslinking approach (24). To determine reciprocal contact points between the two proteins, we

introduced cysteines at positions on either protein that displayed strong photoactivatable crosslinks, and

monitored disulfide bond formation between these MlaA and MlaC cysteine variants in cells. We tested 24

combinations of cysteine residue-pairs from four and six positions on MlaA and MlaC, respectively. Of these

pairs, 11 enabled disulfide bond formation between MlaA and MlaC, indicating that these residue positions

can lie in close proximity when the two proteins interact (**Fig. 2A**). MlaA^{Q195C} and MlaC^{Q151C} did not form

disulfide crosslinks with any partner cysteine variants tested. In contrast, Q205, M212 and F223 on MlaA,

and K128, S172 and T175 on MlaC, each allowed disulfide bond formation with multiple tested positions on

their partner protein, when replaced with cysteine (**Figs. 2A and 2B**). The strongest disulfide crosslink was MlaA^{Q205C}-MlaC^{V171C} (**Figs. 2A and S2**), highlighting perhaps a major interaction mode in the transient complex. Interestingly, Q205 lies directly at the base of the hydrophilic channel of MlaA while V171 is positioned at the entrance of the lipid binding cavity of MlaC (**Fig. 2B**), reflecting an approximate arrangement that could lead to productive lipid transfer.

Electrostatic interactions between the C-terminal tail helix of MlaA and a surface charged patch on MlaC are necessary for MlaC recruitment

We have shown that MlaC interacts with the C-terminal α-helices at the base of MlaA. However, there is presumably an unstructured region beyond these interacting helices (G227-E251; *E. coli* numbering) missing from the reported MlaA crystal structures (32); yet, in the recent structural model predicted by AlphaFold (AF-P76506-F1) (35), the extreme end (N238-E251) of this 'tail' region was confidently modelled as a short α-helix (**Fig. 3A**). To understand if this C-terminal tail helix is also important for MlaC interaction, we made several MlaA deletion constructs, and probed for MlaC interaction. While removing residues after P232 (MlaA^{CTD4}) conferred milder SDS/EDTA sensitivity than cells lacking MlaA (**Fig. S3A**), it had completely abolished disulfide bond formation between MlaA^{Q205C} and MlaC^{V171C} (**Fig. 3B**). In fact, deletion of the last 9 amino acids on MlaA^{D243X} (MlaA^{CTD6}), a highly negatively charged segment on the putative terminal helix, was sufficient to disrupt disulfide crosslinks with MlaC. To test the importance of electrostatics, we also constructed two sets of Arg mutants at a single face of this helix, MlaA$^{D243R/D244R}$ (MlaA^{2DD2R}) and MlaA$^{D247R/E251R}$ (MlaA^{2DE2R}) (**Fig. 3A**), and found that these charged-reversed variants drastically reduced disulfide crosslinks between MlaA^{Q205C} and MlaC^{V171C} (**Fig. 3B**). The different MlaA variants are still able to pull down trimeric OmpC, indicating they preserve overall MlaA structure (**Fig. S4**). We conclude that the negatively-charged residues on the C-terminal tail helix of MlaA are also critical for MlaC interaction.

We next asked if there are specific positively-charged regions on MlaC that interact with the C-terminal tail helix of MlaA. We identified seven clusters of positively-charged residues (Arg/Lys) all around the structure of apo-MlaC (25) (**Fig. 4A**), and successfully converted five of these clusters to aspartates

separately. MlaC$^{K84D/R90D}$ (MlaC1) abolished disulfide crosslinks between MlaA^{Q205C} and MlaC^{V171C}, while

MlaC^{K71D} (MlaC0) and MlaC$^{R143D/R147D}$ (MlaC3) led to partial loss (**Fig. 4B**), confirming the overall

importance of electrostatics. These mutations did not introduce any SDS/EDTA sensitivity (**Fig. S3B**),

however, suggesting they may only impact recruitment efficiency, but not overall lipid transport function.

Remarkably, combining MlaA$^{D243R/D244R}$ (MlaA^{2DD2R}) with MlaC$^{K84D/R90D}$ (MlaC1) fully restored disulfide

bond formation (**Fig. 4C**), implying that residues D243/D244 on MlaA directly contact residues K84/R90 on

MlaC during MlaA-MlaC interaction.

To obtain a possible picture of the MlaA-MlaC interface, we applied AlphaFold2 to predict the

structural model of the MlaA-MlaC complex (35-37). Amazingly, the model not only revealed an arrangement

of MlaA and MlaC that is fully consistent with our in vivo crosslinking data (especially pBpa), it also correctly

predicted the charge-charge interactions between the D243/D244 on MlaA and K84/R90 on MlaC (**Fig. 4D**).

The C-terminal tail helix of MlaA is connected to the rest of the protein via a flexible linker, suggesting that

this tail helix might serve as a bait to recruit apo-MlaC to the OM complex via electrostatic attraction.

Importantly, once MlaC is stably bound and positioned, the hydrophilic channel in MlaA becomes continuous

with the lipid binding cavity of MlaC (**Fig. 4D**), providing a glimpse into the route for PL transfer from the

outer leaflet of the OM to MlaC.

Interactions with MlaC modulate conformational changes in the MlaA channel

Productive MlaA-MlaC interactions would be required for PL transfer from the OmpC-MlaA complex

to MlaC. We hypothesize that such interactions could elicit conformational changes in MlaA to facilitate PL

transfer. To test this idea, we examined the solvent accessibilities, as an indication of possible conformational

states, of specific residues within the MlaA channel in the absence of MlaC (**Fig. 5A**). We have previously

shown using the substituted cysteine accessibility method (SCAM) that many hydrophilic residues in the

MlaA channel are fully accessible to a membrane-impermeable thiol-reactive reagent ((2-sulfonatoethyl)

methanethiosulfonate; MTSES) in wild-type cells. One exception is K184, which exhibited partial MTSES

accessibility when substituted with cysteine; within the MlaA population in wild-type cells, K184C is

accessible to MTSES in some MlaA molecules but not others, in comparable proportions (**Fig. 5B**).

Interestingly, removing MlaC resulted in K184C becoming MTSES-inaccessible in the entire population of MlaA (**Fig. 5B**), an effect not seen for other channel residues tested (**Fig. S5**). This shift in accessibility of K184C was reversed by expressing wild-type MlaC in trans, but not the charge-reversed MlaC1 variant that disrupted MlaA-MlaC interactions (**Fig. 5C**). This suggests that interaction with MlaC is required for MlaA to adopt the state where K184C is solvent accessible. Furthermore, we observed that K184C in MlaA also became MTSES-inaccessible in the Δ*mlaD* strain (**Fig. 5B**). While MlaC is still present in this strain, it is likely bound to PLs (i.e. holo form), given that off-loading of MlaC is not possible in the absence of an intact ABC transporter in the IM. Overall, our results indicate that the binding and release of apo-MlaC may influence the solvent accessibilities and conformational states of the MlaA channel.

We have also considered the possibility that shifts in MTSES-accessibility could be due to OM asymmetry defects in these *mla* strains. This is unlikely, however, since K184C MTSES-accessibility in other OM-defective strains (Δ*tolB,* Δ*pal*) (11) was similar to that in wild-type cells (**Fig. S5A**). Supporting this observation, the overexpression of phospholipase PldA, which can rescue OM asymmetry defects in *mla* mutants (17), was unable to reverse the observed shifts in accessibility of K184C in both Δ*mlaC* and Δ*mlaD* strains (**Fig. S5B**). We conclude that accessibility changes in K184 in MlaA are not due to the general loss of lipid asymmetry in the OM.

Non-functional MlaA variants adopt conformational states that no longer respond to interactions with MlaC

It is intriguing that K184C exhibits accessibility changes that may reflect different conformational states in the MlaA channel, possibly due to functional interactions with apo-MlaC. We next asked whether this same residue can also help us discern the functionality of MlaA itself. We first examined the MTSES-accessibility of K184C in known MlaA variants. MlaA[G141P/G143P/G145P] (MlaA[3G3P]) is a non-functional variant believed to have restricted flexibility in a β-hairpin loop that may gate the MlaA hydrophilic channel (31). MlaA[ΔN43F44] (MlaA*) is a gain-of-function variant that is thought to have poor control of loop dynamics, resulting in PL flipping from the inner to the outer leaflet of the OM (8, 31, 38). Interestingly, K184C in the entire population of MlaA[3G3P] or MlaA* appeared to be fully accessible to MTSES in cells, contrary to that

observed in wild-type MlaA (**Fig. 5D**). Furthermore, removing MlaC or MlaD did not alter the accessibility of K184C in both MlaA variants, suggesting that MlaA^{3G3P} and MlaA* are in conformational states that cannot respond to interactions with apo-MlaC. Both MlaA^{3G3P} and MlaA* are therefore non-functional in facilitating retrograde PL transfer to MlaC.

An outstanding question in the OmpC-Mla system relates to why only OmpC, but not OmpF, appears to function together with MlaA. Under high osmolarity conditions, OmpC is produced at higher levels than OmpF (18). Therefore, to gain insights into the role of porins in the pathway without confounding effects due to expression levels, we went on to examine the solvent accessibility of MlaA K184C in the $\Delta ompC \Delta ompF$ double mutant complemented in trans with either *ompC* or *ompF*. Interestingly, removing both OmpC and OmpF completely shifted the MlaA$_{K184C}$ population to a MTSES-accessible state (**Fig. 5E**), and expressing OmpC or OmpF in trans (at comparable levels, **Fig. S6**) restored the population of MlaA$_{K184C}$ to liken that in wild-type cells. These results indicate that free MlaA may not adopt a native conformational state in the OM, and that interactions with either porin is required yet sufficient for scaffolding MlaA. We further noted that the MlaA$_{K184C}$ population in just the $\Delta ompC$ mutant was already fully MTSES-accessible, and this observation was independent of the presence of MlaC (**Fig. 5F**). Overall, MlaA in strains lacking OmpC alone may also mostly not be bound to porins (owing to low levels of OmpF), and therefore largely adopts a non-functional conformational state, mirroring what we have observed in the MlaA variants. We conclude that interactions with trimeric porins allow MlaA to adopt functional states important for lipid transport.

Discussion

How the OmpC-MlaA complex transfers PLs from the outer leaflet of the OM to MlaC, as part of the role of the OmpC-Mla system in maintaining lipid asymmetry, is not known. In this study, we have mapped the interacting surfaces between MlaA and MlaC, resolved the molecular architecture of the transient MlaA-MlaC complex, as well as revealed conformational changes of MlaA during PL transfer to MlaC. Using molecular cross-linking, we have shown that MlaC binds MlaA at its periplasmic face in a manner where the lipid binding cavity of MlaC is aligned with the hydrophilic channel of MlaA to create a continuous path for PLs. We have also demonstrated that electrostatic interactions between the C-terminal tail helix of MlaA and

a charged patch on MlaC are required for (initial) recruitment. Finally, we have uncovered that wild-type MlaA adopts distinct conformational states that is likely modulated by interactions with (apo)-MlaC. Our work provides a mechanistic glimpse of how PLs may be transferred across the MlaA channel into the MlaC binding pocket for onward shuttling to the IM.

The requirement of electrostatic interactions between MlaA and MlaC allows us to infer a mechanism for how MlaA ensures effective recruitment and binding of apo- over holo-MlaC, thereby facilitating overall retrograde transfer of PLs. We have established that the positively-charged patch formed by residues K84/R90 on MlaC, and the negatively-charged C-terminal tail helix residues D243/D244 on MlaA, directly interact (**Fig. 4**), and are important for MlaC recruitment. We note that residues K84/R90 on MlaC are exposed and accessible on apo-MlaC (25). Remarkably, this positively-charged patch is somewhat occluded by movement of a surface loop on MlaC upon PL binding (23) (**Fig. 6**), limiting access by the MlaA tail helix. Such differential accessibility on MlaC likely allows preferential recruitment of apo-MlaC over holo-MlaC at the OM (**Fig. 6**). Post-recruitment, additional interactions between MlaA and MlaC occur at the base of the MlaA hydrophilic channel, stabilizing the complex for productive PL transfer. Once MlaC picks up a PL molecule, however, the charge-charge interaction with the MlaA tail helix is lost due to sterics (**Fig. 6**), leading to possible destabilization of the MlaA-MlaC complex, and release of holo-MlaC from the OM. In a way, the MlaA tail helix, possibly extended almost halfway into the periplasmic space via a flexible linker (25 residues, ~88 Å), serves both as an electrostatic bait for MlaC during recruitment, and also a latch that locks down the transient MlaA-MlaC complex during PL transfer. This could be a fascinating strategy to favor apo-MlaC recruitment/binding, and may ensure that PL transfer takes place in a retrograde fashion.

In the context of MlaC binding and release, we can now also introduce a model for MlaA cycling between distinct conformation states during PL transfer (**Fig. 6**). We have shown that MlaA exists in at least two states in wildtype strains, where K184 is somehow accessible to the aqueous environment (State B), or not (State A). Interestingly, K184 is positioned midway up the MlaA channel in the membrane, and close to the gating loop; we therefore speculate that the MlaA channel may be open in State B, yet closed in State A. In the absence of MlaC, or with disrupted MlaA-MlaC recruitment, all of MlaA appears to become confined to State A (channel closed). Accordingly, apo-MlaC recruitment and binding is likely required to induce MlaA

to shift to the channel-open State B, allowing productive PL transfer. It is conceivable that holo-MlaC then no longer binds MlaA in quite the same way as apo-MlaC, possibly explaining why MlaA also adopts a channel-closed State A in the absence of MlaD, where all of MlaC is presumably in the holo-form. Such coordination between apo-MlaC binding and conformational changes in wild-type MlaA may favor retrograde PL transfer at the OM.

Conformational analysis using K184 accessibility also facilitated our understanding of non-functional variants/states of MlaA. We have found that free MlaA (not in complex with porins), MlaA[3G3P] and MlaA* variants all adopt conformation state(s) where K184 is fully solvent accessible and is no longer responsive to the presence of MlaC (binding) (**Fig. 5**). Since K184 accessibility only produces binary output, however, free MlaA and these variants may well have distinct conformations that still allow K184 to be accessible to the aqueous environment. MlaA[3G3P] is believed to be a loop-locked non-functional mutant, while MlaA* is thought to have lost control of its gating loop (31). Our study suggests that even though MlaA* has gain-of-function possibly in allowing PLs to flip across the OM to the cell surface (8, 38), this variant is in fact non-functional in the native role of MlaA. Consistent with this idea, it has recently been demonstrated that MlaA* is unable to support retrograde PL transfer in vitro (20). We have also now demonstrated that MlaA requires trimeric porins like OmpC and OmpF for scaffolding and possibly achieving functional conformations, thus overall accounting for porin function in the Mla system (18). Curiously, we did not observe changes in K184 accessibility in a strain expressing OmpC[R92A] (**Fig. 5E**), a mutation at the dimeric interface of OmpC trimers (where MlaA binds) that impacts OM lipid asymmetry (31). Therefore, beyond proper scaffolding of MlaA, OmpC may play yet-to-be-identified roles that facilitate PL transport by MlaA.

Ultimately, the exact mechanism by which the OmpC-MlaA complex extracts mislocalized PLs from the outer leaflet of the OM and transfers them to MlaC will require additional resolution into the various conformational states we have revealed herein. Structural information of wild-type and mutant OmpC-MlaA complexes in the context of a native lipid bilayer, with and without MlaC bound, would provide mechanistic insights into the process. Given that OM lipid asymmetry is critical for overall barrier function, novel insights into the OmpC-MlaA-(MlaC) complex will also guide us in developing useful strategies to inhibit the OmpC-Mla pathway in the future.

Acknowledgements

284 **Acknowledgements**

286 J.Y. was supported by the National University of Singapore Graduate School of Integrative Sciences and

287 Engineering Scholarship (ISEP). This work was supported by the Singapore Ministry of Health National

288 Medical Research Council under its Open Fund Individual Research Grant (MOH-000145) (to S.S.C.). The

289 authors declare no conflict of interest.

References

1. Kamio Y & Nikaido H (1976) Outer membrane of Salmonella typhimurium: accessibility of phospholipid head groups to phospholipase c and cyanogen bromide activated dextran in the external medium. *Biochemistry* 15(12):2561-2570.

2. Funahara Y & Nikaido H (1980) Asymmetric localization of lipopolysaccharides on the outer membrane of Salmonella typhimurium. *J Bacteriol* 141(3):1463-1465.

3. Nikaido H (2003) Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev* 67(4):593-656.

4. Okuda S, Sherman DJ, Silhavy TJ, Ruiz N, & Kahne D (2016) Lipopolysaccharide transport and assembly at the outer membrane: the PEZ model. *Nat Rev Microbiol* 14(6):337-345.

5. Konovalova A, Kahne DE, & Silhavy TJ (2017) Outer Membrane Biogenesis. *Annu Rev Microbiol* 71:539-556.

6. Okuda S & Tokuda H (2011) Lipoprotein sorting in bacteria. *Annu Rev Microbiol* 65:239-259.

7. Yeow J & Chng SS (2022) Of zones, bridges and chaperones - phospholipid transport in bacterial outer membrane assembly and homeostasis. *Microbiology (Reading)* 168(4).

8. Grimm J*, et al.* (2020) The inner membrane protein YhdP modulates the rate of anterograde phospholipid flow in Escherichia coli. *Proc Natl Acad Sci U S A* 117(43):26907-26914.

9. Ruiz N, Davis RM, & Kumar S (2021) YhdP, TamB, and YdbH Are Redundant but Essential for Growth and Lipid Homeostasis of the Gram-Negative Outer Membrane. *mBio* 12(6):e0271421.

10. Douglass MV, McLean AB, & Trent MS (2022) Absence of YhdP, TamB, and YdbH leads to defects in glycerophospholipid transport and cell morphology in Gram-negative bacteria. *PLoS Genet* 18(2):e1010096.

11. Shrivastava R, Jiang X, & Chng SS (2017) Outer membrane lipid homeostasis via retrograde phospholipid transport in Escherichia coli. *Mol Microbiol* 106(3):395-408.

12. Jia W*, et al.* (2004) Lipid trafficking controls endotoxin acylation in outer membranes of Escherichia coli. *J Biol Chem* 279(43):44966-44975.

13.   Wu T*, et al.* (2005) Identification of a multicomponent complex required for outer membrane
biogenesis in Escherichia coli. *Cell* 121(2):235-245.

14.   Dekker N (2000) Outer-membrane phospholipase A: known structure, unknown biological function.
Mol Microbiol 35(4):711-717.

15.   Bishop RE (2005) The lipid A palmitoyltransferase PagP: molecular mechanisms and role in bacterial
pathogenesis. *Mol Microbiol* 57(4):900-912.

16.   Dalebroux ZD, Matamouros S, Whittington D, Bishop RE, & Miller SI (2014) PhoPQ regulates acidic
glycerophospholipid content of the Salmonella Typhimurium outer membrane. *Proc Natl Acad Sci U*
S A 111(5):1963-1968.

17.   Malinverni JC & Silhavy TJ (2009) An ABC transport system that maintains lipid asymmetry in the
gram-negative outer membrane. *Proc Natl Acad Sci U S A* 106(19):8009-8014.

18.   Chong ZS, Woo WF, & Chng SS (2015) Osmoporin OmpC forms a complex with MlaA to maintain
outer membrane lipid asymmetry in Escherichia coli. *Mol Microbiol* 98(6):1133-1146.

19.   Thong S*, et al.* (2016) Defining key roles for auxiliary proteins in an ABC transporter that maintains
bacterial outer membrane lipid asymmetry. *Elife* 5.

20.   Tang X*, et al.* (2021) Structural insights into outer membrane asymmetry maintenance in Gram-
negative bacteria by MlaFEDB. *Nat Struct Mol Biol* 28(1):81-91.

21.   Low WY, Thong S, & Chng SS (2021) ATP disrupts lipid-binding equilibrium to drive retrograde
transport critical for bacterial outer membrane asymmetry. *Proc Natl Acad Sci U S A* 118(50).

22.   Low WY & Chng SS (2021) Current mechanistic understanding of intermembrane lipid trafficking
important for maintenance of bacterial outer membrane lipid asymmetry. *Curr Opin Chem Biol*
65:163-171.

23.   Ekiert DC*, et al.* (2017) Architectures of Lipid Transport Systems for the Bacterial Outer Membrane.
Cell 169(2):273-285 e217.

24.   Ercan B, Low WY, Liu X, & Chng SS (2019) Characterization of Interactions and Phospholipid
Transfer between Substrate Binding Proteins of the OmpC-Mla System. *Biochemistry* 58(2):114-119.

25.   Hughes GW, *et al.* (2019) Evidence for phospholipid export from the bacterial inner membrane by the

343   Mla ABC transport system. *Nat Microbiol* 4(10):1692-1705.

344   26.   Chi X, *et al.* (2020) Structural mechanism of phospholipids translocation by MlaFEDB complex. *Cell*

345   *Res* 30(12):1127-1135.

346   27.   Coudray N, *et al.* (2020) Structure of bacterial phospholipid transporter MlaFEDB with substrate

347   bound. *Elife* 9.

348   28.   Zhang Y, Fan Q, Chi X, Zhou Q, & Li Y (2020) Cryo-EM structures of Acinetobacter baumannii

349   glycerophospholipid transporter. *Cell Discov* 6(1):86.

350   29.   Mann D, *et al.* (2021) Structure and lipid dynamics in the maintenance of lipid asymmetry inner

351   membrane complex of A. baumannii. *Commun Biol* 4(1):817.

352   30.   Zhou C, *et al.* (2021) Structural Insight into Phospholipid Transport by the MlaFEBD Complex from

353   P. aeruginosa. *J Mol Biol* 433(13):166986.

354   31.   Yeow J, *et al.* (2018) The architecture of the OmpC-MlaA complex sheds light on the maintenance of

355   outer membrane lipid asymmetry in Escherichia coli. *J Biol Chem*.

356   32.   Abellon-Ruiz J, *et al.* (2017) Structural basis for maintenance of bacterial outer membrane lipid

357   asymmetry. *Nat Microbiol* 2(12):1616-1623.

358   33.   Huang YM, *et al.* (2016) Molecular dynamic study of MlaC protein in Gram-negative bacteria:

359   conformational flexibility, solvent effect and protein-phospholipid binding. *Protein Sci* 25(8):1430-

360   1437.

361   34.   Chin JW, Martin AB, King DS, Wang L, & Schultz PG (2002) Addition of a photocrosslinking amino

362   acid to the genetic code of Escherichiacoli. *Proc Natl Acad Sci U S A* 99(17):11020-11024.

363   35.   Jumper J, *et al.* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*

364   596(7873):583-589.

365   36.   Evans R, *et al.* (2021) Protein complex prediction with AlphaFold-Multimer.

366   *bioRxiv*:2021.2010.2004.463034.

367   37.   Mirdita M, *et al.* (2022) ColabFold: making protein folding accessible to all. *Nat Methods*.

368   38.   Sutterlin HA, *et al.* (2016) Disruption of lipid homeostasis in the Gram-negative cell envelope activates

369   a novel cell death pathway. *Proc Natl Acad Sci U S A* 113(11):E1565-1574.

  Figure legends

Figure 1. MlaA contacts MlaC in cells via C-terminal α-helices at the periplasmic opening of its hydrophilic channel. (A) Representative immunoblots showing UV-dependent formation of strong crosslinks between MlaA and MlaC in Δ*mlaA* cells expressing MlaA-His substituted with *p*Bpa at indicated positions from the pCDF plasmid. Additional but unidentified MlaA*p*Bpa-His crosslinked adducts were detected and denoted with an asterisk (*). (B) Membrane (*left*) and periplasmic (*right*) views of cartoon representations of the crystal structure of *Klebsiella pneumoniae* OmpK36 (*green*)-MlaA (*cyan*) (PDB 5NUP) (32) with positions that crosslink to MlaC highlighted. Residues exhibiting strong and weak photo-crosslinks with MlaC are illustrated in *red* and *pink* sticks, respectively. The OM boundaries are indicated as *gray* dashed lines.

Figure 2. MlaC interacts with MlaA in a manner that juxtaposes the MlaA channel and the MlaC lipid binding

cavity. (A) Representative immunoblots showing formation of disulfide crosslinks between MlaA and MlaC

in Δ*mlaA* Δ*mlaC* cells expressing cysteine-substituted MlaA-His and MlaC-His from the pCDF and pET22/42

plasmids, respectively. Samples were subjected to non-reducing (*top*) or reducing (*bottom*) SDS-PAGE prior to

immunoblotting. (C) Cartoon representation of *Kp*MlaA (*cyan*) (PDB 5NUP) (32) and holo-MlaC (*pink*, PL in

388   *red* spheres) (PDB 5UWA) (23). Residues on MlaC which displayed disulfide crosslinking with MlaA, and

389   vice versa, are illustrated in *red* and *blue* sticks, respectively. MlaC residues mapped to MlaA residues Q205C,

390   M212C and F223C are connected by *beige, blue* and *pink lines* respectively.

Figure 3. Negatively-charged residues on the putative C-terminal tail helix of MlaA are important for MlaC

recruitment. (A) Cartoon representation of the *Ec*MlaA (*teal*) model generated from AlphaFold (AF-P76506-

F1) (35) with the putative C-terminal tail helix coloured *blue*. C-terminal truncation positions (CTDs) are

annotated. MlaC-crosslinking residue Q205 is labelled in *orange* sticks. Charge-reversed residues in

MlaA[2DD2R] and MlaA[2DE2R] are labelled in *red* and *pink* sticks respectively. (B) Representative immunoblots

showing abolishment of disulfide crosslinks between MlaA and MlaC in Δ*mlaA* Δ*mlaC* cells expressing

various C-terminal tail helix mutants of MlaA[Q205C](-His) and MlaC[V171C]-His from the pCDF and pET22/42

plasmids, respectively. Samples were subjected to non-reducing (*left*) or reducing (*right*) SDS-PAGE prior to

immunoblotting.

Figure 4. Positively-charged surface residues on MlaC are responsible for mediating MlaA interaction. (A) Cartoon and surface vacuum electrostatics representations of apo-MlaC (PDB 6GKI) (25) revealing positively-charged patches (*blue surface*). Positively-charged residues (or groups of residues, *blue sticks*) are numbered from 0 to 7. (B) Representative immunoblots showing abolishment of disulfide crosslinks between MlaA and MlaC in Δ*mlaA* Δ*mlaC* cells expressing indicated surface charged-reversed mutants (numbered according to (A)) of cysteine-substituted MlaC-His and MlaA-His from pET22/42 and pCDF plasmids, respectively. (C) Representative immunoblots showing restoration of disulfide crosslinks between MlaA and

MlaC in Δ*mlaA* Δ*mlaC* cells expressing charge-reversed MlaC1-His and MlaA^{2DD2R}-His from pET22/42 and

pCDF plasmids, respectively. Samples were subjected to non-reducing (*top*) or reducing (*bottom*) SDS-PAGE

prior to immunoblotting. (D) Cartoon representations of Alphafold2-predicted MlaA (*teal*)-MlaC (*pink*) model,

revealing the proximity of residues D243/D244 on MlaA, and residues K84/R90 on MlaC illustrated in *blue*

and *red* sticks on MlaA and MlaC respectively (inset). Cutaway surface representation of this MlaA-MlaC

model is shown on the right, revealing the juxtaposition of the hydrophobic lipid-binding cavity of MlaC with

the hydrophilic channel of MlaA.

Figure 5. The MlaA hydrophilic channel exhibits solvent-accessibility changes due to interactions with MlaC

and trimeric porins. Extracellular (*left*) and periplasmic (*right*) views of cartoon representations of the reported

crystal structure of OmpK36 (*green*)-*Kp*MlaA (*cyan*) (PDB 5NUP) (32), showing channel residues substituted

with cysteine (for SCAM) in *sticks*. (B, C) Representative immunoblots showing Mal-PEG5k alkylation of

MlaA^{K184C}-His cysteine variant expressed from the pCDF plasmid, either in (B) Δ*mlaA*, Δ*mlaA* Δ*mlaC*, and

Δ*mlaA* Δ*mlaD* backgrounds, or (C) Δ*mlaA* Δ*mlaC* background also producing wildtype or charge-reversed

K84D/R90D (MlaC1) MlaC variants from the pET23/43-*mlaCB* plasmid. (D) Representative immunoblots

showing Mal-PEG5k alkylation of MlaA^{K184C}-His variant also harboring non-functional 3G3P or gain-of-

function *mlaA** mutations expressed from pET23/42 plasmids in Δ*mlaA*, Δ*mlaA* Δ*mlaC*, and Δ*mlaA* Δ*mlaD*

background strains. (E, F) Representative immunoblots showing Mal-PEG5k alkylation of MlaA^{K184C}-His

cysteine variant expressed from the pCDF plasmid in various porin mutant backgrounds. In (E), OmpC or

OmpF were expressed from pDSW206 plasmids where indicated. In SCAM, cells were labelled with

membrane-permeable (NEM) or impermeable (MTSES) reagents, followed by alkylation with Mal-PEG5k,

which introduces a ~5-kDa mass shift to MlaA^{K184C}-His. The levels of solvent accessibility of K184C in MlaA

in the various strains, i.e. fully, partially, or not blocked by MTSES, are highlighted in *blue*, *orange*, or *red*,

respectively.

Figure 6. A proposed model for MlaC recruitment/binding and accompanying MlaA conformational changes that facilitate retrograde PL transfer by the OmpC-MlaA-(MlaC) complex. At the OM, apo-MlaC is preferentially recruited to MlaA via electrostatic interactions between MlaC surface charge patch and MlaA C-terminal tail helix. Subsequent docking of MlaC to the base of the MlaA channel induces a conformational change in MlaA, from an MTSES-<u>inaccessible</u> state (**State A**, "channel-closed", residue K184 represented by *red star*) to an MTSES-<u>accessible</u> state (**State B**, "channel-open", residue K184 represented by *blue star*), facilitating the transfer of PLs from the outer leaflet of the OM into the lipid binding cavity of MlaC. Holo-MlaC then leaves the OmpC-MlaA complex to shuttle the PL ligand to the IM, thus resetting MlaA conformation. Vacuum electrostatic surface representations of MlaC bound with lipids (PDB 5UWA) (23), and in the unliganded form (PDB 6GKI) (25).