G-Trap Assay II: Characterization of blood Leukocyte Functionality differentiates immune activation and immune suppression in bacteremia patient samples

Peter Simons,[1] Laura Shevy,[2] Virginie Bondu,[1] Angela Wandinger-Ness,[1] Stephen Young,[3] and Tione Buranda[1]*

[1]Department of Pathology, University of New Mexico School of Medicine, Albuquerque, NM 87131, USA.

[2]Division of Infectious Diseases, Department of Internal Medicine, School of Medicine, University of

New Mexico, Albuquerque, NM 87131, USA. [3]Tricore Reference Laboratories, 1001 Woodward Place NE

Albuquerque, NM 87102.

Abstract

Sepsis is a severe organ dysfunction syndrome caused by a dysregulation of the immune system's response to infection. Unfortunately, most infection-causing pathogens aren't routinely detectable in real-time to enable targeted and lifesaving treatment. Thus, clinicians frequently have limited data on which to base treatment decisions. A complete blood count with differential is available within 24 h, and positive culture is only available in ~30% of cases. Furthermore, a blood culture, the traditional gold standard for accurate diagnosis of bacteremia, may take up to five days for results, long after a clinical decision for sepsis management is required. Circulating leukocytes can sense chemotactic signals released by bloodborne pathogens or focal infections not in the bloodstream. Our earlier study showed that pathogen and host immune factors released in the bloodstream stimulated GTP binding of Ras homology (Rho) GTPases such as Rac1 in quiescent endothelial and human leukocytes after exposure to blood plasma from infected patients.[1] In this study, we measured Rac1•GTP as a biomarker of immune functionality of peripheral blood monocytes and polymorphonuclear cells extracted from blood samples drawn for diagnostic use in blood culture assays; from 120 non-infected control patients and serial blood test samples from 28 patients with a confirmed diagnosis of bloodstream infection. 18 cases presented with Rac1•GTP elevation of ≥3 fold above that of control samples. Ten patients with normal or below-normal GTPase activity, accompanied by neutrophilia or pancytopenia. We used Principal Component Analysis to differentiate the 2D spatial distribution of infected patients and negative controls. Measuring differential leukocyte functionality in infected and control patients' blood samples with the G-Trap assay may provide an innovative process for a real-time distinction between infection and non-infectious etiologies.

INTRODUCTION

There are many causes of sepsis, and the key to targeted evaluation and management includes obtaining an accurate patient history of present illness, including potential exposures, medical comorbidities, physical exam findings, laboratory testing serologic studies, biomarkers such as procalcitonin, lactate, C-reactive protein (CRP), microbiologic studies, radiographic imaging, etc. [2-4] Blood cultures are obtained when there is clinical concern for bacteremia or fungemia as presenting cause of infection, such as fever, chills, leukocytosis, leukopenia, or an apparent localizing source of infection (e.g., catheters, IV lines, implanted prosthetic material, central lines). However, traditional blood cultures have limitations, including a long delay of 24 to 120 hours for getting test results. Typically, empiric antibiotics are started while awaiting microbiologic culture data; often even before such data is obtained, limiting culture sensitivity.[5, 6] There are no clear guidelines for narrowing antimicrobial therapy once initiated. The use of empiric antibiotics compromises future antimicrobial efficacy due to the inception of antibiotic resistance.[7, 18]

Alternatives to blood culture tests,[7-9] are designed to improve sensitivity and limit the time lag between blood sample collection and results. However, such methods are prone to confounding presence of contaminants, [10] and the reality that nearly 70% of blood culture results are negative.[11, 12] Relatively new approaches that target host immune responses are under current development. [11, 13-15] Sepsis often manifests as a complex immune dysfunction syndrome with a simultaneous presentation of inflammatory and anti-inflammatory responses.[16, 17] Proinflammatory cytokines catalyze the release of antimicrobial nitric oxide in leukocytes. [18, 19] However, the excessive presence of NO in the circulation suppresses leukocyte mobility by antagonizing GTPase activation. [20] Alternatively, bacteremia linked to sepsis can subvert the immune system by modifying the life span, production, and function of white blood cells/leukocytes responsible for the immune system homeostasis [21-23] Thus, an assay that measures leukocyte functionality provides a direct measure of the level of an immune response to infection.

In a preceding study, [1] we used serial plasma samples from infected patients to simulate immune activation of human leukocytes and culture cells after exposure to inflammatory mediators released upon infection. That study showed that pro-/anti- inflammatory cytokines and novel biomarkers such as chemerin[24] induced GTPase activation in test cells. This study hypothesized that Rho GTPase activation in response to infection[25] is required for immune effector function, such as chemotaxis and tissue extravasation of circulating blood leukocytes during the early and late stages of an infection.[26-33] **(Fig. 1A-C).** Immune activation and dysfunction highlight the need for an assay to assess the functionality of white blood cells, which are an essential component of diagnosing the potential of an infection.

In the present study, we tested the GTPase activity of leukocytes isolated from serial clinical diagnostic samples used to confirm bacterial infection from blood culture. Our study showed that GTPase activity in samples from different patients varied in functionality and magnitude. It indicated that the GTPase activity of leukocytes is a promising biomarker of immune cell function and could potentially be used as a clinical decision support tool (**Fig. 1D**).

RESULTS AND DISCUSSION

GTP binding to Rac1 in patient PBMCs is sensitive to environmental allergies, whereas GTP binding to Rac1 in PMNs is not. We tested peripheral blood samples from 120 non-infected subjects to establish a baseline assessment of GTPase activity in uninfected control patients. The samples were collected over 6 months, from August to December 2019. On average, 12 unique patient samples were tested each week. During the August-October period, we noted that the median level of GTPase activity for PBMCs was variable each week, whereas the GTPase activity of PMN was stable (**Fig. 2A&B**). Therefore, we hypothesized that changes in local pollen counts drove T_h2 cell-mediated allergic response[34, 35] which caused the variation in GTP binding to Rac1 in PBMCs. Consistent with our hypothesis, a plot of the pollen counts (Weather.com) in Albuquerque on the days the blood samples were collected showed a strong and significant Pearson correlation ($\rho = 0.78$, $p = 0.005$) between the median GTPase activity of PBMC samples and pollen counts. We, therefore, included this signal variability within the signal range expected of control samples.

Rac1•GTP levels are significantly upregulated in patients presenting a normal immune response to infection and are depressed in patients with apparent immunosuppression. We assayed for Rac1•GTP in serial blood samples drawn from patients with culture-positive bacterial infections from 28 patients. The peak GTPase activity in PBMCs from 18 out of 28 infected patients was ≥ 3-fold higher than the activity (interquartile range) of the control samples. Lymphocyte and monocyte counts in infected patients were lower than in uninfected controls. However, the difference was not significant. Sepsis profoundly affects all the main lymphocyte subpopulations [36, 37] which undergo increased apoptosis; T-regulatory cells are more resistant to sepsis-induced apoptosis, leading to an increased proportion of T-regulatory cells and an immunosuppressive phenotype. Neutrophil (N) counts were significantly elevated relative to controls (**Fig. 2C**). Overall, Peak Rac1•GTP was lower in PMNs than PBMCs regardless of the higher cell counts in the former.

Rac1•GTP measured in samples from 10 of 28 cases was comparable to controls or at the lower end of the mean quartile range, even though the patients' neutrophil counts were significantly elevated compared to controls (**Fig. 2D**). We assessed the average leukocyte functionality for single cells in each sample population by dividing the Rac1•GTP expression of each patient sample by its cell count (**Fig 3**). The difference between infected and control Rac1•GTP/cell data was significant for PBMCs and not in PMN samples. This result suggested an inverse correlation between neutrophilia and functionality. A notable exception was patient #14, presenting with severe neutropenia (dy2 = 0.09, 1.4k/µl; Table S1), yet yielded a robust GTPase response. We found similar trends for patients # 24, 28, and 31 due to lowered cell counts. We plotted the temporal Rac1•GTP data associated with PBMCs and PMNs from representative patient samples (**Fig 4A-B**). Normalization of GTPase activity to single cells enhanced separation of immune activity levels among the patients according to differences in cell counts (**Fig. 4C-D**), also notably highlighting patient #14's depleted but very active neutrophils (**Fig. 4D**).

We next used Principal Component Analysis (PCA) [38] to determine the relationship between the relative expression of different types of significant leukocyte groups: lymphocytes (L), monocytes (M), and neutrophils (N) and their immune function represented by Rac1•GTP measured in PBMCs and PMNs from patients with confirmed bloodstream infection compared to uninfected controls. (**Fig. 6**). The data are presented using Graphpad prism as a distance biplot of Principal component (PC) scores (patient samples) and *PC loadings* vectors (Rac1 (PBMC), Rac1(PMN), N, L, M, and L+M). The *PC loadings* vectors indicate a proportionate concentration of the immunological variables and correlation with serial patient samples. The coordinates of the vector end or patient samples indicate the weight of the immune variable and the relationship (correlation) to patient samples.

The Rac1(PBMC) and Rac1(PMN) vectors represent the level of functionality of the patient's L, M, and N populations. The magnitude of the cosines of the angles between the vectors is related to mutual correlation, where small angles between two variables represent a strong correlation. Conversely, vectors mutually appearing at ~180° are negatively correlated, and those that seem nearly orthogonal (90°) are likely not correlated. In addition, proximity between *PC scores* and specific *PC loadings* vectors indicates a strong correlation. In **Fig. 6A,** Rap1(PBMC) and Rap1(PMN), represented by the group's most extended vectors, present a strong correlation to each other along the PC1 negative axis. Control samples were grouped near the origin of PC1 and distributed on the negative axis overlapping the L vector. M and N vectors had more significant weight in PC2. The distribution of infected patient samples spanned both PC1 and PC2 axes.

When we examined the average GTPase activity of single cells, the Rac1(PMN) vector was redistributed to the negative PC2 on the opposite end of N and M. The coordinate shift by Rac1(PMN)/cell vector opposite to the N vector indicated a negative correlation between the two variables. This negative correlation between neutrophil counts and GTPase functionality highlighted a potential link between excessive neutrophilia and immune suppression linked to bacteremia. [39-41] Thus, patient samples clustering near the apex of the N vector expressed the cohort-peak neutrophil counts of low GTPase activity; e.g., the position of Patient #17 in the PCA shifts from a position of strong correlation to N and robust GTPase functionality in **Fig. 6A** to relatively weak Rac1(PMN)/cell functionality in **Fig. 6B**. Conversely, neutropenic patients (14 and 24) produced the highest Rac1(PMN)/cell levels. This result indicated the alternative immune impairment due to limited cell counts, unrelated to functionality.

The aggregate responses from the infected and control patients' leukocytes show the controls clustered near the origin and strongly correlated to L and L+M. The even clustering of controls between the PC1 and PC2 axis is consistent with a dynamic immune equilibrium[42] of the uninfected patients. However, this equilibrium is notably absent in a vast majority of the infected patients. To visualize the spatial distribution of patients based on immune functionality, we color-coded patient status depending on whether the Rac1(PBMC)/cell and Rac1(PMN)/cell data were both ≥ 3-fold greater than the controls (red, *aa*), *active l*evel Rac1(PBMC) with *inactive* level Rac1(PMN) (pink, *ai*) and *inactive* Rac1(PBMC) with Rac1(PMN) (dark, *ii*). Most notably, the patients with high numbers of neutrophils (clusters next to N) appeared immune suppressed. Also, the immunosuppressed cluster cases were infected mostly by Klebsiella pneumoniae (KP), Streptococcus pneumoniae (SP), Staphylococcus aureus (SA), Pseudomonas aeruginosa (PA),[43] and Group B streptococcus (GBS)[44] are all pathogens known to thrive under conditions of immune compromise.

Summary

Immune dysregulation in sepsis comprises the intersection of pro- and anti-inflammatory biomarkers [2]; thus, assays designed to test immune functionality will likely produce real-time actionable results.[45, 46] Proinflammatory cytokines are frequently cited as bacteremia biomarkers in the literature. [2, 24, 47-52] In our previous study[1] we showed how this interplay between circulating pro-and anti-inflammatory cytokines impacted the magnitude of the correlated GTPase signals. Thus, GTPase activity in patient leukocytes represents an integrated signaling outcome reflecting the composition of circulating cytokines due to infection. In this way, our assay delineated control patient samples, immune-active and immune-suppressed patients.

A differential diagnosis of bacterial infection can be narrowed by the patient's history and clinical examination. Comorbidities such as various forms of cancer, cytotoxic chemotherapy, bone marrow, or tissue transplant are common causes of immunocompromise featuring neutropenia in the early stages, followed by depressed cell-mediated and humoral immunity. [53] Neutropenia increases patient susceptibility to opportunistic polymicrobial bacteremia (*e.g.,* pseudomonas aeroginosa, E. Coli (patient #14), and Klebsiella pneumoniae). Thus while patient #14's neutrophil functionality was robust, the 100/ml neutrophil count elevated the patient's susceptibility to infection based on a depleted cell count rather than functionality.[54] Establishing immune functionality in patients with comorbidities such as asplenic patients[53] lacking the ability to clear encapsulated bacteria are susceptible to opportunistic infections mainly caused by pneumococcus and Haemophilus influenzae.[54] T- cell immuno-suppression due to corticosteroids, cytokine antagonists, and monoclonals that deplete B-cells used to target connective tissue diseases, e.g., lupus erythematosus and rheumatoid arthritis, also promotes susceptibility to infection. Leukocyte functionality G-Trap tests could be an effective tool for effective diagnosis and narrow prescription of antibiotics based on local antibiograms.[54, 55]

There are limitations to this study that we will address in future work. First, our primary objective was to establish a proof-of-concept assay for interrogating the immune functionality of patients as a biomarker for the rapid assessment of presumptive bacteremia. Thus, our sample cohort was based on the availability of residual diagnostic samples with limited access to patient health records.[3, 4] Our pilot study lacked comprehensive IRB approval to access clinical patient variables that we could have used in aggregate with our current data. Thus, we could not characterize the disease's trajectory and provide individualized insight into intervention strategy. Another limitation to be addressed in a future study is to characterize cytokine composition of the plasma, which can be used to cross-check leukocyte functionality. Next, regarding our data quality, some variables have missing data. Specifically, the CBC with differential test was not always deemed necessary at the clinical level of our data set. Thus, such data are left blank. However, we established our present objective of using GTPase activity as a biomarker for delineating uninfected and infected patients.

Materials and Methods

Effector Proteins. GST-effector chimeras used for the studies are as follows: p21 activated kinase protein binding domain (PAK-1 PBD), a Rac1 effector, were obtained from Millipore Sigma. Rhotekin-RBD, a Rho effector protein, was purchased from Cytoskeleton (Denver, CO).

Antibodies. Monoclonal Rac1 antibodies (Cat. # ARC03) were purchased from Cytoskeleton.

Buffers. 2X RIPA buffer: 100 mM Tris titrated with HCl to pH 7.4, 300 mM NaCl, 2 mM EDTA, 2 mM NaF, 2 mM Na$_3$VO$_4$, 2% NP-40, 0.5% sodium deoxycholate, and just before adding to the culture medium, 2 mM PMSF and 2X protease inhibitors. HHB buffer: 7.98 g/L HEPES (Na salt), 6.43 g/L NaCl, 0.75 g/L KCl, 0.095 g/L MgCl$_2$ and 1.802 g/L glucose. HPSMT buffer, an intracellular mimic: 30 mM HEPES, pH 7.4, 140 mM KCl, 12 mM NaCl, 0.8 mM MgCl$_2$, 0.01% Tween-20.

Study design

The UNM HSC Institutional Review Board approved this study (UNM IRB#18-068). The study was divided into two patient populations. Our first goal was to collect residual diagnostic blood samples from patients admitted to UNM for a non-infectious disease cause. Residual peripheral blood test samples were collected from 120 subjects (50% female) presumed to be free of infection. Our second goal was to collect residual serial blood samples from the clinical lab after a positive confirmation of bacterial culture. In this way, serial blood test samples from 36 patients with a confirmed diagnosis of bloodstream infection were collected after standard of care (SOC) testing at TriCore Reference Laboratories (TRL). TRL provided the investigators with available complete blood counts with differential (CBC with diff.) for each subject. We did not include samples from 8 patients for lack of day 0-1 sample, limited availability, or sample loss.

Isolation of Peripheral blood mononuclear cells (PBMCs) and Polymorphonuclear cells (PMNs) diagnostic samples. Whole blood samples were kept at 4 °C. Peripheral blood mononuclear cells (PBMC) consisting a mixture of lymphocytes (T cells, B cells, and NK cells) and monocytes; and polymorphonuclear cells (PMNs) comprising primarily of neutrophils with a small fraction of eosinophils, basophils, and mast cells, were isolated using a Ficoll-based density gradient. The layers containing the PBMC and PMNs were carefully recovered in 800 µl volumes, mixed with 800 µl PBS, centrifuged, and removed the excess solution, leaving the pellet in 25 µl volume.

Functional assessment of PBMCs and PMNs with Rac1•GTP binding assays. The pelleted cells are lysed by adding 25 µl 2X RIPA at 4°C and centrifuged at 4°C in a cold room. 45 µl of cleared lysate was then analyzed for GTPase activity with G-Trap beads, functionalized with PAK-1 effector beads. [56] We used the CBC with diff. data to determine how much Rac1•GTP was produced by single cells.
Statistical Analysis. We used GraphPad Prism 9.3.0 (GraphPad Software, La Jolla, CA) for statistical analysis. Principal component analyses and Pearson correlation coefficients were derived from the study of serial samples for each patient to visualize clustering between Rac-1 (PBMC), Rac1(PMN), lymphocytes (L), monocytes (M), and neutrophils (N).

Acknowledgments: This research was funded by NIH National Center for Research Resources, the National Center for Advancing Translational Sciences: UL1TR001449, and the UNM Department of Pathology.
Conflict of interest: TB, AWN, PS, and VB have awarded (10261084, 10962541) or pending (20210231683) patents related to the G-Trap assay.

Fig. 1A. Bloodstream bacterial infection is detected by host innate immune cells, including neutrophils, monocytes, and dendritic cells, which activate adaptive lymphocyte effector functions. Changes in leukocyte counts due to infection are routinely measured by a complete blood count with differential (CBC with diff). However, various immune response biomarkers have proven too variable to be definitive diagnostically. The G-Trap assay measures levels of Rac1•GTP, among other activated GTPases, in lysates of circulating white blood cells (WBC). Active GTPases serve as sensors for infection due to their roles in triggering WBC to migrate to sites of infection and kill pathogens. **B.** Bloodstream bacterial infection is detected by white blood cells. Active GTPases serve as sensors for infection due to their roles in triggering WBC to migrate to sites of infection and kill pathogens. Changes in white blood cell (WBC) counts due to infection are routinely measured as an indicator of infection. (image adapted from https://together.stjude.org/en-us/care-support/immunity-illness-infection.html) **C.** Separation of whole blood components allows isolation of WBC, which are then analyzed using the G-Trap assay. Image of Blood cells Retrieved April 15, 2022, from (https://www.visiblebody.com/learn/biology/blood-cells/blood-overview) **D.** Rac1•GTP is a sensitive biomarker of pathogen-activated white blood cells. Serial analysis of blood samples of representative patients presenting an active immune response: two patients showed a failure of antibiotic treatment one patient was responsive to treatment. One patient indicated immune paralysis indicated by below normal GTPase activity. **E**. Summary of the G-Trap assay protocol.

Fig. 2. A. Correlated changes in GTPase activity in PBMCs and pollen count on the day of blood sample collection. Each day represents 12 samples. Pearson correlation (ρ = 0.78, p = 0.05). **B.** Overlapping histograms of Rac-1 GTPase activity measured with G-Trap in leukocytes from TriCore diagnostic blood draws of uninfected patients. The fluorescence value for each patient is represented as a Median Chanel fluorescence (MCF) of the G-Trap measurement. The patient population's median values for polymorphonuclear neutrophils (PMN) and peripheral blood mononuclear cells (PBMC) are 1334 and 2220, respectively. **C.** Plot of Rac1•GTP measured in cell lysates of normal infection positive patient leukocytes (+PBMC and +PMN) and controls. The plot of Rac1•GTP measured in cell lysates of infection positive patient leukocytes (+PBMC and +PMN) and controls. **D.** Plot of Rac1•GTP measured in cell lysates of infection positive immunosuppressed patient leukocytes (+PBMC and +PMN) and controls. Levels of Rac1•GTP were compared using unpaired t-test; ** $p< 0.01$, * $p< 0.05$.

Figure 3. Immune functionality of PBMCs and PMNs per unit cell. A. Patient samples presenting GTPase activity above at ≥3 fold threshold above the median value of the control. **B**. Patient samples presenting GTPase activity at <3 fold median value of the control. The arrows indicate robustly active cellular immune responses of severely neutropenic (14 and 24) or lymphopenic (28 and 31) patients. Levels of Rac1 were compared by ordinary one-way ANOVA with Tukey's correction for multiple comparison tests; **** $p< 0.0001$. Please refer to Supplementary Table 1 for raw data reference.

Fig. 4 Representative graphs of Rac1•GTP measured in serial lysates of PBMCs and PMNs from patients presenting GTPase activity ≥3fold above controls. A&B. Plots of Rac1•GTP in measured in PBMC and PMN lysates versus sample-collection day post-admission and before positive culture result. Each patient is identified numerically. Infecting pathogen abbreviations in parentheses and (G+ or G-) refer to Gram staining of the organism: (PM) Pasteurella multocida (G-); (SA) Staphylococcus aureus (G+), (MRSA) methicillin-resistant Staphylococcus aureus (G+); (EC) Escherichia coli (G-). **C&D.** Rac1•GTP normalized to cell count for the same patient group.

Fig. 6. Principal Component Analysis (PCA) reveals different spatial distributions between uninfected control and blood culture-positive patients. **A.** Spatial distribution of infected patients (pink) relative to controls (green data points) The immune variables measured for each patient, Rac1 (PBMC), Rac1(PMN), L, L+M, M, and N, are denoted as blue PC loadings vectors. Rac1 (PBMC) and Rac1(PMN) are measures of immune functionality of the total count of peripheral blood monocytes (PBMC) and polymorphonuclear cells (PMN), respectively. The magnitude of loadings vectors or data point coordinates indicates the weight of the variable and mutual relationships. In the following graph, divided circles around Patients #14 and #17 illustrate the opposite effects of neutropenia and neutrophilia when cell functionality is normalized to single cells. **B.** PCA plot showing Rac1•GTP produced by singles for PBMCs: Rac1 (PBMC)/(L+M) and PMN: Rac1(PMN)/(N). Because of the high N count, Patient 17's position in the PCA plot moves closer to controls, indicating diminished functionality of single cells in patients with neutrophilia. **C.** PCA plot showing GTP bound Rac1 normalized to cell count for PBMCs: Rac1 (PBMC)/(L+M) and PMN: Rac1(PMN)/(N). Plot identifies patients with the Rac1•GTP mediated response of PBMC and PMN (*aa*), active PBMC and inactive PMN (*ai*), and patients with a globally inactive immune response (*ii*). The original Biplot and PC scores figures were produced in GraphPad Prism version 9.3.1.

Supplemental Table S1. Raw Data Table of GTPase Activity and WBC blank spots indicate missing CBC with diff. data The Case ID letters refer to infecting pathogen, and (G±) is gram-positive or negative. (PM) Pasteurella multocida (G-); (SA) Staphylococcus aureus (G+), (MRSA) Methicillin-resistant Staphylococcus aureus (G+); (PA) Pseudomonas aeruginosa (G-); (SP) Streptococcus pyogenes (G+); (KP) Klebsiella pneumoniae (G-); (SL) Staphylococcus lugdunensis (G+); (PA) Pseudomonas aeruginosa (G-); EC Escherichia coli (G-); (GBS) Group B Streptococcus; (S) Salmonella (G-); (SC) Streptococcus constellatus (G+).
* representative controls

Case ID	Rac1(PBMC)	Rac1 (PMN)	WBC	L	N	M
9EC1	11120	13528.99	34			
9EC2	7642.001	12486	7.98			
10SA1	6437.998	3175.	7.8	1.7	5.6	0.5
10SA2	10989	3454	6.86			0.6
11EC&SC-1	12673	2423	5.2			
12GBS1	7708	6115	3.4			0.3
12GBS2	4960	2868	5.7			
12GBS3	12873	3665.001	6.6	1.7	5.6	0.5
12GBS4	1853	1388	6.9	0.6	5.98	0.3
13EC1	5463.998	1488	8	1	6.7	0.4
13EC2	14974	11746	7.9	1.8	6.0	0.1
14EC1	6613.999	3187	0.9	0.7	0.5	0.1
14EC2	19178	7556.997	1	0.8	0.1	0.1
14EC3	14801.99	3349.999	1.8	0.5	1.4	0.1
15SA1	16317	16887	6.2	1	4.6	0.5
15SA2	12993	21886.01	5.2	1	3.4	0.6
15SA3	25700	2704	3.7	0.9	2.1	0.4
16PM1	50909.01	9788	19.4	0.5	17.1	0.9
16PM2	44474.96	12352	18.5	0.5	17.1	0.9
16PM3	60164.97	13008	17.4	0.5	17.1	1.4
17MRSA1	30180	9249	17.9	1.9	14.1	1.4
17MRSA2	37360	9815	18	2.1	15.1	1.5
6EA1	12682	5249	9.6			
6EA2	9827	5026	11.6			
7EC1	7270	1903	16.2	2.1	15.1	1.5
7EC2	2682	1504	13.4	0.7	11.9	1.3
7EC3	2291	1031	10.2	0.7	11.9	1.3
7EC4	1929	2567	9.9	0.7	10.9	1.5
8EC1	11241	2165	10.7	0.7	9	1
8EC2	11146	3679	7.4	0.8	5.5	0.5
8EC3	5030	5115	4.4			
21PA1	881	801	17.4	0.2	17.4	0.5
21PA2	1062	872	10.8	0.3	9.5	1
21PA3	1006	854	8.1	0.4	7.4	0.2
23KP1	1491	1169	7.1	0.9	5.6	0.6
23KP2	1270	1028	9.7	0.7	8.4	0.4
23KP3	1257	988	7.1	0.9	5.6	0.6
24KP1*	1306	1101	0.2	0.2	0.1	0.1
24KP2*	790.	831	0.4	0.3	0.1	
22SL1	1062	6218	10.2	0.6	8.9	0.6
22SL-2	1143	983				
22SL-3	940	2581				
22SL-4	955	1000				
25SA1	1025	1867	47.7	1	44.8	1.4
25SA2	8486	3084	40.3	0.8	37.1	2.4
25SA3	1976	1379	33.8	1.7	30.8	1.4
25SA4	3839	1145	32	1	30.1	1
25SA5	2338	1487	28.2	1.4	25.4	1.2
26SP1	4716	1652	8.1	0.7	6.7	0.5
26SP2	1588	4145	5.1	1	3.6	0.4
26SP3	1547	1399	4.6	0.8	3.4	0.3
27SP1	11295	12714	28	0.8	25.5	1.4
27SP2	6320	3691	29.5	4.4	24.5	0.6

Case ID	Rac1(PBMC)	Rac1 (PMN)	WBC	L	N	M
27SP3	6733	3463.	15.7	2.4	12.9	0.3
28EC1	4532	1105	4.1	0.1	3.6	0.1
28EC2	3406	3576	15.4			
28EC3	2496	4609	10.3			
28EC4	7905	2392	5.6			

Case ID	Rac1(PBMC)	Rac1 (PMN)	WBC	L	N	M
28EC5	3173.	2313	4.2			
28EC6	3904.	1501				
29SP1	2880	2230	18.9	0.6	18.1	0.2
29SP3	3362.	1882	11.6	0.7	10.1	0.2
29SP4	1918	1245				
29SP5	5945.	1569	8.1	1.3	5.8	0.5
30SA1	9476.	2934	10.7	0.5	9.2	1.1
30SA2	8895	3594	11.6	0.9	9.2	1.5
31SA1	3002.	6002.	0.1	0.1		
31EC2	4281.	3009.				
31EC3	2568	8990.				
31EC4	1622	3856.	0.19			
31EC5	4102.	1017				
31EC6	1885	10006	0.1			
32SA1	4216.	1203	18.6	0.8	17.6	0.2
32SA2	1833	2366	24.3	0.3	23.3	0.7
32SA3	2346	1887	14.3	0.2	13.9	0.2
32SA4	2415	902	10.2	0.6	8.8	0.2
32SA5	2154	997.				
33SA	4670.	1780	8.6	1.4	6.2	0.9
34SP1	10787	5908.	15.3	0.5	13	0.6
34SP2	7977.	5881.	14.1			
34SP3	10038	4835.	12.6			
34SP4	8227.	5179	11.7			
34SP5	9071	6561.	10.8			
35EC1	4974.	5927	11.2	2	7.8	1.4
35EC3	4728.	3412	8			
35EC4	3372	2518	13.2			
36SP1	3164	1236	26.9	0.5	25.6	0.5
36SP2	5972.	2133	11.9			
36SP3	4261.	1779				
36SP4	3329.	2267				
*	2459	3055	7.2	2.8	3.4	0.7
*	3750	3297	6	1	4.1	0.7
*	3669	3727	6.9	1.9	6.1	0.8
*	2473	2159	4.7	2.2	2.3	0.7
*	2203	2073	6.1	2.4	3	0.5
*	7999	2373	5.3	2.1	2.5	0.4
*	2290	3077	5.9	1.2	4	0.6
*	3160	3440	5.8	1.8	3.4	0.4
*	5272	3762	7	2.7	3.7	0.5
*	2496	1234	6.8	2.2	3.9	0.5
*	5846	1431	6.3	1.4	4.2	0.3
*	2293	1353	4.9	1.6	2.7	0.4
*	2023	1241	6.8	2.5	3.3	0.8
*	1660	1145	4.4	1.5	2.4	0.4
*	2570	1518	5	1.2	3.1	0.5
*	2731	1214	6.4	2.1	3.4	0.6
*	1927	964	5.2	2.5	2.1	0.5
*	3074	1395	5.7	2.4	2.9	0.3
*	2295	1536	4.8	1.4	2.9	0.4
*	2315	1391	6.1	2	3.6	0.4
*	1804	1071	5.3	1.7	3.0	0.5

References

1. Clark, R. *et al.* G-Trap Assay I: Small GTPases as sensitive immune response biomarkers for active bacteremia. (2022).
2. Taneja, I. *et al.* Combining Biomarkers with EMR Data to Identify Patients in Different Phases of Sepsis. *Scientific reports* **7**, 10800 (2017).
3. Vincent, J.L. *et al.* Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *The Lancet. Respiratory medicine* **2**, 380-386 (2014).
4. Cohen, J. *et al.* Sepsis: a roadmap for future research. *The Lancet. Infectious diseases* **15**, 581-614 (2015).
5. Coburn, B., Morris, A.M., Tomlinson, G. & Detsky, A.S. Does this adult patient with suspected bacteremia require blood cultures? *JAMA* **308**, 502-511 (2012).
6. Fay, K., Sapiano, M., Gokhale, R. & al., e. Assessment of Health Care Exposures and Outcomes in Adult Patients With Sepsis and Septic Shock. . *JAMA Netw Open* **3**, e206004 (2020).
7. Stevenson, M. *et al.* Sepsis: the LightCycler SeptiFast Test MGRADE(R), SepsiTest and IRIDICA BAC BSI assay for rapidly identifying bloodstream bacteria and fungi - a systematic review and economic evaluation. *Health Technol Assess* **20**, 1-246 (2016).
8. Carroll, K.C. *et al.* Clinical Performance of the Novel GenMark Dx ePlex Blood Culture ID Gram-Positive Panel. *Journal of clinical microbiology* **58** (2020).
9. Wolk, D.M. *et al.* A Multicenter Clinical Study To Demonstrate the Diagnostic Accuracy of the GenMark Dx ePlex Blood Culture Identification Gram-Negative Panel. *Journal of clinical microbiology* **59**, e0248420 (2021).
10. d'Humieres, C. *et al.* Two original observations concerning bacterial infections in COVID-19 patients hospitalized in intensive care units during the first wave of the epidemic in France. *PloS one* **16**, e0250728 (2021).
11. Sweeney, T.E., Liesenfeld, O. & May, L. Diagnosis of bacterial sepsis: why are tests for bacteremia not sufficient? *Expert Rev Mol Diagn* **19**, 959-962 (2019).
12. Mi, M.Y., Klompas, M. & Evans, L. Early Administration of Antibiotics for Suspected Sepsis. *The New England journal of medicine* **380**, 593-596 (2019).
13. McHugh, L. *et al.* A Molecular Host Response Assay to Discriminate Between Sepsis and Infection-Negative Systemic Inflammation in Critically Ill Patients: Discovery and Validation in Independent Cohorts. *PLoS Med* **12**, e1001916 (2015).
14. Ducharme, J. *et al.* A Multi-mRNA Host-Response Molecular Blood Test for the Diagnosis and Prognosis of Acute Infections and Sepsis: Proceedings from a Clinical Advisory Panel. *J Pers Med* **10** (2020).
15. He, Y.D. *et al.* The Optimization and Biological Significance of a 29-Host-Immune-mRNA Panel for the Diagnosis of Acute Infections and Sepsis. *J Pers Med* **11** (2021).
16. Shankar-Hari, M. *et al.* Early PREdiction of sepsis using leukocyte surface biomarkers: the ExPRES-sepsis cohort study. *Intensive care medicine* **44**, 1836-1848 (2018).
17. Pene, F., Pickkers, P. & Hotchkiss, R.S. Is this critically ill patient immunocompromised? *Intensive care medicine* **42**, 1051-1054 (2016).
18. Cunha, F.Q. *et al.* Differential induction of nitric oxide synthase in various organs of the mouse during endotoxaemia: role of TNF-alpha and IL-1-beta. *Immunology* **81**, 211-215 (1994).
19. Bone, R.C., Grodzin, C.J. & Balk, R.A. Sepsis: a new hypothesis for pathogenesis of the disease process. *Chest* **112**, 235-243 (1997).
20. Lee, J.S. *et al.* Mechanisms of nitric oxide interplay with Rho GTPase family members in modulation of actin membrane dynamics in pericytes and fibroblasts. *The American journal of pathology* **166**, 1861-1870 (2005).
21. Bosmann, M. & Ward, P.A. The inflammatory response in sepsis. *Trends. Immunol.* **34**, 129-136 (2012).

22. Flierl, M.A. *et al.* Functions of the complement components C3 and C5 during sepsis. *Faseb Journal* **22**, 3483-3490 (2008).
23. Rittirsch, D., Flierl, M.A. & Ward, P.A. Harmful molecular mechanisms in sepsis. *Nat Rev Immunol* **8**, 776-787 (2008).
24. Karampela, I. *et al.* Circulating Chemerin and Its Kinetics May Be a Useful Diagnostic and Prognostic Biomarker in Critically Ill Patients with Sepsis: A Prospective Study. *Biomolecules* **12** (2022).
25. Lemichez, E. & Aktories, K. Hijacking of Rho GTPases during bacterial infection. *Experimental cell research* **319**, 2329-2336 (2013).
26. Worbs, T., Hammerschmidt, S.I. & Forster, R. Dendritic cell migration in health and disease. *Nat Rev Immunol* **17**, 30-48 (2017).
27. Randolph, G.J., Angeli, V. & Swartz, M.A. Dendritic-cell trafficking to lymph nodes through lymphatic vessels. *Nat Rev Immunol* **5**, 617-628 (2005).
28. Ring, S. *et al.* Regulatory T cell-derived adenosine induces dendritic cell migration through the Epac-Rap1 pathway. *J Immunol* **194**, 3735-3744 (2015).
29. Lagarrigue, F., Kim, C. & Ginsberg, M.H. The Rap1-RIAM-talin axis of integrin activation and blood cell function. *Blood* **128**, 479-487 (2016).
30. Alves-Filho, J.C., Spiller, F. & Cunha, F.Q. Neutrophil paralysis in sepsis. *Shock* **34 Suppl 1**, 15-21 (2010).
31. Ellett, F. *et al.* Diagnosis of sepsis from a drop of blood by measurement of spontaneous neutrophil motility in a microfluidic assay. *Nature biomedical engineering* **2**, 207-214 (2018).
32. Granucci, F. *et al.* The immune response is initiated by dendritic cells via interaction with microorganisms and interleukin-2 production. *The Journal of infectious diseases* **187 Suppl 2**, S346-350 (2003).
33. Paysant, J. *et al.* Long-term incubation with IL-4 and IL-10 oppositely modifies procoagulant activity of monocytes and modulates the surface expression of tissue factor and tissue factor pathway inhibitor. *British journal of haematology* **131**, 356-365 (2005).
34. Arango Duque, G. & Descoteaux, A. Macrophage cytokines: involvement in immunity and infectious diseases. *Frontiers in immunology* **5**, 491 (2014).
35. Panda, S.K. & Colonna, M. Innate Lymphoid Cells in Mucosal Immunity. *Frontiers in immunology* **10**, 861 (2019).
36. Hotchkiss, R.S., Monneret, G. & Payen, D. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol* **13**, 862-874 (2013).
37. Unsinger, J. *et al.* Sepsis-induced apoptosis leads to active suppression of delayed-type hypersensitivity by CD8+ regulatory T cells through a TRAIL-dependent mechanism. *J Immunol* **184**, 6766-6772 (2010).
38. Simons, P. *et al.* Longitudinal Assessment of Cytokine Expression and Plasminogen Activation in Hantavirus Cardiopulmonary Syndrome Reveals Immune Regulatory Dysfunction in End-Stage Disease. *Viruses* **13** (2021).
39. Fabrice Uhel, F. *et al.* Early Expansion of Circulating Granulocytic Myeloid-derived Suppressor Cells Predicts Development of Nosocomial Infections in Patients with Sepsis. *Am J Respir Crit Care Med* **196**, 315-327 (2017).
40. Gabrilovich, D., Nagaraj, S. Myeloid-derived suppressor cells as regulators of the immune system. . *Nat Rev Immunol* **9**, 162–174 (2009).
41. Venet, F. *et al.* Regulatory T cell populations in sepsis and trauma. . *J Leukoc Biol.* **83**, 523-535 (2008).
42. Eberl, G. Immunity by equilibrium. *Nat Rev Immunol* **16**, 524-532 (2016).
43. Koch, R.M. *et al.* Patterns in Bacterial- and Viral-Induced Immunosuppression and Secondary Infections in the ICU. *Shock* **47**, 5-12 (2017).
44. Kline, K.A., Schwartz, D.J., Lewis, W.G., Hultgren, S.J. & Lewis, A.L. Immune activation and suppression by group B streptococcus in a murine model of urinary tract infection. *Infection and immunity* **79**, 3588-3595 (2011).
45. Albert-Vega, C. *et al.* Immune Functional Assays, From Custom to Standardized Tests for Precision Medicine. *Frontiers in immunology* **9**, 2367 (2018).

46. Salyer, C.E., Bergmann, C.B., Hotchkiss, R.S., Crisologo, P.A. & Caldwell, C.C. Functional Characterization of Neutrophils Allows Source Control Evaluation in a Murine Sepsis Model. *J Surg Res* **274**, 94-101 (2022).
47. Ge, Y., Huang, M. & Yao, Y.M. Biology of Interleukin-17 and Its Pathophysiological Significance in Sepsis. *Frontiers in immunology* **11**, 1558 (2020).
48. Flores-Mejia, L.A. *et al.* Function is Dissociated From Activation-Related Immunophenotype on Phagocytes From Patients With SIRS/Sepsis Syndrome. *Shock* **52**, e68-e75 (2019).
49. Matsumoto, H. *et al.* The clinical importance of a cytokine network in the acute phase of sepsis. *Scientific reports* **8**, 13995 (2018).
50. Hong, T.H. *et al.* Biomarkers of early sepsis may be correlated with outcome. *J Transl Med* **12** (2014).
51. Faix, J.D. Biomarkers of sepsis. *Critical reviews in clinical laboratory sciences* **50**, 23-36 (2013).
52. Lvovschi, V. *et al.* Cytokine profiles in sepsis have limited relevance for stratifying patients in the emergency department: a prospective observational study. *PloS one* **6**, e28870 (2011).
53. Di Sabatino, A., Carsetti, R. & Corazza, G.R. Post-splenectomy and hyposplenic states. *Lancet* **378**, 86-97 (2011).
54. Southwick, F.S. Chapter 15: Infections in the Immunocompromised Host. *Infectious Diseases: A Clinical Short Course 4 th Edition*, 409-420 (2020).
55. Southwick, F.S. Chapter 2: Sepsis. *Infectious Diseases: A Clinical Short Course 4 th Edition*, 63-73 (2020).
56. Simons, P.C., Bondu, V., Wandinger-Ness, A. & Buranda, T. Small volume Flow Cytometry-Based Multiplex Analysis of the Activity of Small GTPases *Rho GTPases: Methods and Protocols, Second Edition* (2018).