Genetic exchange in *Leishmania* is facilitated by IgM natural antibodies

Tiago D. Serafim[1#*], Eva Iniguez[1#], Ana Beatriz F. Barletta [2], Johannes S.P. Doehl[1], Mara Short[1], Justin Lack[3], Pedro Cecilio[4] Vinod Nair[5], Maria Disotuar[1], Timothy Wilson[1], Iliano V. Coutinho-Abreu[1], Fabiano Oliveira[1], Claudio Meneses[1], Carolina Barillas-Mury[2], John Andersen[5], José M.C. Ribeiro[5], Stephen M. Beverley[6*], Shaden Kamhawi[1*], Jesus G. Valenzuela[1*].

Affiliations:

1. Vector Molecular Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA.

2. Mosquito Immunity and Vector Competence Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA.

3. NIAID Collaborative Bioinformatics Resource, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA.

4. Vector Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA.

5. Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT, USA.

6. Department of Molecular Microbiology, School of Medicine, Washington University, St. Louis, MO, USA.

*Correspondence to:
jvalenzuela@niaid.nih.gov;stephen.beverley@wustl.edu;skamhawi@niaid.nih.gov;tiago.donatelliserafim@nih.gov

#Equally contributed

Abstract

        Host factors mediating *Leishmania* genetic exchange are not well defined. Here, we demonstrate that IgM antibodies, but not IgG or IgA, facilitate parasite genetic hybridization in vitro and in vivo. IgM induces the gradual and transient formation of a structured parasite clump in a process essential for *L. major* and *L. tropica* hybridization in vitro. Parasite hybrids and 3-nucleated parasites were observed inside this structure, named the *Leishmania* mating clump. IgM was also required for or significantly increased *Leishmania* hybrid formation in vivo. At minimum, we observed a 12-fold increase in the proportion of hybrids recovered from sand flies provided a second blood meal containing IgM compared to controls. Notably, genetic backcross events in sand flies were only observed in the presence of IgM, and were reproducibly recovered, reinforcing the relevance of IgM for *Leishmania* genetic exchange in vivo. The in vitro and in vivo *Leishmania* crosses from these studies resulted in full genome hybrids. *Leishmania* co-option of a host antibody to facilitate mating in the insect vector establishes a new paradigm of parasite-host-vector coevolution that promotes parasite diversity and fitness through genetic exchange.

Main text

Leishmania parasites are vector-borne pathogens transmitted to mammalian hosts by phlebotomine sand flies[1]. Associated with poverty, leishmaniasis is prevalent in developing countries and results in self-limiting, severely disfiguring, or fatal diseases that cause substantial morbidity and mortality worldwide[2]. More than a billion people are at risk of infection, millions are asymptomatic carriers, and the annual incidence of fatal visceral and cutaneous leishmaniases are estimated at 100,000 to 1 million cases, respectively[3]. These statistics place leishmaniasis at the forefront of parasitic diseases of public health importance.

Leishmania parasites are considered mostly asexual, reproducing via clonal propagation[4-6], though the presence of natural hybrids and demonstration of genetic exchange in the sand fly vector[7-10] indicate that genetic exchange is more likely 'facultative' as seen in other eukaryotic microbes[11]. Genetic exchange is a powerful evolutionary route enabling the development of pathogenicity in microbes and offers an experimental tool for dissecting complex phenotypes underlying virulence[12].

Host and parasite factors mediating this process are not well understood. In earlier studies, passage in sand flies was required to undertake this experimentally[10,13,14], but recently, hybrid formation was observed in culture for a few species including L. tropica, albeit following exposure to harsh conditions[10,15,16]. Nevertheless, in vitro hybrid generation is yet to be accomplished for most Leishmania species, including L. major[10,15,16]. Importantly, genetic exchange or the generation of fertile hybrids through backcrosses has only been reported once[8].

In this study, we focused on elucidating host factors that mediate Leishmania genetic exchange, asking whether host blood ingested by sand flies provides some element(s) absent in culture media that promotes fusion and the formation of hybrid parasites. By adding freshly

isolated dog plasma to culture media we succeeded in generating in vitro crosses of two genetically labeled *L. major* strains, WR-SSU-HYG and FV1-FKP40-BSD (Fig. 1a). Hybrid generation was only observed when accompanied by the formation of a structured spherical 'clump' composed of live aggregating parasites (Fig. 1b, top panel; Supplementary Videos 1-3). In standard culture media, in the absence of dog plasma, the parasite clump did not form (Fig. 1b, bottom panel; Supplementary Videos 6,7) and hybrids were not recovered. The parasite clump forms gradually and is maintained for 24-36h (Supplementary Video 4) before it starts to dissociate releasing viable parasites.

Purification and fractionation of dog plasma using a 2 step-HPLC followed by MS-MS and in vitro incubation of parasites with the HPLC fractions revealed that IgM was the molecule responsible for the formation of the parasite clump (Supplementary Fig. 1). Parasite clumping was retained in HPLC-eluted fractions of a commercially available IgM (Fig. 1c) but was absent with a commercial human IgA or IgG (Fig. 1d), emphasizing the unique role of IgM in parasite clumping. Importantly, fetal bovine serum (FBS) does not induce clumping (Fig. 1d) and it does not contain IgM (Fig. 1e). Pertinently, the process of IgM-induced clump formation (Fig. 1f, Supplementary Fig. 2a) is distinct from other aggregation phenomena such as with peanut agglutinin (PNA), where the parasites agglutinate immediately and irreversibly, forming a disorganized flat mesh-like structure of immotile parasites (Fig. 1g; Supplementary Fig. 2b; Supplementary Video 5). The composition and temporal formation and dissociation of the *Leishmania* parasite clump is also different from other reported structures, including the *Leishmania* rosettes[17]. Furthermore, pentameric IgM was required (Fig. 1h; Supplementary Fig. 3), as IgM monomers failed to clump the parasites (Fig. 1i, Supplementary Fig. 3). Of relevance,

IgM-induced parasite clumping in eight diverse *Leishmania* species (Fig. 1j) strongly suggesting

that parasite clumping upon contact with blood is a general response in *Leishmania*.

To directly establish the contribution of the IgM-induced clump to genetic exchange in

vitro, we used 3 genetically marked *L. major* parasite strains with independent selectable markers

(WR-SSU-HYG [parental line 1], FV1-FKP40-BSD [parental line 2], and FV1-FTL-SAT

[parental line 3]) in various crosses. Stationary phase promastigotes from two parental lines were

mixed at a 1:1 ratio and cultured for 3 days, with or without IgM, in the absence of selection drugs,

and then grown in the presence of selective drugs for up to 30 days (Fig. 1k, l). Hybrids were

observed for all parental combinations, but only when the culture was supplemented with IgM

(Fig. 1k, l).

Collectively, our data demonstrate that IgM is critical for the generation of *L. major* hybrids

in vitro through its induction of a structured and transient living parasite clump. Furthermore,

without any other manipulation to the culture, in vitro hybrids from *L. tropica* crosses were only

observed in the presence of IgM under our experimental conditions (Supplementary Fig. 4).

Another breakthrough of this study was to have accomplished something sought after for many

years, the formation of *L. major* hybrids in vitro.

Our studies addressed the hypothesis that formation of the IgM-dependent parasite clump

promotes a closer contact between parasites, and this, in turn, increases the probability of

Leishmania hybrid formation. Using confocal microscopy, we observed that IgM was abundantly

distributed throughout the parasite clump in close association with *Leishmania* parasites, forming

the distinct spherical structure observed in vitro (Fig. 2a). Transversal sections provided further

evidence that *Leishmania* parasites lie in close proximity to each other within the parasite clump

and revealed that IgM localized to distinct pockets between parasites (Fig. 2b) and coated their

surface (Fig. 2c), potentially contributing to interactions holding the structure together. We then

used scanning electron microscopy to establish the similarity in the organization of closely

interacting parasites within the *L. major* (Fig. 2d, upper panel) and *L. tropica* (Fig. 2d, lower panel)

clumps. Importantly, we located hybrid events inside a *L. tropica* parasite clump using two parental

lines each harboring either a green or red nucleotide analog, only detected in the nucleus (Fig. 2e,

left and middle panels). Double labeled nuclei (yellow), signifying a hybridization event, or a

Leishmania hybrid, were only detected inside the parasite clump by confocal microscopy (Fig. 2e,

right panel). Promastigotes with 3 nuclei (Fig. 2f, red arrows) indicative of fusion events were also

observed inside the parasite clump by transmission electron microscopy. Our data provide direct

evidence that IgM permeates the parasite clump and coats the parasites and establish that

Leishmania fusion and hybrid formation take place within the parasite clump. Based on these

findings and their relevance in parasite fusion and hybrid formation, we termed this structure the

Leishmania mating clump (LMC).

Eukaryotic class II fusogens such as HAP2, intrinsic molecules identified in

Chlamydomonas and *Plasmodium*, are involved in membrane fusion between gametes. In contrast,

IgM is an extrinsic molecule originating from the mammalian host that brings *Leishmania* parasites

together to promote fusion, exposing a novel mechanism co-opted by a eukaryote to facilitate the

early key steps of genetic exchange. We hypothesize that the LMC creates an environment,

nutritional, physical, and biochemical that promotes parasite fusion and hybrid formation. It will

be important to further investigate whether the interaction between IgM and the parasite is

activating a signaling pathway, or whether the tight parasite scaffolding within the LMC is creating

a physical or nutritional stress that facilitates genetic exchange. Nevertheless, it is important to

emphasize that the addition of IgM to *Leishmania* culture media mimics natural exposure of

Leishmania to IgM in sand fly blood meals. Every time a sand fly takes a blood meal, a *Leishmania* parasite that resides in the sand fly gut will interact with IgM-present in the blood of all vertebrate animals[18-20].

The ubiquity of IgM in vertebrate blood[18-21] combined with the multiple and promiscuous (Supplementary Table 1) blood-feeding behavior of the sand fly[22] led us to hypothesize that this antibody is a major facilitator for the initiation of genetic exchange in vivo, in the sand fly gut. Experiments were performed using conditions that mimicked the natural progression of sand fly infections, including incorporation of a second blood meal that is needed to prevent loss of naturally acquired parasites[22]. Having first established that IgM persists in the sand fly midgut for at least 24 hours after each blood meal (Fig. 3a), we performed an experiment where we allowed sand flies to feed on a cutaneous leishmaniasis lesion initiated by two *L. major* parental lines (Fig. 3b, BM1), to pick the parasites up as part of a natural blood meal [23,24]. We then followed the natural sand fly feeding behavior by offering them a second uninfected blood meal six days after feeding on cutaneous lesions (Fig. 3b, BM2). The second blood meal was either devoid of IgM or supplemented with 500 µg/ml of IgM (Fig. 3b, BM2), a concentration at or below those reported in the blood of different animals (Supplementary Table 2). Additionally, a control group was maintained on a single infected blood meal (Fig. 3b, dotted arrow). Fourteen days after the initial infection, sand fly guts from each group were dissected and plated individually to screen for parasite hybrids (Fig. 3b). For the group initially infected with parental lines combination 1x2, IgM increased the proportion of hybrid formation by up to 12-fold in sand flies getting a second blood meal containing IgM compared to sand flies given a second blood meal devoid of IgM or sand flies given a single infectious blood meal (Fig. 3c, d, Supplementary Fig. 5a). In these two control groups, we observed a significantly lower proportion of hybrids as was previously

reported[13,14]. The similar proportion of hybrids observed in flies given a second blood meal devoid

of IgM or sand flies given a single infectious blood meal suggest that this may be a basal level of

chance mating within the confined space of the sand fly midgut. It is worth noting that both control

group scenarios would not occur in nature since an infected sand fly will always take a second

blood meal 5-6 days after infection[1,22]. Strikingly, control groups lacking IgM did not generate any

hybrids in parental lines combinations 1x3 and 2x3, further emphasizing the importance of IgM in

Leishmania hybrid formation in vivo (Fig. 3c, d, Supplementary Fig. 5a). Of note, the parasite

load, and percent metacyclics were similar in all experimental groups (Supplementary Fig. 5b-g).

The pronounced effect of IgM on parasite hybrid formation when provided to sand flies six days

after naturally acquired infections (BM2) suggests that proliferative stages are prone to genetic

exchange. We argue that hybrids may form soon after a sand fly takes a blood meal but can only

be detected later in the infection cycle after undergoing multiple cycles of division[1,23]. Collectively,

the above data establish that IgM facilitates *Leishmania* hybrid formation inside the sand fly gut

after naturally acquired infections, likely by bringing parasites closer together within a LMC that

promotes fusion and hybrid generation. This is supported by the observation of structures similar

to in vitro LMCs within the midgut (Supplementary Fig. 6; Supplementary Videos 7-12).

Corroborating previous findings[10], we observed hybrid formation only in sand flies and did

not find them in *Leishmania* skin lesions or draining lymph nodes containing 2 different parasite

lines (Supplementary Fig. 7). In the mammalian host, the *Leishmania* parasite is intracellular,

residing in a parasitophorous vacuole inside macrophages[25,26]. It may be that a lack of opportunity

for IgM to interact with amastigotes accounts for the absence of hybridization in the mammalian

host. Whole-genome sequencing of the double drug resistant hybrids (Supplementary Fig. 8)

showed that the crosses in sand flies and those generated in vitro were full genome hybrids (Fig.

3e; Supplementary Fig. 8-10). Moreover, aneuploidy commonly seen in cultured parasites[10,13,14]

were observed in this study at comparable levels (Supplementary Fig. 9, 10).

For use in positional cloning, generation of recombinant progeny by mating of hybrids to

each other or to parental lines is required. Previous studies showed that the frequency of this

progeny in *L. major* was considerably lower than it was for initial F1 hybrid formation, greatly

hindering its utility[8]. To test whether insights from IgM-mediated mating could improve F1 hybrid

recovery as well as confirm their fertility, we performed a backcross in the presence of IgM. This

involved crossing a HYG/BSD double resistant (1x2) F1 hybrid with two clones of *L. major*

parental line 2 bearing a SAT marker to allow identification of mating progeny.   Sand flies were

allowed to feed on a cutaneous lesion initiated by mixed infections of the (1x2) F1 hybrid and

parental line 2' or 2" of *L. major* (Fig. 4a, BM1), and then given a second uninfected blood meal

six days later (BM2), either supplemented with 500 µg/ml of IgM or devoid of IgM (Fig. 4a, BM2).

We consistently observed hybrid formation of this backcross, but only in sand flies fed with blood

containing IgM (Fig. 4b, c; Supplementary Fig. 11a). Hybrid generation was not influenced by

parasite density or percent metacyclics that were similar in all experimental groups

(Supplementary Fig. 11b-g). A diagram outlining the approach to the generation of backcrosses is

provided in Supplementary Fig. 12.

Whole-genome sequencing of the triple drug resistant hybrids showed that the second-

generation backcross progeny in sand flies bore the expected patterns of recombination between

the F1, and 3rd parent genomes also observed in previously reported backcrosses[8]. Importantly,

no 'backcross progeny' were obtained with sand flies given a second blood meal devoid of IgM,

providing conclusive evidence of its importance in *Leishmania* genetic exchange in a natural

setting. Our data strongly suggest that IgM facilitates fertile hybrid formation and enhances their

frequency in sand flies consequently promoting *Leishmania* diversity. Importantly, the reproducible and increased recovery of backcrosses opens the way for more in-depth genetic analysis of *Leishmania* parasites.

Historically, aggregation of *L. donovani* by IgM in culture was reported once as a standalone observation[27]. Our work provides direct experimental evidence of a critical role for IgM in LMC formation that promotes the first key steps of *Leishmania* genetic exchange in vitro and in vivo after a naturally acquired infection. Moreover, our data describe a new biological function for an antibody by implicating IgM in genetic exchange in a parasite. The exploitation of host IgM by *Leishmania* is of significant consequence. IgM is an evolutionarily conserved natural antibody that is always present in the blood of all vertebrates after birth[19-21], independent of the immune status of the animal[28-33]. *Leishmania* evolved to exploit the pentameric properties of host IgM and the biological feeding pattern of sand flies to enhance the likelihood of mating. Sand flies ingest blood every 5-6 days to produce eggs[1]. We hypothesize that when a *Leishmania*-infected sand fly takes a subsequent blood meal, the periodic influx of IgM encounters parasites that have evolved to be at a stage that provides the density required to initiate the formation of the LMC in the confinement of the gut, a critical first step of genetic exchange that promotes fusion and hybrid generation. As such, in addition to metacyclic promastigotes, this implicates the two major proliferative stages, leptomonad and retroleptomonad promastigotes, as potential participants in the process of genetic exchange.

The fact that the *Leishmania* parasite co-opted a natural antibody from its mammalian host to advance its reproductive fitness uncovers a critical step in genetic exchange, providing an insight into the intimate evolutionary relationship between an insect vector, a parasite, and a mammalian host.

Main References

1 Serafim, T. D. *et al.* Leishmaniasis: the act of transmission. *Trends Parasitol* **37**, 976-
987, doi:10.1016/j.pt.2021.07.003 (2021).
2 Burza, S., Croft, S. L. & Boelaert, M. Leishmaniasis. *Lancet* **392**, 951-970,
doi:10.1016/S0140-6736(18)31204-2 (2018).
3 Alvar, J. *et al.* Leishmaniasis worldwide and global estimates of its incidence. *PloS one*
7, e35671, doi:10.1371/journal.pone.0035671 (2012).
4 Tibayrenc, M. & Ayala, F. J. Is Predominant Clonal Evolution a Common Evolutionary
Adaptation to Parasitism in Pathogenic Parasitic Protozoa, Fungi, Bacteria, and Viruses?
Adv Parasitol **97**, 243-325, doi:10.1016/bs.apar.2016.08.007 (2017).
5 Tibayrenc, M. & Ayala, F. J. The clonal theory of parasitic protozoa: 12 years on. *Trends*
Parasitol **18**, 405-410, doi:10.1016/s1471-4922(02)02357-7 (2002).
6 Tibayrenc, M., Kjellberg, F. & Ayala, F. J. A clonal theory of parasitic protozoa: the
population structures of Entamoeba, Giardia, Leishmania, Naegleria, Plasmodium,
Trichomonas, and Trypanosoma and their medical and taxonomical consequences. *Proc*
Natl Acad Sci U S A **87**, 2414-2418, doi:10.1073/pnas.87.7.2414 (1990).
7 Rogers, M. B. *et al.* Genomic confirmation of hybridisation and recent inbreeding in a
vector-isolated Leishmania population. *PLoS Genet* **10**, e1004092,
doi:10.1371/journal.pgen.1004092 (2014).
8 Inbar, E. *et al.* Whole genome sequencing of experimental hybrids supports meiosis-like
sexual recombination in Leishmania. *PLoS Genet* **15**, e1008042,
doi:10.1371/journal.pgen.1008042 (2019).
9 Van den Broeck, F. *et al.* Ecological divergence and hybridization of Neotropical
Leishmania parasites. *Proc Natl Acad Sci U S A* **117**, 25159-25168,
doi:10.1073/pnas.1920136117 (2020).
10 Akopyants, N. S. *et al.* Demonstration of genetic exchange during cyclical development
of Leishmania in the sand fly vector. *Science* **324**, 265-268, doi:10.1126/science.1169464
(2009).
11 Stelzer, C. P. & Lehtonen, J. Diapause and maintenance of facultative sexual
reproductive strategies. *Philos Trans R Soc Lond B Biol Sci* **371**,
doi:10.1098/rstb.2015.0536 (2016).
12 Kwok, A. J., Mentzer, A. & Knight, J. C. Host genetics and infectious disease: new tools,
insights and translational opportunities. *Nat Rev Genet* **22**, 137-153, doi:10.1038/s41576-
020-00297-6 (2021).
13 Inbar, E. *et al.* The mating competence of geographically diverse Leishmania major
strains in their natural and unnatural sand fly vectors. *PLoS Genet* **9**, e1003672,
doi:10.1371/journal.pgen.1003672 (2013).
14 Romano, A. *et al.* Cross-species genetic exchange between visceral and cutaneous strains
of Leishmania in the sand fly vector. *Proc Natl Acad Sci U S A* **111**, 16808-16813,
doi:10.1073/pnas.1415109111 (2014).
15 Louradour, I., Ferreira, T. R., Ghosh, K., Shaik, J. & Sacks, D. In Vitro Generation of
Leishmania Hybrids. *Cell Rep* **31**, 107507, doi:10.1016/j.celrep.2020.03.071 (2020).

16　Louradour, I. *et al.* Stress conditions promote Leishmania hybridization in vitro marked
by expression of the ancestral gamete fusogen HAP2 as revealed by single-cell RNA-seq.
Elife **11**, doi:10.7554/eLife.73488 (2022).
17　Walters, L. L. Leishmania differentiation in natural and unnatural sand fly hosts. *J*
Eukaryot Microbiol **40**, 196-206, doi:10.1111/j.1550-7408.1993.tb04904.x (1993).
18　Panda, S. & Ding, J. L. Natural antibodies bridge innate and adaptive immunity. *J*
Immunol **194**, 13-20, doi:10.4049/jimmunol.1400844 (2015).
19　Mashoof, S. & Criscitiello, M. F. Fish Immunoglobulins. *Biology (Basel)* **5**,
doi:10.3390/biology5040045 (2016).
20　Keyt, B. A., Baliga, R., Sinclair, A. M., Carroll, S. F. & Peterson, M. S. Structure,
Function, and Therapeutic Use of IgM Antibodies. *Antibodies (Basel)* **9**,
doi:10.3390/antib9040053 (2020).
21　Boehm, T., Iwanami, N. & Hess, I. Evolution of the immune system in the lower
vertebrates. *Annu Rev Genomics Hum Genet* **13**, 127-149, doi:10.1146/annurev-genom-
090711-163747 (2012).
22　Serafim, T. D. *et al.* Sequential blood meals promote Leishmania replication and reverse
metacyclogenesis augmenting vector infectivity. *Nat Microbiol* **3**, 548-555,
doi:10.1038/s41564-018-0125-7 (2018).
23　Burkett-Cadena, N. D. in *Medical and Veterinary Entomology (Third Edition)* (eds Gary
R. Mullen & Lance A. Durden) 17-22 (Academic Press, 2019).
24　Volfova, V. & Volf, P. The salivary hyaluronidase and apyrase of the sand fly
Sergentomyia schwetzi (Diptera, Psychodidae). *Insect Biochem Mol Biol* **102**, 67-74,
doi:10.1016/j.ibmb.2018.09.010 (2018).
25　Chaves, M. M. *et al.* The role of dermis resident macrophages and their interaction with
neutrophils in the early establishment of Leishmania major infection transmitted by sand
fly bite. *PLoS Pathog* **16**, e1008674, doi:10.1371/journal.ppat.1008674 (2020).
26　Lee, S. H. *et al.* Mannose receptor high, M2 dermal macrophages mediate nonhealing
Leishmania major infection in a Th1 immune environment. *J Exp Med* **215**, 357-375,
doi:10.1084/jem.20171389 (2018).
27　Navin, T. R., Krug, E. C. & Pearson, R. D. Effect of immunoglobulin M from normal
human serum on Leishmania donovani promastigote agglutination, complement-mediated
killing, and phagocytosis by human monocytes. *Infect Immun* **57**, 1343-1346,
doi:10.1128/iai.57.4.1343-1346.1989 (1989).
28　Maddur, M. S. *et al.* Natural Antibodies: from First-Line Defense Against Pathogens to
Perpetual Immune Homeostasis. *Clin Rev Allergy Immunol* **58**, 213-228,
doi:10.1007/s12016-019-08746-9 (2020).
29　Khasbiullina, N. R. & Bovin, N. V. Hypotheses of the origin of natural antibodies: a
glycobiologist's opinion. *Biochemistry (Mosc)* **80**, 820-835,
doi:10.1134/S0006297915070032 (2015).
30　Bendelac, A., Bonneville, M. & Kearney, J. F. Autoreactivity by design: innate B and T
lymphocytes. *Nat Rev Immunol* **1**, 177-186, doi:10.1038/35105052 (2001).
31　Hayakawa, K. *et al.* Positive selection of natural autoreactive B cells. *Science* **285**, 113-
116, doi:10.1126/science.285.5424.113 (1999).
32　Haury, M. *et al.* The repertoire of serum IgM in normal mice is largely independent of
external antigenic contact. *Eur J Immunol* **27**, 1557-1563, doi:10.1002/eji.1830270635
(1997).

33   Hayakawa, K., Hardy, R. R. & Herzenberg, L. A. Peritoneal Ly-1 B cells: genetic
control, autoantibody production, increased lambda light chain expression. *Eur J*
Immunol **16**, 450-456, doi:10.1002/eji.1830160423 (1986).

Main figures

Fig 1. IgM facilitates *Leishmania* parasites clumping which promotes hybrid formation in

vitro. (**a**) *Leishmania* hybrids genotyping by PCR targeting parental selectable drug markers HYG

(Hygromycin) and BSD (Blasticidin). (**b**) Phase contrast image of *L. major* in culture media

containing 20% fetal bovine serum (FBS) without (bottom panel) or with 5% adult inactivated dog

serum (top panel). Images taken 30 min after cells were seeded in fresh media. Scale bars = 20

μm. (**c**) Human IgM eluted from a HiTrap® IgM column. *Leishmania* clumping activity was

observed in fractions F4-F8 corresponding to elution of IgM. (**d**) Presence (+) or absence (-) of *L.*

major clumping activity in inactivated adult sera from different vertebrate species, FBS, or

immunoglobulins (IgM, IgG, IgA) purified from adult bovine or human sera. (**e**) Western blot to

detect IgM in FBS and adult bovine serum (ABS). Serum purified bovine IgM, positive control;

serum albumin, loading control. (**f**) Phase contrast image of *L. major* in culture media containing

20% FBS with bovine IgM at 50 µg/mL after 24h and (**g**) with peanut agglutinin (PNA) at 50

µg/mL after 24h. Scale bars = 50 µm. (**h**) Phase contrast image of *L. major* in culture media

containing 20% FBS with bovine IgM (pentameric form) at 50 µg/mL after 24h or (**i**) with bovine

IgM (monomeric form) at 50 µg/mL after 24h. (**j**) Clumping activity of various *Leishmania* species

in the presence of bovine IgM at 50 µg/mL. (**k**) (top) Workflow for in vitro crossing of *L. major*

parental lines. Mating wells were established with 50 µg/mL IgM in complete Schneider's media.

Double drug resistant hybrid parasites were cloned before genotyping. (bottom) Summary of data

from in vitro crossing of parental lines. 1x2 (n=30), 1x3 (n=40), 2x3 (n=7). (**l**) *Leishmania major*

hybrids genotyping by PCR targeting parental selectable drug markers HYG (Hygromycin), BSD

(Blasticidin) and SAT (Nourseothricin). Parental 1, WR-SSU-HYG; Parental 2, FVI-FKP40-

BSD; parental 3, FVI-FTL-SAT; ntc, no template control; L, 1kb plus ladder (Invitrogen).

Fig 2. Characterization of the *Leishmania* mating clump. (a) Confocal immunofluorescence

wide Z and tile images showing the association of IgM with the *L. major* mating clump after 24h

of culture. Dapi (nuclei, kinetoplast) blue; IgM, red. Scale bar = 40 µm. Bottom rectangles for each

panel represent XZ slices. Scale bar = 30 µm. **(b)** Confocal immunofluorescence of a 5 µm

transversal section of the *L. major* mating clump. Dapi (nuclei; kinetoplast), blue; alpha-tubulin,

green; IgM, red. IgM localizes between cells within the clump (right panel). Scale bars = 5 µm.

Bottom rectangles for each panel represent XZ slices. Scale bars = 3 µm. **(c)** IgM colocalizes on

the surface of parasites (arrows in inset). Scale bar = 5 µm. Bottom rectangles represent XZ slices.

Scale bar = 3 µm. **(d)** SEM of a *L. major* (upper panel) and *L. tropica* (lower panel) mating clump

showing the 3D organization of promastigote forms. Scale bar: 15 µm. **(e)** Confocal

immunofluorescence of a 5 µm transversal section of the *L. tropica* mating clump. *L. tropica*

promastigotes were grown to stationary phase in the presence of EdU or BrdU then extensively

washed and used at 1:1 ratio to assemble the LMC with IgM. EdU, green; BrdU, red. White arrows

point to yellow nuclei indicative of fusion and exchanged genetic material. Scale bars, 3 µm. **(f)**

TEM of a *L. tropica* mating clump showing cells with 3 nuclei. Arrow, individual nucleus within

a parasite cell body. Scale bar: 5 µm.

Fig 3. The impact of IgM on *Leishmania* hybrid formation in the sand fly vector. **(a)** Immunoblot detecting IgM in the sand fly midgut after multiple blood meals. Sand flies were provided an artificial uninfected blood meal with or without IgM. Arrows indicate the timing of the first (BM1, 0 days) and second (BM2, 6 days) blood meals. Pools of 15 midguts were used. 7.5 µg of total protein was loaded per lane. **(b)** Experimental design for the generation of in vivo crosses after sand flies acquire parasites by feeding on a footpad lesion containing two *L. major* parental lines. Bovine IgM was used. **(c)** Percent of sand flies harboring hybrids for 3 different combinations of parental lines (1x2; 1x3; 2x3) in single fed sand flies or in sand flies fed twice in the presence or absence of IgM. Colors indicate independent experiments. **(d)** Cumulative data

from independent experiments for infection status and estimated frequency of hybridization per

fly. 1x2 (n=4), 1x3 (n=3), 2x3 (n=3). Per each biological replicate ~ 20 infected flies per group.

Fisher's exact test, p < 0.05 is considered significant. Pairwise comparison indicate source of group

significance: a, no difference; b, significant difference. **(e)** Whole-genome analysis of in vitro- and

in vivo-generated hybrids for parental line combination 1x2. Biparental ancestry was confirmed

across the whole genome for all crosses. Parental 1, WR-SSU-HYG; Parental 2, FVI-FKP40-BSD;

Parental 3, FVI-FTL-SAT. *P < 0.05, **P ≤ 0.01, ***P ≤ 0.001.

444
Fig 4. The impact of IgM on *Leishmania* genetic exchange in the sand fly vector. **(a)**

Experimental design for the generation of in vivo backcrosses after sand flies acquire parasites by

feeding on a footpad lesion containing one non-hybrid *L. major* parental line and one hybrid *L.*

major parental line. A (1x2) F1 hybrid was injected at 1:1 proportion with the parental 2 harboring

different resistance markers (parental 2'and parental 2"). Bovine IgM was used. **(b)** Percent of

sand flies harboring hybrids in single fed sand flies or in sand flies fed twice in the presence or

absence of IgM. Colors indicate independent experiments. **(c)** Cumulative data from independent

experiments for infection status and estimated frequency of hybridization per fly. (1x2) x 2' [n=3];

(1x2) x 2'' (n=2). Per each biological replicate ~ 30 infected flies per group. Fisher's exact test, p

< 0.05 is considered significant. Pairwise comparison indicate source of group significance: a, no

difference; b, significant difference. **(d)** Whole-genome analysis of in vivo-generated backcross

hybrids for parental line combination 1x2. Where the biparental ancestry was consistent across

each chromosome, recombination events have broken this up in the backcrosses, making parental

copy number discontinuous always in the same direction through parental 2. Parental 1,WR-SSU-

HYG; Parental 2, FVI-FKP40-BSD; Parental 2', FVI-FKP40-SAT; Parental 2'', FVI-SSU-SAT.

*P < 0.05, **P ≤ 0.01.

Methods

Ethics statement

All animal experimental procedures were reviewed and approved by the National Institute of Allergy and Infectious Diseases (NIAID) Animal Care and Use Committee under animal protocol LMVR4E. The NIAID DIR Animal Care and Use Program complies with the Guide for the Care and Use of Laboratory Animals and with the NIH Office of Animal Care and Use and Animal Research Advisory Committee guidelines. Detailed NIH Animal Research Guidelines can be accessed at https://oma1.od.nih.gov/manualchapters/intramural/3040-2/.

Animals

Six to eight weeks old female BALB/c and Swiss Webster mice were obtained from Envigo and Charles River laboratories, respectively. Animals were housed under pathogen-free conditions at the NIAID Twinbrook animal facility, Rockville, MD.

Parasites

Leishmania parasites used in this study: *Leishmania major* (WR 2885) isolated from a soldier deployed to Iraq[34]; *Leishmania major* (MHOM/IL/80/Friedlin); *Leishmania tropica* (MHOM/SU/74/K27 - ATCC 50129); *Leishmania braziliensis* (MHOM/BR/75/M2903 - ATCC 50135); *Leishmania infantum* (MCAN/BR/09/52) isolated from a dog spleen in Natal, Brazil[35]; *Leishmania donovani* (MHOM/SD/62/1S); *Leishmania tarentolae* (Tar II – ATCC 30143); *Leishmania mexicana* (MNYC/BZ/62/m379); *Leishmania amazonensis* (IFLA/BR/67/PH8). Parasites were grown at 27°C in Schneider's Drosophila Insect Medium (Lonza BioWhittaker, 04-351Q) supplemented with 20% heat inactivated fetal bovine serum (FBS) (Gibco, 16140071) and 250 µM of adenosine (MilliporeSigma, A9251). Parasites were maintained by serial passage in BALB/c mice or Golden Syrian hamsters, depending on dermotropic or viscerotropic parasite

497 species characteristics. Promastigotes were kept in culture for a maximum of 10 passages. *L. major*

498 metacyclic promastigotes were purified by peanut agglutinin (PNA) (Vector Laboratories, L-1070)

499 and used for in vitro experiments and infection of mice footpads as described[36].

***Leishmania* parental lines**

The various *Leishmania* lines used in crossing experiments were constructed as detailed below.

For parental line 1, *L. major* – WR-SSU-HYG/GFP, *L. major* (WR 2885) was used as background

for generation of a hygromycin B (HYG) parental line that also harbors the EGFP gene. The EGFP

gene was amplified by PCR using the mEGFP-C1 plasmid as a template for the forward primer 5-

ACAGCGAGATCTATGGTGAGCAAGGGCGAGGA-3,   and   reverse   primer   5-

ACAGCGGGTACCCTTGTACAGCTCGTCCATGC-3. The forward and reverse primers add the

BgIII or KpnI restriction sites to the amplicon, respectively. The PCR product was cloned into the

Lexsy_hyg2 plasmid (Jena Bioscience, EGE-232). The linearized construct (2 μg) was used to

transfect mid-log phase culture promastigotes on a 4D-NucleofectorTM X unit (Lonza) using pulse

code FI-115 (T cell, Human, unistim. HE). Transfected parasites bearing rRNA locus-integrated

EGFP were selected for growth in the presence of 100 μg/mL Hygromycin B (Goldbio, H270).

Clones were selected by limiting dilution and assessed for chromosome integration. Parental line

2, *L. major* – FVI-FKP40-BSD[37] and Parental line 3, *L. major* – FVI-FTL-SAT[38] were generated

as previously described. For generation of parental line 2', *L. major* – FVI-FTL-SAT, we followed

the procedures described for line 2, but used a heterozygous of the gene FKP40 with a construct

harboring Nourseothricin (SAT). For parental line 2'', *L. major* – FVI-SSU-SAT/RFP, *L. major*

(MHOM/IL/80/Friedlin/FV1) was used as background for the generation of a SAT parental line

that also harbors the mRuby gene. The mRuby gene was amplified by PCR using the mRuby-C1

plasmid   as   template   for   the   forward   primer   5-

ACAGCGAGATCTATGAACAGCCTGATCAAAGA-3,   and   reverse   primer   5-

ACAGCGGCGGCCGCTCACAGATCCTCTTCAGAGATGAGTTTCTGCTCCCCTCCGCCC

AGGCCGGC-3. The forward and reverse primers add the BgIII or NotI restriction sites to the

amplicon, respectively. The PCR product was cloned into the Lexsy_sat2 plasmid (Jena

Bioscience, EGE-234). The linearized construct (2 µg) was used to transfect mid-log phase culture

promastigotes on a 4D-NucleofectorTM X unit (Lonza) using pulse code FI-115 (T cell, Human,

unistim. HE). Transfected parasites bearing rRNA locus-integrated mRuby were selected for

growth in the presence of 100 µg/mL Nourseothricin (Goldbio, N-500-100). Clones were selected

by limiting dilution and assessed for chromosome integration. *L. tropica* (K27 – ATCC 50129)

was used as background for the generation of parental line 4, *L. tropica* - K27-SSU-HYG/GFP and

parental line 5, *L. tropica* - K27-SSU-SAT, respectively. Parental line 4 was made following the

same steps as described for Parental line 1. Parental line 5 was made the same way as parental line

2'' but without the addition of any fluorescent gene. mEGFP-C1 and mRuby-C1 were a gift from

the Michael Davidson lab (Addgene plasmid # 54759, http://n2t.net/addgene:54759,

RRID:Addgene_54759; Addgene plasmid # 54552, http://n2t.net/addgene:54552,

RRID:Addgene_54552).

In vitro *Leishmania* clumping activity

Various Leishmania species were cultured in complete Schneider's medium in 25 cm^2 flasks.

4×10^6 /mL stationary phase promastigotes were dispensed into a well of a 24-well plate in the

presence of 50 µg/mL of bovine IgM to test clumping activity. For *L. major*, sera from different

animal sources were also tested for clumping activity in culture. Naïve beagle dog and germ-free

mouse blood were provided by the Division of Veterinary Resources (DVR/NIH) under animal

use protocol LMVR4E. Serum was obtained by centrifugation at 1000xg/10 min, inactivated at

56°C for 30 min, filtered sterilized through a 0.22 µm PES membrane (Millipore MILLEX GP,

SGLP033RB) and stored at -80 in aliquots and used without thaw/freeze cycles. Sera acquired

from commercial vendors are: Baboon (MilliporeSigma, S6884); Bovine -adult (MilliporeSigma,

B9433); Bovine -fetal (Gibco, 16140071. R&D, S12495, S11595); Chicken (MilliporeSigma,

C5405); Dog (BioChemed Services, TD092321NIH); Donkey (MilliporeSigma, D9663); Goat

(MilliporeSigma, G9026); Guinea Pig (MilliporeSigma, G9774); Horse (MilliporeSigma, H1270);

Human (MilliporeSigma, H4522, H6914, P2918, S7023, S1-M); Mouse (MilliporeSigma, S25-M,

S3509, S7273); Porcine (MilliporeSigma, P9783); Rabbit (MilliporeSigma, R4505); Rat

(MilliporeSigma, S7648); Sheep (MilliporeSigma, S2263). Sera were heat inactivated at 56ºC for

30 min and filtered sterilized before usage at 20% final concentration.

In vitro *Leishmania* mating clump assembly for recovery of hybrids

Parental lines of *L. major* were cultured in complete Schneider's medium in 25 cm^2 flasks. 3x10^6

stationary phase promastigotes from each parental were mixed in a well of a 24-well plate with 1.5

mL of fresh complete Schneider's medium + 50 µg/mL of bovine IgM, empirically determined as

the optimal concentration that supports the formation and dissociation of the mating clump in vitro.

Mating wells were maintained at 27ºC for 3 days and then diluted 1:10 in fresh media in a 25 cm^2

flask containing selective drugs. Cultures were followed for up to 30 days to allow for double-drug

resistant cell lines to grow. Selective drugs were combined according to the parental lines used per

experiment. The concentration used for each selective drug was as follows: Hygromycin B - 100

µg/mL (Goldbio, H270); Blasticidin - 10 µg/mL (Goldbio, B-800-500); Nourseothricin - 100

µg/mL (Goldbio, N-500-100). Once resistant cells grew, the polyclonal culture was cloned by

plating 0.5 cell/well in a 96-well plate under double-drug pressure. After clone selection,

genotyping was carried out using primers as displayed in Supplementary Table 3. Clone cultures

were also a source of genomic DNA for deep sequencing. An illustration of the experimental

design for in vitro *Leishmania* genetic exchange is provided in Fig. 1g. Frequencies of

hybridization were estimated by: $\frac{\# \, positive \, crosses}{(Initial \, \# \, of \, one \, parental)*(\# \, trial \, crosses)}$

Mice infection

Metacyclic promastigotes were harvested from stationary phase cultures and purified using peanut

agglutinin (Vector Laboratories, L-1070) as previously described[36]. Purified *L. major* metacyclic

promastigotes ($1x10^5$) of each parental line were mixed at a 1:1 ratio and injected into a BALB/c

mouse footpad. Animals that developed a swollen non-ulcerated (<5 mm thickness) footpad lesion

at weeks 4 to 6 post-injection were used to feed sand flies. After sand fly feeding on infected

animals, amastigotes were harvested from the infected footpads and skin draining lymph nodes as

previously described[39].

Sand fly infection

Four to six days old female *Lutzomyia longipalpis* (Jacobina strain) sand flies were infected by

feeding on mice footpad lesions of BALB/c mice. For first (F1) and second (backcross) generation

crosses, culture purified metacyclic promastigotes of two parentals were injected in the mouse

footpad at a 1:1 proportion. In F1 crosses experiments, mice were injected with parentals 1x2, 1x3

and 2x3. For backcross experiments mice were injected with a (1x2 F1 hybrid) x 2 harboring a

different resistance marker. Illustration of the experimental design for natural feeding on mice is

provided in Fig. 3b & 4a. Briefly, mice were anesthetized by intraperitoneal injection of a mixture

589 of ketamine (100mg/kg) and xylazine (10mg/kg). The infected footpad was then inserted in a

590 cardboard pint containing the sand flies. Sand flies were allowed to feed for 2 hours in the dark.

591 After feeding, blood-fed female sand flies were sorted out and maintained at 26°C with 75%

592 humidity and 30% sucrose solution *ad libitum*. *Lutzomyia longipalpis* is not the natural vector of

593 *L. major* but it is a permissive sand fly species that has been shown to support the complete

594 development of several *Leishmania* species from natural parasite acquisition to successful

595 transmission by bite[40, 41,42]. Importantly, midgut development related to parasites numbers and

596 stages of differentiation in *Lu. longipalpis*/*L. major* infections was similar to those reported for

597 *Phlebotomus papatasi / L. major* and *Lu. longipalpis / L. infantum* natural occurring

598 combinations[43].

Preparation of the second blood meal provided to sand flies

601 For the 2nd blood meal given six days post-infection to a proportion of the sand flies, defibrinated

602 rabbit blood (Spring Valley Laboratories) was centrifuged at 1000xg for 10 min and the normal

603 serum containing IgM was discarded. Red blood cells (RBCs) were washed three times with 10mL

604 of PBS (Lonza BioWhittaker, BE17-516F) followed by centrifugation at 1000xg for 10 min. RBCs

605 were washed an additional time with FBS (Gibco, 16140071). Blood was reconstituted by adding

606 FBS at the same volume of the serum initially removed. Sand flies were allowed to feed for 2 hours

607 on a glass feeder containing reconstituted blood supplemented or not with bovine IgM

608 (MilliporeSigma, I8135) at 500 µg/mL. Unfed and blood-fed flies were sorted out after feeding

609 and kept on 30% sucrose solution *ad libitum* for a total of 14 days after the initial infected blood

610 meal. Females infected by feeding on mice lesions only were used to represent sand flies that have

611 taken a single infected blood meal.

Sand fly midgut counts

To assess maturity of the infection, *Leishmania*-infected sand fly midguts were individually transferred to a 1.7mL microtube (Denville Scientific, C2172) filled with 30μL of PBS. Midguts were then ground using a disposable pellet mixer and a cordless motor (Kimble, 7495400000). For counting, promastigotes were slowed down by a further 1:5 or 1:10 dilution in formalin/PBS at 0.005% final concentration. Then, 10 μL of the sample were loaded onto a Neubauer improved chamber (Incyto, DNC-NO1) and parasites were counted under a phase contrast Axiostar plus microscope (Zeiss) at 40X magnification. The number of parasites per midgut was back-calculated based on the total sample volume and subsequent dilution.

In vivo recovery of hybrids

Only midguts of sand flies that developed a mature infection, recognized by an opaque anterior midgut distended by the presence of a mass of parasites[43], were plated in search for hybrids. Infected midguts were dissected in PBS on microscope slides using tweezers and fine needles. Each midgut was transferred to a 1.7mL microtube (Denville Scientific, C2172) filled with 50 μL of complete Schneider's medium and homogenized using a pestle with a cordless motor. One midgut was plated per well in a 96-well plate (Alkali Scientific, TP9096) at a final volume of 100 μL. An antibiotic cocktail (ABTs) that has no effect on parasite growth (Supplementary Fig. 13) and viability was included in complete media to control contamination with gut microbiota. ABTs: Penicillin-Streptomycin (Gibco, 100 U/mL); Gentamicin (Sigma, 50 μg/mL); Caspofungin (Sigma, 15 μg/mL); 5-fluorocytosine (Sigma, 30 μg/mL). For each independent experiment, a minimum of 20 infected midguts per group were collected. After plating, wells were allowed to stabilize and grow for 24h prior to the addition of 100 μL/well of complete Schneider's media

containing selective drugs (2x concentrated). Cultures were followed for up to 30 days for double

or triple-drug resistant cell lines to grow. The concentration of selective drugs, cloning, and

genotyping procedures were performed as described for in vitro recovery of hybrids. Frequencies

of hybridization were estimated by: $\frac{\text{\# positive crosses}}{(Median\ \#\ of\ parasites\ per\ fly)*(\#\ trial\ crosses)}$.

IgM identification from plasma by HPLC

Canine plasma (1.5-2 mL) was inactivated at 56°C for 30 min. Inactivated plasma was passed

through a 0.22 µm PES membrane (Millipore MILLEX GP, SGLP033RB) and diluted to 7 mL in

PBS containing 5 mM EDTA (buffer A). The inactivated, diluted plasma was then applied to a

Sephacryl S200 (16/60, GE Healthcare) gel filtration column equilibrated with buffer A. Plasma

components were eluted at 1 mL/min, and 2 mL fractions were collected and tested for *Leishmania*

clumping activity. Active fractions were then combined, concentrated, and diluted several times

with 10 mM sodium phosphate (pH 6.0) buffer (buffer B) before application to an SP Sepharose

(16/10, GE Healthcare) ion exchange column equilibrated with buffer B. Proteins were eluted with

a gradient of 0-1 M NaCl in buffer B. Activity was distributed throughout the major peak,

particularly at the leading edge. These fractions were pooled and chromatographed on a Superdex

200 (10/300, GE Healthcare) gel filtration column equilibrated with buffer A, resulting in further

purification of the major component that produced bands on SDS-PAGE gels of approximately 70

and 25 kDa consistent with the heavy and light chains of IgM. This sample was active in

Leishmania clumping assays and was analyzed after tryptic digest by electrospray ionization mass

spectrometry. As an additional means of purification, pooled fractions of IgM purified by gel

filtration were diluted in 20 mM NaPO4 pH 7.5 (buffer C) containing 0.8 M ammonium sulfate

and applied to a 1 mL HiTrap IgM purification HP column (GE healthcare) equilibrated with buffer

C containing 0.8 M ammonium sulfate. The column was washed until the absorbance reached

baseline levels. IgM was then eluted stepwise in buffer C and 0.5 mL fractions were collected.

Activity in the *Leishmania* clumping assay closely aligned with the peak of absorbance.

Commercial IgM preparation for in vitro and in vivo experiments.

Commercially available IgM antibodies used in this study were obtained from MilliporeSigma and

contained preservatives such as sodium azide. To avoid any potential toxicity to the parasites,

removal of preservatives was undertaken. IgM from human (MilliporeSigma, I8260) and bovine

(MilliporeSigma, I8135) sources were purified by passing them through a Sephacryl-S-200 gel

filtration column (Cytiva) equilibrated with either buffer A minus EDTA or PBS. For IgM

equilibrated with buffer A minus EDTA, further purification steps using a HiTrap IgM purification

HP column are given above. Protein-containing fractions eluted with PBS were pooled and the

volume reduced to the desired concentration by ultrafiltration.

IgM stability in blood fed sand flies and its absence in fetal serum.

To estimate how long IgM is maintained in the midgut after blood feeding, *Lu. longipalpis* were

provided two uninfected blood meals on days 0 and 6, with or without 500 µg/mL of bovine IgM.

Midguts were dissected daily on days 0 to 10 (*n* = 15) post initial blood meal and collected in 50µL

of RIPA lysis buffer (Thermo-Fisher Scientific, 89901) supplemented with phenylmethylsulfonyl

fluoride (PMSF) (MilliporeSigma, 93482) and Halt Protease Inhibitor Cocktail (Thermo-Fisher

Scientific, 87786). Samples were gently macerated with a disposable pellet mixer and a cordless

motor, then vortexed and centrifuged at 12,000 rpm for 10 min at 4°C. The supernatant was

collected, and its protein content was quantified by BCA (Thermo-Fisher Scientific, 23227). A

total of 7.5 µg of protein was loaded in a 15-well NuPAGE 4%–12% Bis-Tris protein precast gel

(Thermo-Fisher Scientific, NP0335BOX), run under reducing conditions by the addition of the

NuPAGE Sample Reducing Agent with DL-Dithiothreitol (DTT) (Thermo-Fisher Scientific,

NP0009), and then denaturation at 95°C for 5 min. Proteins were transferred to a 0.45mm

nitrocellulose iBlot 2 Transfer Stack membrane using the iBlot 2 Gel Transfer device (Thermo-

Fisher Scientific, IB23001). To check for successful loading of antigens, No-Stain Protein

Labeling Reagent (Invitrogen, A44717) was used following the membrane manufacture's

protocol. Next, membranes were blocked overnight at 4°C in 5% non-fat milk diluted in Tris-

Buffered Saline, 0.1% TWEEN 20 (TBST) followed by 3 hours of incubation at RT with sheep

anti-bovine IgM (Bio-Rad, AAI179) diluted at 1:1000. Membranes were washed three times with

TBST followed by incubation for 30 min with rabbit anti-sheep IgG H+L secondary antibody

conjugated to horse radish peroxidase (HRP) (Abcam, Ab6747) at a 1:5000 dilution. Membranes

were washed three times with TBST and blots were developed by the addition of SuperSignal West

Pico PLUS Chemiluminescence substrate (Thermo-Fisher Scientific, 34579). The signal was

captured after 2 min using the Azure c600 chemiluminescence imager (Azure Biosystems).

Membranes were stripped by adding 10 ml of Restore Western Blot Stripping Buffer (Thermo-

Fisher Scientific, 21059) for 15 minutes at 37°C, followed by two washes with TBST. Blots were

re-blocked overnight and incubated with recombinant rabbit anti-bovine serum albumin antibody

(Abcam, ab192603) at 1:1000 dilution for 3 hours, followed by a 30 min incubation with HRP-

conjugated goat anti-rabbit IgG H+L secondary antibody (Cell Signaling Technology, 7074S) at

1:1000 dilution. Blots were re-imaged as detailed above. To access the presence of IgM in fetal

serum, NuPAGE 4%–12% Bis-Tris protein precast gels (Thermo-Fisher Scientific, NP0335BOX)

were loaded with 1 µg of either Bovine IgM (Bio-Rad, AAI179), fetal bovine serum (Thermo-

Fisher Scientific, 16140-071) or adult bovine serum (Millipore-Sigma, B9433) under reducing

conditions followed by denaturation at 95°C for 5 min. The pattern of the proteins was visualized

by Coomassie blue staining using the eStain™ L1 protein staining system. For immunoblotting,

proteins were transferred onto nitrocellulose membranes, blocked, and probed with sheep anti-

bovine IgM at 1:1000 followed by a 30 min incubation with HRP-conjugated anti-sheep IgG H+L

secondary antibody (Abcam, Ab6747) diluted at 1:5000. Membranes were stripped and incubated

with recombinant rabbit anti-bovine serum albumin antibody at 1:1000 followed by a 30 min

incubation with HRP-conjugated goat anti-rabbit IgG H+L secondary antibody (Cell Signaling

Technology, 7074S) at 1:1000 and imaged after addition of the chemiluminescence substrate.

IgM depolymerization

To obtain monomeric IgM antibodies, 1 mg of bovine IgM (MilliporeSigma, I8135) was subjected

to a digestion buffer exchange (50 mM Tris, 150 mM NaCl, 10 mM $CaCl_2$, 0.05% NaN_3; pH 8.0)

- final concentration of 1 mg/ml. Afterward, 1 ml of IgM was incubated in the presence of 6 mg

2-mercaptoethylamine at 37 °C for 30 minutes in the dark. The reduced IgM was further alkylated

in the presence of 16 mg iodoacetamide, followed by purification using a Superdex 200 Increase

(10/300 GL; GE Healthcare) equilibrated with Buffer A minus EDTA. After size exclusion, a clear

separation between the pentameric (peak 1 – P1) and monomeric (peak 3 – P3) IgM fractions were

observed via native gel electrophoresis (NativePAGE Novex Bis-Tris Gel system; Life

Technologies, Carlsbad, CA, USA). All fractions corresponding to defined peaks were pooled

together, adjusted to a suitable concentration and frozen at -20 °C until further use.

Mass spectrometry

Protein identification was carried out with a standard bottom-up method. The sample was reduced

with DTT, alkylated with iodoacetamide, then digested using Lys-C followed by trypsin. The

digested sample was cleaned using C18 tips (Agilent, OMIX10) and subjected to the LC-MS

(EASY nLC 1200 and Orbitrap Fusion Lumos mass spectrometer, Thermo Scientific). The

peptides were separated using a PepMap 100 C18 reversed phase column, and a standard data-

dependent acquisition was performed. The survey MS1 (Top Speed mode, 3 seconds) was done

using the Orbitrap mass analyzer, and the ddMS2 was done using the Linear Ion Trap where the

fragmentation was done with CID. Dynamic exclusion was set for the duration of 20 seconds. The

data analysis was done using PEAKS Studio software (Bioinformatcs Solutions Inc.). The database

used is the UniProt proteome UP000002254 (Dog, taxonomy id 9615). The number of identified

peptides, and protein abundance in the sample were confirmed by the PEAKS scores.

Video recording and stereomicroscopic imaging

741 Midgut videos were taken using an iPhone 10 camera connected to the Stemi 508 stereomicroscope

742 (Zeiss) ocular by a microscope mount (iDu Optics, LabCam iPhone 10 microscope adapter with

743 built-in 30mm 10X WF lens). Phase contrast images and videos were recorded using a K5 camera

744 coupled to a Thunder Imager microscope (Leica). Raw files were opened using Fiji ImageJ

745 software[44] and exported at 25 to 90 frames per second.

Immunofluorescence of in vitro *Leishmania* mating clumps

After 24 hours of formation in vitro, whole Leishmania mating clumps were washed twice with PBS before fixation with 4% Paraformaldehyde (PFA) overnight at 4ºC. Fixation was followed by permeabilization with 0.1% PBS Triton for 10 minutes, then 0.1 M glycine in PBS solution for 10 minutes, and a final blocking step with 0.1% Triton/0.1% BSA/ PBS for an hour at room temperature. Mating clumps were then incubated at 4ºC overnight with mouse anti-alpha-tubulin monoclonal antibody (MilliporeSigma, T6199) conjugated to Alexa 555 (1:1000), and sheep anti-bovine IgM (Bio-Rad, AAI179) conjugated to Alexa 647 (1:1000) diluted in blocking buffer. Anti-Bovine IgM antibody was conjugated to an Alexa 647 fluorophore using the READILINK™ 647/674 Antibody Labeling Kit (Bio-Rad, 1351006) following the manufacturer's instructions. The next day, mating clumps were washed three times with blocking buffer, ten minutes each, and then washed two times with permeabilization buffer, both at room temperature. Clumps were counterstained with 2 µM of Hoechst 33342 (Thermo Fisher Scientific, 62249) in permeabilization buffer for 30 minutes at room temperature, followed by a quick rinse of 0.1% Triton/PBS. Clumps were placed in a u-slide angiogenesis (IBIDI, 81506) well in Prolong gold antifade medium (Thermo Fisher Scientific, P10144) for imaging. For the sections, mating clumps were fixed in 4% paraformaldehyde overnight at 4ºC, processed on the Leica ASP 6025, and then embedded in histology grade paraffin. All samples were sectioned at 3 µm. Mating clump section pretreatment was performed on the Leica Bond RX. Sections were baked for 30 minutes at 60ºC and then dewaxed for 30 seconds in Bond Dewax solution (Leica, AR9222) heated to 72ºC. Sections were rehydrated with absolute ethanol washes followed by 1x ImmunoWash (StatLab, ACR-015), then subjected to a 20-minute treatment with Epitope Retrieval Solution 1 (Leica, AR9961) heated to 100ºC. Slides were then washed with PBS and stained. Steps for immunofluorescence of sections

were based on Zaqqout et al. 2020[45]. Briefly, slides were permeabilized with PBS/gelatin/Triton

0.25%, twice for 10 minutes each at room temperature, followed by blocking with 5% BSA/PBS

for one hour at room temperature in a humid box. The primary antibodies listed above were

incubated in a blocking solution and kept overnight in a humid box at 4ºC. The next day, slides

were washed three times with PBS and then with permeabilization solution for 10 minutes at room

temperature, and counterstained with 2 µM of Hoechst 33342 in the same buffer for 30 minutes at

room temperature followed by a quick rinse with 0.1% Triton/PBS. Another final rinsing step with

10mM $CuSO_4$/ 50mM NH_4Cl solution for 10 minutes and a final short rinse with water was

performed. Slides were closed using Prolong gold antifade mounting media and a coverslip.

Confocal Microscopy

Images were captured using a Leica TCS SP8 DMi8 inverted fluorescence confocal microscope

(Leica Microsystems) equipped with a photomultiplier tube/hybrid detector and a Leica DFC345

Monochrome camera. Images were taken using sequential acquisition, variable z-steps, mosaic

size, and integration. Images were viewed with the 40x, 63x, and 100x oil immersion objective

(zoom factor of 2, 3, or 4), and data were collected using the Leica Application Suite X software

platform. White light laser and specific emission and excitation range were applied depending on

the fluorophore used. The following spectra for excitation/emission were used: 488/520 for EdU

and BrdU; 555/580 for alpha-tubulin; 594/620 for BrdU and EdU; and 647/665 for IgM. DAPI

was excited using a 405-nm diode laser. Images were taken using sequential acquisition and

variable z-steps. Whole mating clumps were imaged in the Z-wide mode (8-bit) and tile mode to

capture the entire clump. Individual fields were merged and stitched together to mount the

composite of the whole clump. Image processing was performed using Imaris 9.2.1 (Bitplane,

Oxford Instruments). Maximum intensity projection (MIP) is represented in xy pictures, and cross-

sections of between 2-3 µm were obtained with the ortho-slicer feature from Imaris.

BrdU and EdU staining

Leishmania promastigotes were incubated with 10 µM of EdU - 5-Ethynyl-2'-deoxyuridine

(Abcam, ab146186) or BrDu - 5-bromo-2'-deoxyuridine (Abcam, ab142567) for 4 days to reach

the stationary growth phase. Parasites were then washed twice with PBS and an equal proportion

of each of the labeled promastigotes was used for mating clump assembly. Mating clumps were

fixed and processed for sections as described above. Three-micrometer sections were washed with

PBS, permeabilized with 0.2% Triton/PBS for 10 minutes, then incubated with 0.1M glycine for

10 minutes at room temperature. Samples were then blocked with 0.5% BSA/PBS for 10 minutes

and then briefly washed with PBS before the Click reaction was performed. We used the Click-it

EdU Cell Proliferation Kit to image incorporated Edu with Alexa Fluor 488 dye (Invitrogen,

C10337). The reaction was prepared following the manufacturer's instructions and incubated with

the samples for 30 minutes at room temperature in a humid and dark chamber. After the click

reaction, sections were washed three times with PBS and blocked with 0.5% BSA/PBS for one

hour at room temperature. Primary mouse monoclonal anti-BrdU (MOBU-1) (Invitrogen, B35128)

(1:100) was incubated in blocking buffer overnight at 4ºC. The next day, sections were washed

with blocking buffer three times, 10 minutes each, and incubated with goat anti-mouse Alexa 594

secondary antibody (Invitrogen, A-11005) at 1:1000 in 0.1% Triton/PBS for 2 hours at room

temperature. Slides were washed three times with 0.1%Triton/PBS for 10 minutes each and then

counterstained with 2 µM of Hoechst 33342 in 0.1% Triton/ PBS for 30 minutes at room

temperature. A final rinse was done with 10mM $CuSO_4$/ 50mM NH_4Cl solution for 10 minutes

and followed by a quick rinse with water before the slides were closed with Prolong gold antifade

mounting media and a coverslip.

Genome sequencing

Genomic DNA was extracted from cloned parasite culture pellets using the E.Z.N.A.® DNA/RNA

Isolation Kit (Omega Biotek, R6731) and following the manufacturer's recommendations for

cultured cells. Sample concentration was measured by a DS-11+ spectrophotometer (DeNovix

Inc.). A minimum of 3 µg of DNA was sent for sequencing (Novogene Co.). Genomic DNA was

randomly fragmented by sonication, then DNA fragments were end polished, A-tailed, and ligated

with full-length adapters for Illumina sequencing followed by further PCR amplification with P5

and indexed P7 oligos. The PCR products were purified with the AMPure XP system and libraries

were checked for size distribution using an Agilent 2100 Bioanalyzer (Agilent Technologies) and

quantified by real-time PCR (to meet the criteria of 3 nM). Qualified libraries were fed into

Illumina sequencers after pooling to obtain effective concentrations and the expected data volume.

Genomic analysis

For Supplementary Fig. 8, raw sequencing reads were assessed for quality using the FastQC

software. Fastq files were imported and processed using Geneious prime software v2021.2.2

(https://www.geneious.com). Low quality reads were trimmed from fastq files (<20) and

contaminating adapter primer sequences were removed using BBDuk trimmer plugin. Trimmed

reads were mapped to the reference genome TriTrypDB-52_LmajorFriedlin using bowtie 2 plugin.

BAM files with aligned reads were analyzed with qualimap v2.2.1[46]. The median genomic

coverage of all samples was 33.1X. Using bowtie 2, unmapped reads were aligned to an artificial

chromosome with selective markers. The artificial chromosome was created by concatenating the

resistance genes integrated into the parental lines, HYG (Hygromycin), BSD (Blasticidin) and

SAT (Nourseothricin), buffered by a 1.2kb (150bp repeated 8 times) sequence from mouse (*Mus*

musculus) chromosome 1. For Fig. 3e, 4d and Supplementary Fig. 9, raw sequencing reads were

trimmed using Trimmomatic v0.39[47] and mapped using BWA-MEM v07.17[48] to the TriTrypDB-

52_LmajoprFriedlin reference genome with the resistant gene sequences inserted as additional

contigs. PCR duplicates were marked using Samblaster v0.1.2.5[49] and germline variants were

called jointly across all samples using the HaplotypeCaller tool from GATK v4.1.9.0 and using

default settings. Resulting variants were then filtered down to only biallelic sites with no missing

data across all samples. The genomic insertion locations for each resistance gene were validated

by examining split reads from each sample, and identifying the location of reads with one pair

member mapped to the construct and the other pair member mapped to the human genome. Any

locations with less than 3 supporting split read pairs were excluded from consideration. To quantify

the allelic copy number of each contributing parent in F1 offspring, resulting BAM files were first

used as input to the PAINT software suite[50] to generate chromosome-level somy levels

(Supplementary Fig. 10). A custom perl script was then used to execute the following steps with

the multisample VCF and chromosome somy estimates as input: 1) identify homozygous SNP

differences between parents for each cross, ignoring sites at <20X depth in both parental strains;

2) for each SNP identified in step 1, calculate the frequency of each allele based on read counts

supporting each allele (e.g., allele1 proportion = (# reads supporting allele 1 at position1)/(total

sequencing depth at position 1); 3) for a given F1, multiply the proportion of each allele by the

chromosome-specific somy to get the copy number estimate for each parent. For the backcross

hybrids analysis, the same procedure outlined above was used, and recombination breakpoints

865 were identified as regions where parental copy number along a chromosome shifted by >=0.8 (to

866 allow for 20% variance in absolute copy number estimation) across >=10 kilobases. All resulting

867 copy number breakpoint calls were visually confirmed (e.g., Fig. 4d). The script used to execute

868 the steps above is available here: https://github.com/jlac/Leishmania_allelic_copy_number .

870 **Scanning Electron Microscopy (SEM)**

871 Leishmania mating clumps were fixed in 2.5% Glutaraldehyde in 0.1M Sodium Cacodylate Buffer,

872 pH 7.4 (Electron Microscopy Sciences, 15960). Post fixation, the clumps were rinsed in 0.1M

873 sodium cacodylate buffer and fixed with 1% OsO4 reduced with potassium ferrocyanide in buffer.

874 After a couple of washes in water, the clumps were dehydrated in 100% ethanol and dried using

875 multiple exchanges of hexamethyldisilazane (HMDS). On the following day, clumps were

876 mounted on SEM pins using carbon adhesive tabs. Gentle pressure was applied to split open

877 clumps to reveal their content, sputter coated with 5nm of Iridium using an EMS 300TD Quorum

878 sputter coater, and imaged on a SU 8000 SEM.

880 **Transmission Electron Microscopy (TEM)**

881 *Leishmania* mating clumps were fixed as per SEM protocol. After postfixing the organoids with

882 1% reduced OsO_4 and dehydrating with 100% ethanol, the clumps were infiltrated and embedded

883 with 100% Spurr's resin in a flat mold. The flat mold with clumps were excised and mounted on

884 BEEM™ capsules for trimming. Ultrathin sections were collected using a Leica UC7

885 ultramicrotome and imaged on an 80kV Hitachi 7800 TEM using an AMT™ bottom mount

886 camera.

Statistical analysis

All data manipulation and statistical analysis was performed in R (v. 4.1.1) with custom code written in Rstudio (v. 1.4.1717). Functions from the following packages were used in our scripts: abind[51], chron[52], data.table[53], e1071[54], emmeans[55], FedData[56], gsubfn[57], gt[58], ijtiff[59], janitor[60], lubridate[61], MASS[62], plyr[63], rcompanion[64], readxl[65], reshape[66], R.utils[67], rlist[68], Rmisc[69], rstatix[70], stats[71], stringr[72], tidyverse[73] (contains: dplyr, ggplot2, forcats, tibble, readr, stringr, tidyr, purr), utils[71], WRS2[74], xlsx[75]. To compare hybrid formation success between the different experimental conditions (BM1, BM2/-IgM, BM2/+IgM) with the different parental line combinations (1x2, 1x3 & 2x3) as detailed in figures 3d and 4c, we applied an expanded Fisher's exact test that could cope with r x n matrices over a chi-square test due to the occurrence of low expected values (<5). Since our data matrices exceed the standard 2x2 format, we followed the Fisher's exact test with a pairwise test with the "pairwsieNominalIndependence" function from the R package rcompanion[64] for 2-dimensional matrices, in which at least one dimension exceeded 2 levels. The output of this test was converted into a letter code using the "cldList" function from the R package rcompanion[64] for improved test output representation; equal letters indicate no statistically significant difference, while different letters indicate a statistically significant difference (<0.05).

Methods references

34   Oliveira, F. *et al.* A sand fly salivary protein vaccine shows efficacy against vector-transmitted cutaneous leishmaniasis in nonhuman primates. *Sci Transl Med* **7**, 290ra290, doi:10.1126/scitranslmed.aaa3043 (2015).
35   Aslan, H. *et al.* A new model of progressive visceral leishmaniasis in hamsters by natural transmission via bites of vector sand flies. *J Infect Dis* **207**, 1328-1338, doi:10.1093/infdis/jis932 (2013).
36   Sacks, D. L. & Perkins, P. V. Identification of an infective stage of Leishmania promastigotes. *Science* **223**, 1417-1419 (1984).

37 Guo, H. *et al.* Genetic metabolic complementation establishes a requirement for GDP-
fucose in Leishmania. *J Biol Chem* **292**, 10696-10708, doi:10.1074/jbc.M117.778480
(2017).
38 Vickers, T. J., Murta, S. M., Mandell, M. A. & Beverley, S. M. The enzymes of the 10-
formyl-tetrahydrofolate synthetic pathway are found exclusively in the cytosol of the
trypanosomatid parasite Leishmania major. *Mol Biochem Parasit* **166**, 142-152,
doi:10.1016/j.molbiopara.2009.03.009 (2009).
39 Afonso, L. C. & Scott, P. Immune responses associated with susceptibility of C57BL/10
mice to Leishmania amazonensis. *Infect Immun* **61**, 2952-2959,
doi:10.1128/iai.61.7.2952-2959.1993 (1993).
40 Cecilio, P. *et al.* Exploring Lutzomyia longipalpis Sand Fly Vector Competence for
Leishmania major Parasites. *J Infect Dis* **222**, 1199-1203, doi:10.1093/infdis/jiaa203
(2020).
41 Rogers, M. B. *et al.* Genomic confirmation of hybridisation and recent inbreeding in a
vector-isolated Leishmania population. *PLoS Genet* **10**, e1004092,
doi:10.1371/journal.pgen.1004092 (2014).
42 Dostalova, A. & Volf, P. Leishmania development in sand flies: parasite-vector
interactions overview. *Parasit Vectors* **5**, 276, doi:10.1186/1756-3305-5-276 (2012).
43 Serafim, T. D. *et al.* Sequential blood meals promote Leishmania replication and reverse
metacyclogenesis augmenting vector infectivity. *Nat Microbiol* **3**, 548-555,
doi:10.1038/s41564-018-0125-7 (2018).
44 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat
Methods* **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
45 Zaqout, S., Becker, L. L. & Kaindl, A. M. Immunofluorescence Staining of Paraffin
Sections Step by Step. *Front Neuroanat* **14**, 582218, doi:10.3389/fnana.2020.582218
(2020).
46 Okonechnikov, K., Conesa, A. & Garcia-Alcalde, F. Qualimap 2: advanced multi-sample
quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292-294,
doi:10.1093/bioinformatics/btv566 (2016).
47 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).
48 Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997 (2013).
49 Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural variant
read extraction. *Bioinformatics* **30**, 2503-2505, doi:10.1093/bioinformatics/btu314
(2014).
50 Shaik, J. S., Dobson, D. E., Sacks, D. L. & Beverley, S. M. Leishmania Sexual
Reproductive Strategies as Resolved through Computational Methods Designed for
Aneuploid Genomes. *Genes (Basel)* **12**, doi:10.3390/genes12020167 (2021).
51 Plate, T. & Heiberger, R. abind: Combine Multidimensional Arrays. R package version
1.4-5. *https://CRAN.R-project.org/package=abind* (2016).
52 James, D. & Hornik, K. chron: Chronological Objects which Can Handle Dates and
Times. R package version 2.3-56. *https://cran.r-
project.org/web/packages/chron/index.html* (2020).
53 Dowle, M. & Srinivasan, A. data.table: Extension of \`data.frame\`. R package version
1.14.0. *https://CRAN.R-project.org/package=data.table* (2021).

961	54	Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A. & Leisch, F. e1071: Misc
962		Functions of the Department of Statistics, Probability Theory Group (Formerly: E1071),
963		TU Wien. R package version 1.7-8. *https://CRAN.R-project.org/package=e1071* (2021).
964	55	Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
965		version 1.6.2-1. *https://CRAN.R-project.org/package=emmeans* (2021).
966	56	Bocinsky, R. K. FedData: Functions to Automate Downloading Geospatial Data
967		Available from Several Federated Data Sources. R package version 2.5.7.
968		*https://CRAN.R-project.org/package=FedData* (2019).
969	57	Grothendieck, G. gsubfn: Utilities for Strings and Function Arguments. R package
970		version 0.7. *https://CRAN.R-project.org/package=gsubfn* (2018).
971	58	Iannone, R., Cheng, J. & Schloerke, B. gt: Easily Create Presentation-Ready Display
972		Tables. R package version 0.3.1. *https://CRAN.R-project.org/package=gt* (2021).
973	59	Nolan, R. & Padilla-Parra, S. ijtiff: An R package providing TIFF I/O for ImageJ users.
974		*Journal of Open Source Software.* **3(23)**, 663, doi:10.21105/joss.00633 (2018).
975	60	Firke, S. janitor: Simple Tools for Examining and Cleaning Dirty Data. R package
976		version 2.1.0. *https://CRAN.R-project.org/package=janitor* (2021).
977	61	Grolemund, G. & Wickham, H. Dates and Times Made Easy with lubridate. *Journal of*
978		*Statistical Software, 40(3), 1-25. https://www.jstatsoft.org/v40/i03/* (2011).
979	62	Venables, W. N. & Ripley, B. D. Modern Applied Statistics with S. Fourth Edition. .
980		*Springer, New York. ISBN 0-387-95457-0* (2002).
981	63	Wickham, H. The Split-Apply-Combine Strategy for Data Analysis. *Journal of Statistical*
982		*Software, 40(1), 1-29. http://www.jstatsoft.org/v40/i01/.* (2011).
983	64	Mangiafico, S. rcompanion: Functions to Support Extension Education Program
984		Evaluation. R package version 2.4.1. *https://CRAN.R-project.org/package=rcompanion*
985		(2021).
986	65	Wickham, H. & Bryan, B. readxl: Read Excel Files. R package version 1.3.1.
987		*https://CRAN.R-project.org/package=readxl* (2019).
988	66	Wickham, H. Reshaping Data with the reshape Package. *Journal of Statistical Software*
989		**21**, 1 - 20, doi:10.18637/jss.v021.i12 (2007).
990	67	Bengtsson, H. R.utils: Various Programming Utilities. R package version 2.10.1.
991		*https://CRAN.R-project.org/package=R.utils* (2020).
992	68	Ren, K. rlist: A Toolbox for Non-Tabular Data Manipulation. R package version 0.4.6.1.
993		*https://CRAN.R-project.org/package=rlist* (2016).
994	69	Hope, R. M. Rmisc: Rmisc: Ryan Miscellaneous. R package version 1.5. *https://CRAN.R-*
995		*project.org/package=Rmisc* (2013).
996	70	Kassambara, A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package
997		version 0.7.0. *https://CRAN.R-project.org/package=rstatix* (2021).
998	71	Team, R. C. R: A language and environment for statistical computing. R Foundation for
999		Statistical Computing, Vienna, Austria. *https://www.R-project.org/* (2021).
1000	72	Wickham, H. stringr: Simple, Consistent Wrappers for Common String Operations. R
1001		package version 1.4.0. *https://CRAN.R-project.org/package=stringr* (2019).
1002	73	Wickham, H. *et al.* Welcome to the tidyverse. *Journal of Open Source Software, 4(43),*
1003		*1686, https://doi.org/10.21105/joss.01686* (2019).
1004	74	Mair, P. & Wilcox, R. Robust statistical methods in R using the WRS2 package.
1005		*Behavior Research Methods* **52**, 464-488, doi:10.3758/s13428-019-01246-w (2020).

75 Dragulescu, A. & Arendt, C. xlsx: Read, Write, Format Excel 2007 and Excel 97/2000/XP/2003 Files. R package version 0.6.5. *https://CRAN.R-project.org/package=xlsx* (2020).

End notes:

Acknowledgments: We would like to thank Yvonne Rangel Gonzalez, Roberto Salas Carrillo, and Brittany Mills from VMBS, NIAID for technical support; Serena Doe and Diane Akueson from VMBS, NIAID for sand fly insectary support, Dr. Aline de Souza for critical reading of the manuscript. The authors thank Glenn A. Nardone, Motoshi Suzuki, and Lisa R. Olano of the Research Technologies Branch, NIAID, for mass spectrometry analysis. This research was supported by the Intramural Research Program of the NIH, National Institute of Allergy and Infectious Diseases, and NIH grant R01-AI29646 to S.M.B.

Author contribution: T.D.S. developed the hypothesis. T.D.S., S.K., J.G.V, S.M.B., I.V.C.A., C.B.M., J.A. and F.O. contributed to experimental design. E.I. planned the experiments. T.D.S., E.I., J.D., A.B.F.B., J.A., M.D., V.N., M.S., P.C. and T.W. performed the experiments. T.D.S., J.M.C.R, J.S.P.D, J.L., A.B.F.B., J.G.V., S.K., S.M.B. and E.I. Analyzed the data. C.M. performed sand fly insectary work. J.G.V., S.K. and S.M.B. supervised the project. All authors wrote the manuscript.

Competing interests: The authors declare no competing interests.

Data availability: All data is present in the main text or the supplementary materials. Codes generated on this work used for copy number bioinformatics analysis on https://github.com/jlac/Leishmania_allelic_copy_number. Whole-genome sequencing raw reads files are deposited on SRA database under BioProject PRJNA848835. https://dataview.ncbi.nlm.nih.gov/object/PRJNA848835?reviewer=g3jklmr2bpu6ehm01aagjr68jv.