Title:
High overexploitation risk due to management shortfall in highly traded requiem sharks

Authors:
C. Samantha Sherman[1,2], Eric D. Digel[1], Patrick Zubick[1], Jonathan Eged[1], Alifa B. Haque[3,4], Jay H. Matsushiba[1], Colin A. Simpfendorfer[5], Glenn Sant[2,6], and Nicholas K. Dulvy[1]

Addresses:
      1. Earth to Oceans Research Group, Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada
      2. TRAFFIC International, Cambridge, United Kingdom.
      3. Nature-based Solutions Initiative, Department of Zoology, University of Oxford, UK
      4. Department of Zoology, University of Dhaka, Bangladesh
      5. Institute of Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia
      6. Australian National Centre for Ocean Resources and Security, University of Wollongong, NSW, Australia

Author e-mails:
C. Samantha Sherman (sammsherman27@gmail.com), Eric D. Digel (eric_digel@sfu.ca), Patrick Zubick (zubickp@gmail.com), Jonathan Eged (joneged@gmail.com), Alifa B. Haque (alifa.haque@du.ac.bd), Jay H. Matsushiba (jay.matsushiba@gmail.com), Colin A. Simpfendorfer (colin.simpfendorfer@utas.edu.au), Glenn Sant (glenn.sant@traffic.org), Nicholas K. Dulvy (dulvy@sfu.ca)

Running title: Global management shortfall for sharks

Keywords: Carcharhinidae, CITES, fisheries management, international policy, international trade, Shark conservation, shark fin trade, shark meat

Article Type: Letters
Number of words in Abstract: 140
Number of words in Manuscript: 3,018
Number of references: 42
Number of figures and tables: 4 figures, 0 tables (4 total)

Corresponding author full address:
C. Samantha Sherman
sammsherman27@gmail.com
+1 236-688-6457
Simon Fraser University
8888 University Drive
Burnaby, British Columbia
V5A   1S6

ABSTRACT

Much of the international trade in fins (and likely meat too) is derived from requiem sharks (family Carcharhinidae), yet trade in only two of the 56 species is currently regulated. Here, we quantify catch, trade, and the shortfall in fisheries management (M-Risk) for all 56 requiem shark species based on 831 assessments across 30 countries and four Regional Fisheries Management Organisations (RFMOs). Requiem sharks comprise over half (60%) of the annual reported global chondrichthyan catch with most species (86%) identified in the international fin trade. Requiem sharks are inadequately managed by fisheries, with an average M-Risk of half (50%) of an ideal score, consequently 70% of species are threatened globally. The high catch and trade volume and shortfall in management of these iconic species requires a global integrated improvement in fisheries management, supported by regulating international trade to sustainable levels.

1 | INTRODUCTION

While there are promising signs that conservation is working in the oceans, overfishing remains the principal cause of biodiversity loss (Duarte et al. 2020). Fisheries targeting bony fishes are increasingly being brought into sustainability. For example, sustainable fisheries are found in regions and nations with high capacity for management like Regional Fisheries Management Organisations (RFMOs) and developed countries in North America, Europe as well as South Africa, Australia, and New Zealand (Simpfendorfer and Dulvy 2017, Hilborn et al. 2020). Yet, other species taken in these fisheries, notably sharks and rays, continue to decline in part because of their more sensitive life histories (Myers and Worm 2005), but also because of a lack of management (Juan-Jordá et al. 2017, Fordham et al. 2022). Shark, ray, and chimaera (Class Chondrichthyes, hereafter 'sharks') catch is poorly monitored globally, with only one-third of reported catch identified beyond Family level (FAO 2022). Yet locally, sharks provide significant income for small-scale fishers and individual specimens are retained because of the extremely high value of their body parts, such as fins, jaws, skins, and liver oil in the international trade markets (McClenachan et al. 2016, Booth et al. 2021).

The latest 2021 global Red List assessment of sharks reveals that one-third (37.5%) of all species are threatened with an elevated risk of extinction (Dulvy et al. 2021). Notably, many groups, such as tropical sharks and rays, are at very high risk of extinction (Dulvy et al. 2021). A key way to halt and reverse the declines of sharks is through quantifying and improving national fisheries management. Here, we focus on understanding national fisheries management of sharks that are highly threatened, in part due to international trade demand for meat and fins (Dent and Clarke 2015, Niedermüller et al. 2021).

The requiem sharks (family Carcharhinidae) are a recent radiation found throughout tropical and subtropical waters, which have historically dominated the shark catches of coastal fisheries (Lam and Sadovy de Mitcheson 2011). Indeed, this is one of seven priority shark taxa based on the high volume of fisheries catches taken in poorly documented and poorly regulated tropical multispecies fisheries (Dulvy et al. 2017). Further, many species of the family are highly prevalent in the international fin trade (Fields et al. 2017, Cardeñosa et al. 2020).

Here, we ask whether fisheries management is adequate to halt declines and improve the
status of requiem sharks. We estimate the risk due to overexploitation based on the current
state of global fisheries management for all 56 species from the family Carcharhinidae using a
novel rapid assessment of management risk (M-Risk)(Lack et al. 2014, Sherman et al. 2022).
Specifically, we ask three questions: (1) how prevalent are requiem sharks in global catch and
international trade?, (2) are requiem sharks adequately managed across their geographic
range? and which (if any) species are adequately managed?, and (3) which forms of
management are most prevalent and which are lacking?
117

2 | METHODS

2.1 | Selection of management units

Country management units were selected based on their proportional contribution to global shark catch, (2010–2019, inclusive) (FAO 2022). We included the 21 countries that cumulatively report >80% of global Chondrichthyan catch in addition to nine countries with high catch in each FAO fishing region to ensure global representation. In addition to countries, all four tuna Regional Fisheries Management Organisations (RFMOs) were included as management units due to the high volume of shark catch in Areas Beyond National Jurisdictions (total management units = 34) (Tolotti et al. 2015, Murua et al. 2021). Our references to 'management units' refer to the jurisdictions and not specific stocks.

2.2 | M-Risk: Management risk assessment

Species assessments were completed by scoring management risk against 21 measurable attributes, as fully explained in Sherman *et al.* (2022) and described in **Supplemental Material 1**. The 21 Attributes were split into five classes, three scored for all management units (Management System [n=5 Attributes], Fishing Practises & Catch [n=5], Compliance, Monitoring & Enforcement [n=5]), and two that were specific to either a country (n=4) or RFMO (n=2).

2.3 | Scoring of Attributes

All attributes were scored in an ordinal manner with a narrow range of scores to ensure consistency (i.e., 0–3, 0–4, or 0–5) such that higher scores indicated management with a higher likelihood of sustainable outcomes (for full details see **Supplemental Material 1**). Assessments were completed through exhaustive internet searches and national fisheries department websites. Where English was not the official language, searches were completed in the official language and documents were translated using Google translate. To accommodate the potential for ambiguous translations, points were generously awarded. However, if no information for an attribute was found, a precautionary score of zero was given. The final 'Management Score' was expressed as a percentage of the total possible points. This score indicates progress towards the ideal management, therefore, a score greater than 75% could be considered a well-managed fishery for sharks.

2.4 | Additional Catch, Trade, and Species Data

Identification of Carcharhinidae species in international trade was obtained through a thorough literature search (**Supplementary Material 2**). To determine threat level of all Carcharhinidae, the IUCN Red List Status of all species and species distribution maps were sourced after the 2021-03 update of the IUCN Red List website (www.iucnredlist.org)(Dulvy et al. 2021). Shark catch from all countries and RFMOs was sourced from FishStatJ v4.02.07 (FAO 2022), and the Sea Around Us (www.seaaroundus.org)(Pauly et al. 2021). A generalised linear model (GLM) comparing volume of shark catch and mean management score was completed (**Supplementary Material 3**).

3| RESULTS

3.1 | What is the scale of catch and trade of requiem sharks?

Based on the FAO global catch data from 2010–2019, inclusive, 65.5% of Chondrichthyan catch was reported only as Class (Chondrichthyes), subclass (Elasmobranchii), or superorder (Selachimorpha). Carcharhinidae species accounted for 59.6% of shark catch reported below Order, and 44.2% of reported coastal shark catch (excluding Blue Shark, *Prionace glauca*; 438,799 and 325,418 mt per year, respectively)(FAO 2022). Over three-quarters of Carcharhinidae species (minimum 80.4%, n = 45 species, to 85.7%, n = 48) have been documented in the international fin trade either through market sampling or shipment seizures (**Supplementary Material 2**). Based on the under-reporting of shark catch to the FAO by up to four times (Clarke et al. 2006), the true volume of Carcharhinidae catch is likely closer to 1,755,200 metric tonnes per year and possibly higher if the broad catch reporting includes Carcharhinids.

3.2 | Are requiem sharks adequately managed across their geographic range?

Requiem sharks are inadequately managed worldwide, both by nations and RFMOs, and are globally distributed with hotspots in northern Australia and Southeast Asia (**Figure 1a**). Across 831 M-Risk assessments from 30 countries spanning all six inhabited continents and four RFMO fishing grounds, the average management risk was high, with only half of the ideal management in place for requiem sharks (50.0% ± 0.6, n = 831). Regionally, the management shortfall for requiem sharks was greatest in Africa (mean management score 36.3% ± 1.5, n = 88) and Asia (38.6% ± 0.4, n = 332; **Figures 1b and 2a**). Both regions were much lower than the remaining regions and RFMO average: Oceania (67.1% ± 1.2, n = 122), Europe (62.9% ± 1.4, n = 21), North America (60.9% ± 1.1, n = 130), RFMOs (60.5% ± 0.7, n = 63), and South America (56.7% ± 0.7, n = 75) (**Figures 1b,c and 2**).

Some of the biggest fishing nations of the world have the lowest fisheries management scores including Nigeria (21.9% ± 0.0, n = 12), India (30.4% ± 0.4, n = 25), Yemen (31.5% ± 0.3, n = 20), and Iran (36.5% ± 0.6, n = 17) (**Figures 1b and 2b,c**). These countries are all among the top 20 shark catching countries globally (Okes and Sant 2019). Much of their catch is exported through non-standardized commodity codes, masking catch details (Shea and To 2017). The countries with the highest scores, and assessments on more than three species, were the USA

(76.8% ± 1.2, n = 18), Australia (74.1% ± 0.5, n = 29), Spain (65.7% ± 2.2, n = 7), and Argentina
(61.0% ± 1.0, n = 5), respectively (**Figures 1b and 2b,c**). Interestingly, these countries
represent four different continents, indicating there are 'pockets' with better management scores
(above 60%) for requiem sharks. Three of these countries (all except Australia) also rank
among the top 20 shark catching countries globally (Okes and Sant 2019), indicating catch
quantity is not a direct result of management. Indeed, a GLM showed no significant correlation
between volume of shark catch in each management unit and their respective management
scores (t = 1.46, df = 32, p = 0.153)(**Figure 2c**).
196

3.3 | Which species are adequately managed?

Strictly speaking, any species with an M-Risk score of less than 100% is likely to be
undermanaged. By that criterion–Across a total of 831 assessments–all species are
undermanaged. The species with the highest mean management score, assessed in three or
more countries, are Blacknose Shark (*C. acronotus*: 62.4% ± 7.0, n = 4), Atlantic Sharpnose
Shark (*Rhizoprionodon terraenovae:* 61.1% ± 7.4, n = 3) and Caribbean Reef Shark (*C. perezi:*
60.3% ± 6.6, n = 3) (**Figure 3**). All three of these species have relatively small distributions
when compared to many of the requiem sharks assessed. All three species occur in the
Western Central and Southwest Atlantic Ocean. Status is a function not only of management,
but also total catch volume, therefore, in regions with higher fishing effort and catch, like the
Western Atlantic (Bell et al. 2017), even higher management scores are not currently effective
at ensuring sustainable catch levels for these species (Worm et al. 2013), hence the Blacknose
Shark and Caribbean Reef Shark are listed as Endangered-EN as per the IUCN Red List.
210

The lowest scoring species (i.e., the *least* managed), assessed in at least three countries, were
the Broadfin Shark (*Lamiopsis temminckii:* 36.1% ± 3.4, n = 3), followed by the Blackspot Shark
(*C. sealei*: 36.3% ± 4.5, n = 3), and Spadenose Shark (*Scoliodon laticaudus:* 36.8% ± 3.5, n = 3)
(**Figure 3**). The first two are listed in threatened categories (EN and Vulnerable-VU,
respectively), with the third being Near Threatened-NT, and all occur in the heavily fished
waters of the Indian Oceans (Blaber et al. 2009, Pauly et al. 2021).
217

218 The variation between the single highest and lowest management score across all management
219 units were 21.9% for all 13 species in Nigeria and 85.0% for Copper Shark (*C. brachyurus*) in
220 the USA, indicating highly variable management for species (over 60%). When considered as
221 an average management score for species, there is over a 40% difference in management of

the highest and lowest scoring requiem shark, indicating a mixture of good and bad management across a species' distribution. Species with intermediate management scores tended to be assessed in a greater number of management units. For example, the Sandbar Shark (*C. plumbeus:* 50.5% ± 3.1), Silky Shark (*C. falciformis:* 53.6% ± 2.9), and Blacktip Shark (*C. limbatus*: 48.4% ± 2.6) were assessed in 22, 27, and 28 different management units, respectively, and are all listed as threatened (EN, VU, VU, respectively) (**Figure 3**). The three highest and lowest scoring species were only assessed in three or four management units, indicating potential regional areas of management concern (i.e., Northern Indian Ocean) or hope (i.e., Western Atlantic).

3.4 | Which aspects of management are most prevalent and lacking?

Across all assessed management units, the attributes with the highest average scores were lower resolution structural attributes, related to general fisheries management, such as: if a regulatory body was present (95.7% ± 2.5), engagement with CITES (93.3% ± 2.8) or Ecosystem Based Fisheries Management (EBFM) for RFMOs (91.7% ± 2.9), followed by IUU management (84.8% ± 4.7) (**Figure 4a**). Attributes with lower average scores were the harder to implement, species-specific fisheries operational attributes including: understanding of the species' stock status (18.3% ± 4.9), species-specific compliance measures to reduce fishing mortality (21.0% ± 3.8), and taxonomic resolution of landing limits in place or if any limits existed (21.2% ± 4.5) (**Figure 4a**). These attribute scores indicate that while countries and RFMOs have the foundations in-place to incorporate management, this is not yet being directed towards significant catches of threatened species, like the requiem sharks. The pattern of attribute scoring of species was similar to that for management units, with greater scores for structural attributes and lower scores for fisheries operational attributes (**Figure 4b**). For some species, like the CITES listed Oceanic Whitetip and Silky Sharks (*C. longimanus* and *C. falciformis*), landing limits existed more often, and at higher resolution than the average requiem shark (65.6% ± 8.5 and 49.6% ± 9.5, respectively, vs. 20.0% ± 2.9 average of all requiem sharks).

4 | DISCUSSION

We have three key findings. First, the global catch of requiem sharks is almost two-thirds of the global reported catch of all sharks reported to a higher resolution than Order with a volume likely over 1.75 million metric tonnes per year. Requiem sharks are also highly represented in international trade, with up to 86% of species in the Carcharhinidae family identified. Second, the management risk scores for 56 species of requiem sharks from 30 nations and four RFMOs demonstrate they are undermanaged with only half of the ideal management in place across all species. M-Risk is lower in Africa and Asia and higher in Oceania, but there are better managed countries found on each continent, representing pockets of hope. Third, management deficits typically exist in species-specific regulations, while general attributes often scored higher. Taken together, these three catch, trade, and management shortfall issues have resulted in high levels of overfishing of the Carcharhinidae family and an elevated risk of extinction. We next consider the following issues (1) how to improve domestic management, (2) improving global fisheries management, (3) the emerging opportunity that trade regulation presents, and (4) the benefits of trade regulation of requiem sharks.

Globally, requiem sharks have minimal species-specific fisheries regulations, including a lack of landing limits, and catch reporting generalised to the genus, class, or often not reported at all. The recognition that many shark species, including requiem sharks, are threatened by overfishing, has resulted in increasing national and international efforts to improve their management. However, there is little evidence that these efforts have been able to address the issue in a significant way (Davidson and Dulvy 2017). In addition to a lack of domestic measures, documentation meant to ensure internationally traded species are caught in a sustainable manner, like catch documentation schemes, are not used for requiem sharks (Mundy-Taylor et al. 2014). Domestic and international regulations have increased in the past decade for shark finning, but appear to have had little effect on reducing the volumes of fins traded (Lawson and Fordham 2018, Ferretti et al. 2020). In addition to the fin trade, there has been a recent increase in the international shark meat trade (Dent and Clarke 2015, Liu et al. 2021). This may be adding to demand and, therefore, incentive for fishers to land sharks, thus additional constraints are urgently needed to ensure catches are sustainable. Management units with higher shark catch volumes are not necessarily better managed, but can be used to

provide priority areas for management improvement (i.e., Indonesia, which has high shark catch
but a lower management score).

Globally widespread species of sharks are subject to fragmented management, requiring a
coherent response (Dulvy et al. 2017). The wide-ranging requiem sharks suffer from a
patchwork of management and might have lower extinction risk only if a significant part of their
range lies within the waters of better managed countries like the USA and Australia, e.g.,
Blacktip Reef Shark (*C. melanopterus*), which is well-managed in Australia, but less so
elsewhere in its range (**Figures 1 and 2**). Smaller range species may be at very high risk of
extinction if a significant fraction of their range lies in countries with moderate to high
management risk, such as the Broadfin Shark. The broad distributions of many species, and
large volume of requiem sharks in global trade means the need for well-managed fisheries is
pertinent to all range states. Traded shark products often travel through several different
countries being aggregated, processed and then reimported for sale to the general public. This
results in issues of traceability and species identification (Haque and Spaet 2021, Niedermüller
et al. 2021). The onus should be on both importers and exporters to ensure traded products are
fished sustainably and legally acquired. Introducing national and/or fishery landing limits for
requiem sharks, although just a domestic measure, would have a further reaching benefit, due
to the volume of this species group in international trade. These types of measures can only be
regulated at a domestic scale, making them difficult to implement due to the area covered by
fishing operations. However, this must be completed coherently in all countries to ensure future
sustainability of requiem shark catches. There is an immediate need for global cooperation in
the monitoring, sustainable catch, and trade of these species, which can be achieved through
international agreements such as CITES, CMS, and adherence to the FAO Guidelines for
Responsible Fish Trade.

Listing species on Appendix II of CITES ensures that specimens cannot be traded unless they
are legally acquired from a sustainable fishery (i.e., have a positive Non-Detriment Finding
(NDF)), and both exporting and importing countries have permits (Fernando et al. 2022).
However, a lack of global standards and requirements for NDFs, in addition to the lack of
consideration of cumulative effects across a species' full range, may reduce the usefulness of
NDFs. The development of tools to assess whether a fishery is detrimental or otherwise has
driven investment in monitoring and management of CITES Appendix II listed sharks (Martin
2007). There are promising signs of progress, even though measuring effectiveness is

challenging and the listings have been comparatively recent relative to the long generation
lengths and recovery times of sharks (Friedman et al. 2018). Appendix II CITES listings have
led to detectable progress in governance, particularly in major shark fishing nations like
Indonesia and Malaysia (Friedman et al. 2018). In Indonesia, implementation of listings resulted
in reduced catch and trade of manta rays (*Mobula alfredi* and *M. birostris*)(Booth et al. 2020).
Here, we identify higher scores in species-specific attributes, like landing limits, for the CITES-
listed Oceanic Whitetip and Silky Sharks (*C. longimanus* and *C. falciformis*). For less easily
identifiable species, like most other requiem sharks, their trade from Indonesia, the top shark
catching country in the world, is reported to Family, combining all Carcharhinidae, which
contribute 51% to shark production (Prasetyo et al. 2021). This broad level of reporting may be
masking serial depletions. Additionally, difficulties distinguishing members of the Carcharhinidae
family may mean customs officials are not able to identify listed Carcharhinidae species in
shipments (Partin et al. 2022). However, both the obligations, created through CITES listing, to
have species-specific positive NDFs, and to report species-specific trade data would greatly
improve our ability to discern species-specific fishing mortality using trade volume data
(Janssen and Leupen 2019). Evidence of improved species status after CITES listings is
minimal, but has been observed for mammals (Mialon et al. 2022). However, in some cases,
anticipation of CITES listing and ratification may spur traders to increase trade volume prior to
the need for an NDF (Mialon et al. 2022). Despite this initial increase, regulations moving
forward would allow for species control and recovery, therefore, long-term sustainability is only
possible through immediate action.

335
336

Policy relevance
Requiem sharks are heavily fished and traded, with at least 44, and up to 48 species of the
family Carcharhinidae documented in international markets and fin seizures (**Supp. Data S2**).
Most requiem sharks are threatened and indeed the Carcharhinidae family is in the top ten most
threatened families of sharks and rays, with over two-thirds (69.6%, 39 of 56) of species
threatened (Dulvy et al. 2021). Here, we show that this high level of threat, from overfishing and
unregulated international trade, results from a shortfall in national and international
management. We conclude that listing requiem sharks on CITES Appendix II provides a
mechanism for tackling the management deficits for requiem sharks.

346
347

ACKNOWLEDGEMENTS

The authors thank Brittany Finucci for her comments on study design. We also thank the
following people for help finding or understanding legislation in various countries: Maite Pons
(Argentina), David Kulka (Canada), Sushmita Mukherji and Zoya Tyabji (India), Aiko Matsushiba
(Japan), Sara A.A. Al Mabruk (Libya), Gonzo Araujo and Cat McCann (Malaysia), Brittany
Finucci (New Zealand), Leontine Baje and Michael Grant (Papua New Guinea), Andrew Bickell
(Sri Lanka), Crystal McRae (Taiwan), and Tobey Curtis and John Carlson (USA). This work was
supported by a grant to GS from the Shark Conservation Fund, a philanthropic collaborative that
pools expertise and resources to meet the threats facing the world's sharks and rays. The Shark
Conservation Fund is a project of Rockefeller Philanthropy Advisors. NKD was supported by the
Discovery and Accelerator grants from Natural Science and Engineering Research Council and
the Canada Research Chair program. The authors have no conflicts of interest to declare.
362

REFERENCES

Bell, J. D., R. A. Watson, and Y. Ye. 2017. Global fishing capacity and fishing effort from 1950
to 2012. Fish and Fisheries 18:489-505.
Blaber, S. J. M., C. M. Dichmont, W. White, R. Buckworth, L. Sadiyah, B. Iskandar, S.
Nurhakim, R. Pillans, R. Andamari, Dharmadi, and Fahmi. 2009. Elasmobranchs in
southern Indonesian fisheries: the fisheries, the status of the stocks and management
options. Reviews in Fish Biology and Fisheries 19:367-391.
Booth, H., S. Pooley, T. Clements, M. I. H. Putra, W. P. Lestari, S. Lewis, L. Warwick, and E. J.
Milner-Gulland. 2020. Assessing the impact of regulations on the use and trade of
wildlife: An operational framework, with a case study on manta rays. Global Ecology and
Conservation 22:e00953.
Booth, H., D. Squires, I. Yulianto, B. Simeon, Muhsin, L. Adrianto, and E. J. Milner-Gulland.
2021. Estimating economic losses to small-scale fishers from shark conservation: A
hedonic price analysis. Conservation Science and Practice 2021:e494.
Cardeñosa, D., A. T. Fields, E. A. Babcock, S. K. H. Shea, K. A. Feldheim, and D. D. Chapman.
2020. Species composition of the largest shark fin retail-market in mainland China.
Scientific Reports 10:12914.
Clarke, S. C., M. K. McAllister, E. J. Milner-Gulland, G. P. Kirkwood, C. G. J. Michielsens, D. J.
Agnew, E. K. Piktich, H. Nakano, and M. S. Shivji. 2006. Global estimates of shark
catches using trade records from commercial markets. Ecology Letters 9:1115-1126.
Davidson, L. N. K., and N. K. Dulvy. 2017. Global marine protected areas to prevent extinctions.
Ecology and Evolution 1:1-6.
Dent, F., and S. Clarke. 2015. State of the global market for shark products. 590.
Duarte, C. M., S. Agusti, E. Barbier, G. L. Britten, J. C. Castilla, J.-P. Gattuso, R. W. Fulweiler,
T. P. Hughes, N. Knowlton, C. E. Lovelock, H. K. Lotze, M. Predragovic, E.
Poloczanska, C. Roberts, and B. Worm. 2020. Rebuilding marine life. Nature 580:39-51.

Dulvy, N. K., N. Pacoureau, C. L. Rigby, R. A. Pollom, R. W. Jabado, D. A. Ebert, B. Finucci, C.
M. Pollock, J. Cheok, D. H. Derrick, K. B. Herman, C. S. Sherman, W. J. VanderWright,
J. M. Lawson, R. H. L. Walls, J. K. Carlson, P. Charvet, K. K. Bineesh, D. Fernando, G.
M. Ralph, J. H. Matsushiba, C. Hilton-Taylor, S. V. Fordham, and C. A. Simpfendorfer.
2021. Overfishing drives over one-third of all sharks and rays toward a global extinction
crisis. Current Biology 31:1-15.
Dulvy, N. K., C. A. Simpfendorfer, L. N. K. Davidson, S. V. Fordham, A. Bräutigam, G. Sant, and
D. J. Welch. 2017. Challenges and priorities in shark and ray conservation. Current
Biology 27:R565-R572.
FAO. 2022. FishStatJ.in T. Berger, F. Sibeni, and F. Calderini, editors. FAO, Rome.
Fernando, D., C. Rigby, and G. Sant. 2022. The Convention on International Trade in
Endangered Species of Wild Fauna and Flora (CITES) and Sharks. Pages 12-21 Shark
News: Newsletter of the IUCN SSC Shark Specialist Group. IUCN SSC Shark Specialist
Group (SSG), Dubai, United Arab Emirates.
Ferretti, F., D. M. P. Jacoby, M. O. Pfleger, T. D. White, F. Dent, F. Micheli, A. A. Rosenberg, L.
B. Crowder, and B. A. Block. 2020. Shark fin trade bans and sustainable shark fisheries.
Conservation Letters 13:e12708.
Fields, A. T., G. A. Fischer, S. K. H. Shea, H. Zhang, D. L. Abercrombie, K. A. Feldheim, E. A.
Babcock, and D. D. Chapman. 2017. Species composition of the international shark fin
trade assessed through a retail-market survey in Hong Kong. Conservation Biology
32:376-389.
Fordham, S. V., J. M. Lawson, O. Koubrak, and M. R. Cronin. 2022. Elasmobranch
Conservation Policy: Progress and Priorities. Pages 689-713 in J. Carrier, C. A.
Simpfendorfer, M. R. Heithaus, and K. A. Yopak, editors. Biology of Sharks and Their
Relatives. CRC Press; Taylor and Francis.
Friedman, K., S. Gabriel, O. Abe, A. A. Nuruddin, A. Ali, R. B. R. Hassan, S. X. Cadrin, A.
Cornish, T. D. Meulenaer, Dharmadi, Fahmi, L. H. T. Anh, D. Kachelriess, L. Kissol Jr.,
T. Krajangdara, A. R. Wahab, W. Tanoue, C. Tharith, F. Torres Jr., W. Wanchana, S.
Win, K. Yokawa, and Y. Ye. 2018. Examining the impact of CITES listing of sharks and
rays in Southeast Asian fisheries. Fish and Fisheries 19:662-676.
Haque, A. B., and J. L. Y. Spaet. 2021. Trade in threatened elasmobranchs in the Bay of
Bengal, Bangladesh. Fisheries Research 243:106059.
Hilborn, R., R. O. Amoroso, C. M. Anderson, J. K. Baum, T. A. Branch, C. Costello, C. L. de
Moor, A. Faraj, D. Hively, O. P. Jensen, H. Kurota, L. R. Little, P. Mace, T. McClanahan,
M. C. Melnychuk, C. Minto, G. C. Osio, A. M. Parma, M. Pons, S. Segurado, C. S.
Szuwalski, J. R. Wilson, and Y. Ye. 2020. Effective fisheries management instrumental
in improving fish stock status. Proceedings of the National Academy of Science
117:2218-2224.
Janssen, J., and B. T. C. Leupen. 2019. Traded under the radar: poor documentation of trade in
nationally-protected non-CITES species can cause fraudulent trade to go undetected.
Biodiversity and Conservation 28:2797-2804.
Juan-Jordá, M. J., H. Murua, H. Arrizabalaga, N. K. Dulvy, and V. Restrepo. 2017. Report card
on ecosystem-based fisheries management in tuna regional fisheries management
organizations. Fish and Fisheries.
Lack, M., G. Sant, M. Burgener, and N. Okes. 2014. Development of a rapid management-risk
assessment method for fish species through its application to sharks: framework and
results. TRAFFIC.
Lam, V. Y. Y., and Y. Sadovy de Mitcheson. 2011. The sharks of South East Asia – unknown,
unmonitored and unmanaged. Fish and Fisheries 12:51-74.

Lawson, J. M., and S. V. Fordham. 2018. Sharks Ahead: Realizing the Potential of the
Convention on Migratory Species to Conserve Elasmobranchs. Shark Advocates
International; The Ocean Foundation, Washington, DC, USA.
Liu, C. J. N., S. Neo, N. M. Rengifo, I. French, S. Chiang, M. Ooi, J. M. Heng, N. Soon, J. Y.
Yeo, H. Z. Bungum, K. Ota, A. A. Koul, Y. H. Poh, and B. J. Wainwright. 2021. Sharks in
hot soup: DNA barcoding of shark species traded in Singapore. Fisheries Research
241:105994.
Martin, J. B. 2007. The Price of Fame: CITES Regulation and Efforts Towards International
Protection of the Great White Shark. George Washington University.
McClenachan, L., A. B. Cooper, and N. K. Dulvy. 2016. Rethinking trade-driven extinction risk in
marine and terrestrial megafauna. Current Biology 26:1-7.
Mialon, H. M., T. Klumpp, and M. A. Williams. 2022. International trade and the survival of
mammalian and reptilian species. Science Advances 8:eabh3442.
Mundy-Taylor, V., V. Crook, S. Foster, S. Fowler, G. Sant, and J. Rice. 2014. CITES Non-
detriment findings guidance for shark species: A framework to assist Authorities in
making Non-detriment findings (NDFs) for species listed in CITES Appendix II. Germany
Federal Agency for Nature Conservation (Bundesamt fur Naturschutz, BfN); TRAFFIC.
Murua, H., S. P. Griffiths, A. J. Hobday, S. C. Clarke, E. Cortés, E. L. Gilman, J. Santiago, H.
Arrizabalaga, P. de Bruyn, J. Lopez, A. M. Aires-da-Silva, and V. Restrepo. 2021. Shark
mortality cannot be assessed by fishery overlap alone. Nature 595:E4-16.
Myers, R. A., and B. Worm. 2005. Extinction, survival or recovery of large predatory fishes.
Philosophical Transactions of the Royal Society B-Biological Sciences 360:13-20.
Niedermüller, S., G. Ainsworth, S. de Juan, R. Garcia, A. Ospina-Alvarez, P. Pita, and S.
Villasante. 2021. The shark and ray meat network: a deep dive into a global affair. WWF.
Okes, N., and G. Sant. 2019. An overview of major shark traders, catchers and species.
TRAFFIC International, Cambridge, UK.
Partin, T. G., A. J. Gonzalez, and M. K. Burnham-Curtis. 2022. Sharks on a plane: Large shark
fin seizure shines light on shark exploitation. Forensic Science International: Animals
and Environments 2:100055.
Pauly, D., D. Zeller, and M. L. D. Palomares. 2021. Sea Around Us Concepts, Design and Data.
Prasetyo, A. P., A. D. McDevitt, J. M. Murray, J. Barry, F. Agung, E. Muttaqin, and S. Mariani.
2021. Shark and ray trade in and out of Indonesia: addressing knowledge gaps on the
path to sustainability. Marine Policy 133:104714.
Shea, K. H., and A. W. L. To. 2017. From boat to bowl: patterns and dynamics of shark fin trade
in Hong Kong - implications for monitoring and management. Marine Policy 81:330-339.
Sherman, C. S., G. Sant, C. A. Simpfendorfer, E. D. Digel, P. Zubick, G. Johnson, M. Usher,
and N. K. Dulvy. 2022. M-Risk: A framework for assessing global fisheries management
efficacy of sharks, rays, and chimaeras. Fish and Fisheries.
Simpfendorfer, C. A., and N. K. Dulvy. 2017. Bright spots of sustainable shark fishing. Current
Biology 27:R83-R102.
Tolotti, M. T., J. D. Filmalter, P. Bach, P. Travassos, B. Seret, and L. Dagorn. 2015. Banning is
not enough: The complexities of oceanic shark management by tuna regional fisheries
management organizations. Global Ecology and Conservation 4:1-7.
Worm, B., B. Davis, L. Kettemer, C. A. Ward-Paige, D. Chapman, M. R. Heithaus, S. Kessel,
and S. H. Gruber. 2013. Global catches, exploitation rates, and rebuilding options for
sharks. Marine Policy 40:194-204.
485
486
Bell, J. D., R. A. Watson, and Y. Ye. 2017. Global fishing capacity and fishing effort from 1950
to 2012. Fish and Fisheries **18**:489-505.

Blaber, S. J. M., C. M. Dichmont, W. White, R. Buckworth, L. Sadiyah, B. Iskandar, S.
Nurhakim, R. Pillans, R. Andamari, Dharmadi, and Fahmi. 2009. Elasmobranchs in
southern Indonesian fisheries: the fisheries, the status of the stocks and management
options. Reviews in Fish Biology and Fisheries **19**:367-391.
Booth, H., S. Pooley, T. Clements, M. I. H. Putra, W. P. Lestari, S. Lewis, L. Warwick, and E. J.
Milner-Gulland. 2020. Assessing the impact of regulations on the use and trade of
wildlife: An operational framework, with a case study on manta rays. Global Ecology and
Conservation **22**:e00953.
Booth, H., D. Squires, I. Yulianto, B. Simeon, Muhsin, L. Adrianto, and E. J. Milner-Gulland.
2021. Estimating economic losses to small-scale fishers from shark conservation: A
hedonic price analysis. Conservation Science and Practice **2021**:e494.
Cardeñosa, D., A. T. Fields, E. A. Babcock, S. K. H. Shea, K. A. Feldheim, and D. D. Chapman.
2020. Species composition of the largest shark fin retail-market in mainland China.
Scientific Reports **10**:12914.
Clarke, S. C., M. K. McAllister, E. J. Milner-Gulland, G. P. Kirkwood, C. G. J. Michielsens, D. J.
Agnew, E. K. Piktich, H. Nakano, and M. S. Shivji. 2006. Global estimates of shark
catches using trade records from commercial markets. Ecology Letters **9**:1115-1126.
Davidson, L. N. K., and N. K. Dulvy. 2017. Global marine protected areas to prevent extinctions.
Ecology and Evolution **1**:1-6.
Dent, F., and S. Clarke. 2015. State of the global market for shark products. 590.
Duarte, C. M., S. Agusti, E. Barbier, G. L. Britten, J. C. Castilla, J.-P. Gattuso, R. W. Fulweiler,
T. P. Hughes, N. Knowlton, C. E. Lovelock, H. K. Lotze, M. Predragovic, E.
Poloczanska, C. Roberts, and B. Worm. 2020. Rebuilding marine life. Nature **580**:39-51.
Dulvy, N. K., N. Pacoureau, C. L. Rigby, R. A. Pollom, R. W. Jabado, D. A. Ebert, B. Finucci, C.
M. Pollock, J. Cheok, D. H. Derrick, K. B. Herman, C. S. Sherman, W. J. VanderWright,
J. M. Lawson, R. H. L. Walls, J. K. Carlson, P. Charvet, K. K. Bineesh, D. Fernando, G.
M. Ralph, J. H. Matsushiba, C. Hilton-Taylor, S. V. Fordham, and C. A. Simpfendorfer.
2021. Overfishing drives over one-third of all sharks and rays toward a global extinction
crisis. Current Biology **31**:1-15.
Dulvy, N. K., C. A. Simpfendorfer, L. N. K. Davidson, S. V. Fordham, A. Bräutigam, G. Sant, and
D. J. Welch. 2017. Challenges and priorities in shark and ray conservation. Current
Biology **27**:R565-R572.
FAO. 2022. FishStatJ.*in* T. Berger, F. Sibeni, and F. Calderini, editors. FAO, Rome.
Fernando, D., C. Rigby, and G. Sant. 2022. The Convention on International Trade in
Endangered Species of Wild Fauna and Flora (CITES) and Sharks. Pages 12-21 Shark
News: Newsletter of the IUCN SSC Shark Specialist Group. IUCN SSC Shark Specialist
Group (SSG), Dubai, United Arab Emirates.
Ferretti, F., D. M. P. Jacoby, M. O. Pfleger, T. D. White, F. Dent, F. Micheli, A. A. Rosenberg, L.
B. Crowder, and B. A. Block. 2020. Shark fin trade bans and sustainable shark fisheries.
Conservation Letters **13**:e12708.
Fields, A. T., G. A. Fischer, S. K. H. Shea, H. Zhang, D. L. Abercrombie, K. A. Feldheim, E. A.
Babcock, and D. D. Chapman. 2017. Species composition of the international shark fin
trade assessed through a retail-market survey in Hong Kong. Conservation Biology
32:376-389.
Fordham, S. V., J. M. Lawson, O. Koubrak, and M. R. Cronin. 2022. Elasmobranch
Conservation Policy: Progress and Priorities. Pages 689-713 *in* J. Carrier, C. A.
Simpfendorfer, M. R. Heithaus, and K. A. Yopak, editors. Biology of Sharks and Their
Relatives. CRC Press
Taylor and Francis.
Friedman, K., S. Gabriel, O. Abe, A. A. Nuruddin, A. Ali, R. B. R. Hassan, S. X. Cadrin, A.
Cornish, T. D. Meulenaer, Dharmadi, Fahmi, L. H. T. Anh, D. Kachelriess, L. Kissol Jr.,

T. Krajangdara, A. R. Wahab, W. Tanoue, C. Tharith, F. Torres Jr., W. Wanchana, S.
Win, K. Yokawa, and Y. Ye. 2018. Examining the impact of CITES listing of sharks and
rays in Southeast Asian fisheries. Fish and Fisheries **19**:662-676.

Haque, A. B., and J. L. Y. Spaet. 2021. Trade in threatened elasmobranchs in the Bay of
Bengal, Bangladesh. Fisheries Research **243**:106059.

Hilborn, R., R. O. Amoroso, C. M. Anderson, J. K. Baum, T. A. Branch, C. Costello, C. L. de
Moor, A. Faraj, D. Hively, O. P. Jensen, H. Kurota, L. R. Little, P. Mace, T. McClanahan,
M. C. Melnychuk, C. Minto, G. C. Osio, A. M. Parma, M. Pons, S. Segurado, C. S.
Szuwalski, J. R. Wilson, and Y. Ye. 2020. Effective fisheries management instrumental
in improving fish stock status. Proceedings of the National Academy of Science
117:2218-2224.

Janssen, J., and B. T. C. Leupen. 2019. Traded under the radar: poor documentation of trade in
nationally-protected non-CITES species can cause fraudulent trade to go undetected.
Biodiversity and Conservation **28**:2797-2804.

Juan-Jordá, M. J., H. Murua, H. Arrizabalaga, N. K. Dulvy, and V. Restrepo. 2017. Report card
on ecosystem-based fisheries management in tuna regional fisheries management
organizations. Fish and Fisheries.

Lack, M., G. Sant, M. Burgener, and N. Okes. 2014. Development of a rapid management-risk
assessment method for fish species through its application to sharks: framework and
results. TRAFFIC.

Lam, V. Y. Y., and Y. Sadovy de Mitcheson. 2011. The sharks of South East Asia – unknown,
unmonitored and unmanaged. Fish and Fisheries **12**:51-74.

Lawson, J. M., and S. V. Fordham. 2018. Sharks Ahead: Realizing the Potential of the
Convention on Migratory Species to Conserve Elasmobranchs. Shark Advocates
International

The Ocean Foundation, Washington, DC, USA.

Liu, C. J. N., S. Neo, N. M. Rengifo, I. French, S. Chiang, M. Ooi, J. M. Heng, N. Soon, J. Y.
Yeo, H. Z. Bungum, K. Ota, A. A. Koul, Y. H. Poh, and B. J. Wainwright. 2021. Sharks in
hot soup: DNA barcoding of shark species traded in Singapore. Fisheries Research
241:105994.

Martin, J. B. 2007. The Price of Fame: CITES Regulation and Efforts Towards International
Protection of the Great White Shark. George Washington University.

McClenachan, L., A. B. Cooper, and N. K. Dulvy. 2016. Rethinking trade-driven extinction risk in
marine and terrestrial megafauna. Current Biology **26**:1-7.

Mialon, H. M., T. Klumpp, and M. A. Williams. 2022. International trade and the survival of
mammalian and reptilian species. Science Advances **8**:eabh3442.

Mundy-Taylor, V., V. Crook, S. Foster, S. Fowler, G. Sant, and J. Rice. 2014. CITES Non-
detriment findings guidance for shark species: A framework to assist Authorities in
making Non-detriment findings (NDFs) for species listed in CITES Appendix II. Germany
Federal Agency for Nature Conservation (Bundesamt fur Naturschutz, BfN)

TRAFFIC.

Murua, H., S. P. Griffiths, A. J. Hobday, S. C. Clarke, E. Cortés, E. L. Gilman, J. Santiago, H.
Arrizabalaga, P. de Bruyn, J. Lopez, A. M. Aires-da-Silva, and V. Restrepo. 2021. Shark
mortality cannot be assessed by fishery overlap alone. Nature **595**:E4-16.

Myers, R. A., and B. Worm. 2005. Extinction, survival or recovery of large predatory fishes.
Philosophical Transactions of the Royal Society B-Biological Sciences **360**:13-20.

Niedermüller, S., G. Ainsworth, S. de Juan, R. Garcia, A. Ospina-Alvarez, P. Pita, and S.
Villasante. 2021. The shark and ray meat network: a deep dive into a global affair. WWF.

Okes, N., and G. Sant. 2019. An overview of major shark traders, catchers and species.
TRAFFIC International, Cambridge, UK.

Partin, T. G., A. J. Gonzalez, and M. K. Burnham-Curtis. 2022. Sharks on a plane: Large shark
fin seizure shines light on shark exploitation. Forensic Science International: Animals
and Environments **2**:100055.
Pauly, D., D. Zeller, and M. L. D. Palomares. 2021. Sea Around Us Concepts, Design and Data.
Prasetyo, A. P., A. D. McDevitt, J. M. Murray, J. Barry, F. Agung, E. Muttaqin, and S. Mariani.
2021. Shark and ray trade in and out of Indonesia: addressing knowledge gaps on the
path to sustainability. Marine Policy **133**:104714.
Shea, K. H., and A. W. L. To. 2017. From boat to bowl: patterns and dynamics of shark fin trade
in Hong Kong - implications for monitoring and management. Marine Policy **81**:330-339.
Sherman, C. S., G. Sant, C. A. Simpfendorfer, E. D. Digel, P. Zubick, G. Johnson, M. Usher,
and N. K. Dulvy. 2022. M-Risk: A framework for assessing global fisheries management
efficacy of sharks, rays, and chimaeras. Fish and Fisheries **23**:1383:1399.
Simpfendorfer, C. A., and N. K. Dulvy. 2017. Bright spots of sustainable shark fishing. Current
Biology **27**:R83-R102.
Tolotti, M. T., J. D. Filmalter, P. Bach, P. Travassos, B. Seret, and L. Dagorn. 2015. Banning is
not enough: The complexities of oceanic shark management by tuna regional fisheries
management organizations. Global Ecology and Conservation **4**:1-7.
Worm, B., B. Davis, L. Kettemer, C. A. Ward-Paige, D. Chapman, M. R. Heithaus, S. Kessel,
and S. H. Gruber. 2013. Global catches, exploitation rates, and rebuilding options for
sharks. Marine Policy **40**:194-204.
610

Figure 1. Global maps of (a) species richness of all requiem sharks (Family Carcharhinidae), (b) average management scores for all 56 species of requiem sharks in each country assessed and, (c) in the high-seas, as managed by the four tuna RFMOS. Average management score across different (c) continents and Regional Fisheries Management Organizations (RFMOs), (d) countries and individual RFMOs, abbreviations - ICCAT: International Commission for the Conservation of Atlantic Tunas; WCPFC: Western and Central Pacific Fisheries Commission; IOTC: Indian Ocean Tuna Commission; and IATTC: Inter-American Tropical Tuna Commission. Error bars indicate 95% confidence intervals.

Figure 2. Average management score across different (a) continents and Regional Fisheries Management Organizations (RFMOs), and countries and individual RFMOs, ordered by (b) average management score, and (c) highest volume of shark catch from top to bottom. Abbreviations - ICCAT: International Commission for the Conservation of Atlantic Tunas; WCPFC: Western and Central Pacific Fisheries Commission; IOTC: Indian Ocean Tuna

Commission; and IATTC: Inter-American Tropical Tuna Commission. Error bars indicate 95% confidence intervals.

Figure 3. Average management score for each species assessed. Error bars indicate 95% confidence intervals.

Figure 4. All average attribute scores and calculated management scores for all 56 assessed species of requiem sharks according to: (a) different management units, and (b) species. Attribute columns are arranged from highest average score to lowest average score. NA values are represented by dark gray squares.