# Sequence-sensitive elastic network captures dynamical features necessary for miR-125a maturation

Olivier Mailhot[1−4]      Vincent Frappier[5]      François Major[2,3,∗]

Rafael Najmanovich[4,∗]

[1]Department of Biochemistry and Molecular Medicine, Université de Montréal, Montreal, Canada

[2]Department of Computer Science and Operations Research, Université de Montréal, Montreal, Canada

[3]Institute for Research in Immunology and Cancer, Université de Montréal, Montreal, Canada

[4]Department of Pharmacology and Physiology, Université de Montréal, Montreal, Canada

[5]Generate Biomedecines, Cambridge, Massachusetts

[∗]To whom correspondence should be addressed.

June 9, 2022

## Abstract

The Elastic Network Contact Model (ENCoM) is a coarse-grained normal mode analysis (NMA) model unique in its all-atom sensitivity to the sequence of the studied macromolecule and thus to the effect of mutations. We adapted ENCoM to simulate the dynamics of ribonucleic acid (RNA) molecules, benchmarked its performance against other popular NMA models and used it to study the 3D structural dynamics of human microRNA miR-125a, leveraging high-throughput data on the maturation efficiency of over 26 000 sequence variants. We also introduce a novel way of using dynamical information from NMA to train multivariate linear regression models, with the purpose of highlighting the most salient contributions of dynamics to function. ENCoM has a similar performance profile on RNA than on proteins when compared to the Anisotropic Network Model (ANM), the most widely used coarse-grained NMA model; it has the advantage on predicting large-scale motions while ANM performs better on B-factors prediction. A stringent benchmark from the miR-125a maturation dataset, in which the training set contains no sequence information in common with the testing set, reveals that ENCoM alone can capture signal beyond the sequence. When training the model using all available data, the dynamical features identified as necessary for miR-125a maturation point to known patterns but also offer new insights into the biogenesis of microRNAs. Our novel approach combining NMA with multivariate linear regression is generalizable to any macromolecule for which relatively high-throughput mutational data is available.

Availability: ENCoM is open source software part of the NRGTEN package available at https://github.com/gregorpatof/nrgten_package

Contact: rafael.najmanovich@umontreal.ca, francois.major@umontreal.ca

CONTENTS

It is now well-established that ribonucleic acid (RNA) molecules possess a diverse range of functions in all domains of life and there is growing interest in understanding and characterizing these functions [1]. However, our structural knowledge of RNA is still scarcer than that of proteins, the latter being based on about two orders of magnitude more structures in the Protein Data Bank (PDB) [2]. RNAs are dynamic and thus exist in an ensemble of conformations that are intimately tied to their function [3]. The computational prediction of RNA conformational ensembles from a known or predicted three-dimensional (3D) structure is appealing. In the case of proteins, a popular method for obtaining such information is coarse-grained normal mode analysis (NMA) using elastic network models (ENMs) [4]. Many different ENMs with various application niches have been developed for proteins, while the few studies scpecifically investigating RNA [5, 6, 7] have used either the Gaussian Network Model (GNM), which predicts isotropic fluctuations [8, 9], or the Anisotropic Network Model (ANM) which predicts directional motions [10].

In the last years, we introduced the Elastic Network Contact Model (ENCoM) for NMA of proteins [11, 12, 13]. ENCoM was shown to perform significantly better than ANM for the prediction of conformational changes. More importantly, to our knowledge ENCoM is the only coarse-grained NMA model that is sensitive to the all-atom context of residues, making it sensitive to the effect of mutations even when they do not change the backbone geometry of the protein [12, 14]. In the present work, we adapted ENCoM to work on RNA molecules and benchmarked its performance on a set of 786 curated RNA structures derived from the RNA-only structures available in the PDB resolved by either nuclear magnetic resonance (NMR) or X-ray crystallography. ENCoM outperforms ANM on the prediction of conformational changes and conformational variance within NMR ensembles but performs slightly worse on experimental B-factors prediction as previously observed in the case of proteins [11].

NMA, whether coarse-grained or all-atom, is an analytical technique that can be used to explore the entire conformational space of a macromolecule at every timescale, often with an associated computational cost tremendously lower than that of, for example, molecular dynamics (MD) simulations [15]. In NMA, a molecule is represented as a system of masses connected by springs with a harmonic potential. For a system of N masses, the normal modes are the 3N eigenvectors of the Hessian matrix, each with its associated eigenvalue. The first six normal modes are trivial motions that represent the translational and rotational degrees of freedom of the system. From the seventh, the modes are ordered with increasing frequency and thus increasing energetic cost to achieve the same amount of deformation. Usually, a small set of the first non-trivial modes is sufficient to describe global motions of biological significance, for example in the case of the hinge-bending motion of the citrate synthase enzyme [16].

In the study of proteins, coarse-grained ENMs generally use one mass per amino acid, located on the $C_\alpha$ atom [4]. Efforts have been made to expand these coarse-grained models to include more information, such as the $C_\beta$ atom [17], or the combination of different levels of coarse-graining [18, 19]. However, single-mass per residue models continue to be the standard for coarse-grained NMA of proteins due to their good preservation of the slowest and most global motions. The two most widely-used models for proteins are arguably ANM [20] and GNM [5] and both use one mass per residue at the $C_\alpha$ atom by default. By contrast, when these models were applied to RNA, they performed significantly better when three masses were assigned for each nucleotide: one for the phosphate group, one for the sugar group and one for the nucleobase (Figure 1) [5, 6, 7]. This improvement can be explained in part by the higher intrinsic flexibility of RNA, having six backbone torsional angles per residue instead of the three found in proteins [21], and by the importance of base pairing in forming specific 3D architectures.

Mature microRNAs (miRs) are small non-coding single-stranded RNAs of about 22 nucleotides. They regulate gene expression by guiding RNA-Induced Silencing Complexes (RISCs)

to complementary regions within messenger RNA (mRNA) sequences, triggering silencing of these targets [22]. The canonical pathway of miR production starts with their primary transcript that forms a hairpin-loop structure of approximately 35 base-pairs (pri-miR). A crucial step is the recognition of the pri-miR byt the DROSHA/DGCR8 heterotrimer, also called Microprocessor, which cleaves the hairpin around the 11th base-pair from the bottom [23].

A single nucleotide polymorphism (SNP) in human miR-125a predisposes to aggressive types of breast cancer. One of the effects of the SNP (G22U) is to prevent cleavage by the Microprocessor [24]. The SNP does not affect the global structure of the minimum free energy (MFE) state. Moreover, we showed that chages in 2D structural dynamics of the 16 possible base pairs at the SNP position correlate well with their maturation efficiency measured from cellular luciferase assays [25].

David Bartel's group has generated high-throughput data about the maturation efficiency of upwards of 50 000 pri-miR-125a sequence variants using an enzymatic assay with purified Microprocessor [26]. We generated all pri-miR-125a variants *in silico* and studied their 3D structural dynamics with both ENCoM and ANM to test whether ENCoM's ability to predict the effect of mutations on function extends to RNA.

## 2 MATERIALS AND METHODS

### 2.1 *Elastic network models*

The ENCoM potential is based on a model called generalized Spring Tensor Model (STeM), which uses a potential function containing four terms: covalent bond stretching, angle bending, dihedral angle torsion and non-bonded (or long-range) interactions [11, 27].

$$
\begin{aligned}
V_{\mathrm{ENCoM}}(\vec{R}, \vec{R_0}) = & \sum_{\mathrm{bonds}} V_1(r, r_0) + \sum_{\mathrm{angles}} V_2(\theta, \theta_0) \\
& + \sum_{\mathrm{dihedrals}} V_3(\phi, \phi_0) + \sum_{i<j-3} V_4(r_{ij}, r_{ij_0}) \\
= & \sum_{\mathrm{bonds}} \alpha_1(r - r_0)^2 + \sum_{\mathrm{angles}} \alpha_2(\theta - \theta_0)^2 \\
+ \sum_{\mathrm{dihedrals}} & \left[ \alpha_3(1 - \cos(\phi - \phi_0)) + \frac{\alpha_3}{2}(1 - \cos 3(\phi - \phi_0)) \right] \\
+ \sum_{i<j-3} & (\beta_{ij} + \alpha_4) \left[ 5\left(\frac{r_{ij_0}}{r_{ij}}\right)^{12} - 6\left(\frac{r_{ij_0}}{r_{ij}}\right)^{10} \right]
\end{aligned}
\tag{1}
$$

In the ENCoM potential, the non-bonded interaction term is modulated according to the surface area in contact between residues by the $\beta_{ij}$ term:

$$
\beta_{ij} = \sum_k^{N_i} \sum_l^{N_j} \epsilon_{T(k)T(l)} S_{kl}
\tag{2}
$$

$\epsilon_{T(k)T(l)}$ represents the interaction between atoms of types $T(k)$ and $T(l)$ while $S_{kl}$ is the surface area in contact between the two atoms, calculated using a constrained Voronoi procedure, as described by McConkey and co-workers [28]. The atom types used are those of Sobolev and co-workers [29], which are divided in eight classes: hydrophilic, acceptor, donor, hydrophobic, aromatic, neutral, neutral-donor and neutral-acceptor. Supplementary Table 1 shows the assigned type of every atom from the four common ribonucleotides.

Each term of the ENCoM potential is modulated by a weight coefficient which together represent the parameters of the model and were explored extensively in the past. ENCoM's performance came out as very robust across a wide range of parameters [11]. For the present

study, we thus used exactly the same parameters as those published for proteins. This makes ENCoM in its present form readily applicable to proteins, RNA and RNA-protein complexes without the need for re-parameterization.

Alternatively, ANM uses a very simple potential where each pair of masses that are closer than a certain distance cutoff are considered to be interacting and are connected with a spring of uniform constant $\gamma$ [30]. The interaction distance cutoff $R_c$ (generally set to 18 Angstroms (Å) for proteins) is thus a parameter of the model, which we have extensively tested in increments of 0.5 Å for the B-factors prediction benchmark. The higher and lower bounds for the cutoffs tested were chosen to show a clear decreasing trend in either direction, confirming that the value for maximal performance has been found.

In addition to the classical cutoff-based ANM, ANMs using spring constants that are distance-dependent have been introduced [31]. We chose to also test this version of the model, called power dependence ANM (PD-ANM), as it was applied to RNA by Zimmermann and Jernigan [6]. Instead of a distance cutoff parameter, all masses are connected by springs in this model. The spring constant varies according to the inverse distance to the $x^{th}$ power, where $x$ is the power dependence, the only parameter of the model. As we did for the cutoff-based ANM (Cut-ANM), we scanned the power dependency parameter in increments of 0.5 for the B-factors benchmark, again making sure to identify a clear decreasing trend in either direction.

In the case of proteins, it is generally believed that ENMs using only one mass per residue situated on the $C_\alpha$ atom still capture the essential low-frequency motions of the molecule [4]. However, since RNA molecules are intrinsically more flexible per residue than proteins, using more masses per residue leads to an increase in the predictive power of the model [5]. For all tested elastic networks, we use a model with three masses per residue, where we assign one mass each for the sugar, base and phosphate groups positioned on the C1', C2 and P atoms, respectively, as already validated in the case of an ANM [7]. We report similar findings with increased performance of ENCoM across all benchmarks when using three masses per residue instead of one (data not shown). Since this was a pre-established fact, we use three masses per nucleotide for all the models tested thereafter in the present work. Figure 1 shows the positioning of the masses in the four standard nucleotides.

The three elastic network models tested in this study were ran using the NRGTEN Python package [32] in which they are all implemented by extending the base ENM class. This ensures there are no discrepancies in the assignment of masses or parsing of PDB files since all these steps are performed by the exact same code for all models.

2.2 *Dataset of RNA structures*

Experimental determination of RNA structure is achieved by many different techniques, of which two are by far the most common: X-ray crystallography [34] which takes a snapshot of the molecule near its lowest energy conformation and solution nuclear magnetic resonance (NMR) [35, 36] in which an ensemble of probable conformations is resolved. In the PDB as of 2022-01-08, there were 982 structures solved by X-ray crystallography and 417 NMR ensembles when restricting the search to entries containing only RNA. We used these data in our benchmarks after applying appropriate filters (described at the end of the current section) in order to compare ENCoM's performance to two popular versions of ANM in the prediction of B-factors and conformational transitions. We kept all chains present in the biological assembly for X-ray structures and all chains in the structural ensemble for NMR experiments.

Our model requires the manual assignment of atom types for every residue we want to consider so that the surface in contact term can be adequately computed. It is thus possible in theory to include every modified residue. In order to simplify the present study, we chose to restrict the model to the four standard ribonucleotides, replacing all modified ribonucleotides with their unmodified analog using the ModeRNA software [37], which contains informa-

**Figure 1: Assignment of masses on the four standard nucleotides, arranged in two base pairs extracted from a standard double helix.** Phosphate atoms are in gold, C1' carbons from the sugar group in light green and C2 carbons from the nucleobase in dark green. All three masses are included in the ENMs tested for the present work. The base pairs shown were extracted from an A-form RNA helix generated using the MC-Fold and MC-Sym pipeline [33].

tion about the 170 modifications present in the MODOMICS database [38]. The atom type assignations for these four standard ribonucleotides are given in Supplementary Table 1.

We also used ModeRNA to add missing atoms where it was the case, for example terminal phosphate groups, so that each residue was complete. This addition ensured that the assignment of the masses to the residues was standard for all residues. The structures for which ModeRNA produced an error or which produced an error for any of the models tested subsequently were removed from the analysis. Moreover, the size of the X-ray resolved RNA structures was restricted to below 300 nucleotides to reduce computational cost. In the case of the NMR resolved RNAs, no size threshold was applied as no molecules bigger than 155 residues have been elucidated this way. Since we were interested in dynamical ensembles solved by NMR, we restricted our analysis to submissions containing at least two models. The list of all PDB codes of the structures kept for the three benchmarks is given in Supplementary Table 2 (38 structures for the B-factors benchmark, 129 structures for the conformational change benchmark, 313 NMR ensembles for the structural variance benchmark). The size distribution of the structures kept for the various benchmarks is given in Supplementary Figure 3.

2.3  *Sequence clustering*

We performed complete linkage clustering [39] on the sequences from our database of structures with a distance threshold of 0.1, ensuring that all pairs of sequences within a given cluster are at least 90% similar. The similarity metric used for calculating distance is derived from the score of a Needleman-Wunsch global alignment [40] between the two sequences, with the following scoring scheme: gap penalty: -1, mismatch penalty: -1, match score: 1.

$$\text{distance}(s,t) = \frac{\text{score}(s,t)}{\min\left[\text{length}(s), \text{length}(t)\right]} \tag{3}$$

For all reported benchmark metrics, we first normalize by sequence cluster before reporting the mean metric. This ensures that molecules which have a lot of conformations sampled in the PDB do not drive the score and instead each RNA or family of RNAs has an equal weight.

## 2.4 B-factors prediction

Experimental B-factors can be predicted from calculated normal modes as mean-square fluctuations (MSF) of individual residues [10]:

$$\text{MSF}_i = \sum_{n=7}^{3N} \frac{E_{n,i,x}^2 + E_{n,i,y}^2 + E_{n,i,z}^2}{\lambda_n} \tag{4}$$

where $E_{n,i}$ represents the $xyz$ diplacement of mass $i$ in the $n^{th}$ eigenvector and $\lambda_n$ the associated eigenvalue of that eigenvector. We also define the Entropic Signature, which scales the square fluctuations by the vibrational entropy of each normal mode:

$$\text{ES}_i = \sum_{n=7}^{3N} S_{\text{vib}n} \left( E_{n,i,x}^2 + E_{n,i,y}^2 + E_{n,i,z}^2 \right) \tag{5}$$

$$S_{\text{vib}n} = \frac{\beta \nu_n}{e^{\beta \nu_n} - 1} - \ln \left( 1 - e^{-\beta \nu_n} \right) \tag{6}$$

$$\nu_n = \frac{1}{2\pi} \sqrt{\lambda_n} \tag{7}$$

where $\beta$ is a thermodynamic scaling factor which allows for varying the relative contributions of high- and low-frequency normal modes to the fluctuations of the individual masses.

The Pearson correlation coefficient is then calculated between the predicted and experimental B-factors. In addition to a resolution filter of 2.5 Å or better, we used only the structures for which no modified nucleobases were initially present and in which all residues were complete. This restriction was applied to ensure that every atom used in the prediction had a corresponding experimental B-factor, and not an extrapolated value from the rebuilding of the modified nucleobases and missing atoms by ModeRNA. The 38 remaining structures were clustered according to their sequence as described giving us 34 clusters and we computed the mean correlation normalized by cluster.

## 2.5 Conformational change prediction

The overlap between two conformations is a measure of the similarity between an eigenvector $\vec{E}_n$ predicted using the start conformation and the displacement vector $\vec{R}$ calculated between the coordinates of the target and start conformations, after both have been superimposed [16, 41].

$$\text{overlap}(\vec{E}_n, \vec{R}) = \frac{|\vec{E}_n \cdot \vec{R}|}{\|\vec{E}_n\| \|\vec{R}\|} \tag{8}$$

For each eigenvector, it has a value between 0 and 1, which describes how well we can reproduce the target conformation by deforming the start conformation along the eigenvector. The cumulative overlap (CO) is also a value between 0 and 1 and describes how well a set of orthogonal motions (eigenvectors) can collectively reproduce the target conformation from the start conformation.

$$\text{CO} = \sqrt{\sum_n \text{overlap}(\vec{E}_n, \vec{R})^2} \tag{9}$$

To ensure that the same conformation is not sampled twice, we reject pairs of conformations for which the root mean squared deviation (RMSD) of the center atoms of each mass is lower

than 2 Å. Clustering was performed as described, leaving us with 129 unique structures forming 240 pairs of conformations with RMSD greater than or equal to 2 Å, divided in 29 sequence clusters. We calculated the cumulative overlap for up to 10% of normal modes both ways for every pair and report the mean cumulative overlap normalized by cluster.

### 2.6  *Non-rotational-translational principal component analysis*

Principal component analysis (PCA) is a statistical technique that is used to transform observations of variables which may be correlated into linearly uncorrelated variables which are called principal components (PCs) [42]. PCA was used here to extract dominant motions apparent within the conformational ensembles obtained from NMR experiments. Each conformation is represented as a vector of length 3N where N is the number of masses in the system (3 per nucleotide). PCA was computed on these vectors using a singular value decomposition (SVD) algorithm [43]. The PCs obtained are analogous to normal modes in that they are the eigenvectors of the covariance matrix of the 3N coordinates from the ensemble of structures. The first PC describes the largest proportion of the variance in the ensemble, and each subsequent component captures the largest proportion of the remaining variance and is orthogonal to the preceding components. However, when there are more than two conformations in the ensemble, rotational and translational motions can be present in the principal components. To our knowledge, this has not been discussed in the context of PCA applied to ensembles of 3D conformations of biomolecules. We thus introduce a correction to obtain principal components without rotational and translational degrees of freedom. First, standard PCs are computed from the ensemble, with all conformations superimposed to the first model. Then, the first six rotational and translational normal modes from that first model are used as the starting basis for Gram-Schmidt orthonormalization of the PCs. This transformation ensures all rotational-translational motions from the PCs are removed and all relevant internal motions are maintained. However, the proportion of variance explained needs to be corrected for each PC according to the amount of rotational-translational variance initially present:

$$c_i = \frac{v_i \left(1 - \sqrt{\sum_{j=1}^{6}(\vec{PC}_i \cdot \vec{RT}_j)^2}\right)}{\sum_n c_n} \tag{10}$$

where $v_i$ is the initial proportion of variance explained by $\vec{PC}_i$, $c_i$ is the corrected proportion of variance explained, and $\vec{RT}_j$ are the six rotational-translational normal modes of the first model in the structural ensemble. The PCs are then reordered in decreasing order of corrected proportion of variance explained. We call this method nrt-PCA (non-rotational-translational PCA) and whenever we refer to PCA or PCs, it is implied that nrt-PCA is used. It ensures the accurate comparison of non-trivial normal modes computed from a representative structure with strictly internal motions apparent from the ensemble.

### 2.7  *Structural ensemble variance prediction*

The root mean square inner product (RMSIP) is a value between 0 and 1 representing how well a set of normal modes can reproduce the motions from a set of PCs:

$$\text{RMSIP} = \sqrt{\frac{1}{N} \sum_{i=1}^{N} \sum_{j=1}^{M} (\vec{E}_i \cdot \vec{PC}_j)^2} \tag{11}$$

where $N$ normal modes are compared against $M$ PCs.

The normalized cumulative overlap (NCO) between the first $N$ normal modes and the first $M$ PCs is given by:

$$\text{NCO} = \sum_{j=1}^{M} \left[ v_j \sqrt{\sum_{i=1}^{N} \text{overlap}(\vec{E}_i, \vec{\text{PC}_j})^2} \right] \tag{12}$$

where $\vec{E}_i$ is normal mode $i$, $\vec{\text{PC}}_j$ is principal component component $j$ and $v_j$ is the proportion of variance explained by component $j$. NCO ensures a value between 0 and 1 as both the normal modes and the components are orthogonal, and $v_j$ sums to 1 over all the components.

For the current study, we reported all NCOs and RMSIPs using from 1 normal mode to 10% of the 3N modes, compared against PCs explaining 99% of the observed variance in the ensemble. The normal modes were computed from the conformation of minimal energy in each NMR ensemble, which by convention is the first one in the experimenters' submission. Clustering was performed as described and we computed the mean NCO/RMSIP normalized by cluster.

### 2.8 *Dataset of miR-125a mutations*

The Fang *et al.* dataset of high-throughput miR-125a maturation efficiency contains all possible mutations for thirteen 6-nucleotide boxes plus all fifteen possibilities at the 2-nucleotide bulge, which in addition to the major allele sequence (WT) amount to 53 251 sequence variants (Figure 2A). We submitted all sequence variants to 2D structure prediction using the accelerated MC-Flashfold implementation [44] of the MC-Fold software [33]. We found that 29 478 of them adopt WT 2D minimum free energy (MFE) structure (Figure 2A). Because NMA assumes the input structure is at equilibrium, we decided to focus on these 29 478 sequence variants. Furthermore, we realized that the first eight mutated boxes account for the vast majority (over 90%) of these variants that adopt the WT MFE structure. We thus restricted our analysis to this set of 26 960 variants. The proportion of mutations at each position leading to the WT MFE structure is plotted in Figure 2B, showing a very clear drop in the proportion of sequence variants adopting the WT MFE structure beyond box 8.

Since the mutations were performed exhaustively for each box, there is a great deal of sequence redundancy in the dataset. For example, every of the three possibilities of mutation at a given position appears in as much as 1024 sequence variants in the full dataset. To prevent the learning of sequence features by the models, which could be guessed by the slight variations in the position of the nucleobase masses, we contructed a hard benchmark which ensures all mutated positions in the testing set were never mutated in the training set. In that hard benchmark, the middle base pair of mutated boxes 1-8 was reserved for the testing set while all sequence variants affecting only the bottom and top base pairs of every box were selected for the training set (illustrated in Figure 2C). This left us with 1849 sequence variants in the training set and 116 in the testing set. For each set, a CSV file containing the selected variants along with their measured maturation efficiency is available as supplementary material.

To generate a 3D model of pri-miR-125a, MC-Sym [33] was run using the WT sequence and 2D MFE structure as input, with default parameters. This generated 67 predicted 3D structures and the medoid structure (shown in Figure 2D) was selected as the template on which all mutations were performed *in silico* using the ModeRNA software [37].

### 2.9 *Multivariate linear regression*

After running the ENMs on the pri-miR-125a 3D models of the variants, each position in the Entropic Signature is standardized and used as an input variable to train a LASSO linear regression model [45] to predict the miR-125a maturation efficiency. The inclusion of the standardized MC-Fold enthalpy of folding in the model was also tested. We restricted ourselves

**Figure 2: MiR-125a 2D MFE structure, mutation boxes, hard benchmark sets and 3D structure.** A) The 14 boxes that were each exhaustively mutated in the Bartel study, shown on the MC-Fold predicted WT MFE structure. B) Proportions of sequence variants containing each position which adopt the WT MFE structure. C) Positions in the hard benchmark testing set shown in red, and positions in the training set in blue. D) The medoid 3D miR-125a structure from the 67 structures predicted by MC-Sym.

to a linear regression model to allow the pinpointing of important regions in the pri-miR-125a structure, and the LASSO model has the added benefit of selecting important features by driving most coefficients to zero when there is redundancy between the input variables.

As a sanity check to ensure a sequence-only model cannot learn anything from our hard benchmark, we also test a very simple sequence vector model. The vector is composed of 4N positions for an RNA sequence of length N where each position can only take values of 1 or -1. For each block of four positions, only one takes the value 1: the first if the nucleotide at the corresponding position in the sequence is an A, the second for a C, the third for a G and the last for a U. In theory, such a model could learn sequence features from a naive 80/20 random split of the whole dataset but should not be able to capture anything from the hard benchmark as the variables which change in the test set were constant across the entire training set.

2.10 *Computational cost*

To assess the computational cost of the tested ENMs, we ran all models on the RNA chain from the Thermus thermophilus 30S ribosomal subunit [46]. We chose such a large structure of 1514 nucleotides (4541 masses) to highlight the asymptotic behaviour of the tested models, as the building and diagonalizing of the Hessian take time proportional to the cube of the number of masses in the system. The use of a large structure makes operations of linear time like the parsing of the PDB file and the assignment of the masses take a smaller proportion of compute time and allows for a more direct comparison of the costs associated with each different potential. We ran PD-ANM and Cut-ANM with the best parameters found within the B-factors prediction benchmark. In addition, we ran Cut-ANM at 5 Å distance cutoff to see the effect of having a sparser Hessian to diagonalize. As everywhere else throughout the present study, ENCoM was used without modifying the default parameters. The hardware used for these tests is the AMD Rome 7532 (2.40 GHz) as part of Calcul Québec's Narval supercomputer, restricted to 1 CPU with the Slurm shcheduler. We repeated the test 10 times for each model to account for variability in the disk read times and other potential discrepancies. We also ran the same protocol using our 3D model of miR-125a (86 nucleotides, 258 masses), this time building

the models and Entropic Signature 100 times per job and reporting the mean compute time for 10 repetitions. This was to illustrate the computational cost of this more modestly sized structure which is also the focus of our high-throughput mutational analysis.

## 3 RESULTS

### 3.1 *Experimental B-factors prediction*

X-ray crystallography gives rise to experimental temperature factors commonly called B-factors, which measure how much each atom oscillates around its equilibrium position in the crystal. The correlation between these fluctuations and those predicted by ENMs is a common benchmark, which we have performed here on X-ray RNA structures of 2.5 Å resolution or less. The clustering step left us with 38 high-resolution structures divided in 34 sequence clusters. To predict the B-factors, we compared the classical mean-square fluctuations (MSF) with the Entropic Signature we introduce (see Materials and methods). Figure 3A shows the Pearson correlation between the predicted and experimental B-factors for the three tested models as a function of the $\beta$ thermodynamic scaling factor. For Cut-ANM and PD-ANM, we also explored their respective parameters extensively and use the ones which give the highest mean correlation (power dependency of 7 for PD-ANM and distance cutoff of 10 Å for Cut-ANM). Supplementary Figure 1 shows the details of the parameter exploration, highlighting the discovery of a clear maximum performance. As stated before, we did not test any other ENCoM parameters than the ones already published for proteins. Figure 3B shows the B-factor benchmark in boxplot format, comparing the performance of the three ENMs using either the Entropic Signature (ES) or the classical MSF as a prediction of experimental B-factors. A paired Wilcoxon signed-rank test [47] is performed between the two different forms of prediction for each model. Both ENCoM and Cut-ANM benefit significantly from the ES predictions as opposed to MSF (p = 0.0074 and p = 0.019, respectively), while the benefits for PD-ANM do not reach statistical significance on this dataset (p = 0.65). Since all models perform better using the ES, paired Wilcoxon signed-rank tests are performed between all pairs of ENMs exclusively for the ES predictions. Only the pairwise comparison between Cut-ANM and PD-ANM is statistically significant at p = 0.0038 despite ENCoM having the worst average performance of the three ENMs. The reason for this is that the two flavors of ANM have a similar performance profile across the 34 sequence clusters, while ENCoM tends to perform well where they don't and vice versa. This pattern is apparent when examining the list of correlations for every of the 38 structures shown in Supplementary Table 3. For the MSF predictions, the mean correlations are 0.42 for ENCoM, 0.43 for Cut-ANM and 0.45 for PD-ANM while for the ES predictions, they are 0.44 for ENCoM, 0.48 for Cut-ANM and 0.46 for PD-ANM. These values seem a little low when compared to the correlations previously obtained with ENCoM and other tools applied to proteins [11], all between 0.54 and 0.60. In addition, B-factor correlations of up to around 0.60 have been reported with ANM applied on RNA in a similar fashion as here, with 3 masses per residue, by Zimmermann and Jernigan [6]. However, these studies were all done with only one target B-factor value per nucleotide even when the models used more masses, whereas we correlated the predicted fluctuations at every position in our systems (3 masses per nucleotide).

For three of the 34 sequence clusters used in the experimental B-factors prediction benchmark, both ENCoM and ANM get very low and sometimes negative correlations (shown in Supplementary Table 3, clusters 15, 26, 28). These results could be due to artifacts such as crystal packing or buffer contents, which are not considered by the models, driving the B-factors. If we exclude the cases from these three clusters as experimental flaws and take the mean of the 31 remaining clusters, the correlations we get for the MSF predictions (values in parentheses are the correlations witout excluding these three clusters) are 0.47 (0.42) for ENCoM, 0.48 (0.43) for Cut-ANM and 0.49 (0.45) for PD-ANM while for the ES predictions,

**Figure 3: Pearson correlation coefficient between predicted and experimental B-factors for ENCoM and ANM.** A) The mean Pearson correlation is shown as a function of the $\beta$ scaling factor for the three ENMs. B) The Entropic Signature using the best scaling factor for each model is compared to the mean square fluctuations (MSF). The p-values from paired Wilcoxon signed-rank tests are shown for pairs of the two types of predictions for every model and for every pair of models, only for the Entropic Signature predictions since they outperform the MSF.

they are 0.49 (0.44) for ENCoM, 0.51 (0.48) for Cut-ANM and 0.49 (0.46) for PD-ANM. These values are closer to what is observed when applying NMA models to proteins [11].

### 3.2 *Prediction of conformational changes*

Another source of information regarding dynamics arising from X-ray crystallography is the crystallization of the same molecule in different conformations due to either varying experimental conditions or the presence of mutations in the sequence of residues. After clustering and applying filters (see Materials and methods), we obtain for this conformational change benchmark 129 structures divided in 29 sequence clusters, forming 240 distinct pairs of conformations with RMSD of at least 2 Å. Each of the clusters contains at least one pair of distinct conformations for a given RNA. We use the 240 pairs of conformations to test the models' ability to predict conformational change. This ability is in many ways more relevant to a biological context, as the conformational variation resulting from different experimental conditions happens on much longer timescales than B-factors and can have functional significance [15]. Figure 4 presents the average cumulative overlap from pairs of different conformations, computed from a single mode up to 10% of the total number of normal modes. The idea behind this as opposed to the more frequent use of a fixed number of modes is that since the number of modes grows linearly with the number of masses in the system ($3N - 6$ nontrivial modes for N masses), it is more logical to use a linear proportion of them. Figure 4B shows the results as boxplots when 5% of the normal modes are used. The average number of modes selected this way for the current benchmark is 28 (data shown in Supplementary Figure 4), which falls slightly outside the range of 3 to 20 modes tested by Zimmermann *et al.* [6]. In a real-case scenario, limiting the analysis to a small number of modes is mandatory because of the exponential growth of the conformational space they represent, however relevant modes could still be selected among a fixed linear proportion of the total.

At 5% of nontrivial normal modes, the mean cumulative overlap is 0.73 for ENCoM, 0.67 for Cut-ANM and 0.69 for PD-ANM. The p-values from a paired Wilcoxon signed-rank test are shown for each pair of models, illustrating that the advantages ENCoM has over both ANMs and the advantage PD-ANM has over Cut-ANM are very significant. This result is consistent

with our previous results benchmarking ENCoM on proteins, where ENCoM performed highest of all the models tested on the conformational change benchmark [11].

**Figure 4: Cumulative overlap between normal modes and conformational changes from X-ray crystallography experiments.** A) Mean cumulative overlap as a function of the proportion of the nontrivial normal modes used. B) Mean cumulative overlap at 5% nontrivial normal modes used, with p-values from paired Wilcoxon signed-rank tests for every pair of models.

### 3.3 *Capturing NMR ensemble variance*

Perhaps the most reliable source of experimental information regarding the 3D structural dynamics of RNA comes from solution NMR experiments. In contrast to X-ray crystallography, NMR experiments give us direct information about the conformational space of the molecule studied over vast timescales. The motions apparent within the structural ensembles resolved using this technique can inform us on the ability of NMA models to recreate the conformational space of a molecule from its lowest energy conformation. After filtering and clustering (see Materials and methods), we are left with 313 solution NMR ensembles in 271 sequence clusters. We test each ENM's ability to capture the internal motions apparent from these ensembles by computing the RMSIP and NCO between PCs accounting for 99% of the ensemble variance and a number of the lowest frequency nontrivial modes computed the ensemble's equilibrium structure. We use from 1 mode up to 10% of the nontrivial normal modes and report the performance across this range in Figure 5, panels A and C (RMSIP and NCO, respectively). Panels B and D present the performance at 5% nontrivial normal modes, which corresponds to an average number of 14 normal modes (see Supplementary Figure 4).

The average RMSIP at 5% normal modes is 0.67 for ENCoM, 0.64 for Cut-ANM and 0.65 for PD-ANM. The average NCO, also at 5% normal modes, is 0.78 for ENCoM, 0.75 for Cut-ANM and 0.76 for PD-ANM. Again, the p-values from Wilcoxon signed-rank tests are shown for every pair of models. For the two metrics, the difference in performance is highly significant between each pair of models ($p < 10^{-14}$). Again, this confirms that ENCoM has the advantage when it comes to predicting conformational changes happening over longer timescales than B-factors.

### 3.4 *Predicting miR-125a maturation efficiency*

To ensure a stringent assessment of whether the ENMs tested can capture the influence of 3D structural dynamics on miR-125a maturation efficiency beyond sequence features, we put together a training set of 1849 sequence variants and a testing set of 116 which have no mutated

**Figure 5: RMSIP and NCO between normal modes and principal components from NMR ensembles.**
A) Root mean square inner product (RMSIP) between a proportion of the nontrivial normal modes and the principal components accounting for at least 99% of the variance apparent within the NMR ensemble. B) RMSIP at 5% nontrivial normal modes, with p-values from paired Wilcoxon signed-rank tests for every pair of models. C) Normalized cumulative overlap (NCO) between a proportion of the nontrivial normal modes and the principal components accounting for at least 99% of the variance apparent within the NMR ensemble. D) NCO at 5% nontrivial normal modes, with p-values from paired Wilcoxon signed-rank tests for every pair of models.

position in common. Figure 6A shows the predictive $R^2$ from LASSO models trained on the training set and applied to the testing set. For the three ENMs, the Etropic Signature (ES, see Materials and methods) is computed for 41 values of the $\beta$ scaling factor centered around the one which gave the best performance on the B-factors benchmark ($e^{0.25}$ for ENCoM, $e^1$ for Cut-ANM, $e^{9.5}$ for PD-ANM), varied logarithmically in increments of 0.25. In addition, the regularization strength $\lambda$ of the LASSO model is tested in $\log_2$ increments from $2^{-15}$ to 1. We also tested a sequence vector (see Materials and methods) to validate that the hard benchmark truly prohibits the learning of sequence features. Apart from these four models on their own, the addition of the MC-Fold enthalpy of folding to the LASSO model is tested and the results are shown on the bottom row of Figure 6A. Interestingly, ENCoM is the only ENM for which the ES alone allows the LASSO model to have a positive predictive $R^2$. The maximum value of 0.11 is reached at $\beta = e^{0.25}$, which is the center value from the swept $\beta$ parameters and the one which gave the best performance in the B-factors prediction benchmark. The MC-Fold enthalpy of folding alone also reaches a modest positive predictive $R^2$ of 0.07. As expected, the sequence vector ends up being a constant predictor and its combination with the MC-Fold energy produces the exact same cloud of points as MC-Fold alone. On their own, Cut-ANM and PD-ANM fail to capture any of the variance in the testing set as illustrated by their predictive

$R^2$ values being zero or less for all parameters scanned. This result was expected as these models are completely agnostic to the chemical nature of the nucleobases. The combination of ENCoM with MC-Fold energy seems synergistic, giving rise to a maximal predictive $R^2$ of 0.22. To assess the significance of this value, we simulated the combination of the MC-Fold alone predictions with random predictions generated as a normal distribution centered around the training set mean, with half its standard deviation. The results of this simulation are shown in Supplementary Figure 5. The p-value for the combination of the different models with MC-Fold being significantly better than just MC-Fold combined with noise are shown in the bottom row of Figure 6B. The combination of the ENCoM ES with MC-Fold leads to a highly significant improvement with $p < 0.0001$.

**Figure 6: Performance of LASSO linear regression on the hard test.** A) Predictive $R^2$ for each model alone or in combination with the MC-Fold enthalpy of folding. For the three ENMs, scaling factors for the dynamical signature were explored around the value which gave the best respective performance in the B-factors benchmark. Predictive $R^2$ values below zero are shown in gray. B) Performance on the test set for the best combination of parameters for every model alone or in tandem with MC-Fold. The bottom row shows the p-value from a simulation combining gaussian noise with the MC-Fold prediction, which corresponds to the probability of these predictive $R^2$ values arising by chance from a pure noise model.

The best predictive performance on our hard benchmark of miR-125a maturation efficiency is thus reached by the combination of the ENCoM ES at $\beta = e^{2.25}$ with the MC-Fold enthalpy of folding, fed into a LASSO linear regression model with regularization strength $\lambda = 2^{-4}$. The coefficients of the resulting model are shown in graph format in Figure 7A and mapped back on the 2D structure of pri-miR-125a in Figure 7B. The highest remaining coefficient from the ENCoM ES is a positive coefficient at the phosphate mass of nucleotide 78. This pattern corresponds to a motif previously identified by Fang *et al.* as the mismatched GHG motif [26]. Strikingly, that position is included in the testing set, meaning that the model learns the importance of flexibility at a position never mutated in the training set. All variables in the model are standardized so the coefficients can be directly compared as indicators of the variables' relative importances. With this model, the MC-Fold energy has a coefficient of -0.19 while the sum of ENCoM coefficients is 0.08. These coefficients point to the model favoring a low enthalpy of folding but an entropically flexible structure at the same time, especially at the GHG motif. All the ENCoM coefficients are positive with these parameters so the sum of absolute coefficients is also 0.08, having roughly half the importance of the MC-Fold energy.

Using the same $\beta$ parameter that led to the most generalizable learning from the combination of ENCoM and MC-Fold, we trained another round of LASSO models, this time using the whole set of 26 960 mutant sequences from boxes 1-8 with WT MFE as the training set. Figure 7C illustrates the training performance of the model as measured by Pearson's R linear correlation, $R^2$ and the square of Pearson's R. For high enough regularization strength, $R^2$ dips below the square of Pearson's R, which points to the model being underpowered due to the high regularization. We thus choose $2^{-8}$ as a good tradeoff between the reduction of model variance and the maintaining of good performance. In Figure 7D, we can see that training $R^2$ stays above 90% of its maximal value and Pearson's R above 95%. The coefficients from this model are shown in panels E and F in a similar fashion as in panels A and B. The lower regularization strength allows interesting patterns to emerge at nucleobase masses throughout the structure while still being high enough to drive most backbone coefficients to zero. Here again, the most important backbone feature is flexibility at the GHG mismatch motif. At this regularization strength, the MC-Fold energy coefficient is -0.23, the sum of ENCoM coefficients is 0.15 and the sum of absolute ENCoM coefficients is 0.59. The overall trend of low enthalpy and high vibrational entropy being favorable for processing is still present, but the ENCoM coefficients have almost three times the relative importance of the MC-Fold enthalpy.

### 3.5 *Computational cost*

The results of the computational cost analysis are given in Supplementary Figure 6. For the Thermus thermophilus 30S ribosomal subunit, on average, ENCoM takes 26.5 minutes, Cut-ANM takes 19.2 and 18.8 minutes for the 18 and 5 Å cutoff values respectively and PD-ANM takes 22.7 minutes. For miR-125a, on average for building one model and computing one Entropic Signature, ENCoM takes 3.33 seconds, Cut-ANM takes 1.63 seconds while the 5 Å version takes 1.56 seconds and PD-ANM takes 2.30 seconds. These results illustrate that for large structures, the difference in CPU cost between the models becomes smaller as more time is spent on solving the Hessian matrix, while for small structures the difference is much more significant since the building of the Hessian takes more time for more complex (ENCoM) or more interconnected (PD-ANM) potentials. Cut-ANM is faster at a lower distance cutoff because it leads to a sparser Hessian which is in turn faster to solve.

**Figure 7: Coefficients of LASSO regression models.** A) Coefficients of the combined ENCoM and MC-Fold model trained on the hard training set with $\beta = e^{2.25}$ and $\lambda = 2^{-4}$. B) ENCoM Entropic Signature coefficients mapped on the 2D structure of pri-miR-125a. The DROSHA cut site and mismatched GHG motif are identified on the structures. C) Training metrics from training the ENCoM-MC-Fold combination at $\beta = e^{2.25}$ on the whole set of 26 960 sequence variants selected for this study. Pearson's R, $R^2$ and the square of Pearson's R are plotted in relation to regularization strength. A red vertical line shows the last regularization strength before $R^2$ starts to diverge from the square of Pearson's R. D) Pearson's R and $R^2$ training performance expressed as a proportion of maximal performance, as a function of LASSO regularization strength. E)-F) Same as A)-B), with the LASSO model trained on all 26 960 sequences and $\lambda = 2^{-8}$.

### 4.1 *Overall performance and parameterization*

In the present work, we have shown gains in performance of ENCoM over ANM when applied to RNA to predict large conformational changes, at the expense of a small loss in the ability to predict B-factors from crystallographic experiments. Namely, ENCoM shows an advantage when it comes to conformational change prediction and NMR ensemble variance prediction. Moreover, this gain in performance is accessible directly from parameters that were optimized for performance on proteins without any re-parameterization, whereas we extensively tested the single parameter for both versions of ANM and used the parameters leading to maximal performance. While the ENCoM model is very robust in performance across vast parameter space, it would be interesting to explore this space in relation with performance on RNA structures. We leave this exploration for future work as our goal would be to find parameters optimizing performance on RNA, proteins and their complexes, a task which is beyond the scope of the present work. For now, the default ENCoM parameters that were already published provide a measurable advantage over other ENMs on RNA structures. Because we used parameters originally optimized for performance on proteins, the model is readily applicable to RNA-protein complexes as it is. We can thus say that while it may be slight, the advantage ENCoM has over Cut-ANM and PD-ANM in the prediction of large conformational changes is very robust. This asset would make ENCoM the preferred model in the case of novel structures for which the optimal parameters for ANM cannot be easily estimated.

Our main interest in extending ENCoM to RNA was to use it to study the effect of mutations as we have already done in the case of proteins [12, 14]. The present works illustrates that ENCoM indeed extends this capability to RNA, which is not the case of the sequence agnostic ANM model.

### 4.2 *Artifacts from crystallographic B-factors*

It might be worthwhile to investigate in more details the nature of the quasi-null correlations obtained for three out of the 34 sequence clusters used for the B-factors prediction benchmark to see if some factor in the experimental conditions or the nature of the RNAs could explain the discrepancy. However, in the case of ENCoM, we are more interested in predicting conformational changes than B-factors, and we have previously shown that parameter optimization for our model offers a trade-off between conformational space and B-factors prediction [11]. Furthermore, B-factors have been shown to adopt biphasic behavior with a different slope of variation for cryogenic versus standard temperatures [48]. Since most X-ray crystallography experiments are conducted at cryogenic temperatures, their B-factors measurements are thus less relevant to biological function. B-factors can also arise from crystal packing and rigid body motions which are irrelevant in a biological context [49].

### 4.3 *Large-scale conformational change prediction*

When it comes to the prediction of large-scale, slow timescale conformational change, whether from pairs of X-ray structures in different conformations or from NMR ensembles, ENCoM has a clear advantage over both Cut-ANM and PD-ANM. This better performance comes as no surprise in view of our previous studies [11] and reinforces the idea that moving the coarse-grained ENMs in the direction of more physical realism is beneficial. One innovative further development that could be added to ENCoM is the inclusion of an electrostatic term. In the case of RNA especially, electrostatic interactions are of primordial importance and capturing their relative strengths within the model would most probably further increase its accuracy.

All coarse-grained ENMs have in common the pseudo-physical nature of their potentials. Their eigenfrequencies do not have defined units, and the same goes for the vibrational entropy values we can compute from them. However, the relative ranking of eigenvalues is very relevant to biological function as the slowest normal modes capture the most global, cooperative motions which are often critical for macromolecular function. In this work we introduced the Entropic Signature to reduce the conformational space from all normal modes to a vector of the same length as the number of masses in the system. In contrast to the classically used mean square fluctuations, the addition of a thermodynamic scaling factor as part of the vibrational entropy calculation allows for varying the relative contributions of slow and fast motions to the dynamical signature. This is of utmost importance considering the pseudo-physical nature of the models, as parameters such as temperature and physical constants do not have a clear meaning for the coarse-grained ENMs.

The hard miR-125a maturation efficiency benchmark illustrates this importance of fitting the scaling factor to the studied system. For the ENCoM dynamical signature alone, only a small range of $\beta$ values gave rise to positive predictive $R^2$. At the same time, the best $\beta$ values are close to the value selected from the B-factors prediction benchmark. Thus, it seems that a small range of values might be appropriate for a large range of different molecules.

To our knowledge, this is the first time 3D structural dynamical information has been used in such a way to train multivariate linear models in predicting the effect of mutations. The method has the advantage of being a very powerful statistical model while maintaining the necessary transparency to allow the identification of important regions and patterns directly on the structure. The drawback is the inability of the model to capture complex relationships between the flexibility of different regions, as every input variable is linearly independent.

While the performance of the LASSO model combining ENCoM with the MC-Fold enthalpy is modest with a predictive $R^2$ of 0.22, we must keep in mind that perfect performance on this hard test is not expected and might not even be possible. Indeed, it cannot be excluded that sequence features are part of what constitutes the processing signal for miRs, so this benchmark must be seen as a test of generalizability of the model rather than an upper bound on its performance. Supplementary Figure 2 shows the testing performance when randomly splitting the data in 80% for training and 20% for testing. As discussed in Materials and methods, this is a naive approach that leads to vast amounts of sequence redundancy between the two sets. However, the performance of all models is strikingly high, with the combination of ENCoM and MC-Fold reaching 0.75 predictive $R^2$ and the combination of both Cut-ANM and PD-ANM with MC-Fold reaching 0.74. Even the sequence vector reaches 0.71 predictive $R^2$ when combined with the MC-Fold energy. As opposed to the hard benchmark we developed, this test can be seen as an upper bound on the models' performance. It might seem surprising at first that ANM can capture some signal on its own. However, this signal is explained by the fact that the nucleobase masses all have slightly different positioning relative to the backbone. The different positions of the masses thus lead to a sequence signal being present in ANM's conformational space, which the LASSO model can learn. Performing the same experiment with only one mass per nucleotide, located at the phosphorous atom, leads to the signal completely disappearing for ANM, since the backbone is fixed and thus the dynamical signatures computed for all variants are identical. As for the appearance of a performance ceiling for this 80/20 train/test split, we hypothesize that noise in the experimental data is causing it. One possibility is that since the data is generated by incubating all mutants in the same test tube before sequencing their barcodes to determine processing efficiency [26], it allows for interactions between different molecules to take place, potentially interfering with sequencing accuracy.

Perhaps the most interesting application of our multivariate linear model applied to ENCoM dynamical signatures is the possibility to map the selected coefficients back to the structure of the studied molecule to gain biological insights. In the present case of miR-125a, the selection of the noncanonical UC base pair from the mismatched GHG motif by the LASSO model as the most important feature from the dynamical signature strikes us as very promising in terms of our approach's ability to lead to relevant insights. A striking finding from our study is that the model favors sequences which have both low folding energy and high vibrational entropy. This finding is at odds with Fang and Bartel's model of pri-miR processing in which the gain of more rigid base pairs throughout the whole stem is associated with higher processing efficiency, with the only exception being the GHG mismatch. However, they analyzed their data with a position-centric view in which the effect of a mutation was deemed to happen at the mutated position only. Our model captures the effect of a mutation on the flexibility of the whole pri-miR, thus enabling the recognition of much finer dynamical patterns happening away from the site of mutation. In fact, our model does not contradict the hypothesis of more rigid base pairs leading to better processing, as this relationship is captured by the MC-Fold enthalpy of folding (which favors canonical, and GC base pairs even more). However, from an evolutionary point of view, a rigid hairpin of 35 GC basepairs minus one mismatch would probably lead to the DNA sequence of the miR gene folding on itself and wreaking havoc on genomic stability when transcribed and replicated. Thus, the maintaining of enough flexibility in the hairpin structure could be a way to prevent the DNA hairpin from forming while still allowing the RNA structure to exist.

When looking at the coefficients for the nucleobase masses in Figure 7F, one apparent pattern is that negative coefficients tend to be overrepresented on base pairs that are already rigid while the reverse is true for positive coefficients. This pattern also points away from rigidity being highly favored, as if it were the case, we should see GU wobble and noncanonical base pairs colored in blue. Instead, the reverse pattern is apparent, with most negative coefficients appearing at GC base pairs and positive coefficients appearing in places where a mismatch is already present like the UG base pair at position 19. There are even two GC base pairs for which more flexibility is favored by the model: at positions 25 and 2. Our results thus point to very complex dynamical patterns being intimately tied to human miR processing and merit further investigation. As part of future work, the LASSO model trained from the whole set of 26 960 sequence variants seems an ideal candidate to predict maturation efficiencies for novel pri-miR-125a variants.

## 4.6   *Conclusions*

We presented a version of ENCoM specifically tailored to work optimally on RNA molecules by allowing the use of three masses per nucleotide in the coarse-graining of the input structure. ENCoM consistently outperforms ANM for large-scale conformational space prediction, in addition to being parameter-free as far as the user is concerned. However, the main advantage of ENCoM is the possibility to estimate the effect of mutations, which ANM is incapable of due to its sequence agnostic nature. Utilizing high-throughput data on the maturation efficiency of miR-125a sequence variants, we show that ENCoM indeed preserves this ability on RNA molecules by the careful construction of a stringent test from the maturation dataset. Moreover, the usage of the ENCoM Entropic Signature inside a multivariate linear model is the first of its kind, to the best of our knowledge. It allows us to point back to regions of biological interest on the structure of pri-miR-125a, namely the mismatched GHG motif. We also challenge the notion that rigid base pairs favor processing at almost every position. Our methodology is easily applicable to proteins, RNAs and their complexes, provided some high-throughput data

about the biological property of interest is available for the studied molecule, along with an experimentally solved structure or a good 3D model.

## 5 ACKNOWLEDGEMENTS AND FUNDING

We thank Marie Papineau, Frédéric Mailhot and Etienne Richan for their careful reading of the manuscript. RJN is part of PROTEO (the Québec network for research on protein function, structure and engineering).

This work was supported by Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery program grants (FM and RN); Genome Canada and Genome Quebec (RN); Compute Canada (RN) and Canadian Institutes of Health Research (CIHR) (FM, grant number MOP-93679). OM is the recipient of a Fonds de Recherche du Québec—Nature et Technologies (FRQ-NT) Doctorate's scholarship; and a Faculté des Études Supérieures et Postdoctorales de l'Université de Montréal scholarship for direct passage to the PhD. VF is the recipient of a National Sciences and Engineering Research Council of Canada (NSERC) postdoctoral fellowship.

Conflict of Interest: none declared.

## REFERENCES

[1] Eric J Strobel, Kyle E Watters, David Loughrey, and Julius B Lucks. RNA systems biology: uniting functional discoveries and structural tools to understand global roles of RNAs. *Current Opinion in Biotechnology*, 39:182–191, June 2016.

[2] Peter W. Rose, Andreas Prlić, Chunxiao Bi, Wolfgang F. Bluhm, Cole H. Christie, Shuchismita Dutta, Rachel Kramer Green, David S. Goodsell, John D. Westbrook, Jesse Woo, Jasmine Young, Christine Zardecki, Helen M. Berman, Philip E. Bourne, and Stephen K. Burley. The RCSB protein data bank: views of structural biology for basic and applied research and education. *Nucleic Acids Research*, 43(D1):D345–D356, nov 2014.

[3] Hashim M Al-Hashimi and Nils G Walter. RNA dynamics: it is about time. *Current Opinion in Structural Biology*, 18(3):321–329, jun 2008.

[4] Ivet Bahar, Timothy R. Lezon, Lee-Wei Yang, and Eran Eyal. Global dynamics of proteins: Bridging between structure and function. *Annu. Rev. Biophys.*, 39(1):23–42, apr 2010.

[5] Lee-Wei Yang, A J Rader, Xiong Liu, Cristopher Jon Jursa, Shann Ching Chen, Hassan A Karimi, and Ivet Bahar. o GNM: online computation of structural dynamics using the gaussian network model. *Nucleic Acids Research*, 34(suppl_2):W24–W31, jul 2006.

[6] Michael T. Zimmermann and Robert L. Jernigan. Elastic network models capture the motions apparent within ensembles of RNA structures. *RNA*, 20(6):792–804, apr 2014.

[7] Giovanni Pinamonti, Sandro Bottaro, Cristian Micheletti, and Giovanni Bussi. Elastic network models for RNA: a comparative assessment with molecular dynamics and SHAPE experiments. *Nucleic Acids Res*, 43(15):7260–7269, jul 2015.

[8] Turkan Haliloglu, Ivet Bahar, and Burak Erman. Gaussian dynamics of folded proteins. *Phys. Rev. Lett.*, 79(16):3090–3093, oct 1997.

[9] Ivet Bahar, Ali Rana Atilgan, and Burak Erman. Direct evaluation of thermal fluctuations in proteins using a single-parameter harmonic potential. *Folding and Design*, 2(3):173–181, jun 1997.

[10] A.R. Atilgan, S.R. Durell, R.L. Jernigan, M.C. Demirel, O. Keskin, and I. Bahar. Anisotropy of fluctuation dynamics of proteins with an elastic network model. *Biophysical Journal*, 80(1):505–515, jan 2001.

[11] Vincent Frappier and Rafael J. Najmanovich. A coarse-grained elastic network atom contact model and its use in the simulation of protein dynamics and the prediction of the effect of mutations. *PLoS Comput Biol*, 10(4):e1003569, apr 2014.

[12] Vincent Frappier and Rafael Najmanovich. Vibrational entropy differences between mesophile and thermophile proteins and their use in protein engineering. *Protein Science*, 24(4):474–483, nov 2014.

[13] Vincent Frappier, Matthieu Chartier, and Rafael J. Najmanovich. ENCoM server: exploring protein conformational space and the effect of mutations on protein function and stability. *Nucleic Acids Research*, 43(W1):W395–W400, April 2015.

[14] Natália Teruel, Olivier Mailhot, and Rafael J. Najmanovich. Modelling conformational state dynamics and its role on infection for SARS-CoV-2 spike protein variants. *PLoS Comput Biol*, 17(8):e1009286, aug 2021.

[15] Katherine Henzler-Wildman and Dorothee Kern. Dynamic personalities of proteins. *Nature*, 450(7172):964–972, dec 2007.

[16] Osni Marques and Yves-Henri Sanejouand. Hinge-bending motion in citrate synthase arising from normal mode calculations. *Proteins*, 23(4):557–560, dec 1995.

[17] Cristian Micheletti, Paolo Carloni, and Amos Maritan. Accurate and efficient description of protein vibrational dynamics: Comparing molecular dynamics and gaussian models. *Proteins*, 55(3):635–645, mar 2004.

[18] Ozge Kurkcuoglu, Robert L. Jernigan, and Pemra Doruker. Loop motions of triosephosphate isomerase observed with elastic networks. *Biochemistry*, 45(4):1173–1182, dec 2005.

[19] José Ramón Lopéz-Blanco, José Ignacio Garzón, and Pablo Chacón. iMod: multipurpose normal mode analysis in internal coordinates. *Bioinformatics*, 27(20):2843–2850, aug 2011.

[20] E. Eyal, G. Lum, and I. Bahar. The anisotropic network model web server at 2015 (ANM 2.0). *Bioinformatics*, 31(9):1487–1489, jan 2015.

[21] Wolfram Saenger. *Principles of Nucleic Acid Structure*. Springer New York, 1984.

[22] Victor Ambros. The functions of animal microRNAs. *Nature*, 431(7006):350–355, sep 2004.

[23] Julia Winter, Stephanie Jung, Sarina Keller, Richard I. Gregory, and Sven Diederichs. Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol*, 11(3):228–234, mar 2009.

[24] Ranhui Duan, ChangHui Pak, and Peng Jin. Single nucleotide polymorphism associated with mature miR-125a alters the processing of pri-miRNA. *Human Molecular Genetics*, 16(9):1124–1131, mar 2007.

[25] Paul Dallaire, Huiping Tan, Keith Szulwach, Christopher Ma, Peng Jin, and François Major. Structural dynamics control the MicroRNA maturation pathway. *Nucleic Acids Res*, page gkw793, sep 2016.

[26] Wenwen Fang and David P. Bartel. The menu of features that define primary MicroRNAs and enable de novo design of MicroRNA genes. *Molecular Cell*, 60(1):131–145, oct 2015.

[27] Tu-Liang Lin and Guang Song. Generalized spring tensor models for protein fluctuation dynamics and conformation changes. *BMC Structural Biology*, 10(Suppl 1):S3, 2010.

[28] B.J. McConkey, V. Sobolev, and M. Edelman. Quantification of protein surfaces, volumes and atom-atom contacts using a constrained voronoi procedure. *Bioinformatics*, 18(10):1365–1373, oct 2002.

[29] Vladimir Sobolev, Rebecca C. Wade, Gert Vriend, and Marvin Edelman. Molecular docking using surface complementarity. *Proteins*, 25(1):120–129, may 1996.

[30] Ivet Bahar and AJ Rader. Coarse-grained normal mode analysis in structural biology. *Current Opinion in Structural Biology*, 15(5):586–592, oct 2005.

[31] Lei Yang, Guang Song, and Robert L. Jernigan. Protein elastic network models and the ranges of cooperativity. *Proc. Natl. Acad. Sci. U.S.A.*, 106(30):12347–12352, jul 2009.

[32] Olivier Mailhot and Rafael Najmanovich. The NRGTEN python package: an extensible toolkit for coarse-grained normal mode analysis of proteins, nucleic acids, small molecules and their complexes. *Bioinformatics*, 37(19):3369–3371, mar 2021.

[33] Marc Parisien and François Major. The MC-fold and MC-sym pipeline infers RNA structure from sequence data. *Nature*, 452(7183):51–55, mar 2008.

[34] Gale Thodes. *Crystallography Made Crystal Clear*. Elsevier, 1993.

[35] Yue Wang, Norma M. Wills, Zhihua Du, Anupama Rangan, John F. Atkins, Raymond F. Gesteland, and David W. Hoffman. Comparative studies of frameshifting and non-frameshifting RNA pseudoknots: A mutational and NMR investigation of pseudoknots derived from the bacteriophage t2 gene 32 mRNA and the retroviral gag-pro frameshift site. *RNA*, 8(8):981–996, aug 2002.

[36] Boris Fürtig, Christian Richter, Jens Wöhnert, and Harald Schwalbe. NMR spectroscopy of RNA. *ChemBioChem*, 4(10):936–962, sep 2003.

[37] Magdalena Rother, Kristian Rother, Tomasz Puton, and Janusz M. Bujnicki. ModeRNA: a tool for comparative modeling of RNA 3d structure. *Nucleic Acids Research*, 39(10):4007–4022, feb 2011.

[38] Magdalena A. Machnicka, Kaja Milanowska, Okan Osman Oglou, Elzbieta Purta, Malgorzata Kurkowska, Anna Olchowik, Witold Januszewski, Sebastian Kalinowski, Stanislaw Dunin-Horkawicz, Kristian M. Rother, Mark Helm, Janusz M. Bujnicki, and Henri Grosjean. MODOMICS: a database of RNA modification pathways—2013 update. *Nucleic Acids Research*, 41(D1):D262–D267, oct 2012.

[39] D. Defays. An efficient algorithm for a complete link method. *The Computer Journal*, 20(4):364–366, apr 1977.

[40] Saul B. Needleman and Christian D. Wunsch. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal of Molecular Biology*, 48(3):443–453, mar 1970.

[41] Jianpeng Ma and Martin Karplus. Ligand-induced conformational changes in ras p21: a normal mode and energy minimization analysis. *Journal of Molecular Biology*, 274(1):114–131, nov 1997.

[42] H. Hotelling. Analysis of a complex of statistical variables into principal components. *Journal of Educational Psychology*, 24(6):417–441, 1933.

[43] G. H. Golub and C. Reinsch. Singular value decomposition and least squares solutions. *Numer. Math.*, 14(5):403–420, apr 1970.

[44] Paul Dallaire and François Major. Exploring Alternative RNA Structure Sets Using MC-Flashfold and db2cm. *Methods in Molecular Biology*, 1490:237–251, 2016.

[45] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, jan 1996.

[46] Ditlev E. Brodersen, William M. Clemons, Andrew P. Carter, Brian T. Wimberly, and V. Ramakrishnan. Crystal structure of the 30 s ribosomal subunit from thermus thermophilus: structure of the proteins and their interactions with 16 s RNA. *Journal of Molecular Biology*, 316(3):725–768, feb 2002.

[47] Frank Wilcoxon. Individual comparisons by ranking methods. *Biometrics Bulletin*, 1(6):80–83, 1945.

[48] Robert F. Tilton, John C. Dewan, and Gregory A. Petsko. Effects of temperature on protein structure and dynamics: x-ray crystallographic studies of the protein ribonuclease-A at nine different temperatures from 98 to 320K. *Biochemistry*, 31(9):2469–2481, mar 1992.

[49] Reza Soheilifard, Dmitrii E Makarov, and Gregory J Rodin. Critical evaluation of simple network models of protein dynamics and their comparison with crystallographic B-factors. *Physical Biology*, 5(2):026008, jun 2008.