Modulation of multiple gene clusters expression by a single PAS-LuxR transcriptional

regulator.

Cláudia M. Vicente[a,b,c], Tamara D. Payero[a,b], Antonio Rodríguez-García[a,b], Eva G. Barreales[a],

Antonio de Pedro[a], Fernando Santos-Beneit[b,d] and Jesús F. Aparicio[a,*]

[a] Area of Microbiology, Faculty of Biology, University of León, 24071 León. Spain.

[b] Institute of Biotechnology INBIOTEC, Parque Científico de León, Avda. Real, nº 1, 24006

León, Spain.

[c] Present address: TBI, CNRS, INRA, INSA, Université de Toulouse, Toulouse, France

[d] Present address: Institute of Sustainable Processes, University of Valladolid. Spain.

17 *Corresponding author. Tel:+34-987-293420; Fax:+34-987-291409; e-mail:

18 jesus.aparicio@unileon.es

ABSTRACT

PAS-LuxR transcriptional regulators are highly conserved enzymes governing polyene macrolide antifungal biosynthesis. PteF is one of such regulators, situated in the polyene macrolide filipin gene cluster from *Streptomyces avermitilis*. Its mutation leads to a drastic filipin production decline, but also to a severe loss of oligomycin production, an ATP-synthase inhibitor of macrolide structure, and a delay in sporulation, thus it has been considered as a transcriptional activator. Transcriptomic analyses were carried out in *S. avermitilis* Δ*pteF* and its parental strain *S. avermitilis* NRRL 8165 as control. Both strains were grown in YEME medium without sucrose, and samples were taken in the exponential and stationary growth phases. 257 genes showed altered expression in the PteF-deleted mutant, most of them in the exponential phase of growth. Surprisingly, despite PteF being an activator of filipin biosynthesis, a majority of the genes affected upon mutation showed overexpression thus suggesting a negative modulation of those genes. Genes affected were related to various metabolic processes, including genetic information processing; DNA, energy, carbohydrate, and lipid metabolism; morphological differentiation; and transcriptional regulation; among others, but particularly to secondary metabolite biosynthesis. Notably, ten secondary metabolite gene clusters out of 38 encoded by the genome, some of them encoding cryptic compounds, showed altered expression profiles in the mutant, suggesting a regulatory role for PteF wider than expected. Transcriptomic results were validated by quantitative reverse transcription polymerase chain reaction. These findings provide important clues to understand the intertwined regulatory machinery that modulates antibiotic biosynthesis in *Streptomyces*.

KEYWORDS

INTRODUCTION

PAS-LuxR regulators are transcriptional factors that combine an N-terminal PAS sensory domain (Hefti et al., 2004) with a C-terminal helix-turn-helix (HTH) motif of the LuxR type for DNA-binding (Santos et al., 2012). The sensor domain is thought to detect a physical or chemical stimulus and regulate, in response, the activity of the effector domain (Möglich et al., 2009). The archetype of this class of regulators, PimM, was first identified in the antifungal pimaricin biosynthetic gene cluster from *Streptomyces natalensis* (Antón et al., 2007). It was characterised as a transcriptional activator of pimaricin biosynthesis since antifungal production was abolished upon gene deletion, and later its mode of action was characterised at the molecular level (Santos-Aberturas et al., 2011a). Since its discovery, homologous regulatory proteins have been found to be encoded in all known biosynthetic gene clusters of antifungal polyketides, and they have been shown to be functionally equivalent, to the extent that the production of pimaricin is restored in *S. natalensis ΔpimM* upon introduction of heterologous regulators of the PAS-LuxR class, such as *nysRIV* (nystatin), *amphRIV* (amphotericin), or *pteF* (filipin) into the strain (Santos-Aberturas et al. 2011b). Furthermore, introduction of a single copy of *pimM* into the amphotericin-producing strain *S. nodosus*, into the filipin-producing strain *S. avermitilis*, or into the rimocidin producing strain *S. rimosus*, boosted the production of all polyenes, thus indicating that these regulators are fully exchangeable (Santos-Aberturas et al. 2011b).

Although given their location in the chromosome, PAS-LuxR regulators were initially considered pathway-specific transcriptional regulators, recent results have shown that they should be considered as regulators with a wider range of implications. The canonical operator of PimM was used to search for putative targets of orthologous protein PteF in the genome of *S. avermitilis*, finding multiple binding sites located inside or upstream from genes involved in different aspects of both primary and secondary metabolism (Vicente et al,. 2015), thus

suggesting that the regulator could govern those processes. Several of these operators were

selected, and their binding to PimM DNA-binding domain demonstrated by electrophoretic

mobility shift assays (EMSAs). As a proof of concept, the biosynthesis of the ATP-synthase

inhibitor oligomycin whose gene cluster included two operators was studied (Vicente et al,.

2015). PteF deleted mutants, which showed a severe loss of filipin production and delayed spore

formation in comparison to that of the wild-type strain (Vicente et al 2014), also showed a

severe loss of oligomycin production and reduced expression of *olm* genes. Gene

complementation of the mutant restored phenotype, thus demonstrating that PteF was able to

co-regulate the biosynthesis of two related secondary metabolites, the polyketide macrolides

filipin and oligomycin (Vicente et al., 2015). This cross-regulation could therefore be extended

to all the clusters where operators were found, which suggests that PAS-LuxR regulators may

affect a plethora of processes previously unforeseen. In this sense, the introduction of PAS-

LuxR regulatory genes into different *Streptomyces* hosts has already proven useful for the

awakening of dormant secondary metabolite biosynthetic genes (Olano et al., 2014; Martínez-

Burgo et al., 2019).

Here we have used microarrays to study the transcriptome of *S. avermitilis* Δ*pteF* mutant

in comparison with that of its parental strain in order to deepen our knowledge about the

processes in which PteF is involved.

MATERIALS AND METHODS

Strains and cultivation

S. avermitilis NRRL 8165 and its mutant *S. avermitilis* Δ*pteF* (Vicente et al., 2014) were

routinely grown in YEME medium (Kieser et al., 2000) without sucrose. Sporulation was

achieved in TBO medium (Higgens et al. 1974) at 30˚C.

Nucleic acid extractions.

RNA was extracted as described elsewhere (Vicente et al., 2014). Briefly, 2 ml from liquid cultures in YEME medium without sucrose were harvested by centrifugation and immediately frozen by immersion in liquid nitrogen. Cells were resuspended in lysis solution [600 µl RLT buffer (RNeasy mini kit; Qiagen); 6 µl 2-mercaptoethanol] and disrupted using sonicator (Ultrasonic processor XL; Misonix Inc.). RNeasy ® Mini kit (Qiagen) was used for RNA isolation using RNase-Free DNase Set (Qiagen) as specified by manufacturer, followed by two consecutive digestions with TURBO^{TM} DNase from Ambion® according to the manufacturer's instructions. Total RNA concentration was determined with a NanoDrop ND-1000 spectrophotometer (Thermo Scientific), and quality and integrity were checked in a Bioanalyzer 2100 apparatus (Agilent Technologies). Total genomic DNA (gDNA) was isolated from stationary phase cultures following the salting-out procedure (Kieser et al., 2000).

Microarray hybridizations.

The microarray experiment was performed using a common reference design (Gadgil et al. 2005). The microarray chip Custom Gene Expression Microarray, 8x15K (Agilent) was customized in order to include different sets of probes as indicated elsewhere (Beites et al., 2014). For each microarray hybridization, 10 pmol of Cy3-labelled cDNA obtained from total RNA were mixed with 80 pmol of Cy5-labelled genomic DNA as the common reference. Labelling, hybridization, washing and scanning conditions were carried out as indicated previously (Rodríguez-García et al. 2007; Guerra et al., 2012). Three biological replicates from independent cultures were made for each experimental condition.

Identification of differentially transcribed genes.

Microarray data were normalized and analysed with the Bioconductor package LIMMA (Linear Models for Microarray Analysis) (Smyth 2004; Smyth et al. 2005). Spot quality weights

were estimated as indicated in the Supplementary section (Tables S1 and S2). Both local and global normalizations were used (Wu et al., 2005). Firstly, weighted medians of $\log_2$ Cy3/Cy5 intensities were calculated for print-tip correction and afterwards global Loess was applied (Smyth and Speed 2003). The normalized $\log_2$ of Cy3/Cy5 intensities is referred in this work as the Mg value, which is proportional to the abundance of transcripts for a particular gene (Mehra et al., 2006). The information from within-array spot duplicates (Smyth et al., 2005) and empirical array weights (Ritchie et al., 2006) were taken into account in the linear models (Smyth 2004). The Mg transcription values of the four experimental conditions were compared using two contrasts, mutant versus wild type, corresponding to the two studied growth phases (exponential and stationary). For each gene, the Mc value is the binary log of the differential transcription between the mutant and the wild strain. The Benjamini-Hochberg (BH) false-discovery rate correction was applied to the p-values. A positive Mc value indicates upregulation, and a negative one, downregulation. For each contrast a result was considered as statistically significant if the BH-corrected p-value was <0.05. In certain occasions, however, when the transcription profile of a gene matched that of genes statistically significant and functionally related, or for comparison with previous published results obtained by RT-qPCR or by EMSA assays (Vicente et al., 2014, 2015), we used an uncorrected p-value with a level of significance <0.05.

The microarray data are deposited in the National Center for Biotechnology Information-Gene Expression Omnibus under accession number GSE185887.

Assessment of filipin and oligomycin production.

Filipin production was quantified as described elsewhere (Barreales et al., 2020), whereas oligomycin was measured following the procedure described by Vicente et al (2015).

Reverse transcription-quantitative PCR.

Reverse transcription of total RNA was performed on selected samples with 5 µg of RNA and 12.5 ng/µl of random hexamer primer (Invitrogen) using SuperScript™ III reverse transcriptase (Invitrogen) as described previously (Barreales et al., 2018). Reactions were carried out on two biological replicates with three technical replicates each and appropriate controls were included to verify the absence of gDNA contamination in RNA and primer-dimer formation. Primers (see Table S3) were designed to generate PCR products between 97 and 153 bp, near the 5´ end of mRNA. The PCR reactions were initiated by incubating the sample at 95°C for 10 min followed by 40 cycles at 95°C for 15 s, 62-70°C (depending of the set of primers used) for 34 s, and 72 °C for 30 s. To check the specificity of real-time PCR reactions, a DNA melting curve analysis was performed by holding the sample at 60°C for 60 s followed by slow ramping of the temperature to 95°C. Baseline and threshold values were determined by the StepOnePlus software. C_t values were normalized with respect to *rrnA1* mRNA (encoding 16S rRNA). Relative changes in gene expression were quantified using the Pfaffl method (2001) and the REST© software (Pfaffl et al., 2002). The corresponding real-time PCR efficiency (E) of one cycle in the exponential phase was calculated according to the equation $E = 10^{[-1/\text{slope}]}$ (Rasmussen, 2000) using 5-fold dilutions of genomic DNA ranging from 0.013 to 40 ng (n=5 or 6 with three replicates for each dilution) with a coefficient of determination $R^2 >$ 0.99 (Fig. S1).

RESULTS AND DISCUSSION

Identification of genes with an altered expression profile in *S. avermitilis ΔpteF* mutant.

S. avermitilis ΔpteF and its parental strain *S. avermitilis* NRRL 8165 were grown in YEME medium without sucrose, and samples were taken at the end of the exponential and at the middle of the stationary growth phases (Fig. 1). Transcriptomic analysis was performed by

microarray hybridization to assess the genes with an altered expression in the mutant when compared with the parental strain. Genomic DNA was used as a universal reference for all hybridizations. A result was considered as statistically significant if the BH-corrected *p*-value was <0.05. It is worth noting that these conditions are quite stringent, given that genes that constitute direct targets of PteF (e.g. the filipin polyketide synthases *pteA1* and *pteA2*; Vicente et al., 2014) are not statistically significant. With this criterion, microarrays analysis showed significant differences (with a fold change above or below +/-2) in the expression of 208 genes of the *pteF*-negative mutant at the end of the exponential phase, and 99 at the stationary phase of growth (Table 1; Fig. 2).

Surprisingly, the lack of PteF resulted in the overexpression of a majority of the differentially transcribed genes, at both sampling times, thus indicating that this regulator acts as a negative modulator for those genes expression. This was unexpected given that PteF is an activator of both the antifungal filipin (Vicente et al., 2014) and the ATP-synthase inhibitor oligomycin (Vicente et al., 2015) biosynthesis.

These genes were related to different cellular processes, including genetic information processing; energy, carbohydrate, and lipid metabolism; DNA replication and repair; morphological differentiation; and transcriptional regulation, among others, but particularly to secondary metabolite biosynthesis (Table 1).

Genes involved in genetic information- and protein-processing, and amino acid metabolism.

This group includes 24 genes that showed differential transcription in at least one of the sampling times (Table 1). These genes code for enzymes involved in amino acid metabolism (7 genes), proteins involved in transcription (8 genes, including 5 sigma factors), the ribosomal protein L28 (*SAV2675*), two putative acetyltransferases of ribosomal proteins (*SAV703* and *SAV758*), and enzymes involved in protein processing (5 genes) (Table S4).

Interestingly, while sigma factors *sig10* (*SAV898*), *sig13* (*SAV997*), and *sig60*

(*SAV213*), and ribosomal proteins acetyltransferases *SAV703* and *SAV758* showed increased

transcription levels in the mutant, *sig32* (*SAV3888*), *sig40* (*SAV4561*), the L28 ribosomal

protein encoding gene *rpmB1*, and the *whiB*-like transcriptional factor *wblE* were clearly

underexpressed in the mutant. The Wbl family of transcriptional factors is exclusive of

actinobacteria and their members have been correlated with diverse roles in morphological

differentiation and secondary metabolism (Fowler-Goldsworthy et al., 2011; Bush 2018).

Noteworthy, genes *rocA* (*SAV2723*) and *putA* (*SAV2724*), that encode delta-1-pyrroline-

5-carboxylate dehydrogenase and proline dehydrogenase, respectively, and that have been

related to proline catabolism (Menzel and Roth, 1981), and *rocD2* (*SAV7112*) and *SAV4551*,

which encode putative ornithine aminotransferases, and are also involved in proline metabolism

were underexpressed in the mutant, while *leuB* (*SAV2718*) which has been involved in valine,

leucine, and isoleucine biosynthesis biosynthesis, *paaI* (*SAV1986*) that encodes a phenylacetic

acid thioesterase, and putative cysteine desulfurase *SAV1061* were overexpressed.

<u>Genes involved in nucleotide and vitamin metabolism, and DNA replication, recombination,</u>

<u>and repair</u>.

Eighteen genes falling into this category were found to be differentially transcribed in

the mutant (Table 1). Ten of them are involved in DNA replication, recombination and repair.

Of these, seven putative transposases belonging to different families showed enhanced

transcription in the mutant. Additionally, two genes involved in DNA repair, *ku2* (*SAV879*)

which is probably involved in non-homologous DNA end-joining (Zhang et al., 2012), and

uvrD1 (*SAV3463*) that codes for a putative ATP-dependent helicase, were also upregulated.

Conversely, *int12* (*SAV4626*), which encodes a tyrosine-family recombinase/integrase, showed

reduced transcription levels at stationary phase.

The remaining genes were differentially transcribed only in the exponential phase. Four

genes are involved in vitamin metabolism, three of them with lower transcription in the mutant,

including cobalamin methylase *cobJ* (*SAV6407*) and adenosyltransferase *cobA* (*SAV6413*), and

alkaline phosphatase *phoA* (*SAV5915*), which besides being part of the PhoRP two-component

system (Sola-landa et al., 2008) is also involved in folate metabolism. The fourth gene, *thiC*

(*SAV4265*) is a thiamine biosynthesis protein (Table S4). The rest of the genes are involved in

purine metabolism, including *pgmA*, *purA*, and *purN*, all with enhanced transcription, and *cpdB*,

with lower transcription.

<u>Carbohydrate metabolism genes</u>.

Thirteen genes fall into this category, including four most likely belonging to the same

operon (*SAV1009*, *galE5*, *mpg2*, *SAV1014*) and putatively involved in galactose metabolism,

and showing enhanced transcription in the mutant. Other genes involved in the metabolism of

this sugar were the alpha-galactosidase *agaB1* (*SAV1082*), which was underexpressed in the

mutant, and the phosphoglucomutase *pgmA* (*SAV803*), which showed the opposite behavior.

Interestingly, three genes of the tricarboxylic acid/glyoxylate cycle (citrate synthase *citA2*,

citrate lyase *citE2*, and methylmalonyl-CoA mutase *meaA1*) were overexpressed in the mutant

(Table S4).

<u>Lipid metabolism genes</u>.

Nine genes related to lipid metabolism were differentially transcribed. These include

the putative 3-oxoacyl-ACP synthase II *fabB2* (*SAV2944*), the acyl carrier protein *fabC4*

(*SAV217*), the enoyl-CoA hydratase *echA1* (*SAV492*), and the acetyl/propionyl CoA

carboxylase alpha subunit *accA2* (*SAV3866*), which are all presumably involved in fatty acid

biosynthesis, and the 1-acylglycerol-3-phosphate O-acyltransferase *plsC1* (*SAV1485*)

putatively involved in glycerophospholipid biosynthesis, among others. Interestingly, all these

genes showed increased transcription in the mutant during the exponential phase except *fabB2*

which was underexpressed (Table S4). However, during the stationary phase *fabB2* also showed

enhanced transcription.

Noteworthy, the direct binding of the PteF orthologue PimM to the promoters of two of

these genes was already demonstrated (Vicente et al., 2015), thus they have been included in

Table S4 although they did not meet the statistical criteria. These were the acyltransferase *plsC1*

(Yao and Rock 2013) whose transcription was increased in the mutant (Mc 0.88, uncorrected *p*

value 0.0471), and *fabB2* whose transcription was reduced (Mc -0.84, uncorrected p-value

0.0410 in t1) or increased (Mc 1.12, *p*-value 0.0048 in t2) depending on the growth phase.

<u>Energy production genes</u>.

Only three genes belonging to this group were found to be differentially transcribed in

the mutant. All of them involved in oxidative phosphorylation and with reduced transcription

in the mutant, two belonging to the operon *nuo* (*nuoJ1* and *nuoK1*), and the ATP synthase *atpF*

(Table S4). Interestingly, all the genes belonging to the *nuo* operon (*SAV4837-SAV4850*),

although in several cases not meeting the statistical criteria, showed the same decreased

transcription profile in the mutant.

<u>Transport and external signals processing</u>.

This group includes 25 genes that showed differential transcription in at least one of the

sampling times (Table 1). Interestingly, twelve of them code or participate in the formation of

ATP-binding cassette transporters (Table S4). Of these, four are putatively involved in sugar

transport (*SAV1804*, *SAV2246*, *SAV2247*, and *SAV2609*) and showed reduced transcription in

the mutant.

Four transporters belonging to the major facilitator superfamily showed differential

transcription in the mutant, *SAV2455* with reduced transcription, and *SAV610*, the sulfate

transporter *SAV4600, and SAV6941* with enhanced transcription.

Noteworthy, in agreement with the enhanced transcription of *SAV610*, the genes *fecC1*

(*SAV600*) and *fecB* (*SAV602*) which constitute part of a putative ABC transporter

iron(III)/siderophore transport system were also overexpressed. Based on protein similarity,

SAV600-602 could constitute an ABC transport system homologous to the system FecBCD

from *E. coli* involved in iron dicitrate transport (Staudenmaier et al., 1989). *SAV602* and

SAV610 genes flank a gene cluster involved in the biosynthesis of the siderophore nrp6 whose

expression is also upregulated in the mutant (see below and Tables 2 and S4). Altogether these

results suggest that the ABC system SAV600-602 and the transporter SAV610, would be

involved in iron transport using the siderophore nrp6. These transcriptomic results are further

supported by the direct binding of PimM to the promoters of *SAV602* and *SAV610* (Vicente et

al., 2015).

Genes involved in cell envelope biosynthesis and morphological differentiation.

This group includes eleven genes that showed differential transcription in at least one

of the sampling times. These genes code for enzymes involved in cell envelope biosynthesis

(the N-acetylmuramoyl-L-alanine amidase *ampD1*), and morphological differentiation (8

genes). The latter are particularly interesting because in *Streptomyces* morphological

differentiation is usually accompanied by physiological differentiation (McCormick and

Flärdh, 2012). The differential expression of genes involved in morphological differentiation

was somehow expected given that *S. avermitilis ΔpteF* mutants show a delay in spore formation

(Vicente et al., 2014).

Our results indicate that the transcriptional regulators *wlbE* and *bldC* that are associated

to deficient phenotypes in spore formation (*white*) and in aerial mycelium development (*bald*),

respectively, are underexpressed in the mutant. Similarly, the secreted subtilisin inhibitor *sit2* involved in morphological differentiation via *sigU* in *S. coelicolor* (Gordon et al., 2008), and *SAV2505* that encodes a DNA-binding protein orthologous to *S. lividans* transcriptional regulator ClgR which controls the expression of ATP-dependent protease Clp involved in morphological differentiation (Bellier et al., 2006), are also downregulated (Table S4). Interestingly, *clpC1* gene had also been proposed as direct PteF molecular target given PimM binding to its coding region (Vicente et al., 2015).

Conversely, the gene *ctpB* that encodes a cation-transporting P-type ATPase involved in *Bacillus subtilis* sporulation activation (Campo and Rudner, 2007), the gene *mreC* needed for spore cell-wall synthesis in *S. coelicolor* (Kleinschnitz et al., 2011), and both *kipI*, and its antagonist *kipA*, which have been involved in sporulation control in *B. subtilis* (Wang et al., 1997; Jacques et al., 2011), showed enhanced transcription in the mutant (Table S4).

Regulatory genes.

As described here, a large set of genes with diverse functions are under the control of PteF, including several regulatory genes listed in the categories described above. This prompted us to analyze other possible transcriptional regulators differentially expressed in the mutants, as these could be mediators of the regulatory control. A complete list of regulatory genes whose expression is affected in the mutant is presented in Table S4.

A total of 31 transcriptional regulators showed a significant differential transcription in the mutant when compared with the parental strain. Such a large number reflects the pleiotropic nature of PAS-LuxR regulators (Vicente et al., 2014, 2015; Aparicio et al., 2016), and probably justifies all the biological processes affected by the mutation (see functional categories listed above).

Among the regulators controlled by PteF, is interesting to highlight eight directly involved in diverse secondary metabolites biosynthesis control, namely: *avaL2* (*SAV2268*) and

avaL1 (*SAV2270*), both TetR-family regulators putatively involved in the biosynthesis of a γ-butyrolactone (Ikeda et al., 2014); *avaR1* (*SAV3705*), which encodes the avenolide receptor protein (Kitani et al., 2011; Wang et al., 2014; Zhu et al., 2016); *olmRII* (*SAV2901*) and *olmRI* (*SAV2902*), both LuxR-family positive regulators of macrolide oligomycin biosynthesis (Yu et al., 2012); *pteR* (*SAV410*), the SARP-LAL regulator of the polyene macrolide filipin biosynthesis (Ikeda et al., 2014; Vicente et al., 2014; Payero et al., 2015); *aveR* (*SAV935*), a LAL-family positive regulator of avermectin biosynthesis (Kitani et al., 2009); and *SAV2989*, a MarR-family transcriptional regulator from the neopentalenolactone biosynthetic cluster (Ikeda et al., 2014). All these regulatory genes showed decreased transcription in the mutant, except for *pteR* and *aveR* that were overexpressed (Tables 2 and S4).

Interestingly, the expression of *olmRI* and *olmRII* genes had already been proven to be negatively affected by the lack of PteF (Vicente et al., 2015). Furthermore, *pteF*-deletion mutants showed a severe loss of oligomycin production, whereas gene complementation of the mutant restored parental-strain phenotype, and gene duplication in the wild-type strain boosted oligomycin production (Vicente et al., 2015). Similarly, *pteR* has also been reported as a PteF molecular target, via the action of another hierarchical regulator which would be activated by PteF (Vicente et al., 2014).

Besides the abovementioned regulators, it is also noteworthy the identification of *SAV2301*, that codes for a RedD orthologue, the transcriptional activator of the undecylprodigiosin pathway in *S. coelicolor* (Narva and Feitelson, 1990), *bldC* (*SAV4130*), a MerR-family regulator involved in morphological differentiation and secondary metabolite production in *S. coelicolor* (Hunt et al., 2005), and *cutS* (*SAV2404*), a sensor kinase involved in actinorhodin biosynthesis in *S. lividans* (Chang et al., 1996), all of them being down-regulated in the mutant (Table S4).

Secondary metabolite genes.

The functional group more clearly affected by *pteF* deletion was that of genes involved

in secondary metabolite biosynthesis (Table 1). In this category, when one or more genes

critical for metabolite biosynthesis were found statistically significant, the transcription of other

genes belonging to the same cluster with uncorrected p-values < 0.05 was also considered

significant. Following this broader criterion, sixty one genes belonging to this group, regardless

of regulatory genes mentioned above, showed a significant differential transcription in the

mutant when compared with the parental strain in at least one of the sampling times (Table S4).

Noteworthy, almost all genes were detected at the exponential growth-phase. In particular,

those related to secondary metabolism precursor biosynthesis were only detected at this

sampling time. These were: the ornithine aminotransferases *rocD3* (*SAV2285*) and *rocD2*

(*SAV7112*), and the proline dehydrogenase *putA* (*SAV2724*), which were underexpressed; and

the phosphoglucomutase *pgmA* (*SAV803*), the 3-isopropylmalate dehydrogenase *leuB*

(*SAV2718*), the phosphoribosylglycinamide formyltransferase *purN* (*SAV3445*), and the

putative citrate synthase *citA2* (*SAV3859*), which were overexpressed.

But the most striking results of microarray analyses were the identification of

differential transcription in 67 genes (including regulatory genes) belonging to 10 out of the 38

putative secondary metabolite gene clusters encoded by *S. avermitilis* genome (Ikeda et al.,

2014). Table 2 includes the transcriptional values of genes belonging to differentially expressed

secondary metabolite gene clusters. For gene cluster boundaries definition we used StrepDB

database (http://strepdb.streptomyces.org.uk) in conjunction with information described by

Ikeda et al. (2014).

The secondary metabolites whose biosynthesis would be affected by *pteF* deletion were

of different nature, and included: the polyketides filipin (*pte*), oligomycin (*olm*), avermectin

(*ave*), and the product of *pks3*; the non-ribosomal peptides nrp3 and the siderophore nrp6; the

vibrioferrin-like polyhydroxycarboxylate siderophore *avs*; the terpenoid

neopentalenoketolactone (*ptl*); the γ-butyrolactone (*gbl*); and melanin (*melC-1*).

In all these clusters, the differential transcription of at least one key biosynthetic gene

was observed. The number of genes affected were: eleven in the *nrp6* cluster (out of 12), ten

(out of 13 and 14 respectively) in the case of the filipin and oligomycin clusters, eight (out of

11) in the case of the *pk3* cluster, seven in the case of the avermectin (out of 19) cluster, six in

the *nrp3* cluster (out of 10), six in the *ptl* cluster (out of 14), five (out of 5) in the *gbl* cluster,

and two in the *avs* (out of 4) and melanin *melC-1* (out of 2) clusters (Table 2).

Furthermore, a closer look at the transcription of the remaining genes of each of these

clusters revealed that most of the genes of a given cluster, followed the same tendency. Figure

3 shows the transcription profiles of secondary metabolite gene clusters genes affected by the

mutation including regulatory genes, and Table 2 the transcription values observed for each of

the genes.

Seven of the secondary metabolite gene clusters showed an overall reduced

transcription, including filipin *pte*, oligomycin *olm*, neopentalenoketolactone *ptl*, and melanin

melC-1 clusters, the silent cluster for γ-butyrolactone *gbl*, and the cryptic gene clusters *pk3* and

nrp3. On the opposite, three gene clusters showed overall enhanced transcription, including the

macrolide avermectin *ave*, the siderophore *avs*, and the cryptic non-ribosomal peptide *nrp6*

(Fig. 3).

Interestingly, besides the genes mentioned above, all the genes belonging to the clusters

coding for the terpenoid albaflavenol/albaflavenone (*ezs*), and the cryptic polyketide *pk4* also

followed the same tendency. In these cases, transcription values did not meet the statistical

criteria, but their uncorrected *p*-values were < 0.05 in all instances (Table 2). In the case of the

ezs genes (SAV3031-3032) they showed an average of two fold more transcription in the

mutant, whereas *pk4* genes (*SAV7184-7186*) showed between 7 and 9 fold less transcription

than in the parental strain.

Filipin and Oligomycin production are strongly reduced in *S. avermitilis ΔpteF*.

Although many of the metabolites whose biosynthesis would be affected by *pteF* deletion are of unknown structure (cryptic) and others are not produced under laboratory conditions (silent) (Ikeda et al., 2014), the production of two of them could be readily monitored in *S. avermitilis ΔpteF*. These were the antifungal pentaene filipin which is encoded by the *pte* cluster where the regulator is situated, and the ATP-synthase inhibitor oligomycin which is encoded by the *olm* cluster (Fig. 1). In both cases, production of the secondary metabolite was strongly reduced upon inactivation of the regulatory gene *pteF*. This is in agreement with the reduced transcription of most biosynthetic genes of both clusters (Fig. 3). The exceptions were the discrete thioesterase *pteH*, the cholesterol oxidase *pteG*, and the SARP-LAL regulator *pteR* of the filipin cluster, which were overexpressed. These results corroborate our previous observations by RT-qPCR (Vicente et al., 2014; Vicente et al., 2015).

Validation of microarray results by using quantitative RT-PCR.

Quantitative RT-PCR was used on reversed transcribed RNA samples to confirm that differential expression indicated by the microarray data was supported by an independent method. The selected genes covered a wide range of expression, including up-regulation and down-regulation. Twelve genes were validated including genes for the biosynthesis of filipin (*pteC*, *pteB*, *pteR*, *pteG*), oligomycin (*olmRI, olmRII, olmB*), avermectin (*aveR*), the isomerase of *pk3* cluster (*SAV2273*), one ABC transporter of the *nrp6* cluster (*fecB*), the alpha galactosidase *agaB1*, and the heat shock internal membrane protease *htpX1* (*SAV4891*).

Overall, the RT-qPCR data and microarray data showed a good concordance (Fig. S2). The range of dynamics for relative $\log_2$ fold change obtained from RT-qPCRs (-6.53 to +7.54) was higher than that obtained from Mc values from microarrays (-7.24 to +2.94), indicating that RT-qPCRs are more sensitive. This probably reflects on the Pearson's correlation

coefficient (R^2) for the plot, resulting in a lower value than what could be expected.

Nevertheless, the obtained value ($R^2 = 0.892$) still indicates a good correlation of results.

Concluding remarks.

Up to date, PAS-LuxR regulator-encoding genes have been found only in polyene

macrolide gene clusters, thus constituting a landmark of these type of clusters. In this context,

they are transcriptional activators essential for the biosynthesis of the polyene encoded within

the cluster. Their expression is a bottleneck in the biosynthesis of the antifungal and thus

polyene production is easily incremented upon gene dosage increase (Aparicio et al., 2016).

Additionally, heterologous gene complementation of mutants restores strain ability to produce

the antifungal compound, thus proving that these regulators are highly conserved (Santos-

Aberturas et al. 2011b). Recently, we have obtained evidence indicating that although these

regulators were initially thought to be pathway-specific, they actually are regulatory proteins

with a wider range of connotations in addition to polyene biosynthesis. Thus, PteF, the regulator

of filipin biosynthesis, was proven to control oligomycin production in *S. avermitilis* (Vicente

et al., 2015). This prompted us to propose that introduction of PAS-LuxR regulatory genes into

Streptomyces species could prove useful for the awakening of dormant secondary metabolite

biosynthetic genes (Vicente et al., 2014; 2015). This hypothesis was confirmed when PimM,

the archetype of PAS-LuxR regulators was introduced into *S. albus* J1074, and production of

the hybrid non ribosomal peptide-polyketide antimycin was activated (Olano et al., 2014).

Recently, a similar result has been described in *S. albus* S4, where a PimM orthologue (the

candicidin regulator FscRI) has been identified as required for antimycin production (McLean

et al., 2016).

Here we have studied the transcriptome of *S. avermitilis* Δ*pteF* mutant in comparison

with that of its parental strain. Our results corroborate our previous observations (Vicente et al.,

2014, 2015), reinforcing the idea that PAS-LuxR regulators control many different cellular

processes of bacterial metabolism at the transcriptional level, but in particular stress the

importance of PAS-LuxR involvement on secondary metabolite biosynthesis.

Notably, ten (or twelve if we include *ezs* and *pk4* gene clusters) out of the 38 putative

secondary metabolite gene clusters encoded by *S. avermitilis* genome (Ikeda et al., 2014)

showed altered expression in the mutant. In some instances, the modified expression of

biosynthetic genes of a given cluster could be explained by the effect of the mutation on the

expression of one or more cluster-situated regulators. This is the case of the *aveR* regulator of

the avermectin *ave* cluster, the regulators *avaL1* and *avaL2* of the γ-butyrolactone *gbl* cluster,

the oligomycin regulators *olmRI* and *olmRII*, and the MarR regulator (*SAV2989*) of the

pentalenolactone *ptl* cluster. AveR, the transcriptional activator of avermectin biosynthesis

(Kitani et al., 2009), is overexpressed four-fold in the mutant and concomitantly the remaining

genes of the *ave* cluster showed enhanced transcription. Conversely, OlmRI and OlmRI,

positive regulators of oligomycin biosynthesis (Yu et al., 2012), showed decreased transcription

in the mutant (Mc values -1.56 and -1.47 respectively), and so did the remaining genes of the

cluster. It is not known whether AvaL1 and AvaL2 are positive regulators, but it is conceivable

given that they show reduced transcription values upon mutation of the *pteF* gene (fold changes

of 6.4 and 8.9, respectively) together with the remaining genes of the *gbl* cluster, including the

γ-butyrolactone synthase *avaA*. Both AvaL1 and AvaL2 show convincing similarity to γ-

butyrolactone receptor proteins, and although these proteins normally act repressing

transcription of the synthase gene (Zou et al., 2014; Zhou et al., 2015; Barreales et al., 2020),

there are cases that display the opposite behavior, like FarA from *S. lavendulae* that activates

the transcription of the synthase *farX* (Kitani et al., 2010). The same occurs with the MarR

regulator of the *ptl* cluster (Ikeda et al., 2014) whose transcription is diminished (2-fold) in the

mutant as well as that of all *ptl* genes. In the remaining gene clusters there are no cluster-situated

regulatory genes, thus the effect of the mutation must be explained either by a direct action of

PteF on key biosynthetic genes or via the action of other regulatory genes. In this sense, 28

regulatory genes not situated in the clusters indicated above, most of them with unknown

function, were differentially expressed upon mutation of *pteF* (Table S4).

To our knowledge, this is the second time a genome-wide transcriptomic study is

conducted to describe the pleiotropic nature of a cluster-situated regulator, that of the regulator

of lincomycin biosynthesis LmbU from *S. linconensis* (Li et al., 2020). Cross-regulation of

disparate natural-product biosynthetic gene clusters by a cluster-situated regulator has already

been described by several groups although not in genome-wide studies (Santamarta et al., 2011;

Vicente et al., 2015; McLean et al., 2016). Moreover, the ability of some of these regulators to

modulate the effects of regulators that act more globally (Huang et al., 2005), as well as the

competition between global regulators (Santos-Beneit et al., 2009), have also been reported.

Our findings go beyond, and indicate that PAS-LuxR regulators should be considered wide

domain regulators. They affect the expression of multiple genes involved in both primary and

secondary metabolism.

The findings reported here should provide important clues to understand the intertwined

regulatory machinery that modulates antibiotic biosynthesis in *Streptomyces*, and suggest that

the heterologous expression of PAS-LuxR regulators is likely to represent a powerful general

strategy for novel bioactive natural product discovery.

ACKNOWLEDGEMENTS

This work was supported by the Spanish Ministerio de Economía y Competitividad (Grant

BIO2013-42983-P to JFA), F.P.U. fellowships of the Ministerio de Educación, Cultura y

Deporte (AP2007-02055 to TDP, FPU13/01537 to AP), a contract from the Junta de Castilla y

León co-financed by the European Social Fund (to EGB), and a fellowship from the Portuguese

Fundação para a Ciência e a Tecnologia (SFRH/BD/64006/2009 to CMV).

CONFLICT OF INTEREST:

The authors declare that they have no conflict of interest.

REFERENCES

Antón N, Santos-Aberturas J, Mendes MV, Guerra SM, Martín JF, Aparicio JF. 2007. PimM, a PAS domain positive regulator of pimaricin biosynthesis in *Streptomyces natalensis*. Microbiology 53, 3174-3183. doi: 10.1099/mic.0.2007 /009126-0

Aparicio JF, Barreales EG, Payero TD, Vicente CM, de Pedro A, Santos-Aberturas J. 2016. Biotechnological production and application of the antibiotic pimaricin: biosynthesis and its regulation. Appl Microbiol Biotechnol 100, 61-78. doi: 10.1007/s00253-015-7077-0

Barreales EG, Payero TD, Jambrina E, Aparicio JF. 2020. The gamma-butyrolactone system from Streptomyces filipinensis reveals novel clues to understand secondary metabolism control. Appl Environ Microbiol 86, e00443-20. doi: 10.1128/AEM.00443-20

Barreales EG, Vicente CM, de Pedro A, Santos-Aberturas J, Aparicio JF. 2018. Promoter engineering reveals the importance of heptameric direct repeats for DNA-binding by SARP-LAL regulators in Streptomyces natalensis. Appl Environ Microbiol 84, e00246-18. doi: 10.1128/AEM.00246-18

Beites T, Rodríguez-García A, Santos-Beneit F, Moradas-Ferreira P, Aparicio JF, Mendes MV. 2014. Genome-wide analysis of the regulation of pimaricin production in *Streptomyces natalensis* by reactive oxygen species. Appl Microbiol Biotechnol 98, 2231–2241. doi: 10.1007/s00253-013-5455-z

Bellier A, Gominet M, Mazodier P. 2006. Post-translational control of the *Streptomyces lividans* ClgR regulon by ClpP. Microbiology 152, 1021-1027. doi: 10.1099/mic.0.28564-0

Bush M. 2018. The actinobacterial WhiB-like (Wbl) family of transcription factors. Mol Microbiol 110, 663-676. doi: 10.1111/mmi.14117

Campo N, Rudner DZ. 2007. SpoIVB and CtpB are both forespore signals in the activation of the sporulation transcription factor sigmaK in *Bacillus subtilis*. J Bacteriol 189, 6021–6027. doi: 10.1128/JB.00399-07

Chang HM, Chen MY, Shieh YT, Bibb MJ, Chen CW. 1996. The *cutRS* signal transduction system of *Streptomyces lividans* represses the biosynthesis of the polyketide antibiotic actinorhodin. Mol Microbiol 21, 1075-1085.

Fowler-Goldsworthy K, Gust B, Mouz S, Chandra G, Findlay KC, Chater KF. 2011. The actinobacteria-specific gene *wblA* controls major developmental transitions in *Streptomyces coelicolor* A3(2). Microbiology 157, 1312–1328. doi: 10.1099/mic.0.047555-0

Gadgil M, LianW, Gadgil C, Kapur V, HuWS. 2005. An analysis of the use of genomic DNA
as a universal reference in two channel DNA microarrays. BMC Genomics 6, 66.
doi:10.1186/1471-2164-6-66

Gordon ND, Ottaviano GL, Connell SE, Tobkin GV, Son CH, Shterental S, Gehring AM. 2008.
Secreted-protein response to sigmaU activity in *Streptomyces coelicolor*. J Bacteriol 190,
894–904. doi: 10.1128/JB.01759-07

Guerra SM, Rodríguez-García A, Santos-Aberturas J, Vicente CM, Payero TD, Martín JF,
Aparicio JF. 2012. LAL regulators SCO0877 and SCO7173 as pleiotropic modulators of
phosphate starvation response and actinorhodin biosynthesis in *Streptomyces coelicolor*.
PLoS One 7, e31475. doi: 10.1371/journal.pone.0031475

Hefti MH, Francoijs K-J, de Vries SC, Dixon R, Vervoort J. 2004. The PAS fold. A redefinition
of the PAS domain based upon structural prediction. Eur J Biochem 271, 1198-1208. doi:
10.1111/j.1432-1033.2004.04023.x

Higgens CE, Hamill RL, Sands TH, Hoehn MM, Davis NE. 1974. The occurrence of
deacetoxycephalosporin C in fungi and streptomycetes. J Antibiot 27, 298-300.

Huang J, Shi J, Molle V, Sohlberg B, Weaver D, Bibb MJ, Karoonuthaisiri N, Lih CJ, Kao CM,
Buttner MJ, Cohen SN. 2005. Cross-regulation among disparate antibiotic biosynthetic
pathways of *Streptomyces coelicolor*. Mol Microbiol 58:1276–1287. doi:10.1111/j.1365-
2958.2005.04879.x

Hunt AC, Servín-González L, Kelemen GH, Buttner MJ. 2005. The *bldC* developmental locus
of *Streptomyces coelicolor* encodes a member of a family of small DNA-binding proteins
related to the DNA-binding domains of the MerR family. J Bacteriol 187, 716-728. doi:
10.1128/JB.187.2.716-728.2005

Ikeda H, Shin-ya K, Omura S. 2014. Genome mining of the *Streptomyces avermitilis* genome
and development of genome-minimized hosts for heterologous expression of biosynthetic
gene clusters. J Ind Microbiol Biotechnol 41, 233-250. doi: 10.1007/s10295-013-1327-x

Jacques DA, Langley DB, Hynson RMG, Whitten AE, Kwan A, Guss JM, Trewhella J. 2011.
A novel structure of an antikinase and its inhibitor. J Mol Biol 405, 214–226. doi:
10.1016/j.jmb.2010.10.047

Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA. 2000. Practical *Streptomyces*
genetics. John Innes Foundation, Norwich

Kitani S, Ikeda H, Sakamoto T, Noguchi S, Nihira T. 2009. Characterization of a regulatory
gene, *aveR*, for the biosynthesis of avermectin in *Streptomyces avermitilis*. Appl Microbiol
Biotechnol 82, 1089-1096. doi: 10.1007/s00253-008-1850-2

Kitani S, Doi M, Shimizu T, Maeda A, Nihira T. 2010. Control of secondary metabolism by
farX, which is involved in the gamma-butyrolactone biosynthesis of *Streptomyces*
lavendulae FRI-5. Arch Microbiol 192, 211-220. doi: 10.1007/s00203-010-0550-3.

Kitani S, Miyamoto KT, Takamatsu S, Herawati E, Iguchi H, Nishitomi K, Uchida M,
Nagamitsu T, Omura S, Ikeda H, Nihira T. 2011. Avenolide, a *Streptomyces* hormone
controlling antibiotic production in *Streptomyces avermitilis*. Proc Natl Acad Sci U S A.
108, 16410-16415. doi: 10.1073/pnas.1113908108

Kleinschnitz E-M, Heichlinger A, Schirner K, Winkler J, Latus A, Maldener I, Wohlleben W,
Muth G. 2011. Proteins encoded by the *mre* gene cluster in *Streptomyces coelicolor* A3(2)

cooperate in spore wall synthesis. Mol Microbiol 79, 1367–1379. doi: 10.1111/j.1365-
2958.2010.07529.x

Martínez-Burgo Y, Santos-Aberturas J, Rodríguez-garcía A, Barreales EG, Tormo JR, Truman
AW, Reyes F, Aparicio JF, Liras P. 2019. Activation of Secondary Metabolite Gene
Clusters in Streptomyces clavuligerus by the PimM Regulator of Streptomyces natalensis.
Front Microbiol; 10:580. doi: 10.3389/fmicb.2019.00580

McCormick JR, Flärdh K. 2012. Signals and regulators that govern Streptomyces development.
FEMS Microbiol Rev 36, 206-231. doi: 10.1111/j.1574-6976.2011.00317.x

McLean TC, Hoskisson PA, Seipke RF. 2016. Coordinate regulation of antimycin and
candicidin biosynthesis. mSphere 1:e00305-16. doi: 10.1128/mSphere.00305-16.

Mehra S, Lian W, Jayapal KP, Charaniya SP, Sherman DH, Hu WS. 2006. A framework to
analyze multiple time series data: A case study with *Streptomyces coelicolor*. J Ind
Microbiol Biotechnol 33, 159–172. doi: 10.1007/s10295-005-0034-7

Menzel R, Roth J. 1981. Purification of the *putA* gene product. A bifunctional membrane-bound
protein from *Salmonella typhimurium* responsible for the two-step oxidation of proline to
glutamate. J Biol Chem 256, 9755-9761.

Möglich A, Ayers RA, Moffat K. 2009. Structure and signaling mechanism of Per-ARNT-Sim
domains. Structure 17, 1282-1294. doi: 10.1016/j.str.2009.08.011

Narva KE, Feitelson JS. 1990. Nucleotide sequence and transcriptional analysis of the *redD*
locus of *Streptomyces coelicolor* A3(2). J Bacteriol 172, 326-333.

Olano C, García I, González A, Rodriguez M, Rozas D, Rubio J, Sánchez-Hidalgo M, Braña
AF, Méndez C, Salas JA. 2014. Activation and identification of five clusters for secondary
metabolites in *Streptomyces albus* J1074. Microb Biotechnol 7, 242-256. doi:
10.1111/1751-7915.12116

Payero TD, Vicente CM, Rumbero Á, Barreales EG, Santos-Aberturas J, de Pedro A, Aparicio
JF. 2015. Functional analysis of filipin tailoring genes from *Streptomyces filipinensis*
reveals alternative routes in filipin III biosynthesis and yields bioactive derivatives. Microb
Cell Fact 14:114. doi: 10.1186/s12934-015-0307-4.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res 29, e45. doi:10.1093/nar/29.9.e45

Pfaffl, M.W., Horgan, G.W., Dempfle, L. 2002. Relative expression software tool (REST) for
group-wise comparison and statistical analysis of relative expression results in real-time
PCR. Nucleic Acids Res 30, e36. doi: 10.1093/nar/30.9.e36

Rasmussen, R. 2000. Quantification on the LightCycler. In: Meuer S, Wittwer C, Nakagawara
K, editors. Rapid Cycle Real-Time PCR, Methods and Applications. Springer Press,
Heidelberg. pp. 21-34.

Ritchie M, Diyagama D, Neilson J, van Laar R, Dobrovic A, Holloway A, Smyth GK. 2006.
Empirical array quality weights in the analysis of microarray data. BMC Bioinformatics 7,
261. doi: 10.1186/1471-2105-7-261

Rodríguez-Garcia A, Barreiro C, Santos-Beneit F, Sola-Landa A, Martín JF. 2007. Genome-
wide transcriptomic and proteomic analysis of the primary response to phosphate limitation
in *Streptomyces coelicolor* M145 and in a DeltaphoP mutant. Proteomics 7, 2410–1429.
doi:10.1002/pmic.200600883

Santos CL, Correia-Neves M, Moradas-Ferreira P, Mendes MV. 2012. A walk into the LuxR
regulators of Actinobacteria: phylogenomic distribution and functional diversity. PLoS
One 7:e46758. doi: 10.1371/journal.pone.0046758
Santos-Aberturas J, Vicente CM, Guerra SM, Payero TD, Martín JF, Aparicio JF. 2011a.
Molecular control of polyene macrolide biosynthesis: direct binding of the regulator PimM
to eight promoters of pimaricin genes and identification of binding boxes. J Biol Chem
286, 9150-9161. doi: 10.1074/jbc.M110.182428
Santos-Aberturas J, Payero TD, Vicente CM, Guerra SM, Cañibano C, Martín JF, Aparicio JF.
2011b. Functional conservation of PAS-LuxR transcriptional regulators in polyene
macrolide biosynthesis. Metab Eng 13, 756-767. doi: 0.1016/j.ymben.2011.09.011
Santos-Beneit F, Rodríguez-García A, Sola-Landa A, Martín JF. 2009. Cross-talk between two
global regulators in *Streptomyces*: PhoP and AfsR interact in the control of *afsS*, *pstS* and
phoRP transcription. Mol Microbiol 72, 53-68. doi: 10.1111/j.1365-2958.2009.06624.x.
Smyth GK. 2004. Linear models and empirical bayes methods for assessing differential
expression in microarray experiments. Stat Appl Genet Mol Biol 3, 3. doi:10.2202/1544-
6115.1027
Smyth GK, Michaud J, Scott HS. 2005. Use of within-array replicate spots for assessing
differential expression in microarray experiments. Bioinformatics 21, 2067–2075.
doi:10.1093/bioinformatics/bti270
Smyth GK, Speed TP. 2003. Normalization of cDNA microarray data. Methods 31, 265–273.
doi:10.1016/S1046-2023(03)00155-5
Sola-Landa A, Rodríguez-García A, Apel AK, Martín JF. 2008. Target genes and structure of
the direct repeats in the DNA-binding sequences of the response regulator PhoP in
Streptomyces coelicolor. Nucleic Acids Res 36, 1358-1368. doi: 10.1093/nar/gkm1150.
Staudenmaier H, Van Hove B, Yaraghi Z, Braun V. 1989. Nucleotide sequences of the
fecBCDE genes and locations of the proteins suggest a periplasmic-binding-protein-
dependent transport mechanism for iron(III) dicitrate in *Escherichia coli*. J Bacteriol 171,
2626-2633. doi: 10.1128/jb.171.5.2626-2633.1989
Vicente CM, Santos-Aberturas J, Payero TD, Barreales EG, de Pedro A, Aparicio JF. 2014.
PAS-LuxR transcriptional control of filipin biosynthesis in *S. avermitilis*. Appl. Microbiol.
Biotechnol 98, 9311-9324. doi: 10.1007/s00253-014-5998-7
Vicente CM, Payero TD, Santos-Aberturas J, Barreales EG, de Pedro A, Aparicio JF. 2015.
Pathway-specific regulation revisited: cross-regulation of multiple disparate gene clusters
by PAS-LuxR transcriptional regulators. Appl Microbiol Biotechnol 99, 5123-5135. doi:
10.1007/s00253-015-6472-x
Wang L, Grau R, Perego M, Hoch JA. 1997. A novel histidine kinase inhibitor regulating
development in *Bacillus subtilis*. Genes Dev 11, 2569–2579. doi: 10.1101/gad.11.19.2569
Wang JB, Zhang F, Pu JY, Zhao J, Zhao QF, Tang GL. 2014. Characterization of AvaR1, an
autoregulator receptor that negatively controls avermectins production in a high
avermectin-producing strain. Biotechnol Lett 36, 813-819. doi: 10.1007/s10529-013-1416-
y
Wu W, Xing EP, Myers C, Mian IS, Bissell MJ. 2005. Evaluation of normalization methods
for cDNA microarray data by k-NN classification. BMC Bioinformatics 6, 191. doi:
10.1186/1471-2105-6-191

Yao J, Rock CO. 2013. Phosphatidic acid synthesis in bacteria. Biochim Biophys Acta. 1831,
495-502. doi: 10.1016/j.bbalip.2012.08.018

Yu Q, Bai L, Zhou X, Deng Z. 2012. Inactivation of the positive LuxR-type oligomycin
biosynthesis regulators OlmRI and OlmRII increases avermectin production in
Streptomyces avermitilis. Chinese Sci. Bull. 57, 869-876. doi: 10.1007/s11434-011-4865-
5

Zhang X, Chen W, Zhang Y, Jiang L, Chen Z, Wen Y, Li J. 2012. Deletion of *ku* homologs
increases gene targeting frequency in *Streptomyces avermitilis*. J Ind Microbiol Biotechnol
39, 917-925. doi: 10.1007/s10295-012-1097-x.

Zhou ZX, Xu QQ, Bu QT, Liu SP, Yu P, Li YQ. 2015. Transcriptome-guided identification of
SprA as a pleiotropic regulator in *Streptomyces chattanoogensis*. Appl Microbiol
Biotechnol 99:1287-1298. doi: 10.1007/s00253-014-6132-6

Zhu J, Sun D, Liu W, Chen Z, li J, Wen Y. 2016. AvaR2, a pseudo c-butyrolactone receptor
homologue from Streptomyces avermitilis, is a pleiotropic repressor of avermectin and
avenolide biosynthesis and cell growth. Mol Microbiol 102, 562-578.
doi:10.1111/mmi.13479.

Zou Z, Du D, Zhang Y, Zhang J, Niu G, Tan H. 2014. A γ-butyrolactone-sensing
activator/repressor, JadR3, controls a regulatory mini-network for jadomycin biosynthesis.
Mol Microbiol 94, 490-505. doi: 10.1111/mmi.12752.

Modulation of multiple gene clusters expression by a single PAS-LuxR transcriptional regulator.

Cláudia M. Vicente[a,b,c], Tamara D. Payero[a,b], Antonio Rodríguez-García[a,b], Eva G. Barreales[a], Antonio de Pedro[a], Fernando Santos-Beneit[b,d] and Jesús F. Aparicio[a,*]

[a] *Area of Microbiology, Faculty of Biology, University of León, 24071 León. Spain.*

[b] *Institute of Biotechnology INBIOTEC, Parque Científico de León, Avda. Real, nº 1, 24006 León, Spain.*

[c] Present address: *TBI, CNRS, INRA, INSA, Université de Toulouse, Toulouse, France*

[d] Present address: *Institute of Sustainable Processes, University of Valladolid. Spain.*

*Corresponding author. Tel:+34-987-293420; Fax:+34-987-291409; e-mail: jesus.aparicio@unileon.es

Figures 1-3, Tables 1 and 3

Fig. 1.- Growth and antibiotic production in YEME medium without sucrose. Strains *S. avermitilis* wt (black), and Δ*pteF* mutant (gray). A) Growth curves. B) Filipin production. (C) Oligomycin production. Arrows indicate RNA samples harvesting times.

A

B

Fig. 2.- Differentially expressed genes in the mutant strain Δ*pteF*. Volcano plots show differential gene expression distribution during exponential phase (A) and stationary phase (B). Statistically significant genes are shown in green ($\log_{10}$ p-value ≥ 1.3).

Fig. 3.- Transcription profiles of secondary metabolite gene clusters genes in *S. avermitilis* *ΔpteF*. Only clusters whose transcription was affected by the mutation are included. All the genes of a given cluster are shown in the plot, including regulatory genes. Coloured squares are the plots of differential transcription values for individual genes in the mutant. *pte*, filipin (red); *ave*, avermectin (orange); *olm*, oligomycin (yellow); *pk*, polyketide (gray); *nrp*, non-ribosomal peptide (green); *ptl*, neopentalenoketolactone (purple); *geo*, geosmin (teal); *ezs*, albaflavenol/albaflavenone (white); *gbl*, γ-butyrolactone (dark blue); *avs*, vibrioferrin-like siderophore (light blue); *melC*-1, melanin (black).

Table 1: Differential transcription and functional classification of genes affected by *pteF* deletion.

	Genes underexpressed[1]	Genes overexpressed[a]	Total
Exponential phase (t1)	63	145	**208**
Stationary phase (t2)	35	64	**99**
	Identified genes[2]		
Function	t1	**t2**	
Genetic information- and protein-processing, and amino acid metabolism	20 (7 ↓; 13 ↑)	11 (5 ↓; 6 ↑)	
Nucleotide and vitamin metabolism, DNA replication, recombination and repair	16 (4 ↓; 12 ↑)	4 (1 ↓; 3 ↑)	
Carbohydrate metabolism	13 (3 ↓; 10 ↑)	1 (1 ↓)	
Lipid metabolism	8 (1 ↓; 7 ↑)	4 (4 ↑)	
Energy production	2 (2 ↓)	1 (1 ↓)	
Transport and external signals processing	20 (12 ↓; 8 ↑)	8 (4 ↓; 4 ↑)	
Cell envelope biosynthesis and morphological differentiation	9 (4 ↓; 5 ↑)	5 (4 ↓; 1 ↑)	
Regulation	27 (12 ↓; 15 ↑)	12 (3 ↓; 9 ↑)	
Secondary metabolism	60 (34 ↓; 26 ↑)	6 (2 ↓; 4 ↑)	
Miscellaneous	38 (11 ↓; 27 ↑)	19 (1 ↓; 18 ↑)	

---

[1] Only statistically significant genes with a fold-change value equal higher to +/-2 are included.
[2] All identified genes were accounted for

Table 2: Transcriptional values of genes belonging to differentially expressed secondary metabolite gene clusters in *S. avermitilis ΔpteF* when compared to its parental strain. Differential transcription values *Mc* were obtained by subtracting mutant *Mg* values from parental strain *Mg* values. Only clusters whose transcription was affected by the mutation are included. The *p*-values are indicated in bold type when found statistically significant (see Materials and methods). For simplicity, designations ''putative'' have been removed.

Gene		Description	Mc	Corrected p value	p value
Filipin cluster (*pte*)					
SAV407	*pteH*	thioesterase	0.95	0.1395	**0.0076**
SAV408	*pteG*	cholesterol oxidase	**2.05**	**0.0025**	0.0000
SAV410	*pteR*	SARP-family transcriptional regulator	**2.18**	**0.0009**	0.0000
SAV411	*pteE*	ferredoxin	**-1.10**	0.2533	**0.0233**
SAV412	*pteD*	cytochrome P450 monooxygenase	**-2.74**	**0.0005**	0.0000
SAV413	*pteC*	cytochrome P450 monooxygenase	**-2.33**	**0.0004**	0.0000
SAV414	*pteB*	dehydrogenase	**-2.34**	**0.0000**	0.0000
SAV415	*pteA5*	modular polyketide synthase	**-1.01**	0.1136	**0.0054**
SAV416	*pteA4*	modular polyketide synthase	**-1.28**	**0.0095**	0.0002
SAV417	*pteA3*	modular polyketide synthase	-0.64	0.3514	**0.0457**
SAV418	*pteA2*	modular polyketide synthase	-0.87	0.3639	0.0506
SAV419	*pteA1*	modular polyketide synthase	-0.80	0.4388	0.0752
Non-ribosomal peptide-6 (*nrp6*)					
SAV600	*fecC1*	ABC transporter iron(III)/siderophore transport system ATP-binding protein	**2.40**	**0.0003**	0.0000
SAV601	*fecD1*	ABC transporter iron(III)/siderophore permease	0.75	0.5625	0.1290
SAV602	*fecB*	ABC transporter iron(III)/siderophore-binding protein	**2.95**	**0.0006**	0.0000
SAV603	*nrps6*	non-ribosomal peptide synthetase	0.68	0.3119	**0.0342**
SAV604		hypothetical protein	**1.18**	**0.0224**	0.0005
SAV605	*fadD2*	acyl-CoA synthetase	**1.20**	**0.0049**	0.0001
SAV606		hypothetical protein	0.71	0.3525	**0.0464**
SAV607		taurine catabolism dioxygenase	**1.15**	**0.0017**	0.0000

SAV608	*fabC2*	acyl carrier protein	**1.02**	0.1136	**0.0054**
SAV609	*fabH4*	3-oxoacyl-ACP synthase III	**1.15**	0.1182	**0.0058**
SAV610		MFS transporter protein	**1.19**	**0.0166**	0.0004
SAV611		beta-hydroxylase	**1.64**	**0.0005**	0.0000
Avermectin cluster (*ave*)					
SAV935	*aveR*	LuxR-family transcriptional regulator	**2.00**	**0.0049**	0.0001
SAV936	*aveF*	C-5 ketoreductase	0.59	0.6016	0.1518
SAV937	*aveD*	C5-O-methyltransferase	0.43	0.6148	0.1603
SAV938	*aveA1*	type I polyketide synthase	**1.18**	0.1820	**0.0122**
SAV939	*aveA2*	type I polyketide synthase	0.39	0.7264	0.2537
SAV940	*aveC*		0.00	0.9993	0.9950
SAV941	*aveE*	cytochrome P450 monooxygenase	**1.04**	0.1376	**0.0074**
SAV942	*aveA3*	type I polyketide synthase	0.49	0.6328	0.1725
SAV943	*aveA4*	type I polyketide synthase	0.73	0.4435	0.0764
SAV944[3]	*orf-1*	reductase	0.14	0.9330	0.6990
SAV945	*aveBI*	dTDP-L-oleandrose transferase (glycosyltransferase)	0.70	0.6052	0.1543
SAV946	*aveBII*	dTDP-glucose 4.6-dehydratase	0.23	0.8027	0.3477
SAV947	*aveBIII*	glucose-1-phosphate thymidyltransferase	**1.08**	0.2027	**0.0154**
SAV948	*aveBIV*	dTDP-4-keto-6-deoxy-L-hexose 4-reductase	-0.28	0.8828	0.5032
SAV949	*aveBV*	dTDP-4-keto-6-deoxyhexose 3.5-epimerase	0.68	0.5387	0.1160
SAV950	*aveBVI*	dTDP-4-keto-6-deoxy-L-hexose2.3-dehydratase	0.84	0.4308	0.0711
SAV951	*aveBVII*	dTDP-6-deoxy-L-hexose 3-O-methyltransferase	**2.01**	**0.0020**	0.0000
SAV952	*aveBVIII*	dTDP-4-keto-6-deoxy-L-hexose 2.3-reductase	**1.83**	**0.0049**	0.0001
SAV953	*aveG*	thioesterase	**2.20**	**0.0018**	0.0000
Melanin cluster (*melC-1*)[4]					
SAV1136	*melC1*	tyrosinase co-factor protein	**-1.68**	0.0776	**0.0015**
SAV1137	*melC2*	tyrosinase	**-1.85**	**0.0078**	0.0000
γ-butyrolactone cluster (*gbl*)					

---

[3] Not involved in avermectin biosynthesis
[4] Values from stationary phase (t2) analysis

SAV2266	*avaC*	phosphatase	-0.98	0.0794	**0.0030**
SAV2267	*avaB*	oxidoreductase	**-1.20**	0.2777	**0.0279**
SAV2268	*avaL2*	TetR-family transcriptional regulator	**-3.15**	**0.0078**	0.0001
SAV2269	*avaA*	gamma-butyrolactone biosynthesis protein	**-1.59**	0.1456	**0.0083**
SAV2270	*avaL1*	TetR-family transcriptional regulator	**-2.68**	**0.0035**	0.0000
Polyketide-3 cluster (*pk3*)					
SAV2272		hypothetical protein	**-2.45**	0.2000	**0.0150**
SAV2273		isomerase	**-7.24**	**0.0208**	0.0005
SAV2274		secreted protein	**-5.02**	**0.0473**	0.0014
SAV2275		transmembrane efflux protein	**-4.32**	0.1124	**0.0053**
SAV2276		3-oxoacyl-ACP synthase III	**-4.24**	0.1242	**0.0063**
SAV2277		thioesterase	**-3.62**	0.2222	**0.0183**
SAV2278		F420-dependent dehydrogenase	**-3.52**	0.3037	**0.0327**
SAV2279		acyl-CoA synthetase	**-2.97**	0.3522	**0.0462**
SAV2280	*pks3-1*	modular polyketide synthase	**-1.20**	0.6288	0.1702
SAV2281	*pks3-2*	modular polyketide synthase	**-1.90**	0.5893	0.1436
SAV2282	*pks3-3*	acyl carrier protein	**-2.03**	0.3974	0.0593
Oligomycin cluster (*olm*)					
SAV2890	*ccrA1*	crotonyl-CoA reductase	-0.42	0.4368	0.0743
SAV2891		hypothetical protein	**-1.18**	0.2280	**0.0193**
SAV2892	*olmA4*	modular polyketide synthase	-0.33	0.4760	0.0913
SAV2893	*olmA5*	modular polyketide synthase	-0.22	0.6421	0.1789
SAV2894	*olmB*	cytochrome P450 monooxygenase	**-1.09**	0.1512	**0.0087**
SAV2895	*olmA7*	modular polyketide synthase	-0.64	0.0746	**0.0028**
SAV2896	*olmA6*	modular polyketide synthase	-0.74	0.1443	**0.0080**
SAV2897	*olmA3*	modular polyketide synthase	-0.32	0.3610	**0.0493**
SAV2898	*olmA2*	modular polyketide synthase	-0.31	0.5785	0.1377
SAV2899	*olmA1*	modular polyketide synthase	-0.61	0.1817	**0.0121**
SAV2900		P450-like protein	-0.57	0.2474	**0.0224**
SAV2901	*olmRII*	LuxR-family transcriptional regulator	**-1.47**	0.0712	**0.0026**

SAV2902	*olmRI*	LuxR-family transcriptional regulator	**-1.56**	**0.0006**	0.0000
SAV2903	*olmC*	Thioesterase	**-1.86**	0.1235	**0.0062**
		Neopentalenolactone cluster (*ptl*)			
SAV2989		MarR-family transcriptional regulator	**-1.06**	**0.0487**	0.0015
SAV2990	*gap1*	glyceraldehyde-3-phosphate dehydrogenase	**-1.36**	0.2469	**0.0222**
SAV2991	*ptlH*	1-deoxypentalenic acid 11-beta hydroxylase	-0.93	0.1529	**0.0089**
SAV2992	*ptlG*	transmembrane efflux protein	-0.37	0.9068	0.5988
SAV2993	*ptlF*	1-deoxy-11beta-hydroxypentalenic acid dehydrogenase	-0.76	0.2639	**0.0251**
SAV2994	*ptlE*	Baeyer-Villiger monooxygenase	**-1.30**	**0.0087**	0.0001
SAV2995	*ptlD*	dioxygenase	**-1.21**	0.0962	**0.0041**
SAV2996	*ptlC*	hypothetical protein	**-1.07**	0.5367	0.1153
SAV2997	*ptlB*	farnesyl diphosphate synthase	-0.96	0.5690	0.1322
SAV2998	*ptlA*	pentalenene synthase	**-1.34**	0.5592	0.1269
SAV2999	*ptlI*	pentalenene C13 hydroxylase; cytochrome P450	**-1.24**	0.4943	0.0993
SAV3000	*ptlR*	AraC-family transcriptional regulator	-0.58	0.5379	0.1158
SAV3001	*ptlJ*	lyase	-0.31	0.6109	0.1579
SAV3002	*ptlL*	hypothetical protein	-0.03	0.9754	0.8867
		Albaflavenol/albaflavenone cluster (*ezs*)			
SAV3031	*cyp14*	epi-isozizaene hydroxylase (cytochrome P450 monooxygenase)	**1.00**	0.2918	**0.0301**
SAV3032	*ezs*	epi-isozizaene synthase (sesquiterpene cyclase)	**1.10**	0.1621	**0.0097**
		Non-ribosomal peptide-3 cluster (*nrp3*)			
SAV3155		MbtH-like protein	**-1.18**	0.2696	**0.0261**
SAV3156	*nrps3-1*	non-ribosomal peptide synthetase	**-1.56**	0.0867	**0.0035**
SAV3157		export protein	-0.66	0.6407	0.1773
SAV3158	*nrps3-2*	non-ribosomal peptide synthetase	-0.84	0.6149	0.1623
SAV3159	*nrps3-3*	non-ribosomal peptide synthetase	-0.93	0.5290	0.1123
SAV3160		aminotransferase	**-1.13**	0.4384	0.0749
SAV3161	*dapF2*	diaminopimelate epimerase	**-1.03**	0.5045	**0.1029**
SAV3162		hypothetical protein	**-1.52**	0.1668	**0.0101**

SAV3163		hypothetical protein	**-1.24**	0.3227	**0.0384**
SAV3164		hypothetical protein	**-1.59**	**0.0003**	0.0000
Polyketide-4 cluster (*pk4*)					
SAV7184	*pks4*	modular polyketide synthase	**-3.22**	0.2980	**0.0315**
SAV7185		UDP-glucose:sterol glucosyltransferase	**-3.05**	0.2753	**0.0271**
SAV7186	*cyp26*	cytochrome P450 hydroxylase	**-2.88**	0.2419	**0.0213**
Polyhydroxycarboxylate siderophore cluster (*avs*)					
SAV7320	*avsA*	siderophore synthetase component	**1.11**	**0.0295**	0.0007
SAV7321	*avsB*	siderophore synthetase component	0.77	0.2474	**0.0223**
SAV7322	*avsC*	siderophore synthetase component	0.63	0.4654	0.0862
SAV7323	*avsD*	diaminopimelate decarboxylase	0.25	0.8125	0.3606