Lineage-specific genes are clustered with allorecognition loci and respond to G × E factors regulating the switch from asexual to sexual reproduction in *Neurospora*

Zheng Wang[1], Yaning Wang[1,2], Takao Kasuga[3], Yen-Wen Wang[1], Francesc Lopez-Giraldez[4], Yang Zhang[5], Zhang Zhang[5], Caihong Dong[2], Anita Sil[6], Frances Trail[7], Oded Yarden[8], and Jeffrey P. Townsend[1, 9*]

[1]*Department of Biostatistics, Yale School of Public Health, New Haven, Connecticut 06511, USA*

[2]*Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*

[3]*College of Biological Sciences, University of California, Davis, CA95616, USA*

[4]*Yale Center for Genomic Analysis, New Haven, Connecticut 06511, USA*

[5]*National Genomics Data Center, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China*

[6]*Department of Microbiology and Immunology, University of California, San Francisco, CA94115, USA*

[7]*Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, MI 48824, USA*

[8]*Department of Plant Pathology and Microbiology, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, Israel*

[9]*Department of Ecology and Evolutionary Biology, Program in Microbiology, and Program in Computational Biology and Bioinformatics, Yale University, New Haven, Connecticut 06511, USA*

* Correspondence: jeffrey.townsend@yale.edu;

Abstract:

Lineage-specific genes (LSGs) have long been postulated to play roles in the establishment of

genetic barriers to intercrossing and speciation. However, there is a lack of working hypotheses

as to how they might play that role. In the genome of *Neurospora crassa*, most of the 670

Neurospora LSGs that are aggregated adjacent to the telomeres are clustered with 61% of the

HET-domain genes, which regulate self-recognition and define vegetative incompatibility

groups. Among the 342 LSGs that are dynamically expressed during both asexual and sexual

phases, 64% were detectable on unusual carbon sources such as furfural and HMF—wildfire-

produced chemicals that are a strong inducer of sexual development. Expression of a significant

portion of the LSGs was sensitive to light and temperature, factors that regulate the switch from

asexual to sexual reproduction. Furthermore, expression of the LSGs was significantly affected

in the knockouts of *adv-1* and *pp-1* that regulate hyphal communication, and expression of more

than one quarter of the LSGs was affected by perturbation of the mating locus. Accordingly, we

propose a gene-by-environment interaction model encouraging further investigation of the roles

of LSGs and HET-domain genes in speciation in *Neurospora*. This gene-by-environment

interaction model emphasizes the roles of the LSGs in response to genetic and environmental

factors, leading to the regulation of the switch from the asexual growth and fusion, such that

vegetative incompatibility governed by allorecognition promotes allelic homogeneity, sexual

reproduction, and outbreeding, whereas VI repression and meiotic recombination promotes

allelic polymorphism.

Introduction

Since the emergence of life, molecular evolution has contributed to the accumulation of

novel and diverse features in all kinds of organisms. Two fundamental components of that

molecular evolutionary novelty are new genes and novel gene functions, which have long been

considered to be emergent properties of gene duplication and rearrangement. Nevertheless,

genomes often harbor numerous orphan genes or lineage specific genes (LSGs)—novel genes

that have no homologues in distantly- or closely-related lineages and that cannot be tracked to

ancestral lineages. These LSGs, including de novo genes that evolve from previous non-coding

DNA and non-genic elements [1], manifest in large numbers across a diversity of organisms,

such that they represent nearly one-third of the genes in all genomes, including phages, archaea,

bacteria, and eukaryotic organisms [2]. Three key challenges have thus been the focus of studies

of LSGs: how to identify LSGs, how to track their evolutionary histories, and perhaps most

importantly, how these genes are integrated into pre-existing gene interaction networks [3].

Accurate identification of LSGs can, at times, be difficult. They originate via three

evolutionary processes: (1) rapid gene evolution, which is refractory to tracking homology on the

basis of sequence conservation; (2) intragenomic gene loss and gain or horizontal gene transfer,

which can convey higher fitness in response to genetic/environmental changes; and

(3) accumulated mutations that establish novel function, evolving slowly but long enough on an

independent lineage that the gene phylogeny cannot be tracked back to its distant ancestor or to

related lineages [4–6].

Studies of the genomic characteristics of LSGs in several model organisms have revealed

the likely origins of LSGs from gene duplication, non-coding sequences, as well as fast-

evolution at conserved genomic positions. One example supporting the frequency of origin by

rapid divergence after gene duplication and rearrangement can be found in yeast, where the

presence of 55% to 73% percent of the LSGs can be explained by sufficient divergence from

sister species [6]. Some lineage-specific protein-coding genes might have directly evolved from

non-coding regions in the genome [7], as has been described in the tests of fruit flies [8,9]. One

hundred seventy-five *de novo* genes in Asian rice corresponded to recognizable non-genic

sequences in closely related species [10]. These investigations using model organisms confirmed

unique characteristics of these *de novo* genes, making good frameworks for investigating LSGs

in other species. However, linkages between these revealed genomic characteristics and the

integrative functions of LSGs remain unclear. Therefore, systematic approaches that combine

study of comparative genomics with functional assays of gene expression and gene-perturbation

phenotypes using well-established model systems are critical to integrate the investigation of the

LSGs' origination and function.

The most frequently used approach to identify LSGs is using phylostratigraphy [11,12].

Precise identification of the origins of *de novo* genes using a phylostratigraphic approach is

critically dependent on accurate gene annotation and extensive comparison among proper

representative genomes [13]. It is difficult to distinguish whether genes with no homologues in

closely related lineages are true LSGs as opposed to lacking homologues in closely related

lineages that have few genomes available. An alternative to phylostratigraphy is gene synteny,

which compares each gene's position relative to its neighbors. A recent study suggested that if

the neighbors of a gene are in a conserved order in other species, then the gene is likely to

correspond to whatever is at the orthologous position in the other species as well—even if the

sequences do not match [14].

With detailed pangenomic analyses, identification of LSGs, including *de novo* protein-

coding genes, need to be further verified with a systematic approach focusing on possible

functional novelty and genetic signals that may be associated with such a novelty [3]. Systematic

assessment of the putative LSG function can track their behaviors during the growth and

development and verify their possible roles by examination of corresponding knockdown or

knockout phenotypes [3]. LSGs are naturally thought to be important to species- or genus-level

adaptations of development to taxon-specific ecology.

The well-annotated model species in the genus *Neurospora*, *N. crassa, N. tetrasperma* and

N. discreta of the class Sordariomycetes, provide a set of three closely-related genomes enabling

investigation of possible genetic novelties associated with recent and rapid ecological

divergences [15], such as responses to nutrients and other environmental factors and

developmental divergences such as heterothallic and pseudohomothallic outcrossing during

sexual reproduction as well as incompatibility during vegetative growth. *Neurospora* species are

highly adapted to the postfire environment, capable of fast asexual growth and reproduction on

simple nutrients and have long been genetic models for eukaryotic metabolic regulation and for

mating, meiosis and morphological development during reproduction [16–18]. In fact, fungi in

the Sordariomycetes exhibit remarkable ecological, biological, and morphological diversity [19–

21]. These fungi also exhibit diverse reproductive modes, including heterothallism,

pseudohomothallism, and homothallism with or without asexual reproduction [18,22–26]. Such

diversities in ecology and development that have evolved in parallel or convergently within

closely related species in a single fungal class, may provide avenues toward an understanding of

the $G \times E$ (gene-environment interaction) impacts of lineage-specific elements on the

evolutionary process. Due to the repeat-induced point mutation (RIP) genome-defense system,

Neurospora is known to lack recent gene duplications—a major source of evolutionary novelty

[27,28]. Nevertheless, comparing representative genomes in prokaryotes, plants & animals,

Basidiomycota, major lineages of Ascomycota, and *Chaetomium globosum,* which is closely

related to *N. crassa*, 2219 orphan genes were identified in *N. crassa* by phylostratigraphy [29]. In

the past decade, many more fungal genomes have been sequenced. Their sequence, along with

the advances in genome sequencing and annotation techniques [30–32], provide a more inclusive

comparison for identifying lineage specific genes. Therefore, to understand how important the

roles that LSGs play in genome-wide regulation during the whole life history, we investigated

signals of recent selection pressures in LSGs, focusing on genes involved in environmental

responses and fertility or mating in *Neurospora*.

We also observed that several LSGs are allorecognition loci, often referred to as *het*

(heterokaryon incompatibility) or *vic* (vegetative incompatibility) genes. As their names indicate,

het or *vic* genes regulate allorecognition during vegetative growth, and only individuals with

compatibility at all of their *het*-loci can fuse and simply expand their colonies [33]. Some HET-

domain genes have pleiotropic effects in sexual development in some fungal species and play

direct roles in reproductive isolation and speciation within sympatry [34]. HET-domain genes

can be predicted with conserved HET domain in protein sequences, and there were 69 HET-

domain genes reported (of which 68 were mapped to the original genome-sequenced strain of *N.*

crassa) [33]. Chromosomal locations that are itinerant over evolutionary time, high duplication

rates and high frequencies of gain and loss are shared traits of HET-domain genes and LSGs,

suggesting the merits of integrative investigations of both gene groups.

Results

Summary—In this study, we identified and verified 670 lineage-specific genes (LSGs) in

Neurospora crassa via BLAST search against inclusive representative genomes using previously

published phylostratigraphy data that was based on a limited set of representative genomes.

Using two clustering approaches, we discovered that over 60% of the 670 LSGs formed clusters

in the telomere regions and clustered with the HET-domain genes. However, most of the LSGs

are not functionally annotated (e.g., with gene ontology terms). Therefore, to assess the possible

functional roles of the LSGs, we analyzed genome-wide gene expression data on *N. crassa*

growing on distinct media at different stages of life history and under different light and

temperature conditions. We observed that nearly half of the LSGs were actively expressed during

asexual and sexual growth. A substantial number (291) LSGs were induced by the presence of

furfural. 158 LSGs were exclusively expressed in furfural cultures, in contrast to only 17 LSGs

that were exclusively expressed in cultures on media supplied with common simple

carbohydrates. We also reported a significant portion of the LSGs being turned on or off by the

changes of light exposure and temperature, conditions that are critical for *N. crassa* asexual and

sexual reproduction. We further examined expression of the LSGs and HET-domain genes in

knockouts of two transcription factors, *adv-1* and *pp-1*. Both transcription factors play multiple

roles in asexual and sexual development in *N. crassa*. In addition, we sequenced and analyzed

genome-wide gene expression in a loss-function mutant at the mating locus *mat 1-2-1* in a *mat a*

strain during the crossing. The LSGs were more likely to be affected by gene-manipulation than

other genes, compared with other non-LSG genes and HET-domain genes. They were more

likely to be turned on or turned off completely, rather than being turned up or down slightly.

Finally, we examined asexual and sexual growth phenotypes for 367 available KO strains of the

Neurospora LSGs. We identified two LSGs with abortive sexual reproduction, and several LSGs

with minor phenotypes in response to high temperature or mycelium morphology.

670 LSGs were identified in Neurospora genomes. We defined *Neurospora* lineage-specific

genes (LSGs) as *N. crassa* genes exhibiting homology to only to genes found in sequences from

species within the genus *Neurospora [11]*. To identify *Neurospora* LSGs, phylostratigraphy was

performed on representative taxa for major fungal lineages and several non-fungal reference

genomes, and 1872 *N. crassa* genes were identified as putative LSGs (**Fig. 1, Table S1**). Within

these 1872 *N. crassa* genes, a total of 695 genes are shared between the genomes of *Neurospora*

and the sole species *S. macrospora* in the sister genus (**Table S1**). Further reciprocal-BLAST

searches were made for the 1872 genes against available Sordariomycetes genomes, including

genomes of *Neurospora* closely related species in *Podospora*, *Pyricularia* and *Ophiostoma* as

well as species within the genus including *N. tetrasperma* and *N. discreta* at the National Center

for Biotechnology (NCBI) and FungiDB genome database. We identified 670 genes that are

N. crassa lineage-specific genes (LSGs) (**Fig. S1**, **Table S1**). There are 7400 single-copy

orthologs shared among the genomes of *N. crassa*, *N. discreta,* and *N. tetrasperma* (**Fig. S1A**).

Among the 670 LSGs identified in the *N. crassa* genome, 241 are unique to *N. crassa* and 405

are shared between *N. crassa* and *N. tetrasperma* with 248 showing no orthologs in *N. discreta*.

181 *Neurospora* LSGs are shared between *N. crassa* and *N. discreta* with 26 showing no

orthologs in *N. tetrasperma* (**Fig. S1B**). Because of the special status of *N. crassa* as a model

species, here we specifically investigate and report as LSGs those that are specific to *N. crassa* in

this study. With more genomes in this fungal class being sequenced and annotated and more non-

classified genes in the phylostratigraphy being analyzed, these numbers are expected to be

slightly changed. Of the 670 *Neurospora* LSGs, 515 have at least one intron (average: ~2 introns,

maximum: 8 introns in NCU07480). These LSGs encoded proteins ranging from 26 to 1310

amino acids (NCU05561 and NCU04852, respectively). The average LSG length (~192 amino

acids) was significantly shorter than the average length of the non- *Neurospora* lineage-specific

genes (non-LSG genes, ~528 amino acids).

Figure 1. Systematics of *Neurospora* lineage-specific genes (LSGs) and potential roles in fungal growth and development. (**A**) Genomic phylostratigraphy of lineage-specificity classifications of predicted protein- coding genes (enumerated at ancestral nodes: 3279 **E**ukaryote-core, 1035 **D**ikarya-core, 113 **A**scomycota-core, 2557 **P**ezizomycotina-specific, 157 **S**ordariomycetes-specific, and 1871 **N**eurospora- specific) that are present within the *N. crassa* genome; (**B**) Life history of *N. crassa*. These developmental processes have been transcriptionally profiled [dashed red arrows; 35].

Figure S1. Genome-wide genes and *Neurospora* LSGs compared within the three *Neurospora* species *N. crassa*, *N. tetrasperma*, and *N. discreta*. (**A**) Comparative genomic protein-coding gene content among *N. crassa*, *N. discreta* and *N. tetrasperma*, centering shared single-copy orthologs within the three species. (**B**) Some *Neurospora* LSGs in the *N. crassa* genome are shared within *N. tetrasperma* and *N. discreta* genomes.

LSGs are aggregated in the telomere regions and clustered with the HET-domain genes.

Neurospora LSGs are distributed in all seven chromosomes of the genome, with paralogs from

duplicates often clustered together (**Fig. 2**). Window-free maximum-likelihood model averaging

of the gene-regionalized probability of *Neurospora* LSGs and HET-domain genes revealed that

LSGs were clustered, with significant ($P < 0.05$) clustering on chromosomes I, II, III, IV, and V.

Large LSGs clusters are typically aggregated toward the telomeres of each chromosome and

frequently contain large non-coding spaces, especially in chromosomes I, III, IV, V, VI, and VII

(**Fig. 2 A–G, Table S2**). Detailed clustering revealed with the Cluster `Locator` [36] tallied 67%

of LSGs as present in clusters with a `max-gap` of five (48% with a `max-gap` of one, i.e.,

separated by one gene). About 30% of LSGs were in clusters with more than four genes

(**Tables 1 & S3**), including six clusters hosting 9, 10, 16, and 23 genes.

A total of 68 *Neurospora* genes with HET domains (HET-domain genes) were identified

and mapped to the *N. crassa* FGSC2489 strain [Figure 1 in reference 33]. Many of these HET-

domain genes exhibited nonrandom distributions and were clustered near the end of the linkage

groups, largely overlapping with clusters of LSGs (**Fig. 2**). A previous study demonstrated that

another HET-domain gene, NCU03125 (*het-C*), plays a role in vegetative incompatibility [37].

42 of the 69 HET-domain genes were clustered with at least one LSG within a range of five

genes (`max-gap = 5`), and 23 and 14 HET-domain genes were clustered with at least one LSG

within a range of three (`max-gap = 1`) or two genes (`max-gap = 0`) separately (**Fig. 2A–G,**

Tables 1 and **S4**). In fact, many of HET-domain genes were only syntenic within *Neurospora*

and very closely related species in the Sordariales, and most HET-domain genes were

surrounded by *Neurospora* LSGs and comparatively "young" genes (**Fig. 2H, Table S5**).

Table 1. LSGs and HET-domain genes are present in clusters in the *N. crassa* genome.

Cluster size	Lineage-specific genes (LSGs)[a]			LSG and HET-domain genes[b]		
	# Clusters	# Genes	% Genes	# Clusters	# Genes	% Genes
2	57	114	17.1%	63 (15 *het*)	126	17.1%
3	28	84	12.6%	28 (7 *het*)	84	11.4%
4	8	32	4.8%	12 (6 *het*)	48	6.5%
5	14	70	10.5%	13 (3 *het*)	65	8.8%
6	5	30	4.5%	11 (7 *het*)	66	8.9%
7	3	21	3.2%	2	14	1.9%
8	2	16	2.4%	2	16	2.2%
9	3	27	4.1%	1	9	1.2%
10–11	1	11	1.7%	2(2 *het*)	21	2.9%
16–18	1	16	2.4%	2(2 *het*)	35	4.8%
23	1	23	3.5%	1	23	3.1%
Total	123	444	66.8%	129 (42 *het*)	474 (42 *het*)	64.4%

211 [a] 670 genes; [b] 736 genes (3 *HET-domain* genes are LSGs).

Figure 2. Identification of LSG and LSG-Het clusters in the *N. crassa* genome. Regionalized probability of the lineage-specific status of a gene inferred using MACML, a window-free maximum-likelihood model-averaging approach, heat maps of clustering, and dot-plot of LSGs distributed across (**A**) chromosome I, (**B**) chromosome II, (**C**) chromosome III, (**D**) chromosome IV, (**E**) chromosome V, (**F**) chromosome VI, and (**G**) chromosome VII. Model-averaged profiles (blue: low clustering of LSGs vs non-LSGs; red: high clustering of LSGs vs non-LSGs; gray shading: 95% model uncertainty interval) quantify the window-free regionalized probability that a gene is an LSG. The heat maps quantify LSG-density across chromosomal windows of 10,000 base pairs. Dots located in accompanying chromosomal heat maps correspond to LSGs (clustered: green [$P < 0.01$], and non-clustered [$P \geq 0.01$]: orange) and clustered HET-domain genes (purple triangles). (**H**) Lineage-specificity at four taxonomic depths (greener color intensity corresponds to lineage-specificity at successively lesser taxonomic depths: lineage specificity in *Neurospora* > Sordariales > Sordariomycetes > Ascomycetes; thin orange circle margin indicates synteny within *Neurospora*) of five neighbor genes on either side of 69 HET-domain genes (**Table S5**) reported in the original genome-sequenced *N. crassa* strain [[FGSC2489; 38].

Many LSGs and HET-domain genes are dynamically expressed in response to developmental

and environmental changes. Genome-wide gene expression was measured in key stages of the

N. crassa life cycle. LSGs and non-LSG genes generally exhibited substantial differences in the

numbers of genes that were effectively expressed and silenced (**Figs. 1**, **S2, S3**; **Tables 2, S6**).

During sexual development on Synthetic Crossing Medium (SCM) and asexual growth on Bird

Medium (BM), similar trends were observed for proportions of LSGs and non-LSG genes that

exhibited measurable expression, but an increased proportion of LSGs were expressed during the

early hyphal branching on Maple Sap Medium (MSM; **Fig. S2**). At one or more time points

during perithecial development, 238 genes exhibited at least 5-fold ($P < 0.05$) expression

changes during perithecial development, indicating potential roles in the regulation of sexual

reproduction. 35 genes exhibited no measurable expression in any sampled life history stages,

suggesting either function only in unusual circumstances or mis-annotation as expressed genes

(**Table S6**). During conidial germination and early asexual growth, respectively, 95 and 137

LSGs exhibited at least 5-fold ($P < 0.05$) expression changes for cultures on BM and MSM, and

213 and 181LSGs were expressed during none or only one of the four sampled stages for

cultures on BM or MS separately. Expression of 342 LSGs was detected in at least two sampled

stages during sexual reproduction and during asexual growth on asexual specific BM and MSM

(**Fig. 3**).

Figure S2. Proportion of *Neurospora* LSGs (blue) and non-LSG genes (orange) that were expressed at sampling points from sexual development on SCM and conidial germination and growth on BM and MSM. (**A**) Sexual development from protoperithecia (starting stage) to mature perithecia at 144 h [22]. (**B**) Asexual growth from conidial germination to the first hyphal branching on Bird medium supporting only asexual development. (**C**) Asexual growth from conidial germination to the first hyphal branching on maple sap medium supporting both asexual and sexual reproduction [conidial germination; 23].

Figure S3. Gene expression dynamics of LSGs across sexual development from protoperithecia (starting stage) to mature perithecia at 144 h [22] and asexual growth from conidial germination to the first hyphal branching, on Bird medium supporting only asexual development, and on a maple sap medium supporting both asexual and sexual reproduction [conidial germination; 23]. Heatmap was generated using the ClustVis web tool.

Table 2. Transcriptomics analysis results on *N. crassa* responses to environmental factors.

Experiments[1]	LSGs (670)		Het (66[2])		other		Notes[3] LSG vs other (*P*)
	expressed	silenced	expressed	silenced	expressed	silenced	
25C BM	457	213 (82)	66	0	8668	326 (122)	*P*<0.01
25C MSM	489	181 (52)	66	0	8698	296 (113)	*P*<0.01
25C SCM	484	186 (49)	66	0	8764	230 (77)	*P*<0.01
37C BM	350	342 (84)	55	2 (1)	8616	710 (260)	*P*<0.01
Crop residue[4]	149	521	52	14	7497	1497	*P*<0.01
Furfural[4]	291 (31)	366 (32)	50 (2)	11 (4)	7763 (147)	968 (63)	*P*<0.01; *P*<0.01 (*P*<0.01)
No carb[4]	184	486	52	14	7643	1351	*P*<0.01
Light[5]	224 (97)	381 (6)	25 (10)	31 (1)	7366 (381)	1701 (42)	*P*<0.01; *P*<0.01 (*P*<0.13)
Dark[5]	133 (6)	472 (97)	16 (1)	40 (10)	7027 (42)	2040 (381)	*P*<0.01; *P*<0.13 (*P*<0.01)

[1] In this multi-point gene expression data, only genes that were either expressed in at least two of the sampled points or not expressed at all in all sampled points were reported and considered as significantly active or silenced across the sampled process. For cultures sampled at multiple timepoints, some genes were expressed or silenced only at one of the sampled points (tallied in parentheses).

[2] Three predicted HET-domain genes—NCU03378 (*tol*), 07596, and 10839—are also identified as lineage-specific genes (LSGs).

[3] Chi-square test *P* values regarding the proportion are provided for comparison between lineage-specific genes and non-HET non-lineage-specific genes that are color-coded in the rows.

[4] The experiment compared *N. crassa* growing on media supplied with crop residue or furfural as carbon resources reporting expression of 9449 genes. Numbers of genes being expressed or silenced for cultures on the crop residue media, furfural media, and zero carbon medium, were reported. (Numbers in parentheses are the subset genes that are exclusively expressed and exclusively silenced in cultures on furfural media).

[5] The light-exposure experiment tested dark and four light-exposure timepoints: 15-, 60-, 120-, and 240-minutes reporting expression of 9728 genes. The number of genes expressed or silenced in the light or in the dark are reported (numbers in parentheses are the subset genes that are exclusively expressed or exclusively silenced).

Figure 3. Differential expression of LSGs in *N. crassa* growth during three developmental processes [22,23] and on several media each supplying distinct carbon resources [39,40]. 342 LSGs were expressed at measurable levels in at least two stages in each of the three developmental processes (**Tables S6–S7**), including eight stages of sexual development on SCM (salmon pink, totaling 484 LSGs with measurable expression), four stages of asexual growth on BM (beige, totaling 457 LSGs with measurable expression) and four stages of asexual-sexual growth on MSM (light blue, totaling 489 LSGs with measurable expression). Among the 342 LSGs (arrow-linked), 231 were detectably expressed when cultured on either 2-furaldehyde furfural and/or 5-hydroxymethyl furfural (HMF), substrates that promote sexual development (purple, total 291 LSGs); and 128 were detectably expressed on sucrose and/or residues of at least one of five common crop straws (barley, corn, rice, soybean, and wheat; **Tables S6–S7**), substrates that support asexual growth and sporulation (light green, totaling 150 LSGs).

Genomic gene expression was also assayed for *N. crassa* growing on seven different

carbon conditions including, absence of carbon, solo glucose carbon source, and a complex

carbon source with five commonly available crop residues, including barley, corn, rice, soybean,

and wheat straws, found in the field [39]. Expression of 464 LSGs was too low to be detected

under any of these conditions; this large portion of *Neurospora* LSGs is not required for

vegetative growth associated with carbon metabolism regulation (**Tables S7, S8**). Expression of

22 LSGs required that at least one type of carbon resource was present in the media, while

expression of 56 other LSGs was only detected in the absence of carbon. Revisiting expression

data collected from experiments investigating *N. crassa*'s tolerance to furfural [40] identified

that 245, 239, 257, and 232 LSGs exhibited measurable levels of expression in the simple carbon

cultures, DMSO (the carbon blank control), furfural, and HMF treatments (**Tables S7–S8**).

Within the 291 genes expressed in either furfural or HMF cultures or both, 61 were not

expressed in the wild-type condition. Furfural is derived from lignocellulosic biomass and

enriched in a post-fire environment. *N. crassa* sexual spore germination can be induced by

furfural presence [41,42]. Furfural also inhibits conidia germination. Compared with cultures

under wild-type conditions, 12 LSGs exhibited a 3-fold or higher expression in response to

furfural, with NCU09604, 07323, and 01153 showing 6- to 22-fold up-regulation in furfural

cultures.

Environmental factors, including light and temperature that regulate fungal growth and

development, also dramatically affect expression of LSGs (Table S8 for more details). *N. crassa*

genome is equipped with a set of light sensors responding to different light spectrums, duration

and intensities [43–48], and *Neurospora* LSGs exhibit sensitive responses to light conditions.

When *N. crassa* cultures were exposed to light up to 4 h, genes were classified as short light

responsive genes or long light responsive genes based on their expression profile changes [46].

Revisiting the previous data disclosed that out of 488 genes induced by light stimulus, 106 were

LSGs (significantly enriched, $P < 0.01$), 59 of which were in the predicted clusters, including all

genes in three 2-LSGs clusters, including cluster #25, 61, and 81 as well as three genes in a 4-

gene cluster #127. Among 49 genes whose expression halted upon exposure to light, six were

LSGs, including two genes NCU05052 and 05058 in a 3-gene cluster (**Fig. 4A, Table S9**).

During conidial germination at a high temperature of 37 C on BM, expression of 270 LSGs was

completely inhibited, a substantial enrichment ($P < 0.01$; a total of 941 out of 10592 genes

exhibited no measurable expression). There were 148 LSGs exhibiting no measurable expression

at 25 C and 37 C. There were 152 LSGs exhibiting detectable expression in cultures at 25 C but

being turned off at 37 C, 100 of which were clustered LSGs, including 24 clusters with more

than 2 genes that were turned off at 37 C. In contrast, 13 genes exhibited detectable expression in

cultures at 37 C, but not at 25 C (**Fig. 4B, Table S9**). The LSG cluster #25 of NCU02144–02145

was the only LSG cluster that was silent in dark conditions or at 37 C.

Figure 4. Genome-wide expression responses of LSGs, HET-domain, and other genes to shifting of culture
environment. (**A**) Stacked proportions of 670 LSGs, 68 HET-domain, and other genes in the *N. crassa* genome,
exhibited expression only in the dark (gray) or only in response to 15 to 240 minutes of light exposure (yellow).
(**B**) Stacked proportions of 670 LSGs genes, 68 HET-domain genes, and other non-LSG genes in the *N. crassa*
genome that are expressed across 4 stages of conidial germination and vegetative growth only at 25 C (blue) or
37 C (red) on Bird medium.

Expression of HET-domain genes as a whole exhibited no clear patterns in response to

environmental conditions or developmental stages (**Fig. 4A–B, Tables S8–9**). However, 11

HET-domain genes were not expressed in cultures on furfural or HMF, and 28 HET-domain

genes were expressed neither in the dark nor during a shift from the dark to light for a duration of

up to 2 h. Both of these conditions are known to promote sexual reproduction in *N. crassa*. More

genes () were significantly up-regulated (5968 vs. 2935; *P* < 0.05) during the first branching of

the germ tube on MSM, which supports both asexual and sexual development, than those on BM,

which is specifically designed for promoting asexual reproduction and inhibiting sexual

development. Accordingly, many more HET-domain genes were significantly up-regulated (46

vs. 15; *P* < 0.05) on MSM than on BM during that first branching stage.

Some clustered LSGs and HET-domain genes exhibit coordinate expression. The majority of

LSGs were clustered into physically linked groups—64% with *het* or HET-domain genes

(**Table S4**). Forty-two HET-domain genes clustered with at least one LSG, including HET-

domain LSGs NCU03378, 07596, and 10839. Other than these three HET-domain LSGs, all

clustered HET-domain genes exhibited measurable expression in at least three out of the four

sampled stages in conidia germination as well as six out of eight stages sampled during sexual

development.

Coordinated expression among genes within the clusters during *N. crassa* asexual and

sexual growth and development was not common. Among 26 cases where gene-expression

dynamics were highly coordinated (at least one of the pairwise correlation coefficients were

greater than 0.5 among LSGs in the cluster; **Table S10**) across several developmental stages, two

are notable (**Fig. 5**). The first is cluster #69 of *het-14* and two LSGs (NCU07510 and 08191).

Genes in the cluster with *het-14* exhibited highly coordinated expression during sexual

development, even with a large non-coding sequence of over 15000 bp separating *het-14* and

NCU07510 in that cluster (**Fig. 5A–C**). Interestingly, the non-LSG genes located within the

cluster exhibited expression dynamics that were similar to the clustered orphan and HET-domain

genes during those stages. The second notable case where gene-expression dynamics were highly

coordinated across several developmental stages is the cluster #117 of a HET-domain gene

(NCU11054) and four LSGs (NCU03467, 03469, 03474, and 16509). Genes in the cluster #117

exhibited highly coordinated expression during asexual growth on MSM (**Fig. 5D–F**).

Expression was also observed to be coordinated in a few other clusters, such as among the LSGs

clustered with the HET-domain genes NCU07335, 10142, and 16851 during conidial

germination and asexual growth on BM and on MSM (**Fig. S4**): three clusters exhibited

coordinated expression across sexual development (**Fig. S4A–C**), eight exhibited coordinated

expression across conidial germination and asexual growth on Bird medium (**Fig. S4D–K**), and

15 exhibited coordinated expression across conidial germination and asexual growth on maple

sap medium (**Fig. S4L–Z**).

Figure 5. Expression profiles of two LSG-*het* gene clusters across asexual and sexual growth in *N. crassa*. expression profiles are plotted for LSG and HET-domain genes clustered with *het-14* (color coded, with 95% credible intervals)—as well as the non-LSG genes NCU08189 and NCU08910 located within the cluster (grey dashed)—cultured through (**A**) eight stages of sexual development on synthetic crossing medium, (**B**) four stages of conidial germination and asexual growth on Bird medium, and (**C**) four stages of conidial germination and

asexual growth on maple sap medium. Expression profiles for genes clustered with *het-14* (color coded, with 95% credible intervals), including the five non-LSG genes NCU03468, 03470, 03471, 03472, and 03473 (grey dashed) within the cluster, across (**D**) eight stages of sexual development on synthetic crossing medium, (**E**) four stages of conidial germination and asexual growth on Bird medium, and (**F**) four stages of conidial germination and asexual growth on maple sap medium.

Two of the three LSGs clusters, including the cluster #24 (**Fig. S4A**) and the cluster #131

(**Fig. S4C**), exhibited coordinated expression among the genes within each cluster during sexual

development, and the coordinate expression was observed during the early stages of sexual

development before meiosis (about 48–72 h after crossing). Of the eight clusters, expression

during asexual growth on BM was coordinately down-regulated in seven of them (**Fig. S4D–K**).

The exception was cluster #50 of HET-domain NCU07335 and LSG 07336; expression of these

genes was up-regulated during sexual development (**Fig. S4I**). Coordinate expression during

asexual growth on MSM exhibited an opposite pattern: 12 out of 15 clusters exhibited up-

regulated expression patterns toward the extension of the first hyphal branch (**Fig. S4L–Z**),

including two LSG-*het* clusters: the cluster #50 (**Fig. S4S**) and the cluster #51 of NCU16851

(HET-domain), 07316, 07317, and 07323 (**Fig. S4T**). In fact, coordinate expression was not only

observed between LSGs and HET-domain genes that were clustered together, but also observed

between LSGs in the clusters and neighboring non-LSGs (**Table S10**). Furthermore, genes in

cluster #112 exhibited no measurable expression in least at three out of the four sampled stages

across conidial germination and across asexual growth, and genes in cluster #172 exhibited no

measurable expression in at least six out of eight sampled stages in sexual development in

N. crassa.

Figure S4. Expression profiles of 21 LSGs clusters and LSG-*het* gene clusters (Table S4) across asexual and sexual growth in *N. crassa*. Expression and 95% credible intervals for (**A–C**) genes in clusters 24, 37, and 131 during sexual development, (**D–K**) genes in clusters 121, 128, 133, 88, 62, 50, 51 and 8 during conidial germination and asexual growth on Bird medium, and (**L–Z**) genes in clusters 22, 24, 33, 117, 63, 121, 65, 50, 51, 125, 87, 1, 8, 109 and 110 during conidial germination and asexual growth on maple sap medium.

Cell communication transcription factors affect expression of LSG and HET-domain genes.

The transcription factors *adv-1* and *pp-1* play multiple roles in asexual and sexual development,

cell growth and fusion, and cell communication in *N. crassa* and closely related fungi [49–53].

Functions and regulatory networks involving *adv-1* and *pp-1* were systematically investigated

[53], and in that study 155 genes were identified that were likely positively regulated by both

transcription factors, including two HET-domain genes (NCU03494 and 09954) and one LSG

(NCU17044). We reanalyzed the RNA sequencing data collected during conidial germination

from knockout mutants of *adv-1* and *pp-1* and from wild type strain [53]. In general, knocking

out the two transcription factors had a substantial impact on the activities of LSGs (**Figs. 6, S5,**

Tables 3, S11, S12). Unlike HET-domain genes and other non-LSG genes, a significantly large

portion of 173 LSGs exhibited no expression in the knockout mutants and the wild type strain,

including 196 not expressed in both *Δpp-1* and the wild type, and 195 not expressed in both

Δadv-1 and wild type. At the same time, significant numbers of LSGs were turned on or off by

the mutations to the two transcription factors. Namely, the same number of 44 LSGs that were

expressed in wild type were inactivated in the mutant strains *Δpp-1* or *Δadv-1*, and 23 LSGs were

inactivated in both the mutant strains *Δpp-1* and *Δadv-1*. Among the 23 LSGs, 20 were within the

predicted LSG-het clusters, including NCU04700 and 04710 that clustered with HET-domain

gene NCU04694 and three core genes NCU08822, 08829, and 08830 in a six-gene cluster. At the

same time, expression of significant numbers of LSGs (53 and 54 separately) went from

undetectable to detectable in the *Δpp-1* or *Δadv-1* knockout strains, with expression of 31 LSGs

becoming detectable in both knockout mutants. However, only 14 out of the 31 newly detectably

 expressed genes were clustered LSGs. Therefore, knocking out these two transcription factors

 did not positively affect the LSG-*het* gene clusters.

Figure 6. Impacts of transcription factor (TF) deletions on expression of HET-domain and LSGs within the *N. crassa* genome. Comparatively larger portions of LSGs are expressed exclusively in the mutants (dark red) or in the wildtype strains (dark blue). Expression profiles were classified in five categories (dark red: only measurable expression in mutant; light red: higher in mutant with $P < 0.05$; light grey: no significant difference with $P \geq 0.05$; dark grey: not measurable in either mutant or wildtype; light blue: significantly higher in wildtype with $P < 0.05$; dark blue: only present in wildtype;), and comparative portions of each category in HET-domain genes, LSGs, and other genes in the genome were reported. Data from Fischer et al. (2018) were reanalyzed to assess the impacts of *pp-1* and *adv-1* knockout mutants.

Table 3. Transcriptomics analysis results on *N. crassa* responses to genetic manipulation.

Experiments[1]	LSGs (670)		Het (66[2])		Other (8994)		Notes[3]
	expressed	silenced	expressed	silenced	expressed	silenced	LSG vs other (P)
Δpp-1	89 (106)	240 (249)	9 (21)	0 (2)	3582 (2921)	400 (457)	*P<0.01(P<0.01); P<0.01(P<0.01)*
Δadv-1	93 (90)	239 (249)	16 (21)	2 (2)	3207 (3055)	425 (457)	*P<0.01(P<0.01); P<0.01(P<0.01)*
mat a (1-2-1)	155 (169)	70 (366)	17 (41)	0 (0)	3957 (3966)	898 (913)	*P<0.01(P<0.01); P>0.01(P<0.01)*
Δpp-1 & Δadv-1[4]	49 (53)	196 (249)	12 (16)	0 (2)	1989 (1685)	303 (457)	*P<0.01(P<0.01); P<0.01(P<0.01)*
Δpp-1 & mat a[4]	26 (35)	202 (206)	2 (11)	0 (0)	1645 (1236)	237 (310)	*P<0.01(P<0.01); P<0.01(P<0.01)*
Mat a & Δadv-1[4]	32 (22)	197 (206)	5 (12)	0 (0)	1524 (1384)	259 (310)	*P<0.01(P<0.01); P<0.01(P<0.01)*

[1] Comparisons were performed between mutant and wild type. Expressed genes refer to those that were well measured in each experiment. Numbers of genes with significantly higher (or significantly lower when in parentheses) expression levels, with a LOX $P < 0.01$, and adjusted LOX $P < 0.05$, in the knockouts or the mutant were provided. Silenced genes refer to those without measurable expression, and numbers of silenced genes in mutant and in wild-type (in the parentheses) were reported.

[2] Three predicted HET-domain genes, NCU03378 (*tol*), 07596, and 10839, were also identified as LSGs.

[3] Chi-square test P values for deviation between counts of LSGs and non-LSG genes from marginal expectations for genes exhibiting significantly higher (red), or lower (blue) expression level, and for genes being silent in the

mutant (numbers in parentheses for silenced genes are the subset genes that exclusively silenced in the mutants). [4] Numbers of genes exhibiting significantly different expression between the wild type and co-existed in both mutants listed in the row (not a double-gene mutant). Specific gene IDs are listed in **Table S12**.

Twenty-seven HET-domain genes are expressed at higher level in the wildtype strain

than in both mutants (41 for *Δpp-1*, and 35 for *Δadv-1*), including four HET-domain genes

(NCU03494, 06583, 09037, and 09954) exhibited more than five-fold higher expression in the

wild type strain than in the *Δpp-1* and/or *Δadv-1* strains. However, NCU03494, 06583, and

09037 are HET-domain genes not clustered with any LSGs. For LSGs and HET-domain genes

with well measured expression in mutants and wild type, many genes exhibited similar up- or

down-regulation between the *Δpp-1* and *Δadv-1* compared with their expression in the wild type

(**Fig. S5**).

Figure S5. Heatmaps demonstrating divergent expression between wild-type strains and deletions of *pp-1* and *adv-1* in their expression of HET-domain and LSGs in conidial germlings, and in mutants of *mat 1-2-1* in fertilized

protoperithecia (crossing). (**A**) Expression divergence of HET-domain genes in three mutants *vs*. wild type, expression levels sampled in crossing were scaled according to the wild-type expression measured in germlings, and HET-domain genes with 0 measurements in wild type were excluded; (**B**) Expression divergence of LSGs in three mutants *vs*. wild type, expression levels sampled in crossing were scaled accordingly to the wild type expression measured in germlings, and LSGs with 0 measurements in wild type were excluded.

N. crassa mating loci regulate crossing and sexual development, and opposite mating-type

strains are ordinarily vegetatively heterokaryon-incompatible [54–57]. Transcriptome profiles

were compared between six-day cultures of the wildtype strain and a mating locus *mat 1-2-1*

mutant that has lost mating function (FGSC#4564, *mat a*[m1]s-3B cyh-1) on synthetic crossing

medium (SCM). Of 9758 measured genes, a total of 836 genes exhibited undetectable expression

only in either the mutant or wildtype (**Figs 6, S5**; **Tables S11, S12**), including 179 LSGs (109

expressed and 70 undetected in the *mat 1-2-1* mutant) and 657 non-LSG genes (336 expressed

and 321 undetected in the *mat 1-2-1* mutant). For LSGs, lack of detectable expression occurring

only in the wildtype or only in the mutant was significantly enriched ($P < 0.00001$, chi-squared

test). Of the 109 LSGs that were exclusively expressed in the *mat 1-2-1* mutant, 71 were located

within 55 predicted LSG-*het* clusters. Of the 70 LSGs that were only expressed in the wild type,

59 were located within 47 predicted LSG-*het* clusters. Only 17 clusters were common between

the two groups, and larger clusters with more than three genes presented behaved differently

between the *mat 1-2-1* mutant and wild type, with three genes of five-gene cluster #51

(NCU07306–07323) being inactivated in the mutant, while four genes of seven-gene cluster

#124 (NCU05480–06949) and five genes of the nine-gene cluster #94 (MCU07144–07152)

being activated in the mutant. Some two-gene clusters exhibited coordinated expression that was

detectable only in the mutant or the wild type. There were 53 LSGs expressed significantly

higher ($P < 0.05$) in the mating type mutant, and 111 expressed significantly higher in the wild

type. However, mutations in the mating locus have no such impacts on basic activities for HET-

domain genes, with 18 and 41 HET-domain genes being expressed significantly higher in the

mating locus mutant, and 41 HET-domain genes being expressed significantly higher in the wild

type (**Figs. 6, S5**).

Binding-site enrichment analysis using CiiiDER [58] identified no significant enrichment

of binding sites for *mat 1-2-1*, *adv-1*, or *pp-1* in the upstream 5000 bp of LSGs that exhibited

activity divergence in the mutants, specifying LSGs that whose expression was unchanged

between the mutant and wildtype as the background. LSGs and HET-domain genes were not

significantly enriched in genes that are potentially bound by *adv-1* or *pp-1*, based on data from

DAP-seq [SRP133627 from 53]. However, knocking out a key transcription factor encoding

gene, *ada-6* [59]—whose product regulates asexual and sexual growth—inhibited expression of

30 LSGs and promoted expression of 25 LSGs during conidiation and protoperithecial

production. Therefore, a substantial number of LSGs are at least peripherally involved in those

regulatory networks, abundant enough that they may play fine regulatory roles, and sparsely

distributed enough to impact diverse regulatory pathways.

Selection profiles of LSGs—an original study reported 135,000 SNPs in transcriptomic sequence

from 48 individuals [60], yet a total of 1,086,579 SNPs is reported from just 26 *N. crassa* isolates

whose transcriptomic sequence data are available in FungiDB. The discrepancy likely arises as a

consequence of the exclusion of singletons from the original data [60], which were explicitly left

unreported in the original paper, but were frequently observed (C. Ellison personal

communication, July 13, 2020). 97.8% (655 out of 670) LSGs feature at least one non-

synonymous SNP, which is moderately but statistically significantly higher (Fisher exact test, P

< 0.01) than the 91.6% (8271 out of 9027) non-LSG genes, as revealed by the inclusive SNP

data in FungiDB (**Fig. S6**). Unlike LSGs that generally encode less than 250 amino acids, most

HET-domain genes encode more than 300. Non-synonymous SNPs were observed at similar

frequencies between LSGs and HET-domain genes with similar gene length (**Fig. 7**). Excluding

singletons from the FungiDB data markedly reduced the number of SNPs—only eight

Neurospora LSGs then featured SNPs (1.3%) compared to 4192 non-LSG genes (45.9%).

Therefore, SNP singletons are abundant within *Neurospora* LSGs. A high density of SNP

singletons in short LSGs corresponds to a high allelic polymorphism of LSGs, present in a

comparative low proportion in the population, implying high linkage disequilibrium in LSGs.

Figure S6. SNPs per nucleotide of transcript—including singletons—for 9756 *N. crassa* genes (LSGs, red; non-LSG, blue) from 26 sequenced strains in FungiDB.

Figure 7. Nonsynonymous SNP distribution comparison between LSGs (blue circles) and HET-domain genes (red circles) from *N. crassa* population genetic sequencing [60]. Nonsynonymous SNPs are totaled across all 26 *N. crassa* strains deposited at FungiDB.

Of 603 *Neurospora* LSGs with SNPs data, including those with singleton SNPs, only

NCU07210 exhibited statistically significant strong gene-regional positive selection during the

divergence of *N. crassa* from its sister species *N. tetrasperma* (**Fig. S7A**). No abnormal

phenotype resulting from the knockout of this gene (*mat-a* strain FGSC14291) was observed in

our assays. For 695 LSGs shared within *Neurospora-Sordaria*, 16 genes exhibited statistically

significant evidence of strong gene-regional positive selection, and five exhibited statistically

significant evidence of moderately strong gene-regional positive selection (**Table S13**). Five of

16 strongly positively selected genes were identified in six biological processes related to the

response to stress or stimulus (**Table S14**), including the hypothetical protein coding gene

NCU02932 with the ortholog being annotated as a regulator of ATP-sensitive K^+ channels in

N. discreta and likely play roles in stress responses (**Fig. S7B**). Further exploration suggested

that positively selected *Neurospora-Sordaria* LSGs NCU05395, 07618, 09562, and 09693 are

likely involved in the regulation of sexual development [61]. Furthermore, regulation of the

moderately positively selected genes NCU01306, 00496, and 00748 is likely associated with

conidiation in *N. crassa* [59,62,63].

Figure S7. Window-free regionalized nucleotide site-specific selection profiles in two LSGs showing statistically significant gene-regional positive selection along the divergence between *Neurospora crassa* and its sister species *N. tetrasperma*. Scaled selection coefficients (model-averaged γ) are plotted (red: lower bound of γ > 4; blue: lower bound of γ ≤ 4; gray band: 95% model uncertainty interval) by nucleotide position for (**A**) NCU07210, a *Neurospora* LSG, and (**B**) NCU02932, a *Neurospora-Sordaria* LSGs.

KO phenotypes of LSGs. In an examination of phenotypes of crosses of 367 available KO strains

of the *Neurospora* LSGs to the KO of the opposite mating type (or the WT of the opposite

mating type if the KO of the opposite mating type was not available), two *Neurospora* LSGs,

NCU00176 and 00529, and one *Neurospora-Sordaria* LSG, NCU00201, showed a distinct

knockout phenotype in sexual development (**Fig. 8**). All these knockout mutants exhibited

arrested development with the protoperithecia failing to develop into perithecia. Interestingly,

both NCU00176 and NCU00201 were expressed at significantly higher levels in protoperithecia

than in mycelia after crossing. Expression of NCU00529 was observed only in the late stage of

conidial germination on maple medium. NCU00529, 00530, and 00531 are homologs, and

NCU00529 and 00530 exhibited no expression in sexual development and conidial germination.

No abnormal phenotypes were identified for NCU00531 knockout mutants (FGSC13078 and

13079), and no knockout mutants for NCU00530 are available. Knockouts of NCU00375,

00384, 00485, and 00491 (*Neurospora* LSGs) and NCU01623, 05395, 07618 (*Neurospora-*

Sordaria LSGs) have been reported to exhibit minor phenotypic anomalies during asexual

growth, especially at 37 C (Fungidb.org/fungidb). We examined knockout mutants of these

genes and confirmed increased pigment production in NCU00485 and dense and slow growth in

NCU00491 and 016223 at 37 C.

Figure 8. Knockout mutants of *Neurospora crassa* genes NCU00176, NCU00201, and NCU00529 produced arrested protoperithecia that failed to develop into perithecia and to produce sexual spores on SCM medium. Knockout cross ΔNCU00176 (FGSC12195 × 12196) exhibited (**A**) small protoperithecia (scale bar: 1 mm) and (**B**) squashed protoperithecia exhibiting an abortive ascogenous center (scale bar: 10 μm). Knockout cross ΔNCU00201 (FGSC18867 × 18868) exhibited (**C**) normal-sized protoperithecia (scale bar: 1 mm) with (**D**) abortive ascogenous centers (scale bar: 10 μm). Knockout cross ΔNCU00529 (FGSC13076 × 13077) exhibited (**E**) normal-sized protoperithecia with (**F**) abortive ascogenous centers.

Discussion

In this study, we systematically investigated LSGs in *Neurospora* genomes to determine

their location and organization on chromosomes and possible roles in biology and ecology.

Using genomic phylostratigraphy and reciprocal BLAST searching, we identified 670 genes that

are *Neurospora* LSGs, only shared within *Neurospora* species. More than 63% of the 670

Neurospora LSGs form clusters of 2–21 LSGs, interspersed with HET-domain putative

allorecognition genes. Many of the larger clusters aggregate near the telomeres of the seven

chromosomes. A majority of *Neurospora* LSGs are actively regulated in response to carbon

sources, light, and temperature changes that promote sexual or asexual reproduction. Knockout

mutant strains were phenotyped for 367 *Neurospora* LSGs, and arrested protoperithecia were

observed for mutants of three *Neurospora* LSGs. Our data indicate that regulation of asexual and

sexual reproduction surprisingly empower evolution of *de novo* elements more than the

regulation of core metabolism and cellular biology in *N. crassa*. More specifically, this

regulation is associated with reproduction modes affected by carbon resources—a critical

environmental factor for this post-fire fungus. However, many LSGs were not identified as

essential for *N. crassa* development and biology. Therefore, it is possible that LSGs play key

roles in fine regulation in response to environmental factors inducing reproduction processes that

have been understudied.

Neurospora LSGs and HET-domain genes are tightly clustered. *Neurospora* LSGs and HET-

domain genes exhibit several organizational features on chromosomes. These organizational

features include gene clustering, large non-coding spaces enclosed by flanking condensed coding

regions, high frequencies of gene duplications, and proximity to telomeres. If clustered genes

were derived from relocation and rearrangement, especially for functionally associated genes, it

would probably be easy for these genes to functionally integrate back altogether into the existing

system. Indeed, coordinated expression across clustered LSGs and HET-domain genes was

sometimes detected—especially during *N. crassa* development in response to various

environmental factors. However, LSG-*het* gene clusters and their neighbor genes were syntenic

mainly within very closely related taxa, and many LSGs exhibited no expression under common

laboratory settings—even across a range of nutrient conditions and developmental stages.

Interestingly, heterochromatic interactions of intra- and inter-telomeric contacts were reported as

common in *N. crassa* [64], implying potential *cis*- and *trans*-regulation at chromosomal level of

expression integration in these LSGs enriched regions. It is possible that integrating LSG-*het*

gene clusters into existing regulatory systems temporally and spatially modifies the original

functions that are expected only under common growth conditions. Therefore, specific

experiments conducted in extreme growth conditions, with rare nutrient types, and at

understudied stages of the life cycles should be conducted.

A previous study characterizing chromosome ends in *N. crassa* demonstrated that highly

AT-rich sequences in the telomeres are likely products of RIP and that subtelomeric elements

common in other fungi are absent in *N. crassa*. Telomere repeats are required for H3K27

methylation, which would repress the transcription activities and functionally silent genes in

these regions [65]. More importantly, the telomeric regions have potential significance in niche

adaptation and probably harbor hotspots for novel sequences due to abrupt sequence divergence

involving repeats [66]. Many genes of an annotated *het*-domain also locate near the ends of some

N. crassa chromosomes [33], and 42 of the 69 HET-domain proteins, which account for

heterokaryon incompatibility, are actually clustered with at least one *Neurospora* LSG. In fact,

neighboring regions of HET-domain genes are abundant with "young" genes that lack homologs

in lineages beyond *Neurospora* and closely related species in the Sordariales, and most neighbor

genes are syntenic within Sordariomycetes. Most LSGs were not active in sampled stages in the

N. crassa life cycle. However, a few LSG-*het* gene clusters exhibited coordinate expression

during early sexual development as well as during active hyphal tip growth on the maple sap

medium that support both asexual and sexual developments. Therefore, investigation of gene

activities in the telomeric and sub-telomeric regions during mycelium development and

allorecognition between same and different vegetative compatibility groups will shed light on

possible concerted associations between these two functional groups in heterokaryon

incompatibility, pre-mating process and even during early sexual development in *Neurospora*.

Actually, three LSGs, including NCU03378, 07596, and 10839, were annotated as a *het-*

domain, and NCU03378 was annotated as *tol* [tolerant, 55] and shared several conserved

sequence regions with *het*-6 that is involved in incompatibility in the *N. crassa* population [67].

Functional associations among some HET-domain genes and LSGs within the LSG-*het* clusters

were evidenced with similar gene expression regulation pattern during asexual and sexual

development in *N. crassa*. However, compared with the HET-domain genes that were all

expressed, many LSGs were not expressed in the sampled stages during the *N. crassa* life cycle.

Further investigation on epigenetic modification of gene expression near telomeric

neighborhoods could be critical to capture the right moments of associated functions between

LSGs and HET-doma*in* genes during allorecognition and to understand the evolution of LSGs in

N. crassa.

Neurospora LSGs play roles in response to key regulatory environmental factors. LSGs are

generally not functionally annotated, mainly due to the lack of homologous references in well

studied genomes. To determine the functional roles of *Neurospora* LSGs, we revisited recent

high quality transcriptomics studies on *N. crassa*, covering almost all morphological stages in the

N. crassa life cycle, cultures under different conditions and carbon or nitrogen resources, light

exposure, temperature change, as well as knockout mutants of key regulatory genes [22–

24,39,40,46,59,68–72]. We observed a significant enrichment of clustering of LSGs and HET-

domain genes and, within some of those clusters, highly coordinated regulation in response to

carbon resources, light, and temperature conditions. This study provides evidence that some

LSGs and clustered HET-domain genes are associated with metabolic adaptation to

environmental factors that are critical indicators of successful fungal asexual and sexual growth

and reproduction.

Three hundred forty-two LSGs that were actively expressed during sexual development

were also actively regulated in both asexual growth on two different media, including Bird

medium, which supports only asexual reproduction, and maple sap medium, which supports both

asexual and sexual growth. Nearly two thirds of the 342 genes were actively regulated in samples

collected from the media supplied with carbon resources that promote sexual growth of

N. crassa, supporting their possible roles in sexual development and the asexual-sexual switch.

In favor of their roles in sexual development in *N. crassa*, more than one third of the LSGs were

actively regulated on the presence of specific carbohydrates, including HMF and furfural,

compounds that powerfully stimulate initiation of sexual reproduction. *N. crassa* has been shown

to respond differentially to the two furans and to possess a high tolerance to furfural, which is

present in its natural habitat [40]. It is conceivable that some of the LSGs that are uniquely

expressed upon exposure to HMF or furfural could be further engineered to provide increased

tolerance to atypical carbon resources for *N. crassa*, a trait of significant interest in the pursuit of

robust biofuel production.

Neurospora LSGs were observed to be active only at the hyphal tips without measurable

expression from the colony samples, and thus these genes were considered to be responsible for

environmental sensing and interaction with microbes [73]. Overall, LSGs are probably

associated with reproductive development and growth in response to environmental conditions,

especially carbon resources, light and temperature. Furthermore, we observed that LSGs in

predicted clusters coordinately respond to these environmental factors. How these genes may

possibly contribute together as fine quantitative switches for reproduction requires further

investigation with multiple gene manipulations.

A significant proportion of LSGs are regulated by key developmental transcription factors.

Transcription factors, including *pp-1* and *adv-1* that play key roles in cellular communication

during asexual growth and mating loci that regulate sexual crossing, have been previously

reported [53,54,74]. From the previous transcriptomics data from knockout mutants of *pp-1* and

adv-1 and newly generated transcriptomics data from a mutant of *mat 1-2-1*, we observed that

the expression of a significant portion of LSGs was affected by mutations in these transcription

factors. Interestingly, over 95% of LSGs that were turned off in both *pp-1* and *adv-1* mutants

belonged to predicted clusters, but less than 50% LSGs that were turned on in both mutants

belonged to predicted clusters. The expression of six genes in the NCU07144–07152 LSGs

cluster was actively regulated during sexual development, in knockout mutants of *adv-1* and *pp-1*

that regulate cell communication with knockout phenotypes in sexual development [49,52], as

well as in samples in the absence of carbon. Expression of some LSGs was also affected in

knockout mutants of other regulatory genes, such as *ada-6* and *gul-1* that are critical for sexual

and asexual development in *N. crassa*. The most dramatic impacts to the expression of LSGs

were from the mutation in the mating locus, suggesting likely functional associations of the

mating process and sexual development initiation for LSGs. However, no binding sites of

transcription factors were observed enriched in the promote and up-stream sequences of LSGs

being turned on or off by those factors, and coordinate expression regulation was only detected

in a few LSG-*het* gene clusters. Instead of possible cis or trans regulations, an alternative

explanation for the associations between the LSGs and transcription factors would be that the

LSGs were coordinated with the sampled developmental stages when the transcription factors

were actively engaged. For example, the LSGs exhibited different expression activities during

hyphal branching on natural medium MSM, when cell-to-cell communication is regulated by

adv-1. Therefore, success in further investigations will be aided by focusing on the epigenetics of

the LSG-*het* gene clusters during specific periods of cell-to-cell communication.

Knockout phenotypes suggested that few LSGs play essential roles in Neurospora

development. Our phenotyping of available knockout mutants of 367 LSGs at the Fungal

Genetics Stock Center [75] yielded three genes with a phenotype in sexual development. Within

these three genes, NCU00529 and its homologs NCU00530 and 00531 form a cluster in the

subtelomere of chromosome I. The other two genes with mutant phenotypes in sexual

development, NCU00176 and 00201, exist on chromosome III. Their expression decreased after

protoperithecia. NCU00176 was reported as one of the eight *Neurospora* LSGs that were among

the 300 down-regulated hypothetical proteins in the *gul-1* knockout mutant [76], and GUL-1

plays multiple roles in *N. crassa* hyphal morphology development [77]. We also confirmed

increased pigment production in NCU00485 and dense and slow growth in NCU00491 and

016223 at 37 C.

A recent study reported 40 biologically relevant clusters (BRCs) for 1168 *N. crassa* genes

being phenotyped for 10 growth and developmental processes [78]. Eleven *Neurospora* and 31

Neurospora-Sordaria LSGs were included in this analysis. Interestingly, 4 out of the 11

Neurospora ($P < 0.05$) and 5 out of the 31 *Neurospora-Sordaria* LSGs were concentrated in one

of the 40 BRCs (Cluster 4 of 81 genes) with minimum yeast orthologs and generally no

significant phenotypes. An explanation for the lack of apparent phenotypes may be that these

genes 1) were not investigated under the conditions for the phenotypes; 2) were functionally non-

essential and/or not fully integrated into the regulatory networks, or 3) were functionally

redundant within clusters or with non-LSG paralogs.

LSGs are clustered with allorecognition loci and respond to G × E factors regulating the

switch from asexual to sexual reproduction in Neurospora. The co-location of LSGs and HET-

domain genes may suggest that they are functionally and evolutionarily linked. Our observations

show that the expression of LSGs is significantly altered during the transition from asexual to

sexual reproduction in *N. crassa*, in response to environmental factors such as carbon resources,

light, and temperature, as well as genetic regulators such as cell-to-cell communication

transcription factors and mating types. A substantial proportion of the LSGs are involved in

inducing and promoting sexual development in response to G × E factors. Coincidentally, during

the asexual-sexual transition, allorecognition governed by HET-domain genes, particularly

mating loci, shifts from being active to being repressed, allowing for crossing between different

Vegetative Compatibility groups (VCGs). Direct functional interactions between the two gene

groups would be difficult to investigate using standard transcriptomics approaches, as

allorecognition, which negatively regulates fusion between different VCGs, and sexual

reproduction occur in different developmental phases and under different environmental

conditions. Despite this, we observed limited coordinated responses to environmental and

genetic regulatory factors. Further investigation at the species and population levels, with a focus

on VCGs, is required to determine the evolutionary histories of LSGs and HET-domain genes,

which are linked by their co-location in recent lineages. Therefore, SNP data from various

populations, especially from different VCGs, would be of great interest in understanding the

evolutionary associations between the two gene groups. In fact, balancing selection has been

demonstrated to act on HET-domain genes, and abundant rare singletons have been observed for

LSGs, with a similar occurrence of SNP polymorphisms between HET-domain genes and LSGs

within the same molecular size, based on data from limited *N. crassa* populations

A working model provides a foundation for further investigation of* G × E *interactions of

***LSGs and HET-domain genes in* Neurospora *reproduction*.** The discovery of the co-location of

the LSGs and HET-domain genes at the fast-evolving hotspot in the *N. crassa* genome and of the

LSGs participation in G × E interactions that regulate the asexual-sexual switch inspired us to

develop a model to guide future investigation about possible interactions between LSGs and

HET-domain genes that are challenging in a standard experimental setting for *N. crassa*

molecular genetics (**Fig. 9**). The proposed working model is substantiated by the responses of

LSGs and clustered HET-domain genes to environmental factors, developmental stages, and key

regulatory transcription factors. This model suggests that the G × E regulations from asexual

growth and reproduction shift to sexual reproduction in *N. crassa* connect to activities of both the

LSGs and HET-domain genes.

Figure 9. Hypothesis regarding the roles of LSGs associated with allorecognition and environmental regulation of
reproduction modes. Pink and blue coded for vegetative compatibility groups (VCGs) that are governed by
allorecognition (HET-domain genes), and regulated by the LSGs and environment for being adaptive to different
growth environments E1 and E2 and are incompatible to each other. VCG colonies expand via aggregation and
anastomosis under preferred vegetative growth conditions until the environment turns on the asexual-sexual
switch and shifts into sexual reproduction. Plus (+) and minus (–) represent two opposite mating types. Crossing of
closely-related strains would further promote local segregation of the VCGs, while outbreeding would contribute
to polymorphism in the populations. LSGs may contribute to the consolidation of vegetative incompatibility when
environmental conditions are preferred for asexual growth, as a few HET-domain genes are LSGs. LSG expression in
environmental and genetic manipulations is consistent with roles in the induction of sexual development, as well
as roles in the repression of vegetative incompatibility when environmental conditions are preferred for sexual
reproduction. However, there is a lack of investigation of the complex but interlocking interactions among
environment, LSGs, and VCGs, and their epistatic interactions during fungal asexual and sexual growth. Therefore,
future investigations will reveal whether mutations in LSGs and allorecognition could contribute to parapatric or
even sympatric speciation due to ecological selection, or characterize the accumulation of mutations in LSGs and
HET-domain genes during allopatric and peripatric speciation due to physical barriers.

During asexual growth and reproduction, HET-domain genes restrict cell fusion to be

successful only between mycelia of the same VCG groups. Studies of fungal pathogens on plants

and animals suggest that fast-evolving LSGs, often as clusters, contribute to extensive

chromosomal structure polymorphisms that may drive the aggressiveness during host

colonization and the evolution of virulence [79–82]. When environmental conditions are

favorable for sexual development, LSGs may react and play roles in the initiation of sexual

development and the repression of HET-domain genes. Consistent with our hypothesis,

introgression of *N. crassa het-6* gene (previously identified as *tol*) into *N. tetrasperma* disrupts

the tolerance of both mating types in asexual tissues, thwarting pseudo-homothallism and

causing death of *N. tetrasperma [83]*. Three HET-domain genes are actually LSGs in *N. crassa*.

Nevertheless, the model proposes LSGs-environment associated mechanisms subsequent to

vegetative incompatibility during asexual colonization that enable crossing between compatible

mating types from different VCG backgrounds (**Fig. 9**).

This crossing between compatible mating types from different VCG backgrounds would

induce local homogeneity of VCG, permitting increased colony merging, nutritional exchange,

propagation of signals of disease and predators, and would increase the opportunity for mitotic

recombination via parasexuality. There are no laboratory studies evaluating fitness of VCGs on

controlled genetic backgrounds in different environments providing substantial evidence of

differentially adapted VCGs. However, field studies in *Aspergillus* demonstrated that specific

VCGs can have competitive advantages in specific agro-ecological zones [84]. Furthermore,

A. falvus VCG groups isolated from cotton in the southern USA are genetically isolated, despite

presumed sexual reproduction between the two opposite mating types, each of which was only

detected in one VCG group [85]. Moreover, locally predominant VCGs have been considered

crucial to the formation of isolated, asexual, inbreeding populations in *Fusarium oxysporum*, a

species in which sexual stage and meiotic recombination have been lost, based on an

evolutionary model proposed by Puhalla that was later supported by multiple studies [86,87].

Therefore, our model focusing on LSGs calls for investigation of how vegetative incompatibility

that promotes the local aggregation and anastomosis of identical VCGs would be associated with

specific environmental preferences in nutrients, temperature, light, and osmosis, as well as

parasite evasion, or other competitive advantages. Our model of LSG G × E interaction also

provides a route for experimental investigation of why there are predominant VCGs and rare

VCGs in distinct geographic regions or environmental settings.

It would be also interesting to test if such purifying selection would lead to genomic

homogeneity in allorecognition loci and be further enhanced supposing nearby crossings and

sexual reproduction. Similar with lineage-specific genes in other model systems [e.g. 88,89],

highly frequent SNPs, including SNP singletons, were observed in *N. crassa* LSGs, implying

either weak selection, or recent selection, or both. However, SNP and SNP singletons, often

considered to be at linkage disequilibrium, can be affected by various factors, and SNP

singletons have been assigned functional importance in the human genome [90]. However,

filtered SNP data from limited *N. crassa* population in FungiDB could not provide a full picture

regarding population structures of LSGs. Therefore, our model also suggests that long term and

large scale pangenomics would be critical to understanding whether outbreeding between VCGs

plays a substantial role in promoting allelic diversity in *Neurospora* (**Fig. 9**).

695
Materials and Methods

Mutant and culture conditions—Protoperithecia were sampled for *mat 1-2-1* mutant

(FGSC#4564, *mat a*[m1]s-3B cyh-1). The experiments were performed with macroconidia,

which were harvested from 5-day cultures on Bird medium (BM). 1×10^5 spores were placed

onto the surface of a cellophane-covered synthetic crossing medium (SCM) in Petri dishes

(60 mm, Falcon, Ref. 351007). Dark-colored protoperithecia were abundantly ripen in 6 days

after inoculation. Tissue samples were flash frozen in liquid nitrogen and stored at -80□.

Biological replicates included all tissues collected from multiple plates in one collection process.

Three biological replicates were prepared for each sampled point.

RNA isolation and transcriptome profiling, data acquisition and analysis—Total RNA was

extracted from homogenized tissue with TRI REAGENT (Molecular Research Center) as in

Clark et al. (2008)[91], and sample preparation and sequencing followed our previous works

[23,25,92]. Briefly, mRNA was purified using Dynabeads oligo(dT) magnetic separation

(Invitrogen). RNAseq Library Prep: mRNA was purified from approximately 200 ng of total

RNA with oligo-dT beads and sheared by incubation at 94 C in the presence of Mg (Roche Kapa

mRNA Hyper Prep Catalog # KR1352).

Following first-strand cDNAtA-tailing was performed with dUTP to generate strand-

specific sequencing libraries. Indexed libraries were quantified by qRT-PCR using a

commercially available kit (Roche KAPA Biosystems Cat# KK4854). The quality of cDNA

samples was verified with a bioanalyzer (Agilent Technologies 2100).

The cDNA samples were sequenced at the Yale Center for Genomics Analysis (YCGA).

The libraries underwent 76-bp single-end sequencing using an Illumina NovaSeq 6000 (S4 flow

cell) according to Illumina protocols. Adapter sequences, empty reads, and low-quality

sequences were removed. Trimmed reads were aligned to the *N. crassa* OR74A v12 genome

from the Broad Institute [93] using `HISAT2` v2.1, indicating that reads correspond to the reverse

complement of the transcripts and reporting alignments tailored for transcript assemblers.

Alignments with a quality score below 20 were excluded from further analysis. Reads were

counted for each gene with `StringTie` v1.3.3 and the Python script `prepDE.py` provided in

the package. `StringTie` was limited to report reads that matched the reference annotation.

Sequence data and experiment details were made available (GSE199259) at the GEO database

(https://www.ncbi.nlm.nih.gov/geo/).

Statistical analysis of the sequenced cDNA tallies for each sample was performed with `LOX`

v1.6 [94], ignoring raw reads that mapped ambiguously or to multiple loci.

Identification and verification of Neurospora crassa lineage specific genes (LSGs)—We

applied a two-step strategy to identify and verify *Neurospora* LSGs, including (1) a

phylostratigraphic approach to reveal putative LSGs and (2) confirmation using BLAST against

all genomes available at NCBI and fungal genomes at FungiDB. In the first step, we employed

previously published genomic phylostratigraphy for the *N. crassa* genome that reported over

2000 *N. crassa* orphan genes, discovered by genomic comparisons versus

Chaetomium globosum, *Ascrospora strigata*, *Saccharomyces cerevisiae*, *Phanerochaete*

chrysosporium, *Drosophila melanogaster*, and *Arabidopsis thaliana* [29]. *N. crassa* orphan-gene

status was determined via the Smith-Waterman pairwise similarity of protein-coding sequences

[95,96]. Classifications of genes in *N. crassa* were constructed as mutually exclusive groups

ranked in by their phylostratigraphy, including Euk/Prok-core, Dikarya-core, Ascomycota-core,

Pezizomycotina-specific, *N. crassa*-orphans, and others [11,97]. Putative *Neurospora* LSGs were

also compared with the newest annotation of the *N. crassa* genome. Consequently, the number of

predicted *N. crassa* LSGs based on the representative genomes was narrowed down to 1872

genes.

Many fungal genomes were recently published due to the efforts of the 1000 fungal

genome project launched by the DOE Joint Genome Institute [32]. These fungal genomes—

especially those well-sampled among closely related species—make it possible to be confident

that LSGs are likely the product of de novo gene evolution rather than birth-and-death processes

[98]. To verify that the 1872 genes are not present in species that were not analyzed within the

phylostratigraphic and previous BLAST analyses, we employed BLASTp and tBLASTx to

search in the entire NCBI GenBank database specifying an *E*-value cutoff of 0.05. To utilize

newer genome annotations that were available on GenBank, BLASTp and tBLASTx were used

again in FungiDB with an *E*-value cutoff of 10. These searches included genomes closely related

to the Neurospora genomes, including *Podospora*, *Pyricularia*, *Ophiostoma*, *Chaetomium*, and

Sordaria species. Any gene with a hit that was not from *Neurospora* was removed. This analysis

results in 670 genes herein termed *Neurospora* LSGs.

Due to the frequent duplication history behind *Neurospora* lineage-specific genes, we

enforced a strict expect threshold that identified 670 genes that likely are unique in *Neurospora*.

To understand how these 670 LSGs identified from *N. crassa* are shared among the *Neurospora*

genomes, a 3-way reciprocal `blastp` and a `tblastx` with a PAM30 score matrix were used to

search for homologous genes among the three *Neurospora* genomes, and an *E* value of 1×10^{-10}

was used as a cutoff, and synteny among the orthologs shared within the three *Neurospora*

genomes was further visually checked at the FungiDB. Species-specific genes in *N. tetrasperma*

and *N. discreta* were not investigated in this study, except for some specific cases mentioned in

the text. For uncertain recent duplications, we also relied on the ortholog group identification in

FungiDB, which reported potential homologs in 286 fungal and fungus-like Oomycetes

genomes, including genomes for 35 Sordariomycetes species closely related to *Neurospora*.

When necessary, additional phylogenetic analyses using the sequences downloaded from the

FungalDB homologs groups were pursued to identify lineage-specific genes in *Neurospora.*

Expression and functional analyses LSGs—Genome-wide gene expression was investigated in

N. crassa along multiple stages of its life cycle, including conidial germination on different

media [23] and production of meiotic propagules (ascospores) on synthetic crossing medium

[22]. Transcriptomic data of GSE41484 was reanalyzed with the latest annotation of the

N. crassa genome. The tally for each sample was processed with LOX v1.6 [94] to analyze gene

expression levels across all data points, which uses a Bayesian algorithm to amalgamate different

types of datasets. Gene counts or RPKM reads were analyzed by LOX, reporting relative

expression of each gene normalized to the lowest treatment, 95% confidence intervals for

relative expression, and statistical significance of expression differences. *P* values were adjusted

following the procedure of Benjamini et al [99,100]. To assess environmental impacts on

expression of LSGs, recent available data on 37 C BM from this lab (GSE168995) and a 240-

minute time course of asexual growth in response to darkness and light stimulation [46] were

revisited. To assess possible roles of LSGs in metabolic regulation, transcriptomics data from

mycelia exposed to 5 different carbon resources from crop residues [39] and from mycelia in

response to non-preferred carbon sources such as furfural and 5-hydroxymethyl furfural [HMF;

40] were also examined separately. To assess the gene expression effects of mutations of

transcription factors, including *adv-1*, *pp-1*, and *ada-6*, transcriptomics data from Fisher et al.

(2018) and Sun et al. (2019) [53,59] were also analyzed with LOX v1.6 [94]. Coordinated

expression among genes within the LSG clusters was first manually examined. Twenty-six LSG

clusters and their neighbor genes were selected for computation of pairwise correlation

coefficients using R. Pairs of genes exhibiting coordination coefficients higher than 0.5 were

considered coordinately expressed.

Clustering analyses of LSGs heterogeneous variation sites and profiling of historical selection

—The chromosomal distribution and clustering of LSGs—as well as LSGs and HET-domain

genes—were analyzed with Cluster Locator [36]. Cluster Locator requires a parameter (`Max-`

`Gap`) that specifies the number of genes that can be "skipped" between genes that are considered to

be part of a cluster. We set `Max-Gap = 5, 1`, and `0` (`Max-Gap`. Statistically significant clusters

($P < 0.01$) were reported. To provide a more continuous measure of gene clustering, a vector of

0s and 1s representing non-LSG genes and LSGs was generated as an input sequence for

MACML, a powerful algorithm to profile clustering of discrete ordered data [101]. This

algorithm calculates all likely models of linear clustering by partitioning the entire sequence into

all possible clusters and subclusters, and all models are statistically evaluated for information-

optimality via Akaike Information Criterion [102], 'corrected' Akaike Information criterion

[103], or Bayesian Information Criterion [104]. In this study, weighted likelihoods were

computed based on the conservative Bayesian Information Criterion for each model. LSG cluster

probabilities were calculated as weighted averages of models, and ninety-five percent model

uncertainty intervals were calculated by further analyses of the model distributions.

To quantify historical selection occurring during divergence of *N. crassa* from its sister

species *N. tetrasperma*, window-free regionalized nucleotide site-specific selection profiles were

computed on the sequences of LSGs based on the SNP data retrieved from the FungiDB database

[105] using Model Averaged Site Selection via Poisson Random Field (MASS-PRF; [106].

MASS-PRF identifies an ensemble of intragenic clustering models for polymorphic (SNPs in *N.*

crassa) and divergent sites against orthologs in the *N. tetrasperma* genome (FGSC2508, *mat A*

strain, https://mycocosm.jgi.doe.gov/Neute_matA2/Neute_matA2.home) using the Poisson

Random Field equations [107] to estimate selection intensity on a regionalized site-by-site basis.

Knockout strains and phenotype identification—Knockout strains for more than 9600 genes

[52], including deletion cassettes for genes in either of the two mating types, were acquired from

the Fungal Genetic Stock Center [FGSC: 75]. Identified *Neurospora* LSGs were examined for

altered phenotypes during conidia germination on Bird Medium (BM) and sexual development

on Synthetic Crossing Medium (SCM) from protoperithecium differentiation to ascospore

release. Genotype *mat A* strains were assayed for phenotypes when available; otherwise, *mat a*

strains were used. All available KO strains were phenotyped on BM and on SCM with three

replicates. For each investigated strain, 3000–5000 conidia were plated onto 90 mm diameter

plates and monitored, and crossing was conducted between opposite mating types. Three

independent phenotyping experiments were performed with each knockout strain using stored

conidia supplied by the FGSC.

Acknowledgements: We thank the Broad Institute, FungiDB and JGI for making *Neurospora*

related fungi genomic data available. All authors have declared that no competing interests exist.

This study was supported by funding to JPT from the National Institutes of Health R01 grant

AI146584, by the National Science Foundation grant IOS 1457044 to JPT, National Science

Foundation IOS 1456482 to FT, and BSF-2018712 to OY. The funders had no role in the study

design, data collection and interpretation, or the decision to submit the work for publication.

836
837   References:

838   1.   Weisman CM. The Origins and Functions of De Novo Genes: Against All Odds? J Mol

839   Evol. 2022;90: 244–257.

840 2. Tautz D, Domazet-Lošo T. The evolutionary origin of orphan genes. Nat Rev Genet.
841   2011;12: 692–702.

842 3. McLysaght A, Hurst LD. Open questions in the study of de novo genes: what, how and
843   why. Nat Rev Genet. 2016;17: 567–578.

844 4. Su Z, Townsend JP. Utility of characters evolving at diverse rates of evolution to resolve
845   quartet trees with unequal branch lengths: analytical predictions of long-branch effects.
846   BMC Evol Biol. 2015;15: 86.

847 5. Dornburg A, Su Z, Townsend JP. Optimal Rates for Phylogenetic Inference and
848   Experimental Design in the Era of Genome-Scale Data Sets. Syst Biol. 2019;68: 145–156.

849 6. Weisman CM, Murray AW, Eddy SR. Many, but not all, lineage-specific genes can be
850   explained by homology detection failure. PLoS Biol. 2020;18: e3000862.

851 7. Ruiz-Orera J, Hernandez-Rodriguez J, Chiva C, Sabidó E, Kondova I, Bontrop R, et al.
852   Origins of De Novo Genes in Human and Chimpanzee. PLoS Genet. 2015;11: e1005721.

853 8. Begun DJ, Lindfors HA, Thompson ME, Holloway AK. Recently evolved genes identified
854   from *Drosophila yakuba* and *D. erecta* accessory gland expressed sequence tags.
855   Genetics. 2006;172: 1675–1681.

856 9. Begun DJ, Lindfors HA, Kern AD, Jones CD. Evidence for de novo evolution of testis-
857   expressed genes in the *Drosophila yakuba/Drosophila erecta* clade. Genetics. 2007;176:
858   1131–1137.

859 10. Zhang L, Ren Y, Yang T, Li G, Chen J, Gschwend AR, et al. Rapid evolution of protein
860   diversity by de novo origination in *Oryza*. Nat Ecol Evol. 2019;3: 679–690.

861 11. Cai JJ, Woo PCY, Lau SKP, Smith DK, Yuen K-Y. Accelerated evolutionary rate may be
862   responsible for the emergence of lineage-specific genes in ascomycota. J Mol Evol.
863   2006;63: 1–11.

864 12. Domazet-Loso T, Brajković J, Tautz D. A phylostratigraphy approach to uncover the
865   genomic history of major adaptations in metazoan lineages. Trends Genet. 2007;23: 533–
866   539.

867 13. Casola C. From De Novo to "De Nono": The Majority of Novel Protein-Coding Genes
868   Identified with Phylostratigraphy Are Old Genes or Recent Duplicates. Genome Biol Evol.
869   2018;10: 2906–2918.

870 14. Vakirlis N, Carvunis A-R, McLysaght A. Synteny-based analyses indicate that sequence
871   divergence is not the main source of orphan genes. Elife. 2020;9. doi:10.7554/eLife.53500

872 15. Gladieux P, De Bellis F, Hann-Soden C, Svedberg J, Johannesson H, Taylor JW.
873   *Neurospora* from Natural Populations: Population Genomics Insights into the Life History of
874   a Model Microbial Eukaryote. In: Dutheil JY, editor. Statistical Population Genomics. New
875   York, NY: Springer US; 2020. pp. 313–336.

16. Davis RH, Perkins DD. Timeline: *Neurospora*: a model of model microbes. Nat Rev Genet.
2002;3: 397–403.

17. Mitchell MB. A model predicting characteristics of genetic maps in *Neurospora crassa*.
Nature. 1965;205: 680–682.

18. Ebbole D. *Neurospora*: a new (?) model system for microbial genetics: Neurospora 2000,
Asilomar, CA, USA, 9–12 March 2000. Trends Genet. 2000;16: 291–292.

19. Zhang N, Wang Z. 3 Pezizomycotina: Sordariomycetes and Leotiomycetes. In: McLaughlin
DJ, Spatafora JW, editors. Systematics and Evolution: Part B. Berlin, Heidelberg: Springer
Berlin Heidelberg; 2015. pp. 57–88.

20. Nowrousian M, Würtz C, Pöggeler S, Kück U. Comparative sequence analysis of *Sordaria*
macrospora and Neurospora crassa as a means to improve genome annotation. Fungal
Genet Biol. 2004;41: 285–292.

21. Zámocký M, Tafer H, Chovanová K, Lopandic K, Kamlárová A, Obinger C. Genome
sequence of the filamentous soil fungus *Chaetomium cochliodes* reveals abundance of
genes for heme enzymes from all peroxidase and catalase superfamilies. BMC Genomics.
2016;17: 763.

22. Wang Z, Lopez-Giraldez F, Lehr N, Farré M, Common R, Trail F, et al. Global gene
expression and focused knockout analysis reveals genes associated with fungal fruiting
body development in *Neurospora crassa*. Eukaryot Cell. 2014;13: 154–169.

23. Wang Z, Miguel-Rojas C, Lopez-Giraldez F, Yarden O, Trail F, Townsend JP. Metabolism
and Development during Conidial Germination in Response to a Carbon-Nitrogen-Rich
Synthetic or a Natural Source of Nutrition in *Neurospora crassa*. MBio. 2019;10.
doi:10.1128/mBio.00192-19

24. Lehr NA, Wang Z, Li N, Hewitt DA, López-Giráldez F, Trail F, et al. Gene expression
differences among three *Neurospora* species reveal genes required for sexual reproduction
in *Neurospora crassa*. PLoS One. 2014;9: e110398.

25. Wang Z, López-Giráldez F, Wang J, Trail F, Townsend JP. Integrative Activity of Mating
Loci, Environmentally Responsive Genes, and Secondary Metabolism Pathways during
Sexual Development of *Chaetomium globosum*. MBio. 2019;10. doi:10.1128/mBio.02119-
19

26. Corcoran P, Anderson JL, Jacobson DJ, Sun Y, Ni P, Lascoux M, et al. Introgression
maintains the genetic integrity of the mating-type determining chromosome of the fungus
Neurospora tetrasperma. Genome Res. 2016;26: 486–498.

27. Galagan JE, Selker EU. RIP: the evolutionary cost of genome defense. Trends Genet.
2004;20: 417–423.

28. Gladyshev E. Repeat-Induced Point Mutation and Other Genome Defense Mechanisms in
Fungi. Microbiol Spectr. 2017;5. doi:10.1128/microbiolspec.FUNK-0042-2017

29. Kasuga T, Mannhaupt G, Glass NL. Relationship between phylogenetic distribution and

genomic features in *Neurospora crassa*. PLoS One. 2009;4: e5286.

30. Haridas S, Salamov A, Grigoriev IV. Fungal Genome Annotation. Methods Mol Biol.
2018;1775: 171–184.

31. Grigoriev I. Fungal Genomics Program. Lawrence Berkeley National Lab.(LBNL), Berkeley,
CA (United States); 2012. Available: https://www.osti.gov/biblio/1165591

32. Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, et al. MycoCosm portal: gearing
up for 1000 fungal genomes. Nucleic Acids Res. 2014;42: D699–704.

33. Zhao J, Gladieux P, Hutchison E, Bueche J, Hall C, Perraudeau F, et al. Identification of
Allorecognition Loci in *Neurospora crassa* by Genomics and Evolutionary Approaches. Mol
Biol Evol. 2015;32: 2417–2432.

34. Ament-Velásquez SL, Vogan AA, Granger-Farbos A, Bastiaans E, Martinossi-Allibert I,
Saupe SJ, et al. Allorecognition genes drive reproductive isolation in *Podospora anserina*.
Nat Ecol Evol. 2022;6: 910–923.

35. Wang Z, Gudibanda A, Ugwuowo U, Trail F, Townsend JP. Using evolutionary genomics,
transcriptomics, and systems biology to reveal gene networks underlying fungal
development. Fungal Biol Rev. 2018;32: 249–264.

36. Pazos Obregón F, Soto P, Lavín JL, Cortázar AR, Barrio R, Aransay AM, et al. Cluster
Locator, online analysis and visualization of gene clustering. Bioinformatics. 2018;34:
3377–3379.

37. Saupe SJ, Kuldau GA, Smith ML, Glass NL. The product of the *het-C* heterokaryon
incompatibility gene of *Neurospora crassa* has characteristics of a glycine-rich cell wall
protein. Genetics. 1996;143: 1589–1600.

38. Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, et al. The genome
sequence of the filamentous fungus *Neurospora crassa*. Nature. 2003;422: 859–868.

39. Wang B, Cai P, Sun W, Li J, Tian C, Ma Y. A transcriptomic analysis of Neurospora crassa
using five major crop residues and the novel role of the sporulation regulator *rca-1* in
lignocellulase production. Biotechnol Biofuels. 2015;8: 21.

40. Feldman D, Kowbel DJ, Cohen A, Glass NL, Hadar Y, Yarden O. Identification and
manipulation of genes involved in sensitivity to furfural. Biotechnol Biofuels. 2019;12: 210.

41. Eilers FI, Sussman AS. Conversion of furfural to furoic acid and furfuryl alcohol by
Neurospora ascospores. Planta. 1970;94: 253–264.

42. Emerson MR. Chemical Activation of Ascospore Germination in *Neurospora crassa*. J
Bacteriol. 1948;55: 327–330.

43. Kritsky MS, Belozerskaya TA, Sokolovsky VY, Filippovich SY. Photoreceptor Apparatus of
the Fungus *Neurospora crassa*. Mol Biol. 2005;39: 514–528.

44. Wang Z, Wang J, Li N, Li J, Trail F, Dunlap JC, et al. Light sensing by opsins and fungal

ecology: NOP-1 modulates entry into sexual reproduction in response to environmental
cues. Mol Ecol. 2018;27: 216–232.

45. Wang Z, Li N, Li J, Dunlap JC, Trail F, Townsend JP. The Fast-Evolving *phy-2* Gene
Modulates Sexual Development in Response to Light in the Model Fungus *Neurospora*
crassa. MBio. 2016;7: e02148.

46. Wu C, Yang F, Smith KM, Peterson M, Dekhang R, Zhang Y, et al. Genome-wide
characterization of light-regulated genes in *Neurospora crassa*. G3. 2014;4: 1731–1745.

47. Chen C-H, Ringelberg CS, Gross RH, Dunlap JC, Loros JJ. Genome-wide analysis of light-
inducible responses reveals hierarchical light signaling in *Neurospora*. EMBO J. 2009;28:
1029–1042.

48. Káldi K, González BH, Brunner M. Transcriptional regulation of the *Neurospora* circadian
clock gene *wc-1* affects the phase of circadian output. EMBO Rep. 2006;7: 199–204.

49. Dekhang R, Wu C, Smith KM, Lamb TM, Peterson M, Bredeweg EL, et al. The *Neurospora*
Transcription Factor ADV-1 Transduces Light Signals and Temporal Information to Control
Rhythmic Expression of Genes Involved in Cell Fusion. G3. 2017;7: 129–142.

50. Fu C, Iyer P, Herkal A, Abdullah J, Stout A, Free SJ. Identification and characterization of
genes required for cell-to-cell fusion in *Neurospora crassa*. Eukaryot Cell. 2011;10: 1100–
1109.

51. Lan N, Ye S, Hu C, Chen Z, Huang J, Xue W, et al. Coordinated Regulation of
Protoperithecium Development by MAP Kinases MAK-1 and MAK-2 in *Neurospora crassa*.
Front Microbiol. 2021;12: 769615.

52. Colot HV, Park G, Turner GE, Ringelberg C, Crew CM, Litvinkova L, et al. A high-
throughput gene knockout procedure for *Neurospora* reveals functions for multiple
transcription factors. Proc Natl Acad Sci U S A. 2006;103: 10352–10357.

53. Fischer MS, Wu VW, Lee JE, O'Malley RC, Glass NL. Regulation of Cell-to-Cell
Communication and Cell Wall Integrity by a Network of MAP Kinase Pathways and
Transcription Factors in. Genetics. 2018;209: 489–506.

54. Pöggeler S, Kück U. Comparative analysis of the mating-type loci from *Neurospora crassa*
and *Sordaria macrospora*: identification of novel transcribed ORFs. Mol Gen Genet.
2000;263: 292–301.

55. Newmeyer D. A suppressor of the heterokaryon-incompatibility associated with mating type
in *Neurospora crassa*. Can J Genet Cytol. 1970;12: 914–926.

56. Jacobson DJ. Control of mating type heterokaryon incompatibility by the *tol* gene in
Neurospora crassa and *N. tetrasperma*. Genome. 1992;35: 347–353.

57. Xiang Q, Glass NL. The control of mating type heterokaryon incompatibility by vib-1, a
locus involved in het-c heterokaryon incompatibility in *Neurospora crassa*. Fungal Genet
Biol. 2004;41: 1063–1076.

58. Gearing LJ, Cumming HE, Chapman R, Finkel AM, Woodhouse IB, Luu K, et al. CiiiDER: A
tool for predicting and analyzing transcription factor binding sites. PLoS One. 2019;14:
e0215495.

59. Sun X, Wang F, Lan N, Liu B, Hu C, Xue W, et al. The Zn(II)2Cys6-Type Transcription
Factor ADA-6 Regulates Conidiation, Sexual Development, and Oxidative Stress
Response in. Front Microbiol. 2019;10: 750.

60. Ellison CE, Hall C, Kowbel D, Welch J, Brem RB, Glass NL, et al. Population genomics and
local adaptation in wild isolates of a model microbial eukaryote. Proc Natl Acad Sci U S A.
2011;108: 2831–2836.

61. Leeder AC, Jonkers W, Li J, Glass NL. Early colony establishment in *Neurospora crassa*
requires a MAP kinase regulatory network. Genetics. 2013;195: 883–898.

62. Sun X, Yu L, Lan N, Wei S, Yu Y, Zhang H, et al. Analysis of the role of transcription factor
VAD-5 in conidiation of *Neurospora crassa*. Fungal Genet Biol. 2012;49: 379–387.

63. Greenwald CJ, Kasuga T, Glass NL, Shaw BD, Ebbole DJ, Wilkinson HH. Temporal and
spatial regulation of gene expression during asexual development of *Neurospora crassa*.
Genetics. 2010;186: 1217–1230.

64. Rodriguez S, Ward A, Reckard AT, Shtanko Y, Hull-Crew C, Klocko AD. The genome
organization of *Neurospora crassa* at high resolution uncovers principles of fungal
chromosome topology. G3. 2022;12. doi:10.1093/g3journal/jkac053

65. Jamieson K, McNaught KJ, Ormsby T, Leggett NA, Honda S, Selker EU. Telomere repeats
induce domains of H3K27 methylation in *Neurospora*. Elife. 2018;7.
doi:10.7554/eLife.31216

66. Wu C, Kim Y-S, Smith KM, Li W, Hood HM, Staben C, et al. Characterization of
chromosome ends in the filamentous fungus *Neurospora crassa*. Genetics. 2009;181:
1129–1145.

67. Mir-Rashed N, Jacobson DJ, Dehghany MR, Micali OC, Smith ML. Molecular and
functional analyses of incompatibility genes at het-6 in a population of *Neurospora crassa*.
Fungal Genet Biol. 2000;30: 197–205.

68. Xiong Y, Wu VW, Lubbe A, Qin L, Deng S, Kennedy M, et al. A fungal transcription factor
essential for starch degradation affects integration of carbon and nitrogen metabolism.
PLoS Genet. 2017;13: e1006737.

69. Znameroski EA, Coradetti ST, Roche CM, Tsai JC, Iavarone AT, Cate JHD, et al. Induction
of lignocellulose-degrading enzymes in *Neurospora crassa* by cellodextrins. Proc Natl Acad
Sci U S A. 2012;109: 6012–6017.

70. Coradetti ST, Craig JP, Xiong Y, Shock T, Tian C, Glass NL. Conserved and essential
transcription factors for cellulase gene expression in ascomycete fungi. Proc Natl Acad Sci
U S A. 2012;109: 7397–7402.

71. Coradetti ST, Xiong Y, Glass NL. Analysis of a conserved cellulase transcriptional regulator

reveals inducer-independent production of cellulolytic enzymes in *Neurospora crassa*.
Microbiologyopen. 2013;2: 595–609.

72.   Craig JP, Coradetti ST, Starr TL, Glass NL. Direct target network of the *Neurospora crassa*
plant cell wall deconstruction regulators CLR-1, CLR-2, and XLR-1. MBio. 2015;6: e01452–
15.

73.   Kasuga T, Glass NL. Dissecting colony development of *Neurospora crassa* using mRNA
profiling and comparative genomics approaches. Eukaryot Cell. 2008;7: 1549–1564.

74.   Bobrowicz P, Pawlak R, Correa A, Bell-Pedersen D, Ebbole DJ. The *Neurospora crassa*
pheromone precursor genes are regulated by the mating type locus and the circadian
clock. Mol Microbiol. 2002;45: 795–804.

75.   McCluskey K, Wiest A, Plamann M. The Fungal Genetics Stock Center: a repository for 50
years of fungal genetics research. J Biosci. 2010;35: 119–126.

76.   Herold I, Kowbel D, Delgado-Álvarez DL, Garduño-Rosales M, Mouriño-Pérez RR, Yarden
O. Transcriptional profiling and localization of GUL-1, a COT-1 pathway component, in
Neurospora crassa. Fungal Genet Biol. 2019;126: 1–11.

77.   Herold I, Zolti A, Garduño-Rosales M, Wang Z, López-Giráldez F, Mouriño-Pérez RR, et al.
The GUL-1 Protein Binds Multiple RNAs Involved in Cell Wall Remodeling and Affects the
MAK-1 Pathway in *Neurospora crassa*. Front Fungal Biol. 2021;2.
doi:10.3389/ffunb.2021.672696

78.   Carrillo AJ, Cabrera IE, Spasojevic MJ, Schacht P, Stajich JE, Borkovich KA. Clustering
analysis of large-scale phenotypic data in the model filamentous fungus *Neurospora*
crassa. BMC Genomics. 2020;21: 755.

79.   Plissonneau C, Stürchler A, Croll D. The Evolution of Orphan Regions in Genomes of a
Fungal Pathogen of Wheat. MBio. 2016;7. doi:10.1128/mBio.01231-16

80.   de Jonge R, Bolton MD, Kombrink A, van den Berg GCM, Yadeta KA, Thomma BPHJ.
Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen.
Genome Res. 2013;23: 1271–1282.

81.   Hartmann FE, Sánchez-Vallet A, McDonald BA, Croll D. A fungal wheat pathogen evolved
host specialization by extensive chromosomal rearrangements. ISME J. 2017;11: 1189–
1204.

82.   Wacker T, Helmstetter N, Wilson D, Fisher MC, Studholme DJ, Farrer RA. Two-speed
genome evolution drives pathogenicity in fungal pathogens of animals. Proc Natl Acad Sci
U S A. 2023;120: e2212633120.

83.   Shiu PKT, Glass NL. Molecular Characterization of *tol*, a Mediator of Mating-Type-
Associated Vegetative Incompatibility in *Neurospora crassa*. Genetics. 1999;151: 545–555.

84.   Mitema A, 1 OMICS Research Group, Department of Biotechnology, Vaal University of
Technology, Vanderbijlpark, Africa S, et al. Molecular and Vegetative Compatibility Groups
Characterization of *Aspergillus flavus* Isolates from Kenya. AIMS Microbiol. 2020. pp. 231–

250. doi:10.3934/microbiol.2020015

85.   Grubisha LC, Cotty PJ. Genetic isolation among sympatric vegetative compatibility groups
of the aflatoxin-producing fungus *Aspergillus flavus*. Mol Ecol. 2010;19: 269–280.

86.   Puhalla JE. Classification of strains of *Fusarium oxysporum* on the basis of vegetative
compatibility. Can J Bot. 1985;63: 179–183.

87.   Elias KS. Vegetative Compatibility Groups in *Fusarium oxysporum* f. sp. *lycopersici*.
Phytopathology. 1991;81: 159.

88.   Cai JJ, Petrov DA. Relaxed purifying selection and possibly high rate of adaptation in
primate lineage-specific genes. Genome Biol Evol. 2010;2: 393–409.

89.   Hartfield M, Poulsen NA, Guldbrandtsen B, Bataillon T. Using singleton densities to detect
recent selection in. Evol Lett. 2021;5: 595–606.

90.   Ke X, Taylor MS, Cardon LR. Singleton SNPs in the human genome and implications for
genome-wide association studies. Eur J Hum Genet. 2008;16: 506–515.

91.   Clark TA, Guilmette JM, Renstrom D, Townsend JP. RNA extraction, probe preparation,
and competitive hybridization for transcriptional profiling using *Neurospora crassa* long-
oligomer DNA microarrays. Fungal Genet Rep. 2008;55: 18–28.

92.   Trail F, Wang Z, Stefanko K, Cubba C, Townsend JP. The ancestral levels of transcription
and the evolution of sexual phenotypes in filamentous fungi. PLoS Genet. 2017;13:
e1006867.

93.   Borkovich KA, Alex LA, Yarden O, Freitag M, Turner GE, Read ND, et al. Lessons from the
genome sequence of *Neurospora crassa*: tracing the path from genomic blueprint to
multicellular organism. Microbiol Mol Biol Rev. 2004;68: 1–108.

94.   Zhang Z, López-Giráldez F, Townsend JP. LOX: inferring Level Of eXpression from diverse
methods of census sequencing. Bioinformatics. 2010;26: 1918–1919.

95.   Arnold R, Rattei T, Tischler P, Truong M-D, Stümpflen V, Mewes W. SIMAP—The similarity
matrix of proteins. Bioinformatics. 2005;21: ii42–ii46.

96.   Rattei T, Arnold R, Tischler P, Lindner D, Stümpflen V, Mewes HW. SIMAP: the similarity
matrix of proteins. Nucleic Acids Res. 2006;34: D252–6.

97.   Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO. Assigning protein
functions by comparative genome analysis: protein phylogenetic profiles. Proc Natl Acad
Sci U S A. 1999;96: 4285–4288.

98.   Schmid KJ, Aquadro CF. The evolutionary analysis of "orphans" from the *Drosophila*
genome identifies rapidly diverging and incorrectly annotated genes. Genetics. 2001.
Available: https://www.genetics.org/content/159/2/589.short

99.   Benjamini Y, Bretz F, Sarkar SK. Recent Developments in Multiple Comparison
Procedures. IMS; 2004.

100. Benjamini Y, Heller R, Yekutieli D. Selective inference in complex research. Philos Trans A
Math Phys Eng Sci. 2009;367: 4255–4271.

101. Zhang Z, Townsend JP. Maximum-likelihood model averaging to profile clustering of site
types across discrete linear sequences. PLoS Comput Biol. 2009;5: e1000421.

102. H. Akaike. A new look at the statistical model identification. IEEE Trans Automat Contr.
1974;19: 716–723.

103. Hurvich CM, Tsai C-L. Regression and time series model selection in small samples.
Biometrika. 1989;76: 297–307.

104. Raftery AE, Madigan D, Hoeting JA. Bayesian Model Averaging for Linear Regression
Models. J Am Stat Assoc. 1997;92: 179–191.

105. Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S, Harb OS, et al. FungiDB: an
integrated functional genomics database for fungi. Nucleic Acids Res. 2012;40: D675–81.

106. Zhao Z-M, Campbell MC, Li N, Lee DSW, Zhang Z, Townsend JP. Detection of Regional
Variation in Selection Intensity within Protein-Coding Genes Using DNA Sequence
Polymorphism and Divergence. Mol Biol Evol. 2017;34: 3006–3022.

107. Sawyer SA, Hartl DL. Population genetics of polymorphism and divergence. Genetics.
1992;132: 1161–1176.

1116