The high diversity of *Scedosporium* and *Lomentospora* species and their prevalence in human-disturbed areas in Taiwan

Yin-Tse Huang[1], Yun-Chen Fan[2], Tsu-Chun Hung[1], Chi-Yu Chen[3], Pei-Lun Sun[2,4,*]

1. Department of Biomedical Science and Environment Biology, Kaohsiung Medical School, Kaohsiung, Taiwan.

2. Department of Dermatology, Chang Gung Memorial Hospital, Linkou, Taoyuan, Taiwan

3. Department of Plant Pathology, National Chun-Hsing University, Taichung, Taiwan

4. College of Medicine, Chang Gung University, Taoyuan, Taiwan

* Corresponding author

Key words: *Pseudallescheria*, emerging pathogen, soil fungi, fungal diversity

Correspondence:

Pei-Lun Sun, MD, PhD

Email: sunfungus@gmail.com

Department of Dermatology, Chang Gung Memorial Hospital, Linkou, Taoyuan, Taiwan

Abstract

Scedosporium and *Lomentospora* are important opportunistic pathogens causing localized or disseminated infection in humans. Understanding their environmental distribution is critical for public hygiene and clinical management. We carried out the first environmental survey in urbanized and natural regions in Taiwan. Overall, *Scedosporium* and *Lomentospora* species were recovered in 132 out of 267 soil samples (48.5%) across Taiwan. We morphologically and molecularly identified four *Scedosporium* species and one *Lomentospora* species. All four major clinical relevant species were isolated with high frequency, i.e. *S. apiospermum* (41.7%), *S. boydii* (28.8%), *L. prolificans* (12.3%), and *S. aurantiacum* (4.9%); a clinically minor species, *S. dehoogii*, had moderate incidence (12.3%). These fungal species have high incidence in urban (48%) and hospital (68%) soil samples, and have limited distribution in samples from natural regions (5%). Multivariate analysis of the fungal composition revealed strong evidence of the preferential distribution of these fungi in urban and hospital regions compared to natural sites. In addition, strong evidence suggested that the distribution and abundance of these fungal species are highly heterogeneous in the environment; samples in vicinity often yield varied fungal communities. Our results indicated that these fungal species are prevalent in Taiwan and their occurrences are associated with human activities. Hygiene sensitive places such as hospitals should be particularly aware of the high frequency of the clinical relevant species.

1. Introduction

Scedosporium and *Lomentospora* are two fungal genera that accommodate pathogenic species to immunocompromised and occasionally immunocompetent hosts. These fungal pathogens cause a broad range of clinical manifestation including colonization of respiratory tract, cutaneous infections, allergic reactions, and severe localized or disseminated mycoses. Mortality rates of infected hosts are generally high. In a meta-analysis of 264 infection causes, mortality rate was reported at an average of 22.2% in immunocompromised and of 14% in immunocompetent patients (1). In disseminated infections, mortality rate can be extreme as high as up to 80% (2). Treatments of *Scedosporium* and *Lomentospora* are challenging due to their low susceptibility to many current antifungal drugs such as 5-flucytosine, amphotericin B, and triazole drugs (2). In conjunction with their prevalence in the environment, *Scedosporium* and *Lomentospora* are imperative, emerging pathogens that call for research input in improving the diagnostic methods, developing effective therapies, and systematically surveying their distribution in the environment.

Scedosporium and *Lomentospora* are ubiquitous soil dwellers in temperate regions and are less frequently encountered in tropics (3, 4). Their distribution in the environment is often associated with nutrient-rich substrates such as agricultural and garden soils (3, 5), crude oil-contaminated soil (6), and manure of livestock, poultry or cattle (7–9). In addition, they can be isolated from low aerated and high osmotic substrates such as sewage sludge (10, 11), polluted pond bottoms (12), brackish water (13), coastal tidelands (14), and wastewater treatments (15, 16). Their prevalence in soil environments manifests in the cases of their occurrence in tissues of small wildlifes such as lizards (17), gerbils (18), donkey (19), and pigeons (20). In contrast, surveys of their occurrence in indoor water and air samples indicated that they are limited distributed in these samples (10, 12, 21, 22), while again, are

abundant in the soil of potted plants (21). Due to their prevalence in human active regions,

Scedosporium was proposed as an indicator species for anthropogenic impacts (23). Given

the pathogenic nature of these fungi, their occurrence in the environment are of concern,

especially in hospital and indoor settings, where they are posing potential hazard to people

with chronic respiratory disease and/or immunocompromised (24, 25).

Different *Scedosporium* and *Lomentospora* species pose different clinical relevances. Four

species, *S. boydii*, *S. apiospermum*, *S. aurantiacum* and *L. prolificans* are the most clinically

relevant species that account for most infections in reported cases (26, 27). With a few case

reports, *S. dehoogii* can cause subcutaneous and osteoarticular infections (28, 29). To

effectively distinguish clinical species from other clinically minor species, recent efforts have

made substantial taxonomic revisions and species delimitation of *Scedosporium* and

Lomentospora based on their morphological and molecular distinction (3, 30, 31). To date,

there are 13 distinct *Scedosporium* species: *S. americanum*, *S. apiospermum*, *S. aurantiacum*,

S. boydii, *S. cereisporum*, *S. dehoogii*, *S. desertorum*, *S. haikouense*, *S. hainanense*, *S.*

minutisporum, *S. multisporum*, *S. rarisporum*, *S. sanyaense*. Three species, *S. ellipsoideum*,

and *S. fusoideum*, *S. angustum*, are hypothesized cryptic species in the *S. boydii* species

complex but without provision of proper nomenclature requirements, and therefore invalid

(32). *Lomentospora* accommodates two species including *L. prolificans* and a recently

reported *L. valparaisensis* (33).

Research on their environmental distributions, compared to the clinical studies (e.g.

pathology, diagnosis, antifungal resistance), have been less emphasized. To date, there were a

few systematic surveys conducted in Australia (34), Europe (35, 36), South America (37, 38),

Africa (39), and Southeast Asia (40, 41). These studies have revealed a high variation of

species assemblies among samples, locations, and countries [see reviews such as Luplertlop

(2018) (42) and Rougeron et al (2018)]. This, coupled with a few non-targeted environmental

disclosures of these species in close proximity to human living habitats (5, 22, 43, 44), has

emphasized the importance of recognizing these potential infection sources in the

environment.

Studies of *Scedosporium* and *Lomentospora* species in Taiwan are limited. There were only a

few clinical case reports (45–52) and an environmental survey (25). This study, therefore,

aimed to survey the occurrence of these species in the soil environment across Taiwan. To

assess the risk factor of these species in varied facilities, we compared the species distribution

between natural and urbanized areas. We explicitly tested: (1) Do human-disturbed regions

accommodate more fungal species than in natural areas? (2) Do hygiene-sensitive areas (e.g.

hospital) harbor more fungal species compared to other urbanized regions?

98 2. Materials and Methods

99 2.1 Sampling sites

During 2014 to 2016, we collected soil samples from 79 locations across Taiwan (Table 1).

Sampling sites were selected to represent different levels of human activities (urban and

natural); in particular, we stressed the hospital samples as another sub-category in urban

samples as we wanted to highlight the distribution of these potential opportunistic fungi in the

hygiene sensitive places. In each site,1 – 14 sampling units were randomly chosen. For each

sampling unit, five points with 10 – 20 cm intervals from the center were randomly selected

to represent the fungal community of each unit. Soil was sampled with a metal spatula,

sterilized with 70% ethanol between samples to avoid cross contamination. Soil was sampled

at a depth of 7 – 10 cm beneath the soil surface, put in a sterile plastic bag and gently

squeezed to mix; the pooled soil sample was then filled in a 50 ml sterile Falcon tube. Soil

samples were stored at 4 °C until the fungal isolation.

2.2 Fungal isolation and fungal burden determination

For each soil sampling unit, ten grams of soil was air dried at room temperature for $1 - 2$

days, placed in a 100 ml flask and homogenized with 90 ml sterilized water with $1 - 2$ drops

of Tween 20. The mixture was thoroughly vortexed and left to stand for 15 minutes. 250 µl of

suspension of mixture were inoculated onto five plates of modified *Scedosporium* selective

medium (SceSel+) (53), comprised of malt extract, 6.25 g; maltose, 6.25 g; mono-potassium-

phosphate, 1.25 g; yeast extract, 1.0 g; magnesium sulfate heptahydrate, 0.625 g; soy

peptone, 0.625 g; chloramphenicol, 0.1 g; streptomycin sulfate, 0.1 g; dichloran, 2 mg;

benomyl, 6 mg; agar, 20.0 g; and reverse osmosis water, 990 ml. Plates were incubated at

37°C for $5 - 7$ days and inspected at intervals for the emergence of characteristic colonies of

Scedosporium. All *Scedosporium* colonies on five plates of each sampling unit were counted,

from which the fungal burdens were determined based on the mean value of CFU per gram of

soil dry weight.

2.3 DNA extraction, PCR, sequencing, and phylogeny reconstruction

Fungal tissues ($5 - 20$ mg) were lysed using mechanical beads (mix of 1 mm and 0.5 mm

beads at 1:1 ratio) beating at 1800 rpm for 4 min using a Retsch mixer mill MM400. DNA

extraction of the lysed cells were carried out using ZymoBIOMICS™ (CAT#D4306) with a

ZiXpress 32 automated nucleic acid purification instrument following the manufacturer's

instructions. Quantity and quality of eluted DNA were measured using Qubit dsDNA BR

Assay kit and Qubit Flex Fluorometer 4.0 (Thermo Fisher Scientific) following the

manufacturer's instructions.

For molecular identification, we employed the Multi-Locus Sequence Typing (MLST)

approach (54) and amplified four housekeeping gene loci including internal transcribed

spacer (ITS), β-tubulin (TUB), RNA polymerase II second largest subunit (RPB2), and

calmodulin (CAL). Primer pairs and their references used for PCR amplification are shown in

Table S1 (54–61). Amplification was performed with a two-step barcoding approach

according to Herbold et al. (62). The first-step PCR reaction mixture volume was 25 μl,

consisted of 1 μl of DNA template, 1 μl of each primer, 12.5 μl of 2X Tools Supergreen PCR

master mix (CAT#TTC-PA31-10), and 9.5 μl of ddH$_2$O. Amplification were carried out as

following steps: initial denaturation at 95 °C for 2 min, 25 cycles with denaturation at 95 °C

for 30 s, annealing at 64 – 66 °C for 30 s, primer extension at 72 °C for 1 min, and final

extension step was 7 min at 72 °C. The second-step PCR reaction mixture was 25 μl,

consisted of 1 μl of the first-step DNA product as template, 2.5 μL pre-mixed barcoded

primers, 10.75 μL of 2X Tools Supergreen PCR master mix, and 10.75 μl of ddH$_2$O. The

barcoded amplification products were purified using Ampure Xp Beads with 0.4X product to

bead volume ratio, both procedures followed the manufacturer's protocol. Sequencing was

carried out on a Oxford Nanopore Technologies GridION sequencer using the manufacturer's

library preparation kit.

We reconstructed the phylogeny of the isolated species using their concatenated ITS and β-

tubulin genes. We retried sequences of type specimens of *Scedosporium* and *Lomentospora*

species from NCBI. Sequences of *Petriellopsis africana* (CBS311.72), *Petriella setifera*

(CBS 385.87)*, Petriella sordida* (CBS 144612) were selected as the outgroup. We carried out

the phylogenetic analysis using an online phylogenetic service platform NGPhylogeny.fr

(63). Concatenated sequences were aligned using MAFFT with automatic flavor (64).

Phylogenetic informative regions were selected using Block Mapping and Gathering with

Entropy, a sliding window size of 3, and maximum entropy of 0.5 (65). Phylogenetic trees

were reconstructed using MrBayes (66) with recommended partition parameters. Two

MCMC runs of four chains were executed simultaneously from a random starting tree for

1000000 generations, every 500 generations were sampled resulting in 2000 trees, and 500

trees were discarded during burn-in. Posterior probabilities were estimated from the retained

1501 trees. Trees were visualized using Interactive Tree of Live version 4 (67).

162 2.4 Community analysis

We hypothesized that *Scedosporium* and *Lemontospora* species are more abundant in human-

ditrubed regions than in natural regions. To test this hypothesis, we conducted a distance-

based permutational multivariate analysis of variance (PERMANOVA) between the species

distance matrix of urban and natural samples. This test measures if groups of observations

differ in their composition. From our hypothesis, we expected to see a significant difference

of species composition between urban and natural samples. The test was implemented on a

Euclidean distance matrix using adonis() function of the vegan package for R (68). We then

carried out a PERMANOVA pairwise post hoc test using pairwise.perm.manova() on the

species distance matrix to test the differences of groups pairwisely (69). We further

conducted a homogeneity test on the species community matrix using betadisper() function.

Because the betadisper method is less sensitive to the variances among groups, it can help

justify the significant result from PERMANOVA that is a result of compositional difference

between samples, rather than within group variations. We presented Rsq and P-values and

discussed their inferences using the language of "evidence" following the recommendation in

Muff et al. (70)

To test for the effects of detecting *Scedosporium* and *Lomentospora* species in each location,

location types (i.e. urban, hospital, and natural), and each sampling unit, we conducted a

linear model-based analysis of the recovered species assemblage. We used the manyglm()

function of the mvabund package of R (71) to fit individual negative binomial linear models

for the species abundance of each detected fungal species. This approach provides an overall

multivariate test of the effect of location and location type on the composition of the

recovered fungal assemblage, as well as individual tests of the location, location types, and

sampling units on the abundance of each fungal species with adjusted P-value to account for

multiple comparisons. Three models were tested: (1) the effect of location type, (2) the

effects of location type, location, and their interaction, and (3) the effects of location,

sampling unit, and their interaction.

189 3. Results

190 3.1 Fungal isolation and fungal burden determination

164 *Scedosporium* and *Lomentospora* isolates were recovered from 267 sampling units in 79

locations, including 67 urban sites, six hospitals, and six natural sites across Taiwan (Table

1). The presence and abundance of these fungal species seems to be highly associated with

human activities. They were detected in 48% of urban samples, 68% of hospital samples, and

5% of natural samples (Fig. 1A). Likewise, the highest fungal burden was revealed in

hospital samples with average 20.8 CFU/g, followed by urban samples with 10.29 CFU/g; in

contrast, natural samples had only 0.11 CFU/g.

We identified five *Scedosporium* and *Lomentospora* species based on their morphological

and molecular characteristics, they were *S. apiospermum*, *S. boydii*, *S. aurantiacum*, *S.*

dehoogii, and *L. prolificans*. Of all 164 isolates, the *S. apiospermum* was the most frequently

recovered species (n = 68, 41.7%), followed by *S. boydii* (n = 47, 28.8%), *L. prolificans* (n =

21, 12.3%), *S. dehoogii* (n = 20, 12.3%), and *S. aurantiacum* (n = 8, 4.9%) (Fig. 1B). The

occurrence of these fungal species among locations were highly varied (Fig. 2 & 3). All five

species were detected in urban samples, ranging from one species in 14 locations, two species

in 11 locations, three species in 10 locations, four species in three locations, and all five

species in Tunghai University. All hospital samples revealed the occurrence of these fungal

species, ranging from one species in Linkou Chang Gung Memorial Hospital, two species in

Taichung Tri-service Hospital, three species in four hospitals (Taichung Veterans Hospital,

Taichung Hospital, Taichung Chung Shan Hospital, and China Medical University). In

contrast, *S. apiospermum* was the only species recovered from the natural samples from

Lotus Pond in Nantou.

Notably, the detected five fungal species exhibited some exceptionally high fungal burdens in

varied locations. For example, high fungal burden of *S. apiospermum* were detected in five

sampling locations including the British Consulate at Takow (365 CFU/g), Chishang Hakka

Cultural Park (250 CFU/g), Rushan Visitor Center (186 CFU/g), Meilun (112 CFU/g), and

Taichung Park (106 CFU/g). *Lomentospora prolificans* was high in three locations, they were

Taichung Veterans Hospital (325 CFU/g), Taichung Park (313 CFU/g), and Taichung

Hospital (149 CFU/g). *Scedosporium boydii* was high in Chishang Hakka Cultural Park (250

CFU/g). *Scedosporium aurantiacum* was abundant in Rushan Visitor Center (183 CFU/g),

and *Scedosporium dehoogii* showed it abundance in the British Consulate at Takow (460

CFU/g) (Figure 3).

222 3.2 Fungal community distribution analysis

Our analysis indicated there was an evident variation in the species composition of

Scedosporium and *Lomentospora* within and between locations. Overall, there was strong

evidence for the species composition in different location types (i.e. urban, hospital, and

natural samples), while it accounted for a limited portion of the explained variation (adonis:

df=2, F=5.3980, R^2=0.0375, P=0.001). While explaining much of the compositional

difference, there was no evidence showing the species composition varied among locations

(adonis: df=75, F=1.0991, R^2=0.2870, P=0.245)(Table 2). In a pairwise PERMANOVA

analysis, species composition in natural samples was distinct from that in urban and hospital

samples, while no difference was detected between the urban and hospital samples (Table 3).

The homogeneity test further indicated that species assemblage between natural-urban and

natural-hospital samples were significantly different in their homogeneity; instead, no

dispersion effect was found between urban and hospital samples (Figure 4) (betadisper: df=2,

F=9.8585, P=7.3809e-05). The dissimilarity of species composition was also indicated by the

clustering of natural samples, which were separated from the urban and hospital samples, on

the NMDS plot (Figure 5). Altogether, these results suggested that *Scedosporium* and

Lomentospora had different composition profiles in natural regions compared to human-

dominated regions, i.e. urban and hospital samples.

General linear model-based analysis indicated a evident effect of location types (i.e. urban,

hospital, and natural samples) on the multivariate composition of the recovered *Scedosporium*

and *Lomentospora* assemblages ($P = 0.001$) (Table 4, model 1). In addition, the univariate

test indicated that *S. apiospermum* ($P = 0.006$), *S. boydii* ($P = 0.007$), *L. prolificans* ($P =$

0.007), and *S. dehoogii* ($P = 0.006$) showed moderate-strong evidence of its preferred

distribution in urban regions (including urban and hospital site) (Table 5, model 1). Similar to

the result of PERMANOVA analysis, we found no evidence that the recovered fungal

composition was explained by location ($P = 0.582$) and its interaction term with location type

($P = 0.798$) (Table 4, model 2). In the further test using location and sampling unit as factors

(model 3), we found moderate-strong evidence of the location effect ($P = 0.022$) and strong

evidence of the sampling unit effect ($P = 0.001$) on the recovered fungal composition.

Furthermore, there was strong evidence indicating that *S. apiospermum* is preferentially

recovered from certain sampling units ($P = 0.001$), and moderate-strong evidence for *S.*

boydii (P = 0.032)(Table 5, model 3).

4. Discussion

Scedosporium and *Lomentospora* species were prevalent in soil samples in Taiwan. Their

occurrence in the environment were mostly in human disturbed areas, i.e. in urban and

hospital samples in the present study. In contrast, natural regions had limited distribution of

these fungal species. While detecting *Scedosporium* and *Lomentospora* species in hospital

regions in the present study, our results indicated that these hygiene-sensitive areas, in

general, possessed no more fungal burden than nearby urban regions; lessening the need of

taking precautionary approach at this point.

Scedosporium and *Lomentospora* are prevalent and are heterogeneous distributed in the environment

Overall, half of soil samples in the present study yield *Scedosporium* and *Lomentospora*

species. Their distribution was not only on the main island, but also on the outlying islands

such as Penghu and Kinmen. Although with less density of human population on these

islands, we recovered a comparable amount of *Scedosporium* and *Lomentospora* species as

that on the main island. Whether the fungal community of these species on the outlying island

is endemic or is brought via human activities, and if so the disseminating route, await further

investigations.

Distribution of *Scedosporium* and *Lomentospora* species in the environment appear to be

highly heterogeneous, even sampling sites in vicinity often yielded varied compositions of

these fungal species, as suggested in the homogeneity test (Fig. 4). Notably, we did not

observe apparent patterns explaining the high fungal burden of these species in certain

sampling sites. We found, for example, *S. apiospermum* and *L. prolificans* are abundant in

sampling sites in parks where birds are fed and are routinely roosting in; these are places

where they may accumulate high nitrogen contents that can encourage the population of

Scedosporium and *Lomentospora* species (36, 39). Other than this putative factor, we found

in some tourist visiting sites (e.g. British Consulate at Takow, Chishang Hakka Cultural Park,

and Rushan Visitor Center) and hospitals (e.g. Taichung Veterans Hospital and Taichung

Hospital) where they accommodate exceptionally high fungal burden of these species (Fig.

3). While again, the distribution of *Scedosporium* and *Lomentospora* species are highly

variable in sites of these high fungal burden places. The heterogeneous distribution of these

fungal species in the environment invites analysis of the physical and chemical properties of

soil samples in the future studies.

As in many previous studies, *S. apiospermum* is the most common species in the environment

(34, 36, 38–41). The prevalence of this species in the environment has been repeatedly

revealed in varied countries across continents and climatic zones, and by using varied

selective culturing methods, i.e. dichloran rose bengal chloramphenicol (DRBC) media,

Scedosporium-selective media (SceSel), and Scedo-Select III culture medium (36, 39, 72).

All together, it suggests that *S. apiospermum* appears to be the dominant species in the soil

environment globally regardless of geographical and climatic regions. This, in conjunction

with their prevalence in human disturbed regions (e.g. sewage, wastewater treatment plants,

and landfills), suggests their potential as the indicator species for anthropogenic disturbance

or pollution in areas worldwide (23).

Scedosporium and *Lomentospora* have limited distribution in natural regions

In the present study, we isolated *S. apiospermum* from a natural region. As opposed to previous studies, in which no *Scedosporium* and *Lomentospora* species were isolated from the natural/pristine areas (35, 36). The only three cases of recovering *Scedosporium* from natural regions were samples from Argane forests in Souss-Massa, forests in Maamoura, and beaches in Casablanca, all in Morocco (39, 73, 74). These places, however, appeared to exhibit anthropogenic activities, especially in the highly visited Casablanca beach; it may explain the presence of *Scedosporium* species in these regions. Similar to the case in Morocco, the detected sample in the natural region in Taiwan was from Lotus Pond, where is a recreation place with moderate tourist vistings, the low recovery rate and abundance (16 CFU/g) of *Scedosporium* species was therefore plausible. The other five natural places, Cui-Feng, Daxueshan, Baimaoshan, Kuantaoshan, and Liu-Long-Tou, are relatively high-elevated and with lesser tourist visitings; we therefore did not find the fungi in the present study. Intriguingly, few samples from a recreation area (Moon World) with the Badlands landscape exhibit the distribution of *S. apiospermum* (16 CFU/g) and *L. prolificans* (8 CFU/g). Though with low fungal burden for both species, their presence in such an arid region suggest the tolerance of *Scedosporium* and *Lomemtospora* species to many adverse environments.

Surveys in different countries revealed varied community profiles of *Scedosporium* and *Lomentospora*

The four clinical relevant species, i.e. *S. apiospermum* (41.7%), *S. boydii* (28.8%), *L. prolificans* (12.3%), and *S. aurantiacum* (4.9%), accounted for 87.7% of the recovered fungal species in the present study; the clinically minor species, *S. dehogii*, accounted for the rest

12.3%. This recovered fungal community profile in Taiwan is different from that in different

countries. For example, an environmental survey in Morocco revealed a similar fungal profile

of *S. apiospermum* (56%), *S. aurantiacum* (17%), *S. boydii* (18%), and *S. dehogii* (9%) (39);

in Thailand revealed a profile of *S. apiospermum* (87%), *S. aurantiacum* (7%), *S. dehogii*

(6%) (40); in France revealed a profile of *S. dehogii* (39.4%), *S. aurantiacum* (21.6%), *S.*

boydii (19.7%), and *S. apiospermum* (18.9%) (35); in Australia revealed a profile of *S.*

aurantiacum (54.6%), *L. prolificans* (43%), *S. boydii* (2.1%), *S. dehogii* (0.3%), and *S.*

minutispora (0.004%) (34); and in Austria, a profile of *S. apiospermum* (69.8%), *S. dehoogii*

(19.8%), *S. aurantiacum* (6.3%), *S. boydii* (2.2%), and *S. minutispora* (1.9%) was revealed

(36).

Intriguingly, the disclosed fungal profile of *Scedosporium* and *Lomentospora* in countries

appear to echo the clinical cases attributed to certain species. For example, patients with

chronic lung disease in Australia were mostly colonized by *S. aurantiacum (34)*. Clinical

cases in Austria were mostly attributed to *S. apiospermum* (36). Infections in Taiwan were

diagnosed with *S. apiospermum* and *S. boydii* as the culprits (Sun et al. unpublished). While

discrepancy happens such as the case in France where *S. boydii* is the major cause of

infections but the dominant species was *S. dehoogii* in the environmental study. The exact

link between environmental and clinical strains warrants future studies using a more rigorous

experimental design coupled with approaches such as molecular population genetics.

5. References

1. Seidel D, Meißner A, Lackner M, Piepenbrock E, Salmanton-García J, Stecher M, Mellinghoff

S, Hamprecht A, Durán Graeff L, Köhler P, Cheng MP, Denis J, Chedotal I, Chander J, Pakstis

DL, Los-Arcos I, Slavin M, Montagna MT, Caggiano G, Mares M, Trauth J, Aurbach U,

Vehreschild MJGT, Vehreschild JJ, Duarte RF, Herbrecht R, Wisplinghoff H, Cornely OA.

2019. Prognostic factors in 264 adults with invasive Scedosporium spp. and Lomentospora

prolifícans infection reported in the literature and FungiScope®. Crit Rev Microbiol 45:1–21.

2.   Ramirez-Garcia A, Pellon A, Rementeria A, Buldain I, Barreto-Bergter E, Rollin-Pinheiro R, de

Meirelles JV, Xisto MIDS, Ranque S, Havlicek V, Vandeputte P, Govic YL, Bouchara J-P,

Giraud S, Chen S, Rainer J, Alastruey-Izquierdo A, Martin-Gomez MT, López-Soria LM, Peman

J, Schwarz C, Bernhardt A, Tintelnot K, Capilla J, Martin-Vicente A, Cano-Lira J, Nagl M,

Lackner M, Irinyi L, Meyer W, de Hoog S, Hernando FL. 2018. Scedosporium and

Lomentospora: an updated overview of underrated opportunists. Med Mycol 56:102–125.

3.   Guarro J, Kantarcioglu AS, Horré R, Rodriguez-Tudela JL, Cuenca Estrella M, Berenguer J, de

Hoog GS. 2006. Scedosporium apiospermum: changing clinical spectrum of a therapy-refractory

opportunist. Med Mycol 44:295–327.

4.   Rougeron A, Giraud S, Alastruey-Izquierdo A, Cano-Lira J, Rainer J, Mouhajir A, Le Gal S,

Nevez G, Meyer W, Bouchara JP. 2018. Ecology of Scedosporium Species: Present Knowledge

and Future Research. Mycopathologia 183:185–200.

5.   Babu AG, Kim SW, Yadhav DR, Adhikari M, Kim C, Lee HB, Lee YS. 2014. A New Record of

Pseudallescheria boydii Isolated from Crop Field Soil in Korea. Mycobiology 42:397–400.

6.   García-Peña EI, Hernández S, Favela-Torres E, Auria R, Revah S. 2001. Toluene biofiltration by

the fungus Scedosporium apiospermum TB1. Biotechnol Bioeng 76:61–69.

7.   Bell RG. 1976. The development in beef cattle manure of Petriedllidium boydii (Shear) Malloch,

a potential pathogen for man and cattle. Can J Microbiol 22:552–556.

8.   Alteras I, Evolceanu R. 1969. First isolation of Allescheria Boydii from romanian soil.

Sabouraudia 7:135–137.

9.   Bell RG. 1978. Comparative virulence and immunodiffusion analysis of Petriellidium boydii

(Shear) Malloch strains isolated from feedlot manure and a human mycetoma. Can J Microbiol

24:856–863.

10. Cooke WB, Kabler P. 1955. Isolation of potentially pathogenic fungi from polluted water and

sewage. Public Health Rep 70:689–694.

11. Llanos C, Kjøller A, Kjoller A. 1976. Changes in the flora of soil fungi following oil waste

application. Oikos 27:377.

12. Fisher JF, Shadomy S, Teabeaut JR, Woodward J, Michaels GE, Newman MA, White E, Cook

P, Seagraves A, Yaghmai F, Rissing JP. 1982. Near-drowning complicated by brain abscess due

to Petriellidium boydii. Arch Neurol 39:511–513.

13. Kirk PW. 1967. A comparison of saline tolerance and sporulation in marine and clinical isolates

of Allescheria boydii shear. Mycopathol Mycol Appl 33:65–75.

14. Dabrowa N, Landau JW, Newcomer VD, Plunkett OA. 1964. A SURVEY OF TIDE-WASHED

COASTAL AREAS OF SOUTHERN CALIFORNIA FOR FUNGI POTENTIALLY

PATHOGENIC TO MAN. Mycopathol Mycol Appl 24:136–150.

15. Janda-Ulfig K, Ulfig K, Cano J, Guarro J. 2008. A study of the growth of Pseudallescheria

boydii isolates from sewage sludge and clinical sources on tributyrin, rapeseed oil, biodiesel oil

and diesel oil. Ann Agric Environ Med 15:45–49.

16. Crous PW, Wingfield MJ, Burgess TI, Hardy GESJ, Crane C, Barrett S, Cano-Lira JF, Le Roux

JJ, Thangavel R, Guarro J, Stchigel AM, Martín MP, Alfredo DS, Barber PA, Barreto RW,

Baseia IG, Cano-Canals J, Cheewangkoon R, Ferreira RJ, Gené J, Lechat C, Moreno G, Roets F,

Shivas RG, Sousa JO, Tan YP, Wiederhold NP, Abell SE, Accioly T, Albizu JL, Alves JL,

Antoniolli ZI, Aplin N, Araújo J, Arzanlou M, Bezerra JDP, Bouchara J-P, Carlavilla JR,

Castillo A, Castroagudín VL, Ceresini PC, Claridge GF, Coelho G, Coimbra VRM, Costa LA, da

Cunha KC, da Silva SS, Daniel R, de Beer ZW, Dueñas M, Edwards J, Enwistle P, Fiuza PO,

Fournier J, García D, Gibertoni TB, Giraud S, Guevara-Suarez M, Gusmão LFP, Haituk S,

Heykoop M, Hirooka Y, Hofmann TA, Houbraken J, Hughes DP, Kautmanová I, Koppel O,

Koukol O, Larsson E, Latha KPD, Lee DH, Lisboa DO, Lisboa WS, López-Villalba Á, Maciel

JLN, Manimohan P, Manjón JL, Marincowitz S, Marney TS, Meijer M, Miller AN, Olariaga I,

Paiva LM, Piepenbring M, Poveda-Molero JC, Raj KNA, Raja HA, Rougeron A, Salcedo I,

Samadi R, Santos TAB, Scarlett K, Seifert KA, Shuttleworth LA, Silva GA, Silva M, Siqueira

JPZ, Souza-Motta CM, Stephenson SL, Sutton DA, Tamakeaw N, Telleria MT, Valenzuela-

Lopez N, Viljoen A, Visagie CM, Vizzini A, Wartchow F, Wingfield BD, Yurchenko E, Zamora

JC, Groenewald JZ. 2016. Fungal Planet description sheets: 469-557. Persoonia 37:218–403.

17. Gugnani HC, Okafor JI. 2009. Mycotic Flora of the Intestine and other Internal Organs of

Certain Reptiles and Amphibians with Special Reference to Characterization of Basidiobolus

Isolates: Die Pilzflora des Darmes und anderer innerer Organe von bestimmten Reptilien und

Amphibien unt. Mycoses 23:260–268.

18. Gugnani HC. 1972. Fungi isolated from lungs of small wild animals in India. Mykosen 15:479–

482.

19. El Allawy, Atia, Amer. 1977. Mycoflora of the pharyngeo-tonsillar portion of clinically healthy

donkeys [asses] in Assiut [Egypt]. Assiut Vet Med J.

20. Ramirez R, Robertstad GW, Hutchinson LR, Chavez J. 1976. Mycotic flora in the lower

digestive tract of feral pigeons (Columba livia) in the El Paso, Texas area. J Wildl Dis 12:83–85.

21. Sidot C, Simon P, Bouchara JP, Chabasse D, Urban T, Giniès JL. 2007. 116 Scedosporium

apiospermum. Environmental study in the homes of patients with cystic fibrosis. J Cyst Fibros

6:S29.

22. Aho R, Hirn J. 1981. A survey of fungi and some indicator bacteria in chlorinated water of

indoor public swimming pools. Zentralbl Bakteriol Mikrobiol Hyg B 173:242–249.

23. Al-Yasiri MH, Normand A-C, Mauffrey J-F, Ranque S. 2017. Anthropogenic impact on

environmental filamentous fungi communities along the Mediterranean littoral. Mycoses

60:477–484.

24. Summerbell RC, Krajden S, Kane J. 1989. Potted plants in hospitals as reservoirs of pathogenic

fungi. Mycopathologia 106:13–22.

25. Chen C-H, Lin L-C, Chang Y-J, Liu C-E, Soon M-S, Huang C-S. 2013. Efficacy of copper-silver

ionization for controlling fungal colonization in water distribution systems. J Water Health

11:277–280.

26. Bronnimann D, Garcia-Hermoso D, Dromer F, Lanternier F, French Mycoses Study Group,

Characterization of the isolates at the NRCMA. 2021. Scedosporiosis/lomentosporiosis

observational study (SOS): Clinical significance of Scedosporium species identification. Med

Mycol 59:486–497.

27. Heath CH, Slavin MA, Sorrell TC, Handke R, Harun A, Phillips M, Nguyen Q, Delhaes L, Ellis

D, Meyer W, Chen SCA, Australian Scedosporium Study Group. 2009. Population-based

surveillance for scedosporiosis in Australia: epidemiology, disease manifestations and

emergence of Scedosporium aurantiacum infection. Clin Microbiol Infect 15:689–693.

28. Krauth DS, Barlow BT, Berjohn CM. 2021. Fungal osteomyelitis and septic arthritis in an

immune competent man: The first report of invasive osteoarticular infection due to

Scedosporium dehoogii. Med Mycol Case Rep 33:14–17.

29. Li F-G, Yang Y-P, Li W, Sheng P, Li W, Huang W-M, Fan Y-M. 2017. Spontaneous Remission

of Subcutaneous Scedosporiosis Caused by Scedosporium dehoogii in a Psoriatic Patient.

Mycopathologia 182:561–567.

30. Lackner M, de Hoog GS, Yang L, Ferreira Moreno L, Ahmed SA, Andreas F, Kaltseis J, Nagl

M, Lass-Flörl C, Risslegger B, Rambach G, Speth C, Robert V, Buzina W, Chen S, Bouchara J-

P, Cano-Lira JF, Guarro J, Gené J, Fernández Silva F, Haido R, Haase G, Havlicek V, Garcia-

Hermoso D, Meis JF, Hagen F, Kirchmair M, Rainer J, Schwabenbauer K, Zoderer M, Meyer W,

Gilgado F, Schwabenbauer K, Vicente VA, Piecková E, Regenermel M, Rath P-M, Steinmann J,

de Alencar XW, Symoens F, Tintelnot K, Ulfig K, Velegraki A, Tortorano AM, Giraud S, Mina

S, Rigler-Hohenwarter K, Hernando FL, Ramirez-Garcia A, Pellon A, Kaur J, Bergter EB, de

Meirelles JV, da Silva ID, Delhaes L, Alastruey-Izquerdo A, Li R-Y, Lu Q, Moussa T,

Almaghrabi O, Al-Zahrani H, Okada G, Deng S, Liao W, Zeng J, Issakainen J, Liporagi Lopes

LC. 2014. Proposed nomenclature for Pseudallescheria, Scedosporium and related genera.

Fungal Divers 67:1–10.

31. Gilgado F, Cano J, Gené J, Sutton DA, Guarro J. 2008. Molecular and phenotypic data

supporting distinct species statuses for Scedosporium apiospermum and Pseudallescheria boydii

and the proposed new species Scedosporium dehoogii. J Clin Microbiol 46:766–771.

32. Cortez KJ, Roilides E, Quiroz-Telles F, Meletiadis J, Antachopoulos C, Knudsen T, Buchanan

W, Milanovich J, Sutton DA, Fothergill A, Rinaldi MG, Shea YR, Zaoutis T, Kottilil S, Walsh

TJ. 2008. Infections caused by Scedosporium spp. Clin Microbiol Rev 21:157–197.

33. Crous PW, Wingfield MJ, Burgess TI, Hardy GESJ, Gen'e J, Guarro J, Baseia IG, Garc'ıa D,

Gusmão LFP, Souza-Motta CM, Thangavel R, Adam'ık S, Barili A, Barnes CW, Bezerra JDP,

Bordallo JJ, Cano-Lira JF, de Oliveira RJV, Ercole E, Hubka V, Iturrieta-Gonz'alez I,

Kub'atov'a A, Mart'ın MP, Moreau P-A, Morte A, Ordoñez ME, Rodr'ıguez A, Stchigel AM,

Vizzini A, Abdollahzadeh J, Abreu VP, Adam'ıkov'a K, Albuquerque GMR, Alexandrova

AV, 'Alvarez Duarte E, Armstrong-Cho C, Banniza S, Barbosa RN, Bellanger J-M, Bezerra JL,

Cabral TS, Cabo M, Caicedo E, Cantillo T, Carnegie AJ, Carmo LT, Castañeda-Ruiz RF,

Clement CR, mokov'a A, Conceição LB, Cruz RHSF, Damm U, da Silva BDB, da Silva GA, da

Silva RMF, Santiago ALCM de A, de Oliveira LF, de Souza CAF, D'eniel F, Dima B, Dong G,

Edwards J, F'elix CR, Fournier J, Gibertoni TB, Hosaka K, Iturriaga T, Jadan M, Jany J-L,

Jurjevi, Kola'ık M, Kuan I, Landell MF, Leite Cordeiro TR, Lima DX, Loizides M, Luo S,

Machado AR, Madrid H, Magalhães OMC, Marinho P, Matoec N, Mei A, Miller AN, Morozova

OV, Neves RP, Nonaka K, Nov'akov'a A, Oberlies NH, Oliveira-Filho JRC, Oliveira TGL, Papp

V, Pereira OL, Perrone G, Peterson SW, Pham THG, Raja HA, Raudabaugh DB, Iehulka J,

Rodr\-guez-Andrade E, Saba M, SchauflerovÃ!' A, Shivas RG, Simonini G, Siqueira JPZ,

Sousa JO, Stajsic V, Svetasheva T, Tan YP, Tkalec Z, Ullah S, Valente P, Valenzuela-Lopez N,

Abrinbana M, Viana Marques DA, Wong PTW, Xavier de Lima V, Groenewald JZ. 2018.

Fungal Planet description sheets: 716784. Persoonia - Molecular Phylogeny and Evolution of

Fungi 40:239–392.

34. Harun A, Gilgado F, Chen SC-A, Meyer W. 2010. Abundance of Pseudallescheria/Scedosporium

species in the Australian urban environment suggests a possible source for scedosporiosis

including the colonization of airways in cystic fibrosis. Med Mycol 48 Suppl 1:S70–6.

35. Rougeron A, Schuliar G, Leto J, Sitterlé E, Landry D, Bougnoux M-E, Kobi A, Bouchara J-P,

Giraud S. 2015. Human-impacted areas of France are environmental reservoirs of the

Pseudallescheria boydii/Scedosporium apiospermum species complex. Environ Microbiol

17:1039–1048.

36. Kaltseis J, Rainer J, De Hoog GS. 2009. Ecology of Pseudallescheria and Scedosporium species

in human-dominated and natural environments and their distribution in clinical samples. Med

Mycol 47:398–405.

37. Alvarez E, Sanhueza C. 2016. New record of Scedosporium dehoogii from Chile: Phylogeny and

susceptibility profiles to classic and novel putative antifungal agents. Rev Iberoam Micol

33:224–229.

38. Elizondo-Zertuche M, de J Treviño-Rangel R, Robledo-Leal E, Luna-Rodríguez CE, Martínez-

Fierro ML, Rodríguez-Sánchez IP, González GM. 2017. Molecular identification and in vitro

antifungal susceptibility of Scedosporium complex isolates from high-human-activity sites in

Mexico. Mycologia 109:874–881.

39. Mouhajir A, Poirier W, Angebault C, Rahal E, Bouabid R, Bougnoux M-E, Kobi A, Zouhair R,

Bouchara J-P, Giraud S. 2020. Scedosporium species in soils from various biomes in

Northwestern Morocco. PLoS One 15:e0228897.

40. Luplertlop N, Pumeesat P, Muangkaew W, Wongsuk T, Alastruey-Izquierdo A. 2016.

Environmental Screening for the Scedosporium apiospermum Species Complex in Public Parks

in Bangkok, Thailand. PLoS One 11:e0159869.

41. Luplertlop N, Muangkaew W, Pumeesat P, Suwanmanee S, Singkum P. 2019. Distribution of

Scedosporium species in soil from areas with high human population density and tourist

popularity in six geographic regions in Thailand. PLoS One 14:e0210942.

42. Luplertlop N. 2018. Pseudallescheria/Scedosporium complex species: From saprobic to

pathogenic fungus. J Mycol Med 28:249–256.

43. Zhang Z-Y, Shao Q-Y, Li X, Chen W-H, Liang J-D, Han Y-F, Huang J-Z, Liang Z-Q. 2021.

Culturable Fungi from Urban Soils in China I: Description of 10 New Taxa. Microbiol Spectr

9:e0086721.

44. Han, Zheng, Zhang, Chen, Luo, Wang, Liang, Others. 2017. Two new Scedosporium species

from Guangxi and Hainan. Mycosystem 36:145–153.

45. Hsiue H-C, Ruan S-Y, Kuo Y-L, Huang Y-T, Hsueh P-R. 2010. Invasive infections caused by

non-Aspergillus moulds identified by sequencing analysis at a tertiary care hospital in Taiwan,

2000-2008. Clin Microbiol Infect 16:1204–1206.

46. Lin D, Kamili Q, Lai S, Musher DM, Hamill R. 2013. Cerebral Scedosporium apiospermum

infection presenting with intestinal manifestations. Infection 41:723–726.

47. Lin S-Y, Lee K-M, Chen T-C, Chen Y-H, Lien C-T, Lu P-L. 2013. A Swollen Knee in a 77-

Year-Old Lung Cancer Patient Receiving Antimicrobial Therapy for Pneumonia. Clin Infect Dis

56:1838–1839.

48. Chen T-C, Ho M-W, Chien W-C, Lin H-H. 2016. Disseminated Scedosporium apiospermum

infection in a near-drowning patient. J Formos Med Assoc 115:213–214.

49. Chiang C-H, Hsu C-K, Lee JY-Y, Chang TC, Hsueh Y-Y, Shieh S-J, Chen H-M, Hsu MM-L.

2014. Recurrent Scedosporium apiospermum mycetoma successfully treated by surgical excision

and voriconazole. Dermatologica Sinica 32:29–32.

50. Ko W-H, Tsou Y-J, Ju Y-M, Hsieh H-M, Ann P-J. 2010. Production of a fungistatic substance

by Pseudallescheria boydii isolated from soil amended with vegetable tissues and its

significance. Mycopathologia 169:125–131.

51. Hung CC, Chang SC, Yang PC, Hsieh WC. 1994. Invasive pulmonary pseudallescheriasis with

direct invasion of the thoracic spine in an immunocompetent patient. Eur J Clin Microbiol Infect

Dis 13:749–751.

52. Tan T-Y, Liou C-W, Kung L-C. 2004. Meningitis caused by Pseudallescheria boydii. Chang

Gung Med J 27:228–232.

53. Rainer J, Kaltseis J, de Hoog SG, Summerbell RC. 2008. Efficacy of a selective isolation

procedure for members of the Pseudallescheria boydii complex. Antonie Van Leeuwenhoek

93:315–322.

54. Bernhardt A, Sedlacek L, Wagner S, Schwarz C, Würstl B, Tintelnot K. 2013. Multilocus

sequence typing of Scedosporium apiospermum and Pseudallescheria boydii isolates from cystic

fibrosis patients. J Cyst Fibros 12:592–598.

55. Tedersoo L, Tooming-Klunderud A, Anslan S. 2018. PacBio metabarcoding of Fungi and other

eukaryotes: errors, biases and perspectives. New Phytol 217:1370–1385.

56. White TJ, Bruns T, Lee S, Taylor JW. 1990. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds.

PCR protocols: A guide to the methods and applicationsNew York, NY: Academic Press.

57. Rehner S. 2001. Primers for Elongation Factor 1-α (EF1-α).

http://ocid.nacse.org/research/deephyphae/EF1primer.pdf.

58. Glass NL, Donaldson GC. 1995. Development of primer sets designed for use with the PCR to

amplify conserved genes from filamentous ascomycetes. Appl Environ Microbiol 61:1323–1330.

59. O'Donnell K, Cigelnik E. 1997. Two divergent intragenomic rDNA ITS2 types within a

monophyletic lineage of the fungus *Fusarium* are nonorthologous. Mol Phylogenet Evol 7:103–

116.

60. Liu YJ, Whelen S, Hall BD. 1999. Phylogenetic relationships among ascomycetes: evidence

from an RNA polymerse II subunit. Mol Biol Evol 16:1799–1808.

61. Peterson SW. 2008. Phylogenetic analysis of Aspergillus species using DNA sequences from

four loci. Mycologia 100:205–226.

62. Herbold CW, Pelikan C, Kuzyk O, Hausmann B, Angel R, Berry D, Loy A. 2015. A flexible and

economical barcoding approach for highly multiplexed amplicon sequencing of diverse target

genes. Front Microbiol 6:731.

63. Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S, Gascuel O.

2019. NGPhylogeny.fr: new generation phylogenetic services for non-specialists. Nucleic Acids

Res 47:W260–W265.

64. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol 30:772–780.

65. Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a new

software for selection of phylogenetic informative regions from multiple sequence alignments.

BMC Evol Biol 10:210.

66. Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics 17:754–755.

67. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Res 47:W256–W259.

68.   Oksanen J, Kindt R, Legendre P, O'Hara B. 2007. The vegan package. ecology package 10:631–

637.

69.   Martinez Arbizu. 2017. pairwiseAdonis: Pairwise multilevel comparison using adonis. R

package version 0 0 1.

70.   Muff S, Nilsen EB, O'Hara RB, Nater CR. 2022. Rewriting results sections in the language of

evidence. Trends Ecol Evol 37:203–210.

71.   Wang Y, Naumann U, Wright ST, Warton DI. 2012. mvabund - an R package for model-based

analysis of multivariate abundance data. Methods Ecol Evol 3:471–474.

72.   Pham T, Giraud S, Schuliar G, Rougeron A, Bouchara J-P. 2015. Scedo-Select III: a new semi-

selective culture medium for detection of the Scedosporium apiospermum species complex. Med

Mycol 53:512–519.

73.   Hallouti, Zahidi, Bouharroud. 2017. Diversity of entomopathogenic fungi in Argane forest soil

and their potential to manage Mediterranean fruit fly (Ceratitis capitata). J Pharmacopuncture.

74.   Soussi Abdallaoui M, Boutayeb H, Guessous-Idrissi N. 2007. Flore fongique du sable de deux

plages à Casablanca (Maroc): Analyse et corollaires épidémiologiques. J Mycol Med 17:58–62.

75.   Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics

and parallel computing. Nat Methods 9:772.

76.   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics 30:1312–1313.

77.   Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,

Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and

model choice across a large model space. Syst Biol 61:539–542.

78.   Stöver BC, Müller KF. 2010. TreeGraph 2: combining and visualizing evidence from different

phylogenetic analyses. BMC Bioinformatics 11:7.

Legends

Figure 1. Culture-dependent isolation from soil samples in the present study. (A) Percentage of culture-positive samples of *Scedosporium* and *Lomemtospora* species in different types of sampling sources (urban, hospital, and natural) across Taiwan. (B) Proportion of each isolated *Scedosporium*/*Lomemtospora* species in the 164 isolates.

Figure 2. Mean CFU of *Scedosporium*/*Lomemtospora* species from samples in each location across Taiwan. Names of locations with mean CFU more than 40 are labeled. Circle sizes represent the sampling size in each location. * measured values of CFU/g.

Figure 3. Heatmap showing the mean CFU of each *Scedosporium*/*Lomemtospora* species in each location sampled in the present study; types of sampling sources (urban, hospital, and natural) were labeled next the location name on x-axis.

Figure 4. Homogeneity test on the species assemblage in different types of sampling sources (urban, hospital, and natural); the p-value generated via pairwise comparisons using post hoc t-test were shown above the bars.

Figure 5. NMDS ordination showing the dissimilarities of species assemblage in soil samples from each location.

Table 1. Sampling locations, assigned categories (urban, hospital, and hospital) of each sampling location, geographic coordinate, and sample sizes of each location.

Table 2. Degree of freedom, F-values, determination coefficients, and P-values of tested factors (location type and location) that influence the species assemblage observed in each location.

Table 3. F-values, determination coefficients, and P-values of pairwise test of species

assemblage observed in urban, hospital, and natural regions.

Table 4. Three models using the general linear-model-based analysis tested the occurrence

differences of *Scedosporium* and *Lomentospora* species explained by the effect of location

type (model 1); location type, location and the interaction term (model 2); location, sampling

unit, and the interaction term (model 3). Degree of freedom, deviance, and alpha- corrected

P-values of multivariate tests were shown.

Table 5. Three models using the general linear-model-based analysis tested the occurrence

differences of *Scedosporium* and *Lomentospora* species explained by the effect of location

type (model 1); location type, location and the interaction term (model 2); location, sampling

unit, and the interaction term (model 3). The alpha- corrected P-values of univariate tests

were shown.

Table S1. Primer names and their sequences used in the present study.

A

Culture positive samples

B

Species occurrence frequency

4.9%

41.7%

12.3%

12.3%

28.8%

Species

S. apiospermum

S. boydii

L. prolificans

S. dehoogii

S. aurantiacum

Scedosporium occurrence in locations