Whole blood transcriptome profiles of trypanotolerant and trypanosusceptible cattle
highlight a differential modulation of metabolism and immune response during infection
by *Trypanosoma congolense*

Moana Peylhard [1,2], David Berthier [1,2], Guiguigbaza-Kossigan Dayo [3], Isabelle Chantal [1,2],
Souleymane Sylla [3], Sabine Nidelet [4,5], Emeric Dubois [4,5], Guillaume Martin [6,7], Guilhem
Sempéré [1,2], Laurence Flori [8], Sophie Thévenon [1,2]
7
[1]INTERTRYP, Univ. Montpellier, CIRAD, IRD, Montpellier, France
[2]CIRAD, UMR INTERTRYP, F-34398 Montpellier, France
[3]Centre International de Recherche-Développement sur l'Elevage en zone Subhumide (CIRDES), 01 BP 454,
Bobo-Dioulasso 01, Burkina Faso
[4]Univ. Montpellier, CNRS, INSERM, Montpellier France
[5]Montpellier GenomiX, France Génomique, Montpellier, France
[6]CIRAD, UMR AGAP Institut, F-34398 Montpellier, France
[7]UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France
[8]SELMET, INRAE, CIRAD, Montpellier Supagro, University of Montpellier, Montpellier, France
17
Corresponding author: Sophie Thévenon (sophie.thevenon@cirad.fr https://orcid.org/0000-0001-6059-
5884)

Moana Peylhard https://orcid.org/0000-0001-8574-4567
David Berthier https://orcid.org/0000-0002-3283-6588,
Guiguigbaza-Kossigan Dayo https://orcid.org/0000-0001-7926-5834
Isabelle Chantal https://orcid.org/0000-0002-3429-484X
Souleymane Sylla,
Sabine Nidelet,
Emeric Dubois, https://orcid.org/0000-0002-6052-4178
Guillaume Martin https://orcid.org/0000-0002-1801-7500
Guilhem Sempéré https://orcid.org/0000-0001-7429-2091
Laurence Flori https://orcid.org/0000-0002-7529-8521

Abstract

Animal African trypanosomosis, caused by blood protozoan parasites transmitted mainly by tsetse flies,
represents a major constraint for millions of cattle in sub-Saharan Africa. Exposed cattle include
trypanosusceptible indicine breeds, severely affected by the disease, and West African taurine breeds
called trypanotolerant owing to their ability to control parasite development, survive and grow in enzootic
areas. Until now the genetic basis of trypanotolerance remains unclear. Here, to improve knowledge of the
biological processes involved in trypanotolerance versus trypanosusceptibility, we identified bovine genes
differentially expressed in five West African cattle breeds during an experimental infection by *Trypanosoma*
congolense and their biological functions. To this end, whole blood genome-wide transcriptome of three
trypanotolerant taurine breeds (N'Dama, Lagune and Baoulé), one susceptible zebu (Zebu Fulani) and one
African taurine x zebu admixed breed (Borgou) were profiled by RNA sequencing at four time points, one
before and three during infection. As expected, infection had a major impact on cattle blood transcriptome
regardless of the breed. The functional analysis of differentially expressed genes over time in each breed
confirmed an early activation of the innate immune response, followed by an activation of the humoral
response and an inhibition of T cell functions at the chronic stage of infection. More importantly, we
highlighted overlooked features, such as a strong disturbance in host metabolism and cellular energy
production that differentiates trypanotolerant and trypanosusceptible breeds. N'Dama breed showed the
earliest regulation of immune response, associated with a strong activation of cellular energy production,
also observed in Lagune, and to a lesser extent in Baoulé. Susceptible Zebu Fulani breed differed from other

breeds by the strongest modification in lipid metabolism regulation. Overall, this study provides a better
understanding of the biological mechanisms at work during infection, especially concerning the interplay
between immunity and metabolism that seems differentially regulated depending on the cattle breeds.

Introduction

Animal African Trypanosomosis (AAT) represents a serious impediment to livestock development in
endemic areas of Africa (Alsan, 2015). This vector-borne disease is caused by blood extracellular protozoan
parasites from the *Trypanosoma* genus (e.g., *Trypanosoma congolense*, *T. vivax* and, to a lesser extent, *T.
b. brucei*) mainly transmitted by tsetse flies (genus *Glossina*). It affects about 50 million cattle in 38
countries in humid and sub-humid zones of Africa by causing high morbidity and mortality (Uilenberg,
1998), (Swallow, 2000). The Food and Agriculture Organization of the United Nations estimated its annual
cost at $4.5 billion (Budd, 1999), (Mattioli et al., 2004). Up to now, no vaccine is available and the main
prophylactic and curative measures are based on the reduction of transmission rates through vector
control (Bouyer et al., 2013), and the use of trypanocides in livestock (Meyer et al., 2016).
Interestingly, West African taurine (*Bos taurus taurus*) breeds (AFT), such as long-horn (i.e., N'Dama)
and short-horn breeds (e.g., Somba, Baoulé and Lagune) that have lived in West Africa for about 4000 years
(Payne & Hodges, 1997), (Hanotte et al., 2002), possess the ability to survive and remain productive in
tsetse-infested areas by controlling parasitemia and limiting anemia and body weight loss caused by AAT
(Murray et al., 1984), (CIPEA, 1979). AFT are thus called trypanotolerant. In contrast, both zebu (*Bos taurus
indicus*) cattle, which arrived more recently in Africa (<2000 YBP) (Loftus et al., 1994), (Bradley et al., 1996),
(Hanotte et al., 2002), and European taurine, which were recently introduced to increase African cattle
productivity (Seck et al., 2010), are susceptible to AAT (Roberts & Gray, 1973), (Amene et al., 1991), (Doko
et al., 1997).
The trypanotolerant character is therefore a remarkable example of livestock adaptation to a selective
pressure caused by a pathogenic agent. However, the molecular or biological mechanisms underlying this
trait have puzzled researchers for dozens of years, although some immunological and genetic studies
provided some clues. Indeed, the first immunological studies pointed out a better adaptive immune
response and an earlier monocyte lineage activation in N'Dama compared to African Boran zebu (Authie
et al., 1993), (Taylor et al., 1996), (Sileghem et al., 1993). Nevertheless, according to (Naessens et al., 2003),
the hematopoietic system, which participates in anemia reduction, was not involved in the control of
parasitemia. Moreover, several genetic studies on trypanotolerant and trypanosusceptible breeds revealed
the polygenic architecture of trypanotolerance (Murray et al., 1990), (Trail et al., 1991), (Van der Waaij et
al., 2003), (Hanotte et al., 2003). Further transcriptomic studies (O'Gorman et al., 2006), (O'Gorman et al.,
2009) showed that N'Dama and Boran Zebu exhibited a roughly similar response to infection by
trypanosomes, but subtle differences in response intensity or timing were observed, such as a higher IL6
and IL10 expression in Boran Zebu, or an enhanced B cell activation in N'Dama. By integrating genomic and
transcriptomic data, (Noyes et al., 2011) proposed TICAM1 and ARHGAP15 as two candidate genes for
trypanotolerance. However, these findings have not been confirmed by other studies analyzing gene
coding sequences in several breeds (Alvarez et al., 2015), (Alvarez et al., 2016). More experiments were
performed in mouse models, in which the possibility to knock out candidate genes allows to accurately
assess their contribution to tolerance or pathology (Cnops et al., 2015), (Magez et al., 2006), (Magez et al.,
2007), (Onyilagha et al., 2015). Nevertheless, mice are not natural hosts of livestock trypanosomes, and
important physiological features that differentiate them from cattle (Taylor & Mertens, 1999), (Morrison
et al., 2016) prevent to directly transpose results between the two species.
So far, studies, which investigated immunological, genetic and transcriptomic features of cattle
trypanotolerance, have mainly focused on two breeds, the trypanotolerant N'Dama cattle (long-horn
taurine originating from Fouta-Djallon in Republic of Guinea) and the trypanosusceptible Boran Zebu (an
East-African Zebu). Nevertheless, other West-African taurine cattle are classified as trypanotolerant (CIPEA,
1979), (Akol et al., 1986), (Rege, 1999) among which we recently confirmed, under experimental
conditions, the trypanotolerant status of two West African short-horn breeds, i.e., Lagune and Baoulé
breeds, compared to N'Dama and to the trypanosusceptible Zebu Fulani (Berthier et al., 2015). We also
underlined the intermediate status of Borgou breed, an admixed breed between African short-horn taurine

and African zebu. Biological samples collected during this experiment that allowed a clear characterization
of the trypanotolerant status of these overlooked breeds provide the opportunity to finely investigate
breed-specific modulation of gene expression during infection.
In order to increase knowledge on host-parasite interactions in trypanotolerant and trypanosusceptible
cattle breeds during trypanosomosis, we performed a gene expression profiling of blood cells of five cattle
breeds infected by *T. congolense* using RNA-seq technology, namely the well-characterized trypanotolerant
N'Dama breed, two overlooked trypanotolerant breeds, Baoulé and Lagune, the Borgou crossbred breed,
and one trypanosusceptible breed, the Zebu Fulani. In addition, we performed an in-depth functional
analysis of the differentially expressed genes. More precisely, we looked for i) breed-specific transcriptomic
signatures in blood before infection, ii) main genes and biological functions that responded to infection,
whatever the breed, iii) breed-specific transcriptomic profiles during infection, and iv) basal and dynamic
transcriptomic profiles that could be associated with trypanotolerance.

112   **Material and Methods**

Animals, experimental infection and sampling

A total of 39 animals from five West African cattle breeds, i.e., three AFT comprising N'Dama (NDA, 8
animals), Lagune (LAG, 7 animals), and Baoulé (BAO, 8 animals), Zebu Fulani (ZFU, 8 animals), and Borgou
(BOR, 8 animals), were experimentally infected by intravenous inoculation of 10^5 trypanosomes of the *T.*
congolense savannah IL1180 strain. This experimental infection, conducted at CIRDES (Burkina Faso)
according to a protocol approved by the Burkinabe ethical committee (Project no. A002-2013 / CE-CM),
was described in details in (Berthier et al., 2015).
Cattle blood samples were collected at the jugular vein using Tempus™ Blood RNA Tubes (Applied
Biosystems™, USA), which allowed the immediate blocking of mRNA transcription and degradation, at four
time points: before infection (named DPI.0, DPI for days post-infection) and during the infection at 20 days
post-infection (DPI.20), corresponding roughly to the increase in parasitemia, 30 DPI (DPI.30), around the
peak of parasitemia, and 40 DPI (DPI.40), at the time of the entrance in the chronic phase of the disease.
They were stored 24 hours at +4°C before treatment.

RNA extraction and RNA-seq libraries preparation

RNAs were extracted from blood samples using the Tempus™ Spin RNA Isolation Kit (Applied
Biosystems™, USA) according to the manufacturer's instructions. RNA was finally eluted using 80 µl RNase-
free buffer. Total RNA was quantified using a Nanodrop One (Thermo Fisher Scientific, USA) and its quality
checked on a Bioanalyzer 2100 using RNA 6000 nano kit (Agilent Technologies, USA). Samples with RNA
integrity number >=8.70 were selected.
RNA-seq libraries were constructed from 120 RNA samples obtained from six cattle per breed at four
time points at the MGX platform in Montpellier (France) using the TruSeq Stranded mRNA Library Prep Kit
(Illumina) following the manufacturer's instruction. Briefly, poly-A RNAs were purified using oligo-d(T)
magnetic beads from 1 µg of total RNA. The poly-A+ RNAs were fragmented and reverse transcribed using
random hexamers, Super Script II (Life Technologies) and Actinomycin D. During the second strand
generation step, dUTP substituted dTTP in order to prevent the second strand to be used as a matrix during
the final PCR amplification. Double stranded cDNAs were adenylated at their 3' ends before ligation that
was performed using Illumina's indexed adapters. Ligated cDNAs were amplified following 15 PCR cycles
and PCR products were purified using AMPure XP Beads (Beckman Coulter Genomics). Libraries were
validated using a Bioanalyzer on a DNA1000 chip (Agilent) and quantified using the KAPA Library
quantification kit (Roche).

Sequencing process

Clustering was performed on a cBot and sequencing on a HiSeq2000 (Illumina). After quantification,
the libraries were equimolarly pooled by 12, leading to 10 multiplexes of 12 samples each. Assignment to

the multiplexes was done by random blocking on the time point and the breed, in order for each multiplex
to contain the four time points (3 samples per time point) and the five breeds. Each pool was denatured,
diluted and clustered on three lanes using the Cluster Generation kit v3 (Illumina). Sequencing was
performed using SBS kit v3 (Illumina) in single read 50 nt mode. Raw sequencing data were saved as FASTQ
files. The quality of the data was assessed using FastQC from the Babraham Institute. Potential
contaminants were investigated with the FastQ Screen software (the Babraham Institute).

Reads' alignment to the bovine and trypanosome genomes

Reads were jointly mapped to both reference genomes of bovine (EnsemblDB *Bos taurus* UMD 3.1
release 79) and parasite (TritrypDB *Trypanosoma congolense* IL3000 release 9) using STAR aligner (STAR
2.4.0j, (Dobin et al., 2013)). To this end, mapping was performed in four steps: i) the STAR index was
generated from a unique multi-FASTA file obtained by concatenation of the bovine and trypanosome
sequence files, ii) reads from all libraries were aligned against the indexed reference sequences in order to
identify intron-exon junctions, iii) a new STAR index was generated from the unique multi-FASTA file
obtained by concatenation of the bovine and trypanosome sequence files and information of intron-exon
junctions, iv) a final step of read mapping was performed on the new STAR index. A maximum of three
mismatches were allowed and multi-mapping to up to 20 different positions was permitted according to
the following parameters: --alignIntronMin 50 --alignIntronMax 500000 --outFilterMultimapNmax 20 --
outFilterMismatchNmax 4 --outSAMunmapped Within. Information on reads location on both reference
genomes was contained in final BAM format files.
The percentages of reads that mapped uniquely to the bovine or to the trypanosome genome and reads
that had multiple matches were checked using Picard tools 1.130 (Broad Institute) and Samtools 1.2 (Li et
al., 2009). A large majority of reads was uniquely aligned to the bovine genome (85 to 88% and from 73%
to 82% for the samples sharing the index with the PhiX control, see S1 Table). The joint mapping approach
was validated by the very low number of reads that mapped to both genomes (bovine and trypanosome),
comprised between 17 and 124 (from 0.00004% to 0.0003% of input reads). The percentage of reads
uniquely aligned to the trypanosome genome varied greatly between samples, from 3×10^{-6} to 2.31%, with
a mean of 0.21% (S1 Table). Before the infection (DPI.0), few hundred reads were aligned on the parasite
genome, corresponding to a maximum ratio of 3.8×10^{-5} reads (reads uniquely aligned to the trypanosome
genome/number of uniquely aligned reads) (S1 Fig). The very low rate of sequences assigned to the
parasite genome before infection was considered as negligible and as a background noise of sequencing
technology and mapping algorithm (O'Rawe et al., 2015). During the infection (DPI.20, DPI.30, and DPI.40),
the number of reads assigned to the trypanosome genome increased and was closely related to
parasitemia (S1 Fig). Two animals (BO5 and Z4) did not show any increase in the ratio of reads uniquely
aligned to the trypanosome genome.

Transcript quantification and data normalization

Quantification of gene expression was performed for each library using FeatureCounts (Subread
package 1.4.6-p4 (Liao et al., 2014)). Bovine gene annotation was downloaded from Ensembl 79 (sequence
UMD 3.1). Reads were assigned at the exon level and counts were summarized at the gene level, using
default parameters (-t exon -g gene_id), corresponding to unambiguously assigning uniquely aligned single-
end reads and reversely stranded reads (-s 2).
Following the advised workflow of the Bioconductor package EdgeR 3.18.3 (Robinson et al., 2010), (R
Core Team, 2018), we first removed lowly expressed genes, and kept genes that had more than one count
per million in at least two libraries. Out of the 24,616 bovine genes annotated in Ensembl 79, 13,107 genes
went through the filter. Normalization of count data was performed using the Trimmed Mean of M-values
normalization (Robinson et al., 2010) and dispersion was estimated using a Cox-Reid profile-adjusted
likelihood (Chen et al., 2014), using the Bioconductor package EdgeR 3.18.3 under R 3.4.0 environment
(Robinson et al., 2010), (R Core Team, 2018).

Exploration of the global structuration of bovine count data and sample selection

A two-dimensional scatterplot was launched on the normalized count data to assess the global structuration of 120 samples (Ritchie et al., 2015). It was completed by a Principal Component Analysis (PCA) performed on the log of normalized count per million using mixOmics_6.3.0 (Le Cao et al., 2016) (with center=TRUE and scale=FALSE). The first factorial plan separated the samples according to sampling time point, except for BO5 and Z4 whose samples clustered together (S2 Fig). This observation and the very low ratio of sequences assigned to the trypanosome genome (S1 Fig) supported the hypothesis that the infection process did not occur in BO5 and Z4, in accordance with the phenotypic analysis that revealed transient parasitemia and no anemia in these animals (Berthier et al., 2015). BO5 and Z4 were thus discarded from subsequent analyses. In addition, a Baoulé (BA3) that was detected positive in parasitemia before the experimental infection, contrary to the others that were negative based on diagnostic tests (Berthier et al., 2015), was also excluded from further analyses. The final data set used for statistical and functional analyses contained 108 RNA-seq libraries, corresponding to 27 animals (6 NDA, 6 LAG, 5 ZFU, 5 BOR, and 5 BAO) and four time points per animal.

Differential expression analyses of bovine genes

The differential expression analysis of bovine genes was carried out using the Bioconductor package EdgeR 3.18.3 under R 3.4.0 environment (Robinson et al., 2010), (R Core Team, 2018), which models gene count data according to a negative binomial distribution and moderates the degree of over-dispersion across genes. A generalized linear model (GLM) was fitted for each gene, and tests for determining differential expression were done using a likelihood ratio test (McCarthy et al., 2012). We considered nineteen contrasts to assess differential expressions between two conditions using GLM (Fig 1).

Figure 1. Description of the contrasts used for the differential gene expression analysis. The numbers in the circles represent the days post-infection of cattle sampling, at DPI.0, DPI.20, DPI.30 and DPI.40. Orange arrows represent the contrasts between the breeds at DPI.0, before infection, named BOR.0-ZFU.0, LAG.0-ZFU.0, NDA.0-ZFU.0, and BAO.0-ZFU.0. The blue arrows represent the within-breed contrasts: for each breed, three contrasts were built with DPI.0 as reference, DPI.20-0, DPI.30-0, and DPI.40-0. ZFU, LAG, and BAO pictures by S. Thévenon; BOR picture by G-K. Dayo; NDA picture by D. Berthier.

First, to study baseline differential expression between breeds before infection, we chose ZFU as breed
reference, the unique indicine trypanosusceptible breed whose samples were placed at the extremity of
the second axis in the PCA (S2 Fig). Four contrasts, comparing NDA, LAG, BAO and BOR breeds to ZFU breed
at DPI.0 (Fig 1) were established (namely NDA.0-ZFU.0, LAG.0- ZFU.0, BAO.0-ZFU.0 and BOR.0-ZFU.0).
Second, to assess how each breed reacted to infection, 15 within breed contrasts, corresponding to three
contrasts that assessed differential expression between three post-infection time points and the pre-
infection time point (namely DPI.20-0, DPI.30-0 and DPI.40-0, Fig 1), were considered for each breed.
Because animals were repeatedly sampled, the design matrix was constructed according to a nested
factorial formula, with animals and time points nested within the breed. GLM likelihood ratio tests provided
for each contrast and each gene a logFC (log2-fold change of gene expression between conditions) and a
FDR (False Discovery Rate) corresponding to adjusted p-values for multiple testing using the Benjamini-
Hochberg procedure (Benjamini & Hochberg, 1995). In our analyses, since 19 contrasts were done, a FDR
of 10^{-3} (0.05/19=0.0026 rounded to 0.001) was chosen to identify differentially expressed genes (DEGs).

Functional analysis

The web-based software application Ingenuity® Pathway Analysis (IPA®, Version 43605602, 2018-04-
04) was used to perform the functional analysis of the DEGs, based on the content of the Ingenuity®
Knowledge Base (IKB).
For all contrasts detailed above, among the 13,107 bovine Ensembl identifiers (with their associated
logFC and FDR) that defined the background gene list used in IPA® analyses and that were uploaded into
the software application, 11,316 identifiers were mapped to their corresponding object in IKB®. We then
checked via Ensembl Biomart the existence of human orthologues for the 1,791 bovine identifiers that
were not recognized in IKB® and found 577 human orthologues with a high confidence level and a one-to-
one match. Human Ensembl identifiers for these 577 genes were then used instead of bovine identifiers in
files uploaded into IPA®. At last, 11,893 Ensembl identifiers were mapped to known genes in IKB®.
The functional analysis performed with IPA® identified biological diseases and functions, canonical and
signaling pathways and upstream regulatory molecules that were significantly enriched in our data sets.
Upstream regulators are regulatory molecules that can affect the expression of target DEGs and that may
not have been detected as DEG by RNA-seq (because, e.g., they may be expressed in another tissue than
the sampled one, or at another time than the sampling date, or because they are endogenous biochemical
compounds). Right-tailed Fisher's exact test was used to calculate a p-value determining the probability
that each biological function and/or disease and canonical pathways assigned to our data sets was due to
chance alone. These p-values were adjusted using the Benjamini-Hochberg correction method (B-H) for
diseases and functions and canonical pathways. Diseases and functions, canonical pathways, and upstream
regulators were considered as significant if their corresponding p-values were below 10^{-3} (B-H correction),
10^{-2} (B-H correction) and 10^{-4} respectively, according to the fact that the quantity of information provided
by these functional categories, and thus the number of tests, differed. In addition, based on the logFC of
the DEGs and IKB® information, IPA® inferred activation states (namely "activated" or "inhibited") of
biological functions, pathways and upstream regulators (indicating that the observed up or down-
regulations of the DEGs are mostly consistent with a particular activation state of a biological function or a
regulator) by estimating Z-scores associated with the enriched functions or regulators (Kramer et al., 2014).
A Z-score ≥ 2 corresponded to an inferred significant activation state of a function or a regulator, while a
Z-score ≤ -2 corresponded to an inferred inhibition state. IPA® outputs were visualized using ggplot2 R
package (Wickham, 2016).

257   **Results**

In order to better understand host-parasite interactions in trypanotolerant and trypanosusceptible
cattle breeds during infection, we carried out an overall assessment of the relationships between the
samples, the identification of initially differentially expressed genes between breeds (i.e., before infection),
and the identification of differentially expressed genes in each breed during infection, with a first focus on

common DEGs and the core associated biological processes, followed by an enlightenment on biological
processes specific or prominent to each breed.

Global overview of bovine gene expression data set

265 We first carried out an overall assessment of the relationships between the samples, based on a PCA
266 performed on the logarithm of normalized bovine gene counts that provided a global overview of the cattle
267 transcriptomes according to the sampling time point and the breed (Fig 2).

268
269

Figure 2. Principal components analysis of cattle RNA-seq libraries based on normalized genes counts.
Each point represents a RNA-seq library that corresponds to an animal sampled at a given DPI and that is
plotted on the first two principal components according to its coordinates. Libraries are identified
according to the breed and the sampling date. Arrows link a library to the centroid of the corresponding
breed and sampling date treatment. Each breed is represented by a different shape and by a color
gradient, and each color is graded from light to dark shades corresponding to days post-infection
respectively DPI.0, 20, 30 and 40. ZFU animals are in red and triangle down, BOR animals are in violet and
diamond, LAG animals are in orange and triangle, BAO animals are in blue and circle, and NDA animals
are in green and square. Percentages of variance explanation of the first two components are added.

The first axis that accounted for 16% of the total variation was representative of the infection course,
from DPI.0 to DPI.40. Shifts from DPI.0 to DPI.40 for each breed were roughly parallel. The second axis that
accounted for 10% of the total variation was representative of the breed effect with ZFU count data, at the
bottom, taurine count data (represented by NDA, BAO and LAG breeds) at the top and admixed BOR breed
data, in the middle, suggesting that basal and lasting count differences existed between zebu, taurine and
admixed breeds.

Before infection, a total of 310 genes were detected as differentially expressed (DE) between at least one breed and ZFU (FDR<10^{-3}). During infection, a total of 5,270 differentially expressed genes (DEGs) were identified in at least one contrast, between 41 and 3,839 genes being DE depending on the contrast (Fig 3).

Figure 3. Numbers of genes identified as differentially expressed in the 15 within-breed contrasts. The x-axis represents the 15 contrasts ordered by breed respectively BAO, BOR, LAG, NDA and ZFU, and for each breed, three contrasts according to days post-infection in comparison with before infection, respectively DPI.20-0, 30-0, 40-0. The y-axis represents the number of differentially expressed genes at the threshold of FDR<0.001. The red color corresponds to the part of genes that are differentially expressed and common to, at least, two breeds whatever the sampling time point; the blue color corresponds to the part of genes that are differentially expressed uniquely within a breed.

As expected, the bovine transcriptome was massively modified by trypanosome infection, the total number of DEGs during infection being in descending order 4,344, 2,715, 2,141, 1,753 and 1,696 for NDA, LAG, ZFU, BOR and BAO, respectively. NDA displayed the earliest modulation of its transcriptome, with 1,067 DEGs at DPI.20-0, followed by LAG (370 DEGs), BOR (351 DEGs), ZFU (77 DEGs), and BAO (41 DEGs) (Fig 3). At DPI.30-0, LAG displayed the highest number of DEGs (2,330). At DPI.40-0, NDA showed again the highest number of DEGs (3,839). Many DEGs were common between two or more breeds including the trypanosusceptible ZFU (Fig 3 and S3 Fig). A total of 3,283 genes (62.3% of total number of DEGs, so a majority) were shared between at least two breeds. For instance, 1,823 DEGs were shared between NDA and ZFU, 1,318 between LAG and ZFU, 1,074 between BOR and ZFU, and 1,069 between BAO and ZFU (S3 Fig), whereas the number of DEGs detected exclusively within a breed was 1,272 in NDA, 331 in LAG, 211 in ZFU, 96 in BOR and 77 in BAO. Interestingly, the direction of variation of DEGs, when detected DE in several contrasts, whatever the time point or the breed, was always identical in the aforementioned contrasts (i.e., upregulated with positive logFC or downregulated with negative logFC) except for one gene,

SPARC, which was upregulated in LAG.40-0 but downregulated in BAO.40-0 (S2 Table). The heatmap (Fig
4), performed on the logFCs of 5,270 DEGs in at least one contrast, first clustered contrasts according to
the time of infection, the early date (DPI.20-0) being separated from the others, except for the contrast
BAO.30-0. At later time points, contrasts from a same breed tended to cluster together (LAG.30-0 with
LAG.40-0, BOR.30-0 with BOR.40-0, ZFU.30-0 with ZFU.40-0), although NDA.40-0 and BAO.40-0 were not
close to NDA.30-0 and BAO.30-0 respectively. Fig 4 did not highlight obvious different patterns of gene
expression between breeds.

.

Figure 4. Heatmap on the logFC of 5,270 DEGs in the 15 within-breed contrasts. Columns represent
the within-breed contrasts ordered according to clustering, and rows the genes identified as DE in at least
one contrast. Up-regulated genes (i.e. with positive logFC) are colored in warm colors from light red to
dark red (estimated maximum logFC=11.3), whereas down-regulated genes (i.e. with negative logFC) are
colored in cold colors, from light blue to dark blue (estimated minimum logFC=-8.00). LogFC comprised
between -0.20 and 0.20 are colored in grey.

301 Transcriptome profiling before infection highlights basal differences between breeds.

We then focused on the differences before infection of gene expression levels between each breed and
ZFU taken as trypanosusceptible reference. For the contrasts LAG.0-ZFU.0, NDA.0-ZFU.0, BAO.0-ZFU.0 and
BOR.0-ZFU.0 respectively, 127, 152, 156, and 63 genes were differentially expressed at the FDR threshold
of 10^{-3} (Table S3), showing initial differences between gene counts in ZFU and in the other taurine or
admixed breeds. As expected from PCA results, the number of DEGs between ZFU and BOR was smaller
than those between ZFU and AFT (i.e., NDA, BAO, LAG) at DPI.0. The functional annotation of DEGs for the
four contrasts before infection using IPA® indicated that basal differences were not associated with specific

biological function enrichment, since neither any disease and function nor any canonical pathway was
significantly enriched in these gene sets. However, fifteen upstream regulators, precisely 1, 1, and 15 for
the contrasts LAG.0-ZFU.0, BAO.0-ZFU.0 and BOR.0-ZFU.0 respectively, were identified at a p-value<10^{-4}
(S4 Table). Not any upstream regulator was significantly enriched in NDA.0-ZFU.0. Only BSG, encoding a
plasma membrane protein involved in several molecular functions (i.e., cadherin binding, carbohydrate
binding, mannose binding, monocarboxylic acid transmembrane transporter activity, and protein binding)
was identified as upstream regulator in the three contrasts involving LAG, BAO and BOR. Several upstream
regulators linked to immune response (e.g., IL4, TNF, IFNG, Immunoglobulin, CD3 complex, prostaglandin
E2, IL10RA) and transcription regulators (e.g., HDAC3, FOXA2, STAT3, ATF3) were enriched in the BOR.0-
ZFU.0 gene set.
Among the DEGs before infection, 35, 54, 83 and 18 genes were only DE in LAG.0-ZFU.0, NDA.0-ZFU.0,
BAO.0-ZFU.0 and BOR.0-ZFU.0 contrasts, respectively (S4 Fig). Besides, twelve genes were found DE in the
four contrasts, 36 were shared between the three AFT breeds versus ZFU.0, and 70 were shared between
LAG.0-ZFU.0 and NDA.0-ZFU.0. Among these latter 70 DEGs, all genes except PELI3 were downregulated in
LAG and NDA in comparison to ZFU, among which several genes are known to be linked to immune
response, especially IL2RA, GBP2, PELI3, DCSTAMP, PTX3, and MARCO (S3 Table).

Transcriptomic responses common to all breeds, involving immune response and metabolism, are detected during infection

Genes differentially expressed during infection in all breeds.

To understand the core response of bovine transcriptome to trypanosome infection, we studied the
659 DEGs common to all breeds (namely DEGs in at least one contrast of each breed, S3 Fig). S2 Table gives
detailed information on each gene by indicating whether it was DE, in which breed, and its average FDR
and logFC. Six genes (NR4A1, CCL22, IFI30, CTSZ, KYNU, IL17REL), involved in immune response, were
significantly upregulated in all contrasts, meaning that they were DE within each breed at each time point
during infection (S2 Table). Among the top upregulated genes (i.e., harboring the highest average logFC),
we found HBM, coding for a hemoglobin subunit (average logFC=6.96, average -log10(FDR)=5.7), ARG1
(average logFC=6.90, average -log10(FDR)=6.0), and several genes involved in immune response like
MAPK12, MMP14 and MAPK11. The top downregulated genes (i.e., DEGs displaying the smallest negative
logFC) were UNC5A (average logFC =-2.84, average −log10(FDR)=4.7) followed by OVOS2, ELANE, DAB2,
and BPI. A quick overview of potentially interesting DEGs associated with immune response allowed
highlighting three cytokines (i.e., IL7 upregulated; IL16 and LIF downregulated) and cytokine receptors (e.g.,
IL1R, IL6R, IL7R, IL20RB, all down-regulated), transcription regulators (e.g., NFKB1 and NFKB2 upregulated),
other receptors (e.g., TFRC upregulated), and numerous immune cell antigens, some up-regulated (e.g.,
CD109, CD180, CD19, CD1A, CD22, CD40, CD72, CD79B, and MME syn.CD10), and others down-regulated
(e.g., CD7, CD2, CD226, CD247, CD27, CD3D, CD3E, CD3G, CD40LG, CD99, and ZAP70). Interestingly, several
genes known to be involved in metabolism were up-regulated (e.g., HMGCS1, CYP51A1, FDFT1, IDI1, LSS,
MVD, SQLE).

Enrichment of diseases and functions in the common DEGs.

The functional annotation of DEGs common to the breeds using IPA® identified 164 functions and
diseases significantly enriched with B-H corrected p-values <10^{-3} (S5 Table), and they could be grouped into
24 large categories displayed in Table 1. The major categories were cell-to-cell signaling and interaction,
cellular movement, cellular development, cell death and survival, hematological system development and
function, cellular function and maintenance, and lipid metabolism. We could also highlight cell-mediated
immune response, immunological disease, inflammatory response, and lymphoid tissue structure and
development. More precisely, the top ten diseases and functions in terms of B-H corrected p-values (<10^{-8}
) were: proliferation of lymphocytes, proliferation of lymphatic system cells, proliferation of blood cells,
proliferation of immune cells, lymphopoiesis, synthesis of sterol, synthesis of cholesterol, quantity of
leukocytes, quantity of lymphocytes, and cell movement of lymphatic system cell (S5 Table). IPA® analyses

allowed us to assess the activation or inhibition of the enriched diseases and functions based on the Z-
score, considered significant if its absolute value was larger or equal to 2. Twenty-four diseases and
functions presented significant Z-scores, and surprisingly most had negative Z-scores that could be related
to an inhibition state, for instance: lymphocyte homeostasis, adhesion of lymphocytes, migration of
lymphatic system cell, and T cell development. Only two diseases and functions had positive Z-score,
namely quantity of B-2 lymphocytes and liver lesion.
363

Table 1. Diseases and functions categories enriched in the common DEGs in the within-breed contrasts. The name of diseases and functions categories was indicated, with for each category, the number of significant functions, their average B-H corrected *p*-value, the range of the B-H corrected *p*-values, the average Z-score, and its range. Na: not available.

Diseases and functions category	Number of functions	Average B-H p-value	Range of B-H p-values	Average Z-score	Range of Z-scores
Cell-To-Cell Signaling and Inter-action	28	5.7E-05	7×10^{-4} ; 10^{-8}	-1.78	-2.71 ; 1.13
Cellular Movement	19	1.5E-04	9×10^{-4} ; 5×10^{-9}	-1.45	-2.66 ; 0.54
Cellular Development	18	1.0E-04	7×10^{-4} ; 3×10^{-12}	-0.34	-2.08 ; 1.02
Cell Death and Survival	16	1.5E-04	9×10^{-4} ; 6×10^{-7}	-1.09	-2.36 ; 0.03
Hematological System Development and Function	15	2.5E-04	9×10^{-4} ; 4×10^{-9}	-0.54	-2.07 ; 2.25
Cellular Function and Maintenance	9	2.4E-04	9×10^{-4} ; 2×10^{-7}	-0.87	-2.82 ; 0.14
Lipid Metabolism	8	1.1E-05	6×10^{-5} ; 6×10^{-10}	1.10	0.70 ; 1.63
Cancer	6	2.1E-04	6×10^{-4} ; 5×10^{-6}	-0.05	-0.43 ; 0.33
Cell Morphology	6	1.9E-04	97×10^{-4} ; 3×10^{-5}	-0.73	-1.00 ; -0.47
Cell-mediated Immune Response	6	1.7E-04	5×10^{-4} ; 6×10^{-8}	-2.11	-2.65 ; -1.77
Cell Signaling	5	3.5E-04	9×10^{-4} ; 5×10^{-6}	-0.38	-1.04 ; -0.05
Immunological Disease	5	2.0E-04	7×10^{-4} ; 3×10^{-6}	-1.52	-2.35 ; -0.67
Inflammatory Response	5	1.5E-04	7×10^{-4} ; 5×10^{-8}	-0.72	-2.39 ; 1.21
Lymphoid Tissue Structure and Development	4	2.3E-04	9×10^{-4} ; 5×10^{-8}	-0.43	-1.15 ; 0.11
Connective Tissue Disorders	2	2.2E-05	3×10^{-5} ; 9×10^{-6}	0.05	-0.13 ; 0.23
Embryonic Development	2	5.4E-04	9×10^{-4} ; 2×10^{-4}	-0.58	-1.14 ; -0.02
Inflammatory Disease	2	7.0E-05	8×10^{-5} ; 5×10^{-5}	-0.76	-2.43 ; 0.90
Organismal Development	2	1.1E-04	2×10^{-4} ; 2×10^{-5}	Na	Na
Cellular Growth and Proliferation	1	3.3E-12		-0.21	
Tissue Morphology	1	2.6E-06		-0.16	
Molecular Transport	1	3.2E-04		-0.76	
Gastrointestinal Disease	1	5.3E-04		2.06	
Cardiovascular Disease	1	9.6E-04		Na	
Connective Tissue Development and Function	1	9.6E-04		0.60	

Enrichment of canonical Pathways in the common DEGs.

Twenty-three canonical pathways were significantly enriched in common DEGs with a B-H corrected *p*-
value <10^{-2} (Table 2). This list highlighted several enriched and activated metabolic pathways (Z-score≥2),
e.g., the superpathway of cholesterol biosynthesis, the mevalonate pathway, the TCA (TriCarboxylic acid)
cycle II, and the superpathway of geranylgeranyldiphosphate biosynthesis I. Mitochondrial dysfunction was
also enriched but without inference on the direction of activation. The second class of enriched pathways
concerned the immune response including, for instance, role of NFAT in regulation of the immune
response, primary immunodeficiency signaling, and regulation of IL-2 expression in activated and anergic
T lymphocytes. Two immune response pathways, PI3K signaling in B lymphocytes and B cell receptor
signaling, were estimated as significantly activated.

Table 2. Canonical and signaling pathways enriched in the common DEGs in the within-breed contrasts. The name of the canonical and signaling pathways is indicated with its B-H corrected *p*-value, its Z-score, and the ratio between the number of DEGs and the number of genes in the pathway. Highlighted cells correspond to Z-scores with an inferred activation state.

Canonical and signalling pathways	B-H *p*-value	Z-score	Ratio
Superpathway of Cholesterol Biosynthesis	5.2E-08	3.61	0.52
Mevalonate Pathway I	6.9E-04	2.45	0.60
TCA Cycle II (Eukaryotic)	6.9E-04	2.83	0.40
Superpathway of Geranylgeranyldiphosphate Biosynthesis I (via Mevalonate)	6.9E-04	2.65	0.50
Mitochondrial Dysfunction	1.7E-03	Na	0.15
Role of NFAT in Regulation of the Immune Response	1.7E-03	0.22	0.15
Primary Immunodeficiency Signaling	1.7E-03	Na	0.32
PI3K Signaling in B Lymphocytes	1.7E-03	2.32	0.16
Cholesterol Biosynthesis I	1.7E-03	2.45	0.46
Cholesterol Biosynthesis II (via 24,25-dihydrolanosterol)	1.7E-03	2.45	0.46
Cholesterol Biosynthesis III (via Desmosterol)	1.7E-03	2.45	0.46
Regulation of IL-2 Expression in Activated and Anergic T Lymphocytes	2.4E-03	Na	0.18
Leukocyte Extravasation Signaling	4.5E-03	0.73	0.14
Hematopoiesis from Pluripotent Stem Cells	4.8E-03	Na	0.38
T Cell Receptor Signaling	5.1E-03	Na	0.16
Sirtuin Signaling Pathway	5.2E-03	-0.78	0.12
IL-7 Signaling Pathway	5.5E-03	0.30	0.19
iCOS-iCOSL Signaling in T Helper Cells	5.6E-03	0.33	0.17
CD28 Signaling in T Helper Cells	7.1E-03	-0.30	0.15
B Cell Development	7.1E-03	Na	0.33
Systemic Lupus Erythematosus In B Cell Signaling Pathway	7.1E-03	-0.21	0.12
B Cell Receptor Signaling	7.1E-03	2.68	0.13
Phospholipase C Signaling	8.9E-03	0.24	0.12

Enrichment of upstream Regulators in the common DEGs.

Seventy-three upstream regulators that can affect the expression of target DEGs were considered as
significantly enriched (*P*-value<10^{-4}, S6 Table), and the direction of activation was inferred for some of
them. The list of upstream regulators referred to chemicals, simple protein or protein complexes. Four
endogenous chemicals were identified: cholesterol, which was inferred as inhibited, beta-estradiol,
prostaglandin E2 and D-glucose. Nine cytokines were significantly enriched: TNF, IL15, IL4, IL3, IFNG, IL10,
CSF1, the latter being estimated as activated, and IL2 and IL7, both estimated as inhibited. Detected protein
complexes were also related to immune response (i.e., immunoglobulin, C4BP, TCR, Ige, and NFkB).
Twenty-seven transcription regulators were identified, some involved in metabolism (e.g., the activated
SREBF2, SREBF1, and PPARGC1B), cell cycle (e.g., TP53 inhibited, SP1) or in pleiotropic functions (e.g., SIRT2
activated). Several transcription factors were linked to immune response (e.g., STAT6 and STAT3, TCF3
inhibited, and BCL6 activated).

Specific transcriptomic response of each breed during infection

In order to look for potential breed-specific responses during infection, we performed separate
enrichment analyses of the lists of DEGs of the 15 within-breed contrasts.

Enrichment of diseases and functions in each breed during infection.

The analysis identified 642 diseases and functions significantly enriched in at least one of the 15
contrasts (B-H corrected *p*-values <10^{-3}) (S7 Table). The most enriched contrast was ZFU.40-0 with 365
significant diseases and functions, while no disease and function were enriched in LAG.20-0. As for the
enrichment of the common DEGs, large categories were cellular development and cell cycle, hematological
system development and function, immune response, cell signaling, and lipid metabolism. Among these
642 diseases and functions, 135 displayed significant Z-scores for the corresponding contrasts, 17 being
considered as activated (Z-score≥2), while a majority (118) was inhibited (Z-score≤-2). We confirmed the
results obtained with the analysis of common DEGs, namely a shared inhibition of numerous cellular
functions, especially associated with cell-mediated immune response from 30 DPI (e.g., T cell homeostasis).
Fig 5 represents both the significance level and the inhibition state of 41 diseases and functions harboring
Z-scores smaller than -3 for the 15 contrasts. Few inhibited functions were set up exclusively by one or few
breeds such as the differentiation of mononuclear leukocytes, significantly inhibited in NDA.40-0 only.
Chemotaxis of myeloid cells was inhibited in BAO.40-0, LAG.40-0, and ZFU.40-0 contrasts but was
considered significantly enriched in ZFU only.

Figure 5. Plot of 41 significantly enriched diseases and functions inferred as inhibited in function of the within-breed contrasts. The 15 within-breeds contrasts are shown on the abscissa and the significantly enriched diseases and functions on the ordinate. The size of the circles is inversely proportional to the B-H corrected *p*-value calculated for diseases and functions in the aforementioned contrast. The color gradient of the circle corresponds to the Z-score range, i.e. warm color gradient for positive Z-score>1, cold color gradient for negative Z-score<-1, and gray color gradient for Z-scores ranging between -1 and 1.

Fewer diseases and functions were activated, but they presented more discriminating patterns than
inhibited functions. Fig 6 shows that lipid metabolic functions (synthesis of lipid, synthesis of cholesterol,
steroid metabolism, metabolism of cholesterol, and synthesis of terpenoid) were strongly enriched and
activated in ZFU.20-0 contrast, to a lesser extent in BAO.30-0, and slightly in NDA.20-0, but were not
detected in LAG. Glucose metabolism disorder was enriched in ZFU and BAO, but activated in BAO.30-0
contrast only. Cytopenia was significantly activated in ZFU.40-0 contrast.
The cancer category (i.e., 63 significant functions) and cell cycle functions (i.e., 19 significant functions
like cell cycle progression, mitosis) were especially and precociously enriched in NDA.20-0 (S7 Table). G2/M
phase transition, metabolism of DNA, segregation of chromosomes, and M phase were strongly activated

in NDA.20-0 contrast. This latter was also found activated in BOR.20-0. Interphase was activated in LAG.30-0. BOR.40-0 presented an activation of RNA transcription.

Figure 6. Plot of 17 significantly enriched diseases and functions inferred as activated in function of the within-breed contrasts. The 15 within-breeds contrasts are shown on the abscissa and the significantly enriched diseases and functions on the ordinate. The size of the circles is inversely proportional to the B-H corrected *p*-value of the diseases and functions in the aforementioned contrast. The color gradient of the circle corresponds to the Z-score range, i.e. warm color gradient for positive Z-score>1, cold color gradient for negative Z-score<-1, and gray color gradient for Z-scores ranging between -1 and 1.

Enrichment of canonical pathways in each breed during infection.

A total of 92 canonical pathways were significantly enriched in at least one contrast during infection, with a B-H corrected *p*-value<10^{-2} (S8 Table). The contrast ZFU.40-0 displayed the highest number of significant canonical pathways (48), followed by BAO.30-0 (39), while no significant pathways were detected in LAG.20-0, LAG.40-0, and BAO.20-0. Among the 92 significant canonical pathways, 26 presented also significant Z-scores, measuring their activation state: 7 being considered inhibited and 19 activated (Fig 7).

Figure 7. Plot of 26 significantly enriched canonical pathways inferred as activated or inhibited in function of the within-breed contrasts. The 15 within-breeds contrasts are shown on the abscissa and the significantly enriched diseases and functions on the ordinate. The size of the circles is inversely proportional to the B-H corrected *p*-value of the canonical pathways in the aforementioned contrast. The color gradient of the circle corresponds to the Z-score range, i.e. warm color gradient for positive Z-score>1, cold color gradient for negative Z-score<-1, and gray color gradient for Z-scores ranging between -1 and 1.

The superpathway of cholesterol biosynthesis was found significantly activated in all breeds, at one (LAG and NDA), two (BAO and BOR) or three time points (ZFU), but the set of pathways related to cholesterol and lipid metabolisms was particularly and durably enriched in ZFU, and to a lesser extent in BAO. Likewise, LXR/RXR was activated in ZFU.40-0 and almost in BAO.40-0.

Conversely, oxidative phosphorylation, TCA cycle, mitochondrial dysfunction, sirtuin signaling pathway and cell cycle associated pathways were highly significant in NDA contrasts. Pathways associated with cell energy production, oxidative phosphorylation and TCA cycle, were strongly activated in NDA, and to a lesser extent in LAG (oxidative phosphorylation and TCA cycle) and BAO (TCA cycle). Glycolysis and gluconeogenesis were considered enriched and activated in NDA.20-0, and gluconeogenesis was also significantly activated in BOR.20-0. Cell cycle associated pathways (Estrogen-mediated S-phase Entry, Cyclins and Cell Cycle Regulation, Cell Cycle Regulation by BTG Family Proteins, and mitotic Roles of Polo-

Like Kinase) were activated in NDA.20-0 exclusively. The sirtuin signaling pathway was highly inhibited in
NDA during the infection, and the G1/S checkpoint regulation was inhibited in NDA.20-0.
Some pathways, linked to immune response, were transiently enriched in some breeds, but not in NDA,
and their direction of activation was not able to be inferred (i.e., leukocyte extravasation signaling
significant in ZFU.40-0, LAG.30-0, and BAO.30-0 and BAO.40-0; hematopoiesis from pluripotent stem cells
significant in LAG.30-0, BAO.30-0 and BAO.40-0, and BOR.40-0; natural killer signaling significant in ZFU,
BAO and BOR at 40-0). Two B-cell associated pathways (B Cell Receptor signaling, and PI3K Signaling in B
Lymphocytes) were activated in BAO.40-0. Interestingly, five pathways were significantly inhibited in
ZFU.30-0 and/or ZFU.40-0 (i.e., IL-6 Signaling, NF-kB Signaling, p38 MAPK Signaling, Th2 pathway and the
inflammasome pathway).

Upstream Regulators enriched in each breed during infection.

316 upstream regulators that can affect the expression of target DEGs were considered as significantly
enriched in at least one contrast (P-value$<10^{-4}$, S9 Table). Among them, 95 upstream regulators belonged
to the transcription regulator category, 29 were endogenous chemicals and 26 were cytokines according
to IPA® classification. Upstream regulators found significant in almost all contrasts at the chosen threshold
were TGFB1 (14/15 contrasts), TNF (13/15 contrasts), CSF2, Vegf (12/15 contrasts), TCF3, IL4, PTEN (11/15
contrasts). The contrast ZFU.20-0 presented 115 significant upstream regulators, while only eight and 13
were significantly enriched in BAO.20-0 and LAG.20-0 respectively; NDA.20-0 and BOR.20-0 displayed 86
and 87 significant upstream regulators respectively.
In order to visualize the main activated or inhibited regulators predicted by IPA according to the up- or
downregulation of the target DEGs, Fig 8 presents the estimated Z-scores of 61 upstream regulators with
significant *p*-values ($<10^{-6}$) and Z-scores (|Z-score|≥2) in at least one significant contrast. Among them, 29
upstream regulators, presenting positive Z-scores, were assessed as activated in some contrasts during
infection (top of Fig 8), 25 were roughly inhibited (middle of Fig 8), and 7 showed a dynamic pattern with
rather positive Z-scores at DIP.20 followed by negative Z-scores (bottom of Fig 8).

Figure 8. Plot of 61 significantly enriched upstream regulators inferred as activated or inhibited in function of the within-breed contrasts. The 15 within-breeds contrasts are shown on the abscissa and the significantly enriched diseases and functions on the ordinate. The size of the circles is inversely proportional to the P-value of the upstream regulator in the aforementioned contrast. The color gradient of the circle corresponds to the Z-score range, i.e. warm color gradient for positive Z-score>1, cold color gradient for negative Z-score<-1, and gray color gradient for Z-scores ranging between -1 and 1.

ZFU was characterized by very high Z-scores and levels of significance during infection for SREBF2, SCAP,
and SREBF1. These upstream regulators were also significantly activated in the other breeds but not
constantly during infection. Interestingly, the SREBF2 gene itself was significantly and exclusively up-
regulated in ZFU contrasts. ATP7B was durably activated in ZFU, and temporarily in BAO and BOR. ZFU
displayed constantly inhibited upstream regulators: cholesterol, CYP51A1, ACACB, INSIG1, INSIG2, and
POR.
NDA was distinguished on Fig 8 by strong activation of CSF2, INSR, MYC, HGF, Vegf, and PTGER2. These
upstream regulators were also activated in the other breeds but with smaller intensities and durations. The
high inhibition of RICTOR was a unique feature of NDA, as well as that of KDM5A. The last three breeds,
BAO, BOR, and LAG, did not present striking features. The contrast BAO.30-0 shared similarities with

ZFU.20-0, naming high activation of SREBF1, SREBF2, and SCAP, and inhibition of INSIG1 and POR. BAO and
BOR presented an activation of NFkB (complex). BOR and LAG displayed similar features to NDA (e.g.,
strong activation of CSF2 or FOXO1). LAG only showed a significant activation of TBX21 and an inhibition of
GATA3 at DPI.40-0.
As upstream regulators, several cytokines (IL2, IL5, IL7, and TNF) and the growth factor TGFB1 showed
a dynamic pattern: they presented rather positive Z-scores at DPI.20 and were then assessed as
significantly inhibited at 30 or 40 DPI (bottom of Fig 8). TNF was significantly activated in BOR.20-0 and
significantly inhibited in NDA.40-0 and ZFU.40-0. Likewise, TGFB1 was activated in BAO.20-0 and inhibited
in NDA, ZFU and LAG at DPI.40. IL10 was significantly inhibited in ZFU.30-0.

Some differentially expressed genes between NDA and ZFU before infection are differentially expressed during infection

Because NDA was the breed with the highest total number of DEGs during infection, and ZFU and NDA
presented some distinct enriched biological pathways, we decided to compare DEGs detected in the
contrast NDA.0-ZFU.0 to those identified during infection in the corresponding within-breed contrasts.
Among DEGs identified in the NDA.0-ZFU.0 contrast, 82 (53%) were also found DE during infection either
within NDA or within ZFU. A heatmap performed on the logFC of these 82 genes in the within-breed
contrasts shows that the first level of clustering was due to the separation between ZFU and the other
breeds, and the second level separated the early infection time point (DPI.20-0) from the others (S5 Fig).
The most upregulated genes during infection were MARCO (macrophage receptor with collagenous
structure, significantly upregulated in all breeds except ZFU during infection), and ENSBTAG00000022715
(unannotated gene in Ensembl). Conversely, IGF2 (Insulin-like Growth Factor 2) was downregulated in all
breeds, but especially in NDA, whereas ANKEF1 was significantly downregulated in NDA only.
Among these genes, 55 were involved in canonical pathways or biological functions that were enriched
during infection (S10 Table) and, among them, 25 were present in at least ten functions. Top represented
DEGs were IL2RA, which was involved in 276 functions, IGF2 in 142, GATA1 in 95, GPB2 in 52, and MARCO
in 58 functions. Thirty-three canonical pathways were concerned, among which oxidative phosphorylation,
mitochondrial dysfunction, sirtuin pathway, and Th1 and Th2 pathways. 351 diseases and functions
contained DEGs in NDA.0-ZFU.0, top functions in term of number of genes were in particular associated
with cancer, cell death and survival, cell movement, cell-to-cell signaling and interaction, hematological
disease and inflammatory response. At last, 105 upstream regulators could influence the expression of
genes among which some were DE between ZFU and NDA before infection. Top enriched upstream
regulators included cytokines (i.e., INFG, IL2, TNF, CSF2, IL4), growth factors (TGFB1, HGF), TP53, Vegf and
MYC.

509   **Discussion**

Though bovine trypanotolerance has been described for more than a century (Pierre, 1906), the
biological bases of this phenotype remain poorly understood, due to the complexity of the trait that is
multigenic and multifactorial (dependent on environmental factors), and to the difficulty to perform
experimental infections on cattle, in comparison with model species (Morrison et al., 2016). In order to
increase knowledge on the physio-pathology of *T. congolense* infection in cattle, our study provides the
first whole blood transcriptome profiling using RNA-seq of five bovine breeds, including African taurine
shorthorn and longhorn, during an experimental trypanosome infection. The deep functional analysis of
the differentially expressed genes before and during infection identified new clues about trypanotolerance.
More precisely, our results confirmed and clarified some previous observations about the genes and the
main biological functions modulated during trypanosome infection in cattle (O'Gorman et al., 2006),
(O'Gorman et al., 2009). We observed a massive modification of the bovine transcriptome during *T.*
congolense infection in five West-African breeds, in accordance with the results of these previous
experiments that compared N'Dama and Boran breeds, notwithstanding that the experiments were done
on different cells (whole blood in the present study instead of Peripheral Blood Mononuclear Cells in
previous studies) using different techniques (RNA-seq in this study versus microarrays and/or qRT-PCR in

previous studies). In addition, the six genes (i.e., IL6, CCL2, IL10, IL1RN, IL8, NFKB1) DE in (O'Gorman et al.,
2006) and our study varied in the same direction, e.g., IL6 and NFKB1 were clearly upregulated, while IL1RN
was strongly downregulated in all breeds (except BOR). Among 23 DE genes representing various biological
processes and selected by (O'Gorman et al., 2009) for qRT-PCR validation, sixteen were also DE in the same
direction in our data set (eight upregulated and eight downregulated in both experiments, the remaining
seven being not significantly expressed in our data set or not annotated in Ensembl).

A core transcriptomic response in blood cells of all five West-African breeds infected by _T. congolense_

Our global analysis of gene expression of blood cells and of their associated biological functions
completed previous works by showing that blood cell transcriptomes responded globally similarly to
trypanosome infection whatever the breed. Indeed, all breeds exhibited a parallel shift in their gene
expression profiling. A majority of DEGs (62.3%) were shared among at least two breeds, and fold-changes
of all DEGs common to two or more breeds varied in the same direction, except one gene (i.e., SPARC).
This core cattle transcriptomic response to trypanosome infection is a reminder that cattle, whatever the
breed, experience parasitemia waves and show symptoms (e.g., anemia), albeit at varying intensities and
durations depending on the breed, as observed in experiments (Berthier et al., 2015), (Authie et al., 1993),
(Naessens et al., 2003), (Van der Waaij et al., 2003), or in the field (Trail et al., 1994). The functional
annotation of the DEGs shared between breeds during infection also allowed us to provide a global picture
of the impact of infection on whole blood gene transcription, by identifying hematopoiesis and immune
response, and metabolism as the main gene functions targeted by _T. congolense_ infection.

Modulation of hematopoiesis and immune response during infection

We detected in particular the modulation of gene transcription involved in hematopoiesis during
infection such as HBM, a highly upregulated DEG in all breeds. This increase of HBM expression suggests
that cattle responded quickly to cope with anemia induced by trypanosomes through an important supply
of reticulocytes from the bone marrow into the blood, while very low expression level of hemoglobin
subunits is normally found in the blood of healthy cattle (Correia et al., 2018). Expression of TFRC, an
important gene for erythropoiesis and also for T and B cell development and proliferation (Ned et al., 2003)
was also upregulated.
In addition, many top DEGs detected in all breeds are involved in immune response. Among them, five
(i.e., NR4A1, CCL22, IFI30, CTSZ and KYNU) out of the six genes significantly upregulated in all within-breeds
contrasts during infection are especially expressed in the monocyte lineage and/or dendritic cells (DC),
which initiate the immune response, presenting antigens to T lymphocytes (Wang et al., 2018), (Nowyhed
et al., 2015), (Vulcano et al., 2001), (Arunachalam et al., 2000), (Kos et al., 2005), (Harden et al., 2016),
(Obermajer et al., 2008). These cell types are known to be activated during trypanosomosis in cattle (Anosa
et al., 1999) and mice (Magez et al., 2007), and to play a decisive role in protection or pathology
(Bosschaerts et al., 2009). More specifically, NR4A1, a nuclear receptor with an inhibitory role in Th1 and
Th17 cell differentiation and in CD8 T cell expansion (Wang et al., 2018), (Nowyhed et al., 2015) could have
a protective role during trypanosomosis by modulating inflammation (Morias et al., 2015). As for IL17REL,
it was associated with inflammation and inflammatory diseases in human (Franke et al., 2010).
Additional common upregulated genes and enriched biological functions and upstream regulators
underlined the activation of innate immune response and macrophages, reported in cattle (Sileghem et al.,
1993), (Anosa et al., 1999), (Taiwo & Anosa, 2000), (Naessens, 2006) and mouse (Bosschaerts et al., 2009),
(Kuriakose et al., 2019). Indeed, important genes known to respond to pathogen stimuli and to trigger
immune response were found upregulated in all breeds during infection, such as MAPK11 and MAPK12
(Risco et al., 2012), NFKB1, up-regulated in bovine PBMC during trypanosomosis (O'Gorman et al., 2006),
and NFKB2. Activation of NF-kB complex was shown in endothelial cells and macrophages during _in vitro_
interactions with trypanosomes where it promoted a pro-inflammatory response (Ammar et al., 2013),
(Leppert et al., 2007), but NF-kBp50 encoded by NFKB1 was also able to stimulate the production of the
anti-inflammatory cytokine IL-10 (Bosschaerts et al., 2011). Interestingly, a strongly up-regulated common
DEG was ARG1, which codes for arginase-1, a key enzyme of arginine metabolism mainly expressed in

macrophages and associated with ornithine production and inhibition of NO production (Mills et al., 2000).
ARG1 activity is induced in macrophages of mice infected by *T. brucei* (Gobert et al., 2000) or *T. congolense*
(Noel et al., 2002) and its expression level depends on the genetic background of trypanotolerant and
susceptible mice (Duleu et al., 2004), (Kierstein et al., 2006). In mice infected by *T. congolense*, ARG1 is
involved in immune suppression induced by myeloid-derived suppressor cells that impair proliferation and
INF-g production of CD4 T cells (Onyilagha et al., 2018). Moreover, arginase activity has a direct positive
impact on trypanosome growth through ornithine production (Fairlamb & Cerami, 1992), (De Muylder et
al., 2013). In our data set, ARG1 was found to be highly expressed during infection, similarly to what is
found in mice, but with no straightforward differences between breeds. Activation of DC and macrophages
may also be highlighted by down-regulated genes, like DAB2 (a Clathrin Adaptor Protein) that acts as an
intrinsic negative regulator of immune function and inflammation in those cells (Ahmed et al., 2015), (Hung
et al., 2016).
Macrophage activation was also highlighted by the enrichment of upstream regulators like CSF1 (highly
enriched and activated in the common DEGs) and CSF2 (highly enriched and activated in within-breed
contrasts), which are essential cytokines enhancing macrophage and DC survival and activation (Becher et
al., 2016), IFNG, shown to be essential in the early-stage control of trypanosomosis in the mouse (Magez
et al., 2020), and TNF, which changed from a rather activated state at DPI.20-0 to an inhibition state at
DPI.40-0 recalling the switch from pro-inflammatory macrophages to anti-inflammatory macrophages
highlighted in tolerant mouse models (De Baetselier et al., 2001). Looking globally at the results, common
DEGs were rather associated with an anti-inflammatory balance (e.g., ARG1), as exemplified by the function
inflammation of central nervous system which was inferred as inhibited. Nevertheless, it is speculative to
infer pro-inflammatory or anti-inflammatory macrophages from our data set, as macrophage populations
are particularly plastic in space and time, and genes associated with both types were regulated (e.g., CCL22,
ARG1, IL1R1, PDC1LG2 regulation fitted with anti-inflammatory macrophages, while IL16, NR1D2, DAB2,
IRF5, NFKB fitted with pro-inflammatory macrophages according to (Arango Duque & Descoteaux, 2014),
(Jablonski et al., 2015), (Murray, 2017)).
The functional annotation of DEGs common to the breeds also highlighted the enrichment of the
activation of B cells (with several upregulated common DEGs highly expressed in B cells, like CD19, CD22,
CD40, CD72, CD79B, and CD180), which was previously reported in cattle infected by *T. congolense* (Pinder
et al., 1988), (Naessens & Williams, 1992), (Taylor et al., 1996), (O'Gorman et al., 2009), in human (Boda et
al., 2009), (Lejon et al., 2014) and mouse (Onyilagha et al., 2015). In addition, we noticed an activation of
B Cell Receptor Signaling, PI3K Signaling in B Lymphocytes and Quantity of B-2 cells in which common DEGs
were involved, and an enrichment of Quantity of B Lymphocytes in all breeds from DPI.30.
Common DEGs and their functions were also related to T-cell-mediated immunity. We noticed a
downregulation of genes mainly expressed in T cells, or NK cells, like CD2, CD3D, CD3E, CD3G, CD7, CD27,
CD40L, CD225, CD247, and ZAP70, of IL16, a chemoattractant for CD4 T cells (Wilson et al., 2004), and of
LIF (Leukemia Inhibitory Factor), a pleiotropic cytokine expressed by T cells, especially CD4 T cells in human
and regulatory T cells in mouse (Metcalfe, 2011). T cell functions were inferred as strongly inhibited (e.g.,
T cell development, activation of T lymphocytes, T cell migration), especially from DPI.30. Moreover, IL2,
an upstream regulator of the common DEGs, was significantly inhibited from DPI.30 in NDA, ZFU and LAG,
and from DPI.40 in BAO and BOR, as previously observed in PBMC from infected NDA and Boran Zebu
(O'Gorman et al., 2006). This global result is in line with an impairment of T cell functions as previously
observed in infected cattle ((Flynn & Sileghem, 1991), (Sileghem & Flynn, 1992b)), suggested in HAT (Boda
et al., 2009), and reported in mouse (Uzonna et al., 1998).
If we underlined an inhibition of T cell functions during the chronic phase of the disease, our results
also suggest an activation in the earlier phase, as illustrated by the detection of an early activation of the T
cell co-receptor CD3 as an upstream regulator. Interestingly, CD3 gene expression may be negatively
regulated during T cell activation (Paillard et al., 1990), (Badran et al., 2005), (Krishnan et al., 2001) likely
by a negative feed-back system necessary to control this process. In addition, another surprising result was
the inferred inhibited state of IL7 as an upstream regulator of the common DEGs during infection, though,
at the mRNA level, IL7 expression was significantly and sustainably upregulated and IL7RA was significantly
downregulated, which is expected under T cell activation (Alves et al., 2008). Two hypotheses may be
proposed. First, since trypanosomes are capable of long-term survival in the vertebrate host thanks to

antigenic variation (Jackson et al., 2012), (Matthews et al., 2015), a reduced expression of IL7RA and a
resulting hyporesponsiveness to IL7 could constitute a protective way to avoid T cell over-activation.
Second, IL7 upregulation could be a consequence of T cell lymphopenia (Fry et al., 2001) due to
trypanosome factors (Cnops et al., 2015). Whether this dynamic is due to an immunosuppression induced
by trypanosomes, or a component of the host-parasite equilibrium (Taylor, 1998) to avoid a deleterious
sustained inflammatory response cannot be answered.

Modulation of gene expression involved in metabolism during infection

Among the main functions and biological processes identified as significantly enriched in DEGs detected
in all breeds during infection, functions related to metabolism are clearly highlighted. First, regulation of
lipid metabolism was highly impacted during infection, and could suggest a direct effect of trypanosomes
or a modulation of the host immune response (Kay et al., 2006), (Fessler, 2015). Indeed, several up-
regulated genes in all breeds were involved in cholesterol synthesis (e.g., MVD, IDI1, FDFT1, SQLE, LSS,
CYP51A1), and accordingly, lipid metabolism functions (e.g., synthesis of cholesterol and sterol),
cholesterol biosynthesis and mevalonate pathways, and key upstream regulators of cholesterol synthesis
(SREBF1 and SREBF2, INSR) were strongly enriched and activated in the common DEGs data set, while
cholesterol, as an upstream regulator, was considered inhibited, thus promoting its own synthesis. If
previous studies suggested the involvement of lipid metabolism regulation in the physiopathology of AAT
in mice (Kierstein et al., 2006), in cattle and goat infected by *T. congolense* (Traore-Leroux et al., 1987),
(Meade et al., 2009), (Rajavel et al., 2020), (Ndoutamia et al., 2002), this is the first time that this
involvement has been detected to such an extent. Modification in lipid metabolism could be due to a direct
effect of trypanosomes that are able to internalize lipoproteins (Green et al., 2003), but it could also likely
reflect an activation of the inflammatory response as suggested by (Bouvier-Muller et al., 2017).
Second, the change observed in our study in the energy production of the cells probably reflects the
modification of immune cell energetic metabolism required to support cell growth, proliferation and
effector functions (Donnelly & Finlay, 2015) during infection by *T. congolense*. The tricarboxylic acid (TCA)
cycle, which is a hub for generating energy and building blocks for macromolecule synthesis as well as
releasing intra-cellular signaling molecules (Martinez-Reyes & Chandel, 2020), was indeed highly enriched
and activated in the common DEGs that we detected in our study and in some within-breeds contrasts.
Besides, gluconeogenesis and glycolysis were also assessed as activated during infection. These two
opposite pathways shared many genes whose corresponding proteins may catalyze reactions in two
directions. Nevertheless, since PKLR, involved in a key step for glycolysis, was upregulated in all the breeds
during infection, and that FBP1, responsible for final steps for gluconeogenesis (Lebigot et al., 2015), was
downregulated, we can suppose that glycolysis was actually the activated pathway during infection. These
results are concordant with the absence of gluconeogenesis in blood cells and with the interdependency
of glycolysis and TCA cycle, glycolysis supplying TCA cycle with pyruvate and providing molecules for
synthetizing nucleotides, glycerol and amino acids (Donnelly & Finlay, 2015).

Breed-specific whole blood transcriptomic responses

A focus on how each breed reacted to infection allowed observing some variations in their
transcriptome, in regard to the enrichment and intensity of activation or inhibition of some biological
functions. As previously reported by (O'Gorman et al., 2009), the blood transcriptome of NDA cattle, the
trypanotolerant breed of reference (Murray et al., 1984), (Hanotte et al., 2003), seemed to respond earlier
and more intensely to infection, with a higher number of detected DEGs in this breed during our
experiment in comparison with the other breeds, except at DPI.30-0 when LAG had a little more DEGs. In
addition, NDA harbored a more intense and earlier activation of several upstream regulators, notably
involved in immune response, in comparison with the other breeds, as exemplified by CSF2, the top
activated upstream regulator in NDA at DPI.20-0, that is a key cytokine produced by various cells during an
infection or an inflammation and involved in monocyte, macrophage, granulocyte and DC functions (Van
de Laar et al., 2012), (Hamilton & Achuthan, 2013). The highly estimated activation state of CSF2 in NDA in
our study could reflect an earlier and greater activation of macrophages in this breed, consistent with the

earlier release of co-stimulatory cytokines by NDA monocytes in comparison with Boran Zebu monocytes (Sileghem et al., 1993) and the hypothesis of an earlier pro-inflammatory response in PBMC from NDA (O'Gorman et al., 2006). PTGER2, the receptor of prostaglandin E2, an inflammation mediator (Kawahara et al., 2015) associated with a pro-inflammation profile in a mouse model of Chagas disease caused by *T. cruzi* (Guerrero et al., 2015), was also considered as highly and precociously activated. In NDA.30-0 and NDA.40-0 contrasts, MYC, activated in M2 macrophages (Pello et al., 2012), (Jablonski et al., 2015), and critical in T cell proliferation and growth following activation (Wang et al., 2011), was the most activated upstream regulator. We could thus hypothesize that trypanotolerance in NDA could be linked to the precocity and the chronology of activation state of cells, which would allow an efficient immune response while avoiding immune disorders (Vincendeau & Bouteille, 2006).

In addition, biological processes related to cell cycle and DNA metabolism (e.g., diseases and functions like segregation of chromosomes, metabolism of DNA; canonical pathways like cyclins and cell cycle regulation; upstream regulators like E2F1, FOXM1), were strongly enriched and activated in blood cells of NDA breed at DPI.20-0 and DPI.30-0, and explained the enrichment of functions linked to cancer. These results suggest an early activation of division and proliferation of some blood cell types in NDA (e.g., lymphocytes, (Naessens & Williams, 1992), (Naessens et al., 2003)). Curiously, like in all breeds, the lymphopoiesis function was assessed as strongly inhibited from DPI.30-0. These results could seem paradoxical but the sets of genes associated with lymphopoiesis on the one hand, and with cell cycle on the other hand, are different, the former function comprising DEGs encoding T cells surface antigens, cytokines receptors and signal transducers, and the latter comprising DEGs involved in mitosis and centromere formation.

Lastly, NDA displayed a particularly important shift in energetic metabolism of blood cells, whose fine tuning is associated with cell type and fate (O'Neill et al., 2016). Indeed, a key feature of NDA canonical pathways was the strong enrichment of oxidative phosphorylation and mitochondrial dysfunction, the former being highly activated such as TCA cycle and glycolysis. The fact that these three pathways were inferred as activated corroborates the presence of diverse immune cell populations and/or B cell activation. Indeed, glycolysis is rather associated with pro-inflammatory macrophages (Mills et al., 2016), activated DCs or effector T cells (McGettrick & O'Neill, 2013), while oxidative phosphorylation is rather associated with anti-inflammatory macrophages, Tregs or memory T cells (McGettrick & O'Neill, 2013), (Mills et al., 2016), and both seem increased in activated B cells (Caro-Maldonado et al., 2014). However, the strongest activation of oxidative phosphorylation observed in NDA could also be associated with a predominance of an anti-inflammatory component (Pearce & Pearce, 2013) during the chronic phase of infection at 30 and 40 DPI. Regulation of metabolic processes was also highlighted by the enrichment of the sirtuin signaling pathway, which was particularly significantly inhibited in NDA. This pathway comprises several proteins involved in ubiquitous processes and especially in a coupling between metabolic and stress factors and inflammatory response (Loftus & Finlay, 2016). In the same way, the most inhibited upstream regulator in NDA was the RICTOR protein that belongs to mTORC2 complex. The latter regulates cell metabolism and is involved in numerous functions, including development (Guertin et al., 2006), insulin signaling by promoting lipogenesis and glycogen synthesis (Yoon, 2017), and immune cell functions.

ZFU, which had the most pronounced anemia, the most durable parasitemia, and the lowest leukocytosis among the five breeds studied (Berthier et al., 2015), was characterized by a high enrichment and a strong activation/inhibition of biological functions, pathways and upstream regulators linked to lipid metabolism from the beginning of infection (i.e., synthesis of cholesterol, superpathway of cholesterol biosynthesis), which could interact with immune response. This was exemplified by SREBF2, a major transcription factor involved in cholesterol metabolism (Bommer & MacDougald, 2011), strongly activated as an upstream regulator, and also continuously upregulated in ZFU exclusively. In addition, the LXR/RXR activation pathway was found significantly enriched in ZFU but not in NDA nor LAG. Interestingly, (Morrison et al., 2010) identified this pathway as differentially enriched between mice infected by two *T. brucei* strains provoking distinct phenotypes. This pathway, notably expressed in hepatocytes and macrophages, is involved in lipid metabolism and innate immunity in macrophages (Joseph et al., 2004), (Ahsan et al., 2018), with rather an anti-inflammatory balance (Schulman, 2017). In the same direction, the significant inhibition of the inflammasome pathway at DPI.40 in ZFU only tended to show an inhibition of the inflammatory response at the chronic stage of infection in this breed (Zamboni & Lima-Junior, 2015).

LAG, whose trypanotolerance has been demonstrated (Berthier et al., 2015) (i.e., mild anemia and quick
recovery), showed a similar whole-blood transcriptome profile to NDA during the experiment (Fig 2). If LAG
had less DEGs than NDA, it responded relatively intensely to infection, especially at DPI.30-0, and most
information about functional analyses was found at this date. In LAG, like in NDA, top activated canonical
pathways were linked to cell energy production, i.e., oxidative phosphorylation and TCA Cycle,
mitochondrial dysfunction was enriched, and diseases and functions linked to cell cycle functions (G2/M
phase transition, Interphase, Segregation of chromosomes) were activated. Superpathway of cholesterol
biosynthesis was also activated, but, unlike the other breeds, not any disease and function linked to
cholesterol or lipid metabolism was significant. Top upstream regulators were also shared with NDA, as
CSF2, the most enriched and activated upstream regulator at DPI.20-0, and HGF, Vegf, MYC, INSR and TCF3.
More specific to LAG response, GATA3 and TBX21, reportedly expressed in an opposite way and
respectively associated with Th2 and Th1 cells (Chakir et al., 2003), were significantly detected as upstream
regulators in LAG.40-0. GATA3, down regulated in all breeds at the mRNA level, was assessed as inhibited
in LAG, and TBX21, upregulated in LAG only, as activated. TBX21 was also associated with NK cell
development and effector functions (Deng et al., 2015). In LAG, a pro-inflammatory activation of the Th1
cell subset or NK could be considered, but results from the diseases and functions analysis showed rather
an inhibition of T cells. Nevertheless, TBX21 being also expressed in B-cells precursors, contrary to GATA3
(Harashima et al., 2005) and being required in B cells for IFNG dependent switching in IgG2a production
(Mohr et al., 2010), other biological mechanisms could be involved.
The BAO breed had an unexpected global transcriptomic response in comparison with its
trypanotolerant status, given that this breed did not present a significantly different level of anemia from
those of NDA and LAG during infection (Berthier et al., 2015). Indeed, the differential expression between
the different times after infection and DPI.0 were smaller in BAO than in other breeds, as shown by the
PCA visualization (Fig 2) and the small numbers of DEGs detected in BAO during the infection process. This
discrepancy between its transcriptomic response and its trypanotolerant status was reflected at the
functional analysis. Indeed, the transcriptomic response in BAO showed similar features to that of ZFU
regarding DEGs involved in cholesterol and lipid metabolism, while it resembled NDA and LAG regarding
TCA cycle activation, control of Ig quantities and B Cell Receptor Signaling.
At last, BOR, an admixed breed between AFZ and AFT (Flori et al., 2014), displayed an expected global
transcriptomic response according to its intermediate phenotype between tolerant and susceptible breeds
concerning anemia, parasitemia and leucocyte counts (Berthier et al., 2015). The functional analyses of
DEGs did not reveal specific responses in BOR, except the activation in BOR.40-0 of RNA transcription.
Significant functions were indeed shared with other breeds, such as the M phase and the segregation of
chromosomes at 20 DPI, strongly enriched in NDA, or the canonical pathways linked to cholesterol
metabolism at DPI.30-0 more enriched in ZFU.

Baseline transcriptomic differences between trypanotolerant and susceptible breeds point to genes that
could influence the outcome of infection

The cross-referencing of the DEG between breeds before infection and within breeds during infection
could provide additional information on gene expression and functions associated with trypanotolerance.
Though the PCA supports the hypothesis that a majority of the differences in gene expression before
infection results from demographic history (see Fig 2), some differences could be adaptive (Whitehead &
Crawford, 2006). Indeed, eighty-two genes were DE between NDA and ZFU before infection and also
responded to infection in NDA and/or ZFU. Many of these genes are reported to be involved in functions
highlighted previously, e.g., immune response and metabolism. One hypothesis is that differences in basal
expression of these genes could underline different proportions of cell types or different states of cell
activation between breeds, which could play a role in subsequent pathogenic processes caused by
trypanosomes.
As illustration, several genes are known to be associated with macrophage functions. Among them,
MARCO, whose expression level was higher in ZFU than in NDA and LAG before infection, but was
subsequently significantly upregulated in all breeds except in ZFU, is a pattern recognition receptor on
macrophages surface involved in phagocytosis of various pathogens and the subsequent enhancement of

immune response and chemokines expression (Arredouani et al., 2004), (Bowdish et al., 2009), (Xu et al., 2017). MMD (Monocyte to macrophage differentiation-associated) is a gene expressed during monocyte differentiation and macrophage activation (Liu et al., 2012), and it was similarly upregulated in ZFU.0 in comparison to NDA.0, and upregulated during the infection in NDA, LAG and BOR. SLC11A1 (syn. Nramp1) is also a macrophage gene, and was upregulated in ZFU.0 versus NDA.0, but it was then downregulated in ZFU during infection. The corresponding protein regulates iron homeostasis in macrophages and gene variants are associated with disease susceptibility or resistance (Archer et al., 2015). In addition, IL2RA, which is strongly expressed by Tregs, effector T cells (Banham et al., 2006), NK cells (Esin et al., 2013), (Hamilton et al., 2017), and granulocytes in cattle (Zoldan et al., 2014) was more expressed in ZFU.0 in comparison with the other breeds, and downregulated during infection and particularly in ZFU. IL2RA (syn. CD25) has been the target of research in bovine trypanosomosis, where its impairment was noticed in lymph nodes (Sileghem & Flynn, 1992a). In mice, injection of anti-CD25 antibodies before experimental infections with *T. congolense* led to discordant results related to protection or pathogeny (Okwor et al., 2012), (Guilliams et al., 2007). Other genes harboring basal differences in their expression level were associated with metabolism regulation. For instance, IGF2, which was upregulated in NDA.0 versus ZFU.0 and was downregulated during the infection, codes for a peptide hormone that is involved in metabolism, tissue development and maintenance and is downregulated during under-nutrition or hypoglycemia (Livingstone & Borai, 2014).

In summary, our study provides the first transcriptome profiling of whole blood cells of five West-African bovine breeds during a trypanosome infection using RNA-seq. In order to identify potential similarities among African taurine cattle, overlooked trypanotolerant breeds (i.e., Lagune and Baoule) were considered in addition to N'Dama, which accounts for the majority of the studies about cattle trypanotolerance. We observed that trypanosome infection due to *T. congolense* has a major impact on cattle blood transcriptome, whatever the breed and we provided a global transcriptomic picture of infection. In accordance with previous results, a strong regulation of the immune system functions with an early activation of innate immune response, followed by an activation of humoral response and an inhibition of T cell functions at the chronic stage of infection were detected in all breeds.

Most importantly, we highlighted overlooked features, as a strong modification in lipid metabolism. We mainly noticed an early regulation of the immune response in NDA, associated with a strong activation of energy production by the cell, a strong enrichment and activation of oxidative phosphorylation in NDA and LAG, and an activation of the TCA cycle in AFT breeds, which was not highlighted in ZFU. These differences in cellular energy could be linked in AFT to better functions of some cell types, like M2 macrophages, memory T cells or activated B cells, and could represent a key to decipher trypanotolerance. If some DEGs, functions and biological pathways were shared between AFT breeds during infection, our results highlight also differences in gene expression dynamics in these three trypanotolerant breeds (as exemplified by the singular transcriptomic profile of Baoule). This suggests that AFT breeds, although subjected to the same selective pressure caused by trypanosomes, may have developed different adaptation mechanisms. In addition, the trypanosusceptible breed ZFU presented several canonical pathways linked to inflammation inhibited from DPI.30, which was not observed in the other breeds, and the strongest modification in lipid metabolism regulation. It would be worth exploring other African zebu breeds to confirm if this observation is a global feature of indicine breeds. Some genes, known to be involved in immune response or metabolism, were differently expressed between breeds before infection and within breeds during infection and raise the hypothesis that basal differences between breeds could impact the outcome in infection.

Finally, our study provided new and valuable data to contribute to a better knowledge of African livestock genomics (Kemp, 2019), (Kim et al., 2020), and to decipher the pathogenic process in bovine trypanosomosis due to *T. congolense*. A comparable experiment using *T vivax* or *T. brucei brucei* would be worthwhile to verify whether host responses are similar regardless of the infecting species or not. Overall, interactions between immune response and metabolism deserve to be deeply explored in cattle in order to improve preventive and curative measures of AAT and also other infectious diseases.

Acknowledgements

This work was supported by the CIRAD - UMR AGAP HPC Data Center of the South Green Bioinformatics platform (http://www.southgreen.fr/). We thank Jean Nakhle for having corrected the English version of the manuscript.

Funding

This work was supported by the ANR grant n°2011 JSV6 001 01 (http://www.agence-nationalerecherche.fr/) and by the LabEx ParaFrap (ANR-11-LABX-0024). MGX acknowledges financial support from France Génomique National infrastructure, funded as part of "Investissement d'avenir" program managed by Agence Nationale pour la Recherche (contract ANR-10-INBS-09). The PhD grant of Moana Peylhard was supported by CIRAD PhD grant program and IRD.

Conflict of interest disclosure

The authors declare no conflicts of interest relating to the content of this article. Sophie Thévenon is a recommender for PCI Infections.

Data, script and code availability

Raw sequences data (fastq, count matrix, experimental design) will be publicly available at GEO (https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE197108 upon publication.

Scripts and intermediate tables (DEG analysis, input tables for IPA® analyses, output tables from IPA® analyses) are publicly available in Cirad Dataverse under https://doi.org/10.18167/DVN1/L9SHAX. SNP genotypes of the experimental cattle are publicly available in Cirad Dataverse under https://doi.org/10.18167/DVN1/APTZOC and in WIDDE (http://widde.toulouse.inra.fr/widde/).

Supporting information availability

Tables and figures are in the supplementary tables and figures files. Legends are displayed at the bottom of each tables and figures.

S1 Table. Summary information on sequencing and mapping results.

S2 Table. LogFC and FDR of the 13,107 genes for the 15 within-breed contrasts during infection.

S3 Table. LogFC and FDR of the 13,107 genes for the 4 between-breed contrasts at DPI.0.

S4 Table. Upstream regulators significantly enriched in the between-breed contrasts at DPI.0

S5 Table. Diseases and functions enriched in the common DEGs in the within-breed contrasts.

S6 Table. Upstream regulators enriched in the common DEGs in the within-breed contrasts.

S7 Table. Cross table of the enriched diseases and functions in the 15 within-breed contrasts.

S8 Table. Cross table of the canonical pathways in the 15 within-breed contrasts**.**

S9 Table. Cross table of the upstream regulators in the 15 within-breed contrasts.

S10 Table. Cross table between the functional categories enriched during infection and the genes differentially expressed in the contrast NDA.0-ZFU.0.

S1 Figure. Proportion of uniquely mapped reads on the trypanosome genome depending on time points.

S2 Figure. Principal components analysis of 120 cattle RNA-seq libraries based on normalized genes counts.

S3 Figure. Venn diagram showing the intersection of genes identified as DE in the within-breed contrasts.

S4 Figure. Venn diagram showing the intersection of genes identified as DE in the between-breed contrasts.

S5 Figure. Heatmap on the logFC of a subset of 82 DE genes.

References

Ahmed MS, Byeon SE, Jeong Y, Miah MA, Salahuddin M, Lee Y, Park SS, Bae YS (2015) Dab2, a negative regulator of DC immunogenicity, is an attractive molecular target for DC-based immunotherapy. *Oncoimmunology, 4*, e984550. https://doi.org/10.4161/2162402X.2014.984550

Ahsan F, Maertzdorf J, Guhlich-Bornhof U, Kaufmann SHE, Moura-Alves P (2018) IL-36/LXR axis modulates cholesterol metabolism and immune defense to Mycobacterium tuberculosis. *Sci Rep, 8*, 1520. https://doi.org/10.1038/s41598-018-19476-x

Akol GW, Authie E, Pinder M, Moloo SK, Roelants GE, Murray M (1986) Susceptibility and immune responses of zebu and taurine cattle of West Africa to infection with *Trypanosoma congolense* transmitted by *Glossina morsitans centralis*. *Vet Immunol Immunopathol, 11*, 361-373. https://doi.org/10.1016/0165-2427(86)90038-3

Alsan M (2015) The Effect of the TseTse Fly on African Development. *American Economic Review, 105*, 382-410. https://doi.org/10.1257/aer.20130604

Alvarez I, Perez-Pardal L, Traore A, Fernandez I, Goyache F (2015) Lack of haplotype structuring for two candidate genes for trypanotolerance in cattle. *J Anim Breed Genet*. https://doi.org/10.1111/jbg.12181

Alvarez I, Perez-Pardal L, Traore A, Fernandez I, Goyache F (2016) African Cattle do not Carry Unique Mutations on the Exon 9 of the ARHGAP15 Gene. *Anim Biotechnol, 27*, 9-12. https://doi.org/10.1080/10495398.2015.1053606

Alves NL, van Leeuwen EM, Derks IA, van Lier RA (2008) Differential regulation of human IL-7 receptor alpha expression by IL-7 and TCR signaling. *J Immunol, 180*, 5201-5210. https://doi.org/10.4049/jimmunol.180.8.5201

Amene BM, Chime AB, Anika SM (1991) The production performance of imported friesian cattle under heavy *Trypanosoma* challenge in a rain forest zone of Nigeria. *Br. vet. J., 147*, 274-282. https://doi.org/10.1016/0007-1935(91)90052-O

Ammar Z, Plazolles N, Baltz T, Coustou V (2013) Identification of trans-sialidases as a common mediator of endothelial cell activation by African trypanosomes. *PLoS Pathog, 9*, e1003710. https://doi.org/10.1371/journal.ppat.1003710

Anosa VO, Logan-Henfrey LL, Wells CW (1999) The role of the bone marrow in bovine trypanotolerance II. Macrophage function in *Trypanosoma congolense*-infected cattle. *Comparative Haematology International, 9*, 208-219. https://doi.org/10.1007/BF02585507

Arango Duque G, Descoteaux A (2014) Macrophage cytokines: involvement in immunity and infectious diseases. *Front Immunol, 5*, 491. https://doi.org/10.3389/fimmu.2014.00491

Archer NS, Nassif NT, O'Brien BA (2015) Genetic variants of SLC11A1 are associated with both autoimmune and infectious diseases: systematic review and meta-analysis. *Genes Immun, 16*, 275-283. https://doi.org/10.1038/gene.2015.8

Arredouani M, Yang Z, Ning Y, Qin G, Soininen R, Tryggvason K, Kobzik L (2004) The scavenger receptor MARCO is required for lung defense against pneumococcal pneumonia and inhaled particles. *J Exp Med, 200*, 267-272. https://doi.org/10.1084/jem.20040731

Arunachalam B, Phan UT, Geuze HJ, Cresswell P (2000) Enzymatic reduction of disulfide bonds in lysosomes: characterization of a gamma-interferon-inducible lysosomal thiol reductase (GILT). *Proc Natl Acad Sci U S A, 97*, 745-750. https://doi.org/10.1073/pnas.97.2.745

Authie E, Muteti DK, Williams DJ (1993) Antibody responses to invariant antigens of *Trypanosoma congolense* in cattle of differing susceptibility to trypanosomiasis. *Parasite Immunol, 15*, 101-111. https://doi.org/10.1111/j.1365-3024.1993.tb00589.x

Badran BM, Kunstman K, Stanton J, Moschitta M, Zerghe A, Akl H, Burny A, Wolinsky SM, Willard-Gallo KE (2005) Transcriptional regulation of the human CD3 gamma gene: the TATA-less CD3 gamma promoter functions via an initiator and contiguous Sp-binding elements. *J Immunol, 174*, 6238-6249. https://doi.org/10.4049/jimmunol.174.10.6238

Banham AH, Powrie FM, Suri-Payer E (2006) FOXP3+ regulatory T cells: Current controversies and future perspectives. *Eur J Immunol, 36*, 2832-2836. https://doi.org/10.1002/eji.200636459

Becher B, Tugues S, Greter M (2016) GM-CSF: From Growth Factor to Central Mediator of Tissue Inflammation. *Immunity,* **45**, 963-973. https://doi.org/10.1016/j.immuni.2016.10.026

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol,* **57**, 289-300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Berthier D, Peylhard M, Dayo GK, Flori L, Sylla S, Bolly S, Sakande H, Chantal I, Thevenon S (2015) A comparison of phenotypic traits related to trypanotolerance in five west african cattle breeds highlights the value of shorthorn taurine breeds. *PLoS One,* **10**, e0126498. https://doi.org/10.1371/journal.pone.0126498

Boda C, Courtioux B, Roques P, Pervieux L, Vatunga G, Josenando T, Ayenengoye CR, Bouteille B, Jauberteau MO, Bisser S (2009) Immunophenotypic lymphocyte profiles in human african trypanosomiasis. *PLoS One,* **4**, e6184. https://doi.org/10.1371/journal.pone.0006184

Bommer GT, MacDougald OA (2011) Regulation of lipid homeostasis by the bifunctional SREBF2-miR33a locus. *Cell Metab,* **13**, 241-247. https://doi.org/10.1016/j.cmet.2011.02.004

Bosschaerts T, Guilliams M, Stijlemans B, De Baetselier P, Beschin A (2009) Understanding the role of monocytic cells in liver inflammation using parasite infection as a model. *Immunobiology,* **214**, 737-747. https://doi.org/10.1016/j.imbio.2009.06.010

Bosschaerts T, Morias Y, Stijlemans B, Herin M, Porta C, Sica A, Mantovani A, De Baetselier P, Beschin A (2011) IL-10 limits production of pathogenic TNF by M1 myeloid cells through induction of nuclear NF-kappaB p50 member in *Trypanosoma congolense* infection-resistant C57BL/6 mice. *Eur J Immunol,* **41**, 3270-3280. https://doi.org/10.1002/eji.201041307

Bouvier-Muller J, Allain C, Tabouret G, Enjalbert F, Portes D, Noirot C, Rupp R, Foucras G (2017) Whole blood transcriptome analysis reveals potential competition in metabolic pathways between negative energy balance and response to inflammatory challenge. *Scientific reports,* **7**, 2379

Bouyer J, Bouyer F, Donadeu M, Rowan T, Napier G (2013) Community- and farmer-based management of animal African trypanosomosis in cattle. *Trends Parasitol,* **29**, 519-522. https://doi.org/10.1016/j.pt.2013.08.003

Bowdish DM, Sakamoto K, Kim MJ, Kroos M, Mukhopadhyay S, Leifer CA, Tryggvason K, Gordon S, Russell DG (2009) MARCO, TLR2, and CD14 are required for macrophage cytokine responses to mycobacterial trehalose dimycolate and Mycobacterium tuberculosis. *PLoS Pathog,* **5**, e1000474. https://doi.org/10.1371/journal.ppat.1000474

Bradley DG, MacHugh DE, Cunningham P, Loftus RT (1996) Mitochondrial diversity and the origins of African and European cattle. *Proc Natl Acad Sci U S A,* **93**, 5131-5135. https://doi.org/10.1073/pnas.93.10.5131

Budd LT (1999) *DFID-funded tsetse and trypanosomiasis research and development since 1980.* Economic Analysis (Vol. 2). Department of International Development, UK, London.

Caro-Maldonado A, Wang R, Nichols AG, Kuraoka M, Milasta S, Sun LD, Gavin AL, Abel ED, Kelsoe G, Green DR, Rathmell JC (2014) Metabolic reprogramming is required for antibody production that is suppressed in anergic but exaggerated in chronically BAFF-exposed B cells. *J Immunol,* **192**, 3626-3636. https://doi.org/10.4049/jimmunol.1302062

Chakir H, Wang H, Lefebvre DE, Webb J, Scott FW (2003) T-bet/GATA-3 ratio as a measure of the Th1/Th2 cytokine profile in mixed cell populations: predominant role of GATA-3. *J Immunol Methods,* **278**, 157-169. https://doi.org/10.1016/S0022-1759(03)00200-X

Chen Y, Lun ATL, Smyth GK (2014) Differential expression analysis of complex RNA-seq experiments using edgeR. In: *Statistical Analysis of Next Generation Sequence Data* (ed. Nettleton SDaDS), pp. 51-74. Springer, New York.

CIPEA (1979) *Le bétail trypanotolérant d'Afrique occidentale et centrale* (Vol. 2), Addis Abeba, Ethiopie.

Cnops J, De Trez C, Stijlemans B, Keirsse J, Kauffmann F, Barkhuizen M, Keeton R, Boon L, Brombacher F, Magez S (2015) NK-, NKT- and CD8-Derived IFNgamma drives myeloid cell activation and erythrophagocytosis, resulting in trypanosomosis-associated acute anemia. *PLoS Pathog,* **11**, e1004964. https://doi.org/10.1371/journal.ppat.1004964

Correia CN, McLoughlin KE, Nalpas NC, Magee DA, Browne JA, Rue-Albrecht K, Gordon SV, MacHugh DE (2018) RNA Sequencing (RNA-Seq) reveals extremely low levels of reticulocyte-derived globin gene

transcripts in peripheral blood from horses (*Equus caballus*) and cattle (*Bos taurus*). *Front Genet,* **9**, 278. https://doi.org/10.3389/fgene.2018.00278

De Baetselier PD, Namangala B, Noel W, Brys L, Pays E, Beschin A (2001) Alternative versus classical macrophage activation during experimental African trypanosomosis. *Int J Parasitol,* **31**, 575-587. https://doi.org/10.1016/S0020-7519(01)00170-9

De Muylder G, Daulouede S, Lecordier L, Uzureau P, Morias Y, Van Den Abbeele J, Caljon G, Herin M, Holzmuller P, Semballa S, Courtois P, Vanhamme L, Stijlemans B, De Baetselier P, Barrett MP, Barlow JL, McKenzie AN, Barron L, Wynn TA, Beschin A, Vincendeau P, Pays E (2013) A *Trypanosoma brucei* kinesin heavy chain promotes parasite growth by triggering host arginase activity. *PLoS Pathog,* **9**, e1003731. https://doi.org/10.1371/journal.ppat.1003731

Deng Y, Kerdiles Y, Chu J, Yuan S, Wang Y, Chen X, Mao H, Zhang L, Zhang J, Hughes T, Deng Y, Zhang Q, Wang F, Zou X, Liu CG, Freud AG, Li X, Caligiuri MA, Vivier E, Yu J (2015) Transcription factor Foxo1 is a negative regulator of natural killer cell maturation and function. *Immunity,* **42**, 457-470. https://doi.org/10.1016/j.immuni.2015.02.006

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics,* **29**, 15-21. https://doi.org/10.1093/bioinformatics/bts635

Doko A, Verhulst A, Pandey VS, Van der Stuyft P (1997) Trypanosomose expérimentale à *Trypanosoma brucei brucei* chez les taurins Holsteins et les zébus Bororo blancs. *Revue Elev. Méd. vét. Pays trop.,* **50**, 23-28. https://doi.org/10.19182/remvt.9597

Donnelly RP, Finlay DK (2015) Glucose, glycolysis and lymphocyte responses. *Mol Immunol,* **68**, 513-519. https://doi.org/10.1016/j.molimm.2015.07.034

Duleu S, Vincendeau P, Courtois P, Semballa S, Lagroye I, Daulouede S, Boucher JL, Wilson KT, Veyret B, Gobert AP (2004) Mouse strain susceptibility to trypanosome infection: an arginase-dependent effect. *J Immunol,* **172**, 6298-6303. https://doi.org/10.4049/jimmunol.172.10.6298

Esin S, Counoupas C, Aulicino A, Brancatisano FL, Maisetta G, Bottai D, Di Luca M, Florio W, Campa M, Batoni G (2013) Interaction of Mycobacterium tuberculosis cell wall components with the human natural killer cell receptors NKp44 and Toll-like receptor 2. *Scand J Immunol,* **77**, 460-469. https://doi.org/10.1111/sji.12052

Fairlamb AH, Cerami A (1992) Metabolism and functions of trypanothione in the Kinetoplastida. *Annu Rev Microbiol,* **46**, 695-729. https://doi.org/10.1146/annurev.mi.46.100192.003403

Fessler MB (2015) Regulation of Adaptive Immunity in Health and Disease by Cholesterol Metabolism. *Curr Allergy Asthma Rep,* **15**, 48. https://doi.org/10.1007/s11882-015-0548-7

Flori L, Thevenon S, Dayo GK, Senou M, Sylla S, Berthier D, Moazami-Goudarzi K, Gautier M (2014) Adaptive admixture in the West African bovine hybrid zone: insight from the Borgou population. *Mol Ecol,* **23**, 3241-3257. https://doi.org/10.1111/mec.12816

Flynn JN, Sileghem M (1991) The role of the macrophage in induction of immunosuppression in *Trypanosoma congolense*-infected cattle. *Immunology,* **74**, 310-316

Franke A, Balschun T, Sina C, Ellinghaus D, Hasler R, Mayr G, Albrecht M, Wittig M, Buchert E, Nikolaus S, Gieger C, Wichmann HE, Sventoraityte J, Kupcinskas L, Onnie CM, Gazouli M, Anagnou NP, Strachan D, McArdle WL, Mathew CG, Rutgeerts P, Vermeire S, Vatn MH, group Is, Krawczak M, Rosenstiel P, Karlsen TH, Schreiber S (2010) Genome-wide association study for ulcerative colitis identifies risk loci at 7q22 and 22q13 (IL17REL). *Nat Genet,* **42**, 292-294. https://doi.org/10.1038/ng.553

Fry TJ, Christensen BL, Komschlies KL, Gress RE, Mackall CL (2001) Interleukin-7 restores immunity in athymic T-cell-depleted hosts. *Blood,* **97**, 1525-1533. https://doi.org/10.1182/blood.v97.6.1525

Gobert AP, Daulouede S, Lepoivre M, Boucher JL, Bouteille B, Buguet A, Cespuglio R, Veyret B, Vincendeau P (2000) L-Arginine availability modulates local nitric oxide production and parasite killing in experimental trypanosomiasis. *Infect Immun,* **68**, 4653-4657. https://doi.org/10.1128/IAI.68.8.4653-4657.2000

Green HP, Del Pilar Molina Portela M, St Jean EN, Lugli EB, Raper J (2003) Evidence for a *Trypanosoma brucei* lipoprotein scavenger receptor. *J Biol Chem,* **278**, 422-427. https://doi.org/10.1074/jbc.M207215200

Guerrero NA, Camacho M, Vila L, Iniguez MA, Chillon-Marinas C, Cuervo H, Poveda C, Fresno M, Girones N (2015) Cyclooxygenase-2 and Prostaglandin E2 Signaling through Prostaglandin Receptor EP-2 Favor the Development of Myocarditis during Acute Trypanosoma cruzi Infection. *PLoS Negl Trop Dis,* **9**, e0004025. https://doi.org/10.1371/journal.pntd.0004025

Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, Brown M, Fitzgerald KJ, Sabatini DM (2006) Ablation in mice of the mTORC components raptor, rictor, or mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not S6K1. *Dev Cell,* **11**, 859-871. https://doi.org/10.1016/j.devcel.2006.10.007

Guilliams M, Oldenhove G, Noel W, Herin M, Brys L, Loi P, Flamand V, Moser M, De Baetselier P, Beschin A (2007) African trypanosomiasis: naturally occurring regulatory T cells favor trypanotolerance by limiting pathology associated with sustained type 1 inflammation. *J Immunol,* **179**, 2748-2757. https://doi.org/10.4049/jimmunol.179.5.2748

Hamilton CA, Mahan S, Bell CR, Villarreal-Ramos B, Charleston B, Entrican G, Hope JC (2017) Frequency and phenotype of natural killer cells and natural killer cell subsets in bovine lymphoid compartments and blood. *Immunology,* **151**, 89-97. https://doi.org/10.1111/imm.12708

Hamilton JA, Achuthan A (2013) Colony stimulating factors and myeloid cell biology in health and disease. *Trends Immunol,* **34**, 81-89. https://doi.org/10.1016/j.it.2012.08.006

Hanotte O, Bradley DG, Ochieng JW, Verjee Y, Hill EW, Rege JE (2002) African pastoralism: genetic imprints of origins and migrations. *Science,* **296**, 336-339. https://doi.org/10.1126/science.1069878

Hanotte O, Ronin Y, Agaba M, Nilsson P, Gelhaus A, Horstmann R, Sugimoto Y, Kemp S, Gibson J, Korol A, Soller M, Teale A (2003) Mapping of quantitative trait loci controlling trypanotolerance in a cross of tolerant West African N'Dama and susceptible East African Boran cattle. *Proc Natl Acad Sci U S A,* **100**, 7443-7448. https://doi.org/10.1073/pnas.1232392100

Harashima A, Matsuo Y, Drexler HG, Okochi A, Motoda R, Tanimoto M, Orita K (2005) Transcription factor expression in B-cell precursor-leukemia cell lines: preferential expression of T-bet. *Leuk Res,* **29**, 841-848. https://doi.org/10.1016/j.leukres.2004.12.010

Harden JL, Lewis SM, Lish SR, Suarez-Farinas M, Gareau D, Lentini T, Johnson-Huang LM, Krueger JG, Lowes MA (2016) The tryptophan metabolism enzyme L-kynureninase is a novel inflammatory factor in psoriasis and other inflammatory diseases. *J Allergy Clin Immunol,* **137**, 1830-1840. https://doi.org/10.1016/j.jaci.2015.09.055

Hung WS, Ling P, Cheng JC, Chang SS, Tseng CP (2016) Disabled-2 is a negative immune regulator of lipopolysaccharide-stimulated Toll-like receptor 4 internalization and signaling. *Sci Rep,* **6**, 35343. https://doi.org/10.1038/srep35343

Jablonski KA, Amici SA, Webb LM, Ruiz-Rosado Jde D, Popovich PG, Partida-Sanchez S, Guerau-de-Arellano M (2015) Novel Markers to Delineate Murine M1 and M2 Macrophages. *PLoS One,* **10**, e0145342. https://doi.org/10.1371/journal.pone.0145342

Jackson AP, Berry A, Aslett M, Allison HC, Burton P, Vavrova-Anderson J, Brown R, Browne H, Corton N, Hauser H, Gamble J, Gilderthorp R, Marcello L, McQuillan J, Otto TD, Quail MA, Sanders MJ, van Tonder A, Ginger ML, Field MC, Barry JD, Hertz-Fowler C, Berriman M (2012) Antigenic diversity is generated by distinct evolutionary mechanisms in African trypanosome species. *Proc Natl Acad Sci U S A,* **109**, 3416-3421. https://doi.org/10.1073/pnas.1117313109

Joseph SB, Bradley MN, Castrillo A, Bruhn KW, Mak PA, Pei L, Hogenesch J, O'Connell R M, Cheng G, Saez E, Miller JF, Tontonoz P (2004) LXR-dependent gene expression is important for macrophage survival and the innate immune response. *Cell,* **119**, 299-309. https://doi.org/10.1016/j.cell.2004.09.032

Kawahara K, Hohjoh H, Inazumi T, Tsuchiya S, Sugimoto Y (2015) Prostaglandin E2-induced inflammation: Relevance of prostaglandin E receptors. *Biochim Biophys Acta,* **1851**, 414-421. https://doi.org/10.1016/j.bbalip.2014.07.008

Kay JG, Murray RZ, Pagan JK, Stow JL (2006) Cytokine secretion via cholesterol-rich lipid raft-associated SNAREs at the phagocytic cup. *J Biol Chem,* **281**, 11949-11954. https://doi.org/10.1074/jbc.M600857200

Kemp S (2019) Why science matters. African cattle in the genomic century. In: *The story of cattle in Africa: Why diversity matters* eds Dessie T & Mwai O), pp. 268. International Livestock Research Institute,

Rural Development Administration of the Republic of Korea and the African Union-InterAfrican, Bureau for Animal Resources, Nairobi, Kenya.

Kierstein S, Noyes H, Naessens J, Nakamura Y, Pritchard C, Gibson J, Kemp S, Brass A (2006) Gene expression profiling in a mouse model for African trypanosomiasis. *Genes Immun,* **7**, 667-679. https://doi.org/10.1038/sj.gene.6364345

Kim K, Kwon T, Dessie T, Yoo D, Mwai OA, Jang J, Sung S, Lee S, Salim B, Jung J, Jeong H, Tarekegn GM, Tijjani A, Lim D, Cho S, Oh SJ, Lee HK, Kim J, Jeong C, Kemp S, Hanotte O, Kim H (2020) The mosaic genome of indigenous African cattle as a unique genetic resource for African pastoralism. *Nat Genet,* **52**, 1099-1110. https://doi.org/10.1038/s41588-020-0694-2

Kos J, Sekirnik A, Premzl A, Zavasnik Bergant V, Langerholc T, Turk B, Werle B, Golouh R, Repnik U, Jeras M, Turk V (2005) Carboxypeptidases cathepsins X and B display distinct protein profile in human cells and tissues. *Exp Cell Res,* **306**, 103-113. https://doi.org/10.1016/j.yexcr.2004.12.006

Kramer A, Green J, Pollard J, Jr., Tugendreich S (2014) Causal analysis approaches in Ingenuity Pathway Analysis. *Bioinformatics,* **30**, 523-530. https://doi.org/10.1093/bioinformatics/btt703

Krishnan S, Warke VG, Nambiar MP, Wong HK, Tsokos GC, Farber DL (2001) Generation and biochemical analysis of human effector CD4 T cells: alterations in tyrosine phosphorylation and loss of CD3zeta expression. *Blood,* **97**, 3851-3859. https://doi.org/10.1182/blood.V97.12.3851

Kuriakose S, Onyilagha C, Singh R, Olayinka-Adefemi F, Jia P, Uzonna JE (2019) TLR-2 and MyD88-Dependent Activation of MAPK and STAT Proteins Regulates Proinflammatory Cytokine Response and Immunity to Experimental *Trypanosoma congolense* Infection. *Frontiers in immunology,* **10**, 2673. https://doi.org/10.3389/fimmu.2019.02673

Le Cao K-A, Rohart F, Gonzales I, Dejean S, Gautier B, Bartolo F, Monget P, Coquery J, Yao F, Liquet B. (2016). mixOmics: Omics Data Integration Project. Retrieved from R package version 6.1.1. https://CRAN.R-project.org/package=mixOmics

Lebigot E, Brassier A, Zater M, Imanci D, Feillet F, Therond P, de Lonlay P, Boutron A (2015) Fructose 1,6-bisphosphatase deficiency: clinical, biochemical and genetic features in French patients. *J Inherit Metab Dis,* **38**, 881-887. https://doi.org/10.1007/s10545-014-9804-6

Lejon V, Mumba Ngoyi D, Kestens L, Boel L, Barbe B, Kande Betu V, van Griensven J, Bottieau E, Muyembe Tamfum JJ, Jacobs J, Buscher P (2014) Gambiense human african trypanosomiasis and immunological memory: effect on phenotypic lymphocyte profiles and humoral immunity. *PLoS Pathog,* **10**, e1003947. https://doi.org/10.1371/journal.ppat.1003947

Leppert BJ, Mansfield JM, Paulnock DM (2007) The soluble variant surface glycoprotein of African trypanosomes is recognized by a macrophage scavenger receptor and induces I kappa B alpha degradation independently of TRAF6-mediated TLR signaling. *J Immunol,* **179**, 548-556. https://doi.org/10.4049/jimmunol.179.1.548

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics,* **25**, 2078-2079. https://doi.org/10.1093/bioinformatics/btp352

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics,* **30**, 923-930. https://doi.org/10.1093/bioinformatics/btt656

Liu Q, Zheng J, Yin DD, Xiang J, He F, Wang YC, Liang L, Qin HY, Liu L, Liang YM, Han H (2012) Monocyte to macrophage differentiation-associated (MMD) positively regulates ERK and Akt activation and TNF-alpha and NO production in macrophages. *Mol Biol Rep,* **39**, 5643-5650. https://doi.org/10.1007/s11033-011-1370-5

Livingstone C, Borai A (2014) Insulin-like growth factor-II: its role in metabolic and endocrine disease. *Clin Endocrinol (Oxf),* **80**, 773-781. https://doi.org/10.1111/cen.12446

Loftus RM, Finlay DK (2016) Immunometabolism: Cellular Metabolism Turns Immune Regulator. *J Biol Chem,* **291**, 1-10. https://doi.org/10.1074/jbc.R115.693903

Loftus RT, MacHugh DE, Ngere LO, Balain DS, Badi AM, Bradley DG, Cunningham EP (1994) Mitochondrial genetic variation in European, African and Indian cattle populations. *Anim Genet,* **25**, 265-271. https://doi.org/10.1111/j.1365-2052.1994.tb00203.x

Magez S, Pinto Torres JE, Obishakin E, Radwanska M (2020) Infections With Extracellular Trypanosomes
Require Control by Efficient Innate Immune Mechanisms and Can Result in the Destruction of the
Mammalian Humoral Immune System. *Front Immunol,* **11**, 382.
https://doi.org/10.3389/fimmu.2020.00382

Magez S, Radwanska M, Drennan M, Fick L, Baral TN, Allie N, Jacobs M, Nedospasov S, Brombacher F, Ryffel
B, De Baetselier P (2007) Tumor necrosis factor (TNF) receptor-1 (TNFp55) signal transduction and
macrophage-derived soluble TNF are crucial for nitric oxide-mediated *Trypanosoma congolense*
parasite killing. *J Infect Dis,* **196**, 954-962. https://doi.org/10.1086/520815

Magez S, Radwanska M, Drennan M, Fick L, Baral TN, Brombacher F, De Baetselier P (2006) Interferon-
gamma and nitric oxide in combination with antibodies are key protective host immune factors
during *Trypanosoma congolense* Tc13 Infections. *J Infect Dis,* **193**, 1575-1583.
https://doi.org/10.1086/503808

Martinez-Reyes I, Chandel NS (2020) Mitochondrial TCA cycle metabolites control physiology and disease.
Nat Commun, **11**, 102. https://doi.org/10.1038/s41467-019-13668-3

Matthews KR, McCulloch R, Morrison LJ (2015) The within-host dynamics of African trypanosome
infections. *Philos Trans R Soc Lond B Biol Sci,* **370**. https://doi.org/10.1098/rstb.2014.0288

Mattioli RC, Feldmann U, Hendrickx G, Wint W, Jannin J, Slingenbergh J (2004) Tsetse and trypanosomiasis
intervention policies supporting sustainable animal-agricultural development. *Journal of Food*
Agriculture & Environment, **2**, 310-314

McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq experiments
with respect to biological variation. *Nucleic Acids Res,* **40**, 4288-4297.
https://doi.org/10.1093/nar/gks042

McGettrick AF, O'Neill LA (2013) How metabolism generates signals during innate immunity and
inflammation. *J Biol Chem,* **288**, 22893-22898. https://doi.org/10.1074/jbc.R113.486464

Meade KG, O'Gorman GM, Hill EW, Narciandi F, Agaba M, Kemp SJ, O'Farrelly C, MacHugh DE (2009)
Divergent antimicrobial peptide (AMP) and acute phase protein (APP) responses to *Trypanosoma*
congolense infection in trypanotolerant and trypanosusceptible cattle. *Mol Immunol,* **47**, 196-204.
https://doi.org/10.1016/j.molimm.2009.09.042

Metcalfe SM (2011) LIF in the regulation of T-cell fate and as a potential therapeutic. *Genes Immun,* **12**,
157-168. https://doi.org/10.1038/gene.2011.9

Meyer A, Holt HR, Selby R, Guitian J (2016) Past and Ongoing Tsetse and Animal Trypanosomiasis Control
Operations in Five African Countries: A Systematic Review. *PLoS Negl Trop Dis,* **10**, e0005247.
https://doi.org/10.1371/journal.pntd.0005247

Mills CD, Kincaid K, Alt JM, Heilman MJ, Hill AM (2000) M-1/M-2 macrophages and the Th1/Th2 paradigm.
J Immunol, **164**, 6166-6173. https://doi.org/10.4049/jimmunol.164.12.6166

Mills EL, Kelly B, Logan A, Costa ASH, Varma M, Bryant CE, Tourlomousis P, Dabritz JHM, Gottlieb E, Latorre
I, Corr SC, McManus G, Ryan D, Jacobs HT, Szibor M, Xavier RJ, Braun T, Frezza C, Murphy MP,
O'Neill LA (2016) Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to
Drive Inflammatory Macrophages. *Cell,* **167**, 457-470 e413.
https://doi.org/10.1016/j.cell.2016.08.064

Mohr E, Cunningham AF, Toellner KM, Bobat S, Coughlan RE, Bird RA, MacLennan IC, Serre K (2010) IFN-
{gamma} produced by CD8 T cells induces T-bet-dependent and -independent class switching in B
cells in responses to alum-precipitated protein vaccine. *Proc Natl Acad Sci U S A,* **107**, 17292-17297.
https://doi.org/10.1073/pnas.1004879107

Morias Y, Abels C, Laoui D, Van Overmeire E, Guilliams M, Schouppe E, Tacke F, deVries CJ, De Baetselier P,
Beschin A (2015) Ly6C- Monocytes Regulate Parasite-Induced Liver Inflammation by Inducing the
Differentiation of Pathogenic Ly6C+ Monocytes into Macrophages. *PLoS Pathog,* **11**, e1004873.
https://doi.org/10.1371/journal.ppat.1004873

Morrison LJ, McLellan S, Sweeney L, Chan CN, MacLeod A, Tait A, Turner CM (2010) Role for parasite genetic
diversity in differential host responses to *Trypanosoma brucei* infection. *Infect Immun,* **78**, 1096-
1108. https://doi.org/10.1128/IAI.00943-09

Morrison LJ, Vezza L, Rowan T, Hope JC (2016) Animal African Trypanosomiasis: Time to Increase Focus on
Clinically Relevant Parasite and Host Species. *Trends Parasitol,* **32**, 599-607.
https://doi.org/10.1016/j.pt.2016.04.012
Murray M, Trail JC, D'Ieteren GD (1990) Trypanotolerance in cattle and prospects for the control of
trypanosomiasis by selective breeding. *Rev Sci Tech,* **9**, 369-386.
https://doi.org/10.20506/rst.9.2.506
Murray M, Trail JC, Davis CE, Black SJ (1984) Genetic resistance to African Trypanosomiasis. *J Infect Dis,*
149, 311-319. https://doi.org/10.1093/infdis/149.3.311
Murray PJ (2017) Macrophage Polarization. *Annu Rev Physiol,* **79**, 541-566.
https://doi.org/10.1146/annurev-physiol-022516-034339
Naessens J (2006) Bovine trypanotolerance: A natural ability to prevent severe anaemia and
haemophagocytic syndrome? *Int J Parasitol,* **36**, 521-528
Naessens J, Leak SG, Kennedy DJ, Kemp SJ, Teale AJ (2003) Responses of bovine chimaeras combining
trypanosomosis resistant and susceptible genotypes to experimental infection with *Trypanosoma*
congolense. *Vet Parasitol,* **111**, 125-142. https://doi.org/10.1016/S0304-4017(02)00360-6
Naessens J, Williams DJ (1992) Characterization and measurement of CD5+ B cells in normal and
Trypanosoma congolense-infected cattle. *Eur J Immunol,* **22**, 1713-1718.
https://doi.org/10.1002/eji.1830220708
Ndoutamia G, Mbakasse RN, Brahim A, Khadidja A (2002) Influence de la Trypanosomose à *T. congolense*
sur les paramètres hématologiques, minéraux et protéo-énergétiques chez les chèvres sahéliennes
du Tchad. *Revue Méd. Vét.,* **153**, 395-400
Ned RM, Swat W, Andrews NC (2003) Transferrin receptor 1 is differentially required in lymphocyte
development. *Blood,* **102**, 3711-3718. https://doi.org/10.1182/blood-2003-04-1086
Noel W, Hassanzadeh G, Raes G, Namangala B, Daems I, Brys L, Brombacher F, Baetselier PD, Beschin A
(2002) Infection stage-dependent modulation of macrophage activation in *Trypanosoma*
congolense-resistant and -susceptible mice. *Infect Immun,* **70**, 6180-6187.
https://doi.org/10.1128/IAI.70.11.6180-6187.2002
Nowyhed HN, Huynh TR, Thomas GD, Blatchley A, Hedrick CC (2015) Cutting Edge: The Orphan Nuclear
Receptor Nr4a1 Regulates CD8+ T Cell Expansion and Effector Function through Direct Repression
of Irf4. *J Immunol,* **195**, 3515-3519. https://doi.org/10.4049/jimmunol.1403027
Noyes H, Brass A, Obara I, Anderson S, Archibald AL, Bradley DG, Fisher P, Freeman A, Gibson J, Gicheru M,
Hall L, Hanotte O, Hulme H, McKeever D, Murray C, Oh SJ, Tate C, Smith K, Tapio M, Wambugu J,
Williams DJ, Agaba M, Kemp SJ (2011) Genetic and expression analysis of cattle identifies
candidate genes in pathways responding to *Trypanosoma congolense* infection. *Proc Natl Acad Sci*
U S A, **108**, 9304-9309. https://doi.org/10.1073/pnas.1013486108
O'Gorman GM, Park SD, Hill EW, Meade KG, Coussens PM, Agaba M, Naessens J, Kemp SJ, MacHugh DE
(2009) Transcriptional profiling of cattle infected with Trypanosoma congolense highlights gene
expression signatures underlying trypanotolerance and trypanosusceptibility. *BMC Genomics,* **10**,
207. https://doi.org/10.1186/1471-2164-10-207
O'Gorman GM, Park SD, Hill EW, Meade KG, Mitchell LC, Agaba M, Gibson JP, Hanotte O, Naessens J, Kemp
SJ, MacHugh DE (2006) Cytokine mRNA profiling of peripheral blood mononuclear cells from
trypanotolerant and trypanosusceptible cattle infected with *Trypanosoma congolense*. *Physiol*
Genomics, **28**, 53-61. https://doi.org/10.1152/physiolgenomics.00100.2006
O'Neill LA, Kishton RJ, Rathmell J (2016) A guide to immunometabolism for immunologists. *Nat Rev*
Immunol, **16**, 553-565. https://doi.org/10.1038/nri.2016.70
O'Rawe JA, Ferson S, Lyon GJ (2015) Accounting for uncertainty in DNA sequencing data. *Trends Genet,* **31**,
61-66. https://doi.org/10.1016/j.tig.2014.12.002
Obermajer N, Repnik U, Jevnikar Z, Turk B, Kreft M, Kos J (2008) Cysteine protease cathepsin X modulates
immune response via activation of beta2 integrins. *Immunology,* **124**, 76-88.
https://doi.org/10.1111/j.1365-2567.2007.02740.x
Okwor I, Onyilagha C, Kuriakose S, Mou Z, Jia P, Uzonna JE (2012) Regulatory T cells enhance susceptibility
to experimental *Trypanosoma congolense* infection independent of mouse genetic background.
PLoS Negl Trop Dis, **6**, e1761. https://doi.org/10.1371/journal.pntd.0001761

Onyilagha C, Jia P, Jayachandran N, Hou S, Okwor I, Kuriakose S, Marshall A, Uzonna JE (2015) The B cell
adaptor molecule Bam32 is critically important for optimal antibody response and resistance to
Trypanosoma congolense infection in mice. *PLoS Negl Trop Dis,* **9**, e0003716.
https://doi.org/10.1371/journal.pntd.0003716
Onyilagha C, Kuriakose S, Ikeogu N, Jia P, Uzonna J (2018) Myeloid-Derived Suppressor Cells Contribute to
Susceptibility to *Trypanosoma congolense* Infection by Suppressing CD4(+) T Cell Proliferation and
IFN-gamma Production. *J Immunol,* **201**, 507-515. https://doi.org/10.4049/jimmunol.1800180
Paillard F, Sterkers G, Vaquero C (1990) Transcriptional and post-transcriptional regulation of TcR, CD4 and
CD8 gene expression during activation of normal human T lymphocytes. *EMBO J,* **9**, 1867-1872.
https://doi.org/10.1002/j.1460-2075.1990.tb08312.x
Payne WJA, Hodges J (1997) *Tropical cattle, origins, breeds and breeding policies*. Blackwell Science, Oxford.
Pearce EL, Pearce EJ (2013) Metabolic pathways in immune cell activation and quiescence. *Immunity,* **38**,
633-643. https://doi.org/10.1016/j.immuni.2013.04.005
Pello OM, De Pizzol M, Mirolo M, Soucek L, Zammataro L, Amabile A, Doni A, Nebuloni M, Swigart LB, Evan
GI, Mantovani A, Locati M (2012) Role of c-MYC in alternative activation of human macrophages
and tumor-associated macrophage biology. *Blood,* **119**, 411-421. https://doi.org/10.1182/blood-
2011-02-339911
Pierre C (1906) *L'élevage en Afrique occidentale française*. Gouvernement général de l'Afrique occidentale
française, Inspection de l'agriculture, Paris.
Pinder M, Bauer J, Van Melick A, Fumoux F (1988) Immune responses of trypanoresistant and
trypanosusceptible cattle after cyclic infection with *Trypanosoma congolense*. *Vet Immunol
Immunopathol,* **18**, 245-257. https://doi.org/10.1016/0165-2427(88)90069-4
Rajavel A, Heinrich F, Schmitt AO, Gultas M (2020) Identifying Cattle Breed-Specific Partner Choice of
Transcription Factors during the African Trypanosomiasis Disease Progression Using Bioinformatics
Analysis. *Vaccines (Basel),* **8**. https://doi.org/10.3390/vaccines8020246
Team RC (2018) R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. URL http://www.R-project.org/.
Rege JE (1999) The state of African cattle genetic resources I. Classification framework and identification
of threatened and extinct breeds. *AGRI,* **25**, 1-25
Risco A, del Fresno C, Mambol A, Alsina-Beauchamp D, MacKenzie KF, Yang HT, Barber DF, Morcelle C,
Arthur JS, Ley SC, Ardavin C, Cuenda A (2012) p38gamma and p38delta kinases regulate the Toll-
like receptor 4 (TLR4)-induced cytokine production by controlling ERK1/2 protein kinase pathway
activation. *Proc Natl Acad Sci U S A,* **109**, 11200-11205. https://doi.org/10.1073/pnas.1207290109
Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers differential expression
analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res,* **43**, e47.
https://doi.org/10.1093/nar/gkv007
Roberts CJ, Gray AR (1973) Studies on trypanosome-resistant cattle. II. The effect of trypanosomiasis on
N'dama, Muturu and Zebu cattle. *Trop Anim Health Prod,* **5**, 220-233.
https://doi.org/10.1007/BF02240423
Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinformatics,* **26**, 139-140.
https://doi.org/10.1093/bioinformatics/btp616
Schulman IG (2017) Liver X receptors link lipid metabolism and inflammation. *FEBS Lett,* **591**, 2978-2991.
https://doi.org/10.1002/1873-3468.12702
Seck MT, Bouyer J, Sall B, Bengaly Z, Vreysen MJ (2010) The prevalence of African animal trypanosomoses
and tsetse presence in Western Senegal. *Parasite,* **17**, 257-265.
https://doi.org/10.1051/parasite/2010173257
Sileghem M, Flynn JN (1992a) Suppression of interleukin 2 secretion and interleukin 2 receptor expression
during tsetse-transmitted trypanosomiasis in cattle. *Eur J Immunol,* **22**, 767-773.
https://doi.org/10.1002/eji.1830220321
Sileghem M, Flynn JN (1992b) Suppression of T-cell responsiveness during tsetse-transmitted
trypanosomiasis in cattle. *Scand J Immunol Suppl,* **11**, 37-40. https://doi.org/10.1111/j.1365-
3083.1992.tb01616.x

Sileghem MR, Flynn JN, Saya R, Williams DJ (1993) Secretion of co-stimulatory cytokines by monocytes and
macrophages during infection with *Trypanosoma* (Nannomonas) *congolense* in susceptible and
tolerant cattle. *Vet Immunol Immunopathol,* **37**, 123-134. https://doi.org/10.1016/0165-
2427(93)90060-H

Swallow BM (2000) *Impacts of Trypanosomiasis on African agriculture*. PAAT Technical and scientific series
2. FAO (Food and Agricultural Organization of the United Nations), Rome, Italy.

Taiwo VO, Anosa VO (2000) *In vitro* erythrophagocytosis by cultured macrophages stimulated with
extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and
N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosmoa vivax*. *Onderstepoort*
Journal of Veterinary Research, **67**, 273-287

Taylor KA (1998) Immune responses of cattle to African trypanosomes: protective or pathogenic? *Int J*
Parasitol, **28**, 219-240. https://doi.org/10.1016/S0020-7519(97)00154-9

Taylor KA, Lutje V, Kennedy D, Authie E, Boulange A, Logan-Henfrey L, Gichuki B, Gettinby G (1996)
Trypanosoma congolense: B-lymphocyte responses differ between trypanotolerant and
trypanosusceptible cattle. *Exp Parasitol,* **83**, 106-116. https://doi.org/10.1006/expr.1996.0054

Taylor KA, Mertens B (1999) Immune response of cattle infected with African trypanosomes. *Mem Inst*
Oswaldo Cruz, **94**, 239-244

Trail JC, d'Ieteren GD, Maille JC, Yangari G (1991) Genetic aspects of control of anaemia development in
trypanotolerant N'Dama cattle. *Acta Trop,* **48**, 285-291. https://doi.org/10.1016/0001-
706X(91)90016-D

Trail JC, Wissocq N, d'Ieteren GD, Kakiese O, Murray M (1994) Quantitative phenotyping of N'Dama cattle
for aspects of trypanotolerance under field tsetse challenge. *Vet Parasitol,* **55**, 185-195.
https://doi.org/10.1016/0304-4017(94)00649-W

Traore-Leroux T, Fumoux F, Pinder M (1987) High density lipoprotein levels in the serum of
trypanosensitive and trypanoresistant cattle. Changes during *Trypanosoma congolense* infection.
Acta Trop, **44**, 315-323

Uilenberg G (1998) A field guide for the diagnosis, treatment and prevention of African animal
trypanosmosis. FAO, Rome.

Uzonna JE, Kaushik RS, Zhang Y, Gordon JR, Tabel H (1998) Experimental murine *Trypanosoma congolense*
infections. II. Role of splenic adherent CD3+Thy1.2+ TCR-alpha beta- gamma delta- CD4+8- and
CD3+Thy1.2+ TCR-alpha beta- gamma delta- CD4-8- cells in the production of IL-4, IL-10, and IFN-
gamma and in trypanosome-elicited immunosuppression. *J Immunol,* **161**, 6189-6197

Van de Laar L, Coffer PJ, Woltman AM (2012) Regulation of dendritic cell development by GM-CSF:
molecular control and implications for immune homeostasis and therapy. *Blood,* **119**, 3383-3393.
https://doi.org/10.1182/blood-2011-11-370130

Van der Waaij EH, Hanotte O, Van Arendonk JAM, Kemp SJ, Kennedy D, Gibson A, Teale A (2003) Population
parameters for traits defining trypanotolerance in a F2 cross of N'Dama and Boran cattle. *Livest*
Sci, **84**, 219-230. https://doi.org/10.1016/S0301-6226(03)00078-2

Vincendeau P, Bouteille B (2006) Immunology and immunopathology of African trypanosomiasis. *An Acad*
Bras Cienc, **78**, 645-665. https://doi.org/10.1590/S0001-37652006000400004

Vulcano M, Albanesi C, Stoppacciaro A, Bagnati R, D'Amico G, Struyf S, Transidico P, Bonecchi R, Del Prete
A, Allavena P, Ruco LP, Chiabrando C, Girolomoni G, Mantovani A, Sozzani S (2001) Dendritic cells
as a major source of macrophage-derived chemokine/CCL22 in vitro and in vivo. *Eur J Immunol,*
31, 812-822. https://doi.org/10.1002/1521-4141(200103)31:3<812::AID-
IMMU812>3.0.CO;2-L

Wang LM, Zhang Y, Li X, Zhang ML, Zhu L, Zhang GX, Xu YM (2018) Nr4a1 plays a crucial modulatory role in
Th1/Th17 cell responses and CNS autoimmunity. *Brain Behav Immun,* **68**, 44-55.
https://doi.org/10.1016/j.bbi.2017.09.015

Wang R, Dillon CP, Shi LZ, Milasta S, Carter R, Finkelstein D, McCormick LL, Fitzgerald P, Chi H, Munger J,
Green DR (2011) The transcription factor Myc controls metabolic reprogramming upon T
lymphocyte activation. *Immunity,* **35**, 871-882. https://doi.org/10.1016/j.immuni.2011.09.021

Whitehead A, Crawford DL (2006) Neutral and adaptive variation in gene expression. *Proc Natl Acad Sci U*
S A, **103**, 5425-5430. https://doi.org/10.1073/pnas.0507648103

Wickham H (2016) *ggplot2: Elegant graphics for data analysis*. Springer International Publishing, New York.

Wilson KC, Center DM, Cruikshank WW (2004) The effect of interleukin-16 and its precursor on T
lymphocyte   activation   and   growth.   *Growth   Factors,*   **22**,   97-104.
https://doi.org/10.1080/0897719041001704679

Xu J, Flaczyk A, Neal LM, Fa Z, Eastman AJ, Malachowski AN, Cheng D, Moore BB, Curtis JL, Osterholzer JJ,
Olszewski MA (2017) Scavenger Receptor MARCO Orchestrates Early Defenses and Contributes to
Fungal   Containment   during   Cryptococcal   Infection.   *J Immunol,*   **198**,   3548-3557.
https://doi.org/10.4049/jimmunol.1700057

Yoon MS (2017) The Role of Mammalian Target of Rapamycin (mTOR) in Insulin Signaling. *Nutrients,* **9**.
https://doi.org/10.3390/nu9111176

Zamboni DS, Lima-Junior DS (2015) Inflammasomes in host response to protozoan parasites. *Immunol Rev,*
265, 156-171. https://doi.org/10.1111/imr.12291

Zoldan K, Moellmer T, Schneider J, Fueldner C, Knauer J, Lehmann J (2014) Increase of CD25 expression on
bovine neutrophils correlates with disease severity in post-partum and early lactating dairy cows.
Dev Comp Immunol, **47**, 254-263. https://doi.org/10.1016/j.dci.2014.08.002

1357